The Lifespan Trajectories of Brain Activities Related to Cognitive Control

Zhenghan Li[1,2,3,4], Isaac T. Petersen[5], Lingxiao Wang[6,7,3], Joaquim Radua[8,9,10], Guo-
chun Yang[5,11,4*], Xun Liu[4,12]

[1] Institute of Brain Science and Department of Physiology, School of Basic Medical
Sciences, Hangzhou Normal University, Hangzhou, China

[2] Zhejiang Philosophy and Social Science Laboratory for Research in Early Develop-
ment and Childcare, Hangzhou Normal University, Hangzhou, China

[3] Zhejiang Key Laboratory for Research in Assessment of Cognitive Impairments,
Hangzhou, China

[4] Key Laboratory of Behavioral Science, Institute of Psychology, Chinese Academy of
Sciences, Beijing, China

[5] Department of Psychological and Brain Sciences, University of Iowa, Iowa City, IA,
USA

[6] Centre for Cognition and Brain disorders, The Affiliated Hospital of Hangzhou Nor-
mal University, Hangzhou, Zhejiang Province, China

[7] Institute of Psychological Science, Hangzhou Normal University, Hangzhou,
Zhejiang Province, China

[8] Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), CIBERSAM,
University of Barcelona, Barcelona, Spain

[9] Institute of Psychiatry, Psychology and Neuroscience, King's College London, Lon-
don, United Kingdom

[10] Department of Clinical Neuroscience, Karolinska Institute, Stockholm, Sweden

[11] Cognitive Control Collaborative, University of Iowa, Iowa City, IA, USA

[12] Department of Psychology, University of Chinese Academy of Sciences, Beijing,
China

***Correspondence:** guochun-yang@uiowa.edu

Classification: Biological Sciences/Psychological and Cognitive Sciences

Keywords: Cognitive control; Lifespan; Brain activity; Neuroimaging meta-analysis;
Heterogeneity

Abstract

Cognitive control plays a pivotal role in guiding human goal-directed behavior. While existing studies have documented an inverted U-shaped trajectory of cognitive control both behaviorally and anatomically, little is known about the corresponding changes in functional brain activation with age. To bridge this gap, we conducted a comprehensive meta-analysis of 129 neuroimaging studies using conflict tasks, encompassing 3,388 participants aged from 5 to 85 years old. We have three major findings: 1) The inverted U-shaped trajectory is the predominant pattern; 2) Cognitive control-related brain regions exhibit heterogeneous lifespan trajectories: the frontoparietal control network follows inverted U-shaped trajectories, peaking between 24 and 40 years, while the dorsal attention network demonstrates no clear trajectories; 3) Both the youth and the elderly show weaker brain activities and greater left laterality than young to middle-aged adults. These results reveal the lifespan trajectories of cognitive control, highlighting heterogeneous fluctuations in brain networks with age.

Introduction

The cognitive abilities of human beings dynamically change throughout the entire lifespan, experiencing rapid development in the early stages, and gradual decline in the later period of life. As one of the most fundamental cognitive functions, cognitive control is deeply engaged in various domains of high-level capabilities that humans greatly outperform other species, such as decision making, planning and problem solving[1]. Cognitive control refers to the cognitive processes that enable individuals to manage and regulate their attention, thoughts, and actions, which plays a vital role in goal-directed behavior, allowing us to focus on the target and ignore distractors[2,3]. For instance, cognitive control enables us to concentrate on reading in a library despite the presence of people chatting nearby.

Cognitive control provides fundamental support for normal human behaviors. Young adults typically maintain an optimal mature level of cognitive control[4]. However, the youth (including children and adolescents, less than 18 years) and elderly (60 years and older) individuals may struggle with behavioral problems because of their suboptimal cognitive control system[5-10]. While the state-of-art progress of cognitive/behavioral changes has been well-documented and shaped how the diagnosis of developmental/ageing related disorders[11], the change of related neural system has been under-investigated. Understanding how the neural underpinning of cognitive control changes

over the lifespan can yield valuable insights into the developmental and aging mecha-

nisms of the human brain. This knowledge may assist in customizing cognitive training

strategies based on related brain regions and their activities[12].

Researchers generally believe that cognitive control ability follows an inverted U-

shaped trajectory across the lifespan[5,13,14,17]. This inverted U-shaped trajectory has been

generally supported by behavioral and anatomical evidence. The Eriksen Flanker task

(requiring participants to respond to central stimuli while ignoring flanking distractions)

has been widely utilized to detect cognitive control across the lifespan[15], and results

suggest a clear U-shape trajectory of conflict cost (measured by worsened behavioral

performance, e.g., reaction time, in incongruent compared to congruent conditions)

with age[16-18]. Similar results have been observed in other conflict tasks, such as color-

word Stroop (requiring participants to name the ink color of a word that is incongruent

with the word's semantic meaning)[16]. Recent large-cohort studies have also found that

gray and white matter volumes across all brain regions exhibit overall inverted U-

shaped trajectories with age, with the gray matter volume peaking at early adolescence

and the white matter volume peaking at young adulthood[19,20]. During late adulthood,

normal ageing yields a protracted decline of brain structure, with the volumes of both

gray matter and white matter reduced[14]. Consistently, the gray matter volumes of the

frontal and parietal regions, which are essential in cognitive control tasks[21], have also

been found to increase during early childhood and atrophy in early elderly age[22].

However, it remains largely unknown how brain activities related to cognitive

control change over the lifespan. Previous research has primarily focused on brain ac-

tivities in either youths or elderly adults, rather than examining changes across the en-

tire lifespan. With conflict paradigms, children and adolescents are often found to have

lower brain activity than young adults in frontoparietal regions (refs.[23-28], but see Bunge

et al. [29]). For example, a study utilizing the Flanker task revealed that children aged

8−12 years had reduced activation in dorsolateral prefrontal regions compared to young

adults, suggesting an immature cognitive control system[27]. However, brain activity dif-

ferences in elderly adults as compared to young adults during cognitive control tasks

have been less consistently reported[14]. Some studies have found that elderly adults have

lower neural activity in frontoparietal regions than young adults[30-32], possibly because

elderly adults may be unable to engage in an equal level of control-related activity due

to functional decline. On the other hand, other studies have found that elderly adults

may exhibit greater brain activity in frontoparietal regions than young adults[33,34], pos-

sibly because they have recruited additional brain regions to compensate for their de-

creased efficiency in utilizing control resources. Adding to the debate, it has been pro-

posed that the cognitive control function in elderly adults might not decline at all[35,36].

These conflicting findings underscore the complexity of understanding age-related dif-

ferences in cognitive control.

Few studies have directly tested the change of brain activities related to cognitive

control across the lifespan. One existing study observed a positive association between

the activation of the bilateral prefrontal cortex and age[37]. Given the relatively small

sample size (N = 30), the reliability of these findings is somewhat limited. As a result,

it is difficult to draw a clear conclusion about how brain activities related to cognitive

control change across the entire age range.

One direct way to test the lifespan trajectory of brain activation related to cognitive

control is to conduct a large cohort of neuroimaging study with participants covering a

wide age range. To the best of our knowledge, such studies have not yet been conducted.

An alternative approach is to utilize meta-analyses to combine the results of existing

studies targeting different age groups. Compared to the large cohort studies, meta-anal-

yses are more accessible and resource-saving. In addition, meta-analyses can increase

the statistical power and generalizability by combining various studies, reducing heter-

ogeneity and bias from individual studies' methods, populations, or confounding vari-

ables[38]. Importantly, meta-analyses can reveal patterns not evident in individual studies,

like nonlinear effects or interactions[38]. Several neuroimaging meta-analyses have ex-

amined age-related changes of cognitive control brain activity[31,39,40], but limitations

such as incomplete age coverage and insufficient studies in certain age ranges prevent

them from appropriately answering questions about the lifespan trajectory of cognitive

control functions[31,39,40]. These studies have primarily focused on spatial convergence

and/or diversity of coordinates across different age ranges, offering limited insights into

lifespan trajectories of activity strength. Although a few studies have attempted para-

metric meta-regression to examine the age-related differences in cognitive control-re-

lated brain activity[39,41], they have been constrained by utilizing linear models that may

overlook non-linear trajectory patterns, such as the inverted U-shaped trend.

The goal of this study is to provide a comprehensive examination to reveal the

lifespan trajectory of brain activities responsible for cognitive control. Instead of en-

compassing various aspects of cognitive control, we focus on conflict processing for

several reasons. First, conflict processing reflects the fundamental cognitive control

ability to maintain a goal while avoiding distractions[2]. Second, its mechanisms in young

adults are relatively well-known, with the frontoparietal and cingulo-opercular net-

works engaged[21,42], providing a baseline reference for our study. Third, conflict tasks

with neuroimaging data have been widely applied to both younger and elderly groups,

making a systematic meta-analysis feasible. Lastly, different conflict tasks share key

components of cognitive control, such as conflict monitoring[43] and inhibitory control[8],

which enables us to conduct effect-size-based meta-analyses using the congruency ef-

fect (i.e., the contrast between incongruent and congruent/neutral conditions). Including

other sub-processes of cognitive control may introduce heterogeneity and make effect

sizes incomparable.

Previous research suggests that better cognitive control-related performance is of-

ten associated with greater brain activity, especially in the prefrontal cortex[44]. Therefore,

we hypothesized that brain activities related to cognitive control might follow an in-

verted U-shaped trajectory like that of behavior patterns. Additionally, we hypothesized

that different brain networks may show some heterogeneity in their lifespan trajectories.

Results

Sample Description

A total of 3,611 articles were identified including 3,484 articles searched from the da-

tabase, 111 articles adopted from a previous study[21], and 16 articles searched from the

references of crucial articles. After excluding duplicates and applying exclusion criteria,

119 articles including 129 studies with 3,388 participants and 1,579 brain activation

foci, were included in this meta-analytic study (Supplementary Fig. S1 and Table S1).

The average age of participants ranged from 8 to 74 years, with the individual age rang-

ing from 5 to 85 years. A demonstration of the age distribution for each included study

is presented in Supplementary Fig. S2.

Regions Related to Cognitive Control Identified by both ALE and SDM

To enhance the replicability and robustness of our finding and enable direct comparison

with a previous study[21], we conducted the mean analysis across all studies with the

activation likelihood estimation (ALE) and seed-based d mapping (SDM) approaches

separately. First, we performed the single dataset analysis with the GingerALE software

to explore the brain regions consistently reported in all the included studies (see section

"*Activation Likelihood Estimation (ALE)*" in Methods). Results showed significant ac-

tivation in the frontoparietal regions, including the left dorsolateral prefrontal cortex,

bilateral frontal eye field, right inferior frontal gyrus and bilateral inferior parietal lob-

ule; the cingulo-opercular regions, including the supplementary motor area and bilateral

insula; and other regions, including the left inferior temporal gyri (Fig. 1A and Supple-

mentary Table S2). Second, we calculated the average brain activation based on the

effect sizes reported in all studies using the SDM with permutation of subject images

(SDM-PSI) software (see section "*Mean Analyses Across all Studies*" in Methods).

Similar to the ALE results, we found significant activation in the frontoparietal regions

(left inferior parietal lobule, right inferior frontal gyrus, and right middle frontal gyrus),

the cingulo-opercular regions (left anterior cingulate cortex), and other regions includ-

ing bilateral inferior temporal gyrus, right caudate nucleus, right cerebellum, and left

anterior thalamic projections (Fig. 1B and Supplementary Table S3). The count of

voxels revealed that the overlapped area (3,496 voxels, Fig. 1C) accounted for 96.3%

of the regions from the ALE analysis (3,635 voxels), and 15.0% of the regions from the

SDM analysis (23,240 voxels). This result suggested that SDM analysis could be a re-

liable approach for detecting brain regions, thereby laying a solid foundation for using

this approach in subsequent analyses. In addition, a robustness analysis suggested that

age does not influence the mean results of SDM analysis (Supplementary Fig. S3 and

Table S2).

187   [Fig. 1]

Trajectories of Cognitive Control Regions Identified in the Mean Analysis

Having identified nine brain regions in the mean analysis using SDM-PSI, we pro-

ceeded to explore how the activation levels of these regions change with age. To this

end, we extracted brain activity data from all studies for each identified region. Before

performing the meta-regression, we verified the effectiveness of using mean age as a

predictor with a simulation approach (Supplementary Note S1 and Fig. S4). We then

subjected the extracted data to separate generalized additive model (GAM) analyses,

factoring in the confounding covariates (see section "*Generalized Additive Model*

(GAM) Fitting" in Methods). The analyses (Supplementary Table S4) revealed signifi-

cant age-related changes in the activation levels of 4 out of the 9 regions (Fig. 2, l-ACC,

r-IFG, l-ITG, and r-CN). Visualization of the trajectories suggests that these regions

showed inverted U-shaped patterns. The significance of their inverted U-shaped trajec-

tories was further examined using a two-line test approach[45]. Results suggest that all

four regions involve an increase in activity from childhood to young adulthood (ap-

proximately up to 30 years of age), and three of them (r-IFG, l-ITG and r-CN) showed

consistent decrease in the later stages of age (Supplementary Note S2). The other five

regions (Fig. 2, l-IPL, r-ITG, r-MFG, r-CB and l-ATP) showed no significant age-re-

lated changes. Notably, none of the clusters showed significant publication bias based

on Egger's test (ps > 0.86), and they all showed low between-study heterogeneity (τs <

0.13, Qs < 12.11, I²s < 25%). This indicates that the observed results are not likely

influenced by biased reporting or substantial variability in the included studies. A ro-

bustness analysis further ruled out the potential influence of including unpublished and

non-English studies in this study (Supplementary Note S4 and Fig. S5).

211   [Fig. 2]

Note that some regions are large, containing up to over 9,000 voxels. This could

obscure the distinct trajectories specific to different subregions, which are crucial in

understanding the hierarchical patterns of lifespan trajectories. We address this by using

a more granular approach below.

Detecting Different Trajectories of Whole Brain Activities

To explore various possibilities of lifespan trajectories, we grouped studies by their

mean age into the youth, young to middle-aged, and elderly groups, and then conducted

several contrast analyses (see section "*Contrast Analysis*" in Methods). These analyses

did not reveal any regions that exhibited significantly higher or lower brain activity in

the youth compared to others (the combination of young to middle-aged and older

adults) (Table 1). Similarly, we did not observe any regions with higher or lower brain

activity in older adults compared to others (the combination of the youth and young to

middle-aged adults) (Table 1). These results excluded the possibilities of increase/de-

crease-then-stable and stable-then-increase/decrease trajectories (Fig. 3, panels D, F, G,

and I). In addition, we failed to observe any region showing lower activity in young to

middle-aged adults than others (the combination of the youth and older adults), and

thereby ruled out the possibility of the upright U-shaped trajectory (Table 1, Fig. 3E).

[Fig. 3]

However, we identified greater activity in young to middle-aged adults compared

to others in the frontoparietal regions, including bilateral inferior frontal gyrus and bi-

lateral inferior parietal lobule; the cingulo-opercular regions, including left supplemen-

tary motor area, left insula, and right middle cingulate cortex; and a subcortical re-

gion—right caudate nucleus (Fig. 4 and Table 1). This result essentially supports an

inverted U-shaped trajectory. Notably, none of the clusters showed significant publica-

tion bias based on Egger's test (ps > 0.79), and they all showed low between-study

heterogeneity (τs < 0.17, Qs < 12.21, I^2s < 25%). This indicates that the observed results

are not likely influenced by biased reporting or substantial variability in the included

studies. Consistently, further contrast analyses revealed that the young to middle-aged

adults showed greater activity than both the youth (Supplementary Fig. S6) and the

elderly (Supplementary Fig. S7).

Fitting the Lifespan Trajectories with the GAM

For each region identified from the contrast between young to middle-aged adults and

others, the GAM could fit the data significantly with a smooth curve (Fig. 4), with

degrees of freedom varying from 2.9 to 6.7. Peak ages of the inverted U-shaped trajec-

tories were between 24.5 and 39.4 years. Detailed statistics are shown in Supplementary

Table S4.

Model Simplification of the Lifespan Trajectories

Considering the GAM may overfit the data, we fitted the results with simpler models

commonly adopted in lifespan developmental literature[46-48], including the quadratic,

cubic, square root and quadratic logarithmic models, and estimated the goodness of fit

by comparing their Akaike information criterion (AIC) (see section "*Model Simplifica-*

tion and Model Comparison" in Methods). Results showed that the quadratic model

provided the best fit for capturing the age-related changes in left inferior parietal lobule,

while the square root model demonstrated the best goodness of fit for all the other re-

gions (Supplementary Table S5). We also calculated the peak age for each region based

on the optimal model, and results showed that the peak ages ranged from 33.3 to 40.0

years. In addition, we found the peak ages obtained from the above optimal model (i.e.,

square root or quadratic models) and the GAM are consistent, $r = 0.71$, $p = 0.032$, 95%

CI = [0.09 0.93]. See Fig. 4 and Supplementary Table S6 for details.

261   [Fig. 4]

Fitting the Whole Brain with Square Root and Linear Models

Based on model comparisons, we found that the square root model provided the best

goodness of fit for the age-related change of the brain activities. To supplement the

inverted U-shaped results from the contrast analysis, the square root function was then

submitted to whole-brain meta-regression analyses in SDM-PSI (see section "*Meta-*

regression Analyses" in Methods).

By fitting the activation over the whole brain, we found seven significant brain

regions, including the bilateral inferior frontal gyrus, right angular, left inferior parietal

lobule, right insula, right caudate nucleus, and left anterior thalamic projections (Fig.

5A and Supplementary Table S7). These results further supported the existence of the

inverted U-shaped regions we initially identified (Fig. 4).

In addition, we explored the whole-brain trajectories with a linear meta-regression

(see section "*Meta-regression Analyses*" in Methods). However, no significant regions

were observed (Fig. 5B), even under a more tolerant threshold of uncorrected $p < 0.01$.

276   [Fig. 5]

Dissociated Brain Networks with Distinct Lifespan Trajectories

We note that the inverted U-shaped regions constitute only part of the cognitive control-

related regions (Fig. 2), and the remaining regions show a less clear trajectory. To fur-

ther elucidate the spatial distribution of brain regions following these different trajec-

tory patterns, we used the results from the mean SDM analysis (Fig. 1B) as the mask

and replotted the results of the contrast analysis between young to middle-aged adults

and others with two different thresholds, and then compared them with the Yeo's 7-

network atlas[49] (Fig. 6). Visualization of the spatial distribution patterns revealed a dis-

sociation between the middle frontal gyrus and its adjacent rostral and caudal areas (Fig.

6A). Moreover, the distribution of inverted U-shaped regions was more consistent with

the frontoparietal control network (FPCN), and the distribution of non-U-shaped re-

gions was more closely related to the dorsal attention network (DAN, Fig. 6B). The

count of voxels revealed that a numerically larger portion of the inverted U-shaped re-

gions overlapped with the FPCN (2,538 voxels) than with the DAN (932 voxels), while

the overlap with the cingulo-opercular network (CON) was in between (1,867 voxels).

Conversely, a numerically larger portion of the non-U-shaped regions overlapped with

the DAN (3,009 voxels) than with the FPN (2,447 voxels), while the overlap with the

CON was lower (1,594 voxels).

295   [Fig. 6]

To further investigate the potential functional difference between the two sets of

brain regions, we decoded the related terms with the Neurosynth decoder[41] (see section

"*Neurosynth Decoding Analysis*" in Methods). Results showed that the inverted U-

shaped regions were related to the term "cognitive control"($r = 0.187$), but were less so

to "attentional" ($r = 0.100$) and "monitoring" ($r = 0.072$). Note the keyword "monitor-

ing" refers to the major function of the cingulo-opercular network[21]. On the other hand,

the non-U-shaped regions were related to the term "attentional" ($r = 0.240$) but were

less so to "monitoring" ($r = 0.090$) and "control" ($r = 0.062$) (Supplementary Table S8).

This result suggests that the inverted U-shaped and non-U-shaped regions may be as-

sociated with cognitive control and attention, respectively, which is consistent with the

frontoparietal control network and dorsal attention network as identified in the atlas

overlapping analysis.

Lifespan Trajectory of the Laterality

We also tested how the laterality of the brain activity changes with age (see section

"*Laterality Analysis*" in Methods). We first modeled the laterality trajectory using the

GAM. Results showed a significant model fitting, $F(3.0, 3.7) = 3.49$, $p = 0.012$, $R^2 =$

0.24. Moreover, we fitted the data with the four simplified models (i.e., the quadratic,

cubic, square root, and quadratic logarithmic models). The results showed that a square

root function provided the best goodness of fit, with the $\beta_{\text{sqrt(age)}} = -0.68$ (95% CI =

$[-1.02, -0.33]$), $p < 0.001$. The two-line test suggests the hypothetical peaks from both

models did not reach significance (Supplementary Note S2). A visually upright U-

shaped trajectory indicated the youth and elderly adults tended to be more left-lateral-

ized across the whole brain. A further comparison of the relative levels of laterality

across different age groups revealed that both the youth and the elderly groups exhibited

greater left lateralization than the young to middle-aged adult group (Supplementary

Note S3). This left-lateralized pattern could be illustrated by the brain map estimated

with voxel-wise laterality calculation (Fig. 7).

323   [Fig. 7]

Discussion

The present study yielded three primary findings: 1) Among different possible trajec-tories, only the inverted U-shaped trajectories were reliably observed across the whole brain; 2) The cognitive control-related brain regions exhibit heterogeneous lifespan tra-jectories: the frontoparietal control network (such as the inferior frontal gyrus and in-ferior parietal lobule) follows inverted U-shaped trajectories, peaking between 24 and 40 years, while the dorsal attention network (such as the frontal eye field and superior parietal lobule) demonstrates less clear trajectories with age; 3) The youth and the el-derly demonstrate weaker brain activities and a relatively greater extent of left laterality compared to the young to middle-aged adults. These results provide strong evidence for the existence of cognitive control regions exhibiting inverted U-shaped trajectory, and also show the heterogenous lifespan trajectories in different brain regions.

The Inverted U-shaped Trajectory of Brain Activity Related to Cognitive Control

The main finding is that a wide range of cognitive control regions follow inverted U-shaped lifespan trajectories, but no regions showed decrease-then-stable (Fig. 3D), up-right U-shaped (Fig. 3E), stable-then-increase (Fig. 3F), increase-then-stable (Fig. 3G), stable-then-decrease (Fig. 3I), or linear trajectories (Fig. 3A and 3C).

The greater activation in the frontoparietal control network among young to mid-dle-aged adults compared to the youth and the elderly supports the notion that cognitive

control abilities may not be fully developed in children and may decline in the elderly.

This finding is consistent with the idea that the cognitive control system is most effec-

tive in young adulthood, suggesting a possible correlation between the higher functional

activations in the brain and the superior performance of young adults on cognitive con-

trol tasks[50]. Consistently, a previous study[29] showed that behavioral performance (suc-

cess of interference suppression) is positively correlated with the activity in frontal re-

gions. Similar patterns have been repeatedly reported[51,52], although the opposite results

have also been observed[53].

The inverted U-shaped trajectory of brain activation might be associated with the

development of brain structure and functional changes. First, it may reflect the well-

documented structural changes that occur in these regions across the lifespan, which

include synaptic pruning, myelination, cortical thinning, and white matter matura-

tion[19,54]. For example, the density of dopamine receptors increases during adolescence

and young adulthood and subsequently declines with age[55]. These changes can affect

the efficiency and connectivity of neural circuits within the frontoparietal control net-

work[13,56]. Second, the inverted U-shaped trajectory may also arise from functional

changes resulting from the modulation of neurotransmitters, hormones, and environ-

mental factors[57]. Understanding change patterns of brain structure and function is crit-

ical for developing interventions and treatments aimed at improving cognitive control

abilities across the lifespan.

The present results further revealed that the inverted U-shaped lifespan trajectories

of cognitive control regions are not uniform. The GAM fitting results (Fig. 4, Supple-

mentary Table S4) showed that the subregions exhibited varying degrees of association

with age. Moreover, the model simplification demonstrated different underlying trajec-

tory curves, with most regions showing a skewed shape that could best be fitted with a

square root model, except that one region showed a symmetric quadratic shape. The

quadratic lifespan trajectory has been well-documented in previous studies[19,46,48,58],

while the application of the square root model has been relatively rare[59]. The square

root model can better capture the early peak in the trajectory. We also identified differ-

ent peak ages for those regions, ranging from 24 to 40 years, suggesting that cognitive

control regions may not develop at the same rate.

Hierarchical development trajectories in different brain networks

Previous research has indicated that the attentional orientation function is preserved

during ageing[35,60]. Consistently, we found that the dorsal attention network regions un-

derlying the attentional orientation showed no significant age-related change, in con-

trast to the inverted U-shaped trajectory in frontoparietal control network regions. In

addition, we observed that the supporting regions mediating top-down control with mo-

tor[61] (right cerebellum, Fig. 2) and sensory[62] (left anterior thalamic projections, Fig. 2)

functions also lack the sensitivity to age. The dissociation across regions may reflect

hierarchical associations with age on brain function.

The brain regions are organized in a functional hierarchy, with the frontoparietal

control network at the highest level. It acts as a hub that interacts with other systems,

including the dorsal attention network[63]. The cingulo-opercular network did not present

a clear dissociation of trajectory patterns, possibly suggesting its intermediate position

between the frontoparietal control and dorsal attention networks[64]. During conflict tasks,

these networks function in a hierarchical manner. The frontoparietal control network

maintains task goals and resolves conflicts, the cingulo-opercular network monitors

conflict, and the dorsal attention network directs attention towards task-relevant stim-

uli[65]. Even within the prefrontal cortex itself, a hierarchical organization exists, with

middle frontal areas occupying the peak position[66]. This is in line with our finding that

the middle frontal cortex is dissociated from rostral and caudal frontal regions (Fig. 6A).

In addition, previous research suggests that the frontoparietal control network can be

further divided[67], with the rostral and caudal frontal regions observed in our study align-

ing closely with the sub-network that connects more strongly with the dorsolateral at-

tention network. This may explain why some areas within the frontal region do not

show age-related changes.

Furthermore, different brain regions exhibit different age-related changes. Higher-

order regions typically have more complex lifespan trajectories[58] and reach peaks dur-

ing later periods[56,68]. Specifically, prefrontal control regions are among the last to ma-

ture and one of the earliest to decline[5,13,69]. Therefore, the different lifespan trajectory

patterns among different networks likely reflect their hierarchical positions of age-re-

lated changes.

Implications for the Compensatory and Asymmetry Reduction Theories

Critically, there was no region showing higher activity in the elderly compared to the

young to middle-aged adults, but we observed several regions showing the opposite

(Supplementary Note S3 and Table S9). The results persisted after we controlled the

behavioral congruency effect (Supplementary Note S4 and Fig. S8), thereby ruling out

the possibility of weaker brain activity associated with poor behavioral performance in

the elderly. The observed decrease in brain activation among the elderly might be at-

tributed to several interrelated factors. First, cognitive control regions, especially the

frontal area, tend to shrink with age, leading to a reduction in overall brain volume and

potential loss of synaptic integrity[70]. This shrinkage can impair the brain's ability to

effectively process and manage complex tasks. Another significant factor is the impair-

ment of neurovascular coupling, the relationship between neuronal activity, synaptic

function, and subsequent blood flow, which disrupts the brain's ability to maintain op-

timal function during cognitive tasks[71,72]. Furthermore, the decrease in cerebral blood

flow with age can diminish the delivery of essential nutrients and oxygen to the brain,

impairing its overall functionality[73]. These changes could lead to regional abnormali-

ties, such as blood flow, blood volume, metabolic rate, or BOLD-derived physiologic

proxies like the fractional amplitude of low-frequency fluctuation and regional homo-

geneity[74]. Future studies may validate and extend our study by adopting the age-sensi-

tive regions we observed and testing other measurements, such as resting-state data,

which are more easily collected in large-scale studies involving children and the elderly

compared to task-based activations.

The compensatory theory[10] proposes that the elderly recruit additional brain re-

gions to compensate for age-related cognitive decline, but our results did not show this

pattern. We suggest that the absence of compensatory upregulation in frontoparietal

regions among the elderly observed in our study might be attributed to limited available

resources when cognitive control related brain regions are already fully engaged[75,76].

Previous research has shown that younger adults recruit lower activity in frontoparietal

regions during the congruent condition but significantly greater activities during the

incongruent condition. In contrast, older adults already show a relatively higher activa-

tion during the congruent condition, leaving limited capacity for further increases in

activation during the incongruent condition[32]. This is consistent with our findings,

which are based on the contrast between incongruent and congruent conditions.

Moreover, the nature of the task investigated might influence whether there is an up-

regulation in cognitive control regions with age. Upregulation in the frontal regions

usually compensates for memory and sensory declination due to deficits in the hippo-

campus and sensory cortices[77]. Semantic cognition[78] might also be a target of compen-

sation. However, conflict tasks seem to rely minimally on memory, and involve rela-

tively simple sensory stimuli (e.g., colors and locations) and simple semantic pro-

cessing (e.g., reading a word). As such, conflict tasks may not necessitate compensation

in these functions.

In addition, compensation in older adults may manifest as increased recruitment

of bilateral regions and homologues with age[79]. For example, the hemispheric asym-

metry reduction in older adults (HAROLD) theory[80] suggests that older adults typically

exhibit less lateralization, either as a compensatory response to functional deficits or as

a reflection of neural dedifferentiation. However, we observed that the elderly showed

greater left lateralization compared to young to middle-aged adults. This finding is in-

consistent with the assumption of HAROLD but aligns with the right hemi-ageing

model[81], which posits that the right hemisphere is more vulnerable to age-related de-

cline. Prior research has shown that functional connectivity within the frontoparietal

control network is more disrupted in the right hemisphere than in the left during age-

ing[82]. This suggests that neural resources in the right hemisphere might be more limited

for the elderly, reducing its capacity to compensate for cognitive demands. Stronger

patterns of left laterality were also identified in childhood in the current study, primarily

noticeable within the prefrontal region (Fig. 7), which may reflect the earlier develop-

ment of the left hemisphere compared to the right[83]. In contrast, we found lower lateral-

ization in young adults. It is possible that previous studies showing stronger laterality

in young adults may have been biased by too small sample sizes and the use of non-

quantitative methods for calculation of laterality[84,85]. Moreover, because both left and

right lateralized results were reported in the literature on laterality[81], it is reasonable to

observe low laterality for young to middle-aged adults in the current meta-analysis.

Methodology Implications

By incorporating all the studies, our results demonstrate that the SDM can reliably iden-

tify brain regions as the ALE. However, the SDM has the added advantage of fully

utilizing existing effect size data and coordinates[86], allowing us to compare the relative

activity strength among various age groups, such as the contrast between young to mid-

dle-aged adults and elderly groups. More importantly, this approach allows for meta-

regressions to examine parametric relationships between brain region activity and age,

providing insights into the lifespan trajectories of cognitive control regions.

Limitation of Results

One caveat to consider in this study is the non-uniform distribution of age among the

included studies. Specifically, there is a noticeable gap in the age range of 45 to 60

years. Consequently, the observed age distribution could potentially influence the re-

sults of the regression analysis. We hope that future research could allocate more atten-

tion to the middle-aged period, considering the significant cognitive and neural changes

during this stage, such as the onset of cognitive decline[15,17,87]. In addition, it is crucial

to avoid the occurrence of ecological fallacy[88] (associations observed at the group level

are erroneously assumed to apply to individuals) when interpreting the results of meta-

regression analyses. Therefore, associations between brain activities and age across var-

ious studies do not provide direct insights into the specific age-related changes at the

individual level. Future research incorporating individual-level investigations (e.g., lon-

gitudinal follow-up studies) is crucial to obtaining a more comprehensive understand-

ing of these relationships.

Conclusions

Our meta-analysis adopted advanced meta-regression approaches to chart the lifespan

trajectories of cognitive control brain activities. We observed inverted U-shaped chang-

ing patterns in regions aligned with the frontoparietal control network, with the peaks

occurring between 24 and 40 years. In contrast, the dorsal attention network does not

present a clear age-related trajectory. This dissociation may reflect the hierarchy of

brain development in different regions. No other trajectory patterns were observed,

highlighting the predominance of the inverted U-shaped pattern in the lifespan trajec-

tory of cognitive control. Furthermore, we found the youth and elderly showed a more

asymmetric brain distribution than young to middle-aged adults. In sum, these results

demonstrate the multifaceted nature of age-related changes in cognitive control brain

function.

Methods

Literature Preparation

Literature Search

We report how we determined all data exclusions (if any), all manipulations, and all measures in the study. We first searched both English and Chinese articles on the youth and the elderly from PUBMED, Web of Science and CNKI (China National Knowledge Infrastructure) till 2022. The following search terms were applied in titles, abstracts, table of contents, indexing, and key concepts: *("Stroop" OR "Flanker" OR "Simon" OR "SNARC" OR "Navon" OR "interference" OR "cognitive conflict") AND ("fMRI" OR "functional resonance imaging" OR "functional imaging" OR "neuroimaging" OR "PET") AND ("children" OR "kids" OR "adolescents" OR "teenagers" OR "underage" OR "aged" OR "old" OR "older" OR "elder" OR "elderly" OR "senior" OR "development" OR "developmental" OR "aging" OR "life span").* The above process yielded 3,484 articles. In addition, 111 studies on young to middle-aged adults from a previous meta-analysis study[21] were included in the literature pool, 40 of which were excluded according to the current literature exclusion criteria (see below). Moreover, we screened 16 articles citing or being cited by the crucial literature. After removing duplicates, the literature search identified 2,930 articles.

Exclusion Criteria

We excluded any articles that met one or more of the following predefined exclusion criteria[89]: 1) not in English or Chinese; 2) not including healthy human participants; 3) case study; 4) not empirical study; 5) not functional resonance imaging (fMRI) or positron emission tomography (PET) study; 6) not whole-brain results (i.e., not have covered the whole gray matter); 7) not in Talairach or Montreal Neurological Institute (MNI) space; 8) not reflecting the congruency effect (i.e., contrasts between incongruent and congruent or between incongruent and neutral conditions); 9) not reporting exact mean age of participants.

A total of 119 articles were identified as eligible for inclusion in our meta-analyses. No statistical methods were used to pre-determine sample sizes, but our sample sizes are similar to or larger than those reported in previous publications[31,90]. Supplementary Fig. S1 shows the preferred reporting items for systematic reviews and meta-analyses (PRISMA)[91] flow chart for the literature screening process. The 119 articles included 129 studies (individual contrasts reported in the articles) with 3,388 participants and 1,579 activation foci reported. All studies were published or completed between 1994 and 2022. None of the experiments share the same group of participants. The included studies are written in English (124 studies) and Chinese (5 studies). Of the studies included, 125 were published in peer-reviewed journals, and 4 were master's theses. All included studies reported the task type used, including 74 studies utilizing Stroop-like

task (57%), 25 studies utilizing Simon task (19%), 25 studies utilizing Flanker task (19%), 2 studies utilizing a combination of Simon and Flanker tasks (2%), 1 study utilizing a combination of Simon and Stroop tasks (1%), and 3 studies utilizing multi-source interference task (2%). In addition, the contrasts conducted to reveal brain activations were also reported, with 98 studies (76%) resulting from the contrast of Incongruent trials > Congruent trials, 25 studies (19%) resulting from the contrast of Incongruent trials > Neutral trials, and 6 studies (5%) resulting from the union contrast of Incongruent trials > Congruent trials and Incongruent trials > Neutral trials. Regarding the handedness of participants in the included studies, 89 studies (69%) included right-handed participants only, 6 studies (5%) included both left and right-handed participants, while 34 studies (26%) did not report this information. Furthermore, 78 studies (60%) included only correct response trials, 2 studies (2%) included both correct and incorrect response trials, while 49 studies (38%) did not report this information. A detailed description of these features for each study is available in the Supplementary Table S1. To eliminate the influence of these confounding factors, we included them as covariates in the modeling analyses.

Coding Procedure

A coding manual was formulated to record pertinent study information, including authors, publication dates, experimental tasks, contrasts, and sample demographics (such

as the average age and sample size). To ensure coding accuracy, two authors inde-

pendently coded all studies, with discrepancies resolved through discussion or refer-

ence to the original studies. In instances where studies lacked essential information,

such as peak coordinates for relevant contrasts, participant age averages, or data for

specific age groups, efforts were made to contact the authors via e-mail to obtain the

relevant data. In addition, both coordinates and effect sizes (i.e., Hedge's *g*) were ex-

tracted from each study. Further, Talairach space coordinates were transformed to MNI

coordinates using the Lancaster transform[92].

Meta-Analytic Procedure

Activation Likelihood Estimation (ALE)

In order to obtain a comprehensive understanding of cognitive control-related brain ac-

tivity across all age groups and to replicate a prior study[21], we initially conducted a

single dataset meta-analysis using BrainMap GingerALE software[93] (version 3.0.2,

http://www.brainmap.org). This meta-analytical approach, known as activation likeli-

hood estimation, utilizes the spatial convergence of brain activity across multiple stud-

ies to determine the probability of activation in specific regions. Foci from individual

studies were transformed into a standardized coordinate space and modeled as Gaussian

probability values that accounted for variability in the number of participants in each

study. In situations where foci overlapped across studies, multiple Gaussians were as-

sociated with a single focus, and ALE selected the Gaussian with maximum probability

for each focus[93]. Subsequently, ALE score maps were generated by comparing these

modeled Gaussian distributions with a null distribution that simulated random brain

effects. The null distribution was generated using the same sample size and number of

foci groups as the experimental dataset for 1,000 times[94]. ALE scores were then used

to calculate p-values, which were based on the proportion of values higher than a certain

threshold in the null distribution. This resulted in a statistical ALE map that differenti-

ated true brain effects from random effects. A cluster-defining threshold of $p < 0.001$

and a minimum cluster size of 10 voxels (80 mm^3) were utilized to compute ALE maps,

consistent with the threshold applied in the seed-based d mapping (SDM) approach (see

below).

Seed-based d (Effect Size) Mapping

SDM is an alternative approach to statistically synthesize results from multiple neu-

roimaging experiments[86]. Similar to ALE, SDM employs a coordinate-based random-

effect approach to amalgamate peak coordinate information into a standard space across

several experiments. However, while ALE solely considers the binary feature (i.e., ac-

tive versus inactive) of peak coordinates, SDM takes into account the quantitative effect

size (can be positive or negative) connected to each peak and reconstructs the initial

parametric maps of individual experiments before amalgamating them into a meta-an-

alytic map[95]. Therefore, the use of a distinct algorithm in SDM from ALE allows us to

scrutinize the robustness and replicability of the outcomes obtained via ALE. More im-

portantly, SDM enables the inclusion of covariates in the meta-regression analyses to

reflect the changes in brain function across the lifespan.

We conducted three types of analyses using the SDM approach. Firstly, we esti-

mated the mean activation across all age groups and compared the results with ALE's

single dataset meta-analysis results. Secondly, we conducted contrasts between two

groups of studies (e.g., between young to middle-aged adults and a combination of the

youth and elderly groups) to identify brain regions that showed different levels of ac-

tivity across age. This analysis method served to investigate the hypothesized lifespan

trajectories, such as the inverted U-shaped pattern by elucidating neural activity varia-

tions linked to age. Thirdly, we defined specific models (e.g., linear and square root

models) to fit the whole brain to validate brain regions adhering to the hypothetical

lifespan changing patterns. This type of analysis aimed to explore various lifespan tra-

jectories, recognizing that different brain regions might follow distinct model functions.

See below for the details.

These analyses were conducted using the software of SDM with permutation of

subject images (SDM-PSI) (version 6.22, https://www.sdmproject.com). Effect size

maps were built for the 129 individual experiments. This was accomplished by (a) con-

verting the statistical value of each peak coordinate into an estimate of effect size

(Hedge's *g*) using standard formulas[96] and (b) convolving these peaks with a fully ani-

sotropic unnormalized Gaussian kernel ($\alpha = 1$, FWHM = 20 mm) within the boundaries

of a gray matter template (voxel size = $2 \times 2 \times 2$ mm^3). Imputation (50 times) was con-

ducted for each study separately to obtain a reliable estimate of brain activation maps[95].

In addition, the individual effect size maps were combined using a random-effect gen-

eral linear model. To assess the statistical significance of activations in the resulting

meta-analytic effect size map, 1,000 random permutations of activation peaks within

the gray matter template were compared. Finally, the meta-analytic maps were

thresholded using a voxel-wise family-wise error (FWE) corrected threshold of $p <$

0.001 and a cluster-wise extent threshold of 10 voxels[97].

Mean Analyses Across all Studies

This analysis aimed to characterize the activation distributions of cognitive control-

related brain regions across all studies, which was conducted utilizing the "Mean" func-

tion in SDM-PSI software. In order to verify the reliability of the SDM analysis results,

we compared the results with the single dataset meta-analysis using ALE. Results from

this analysis were further used as regions of interest (ROIs) in the subsequent model

fitting analyses (see section "*Generalized Additive Model (GAM) Fitting*" below). The

possibility of publication bias for resultant clusters was examined using Egger's test[98],

in which any result showing $p < 0.05$ was regarded as having significant publication

bias. Heterogeneity was evaluated using the I^2 index, which quantifies the proportion

of total variability attributable to heterogeneity between studies. A value less than 25%

indicates low heterogeneity among the included studies[99].

Contrast Analyses

To test our hypothesis that cognitive control related brain activities follow an inverted

U-shaped trajectory with age, we categorized each study based on the mean age of par-

ticipants into youth (< 18 years), young to middle-aged adults (18−59 years), and el-

derly (>= 60 years) groups. The age boundaries were determined to minimize age dis-

tribution overlap. We utilized SDM-PSI to perform a contrast analysis between the

group of young to middle-aged adults and the combination of other groups in order to

examine whether there are brain regions that exhibit an inverted U-shaped lifespan tra-

jectory. This was achieved by assigning studies from the young to middle-aged adult

group as 1 and all other studies as −1. This analysis yielded two results, one showing

higher activity in young to middle-aged adults than the youth and elderly groups, and

the other showing the opposite. Like the mean analysis, results from this analysis were

used as ROIs in the subsequent model fitting analyses (see sections "Generalized Ad-

ditive Model (GAM) Fitting" and "Model Simplification and Model Comparison").

In addition, to explore other possible trajectories, such as the increase-and-stable

pattern[5], we conducted contrast analyses between the youth and the combined group of

young to middle-aged and the elderly, as well as between the elderly and the combined

group of youth and young to middle-aged adults. Furthermore, to address the contro-

versies in previous studies, we conducted contrast analyses between older and young to

middle-aged adult groups, and between the youth and young to middle-aged adult

groups, respectively.

Meta-regression Analyses

To better describe the possible lifespan trajectories of the whole brain, we carried out

meta-regression analyses with the age and/or its derivatives as regressors. Two regres-

sions were conducted across the whole-brain, including a linear regression (with only

age as the regressor) and a square root regression (with age and its square root as sepa-

rate regressors). The linear regression aimed to test regions with increasing/decreasing

activity with age, and the square root regression aimed to test regions with the inverted

U-shaped trajectories based on the model fitting analyses (see below).

Data Extraction

Masks were generated for each ROI derived from the mean and contrast analyses in

SDM-PSI as described above. Subsequently, we extracted the effect sizes for each mask.

Fifty values were obtained from the SDM iterations and subsequently averaged for each

study in each region. The iterated variances were also averaged in a similar way. Addi-

tionally, we removed the outliers (beyond 3 standard deviations from the mean) in the

following model fitting analyses.

Generalized Additive Model (GAM) Fitting

To precisely estimate the inverted U-shaped trajectories, we adopted the GAM to fit the

curves. The GAM allows for flexible, nonparametric smoothing of predictor varia-

bles[100], and has been widely used to depict the lifespan trajectories[101,102]. We imple-

mented GAMs using the "mgcv" package[100] in R. For each ROI, we fitted a GAM with

the following formula:

g ~ s(age) + covariates,

where g is the effect size (dependent variable), s(age) represents a smoothing spline of

age (predictor variable), and covariates represent the dummy-coded categorical covari-

ate regressors. These regressors correspond to six aspects of the included studies: (1)

the presence of various conflict types (e.g., Stroop or Simon), (2) the mixed subject

samples based on handedness (e.g., right handed only or both handed), (3) the different

contrasts in reporting congruency effects (e.g., incongruent – congruent or incongruent

– neutral), (4) different trial types regarding whether they excluded error trials, (5) the

use of different types of experimental design (i.e., event-related or block designs), and

(6) the behavioral congruency effects measured by reaction time. Notably, we adopted

median imputation[103] for 9 studies (accounting for 6.98% of the total included studies)

not reporting the behavioral congruency effects, and included an indicator regressor to

account for the potential impact of imputation[104]. The validity of this imputation ap-

proach was confirmed through a robustness analysis (Supplementary Note S4). We in-

corporated these covariates to control for their potential confounding effects related to

age, which could otherwise influence our results. We also adjusted the estimate with a

weight parameter, which was the reciprocal of variance. We used penalized regression

splines, with the amount of smoothing determined automatically based on generalized

cross validation.

For each ROI, we quantified the peak age by choosing the highest prediction of a

fine-grained age scale (1,000 points from 8 to 74 years old). We also calculated the

estimated degree of freedom (EDF) for the smooth curve by summing up the degree of

freedom for each penalized term (i.e., s(age).1 to s(age).9).

Model Simplification and Model Comparison

While the GAM analysis may yield good fitting results on the data, it is important to

acknowledge its potential limitations. One concern is that it can fit the data with high

degree of freedoms (up to 7.0, Supplementary Table S4), which makes it susceptible to

over-fitting and harder to generalize. Another issue is its poor interpretability. Therefore,

we next sought to fit the data with simpler models.

To this end, we used the "metafor" package in R to fit these effect sizes with the

age and its derivatives as predictors. Specifically, we tested four non-linear models (see

below formulas and Fig. 3). Quadratic and cubic models were included based on previ-

ous studies[46,47]; quadratic logarithmic and square-root models were included to capture

the possible skewed trajectory, which would reflect the asymmetric trajectory of devel-

opment and decline of cognitive control[5]. Each model was fitted to each ROI separately.

In addition, we included the same covariates as the GAM analysis in each regression

model. We calculated the Akaike information criterion (AIC) to evaluate the goodness

of fit for each model.

1) Quadratic model

$g \sim age + age^2 + covariates$

2) Cubic model

$g \sim age + age^2 + age^3 + covariates$

3) Quadratic logarithmic model

$g \sim \log(age) + (\log(age))^2 + covariates$

4) Square root model

$g \sim age + \mathrm{sqrt}(age) + covariates$

Neurosynth Decoding Analysis

To investigate the potential functional difference between the two sets of brain regions

reported in section "*Dissociated Brain Networks with Distinct Lifespan Trajectories*"

of Results, we generated two binary maps from the contrast analysis between young to

middle-aged adults and the combination of other groups, and then submitted them to

the Neurosynth decoding system[41]. As the non-U-shaped map is defined as the converse

of the inverted U-shaped map, they were obtained by applying thresholds of $0.1 < p <$

0.9 and $p < 0.05$, respectively. These lenient threshold boundaries were used to mini-

mize the influence of sparsity on the decoding results. The two-boundary threshold was

used in generating the non-U-shaped map, as the original statistical map from the SDM-

PSI analysis was one-tailed, which means the threshold of $p > 0.9$ indicates the upright

U-shaped trend instead of a non-U-shaped trajectory. Additionally, to focus on the brain

regions specifically related to cognitive control, the two maps were masked by using

results from the mean SDM analysis (Fig. 1B). In the decoding results, we deleted terms

that were related to brain regions (e.g., "frontal"), not functional specific (e.g., "task"),

and duplicated (e.g., "attention" was removed if there was already "attentional").

Laterality Analysis

We calculated the laterality based on the effect size of reported brain coordinates from

each study. We computed the sum of effect sizes across coordinates for the left and right

hemisphere, respectively, yielding one global effect size each (i.e., g_L and g_R). Then,

we calculated the index of brain laterality with the following equation[105]:

$$\text{laterality} = \frac{g_L - g_R}{g_L + g_R} \ ,$$

which was then submitted to the GAM and simplified models (i.e., the quadratic, cubic,

square root, and quadratic logarithmic models).

To illustrate the contribution of different brain regions to the age-related change

of laterality, we calculated the laterality for each voxel[78]. We first used the SDM-PSI

to conduct a mean analysis for each age group (i.e., the youth, young to middle-aged

adult and elderly), yielding three z-maps. Then the laterality was computed with the

above equation for each voxel from the left hemisphere. The opposite values were cal-

culated for the right hemisphere. To avoid the bias due to asymmetric brain hemispheres,

we removed voxels without corresponding mirror coordinates. The results were visual-

ized confining to the brain regions estimated from the grand mean analyses (Fig. 1B).

Data Availability

The meta-data that support the findings of this study are available in Zenodo with the identifier doi: 10.5281/zenodo.12727621[106].

Code Availability

All codes conducting the ROI analyses are available in Zenodo with the identifier doi: 10.5281/zenodo.12727621[106].

Acknowledgement

We thank Fergus I.M. Craik, Beatriz Luna, and Haiyan Wu for valuable suggestions on this manuscript. We also thank Jing Yang and Yifan Zhang for the data check of extracted coordinates. Z.L. discloses support for the research of this work from Scientific Research Fund of Zhejiang Provincial Education Department (Y202249966), Starting Research Fund from Hangzhou Normal University (2021QDL079), and STI 2030-Major Projects (2021ZD0201705). I.T.P discloses support for the research of this work from the Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD) (HD098235). G.Y. discloses support for the research of this work from Jiefeng Jiang, and China Postdoctoral Science Foundation (2019M650884). X.L. discloses support for the research of this work from the Ministry of Science and Technology of the People's Republic of China (2021ZD0200505). The funders had no role in study design, data collection and analysis, decision to publish or preparation of the manuscript.

Author Contributions Statement

Conceptualization: Z.L. and G.Y.; Methodology: G.Y.; Formal analysis: G.Y. and Z.L. ; Data curation: Z.L.; Writing original draft: G.Y. and Z.L.; Writing, review, and editing: G.Y., Z.L., I.T.P., L.W., X.L. and J.R.; Funding: Z.L., I.T.P., X.L. and G.Y; Supervision: G.Y.

Competing Interests Statement

The authors declare no competing interests.

Inclusion & Ethics

All collaborators of this study fulfilled the criteria for authorship required by Nature Portfolio journals have been included as authors, as their participation was essential for implementation of the study and improvement of the manuscript. Roles and responsibilities were agreed among collaborators ahead of the research. This work does not include findings that are locally relevant. As a meta-analysis, this study does not involve direct interactions with human participants or the collection of new data. Each of the original studies included in this meta-analysis obtained ethical approval from their respective institutions, and we confirm that we complied with all applicable ethical guidelines and regulations.

794 **Tables**

795 **Table 1.** Brain areas activated in the contrast of one age group versus others (voxel-
796 wise FWE-corrected, $p < 0.001$, with minimum cluster size ≥ 10 voxels) with the SDM-
797 PSI.

Order	# Voxels	Z	p	L/R	MNI coordinate			Anatomical location	BA
					x	y	z		
Young to middle-aged adults > Others									
1	1165	4.713	< 0.001	R	52	16	4	inferior frontal gyrus	48
2	607	4.895	< 0.001	L	−46	16	26	inferior frontal gyrus	48
3	569	4.289	< 0.001	R	54	−48	42	inferior parietal lobule	40
4	215	3.405	< 0.001	L	−8	0	58	supplementary motor area	6
5	202	3.830	< 0.001	L	−36	−54	42	inferior parietal lobule	40
6	106	3.245	< 0.001	R	10	4	6	caudate nucleus	/
7	76	3.231	< 0.001	L	−38	24	0	insula	47
8	33	3.119	< 0.001	L	−36	12	−6	insula	48
9	10	2.794	< 0.001	R	4	−16	44	middle cingulate cortex	23
Young to middle-aged adults < Others									
None									
The youth > Others									
None									
The youth < Others									
None									
The elderly > Others									
None									
The elderly < Others									
None									

Note. The brain regions in the table correspond to the regions in Fig. 4. MNI = Montreal Neurological
Institute, BA = Brodmann area, L = left, R = right.

Figure Legends/Captions

Fig. 1. Overview of significant clusters across all studies regardless of age in the ALE

meta-analysis (A), the SDM meta-analysis (B), and their overlap (C). ALE = activa-

tion likelihood estimation; SDM = seed-based d mapping.

Fig. 2. Lifespan trajectories within regions identified in the mean analysis. l-ACC: left

anterior cingulate cortex, l-IPL: left inferior parietal lobule, r-IFG: right inferior

frontal gyrus, r-ITG: right inferior temporal gyrus, l-ITG: left inferior temporal gyrus,

r-MFG: right middle frontal gyrus, r-CN: right caudate nucleus, r-CB: right cerebel-

lum, l-ATP: left anterior thalamic projections. Scattered plots are the effect sizes as a

function of age, with curves fitted by GAM. The sizes of the scattered dots show the

square root of model weights (1/variance) for each study. Shaded areas around the

curves represent standard errors. Dashed lines indicate peak ages. Panels l-ACC and l-

IPL do not show the peak age due to an insignificant decrease at the later stage (Sup-

plementary Note S2).

Fig. 3. The lifespan trajectories explored in our study. Panels A-C show linear de-

crease, flat, and linear increase patterns, respectively, and were modelled with the lin-

ear function. Panels E and H show the upright and inverted U-shapes, respectively,

and were tested with the contrast between young to middle-aged adults and others, as

well as with the quadratic function. Panels D, F, G, and I show combinations of a sta-

ble period and an increase/decrease period across the lifespan, and were tested with

the contrast between the youth and others, or between the elderly and others. Panels J,

K and L show the variants of inverted U-shaped trajectories, which capture the possi-

bly early peak feature. They were tested with square root, quadratic logarithmic, and

cubic functions, respectively. See Methods for detailed models.

Fig. 4. Brain regions showing inverted U-shaped trajectory patterns. Scattered plots are the effect size as a function of age, with curves fitted by GAM (blue color) and the best simplified model (red color). Shaded areas around the curves represent standard errors. Dashed vertical lines show peak ages estimated from GAM (blue) and simplified model (red). The sizes of the scattered dots show the square root of model weights (1/variance) for each study. r-IFG: right inferior frontal gyrus, l-IFG: left inferior frontal gyrus, r-IPL: right inferior parietal lobule, l-SMA: left supplementary motor area, l-IPL: left inferior parietal lobule, r-CN: right caudate nucleus, l-Insula: left insula, r-MCC: right middle cingulate cortex.

Fig. 5. Significant clusters (voxel-wise FWE-corrected, $p < 0.001$, voxels ≥ 10) show-

ing square root pattern (A) and linear pattern (B) with age in the model fitting.

Fig. 6. Dissociated brain regions based on their trajectory patterns. A) The regions

following inverted U-shaped trajectories (red color) and non-U-shaped trajectories

(blue color). B) The axial view of the same results. The border lines display the fron-

toparietal control network (black), dorsal attention network (white), and their bound-

ary (gray) from Yeo's 7-network atlas[49]. Cingulo-opercular network was not plotted

due to its less clear dissociation among the two maps. The two scatter plots show two

example regions showing the non-U-shaped trajectory, one ([34, 4, 52]) representing a

peak region from the average brain activity analysis (Supplementary Table S3), and

the other ([22, −63, 42]) representing a region displaying a weak age-related change

from the contrast analysis with p between 0.49 and 0.51. The GAM analysis showed

that neither coordinate could be adequately fitted by a smoothed curve, with ps >

0.22.

Fig. 7. The laterality as a function of age. A) The trajectory fitted with a square root

model (red) and the GAM (blue). Higher values mean more left-lateralized and lower

values mean more right-lateralized. B) Visualization of the laterality for each group.

Regions in the left hemisphere show the left laterality, and vice versa.

References

1.   Cohen JD. Cognitive control: Core constructs and current considerations. In:
The Wiley handbook of cognitive control (ed Egner T) (2017).
2.   Miller EK, Cohen JD. An integrative theory of prefrontal cortex function. *Annu*
Rev Neurosci **24**, 167-202 (2001).
3.   Li Z, Goschl F, Yang G. Dissociated Neural Mechanisms of Target and
Distractor Processing Facilitated by Expectations. *J Neurosci* **40**, 1997-1999
(2020).
4.   Constantinidis C, Luna B. Neural Substrates of Inhibitory Control Maturation
in Adolescence. *Trends Neurosci* **42**, 604-616 (2019).
5.   Craik FI, Bialystok E. Cognition through the lifespan: mechanisms of change.
Trends Cogn Sci **10**, 131-138 (2006).
6.   Gupta R, Kar BR. Development of attentional processes in ADHD and normal
children. *Prog Brain Res* **176**, 259-276 (2009).
7.   Casey BJ, Getz S, Galvan A. The adolescent brain. *Dev Rev* **28**, 62-77 (2008).
8.   Diamond A. Executive functions. *Annu Rev Psychol* **64**, 135-168 (2013).
9.   Zanto TP, Gazzaley A. Attention and ageing. In: *The Oxford handbook of*
attention.). Oxford University Press (2014).
10.   Reuter-Lorenz PA, Cappell KA. Neurocognitive Aging and the Compensation
Hypothesis. *Current Directions in Psychological Science* **17**, 177-182 (2008).
11.   Association AP. Diagnostic and statistical manual of mental disorders : DSM-
5.). 5th edn. American Psychiatric Association (2013).
12.   Adnan A, Chen AJW, Novakovic-Agopian T, D'Esposito M, Turner GR. Brain
Changes Following Executive Control Training in Older Adults. *Neurorehabil*
Neural Repair **31**, 910-922 (2017).
13.   Casey BJ, Tottenham N, Liston C, Durston S. Imaging the developing brain:
what have we learned about cognitive development? *Trends Cogn Sci* **9**, 104-
110 (2005).
14.   Grady C. The cognitive neuroscience of ageing. *Nat Rev Neurosci* **13**, 491-505
(2012).
15.   Weintraub S, *et al.* Cognition assessment using the NIH Toolbox. *Neurology* **80**,
S54-64 (2013).
16.   Ferguson HJ, Brunsdon VEA, Bradford EEF. The developmental trajectories of
executive function from adolescence to old age. *Sci Rep* **11**, 1382 (2021).
17.   Li SC, Hammerer D, Muller V, Hommel B, Lindenberger U. Lifespan
development of stimulus-response conflict cost: similarities and differences
between maturation and senescence. *Psychol Res* **73**, 777-785 (2009).

18. Erb CD, Germine L, Hartshorne JK. Cognitive control across the lifespan:
Congruency effects reveal divergent developmental trajectories. *J Exp Psychol*
Gen **152**, 3285-3291 (2023).

19. Brouwer RM*, et al.* Genetic variants associated with longitudinal changes in
brain structure across the lifespan. *Nat Neurosci* **25**, 421-432 (2022).

20. Bethlehem RAI*, et al.* Brain charts for the human lifespan. *Nature*,   (2022).

21. Li Q, Yang G, Li Z, Qi Y, Cole MW, Liu X. Conflict detection and resolution
rely on a combination of common and distinct cognitive control networks.
Neurosci Biobehav Rev **83**, 123-131 (2017).

22. Giedd JN*, et al.* Brain development during childhood and adolescence: a
longitudinal MRI study. *Nat Neurosci* **2**, 861-863 (1999).

23. Crone EA, Dahl RE. Understanding adolescence as a period of social-affective
engagement and goal flexibility. *Nat Rev Neurosci* **13**, 636-650 (2012).

24. Andrews-Hanna JR, Mackiewicz Seghete KL, Claus ED, Burgess GC, Ruzic L,
Banich MT. Cognitive control in adolescence: neural underpinnings and
relation to self-report behaviors. *PLoS One* **6**, e21598 (2011).

25. Veroude K, Jolles J, Croiset G, Krabbendam L. Changes in neural mechanisms
of cognitive control during the transition from late adolescence to young
adulthood. *Dev Cogn Neurosci* **5**, 63-70 (2013).

26. Wilk HA, Morton JB. Developmental changes in patterns of brain activity
associated with moment-to-moment adjustments in control. *Neuroimage* **63**,
475-484 (2012).

27. Konrad K*, et al.* Development of attentional networks: an fMRI study with
children and adults. *Neuroimage* **28**, 429-439 (2005).

28. Marsh R*, et al.* A developmental fMRI study of self-regulatory control. *Hum*
Brain Mapp **27**, 848-863 (2006).

29. Bunge SA, Dudukovic NM, Thomason ME, Vaidya CJ, Gabrieli JD. Immature
frontal lobe contributions to cognitive control in children: evidence from fMRI.
Neuron **33**, 301-311 (2002).

30. Milham MP*, et al.* Attentional control in the aging brain: insights from an fMRI
study of the stroop task. *Brain Cogn* **49**, 277-296 (2002).

31. Yaple ZA, Stevens WD, Arsalidou M. Meta-analyses of the n-back working
memory task: fMRI evidence of age-related changes in prefrontal cortex
involvement across the adult lifespan. *Neuroimage* **196**, 16-31 (2019).

32. Prakash RS, Erickson KI, Colcombe SJ, Kim JS, Voss MW, Kramer AF. Age-
related differences in the involvement of the prefrontal cortex in attentional
control. *Brain Cogn* **71**, 328-335 (2009).

33. Fernandez NB, Hars M, Trombetti A, Vuilleumier P. Age-related changes in
attention control and their relationship with gait performance in older adults
with high risk of falls. *Neuroimage* **189**, 551-559 (2019).

34.   Sebastian A*, et al.* Differential effects of age on subcomponents of response
inhibition. *Neurobiology of aging* **34**, 2183-2193 (2013).

35.   Verhaeghen P, Cerella J. Aging, executive control, and attention: a review of
meta-analyses. *Neurosci Biobehav Rev* **26**, 849-857 (2002).

36.   Verhaeghen P. *The elements of cognitive aging: Meta-analyses of age-related*
differences in processing speed and their consequences. OUP USA (2014).

37.   Wood G, Ischebeck A, Koppelstaetter F, Gotwald T, Kaufmann L.
Developmental trajectories of magnitude processing and interference control:
an FMRI study. *Cereb Cortex* **19**, 2755-2765 (2009).

38.   Hart H, Radua J, Nakao T, Mataix-Cols D, Rubia K. Meta-analysis of functional
magnetic resonance imaging studies of inhibition and attention in attention-
deficit/hyperactivity disorder: exploring task-specific, stimulant medication,
and age effects. *JAMA psychiatry* **70**, 185-198 (2013).

39.   Zhang Z, Peng P, Eickhoff SB, Lin X, Zhang D, Wang Y. Neural substrates of
the executive function construct, age-related changes, and task materials in
adolescents and adults: ALE meta-analyses of 408 fMRI studies. *Dev Sci* **24**,
e13111 (2021).

40.   Long J, Song X, Wang Y, Wang C, Huang R, Zhang R. Distinct neural
activation patterns of age in subcomponents of inhibitory control: A fMRI meta-
analysis. *Front Aging Neurosci* **14**, 938789 (2022).

41.   Yarkoni T, Poldrack RA, Nichols TE, Van Essen DC, Wager TD. Large-scale
automated synthesis of human functional neuroimaging data. *Nat Methods* **8**,
665-670 (2011).

42.   Nee DE, Wager TD, Jonides J. Interference resolution: insights from a meta-
analysis of neuroimaging tasks. *Cogn Affect Behav Neurosci* **7**, 1-17 (2007).

43.   Botvinick MM, Cohen JD, Carter CS. Conflict monitoring and anterior
cingulate cortex: an update. *Trends Cogn Sci* **8**, 539-546 (2004).

44.   Kerns JG, Cohen JD, MacDonald AW, 3rd, Cho RY, Stenger VA, Carter CS.
Anterior cingulate conflict monitoring and adjustments in control. *Science* **303**,
1023-1026 (2004).

45.   Simonsohn U. Two Lines: A Valid Alternative to the Invalid Testing of U-
Shaped Relationships With Quadratic Regressions. *Advances in Methods and*
Practices in Psychological Science **1**, 538-555 (2018).

46.   Zuo XN*, et al.* Growing together and growing apart: regional and sex
differences in the lifespan developmental trajectories of functional homotopy. *J*
Neurosci **30**, 15034-15043 (2010).

47.   Coupe P, Catheline G, Lanuza E, Manjon JV, Alzheimer's Disease
Neuroimaging I. Towards a unified analysis of brain maturation and aging
across the entire lifespan: A MRI analysis. *Hum Brain Mapp* **38**, 5501-5518
(2017).

48. Amlien IK, Sneve MH, Vidal-Pineiro D, Walhovd KB, Fjell AM. The Lifespan
Trajectory of the Encoding-Retrieval Flip: A Multimodal Examination of
Medial Parietal Cortex Contributions to Episodic Memory. *J Neurosci* **38**, 8666-
8679 (2018).

49. Yeo BT, *et al.* The organization of the human cerebral cortex estimated by
intrinsic functional connectivity. *J Neurophysiol* **106**, 1125-1165 (2011).

50. De Luca CR, Leventer RJ. Developmental trajectories of executive functions
across the lifespan. In: *Executive functions and the frontal lobes*). Psychology
Press (2010).

51. Vaidya CJ, Bunge SA, Dudukovic NM, Zalecki CA, Elliott GR, Gabrieli JD.
Altered neural substrates of cognitive control in childhood ADHD: evidence
from functional magnetic resonance imaging. *The American journal of*
psychiatry **162**, 1605-1613 (2005).

52. Robertson BD, Hiebert NM, Seergobin KN, Owen AM, MacDonald PA. Dorsal
striatum mediates cognitive control, not cognitive effort per se, in decision-
making: An event-related fMRI study. *Neuroimage* **114**, 170-184 (2015).

53. Hsu CL, *et al.* Aerobic exercise promotes executive functions and impacts
functional neural activity among older adults with vascular cognitive
impairment. *Br J Sports Med* **52**, 184-191 (2018).

54. , (!!! INVALID CITATION !!!).

55. Brenhouse HC, Andersen SL. Developmental trajectories during adolescence in
males and females: a cross-species understanding of underlying brain changes.
Neurosci Biobehav Rev **35**, 1687-1703 (2011).

56. Gogtay N, *et al.* Dynamic mapping of human cortical development during
childhood through early adulthood. *Proc Natl Acad Sci U S A* **101**, 8174-8179
(2004).

57. Luna B, Marek S, Larsen B, Tervo-Clemmens B, Chahal R. An integrative
model of the maturation of cognitive control. *Annu Rev Neurosci* **38**, 151-170
(2015).

58. Shaw P, *et al.* Neurodevelopmental trajectories of the human cerebral cortex. *J*
Neurosci **28**, 3586-3594 (2008).

59. Grimm KJ, Ram N, Estabrook R. *Growth Modeling: Structural Equation and*
Multilevel Modeling Approaches. Guilford Publications (2016).

60. Verissimo J, Verhaeghen P, Goldman N, Weinstein M, Ullman MT. Evidence
that ageing yields improvements as well as declines across attention and
executive functions. *Nat Hum Behav* **6**, 97-110 (2022).

61. D'Mello AM, Gabrieli JDE, Nee DE. Evidence for Hierarchical Cognitive
Control in the Human Cerebellum. *Curr Biol* **30**, 1881-1892 e1883 (2020).

62. Shine JM, Lewis LD, Garrett DD, Hwang K. The impact of the human thalamus
on brain-wide information processing. *Nat Rev Neurosci* **24**, 416-430 (2023).

63.   Cole MW, Reynolds JR, Power JD, Repovs G, Anticevic A, Braver TS. Multi-
1022   task connectivity reveals flexible hubs for adaptive task control. *Nat Neurosci*
1023   **16**, 1348-1355 (2013).

1024   64.   Zhou Y, Friston KJ, Zeidman P, Chen J, Li S, Razi A. The Hierarchical
1025   Organization of the Default, Dorsal Attention and Salience Networks in
1026   Adolescents and Young Adults. *Cereb Cortex* **28**, 726-737 (2018).

1027   65.   Petersen SE, Posner MI. The attention system of the human brain: 20 years after.
1028   *Annu Rev Neurosci* **35**, 73-89 (2012).

1029   66.   Badre D, Nee DE. Frontal Cortex and the Hierarchical Control of Behavior.
1030   *Trends Cogn Sci* **22**, 170-188 (2018).

1031   67.   Dixon ML*, et al.* Heterogeneity within the frontoparietal control network and
1032   its relationship to the default and dorsal attention networks. *Proc Natl Acad Sci
1033   U S A* **115**, E1598-E1607 (2018).

1034   68.   Sydnor VJ*, et al.* Intrinsic activity development unfolds along a sensorimotor-
1035   association cortical axis in youth. *Nat Neurosci* **26**, 638-649 (2023).

1036   69.   Raz N*, et al.* Selective aging of the human cerebral cortex observed in vivo:
1037   differential vulnerability of the prefrontal gray matter. *Cereb Cortex* **7**, 268-282
1038   (1997).

1039   70.   Fjell AM, Walhovd KB. Structural brain changes in aging: courses, causes and
1040   cognitive consequences. *Rev Neurosci* **21**, 187-221 (2010).

1041   71.   Kaplan L, Chow BW, Gu C. Neuronal regulation of the blood-brain barrier and
1042   neurovascular coupling. *Nat Rev Neurosci* **21**, 416-432 (2020).

1043   72.   Lipecz A*, et al.* Age-related impairment of neurovascular coupling responses: a
1044   dynamic vessel analysis (DVA)-based approach to measure decreased flicker
1045   light stimulus-induced retinal arteriolar dilation in healthy older adults.
1046   *Geroscience* **41**, 341-349 (2019).

1047   73.   Zlokovic BV. Neurovascular pathways to neurodegeneration in Alzheimer's
1048   disease and other disorders. *Nat Rev Neurosci* **12**, 723-738 (2011).

1049   74.   Montala-Flaquer M, Canete-Masse C, Vaque-Alcazar L, Bartres-Faz D, Pero-
1050   Cebollero M, Guardia-Olmos J. Spontaneous brain activity in healthy aging: An
1051   overview through fluctuations and regional homogeneity. *Front Aging Neurosci*
1052   **14**, 1002811 (2022).

1053   75.   Cappell KA, Gmeindl L, Reuter-Lorenz PA. Age differences in prefontal
1054   recruitment during verbal working memory maintenance depend on memory
1055   load. *Cortex* **46**, 462-473 (2010).

1056   76.   Zanto TP, Gazzaley A. Cognitive Control and the Ageing Brain. In: *The Wiley
1057   Handbook of Cognitive Control* (ed Egner T) (2017).

1058   77.   Park DC, Reuter-Lorenz P. The adaptive brain: aging and neurocognitive
1059   scaffolding. *Annu Rev Psychol* **60**, 173-196 (2009).

1060   78.   Hoffman P, Morcom AM. Age-related changes in the neural networks
1061   supporting semantic cognition: A meta-analysis of 47 functional neuroimaging
1062   studies. *Neurosci Biobehav Rev* **84**, 134-150 (2018).

1063   79.   Langenecker SA, Nielson KA, Rao SM. fMRI of healthy older adults during
1064   Stroop interference. *NeuroImage* **21**, 192-200 (2004).

1065   80.   Cabeza R. Hemispheric asymmetry reduction in older adults: the HAROLD
1066   model. *Psychol Aging* **17**, 85-100 (2002).

1067   81.   Dolcos F, Rice HJ, Cabeza R. Hemispheric asymmetry and aging: right
1068   hemisphere decline or asymmetry reduction. *Neurosci Biobehav Rev* **26**, 819-
1069   825 (2002).

1070   82.   Zhang HY, Chen WX, Jiao Y, Xu Y, Zhang XR, Wu JT. Selective vulnerability
1071   related to aging in large-scale resting brain networks. *PLoS One* **9**, e108807
1072   (2014).

1073   83.   Thatcher RW, Walker RA, Giudice S. Human cerebral hemispheres develop at
1074   different rates and ages. *Science* **236**, 1110-1113 (1987).

1075   84.   Cabeza R*, et al.* Age-related differences in neural activity during memory
1076   encoding and retrieval: a positron emission tomography study. *J Neurosci* **17**,
1077   391-400 (1997).

1078   85.   Grady CL, Bernstein LJ, Beig S, Siegenthaler AL. The effects of encoding task
1079   on age-related differences in the functional neuroanatomy of face memory.
1080   *Psychol Aging* **17**, 7-23 (2002).

1081   86.   Albajes-Eizagirre A, Solanes A, Vieta E, Radua J. Voxel-based meta-analysis
1082   via permutation of subject images (PSI): Theory and implementation for SDM.
1083   *Neuroimage* **186**, 174-184 (2019).

1084   87.   Sowell ER, Thompson PM, Toga AW. Mapping changes in the human cortex
1085   throughout the span of life. *Neuroscientist* **10**, 372-392 (2004).

1086   88.   Thompson SG, Higgins JP. How should meta-regression analyses be
1087   undertaken and interpreted? *Stat Med* **21**, 1559-1573 (2002).

1088   89.   Cumpston M*, et al.* Updated guidance for trusted systematic reviews: a new
1089   edition of the Cochrane Handbook for Systematic Reviews of Interventions.
1090   *Cochrane Database Syst Rev* **10**, ED000142 (2019).

1091   90.   Lu J, Yao J, Zhou Z, Wang XT. Age effects on delay discounting across the
1092   lifespan: A meta-analytical approach to theory comparison and model
1093   development. *Psychological Bulletin* **149**, 447-486 (2023).

1094   91.   Page MJ*, et al.* The PRISMA 2020 statement: an updated guideline for reporting
1095   systematic reviews. **88**, 105906 (2021).

1096   92.   Lancaster JL*, et al.* Bias between MNI and talairach coordinates analyzed using
1097   the ICBM-152 brain template. *Human Brain Mapping* **28**, 1194-1205 (2007).

1098   93.   Turkeltaub PE, Eickhoff SB, Laird AR, Fox M, Wiener M, Fox P. Minimizing
1099   within-experiment and within-group effects in activation likelihood estimation
1100   meta-analyses. *Human Brain Mapping* **33**, 1-13 (2012).

1101   94.   Eickhoff SB, Bzdok D, Laird AR, Kurth F, Fox PT. Activation likelihood
1102   estimation meta-analysis revisited. *NeuroImage* **59**, 2349-2361 (2012).
1103   95.   Radua J, Mataix-Cols D. Meta-analytic methods for neuroimaging data
1104   explained. *Biol Mood Anxiety Disord* **2**, 6 (2012).
1105   96.   Hedges LV. Distribution theory for Glass's estimator of effect size and related
1106   estimators. *Journal of Educational Statistics* **6**, 107-128 (1981).
1107   97.   Radua J*, et al.* A new meta-analytic method for neuroimaging studies that
1108   combines reported peak coordinates and statistical parametric maps. *Eur*
1109   *Psychiat* **27**, 605-611 (2012).
1110   98.   Egger M, Smith GD, Phillips AN. Meta-analysis: principles and procedures.
1111   *BMJ (Clinical research ed)* **315**, 1533-1537 (1997).
1112   99.   Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in
1113   meta-analyses. *BMJ (Clinical research ed)* **327**, 557-560 (2003).
1114   100.   Wood SN. mgcv: GAMs and generalized ridge regression for R. *J R news* **1**, 20-
1115   25 (2001).
1116   101.   Zuo XN, He Y, Betzel RF, Colcombe S, Sporns O, Milham MP. Human
1117   Connectomics across the Life Span. *Trends Cogn Sci* **21**, 32-45 (2017).
1118   102.   Bethlehem RAI*, et al.* Brain charts for the human lifespan. *Nature* **604**, 525-533
1119   (2022).
1120   103.   Ochieng' Odhiambo F. Comparative Study of Various Methods of Handling
1121   Missing Data. *Mathematical Modelling and Applications* **5**,   (2020).
1122   104.   Sandhu AT, Kohsaka S, Turakhia MP, Lewis EF, Heidenreich PA. Evaluation
1123   of Quality of Care for US Veterans With Recent-Onset Heart Failure With
1124   Reduced Ejection Fraction. *JAMA Cardiol* **7**, 130-139 (2022).
1125   105.   Seghier ML. Laterality index in functional MRI: methodological issues. *Magn*
1126   *Reson Imaging* **26**, 594-601 (2008).
1127   106.   Yang   G.   guochun-yang/Cognitive_Control_Developmental_Trajectory:
1128   Release_20240711 (Lifespan).). Zenodo (2024).
1129