Expression-based machine learning models for predicting plant tissue identity

Sourabh Palande[1], Jeremy Arsenault[2], Patricia Basurto-Lozada[3], Andrew Bleich[4], Brianna N. I. Brown[4], Sophia F. Buysse[4,5,6], Noelle A. Connors[7], Sikta Das Adhikari[1,8], Kara C. Dobson[5,9], Francisco Xavier Guerra-Castillo[10,11], Maria F. Guerrero-Carrillo[12], Sophia Harlow[7], Héctor Herrera-Orozco[13,14], Asia T. Hightower[4,5], Paulo Izquierdo[15], MacKenzie Jacobs[16,17], Nicholas A. Johnson[5,18], Wendy Leuenberger[5,9], Alessandro Lopez-Hernandez[3,19], Alicia Luckie-Duque[12], Camila Martínez-Avila[20], Eddy J. Mendoza-Galindo[12], David Plancarte[21], Jenny M. Schuster[22,17], Harry Shomer[2], Sidney C. Sitar[15,23,24], Anne K. Steensma[4,17,25], Joanne Elise Thomson[17,22], Damián Villaseñor-Amador[26], Robin Waterman[4,5,6], Brandon M. Webster[4], Madison Whyte[15], Sofía Zorilla-Azcué[27], Beronda L. Montgomery[28], Aman Y. Husbands[29], Arjun Krishnan[30], Sarah Percival[1], Elizabeth Munch[1,31], Robert VanBuren[7,32], Daniel H. Chitwood[1,7,*], Alejandra Rougon-Cardoso[12,33*]

[1]Department of Computational Mathematics, Science and Engineering, Michigan State University, USA
[2]Department of Computer Science and Engineering, Michigan State University, USA
[3]Laboratorio Internacional de Investigación sobre el Genoma Humano (LIIGH), Universidad Nacional Autónoma de México, México
[4]Department of Plant Biology, Michigan State University, USA
[5]Ecology, Evolution, and Behavior Program, Michigan State University, USA
[6]Kellogg Biological Station, Michigan State University, USA
[7]Department of Horticulture, Michigan State University, USA
[8]Department of Statistics and Probability, Michigan State University, USA
[9]Department of Integrative Biology, Michigan State University, USA
[10]Unidad de Investigación Médica en Inmunología e Infectología, Instituto Mexicano del Seguro Social, México
[11]Programa de Posgrado en Ciencias Biológicas, Facultad de Medicina, Universidad Nacional Autónoma de México, México
[12]Laboratory of Agrigenomic Sciences, Escuela Nacional de Estudios Superiores Unidad León, Universidad Nacional Autónoma de México, México
[13]Posgrado en Ciencias Biológicas, Universidad Nacional Autónoma de México, México
[14]Laboratorio de Ecología Evolutiva y Conservación de Anfibios y Reptiles. Facultad de Estudios Superiores Iztacala, Universidad Nacional Autónoma de México, México
[15]Department of Plant, Soil, and Microbial Sciences, Michigan State University, USA
[16]Department of Biochemistry and Molecular Biology, Michigan State University, USA
[17]Molecular Plant Sciences Program, Michigan State University, USA
[18]Genetics and Genome Sciences, Michigan State University, USA
[19]Computational Population Genetics Group, Universidad Nacional Autónoma de México, México
[20]Colección Nacional de Aves, Posgrado en Ciencias Biológicas, Instituto de Biología, Universidad Nacional Autónoma de México, México
[21]Departamento de Botánica, Posgrado en Ciencias Biológicas, Instituto de Biología, Universidad Nacional Autónoma de México, México
[22]Cell and Molecular Biology, Michigan State University, USA
[23]Plant Breeding, Genetics, and Biotechnology, Michigan State University, USA
[24]Crop and Soil Sciences Program, Michigan State University, USA
[25]MSU-DOE Plant Research Laboratory, Michigan State University, USA
[26]Programa de Posgrado en Ciencias Biológicas, Facultad de Ciencias, Universidad Nacional Autónoma de México, México
[27]Programa de Posgrado en Ciencias Biológicas, Escuela Nacional de Estudios Superiores (ENES), Unidad Morelia, Universidad Nacional Autónoma de México, México
[28]Department of Biology, Grinnell College, USA
[29]Department of Biology, University of Pennsylvania, USA
[30]Department of Biomedical Informatics, Center for Health AI, University of Colorado Anschutz Medical Campus, USA
[31]Department of Mathematics, Michigan State University, USA
[32]Plant Resilience Institute, Michigan State University, USA
[33]Plantecc National Laboratory, ENES-León, México

*Corresponding authors
Dr. Alejandra Rougon-Cardoso, arougon@enes.unam.mx
Dr. Daniel H. Chitwood, dhchitwood@gmail.com

ABSTRACT

The selection of *Arabidopsis* as a model organism played a pivotal role in advancing genomic science, firmly establishing the cornerstone of today's plant molecular biology. Competing frameworks to select an agricultural- or ecological-based model species, or to decentralize plant science and study a multitude of diverse species, were selected against in favor of building core knowledge in a species that would facilitate genome-enabled research that could assumedly be transferred to other plants. Here, we examine the ability of models based on *Arabidopsis* gene expression data to predict tissue identity in other flowering plant species. Comparing different machine learning algorithms, models trained and tested on *Arabidopsis* data achieved near perfect precision and recall values using the K-Nearest Neighbor method, whereas when tissue identity is predicted across the flowering plants using models trained on *Arabidopsis* data, precision values range from 0.69 to 0.74 and recall from 0.54 to 0.64, depending on the algorithm used. Below-ground tissue is more predictable than other tissue types, and the ability to predict tissue identity is not correlated with phylogenetic distance from *Arabidopsis*. This suggests that gene expression signatures rather than marker genes are more valuable to create models for tissue and cell type prediction in plants. Our data-driven results highlight that, in hindsight, the assertion that knowledge from *Arabidopsis* is translatable to other plants is not always true. Considering the current landscape of abundant sequencing data and computational resources, it may be prudent to reevaluate the scientific emphasis on *Arabidopsis* and to prioritize the exploration of plant diversity.

INTRODUCTION

Historically, plant biology has focused on inferring genetic, molecular, physiological, and ecological mechanisms. Conventionally, through quantifying phenomena and applying statistics, hypotheses are tested and decisions of most likely scenarios are determined. New technologies and computational approaches have caused a shift from hypothesis- to data-driven research (Mazzocchi, 2015). Moreover, plant biology has embraced the inclusion of machine learning methods in addition to traditional statistical approaches (Ij, 2018). Both a deluge of data and new computational methods have allowed for predictive, rather than inferential, methods. Both statistics and machine learning can be used for inference and prediction, but machine learning methods more often classify and predict on class labels rather than inferring statistical parameters of a population. In plant biology, such predictive approaches underlie the frameworks of phenotyping (Coppens et al., 2017), precision agriculture (Zhang et al., 2002), genomic prediction (Crossa et al., 2014), linking transcriptomic profiles to phenotype (Azodi et al., 2020), and protein structure determination (Jumper et al., 2021). Just as inferential statistics has its limitations, the robustness and ability to extrapolate predictive models are also constrained by the empirical context from which the data originates. Although data-driven research is slowly becoming more theoretical and predictive (Hogeweg, 2011), the creation of universal plant models is hindered by their overwhelming diversity. Not only is the phylogenetic diversity among flowering plants immense (The Angiosperm Phylogeny Group et al., 2016), but plants are exceptionally responsive to their environments (Sultan, 2000) and have evolved symbiotic interactions with and defense mechanisms against innumerable microbes (Mitchell et

101   al., 2006). Furthermore, technical variability in data acquisition makes it difficult to exploit the
102   huge amount of expression data archived in databases. The number of ways we sample
103   molecular profiles from plant tissues and the interaction effects that arise between
104   phylogenetically diverse species with environments, stresses, and biotic interactions is
105   countless and prevents extrapolating results between studies.
106
Due to the clear advantages of studying a single model species, the early days of the genomics
era tended to overlook the importance of prioritizing plant diversity. The candidates considered
for the first sequenced genome were either easily transformable (e.g., species within
Solanaceae; Knapp et al., 2004) or were already used for genetics (e.g., maize; Strable and
Scanlon, 2009), but never was biodiversity considered (Meyerowitz, 2001). Reasons for
choosing *Arabidopsis* as the first sequenced plant genome (Arabidopsis Genome Initiative,
2000) include ease of transformation (Clough and Bent, 1998), its small genome (Bennett et al.,
2003), and life history traits that allow for genetics through crossing, and short generation times
(Meyerowitz, 1987). The justification for initially sequencing the genome of a single model
species was that such focus would allow unprecedented molecular discoveries that could be
translated into other species and improve our understanding of all plants (Bevan and Walsh,
2005). The strategy to focus on a single model species was successful, and *Arabidopsis* is the
most cited plant in the last 20 years, even surpassing key crops and all other plant species
(Marks et al., 2023). Our molecular knowledge in plants was purposefully constructed to focus
on *Arabidopsis* over crops and plant genetic diversity. However, such a choice has little
relevance in a changing climate with dwindling natural resources and vanishing biodiversity that
have become the most pressing concerns of our time. The cultural dynamics that influenced the
choice of *Arabidopsis* as the first sequenced genome are reflected in subsequently sequenced
plant genomes. Plants intrinsic to Indigenous cultures and territories have been sequenced by
colonial powers (Marks et al., 2021; Dyer et al., 2022). While sequencing *Arabidopsis* has
certainly expanded our knowledge of molecular processes, due to such an intense focus, our
understanding in other species remains limited. This leaves us questioning the extent to which
the insights from *Arabidopsis* can be extrapolated to the rest of flowering plants.
130
In the 20 years since the release of the *Arabidopsis* genome sequence (Arabidopsis Genome
Initiative, 2000), the number of sequenced plant genomes has dramatically risen (Michael and
Jackson, 2013; Li and Harkess, 2018; Marks et al., 2021) leading to a greater understanding of
the evolutionary origin and genetic mechanisms underlying numerous traits across the green
lineage. Next-generation sequencing, for example, has enabled unprecedented surveys of
genome-scale features across species, tissue types, environments, and interactions between
plants with abiotic and biotic factors. There are currently over 300,000 public gene expression
datasets spanning thousands of diverse plant species (Lim et al., 2022). Cross-species
comparisons of gene expression across plants have usually been limited by the number of
species analyzed (Proost and Mutwil, 2018) or their sampling breadth. Most studies have
generated datasets from scratch (Julca et al., 2021) instead of leveraging public repositories.
Databases and datasets curating and making vast amounts of gene expression profiles and
their associated metadata have been created. For example, an *Arabidopis* RNA-seq database
(ARS) compiles 20,068 publicly available *Arabidopsis* RNA-Seq libraries (Zhang et al., 2020),

and the Plant Public RNA☐seq Database has ~45,000 maize, rice, wheat, soybean and cotton
samples (Yu et al. 2022). Previously we had curated a dataset of 2,671 publicly available gene
expression profiles from 54 flowering plant species across 7 developmental tissue types and
nine stresses (Palande et al., 2023). More than 20 years after the release of the *Arabidopsis*
genome, not only have we accumulated enough data across plants to ask unprecedented
questions but new computational tools are available that permit comparative approaches to
analyze such massive amounts of data.
152
Here, building upon large, curated databases of *Arabidopsis* (Zhang et al., 2020) and flowering
plant gene expression profiles (Palande et al., 2023), we examine how predictive *Arabidopsis* is
as a model species relative to the rest of the flowering plants and to what degree we can
extrapolate our knowledge from model organisms to diverse plant species. Dimension reduction
through principal component analysis (PCA) reveals that biotic stress response and tissue type
are primary, orthogonal sources of structure in gene expression data from *Arabidopsis*, and
while angiosperm data projected onto this space retains some structure, the regions occupied
between tissue types become less distinct. We next compare the performance of different
machine learning models. The k-nearest neighbor (KNN) method yields precision and recall
values of up to 0.99 with models trained and tested on *Arabidopsis* data. Model performance
drops significantly, with higher precision than recall values, when data from across flowering
plants is tested using models trained on *Arabidopsis* data. Below-ground tissue is more
separated from and predictable than other tissue types, and phylogenetic distance from
Arabidopsis does not appear to influence prediction rates. We end with a discussion of the
implications of our results for the current structure of the plant science community,
acknowledging that the past focus on *Arabidopsis* as a model organism based on decisions
decades ago was effective at that time; however, we now advocate for a shift in approach due
to changing circumstances, particularly in light of the pressing issue of biodiversity loss. We
argue for a more decentralized and inclusive research framework that better encompasses the
diversity of plants and the human cultures that represent them, adapting to current
environmental and scientific challenges.
174
175   **MATERIALS AND METHODS**
176
177   *Datasets*
178
We used two curated databases in this analysis. The first contained 28,165 *Arabidopsis* gene
expression profiles across 37,334 genes (Zhang et al. 2020). The second contained 2,671
flowering plant expression profiles across 6,327 orthogroups (Palande et al. 2023). Metadata
labels for each sample from both of the databases was assigned one of four tissue type labels
(above-ground, below-ground, whole plant, or other). The categories are purposefully
encompassing and chosen to facilitate accurate assignment across the broad categories of
experimental data we analyzed, focusing on above-ground and below-ground tissue identity as
one of the simplest cases to test tissue predictability. After removing samples with missing
metadata and samples with low unique mapped rate (<75%), the *Arabidopsis* database was left
with 19,415 samples. A conserved *Arabidopsis* database was also constructed by keeping only

the genes mapped to the orthogroups from the flowering plant database. The conserved
Arabidopsis database contained the same number of samples, but with much smaller
expression profiles across only the 6,327 orthogroups shared with the angiosperm dataset.
192
193 *Classification models*
194
Classification is a common machine learning task where, given data points belonging to two or
more classes, the goal is to *learn* a function that best differentiates between points from different
classes. Then, given a new data point, the function can be used to decide which class the point
belongs to. The classifier function can be learned in many different ways, leading to various
types of machine learning models. For each classifier model in this study, we employed the
following modeling methods:
201
Linear support vector classifier (SVC): In linear classification, each point is viewed as a vector in
k-dimensional space (Cortes and Vapnik, 1995). The goal is to find *(k-1)*-dimensional
hyperplanes that separate the points belonging to different classes. There are many possible
choices for hyperplanes that can classify the points. A reasonable choice is to find the ones that
maximize the separation between points from different classes. These are known as maximum-
margin hyperplanes. Geometrically, the max-margin hyperplanes are defined by the points that
lie closest to them; therefore, such points are called support vectors.
209
Multi-layer perceptron (MLP): The SVC model assumes that the classes are linearly separable,
which may not be true. MLPs are a class of artificial neural networks (Haykin, 1998) with three
or more layers of "perceptrons" with non-linear activation. An MLP consists of an input and an
output layer, with one or more hidden layers of neurons. We experimented with one and two
hidden-layer MLPs and used rectified linear unit (ReLU) activation in all cases. In ReLU, a
neuron's activation is the weighted sum of its inputs, if the sum is non-negative, and zero
otherwise. Even with this simple nonlinear activation function, MLPs are able to outperform the
linear SVC.
218
Random forest (RF): Random forests (Ho, 1995) perform classification by constructing an
ensemble of decision trees. Each decision tree outputs a class label for the given sample and
the output of the RF is the class label predicted by the majority of the trees. In a decision tree,
each internal node is labeled by an input feature and the leaf nodes are labeled by the class
labels. Starting from the root node, the input set is recursively partitioned into children nodes
using the input feature associated with the node. The recursion ends when all data points in the
node belong to the same class, or some pre-specified termination criteria, such as maximum
depth of the tree, are met. Which feature to split the data on at each level is determined using
information criteria such as gini impurity or entropy that measure how consistent the subsets are
with respect to the class labels after the split.
229
230 *Histogram-based gradient boosting (HGB)*: Gradient boosting (Mason et al., 1999) is another
231 class of methods that uses a large ensemble of decision trees. In histogram-based boosting, the
232 real-valued input features are first discretized into a few (typically 256) bins using histograms.

233 This allows the training algorithm to run much more efficiently and construct a much larger
234 ensemble of decision trees to support the classification.
235
236 *K-nearest neighbor (KNN) classifier*: In KNN classifiers (Cover and Hart, 1967) class labels are
237 assigned based on a majority vote of the K nearest training points. The distance metric and the
238 number of neighbors are specified by the user. In our experiments, correlation distance between
239 the expression profiles was used to train the KNN classifier.
240
241 *Experimental design*
242
To establish the utility of gene expression profiles in predicting tissue type, we trained the
supervised machine learning models to classify the *Arabidopsis* data by tissue types (**Table 1**).
The database was split into training and test sets (70%-30% split). To ensure comparability, all
five models were trained and tested on the same training and test sets. Next, we wanted to
examine how predictive *Arabidopsis* is to the rest of the flowering plants (**Table 2**). To test this,
we used a set of conserved *Arabidopsis* transcripts with orthogroups across angiosperms, split
into training and test sets (70%-30% split) as before. The same five machine learning models
were trained on the conserved *Arabidopsis* training set. The performance of these models was
first tested on the conserved gene *Arabidopsis* test set to make sure that the models were still
able to predict the tissue types with a significantly smaller number of features. We then used the
same models to classify the angiosperm data to test how well they extrapolate to species other
than *Arabidopsis*. Each machine learning model employed in our experiments requires
additional hyperparameters that need to be tuned to optimize model performance. We used the
Bayesian optimization procedure implemented in the hyperopt package in Python (Bergstra et
al., 2013). To gain insights into the functional annotation and enrichment of our gene list, we
performed a Gene Ontology term analysis using the DAVID Functional Annotation Clustering
tool (version 2021) from the web interface http://david.ncifcrf.gov (Huang et al., 2009). We
filtered the 200 genes with the most positive and negative PC1 loading values. The annotation
was performed using TAIR IDs and selecting Gene Ontology terms from levels 3 and 4 of
Molecular Function and Biological Process categories. All data and code to reproduce the
results in this manuscript are available at [https://github.com/PlantsAndPython/arabidopsis-gene-](https://github.com/PlantsAndPython/arabidopsis-gene-expression)
[expression](https://github.com/PlantsAndPython/arabidopsis-gene-expression).
265
266 **RESULTS**
267
268 *Dimension reduction and alignment between* Arabidopsis *and angiosperm gene expression*
269 *datasets*
270
A principal component analysis (PCA) performed on the full dataset of 19,415 *Arabidopsis*
RNAseq samples shows a clear separation by tissue type (**Fig. 1a**). For simplicity, we
categorized samples into bins of above-ground, below-ground, whole plant, and other. The
above-ground, below-ground, and other tissue types are well-separated from each other, but the
below-ground tissue has the least overlap with other tissues. The whole plant tissue type,
composed of different combinations of the other tissues, is not well separated, as we would

expect. The separation of tissues occurs along a gradient defined by PC2, demonstrating that
tissue type is not the primary source of variance in the data. Rather, a small proportion of
samples are strewn across PC1 in an additive, orthogonal manner, preserving the separation of
tissue types defined by PC2. To investigate the underlying cause responsible for the primary
source of variation in the data, we performed GO enrichment on genes with the most extreme
PC1 loading values that are most responsible for defining PC1. In the full *Arabidopsis* dataset
(**Fig. 1a**), high PC1 values, which include a small number of samples that contribute to a
disproportionate amount of variance in the data, are defined by high expression of genes
associated with response to biotic stress and oxidative damage GO terms (**Table S1**). Low PC1

Figure 1: Principal Component Analysis (PCA) of gene expression profiles. PCAs with gene expression profiles colored by above-ground (blue), below-ground (orange), whole plant (red), and other (green) tissue types for **a)** the full *Arabidopsis* dataset, **b)** the conserved *Arabidopsis* data set, **c)** the angiosperm dataset projected onto the conserved *Arabidopsis* PCA from b), and **d)** the same as c), but with conserved *Arabidopsis* gene expression profiles in the background (transparent gray).

286 values, which include a majority of samples across tissues and which we assume arise from
287 plants grown under regular conditions associated with the less stress, are defined by high
288 expression of genes with GO terms associated with biosynthesis, biogenesis, and cell growth.
289 Remarkably, in the full *Arabidopsis* dataset, negative PC1 loading values are enriched for
290 glucosinolate biosynthetic and other metabolic processes (FDR <0.05) .
291
From these large-scale datasets, we developed a predictive model to test if tissue type could be
inferred from expression patterns alone and if this *Arabidopsis*-trained model could be
transferred to other flowering plants. We previously created a set of 6,328 low copy orthogroups
that are deeply conserved across flowering plants (Palande et al., 2023) and used a set of 6,327
Arabidopsis genes corresponding to these orthogroups for all downstream analyses. A PCA
performed on this subset of 6,327 conserved flowering plant genes shows mostly the same
structure as the analysis with all *Arabidopsis* genes included (**Fig. 1b**). However, while the
below-ground tissue type remains distinct from the rest of the data, the above-ground tissue
type overlaps more with whole plant and other tissue types. Note that the sign of principal
components is arbitrary, which explains the "flip" of PC2 values relative to the full set of
Arabidopsis genes. An analysis of the enriched GO terms for PC1 loading values from the
conserved gene PCA reveals that high PC1 values are associated with biotic responses, but
also with anther- and pollen-related GO terms (**Table S1**). Low PC1 values are associated
overwhelmingly with photosynthesis. Because the two datasets have corresponding orthogroup
features, we are able to project the angiosperm dataset onto the PCA defined by the conserved
gene *Arabidopsis* dataset (**Fig. 1c-d**). While the overall structure defining the distributions of
tissue types is maintained in the projected angiosperm data, there is substantial overlap
between above-ground and below-ground tissue types. We conclude that indeed there is
conservation of tissue-specific expression between *Arabidopsis* and the rest of the flowering
plants, but that as expected, the alignment of the underlying structures of gene expression
patterns defining tissue type identity are not identical.
313
314 *Predictive modeling of plant tissue from gene expression*
315
We used supervised learning classifiers to test if gene expression profiles could predict tissue
type in *Arabidopsis* and if these *Arabidopsis* trained models could be applied more broadly to
flowering plants. We first split the *Arabidopsis* data into testing and training sets with samples
split into four classes of above-ground, below-ground, whole-plant, or other as described above.
Models trained on *Arabidopsis* expression data and used to predict tissue type in *Arabidopsis*,
whether the full or conserved gene datasets, achieved high precision and recall scores. The
highest f1-scores (the harmonic mean of precision and recall) for the full and conserved
datasets were achieved using a K-Nearest Neighbors algorithm (KNN) (0.99 and 0.99,
respectively; **Tables 1 and 2**) and the lowest using Linear Support Vector Classification (SVC)
(0.78 and 0.75). Histogram-Based Gradient Boosting (HGB) also achieved high f1-scores (0.98
and 0.97) while the results for Random Forest (RF) (0.83 and 0.86) and Multilayer Perceptron
(MLP) (0.83 and 0.82) were intermediate. When used to predict *Arabidopsis* data, the precision
and recall values for each model were similar to each other, indicating similar positive prediction
value (precision, true positives divided by true positives and false positives) and sensitivity

(recall, true positives divided by true positives and false negatives). The relative prediction rates
of different tissue types to each other were equivalent for the full *Arabidopsis* dataset (**Fig. 2a**).

Figure 2: Confusion matrices using the KNN-classifier. Confusion matrices showing true label identity (vertical axis) and the proportion of samples assigned to predicted label identities (horizontal axis) for **a)** the full *Arabidopsis* dataset and **b)** the angiosperm dataset. Proportion indicated by viridis color scale.

332
The projection of gene expression patterns from across flowering plants onto a PCA using a
conserved set of genes from *Arabidopsis* shows considerable variability (**Fig. 1c-d**). Using
models trained on *Arabidopsis* data and tested on flowering plants, prediction rates are more
similar to each other using different algorithms than *Arabidopsis* alone but perform much worse,
and with higher precision than recall rates (**Table 2**). For KNN, HGB, RF, MLP, and SVC
methods, precision values were 0.73, 0.74, 0.75, 0.73, and 0.70, respectively, whereas the rates
of recall were 0.64, 0.57, 0.57, 0.55, and 0.58. Although these rates are moderately high, they
must be interpreted in the context of using only four tissue type labels. The relatively higher
precision rates compared to recall indicate that when a sample is retrieved, there is a higher
rate of the models calling a true positive (positive prediction value) compared to the fraction of
relevant samples retrieved (sensitivity). The prediction rates across tissue types were not evenly
distributed (**Fig. 2b**). Below-ground tissue was accurately classified, at a rate of 0.96, while
above-ground tissue was only correctly predicted at a rate of 0.64. Other and whole plant tissue
types were classified poorly (0.074 and 0.32, respectively), and almost no samples were
predicted as other tissue type, including other samples themselves. Although the prediction
accuracy varies considerably across plant families (**Fig. 3**), from around 0.4 to 0.8, we could not
identify any phylogenetic signal or find any support that prediction of tissue identity is inversely
correlated with distance of a plant family from *Arabidopsis* in the Brassicaceae.

351
DISCUSSION
353

Arabidopsis-only models are highly accurate

Although we focus on tissue identity in this
study, we note that the strongest source of
variance (PC1) in publicly available *Arabidopsis*
gene expression profiles is a signature
associated with biotic defense (**Table S1**) and
that it acts in an additive, orthogonal manner
with respect to tissue type which is the next
strongest source of variance (PC2). Not only
are higher prediction rates expected for the
Arabidopsis-only models because the same
dataset is being used for training and testing,
but because of the data structure itself that
separates the main factors we are testing—
above and below ground tissues—as visualized
in a PCA (**Fig. 1a-b**). From this perspective, it is
perhaps not surprising that KNN is the best
performing algorithm, based on the overall

Figure 3: Prediction accuracy by plant family. Using KNN-classifier on the angiosperm dataset, the proportion of samples correctly (blue) and wrongly (orange) predicted is shown as a stacked bar plot.

distance-based proximity of gene expression profiles for each label to each other (**Table 1**). The
other methods, based on decision trees or neural networks, by focusing on individual gene
expression values as parameters, fail to account for overall distance. The focus on individual
gene expression values instead of the overall signature or profile is reminiscent of the molecular
biology concept of "biomarkers" to indicate the tissue or stress from which a sample arises. The
outperformance of KNN over other algorithms we tested may suggest that gene expression
signatures (rather than focusing on individual gene expression values) are more valuable to
create models for tissue and cell type prediction.

Arabidopsis *gene expression as a model for other flowering plants may not be the most suitable*
approach

Lower prediction rates are expected when testing a model on different data than its training set
(**Table 2**). However, the lower precision and recall scores when a model trained on *Arabidopsis*
is tested on gene expression samples across the flowering plants undermines the foundational
argument for using model species: that data from *Arabidopsis* would be predictive for plants in
general. This is not to say that there is not substantial conservation of tissue-specific gene
expression patterns. Our own work (Palande et al., 2023) and that of others (Julca et al., 2021)
strongly supports conserved tissue-specific gene expression patterns across flowering plants,
as is true of animals as well (Fukushima and Pollock, 2020). Rather, the ability to leverage and
predict tissue identity from conserved gene expression profiles is diminished when building a
model from a single, arbitrary species.

395
Details of the performance of our model hint at underlying biological considerations when using
model species data. Not all tissue types are equally predictable, and the prediction of below-
ground tissue outperforms other tissue types (**Fig. 2**). We hypothesized that the ability to predict
tissue identity from *Arabidopsis* may be inversely correlated with phylogenetic distance of a
sample from Brassicaceae, but we found no evidence to support this idea (**Fig. 3**). Additionally,
the precision values for predicting tissue type of flowering plant data from *Arabidopsis* are much
higher than recall values (**Table 2**). This may indicate that models are relatively better at calling
samples with conserved tissue-specificity with *Arabidopsis* (a true positive) over those without (a
false negative). These results may also be a product of our classification scheme. For example,
above and whole plant tissues are often more similar to each other than below ground tissue
because they are missing roots, and might more easily be misclassified with each other. The
other category is composed of diverse tissues which may not have clear predictive features.
These factors should be considered when evaluating the classification results (**Fig. 2**).
409
Our results potentially arise not only from genes with evolutionary differences in tissue-specific
expression compared to *Arabidopsis*, but ones that may indeed have conserved expression but
differ in the ways we have culturally constructed our developmental descriptions of plant
species. Such a circumstance might arise when the cell type-specific expression of a gene is
truly conserved, but that evolved differences in functional morphology between species lead us
to apply different tissue descriptors (for example, between an herbaceous annual and a woody
perennial, or a CAM succulent compared to a weedy C3 plant). The misalignment of tissue
labels extends to more quantitative descriptors and to the molecular level, including Gene
Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) terms that ultimately
become biased to plants with sequenced genomes (Provart et al., 2016). For example, in our
analysis of genes corresponding to the most positive and most negative PC1 loading values,
there was a noticeable enrichment of genes associated with the glucosinolate biosynthetic and
metabolic pathways in *Arabidopsis* samples (**Table S1**). However, this enrichment was absent
in broader angiosperm samples, as these compounds are found almost exclusively in
Brassicaceae. Glucosinolates are a diverse group of secondary metabolites that play a critical
role in plant defense against herbivores and pathogens. Beyond their defensive role, they seem
to be involved in growth, development, microbiota interactions, and phosphate nutrition
(Kopriva, 2021). Focusing on a single organism, or small group of model species to predict
attributes of all plants is flawed from both biological (arising from evolutionary novelty) as well as
philosophical (due to semantic, ontological, and cultural differences in how we socially construct
plants) perspectives.
431
432   *Moving forward and embracing plant and cultural diversity*
433
Arabidopsis was selected as a model species unilaterally, over raised objections, decades ago
arising from mostly genetic and molecular biology considerations (Meyerowitz, 1987; Clough
and Bent, 1998; Arabidopsis Genome Initiative, 2000; Bennett et al., 2003; Bevan and Walsh,
2005). Arguments in favor of plant diversity or selecting agricultural or ecological models were
ignored. These past decisions have led to continued focus on *Arabidopsis* and there is

continuing advocacy for a plant model species and to fund *Arabidopsis* research at the expense
of plant diversity to this current day (Provart et al., 2016; Parry et al., 2020). Since then, data
science and computational approaches have begun to grow. Retrospectively, after which
decades of sequencing data across flowering plants has allowed us to objectively ask if the
focus on a single, arbitrary plant allows us to predict the biology of other flowering plants better
than if we had studied all plants equally from the start, the answer is no (**Table 2**). Using a data
science approach and building machine learning models on *Arabidopsis* gene expression data
to predict the tissue identity of gene expression samples from across flowering plants as we
have done here, does not preclude the consideration of other, more important qualitative
arguments against the model species concept that continue to limit the potential of the plant
science community. Beyond just *Arabidopsis*, there is still a focus on agriculturally important
species at the expense of all plants (Marks et al., 2023). More insidiously, the social construct of
plants and their diversity arises from colonialism, evidenced not only by the gaze of the Global
North and the plants we have chosen to research and document and how we do so, but in ways
that can be quantified related to the specific discussion of *Arabidopsis* here, specifically which
plant genomes have been sequenced and by whom (Marks et al., 2021), usually through
extinguishing and stealing the cultural knowledge of Indigenous people (Dwer et al., 2022).
456
457 Useful discoveries and insights have arisen from *Arabidopsis* (Arabidopsis Genome Initiative,
458 2000). Rather than advocating for continued focus and funding for a single model species
459 (Provart et al., 2016; Parry et al., 2020), it is long past due that we address the historical
460 inequities that have led to our current construction of the plant sciences and that we avoid a
461 biased focus and embrace the biological and cultural diversity of the plant world.
462

463 **Data availability:** The code, metadata, and raw datasets from this project are available on a
464 dedicated GitHub page: https://github.com/PlantsAndPython/arabidopsis-gene-expression
465
Acknowledgements: This work was funded primarily by an NSF-NRT training grant (NSF
1828149) which established the Integrated training Model in Plant And Compu-Tational
Sciences (IMPACTS) program at Michigan State University. This grant funded fellows within this
program as well as the project based curriculum for the Plants and Python Course that formed
the backbone of this manuscript. This work is also supported by NSF Plant Genome Research
Program awards IOS-2310355, IOS-2310356, and IOS-2310357, and NSF Plant, Fungal and
Microbial Developmental Mechanisms award IOS-2039489. This project was supported by the
USDA National Institute of Food and Agriculture, and by Michigan State University
AgBioResearch.

475
476 **REFERENCES**
477

Angiosperm Phylogeny Group, Chase, M.W., Christenhusz, M.J., Fay, M.F., Byng, J.W., Judd,
W.S., Soltis, D.E., Mabberley, D.J., Sennikov, A.N., Soltis, P.S. and Stevens, P.F., 2016. An
update of the Angiosperm Phylogeny Group classification for the orders and families of
flowering plants: APG IV. *Botanical Journal of the Linnean Society*, *181*(1), pp.1-20.

Arabidopsis Genome Initiative, 2000. Analysis of the genome sequence of the flowering plant
Arabidopsis thaliana. *Nature*, *408*(6814), pp.796-815.
485
486 Azodi, C.B., Pardo, J., VanBuren, R., de Los Campos, G. and Shiu, S.H., 2020. Transcriptome-
487 based prediction of complex traits in maize. *The Plant Cell, 32*(1), pp.139-151.
488
489 Bennett, M.D., Leitch, I.J., Price, H.J. and Johnston, J.S., 2003. Comparisons with
490 Caenorhabditis (approximately 100 Mb) and Drosophila (approximately 175 Mb) using flow
491 cytometry show genome size in Arabidopsis to be approximately 157 Mb and thus
492 approximately 25% larger than the Arabidopsis genome initiative estimate of approximately 125
493 Mb. *Annals of Botany*, *91*(5), pp.547-557.
494
495 Bergstra, J., Yamins, D. and Cox, D.D., 2013. Hyperopt: A python library for optimizing the
496 hyperparameters of machine learning algorithms. In *Proceedings of the 12th Python in Science*
497 *Conference* (Vol. 13, p. 20).
498
499 Bevan, M. and Walsh, S., 2005. The Arabidopsis genome: a foundation for plant research.
500 *Genome Research*, *15*(12), pp.1632-1642.
501
502 Clough, S.J. and Bent, A.F., 1998. Floral dip: a simplified method for Agrobacterium-mediated
503 transformation of Arabidopsis thaliana. *The Plant Journal, 16*(6), pp.735-743.
504
505 Coppens, F., Wuyts, N., Inzé, D. and Dhondt, S., 2017. Unlocking the potential of plant
506 phenotyping data through integration and data-driven approaches. *Current Opinion in Systems*
507 *Biology*, *4*, pp.58-63.
508
509 Cortes, C. and Vapnik, V., 1995. Support-vector networks. *Machine Learning*, *20*, pp.273-297.
510
511 Cover, T. and Hart, P., 1967. Nearest neighbor pattern classification. *IEEE transactions on*
512 *Information Theory*, *13*(1), pp.21-27.
513
514 Crossa, J., Perez, P., Hickey, J., Burgueno, J., Ornella, L., Cerón-Rojas, J., Zhang, X.,
515 Dreisigacker, S., Babu, R., Li, Y. and Bonnett, D., 2014. Genomic prediction in CIMMYT maize
516 and wheat breeding programs. *Heredity*, *112*(1), pp.48-60.
517
518 Dwyer, W., Ibe, C.N. and Rhee, S.Y., 2022. Renaming Indigenous crops and addressing
519 colonial bias in scientific language. *Trends in Plant Science*.
520
521 Fukushima, K. and Pollock, D.D., 2020. Amalgamated cross-species transcriptomes reveal
522 organ-specific propensity in gene expression evolution. *Nature Communications*, *11*(1), p.4459.
523
524 Haykin, S., 1998. *Neural networks: a comprehensive foundation*. Prentice Hall PTR.
525

Ho, T.K., 1995, August. Random decision forests. In *Proceedings of 3rd international*
conference on document analysis and recognition (Vol. 1, pp. 278-282). IEEE.

Hogeweg, P., 2011. The roots of bioinformatics in theoretical biology. *PLoS Computational*
Biology, *7*(3), p.e1002021.

Huang, D.W., Sherman, B.T. and Lempicki, R.A., 2009. Systematic and integrative analysis of
large gene lists using DAVID bioinformatics resources. *Nature Protocols*, *4*(1), pp.44-57.

Ij, H., 2018. Statistics versus machine learning. *Nat Methods*, *15*(4), p.233.

Julca, I., Ferrari, C., Flores-Tornero, M., Proost, S., Lindner, A.C., Hackenberg, D.,
Steinbachová, L., Michaelidis, C., Gomes Pereira, S., Misra, C.S. and Kawashima, T., 2021.
Comparative transcriptomic analysis reveals conserved programmes underpinning
organogenesis and reproduction in land plants. *Nature Plants*, *7*(8), pp.1143-1159.

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool,
K., Bates, R., Žídek, A., Potapenko, A. and Bridgland, A., 2021. Highly accurate protein
structure prediction with AlphaFold. *Nature*, *596*(7873), pp.583-589.

Kopriva, S. (2021). Glucosinolates revisited—A follow-up of ABR volume 80: Glucosinolates. In
Advances in Botanical Research (Vol. 100, pp. 249-274). Academic Press.

Knapp S, Bohs L, Nee M, Spooner DM., 2004. Solanaceae—a model for linking genomics with
biodiversity. Comparative and Functional Genomics. Apr;5(3):285-91.

Li, F.W. and Harkess, A., 2018. A guide to sequence your favorite plant genomes. *Applications*
in Plant Sciences, *6*(3), p.e1030.

Lim, P.K., Zheng, X., Goh, J.C. and Mutwil, M., 2022. Exploiting plant transcriptomic databases:
resources, tools, and approaches. *Plant Communications*, p.100323.

Marks, R.A., Hotaling, S., Frandsen, P.B. and VanBuren, R., 2021. Representation and
participation across 20 years of plant genome sequencing. *Nature Plants*, *7*(12), pp.1571-1578.

Marks, R.A., Amézquita, E.J., Percival, S., Rougon-Cardoso, A., Chibici-Revneanu, C., Tebele,
S.M., Farrant, J.M., Chitwood, D.H., VanBuren, R., 2023. A critical analysis of plant science
literature reveals ongoing inequities. *Proc Natl Acad Sci USA*

Mason, L., Baxter, J., Bartlett, P. and Frean, M., 1999. Boosting algorithms as gradient descent.
Advances in Neural Information Processing Systems, *12*.

Mazzocchi, F., 2015. Could Big Data be the end of theory in science? A few remarks on the
epistemology of data-driven science. *EMBO Reports*, *16*(10), pp.1250-1255.

Meyerowitz, E.M., 1987. Arabidopsis thaliana. *Annual Review of Genetics*, *21*(1), pp.93-111.

Meyerowitz, E.M., 2001. Prehistory and history of Arabidopsis research. *Plant Physiology*,
125(1), pp.15-19.

Michael, T.P. and Jackson, S., 2013. The first 50 plant genomes. *The Plant Genome*, *6*(2).

Mitchell, C.E., Agrawal, A.A., Bever, J.D., Gilbert, G.S., Hufbauer, R.A., Klironomos, J.N.,
Maron, J.L., Morris, W.F., Parker, I.M., Power, A.G. and Seabloom, E.W., 2006. Biotic
interactions and plant invasions. *Ecology Letters*, *9*(6), pp.726-740.

Palande, S., Kaste, J.A., Roberts, M.D., Aba, K.S., Claucherty, C., Dacon, J., Doko, R.,
Jayakody, T.B., Jeffery, H.R., Kelly, N. and Manousidaki, A., 2023. The topological shape of
gene expression across the evolution of flowering plants. *PLOS Biology*.

Parry, G., Provart, N.J., Brady, S.M., Uzilday, B., Multinational Arabidopsis Steering Committee,
Adams, K., Araújo, W., Aubourg, S., Baginsky, S., Bakker, E. and Bärenfaller, K., 2020. Current
status of the multinational Arabidopsis community. *Plant Direct*, *4*(7), p.e00248.

Proost, S. and Mutwil, M., 2018. CoNekT: an open-source framework for comparative genomic
and transcriptomic network analyses. *Nucleic Acids Research*, *46*(W1), pp.W133-W140.

Provart, N.J., Alonso, J., Assmann, S.M., Bergmann, D., Brady, S.M., Brkljacic, J., Browse, J.,
Chapple, C., Colot, V., Cutler, S. and Dangl, J., 2016. 50 years of Arabidopsis research:
highlights and future directions. *New Phytologist*, *209*(3), pp.921-944.

Sultan, S.E., 2000. Phenotypic plasticity for plant development, function and life history. *Trends
in Plant Science*, *5*(12), pp.537-542.

Strable J, Scanlon MJ., 2009. Maize (Zea mays): a model organism for basic and applied
research in plant biology. *Cold Spring Harb Protoc*. Oct 1;10(2009):pdb-emo132.

Yu, Y., Zhang, H., Long, Y., Shu, Y. and Zhai, J., 2022. Plant public RNA□seq database: a
comprehensive online database for expression analysis of~ 45 000 plant public RNA□seq
libraries. *Plant Biotechnology Journal*, *20*(5), p.806.

Zhang, N., Wang, M. and Wang, N., 2002. Precision agriculture—a worldwide overview.
Computers and Electronics in Agriculture, *36*(2-3), pp.113-132.

Zhang, H., Zhang, F., Yu, Y., Feng, L.I., Jia, J., Liu, B.O., Li, B., Guo, H. and Zhai, J., 2020. A

comprehensive online database for exploring~ 20,000 public Arabidopsis RNA-seq libraries.

Molecular Plant, *13*(9), pp.1231-1233.

Tables
614
615   **Table 1: Classification performance of models trained on the full *Arabidopsis* dataset.**
616

Model	Precision	Recall	f1-score
SVC	0.765131	0.80103	0.777531
MLP	0.843599	0.844979	0.832854
RF	0.845664	0.826609	0.833746
HGB	0.976665	0.976481	0.976319
KNN	0.98921	0.989185	0.989193

617
618   **Table 2: Classification performance of models trained on the conserved *Arabidopsis***
619   **dataset and tested on conserved *Arabidopsis* or Angiosperm datasets.**
620

Model	Test Set	Precision	Recall	f1-score
SVC	Arabidopsis	0.740855	0.778026	0.754276
	Angiosperm	0.695691	0.576189	0.591683
MLP	Arabidopsis	0.822682	0.828155	0.824351
	Angiosperm	0.734603	0.547361	0.611767
RF	Arabidopsis	0.862941	0.864721	0.861927
	Angiosperm	0.747272	0.569075	0.622122
HGB	Arabidopsis	0.971034	0.970987	0.970574
	Angiosperm	0.741902	0.567952	0.640741
KNN	Arabidopsis	0.987804	0.987811	0.987803
	Angiosperm	0.733478	0.643205	0.663313

621
622   **Figure Legends**
623
Figure 1: Principal Component Analysis (PCA) of gene expression profiles. PCAs with
gene expression profiles colored by above-ground (blue), below-ground (orange), whole plant

(red), and other (green) tissue types for **a)** the full *Arabidopsis* dataset, **b)** the conserved
Arabidopsis data set, **c)** the angiosperm dataset projected onto the conserved *Arabidopsis* PCA
from b), and **d)** the same as c), but with conserved *Arabidopsis* gene expression profiles in the
background (transparent gray).
630
631 **Figure 2: Confusion matrices using the KNN-classifier.** Confusion matrices showing true
632 label identity (vertical axis) and the proportion of samples assigned to predicted label identities
633 (horizontal axis) for **a)** the full *Arabidopsis* dataset and **b)** the angiosperm dataset. Proportion
634 indicated by viridis color scale.
635
636 **Figure 3: Prediction accuracy by plant family.** Using KNN-classifier on the angiosperm
637 dataset, the proportion of samples correctly (blue) and wrongly (orange) predicted from
638 *Arabidopsis* data is shown as a stacked bar plot.

Figure 1: Principal Component Analysis (PCA) of gene expression profiles. PCAs with gene expression profiles colored by above-ground (blue), below-ground (orange), whole plant (red), and other (green) tissue types for **a)** the full *Arabidopsis* dataset, **b)** the conserved *Arabidopsis* data set, **c)** the angiosperm dataset projected onto the conserved *Arabidopsis* PCA from b), and **d)** the same as c), but with conserved *Arabidopsis* gene expression profiles in the background (transparent gray).

Figure 2: Confusion matrices using the KNN-classifier. Confusion matrices showing true label identity (vertical axis) and the proportion of samples assigned to predicted label identities (horizontal axis) for **a)** the full *Arabidopsis* dataset and **b)** the angiosperm dataset. Proportion indicated by viridis color scale.

Figure 3: Prediction accuracy by plant family. Using KNN-classifier on the angiosperm dataset, the proportion of samples correctly (blue) and wrongly (orange) predicted is shown as a stacked bar plot.