1 Non-point source pollution from human activities increasing the

2 biotic homogenization in urban river network system

Haizhou Li[1], Jing Fu[1], Xiangyu Fan[2], Jinchen Yu[1], Zhiwei He[3], Jiawang Wu[4], Li Wu[5], Jin

Zhou[1,*].

Email: zhou_jin@foxmail.com

1. East China Sea Fisheries Research Institute, Chinese Academy of Fishery Science,
Shanghai, China.
2. Department of Molecular and Cellular Biology, University of California, Davis, United
States.
3. State Key Laboratory of Marine Geology, Tongji University, Shanghai, China.
4. School of Marine Sciences, Sun Yat-Sen University, Guangdong, China.
5. Laboratory for Coastal Ocean Variation and Disaster Prediction, College of Ocean and
Meteorology, Guangdong Ocean University, Guangdong, China.

Keywords: river network, non-point source pollution, biotic homogenization.
Subtropical lotic ecosystems.

Abstract

Rivers have been a preferred location for human settlements throughout history. Human

activities were believed to affect the assembly and biogeographic pattern of river

microorganisms, while the underlying processes are not sufficiently understood yet.

Comparing the river microbial communities in highly urbanized rivers and undisturbed

natural rivers could expand our understanding of the river ecosystem responses to

environmental changes resulting from human activities. Here, 28 urban and 24 nature

reserve river surface sediment samples were collected in a highly connected river network

spanning ~400 km located in the Qinling Mountains, Northwest China, and their

geobiochemistry profiles and microbial populations were investigated. The results showed

that human activities resulted in significant increases in organic/inorganic nutrients,

microbial population size and diversity in urbanized rivers. The structure of the microbial

communities in the surface sediments had distinct relationships with environmental factors

and their co-occurrence networks exhibited a clear spatial pattern. Redundancy and

correlation analysis together indicate that nitrogen was the major environmental attribute

that affected the community structure and assembly process. High abundances of nitrogen

transformation microorganisms, pathogens, antibiotic-resistant bacteria and

PAHs-degrading bacteria were observed in urbanized areas. The neutral community model

and null model revealed community composition to be predominant governed by homogeneous selection in urban river areas, suggested the non-point source pollution led to significantly increased biotic homogenization.

1. Introduction

Rivers play a crucial role in aquatic ecosystems as they host the highest levels of biodiversity on Earth (Widder et al. 2014). Therefore, rivers have been a preferred location for human settlements throughout history. However, in recent decades, human activities such as rapid urbanization, industrialization and agriculture has discharged multiple anthropogenic contaminants and non-native (exotic) microorganisms into the river through industrial discharge, agricultural runoff, and municipal sewage (Wang et al. 2012, Beattie et al. 2020). These contaminants can be either organic (e.g., antibiotics, polycyclic aromatic hydrocarbons, fertilizers, pesticides, manure) (Singh et al. 2019, Patel et al. 2020, Liu et al. 2023, Xu et al. 2023) or inorganic contaminants (e.g., nitrate, phosphorus, heavy metal) (Pruden et al. 2012, Manyi-Loh et al. 2016, Yang et al. 2022). At the same time, river sediments often act as the recipients of reservoirs for contaminants influenced by human activity inputs.

In highly urbanized rivers, centralized wastewater treatment plants (WWTPs) are one of the most common systems for the treatment of domestic wastewater. Several recent studies have presented evidence that the WWTPs high-concentration effluent reduces the abundance and diversity of benthic and sediment bacterial communities in urban rivers (Drury et al. 2013, Pascual-Benito et al. 2020, Dai et al. 2023). WWTPs are a common source of point-source pollution (Kiedrzyńska et al. 2014). In contrast, the immense industrial, domestic and agricultural activities sewage are considered as the primary contributors to non-point source pollution (Ding and Liu 2019, Beattie et al. 2020). Point-source pollution is easy to identify, but non-point source pollution is harder to identify and harder to address. United States Environmental Protection Agency reports that non-point source pollution is the leading remaining cause of water quality problems, yet there is little information available on the effects of widespread non-point source pollution on benthic microbial communities in river ecosystems (EPA 2022). The potential ecological and health influences of non-point source pollution in river ecosystems require further research and monitoring.

Here, we focus on a large scale river network, which flows out of the Qinling Mountains, where it eventually empties into the Yangtze River (Fig. 1). The Qinling Mountains in

Northwest China's Shaanxi Province are a geographically important north-south boundary in
China, and an important natural barrier for the eastward flow of moisture from the Tibetan
Plateau, providing a significant source of freshwater for downstream human communities.
The distinctive feature of the river network is that it passes numerous cities (a current
population of about 11.06 million) (Cheng and Duan 2021) and national nature reserves.
Establishing nature reserves is an effective measure to avoid human activity disturbances
(Dudley 2008, Xiao et al. 2022). This river network provides a useful model for
understanding the impact of non-point source pollution come from human activity on the
river ecosystem.
80
In this study, the collected river surface sediment samples (0–2 cm deep) were located in
areas with various levels of human activity, ranging from highly urbanized rivers to
undisturbed nature reserve rivers. River sediments are considered as the major habitat of
microbes in the aquatic environment (Gibbons et al. 2014, Battin et al. 2016). Therefore,
sediment-associated microorganisms drive essential ecological processes and contribute
significantly to river biogeochemical cycles (Xie et al. 2016), and their responses to
human-induced pollution can provide insights into long-term and broader ecosystem health.
Here, these collected surface sediments represent diverse long-term non-point source
pollution gradients emanating from human activity, and creating multiple niches in this river
network. We obtained millions of 16S rRNA tags from the river network sediments by deep
sequencing, and attempted to elucidate the relationship between the microbial community
and environmental factors in the river ecosystem.
93

2. Materials and methods

2.1 Study area, sampling procedure, and physicochemical analysis

Sampling was conducted in August-September 2022 within the river network. From each
sampling site on three sampling occasions, water and sediment samples were collected
within approximately 2 square meters at each site. The geographical data (longitude, latitude,
and distance between sampling sites) were recorded by a GPS device (Dataset 1A). The
water quality parameters, including temperature, pH, dissolved oxygen, water clarity of
Secchi depth, turbidity, total nitrogen, nitrate-nitrogen (NO_3^-), ammonium-nitrogen (NH_4^+),
total phosphors (TP), total dissolved phosphorus (TDP), phosphate, permanganate index,
volatile phenol, *chlorophyll*-a concentration were measured according to Environmental
Quality Standards for Surface Water (Ministry of Ecology and Environment of People's

Republic of China, GB3838-2002). Sediment total organic carbon (TOC) was measured by
the Vario E1-III Elemental Analyzer interfaced with a Thermo Finnigan DELTA plus XL
stable isotope ratio mass spectrometer. Each sample was performed in triplicate, and the
values were expressed as the mean (n = 3).

2.2 Sediment cell counts

Immediately upon returning to the laboratory, composite sediment samples were prepared
by the pooling and mixing of 3 replicates (10 g of each replicate) taken from each site (3
composite samples in total). Cell densities were investigated using catalyzed reporter
deposition-fluorescent *in situ* hybridization (CARD-FISH) based on our previous studies (Li
et al. 2020, Li et al. 2023). Cell counting was performed in triplicate, and the cell abundance
were expressed as the mean (n = 3).

2.3 High-throughput sequencing

The frozen sediment samples (n=3) at each site were mixed for DNA extraction according
our previous study method (Li et al. 2015). The V3–V4 hypervariable region of the 16S rRNA
gene was amplified by PCR using bacterial and archaeal universal primers (515F and 806R).
16S rDNA amplicon samples were sequenced using Illumina MiSeq platform (Illumina Inc.).
The raw data were analyzed according to our previous methods (Li et al. 2020, Li et al.
2023).

Here, the "niche breadth" approach (Levins 1968) was used to quantify habitat specialization.
The formula is described below:

$$B_j = \frac{1}{\sum_{i=1}^{N} P_{ij}^2}$$

where B_j is the habitat niche breadth of OTUj in a metacommunity; and P_{ij} is the proportion
of OTUj in a given site i, and N is the total number of sites. A higher B value of OTUs
indicates a wider habitat niche breadth. In contrast, lower *B* values indicate that the OTUs
occurred in fewer habitats and were unevenly distributed (Pandit et al. 2009). The average
of B values from all OTUs in one community was regarded as the community-level B-value
(*Bcom*) (Jiao et al. 2020). We calculated the *B* value based on species abundance and
evenness for urban and nature reserve samples (Salazar 2020).

2.4 Prediction of microbial functions groups

In order to identify potential microbial ecological functions at the studied sites, the obtained OTUs were compared with the FAPROTAX database (Louca et al. 2016) to predict potential metabolic functions of the microbial community. Furthermore, potential functions of bacterial and archaeal communities were also predicted from 16S rRNA gene-derived OTUs using the Phylogenetic Investigation of Communities by Reconstruction of Unobserved States (PICRUSt2) against the Kyoto Encyclopedia of Genes and Genomes database (KEGG) (Douglas et al. 2020).

2.5 Screening for antibiotic resistant bacteria from river sediments

The approach that has been used to evaluate the status of antibiotic resistance in the environment involves the calculation of resistance percentage, which is based on the ratio between the number of bacteria that are able to grow on culture media that are supplemented with antibiotics at doses close to the minimal inhibitory concentration (MIC) and the number of bacteria that grow on antibiotic-free media (Rizzo et al. 2013, Berendonk et al. 2015). The approach of heterotrophic plate count (HPC) method (Munir et al. 2011) was used to determine the concentration of antibiotic resistant bacteria in the collected river surface sediments. 0.1 g sediment samples were dissolved in 10 mL of 0.85% sterilized normal saline, and diluted 10 × fold for immediate processing. 100 µL diluted samples were cultured onto plate count agar medium. Medium contained four-class plates, 50.4 µg/mL sulfamethoxazole, 16 µg/mL tetracycline, 15□µg/mL azithromycin and no antibiotics. The media was added antifungal agent 200 µg/mL cycloheximide according to Gao el al., (2012) and Brooks et al. (2007) method. Total heterotrophic cultivable bacterial population was determined by plate count agar medium without antibiotics. The selected concentrations of antibiotics on the media were determined according to the criteria of the Clinical and Laboratory Standards Institute (CLSI) guidelines (2015) and the previous study (Munir et al. 2011, Gao et al. 2012, Milaković et al. 2020). Plates were incubated for 48–72 h at 37 °C (Brooks et al. 2007, Munir et al. 2011). Duplicate counting was performed, and colony-forming units (CFU) were calculated. The resistance rates defined as the number of bacteria that grew on antibiotic plates divided by the total number of bacteria that grew on plates containing no antibiotics.

Real-time quantitative PCR (qPCR)

Absolute quantification standard curve method was used to quantify the function genes in the sediment samples. The real-time quantitative PCR (q-PCR) (ABI7500, USA) using SYBR 2×RealUniversal PreMix (Tiangen Biotech). qPCR was performed to assess the abundance of the macrolide, tetracycline and sulfonamide resistant genes. The sequences of the primers and the reaction conditions used in this study are presented in Table S1 in the Supporting Information (SI) section.

The PCR reactions were performed in a 20-µL reaction mixture that comprised 10 µL of SYBR®Premix Ex Taq™ II (2 ×), 0.4 µL of each type of primer (10/20 µM), and 1 µL of a DNA template (5 ng). The thermocycling protocol consisted of an initial denaturation for 1 min at 95 °C, followed by 40 cycles of denaturation (95 °C, 10 s) and annealing at the temperatures in Tables S1. The specificity of the PCR products was checked by a melting curve analysis. All qPCR assays were performed in triplicate for each gene,

The q-PCR standards were synthesized by cloning target genes into *Escherichia coli* DH5a using a PMD18-T vector Cloning Kit (TaKaRa Biotechnology, China). Vector concentrations were measured using NanoDrop, and the quantities of target genes per µL of solution were obtained basing on the length of plasmid and target gene sequence. Six-point calibration curves were generated by 10-fold serial dilutions of the plasmid carrying ARGs. All PCR reactions were run in parallel with serially diluted q-PCR standards and DNA-free water as the negative controls.

2.6 Statistical analysis

Null model analysis was carried out using the framework described by Stegen et al. (2012, 2013) to classify community pairs into underlying drivers of species sorting (or selection), dispersal limitation, homogeneous dispersal, and drift. The variation of both phylogenetic diversity and taxonomic diversity was measured using null model-based phylogenetic and taxonomic β-diversity metrics, namely β-nearest taxon index (βNTI) and Bray–Curtis-based Raup–Crick (RC_{Bray}). A significant deviation (i.e., $|\beta NTI| > 2$) indicates the dominance of selection processes. $\beta NTI < -2$ indicates significantly less phylogenetic turnover than expected (i.e., homogeneous selection) while $\beta NTI > +2$ indicates significantly more phylogenetic turnover than expected (i.e., variable selection). Subsequently, RC_{Bray} was used to further partition the pairwise comparisons that were not assigned to selection (i.e., $|\beta NTI| < 2$). The relative influence of homogenizing dispersal was quantified as the fraction of pairwise comparisons with $|\beta NTI| < 2$ and $RC_{Bray} < -0.95$. Dispersal limitation was quantified as the fraction of pairwise comparisons with $|\beta NTI| < 2$ and $RC_{Bray} > 0.95$. The

fractions of all pairwise comparisons with $|\beta NTI| < 2$ and $|RC_{Bray}| < 0.95$ were used to

estimate influence of "undominated" assembly, which mostly consists of weak selection,

weak dispersal, diversification, and/or drift.

A neutral community model was used to determine the contribution of stochastic processes

to microbial community assembly by predicting the relationship between the frequency with

which taxa occur in a set of local communities and their abundance across the wider

metacommunity (Sloan et al. 2006). The model predicts that abundant taxa are more likely

to be dispersed by chance and widespread across metacommunity, while rare taxa would be

lost in different local communities due to ecological drift. In the model, the estimated

migration rate is a parameter for evaluating the probability that a random loss of an individual

in a local community would be replaced by dispersal from the metacommunity, and,

therefore, is a measure of dispersal limitation. Higher m values indicate that microbial

communities are less dispersal limited. The formula is as follows:

$$Freq_i = 1 - I(1/N \mid N*m*p_i, N*m*(1 - p_i))$$

where $Freq_i$ is the occurrence frequency of taxon i across communities; N is the number of

individuals per community; m is the estimated migration rate; p_i is the average relative

abundance of taxon i across communities; and $I()$ is the probability density function of beta

distribution. R^2 indicates the fit of the parameter based on nonlinear least-squares fitting.

2.7 Data available

Sequencing data are stored on the National Genomics Data Center (NGDC) Sequence

Read Archive (PRJCA018972).

3. Results

3.1 Biogeochemical properties of river water and sediment samples

A total of 28 urban river and 24 undisturbed pristine nature reserve river surface (0-2 cm)

sediment samples were collected (Fig. 1). Samples from all 52 locations covered a wide

range of horizontal distance (0.72-376 km). The environmental chemical profiles of river

water showed remarkable differences between the two regions (Fig. 2, Dataset 1A). The

differences mainly included significantly higher concentration of total nitrogen (TN),

ammonium, nitrite, nitrate, total phosphorus (TP), total dissolved phosphorus (TDP),

phosphate, volatile phenol, Chlorophyll a in urban river samples ($P<0.01$). Ammonium and

nitrate were the dominant forms of dissolved nitrogen within the rivers. The total organic

carbon (TOC) content in sediment also showed a significant increasing trend from the
natural river to urban river ($P<0.0001$) (Fig. 2).

3.2 Microbial community abundance, richness and diversity

Microbial abundance ranged from 1.3×10^7 to 1.1×10^9 cells g-1 and 8.0×10^6 to 7.1×10^8
cells g-1 for urban and natural river samples, respectively (Fig. 3A, details in Dataset 2A).
The urban sites exhibited a significantly higher abundance of microorganisms than those
from the natural river samples ($P<0.01$), indicated that the human activities have a great
impact on microbial population.
239
Using universal primers for bacteria and archaea, 3,309,566 high-quality sequences were
generated throughout the river network (Dataset 2A, SI Appendix Fig. S1). These
high-quality sequences were clustered into 18,038 operational taxonomic units (OTUs, 97%
similarity level) (Dataset 2B). Diversity within each samples was calculated at an equivalent
sequencing depth across all samples using OTU-level based on three sediment composite
samples per site (Materials and Methods). A range of 1,628–6,104 chao1 OTUs and
1.89–6.68 Shannon index were observed in the urban samples, and a range of 1,321–5,471
chao1 OTUs and 1.88–6.31 Shannon index were observed in nature reserve samples (Fig.
3B-C). Notably, both the Chao 1 and Shannon demonstrated that microbial richness and
diversity were considerably higher in the urban river compared to the natural river samples
($P < 0.05$). Furthermore, the niche breadths (*Bcom*) (Fig. 3D) exhibited the
narrowest *Bcom* values in urban areas, rather than those in natural river.

3.3 Microbial community structure and composition

80 microbial phyla and 3,570 species were identified in the urban samples, and 74 phyla and
3,371 species were identified in nature reserve samples (Dataset 2C-F). At the phylum level,
the urban and nature reserve samples investigated here shared a high proportion of taxa.
The ten most prevalent phyla in all of sites collectively constituted 95.48% of the total
sequences (Fig. 4A), matching the classical abundance-range relationship (Amend et al.
2013, Ruff et al. 2015). The dominant microbial phyla included Proteobacteria (28.69%),
Firmicutes (28.32%), Actinobacteriota (17.79%), Chloroflexi (4.99%), Bacteroidota (4.51%),
Acidobacteriota (3.51%), Planctomycetota (3.51%), Verrucomicrobiota (2.43%),
Deinococcota (1.01%), Cyanobacteria (0.72%) (Fig 4A, Dataset 2C).
262

Archaea constituted only a minor fraction of the river sediment communities (0.02~4.96%),

and an increase in the diversity and proportion of the archaea was identified in urban areas.

In the urban areas, proportion of the archaea exhibited 1.58 times higher than those in

natural river, and archaeal Chao1 richness and Shannon diversity exhibited 1.47 and 2.36

times higher than those in natural river, respectively.

Although the microbial community of these sites exhibited similarities at the phylum level, 38

order in samples showed significant differences (P<0.01) between urban and nature reserve

river ecosystems (Fig. 4B, SI Appendix Fig. S1B). For example, Bacillales, Clostridiales,

Steroidobacterales were dominant in urban areas, while Erysipelotrichales, Deinococcales,

Alicyclobacillales, Pyrinomonadales, Burkholderiales were dominant in nature reserve areas

(SI Appendix Fig. S2B, Dataset 2C).

A heatmap was generated to represent the 50 most prevalent species (Fig. 5), and most of

these dominant OTUs were classified as unclassified species. Commonly observed with

relatively high abundances in all samples were *Exiguobacterium-sibiricum*, *Exiguobacterium*

sp., and *Acinetobacter lwoffii*. In urban samples, the dominant taxa included Unclassified

Planococcaceae, Unclassified Steroidobacteraceae, Unclassified Anaerolineaceae, and

Unclassified Rhizobiales. On the other hand, the unclassified *Acinetobacter*, unclassified

Pseudomonas, unclassified Sphingomonadaceae, unclassified *Deinococcus*, *Pseudomonas*

parafulva had high abundances in nature reserve samples.

Furthermore, the analysis of similarities showed that the urban samples harbored

significantly dissimilar microbial communities at OTU level (Fig. 3E; PERMANOVA pseudo F

= 2.1455, p.adj = 0.024, 999 permutations; ANOSIM R sqr = 0.0597, P = 0.03). In other

words, our analyses show a greater similarity in the community composition of archaea,

bacteria across the urban river than across the corresponding natural river ecosystems (Fig.

3F).

3.5 Prediction of the functional diversity and metabolic pathways

Microbial community metabolic predictions guided by the FAPROTAX database (Louca et al.

2016) revealed distinctions in the metabolic potential related to nitrogen, carbon, and sulfur

cycling between urban and nature reserve areas (Dataset 4A-D). For nitrogen

transformation, the microorganisms involved in ammonia oxidation, nitrite oxidation, nitrogen

fixation, nitrification at urban rivers was higher than that at nature reserve (ANOVA,

P < 0.001). For carbon transformation, urban areas exhibited a higher abundances of

microbial groups associated with anoxygenic photoautotrophy, oxygenic photoautotrophy,
chloroplasts, and photoautotrophy. Furthermore, the urban areas displayed a higher
abundance of taxa involved in degradation of aromatic compound compared to the nature
reserve (ANOVA, $P < 0.05$) (Fig. 3I). The sulfur cycling displayed heterogeneity among the
samples. As depicted in Fig. 3I, the urban areas exhibited abundant taxa associated with the
dark oxidation of sulfur compounds (ANOVA, $P < 0.05$). Furthermore, the abundance of taxa
involved in sulfate respiration was higher in urban areas.
305
Additionally, the PICRUSt2 database (Douglas et al. 2020) annotated 414 functional KEGG
pathways (Dataset 4C-F). The metabolism associated with carbon fixation, nitrogen,
sulfur, phosphate, antibiotic biosynthesis and xenobiotic biodegradation were investigated.
The presence of antibiotic resistance proteins was also detected (Dataset 4B, D, F). In all
samples, biodegradation pathways encompassing 21 types of xenobiotic compounds were
determined (Dataset 4F-G). The ecological function of xenobiotic biodegradation exhibited a
noticeable increasing trend from the nature reserve to the human-impacted sites. This trend
suggested an increase in pollutants discharge from urban rivers. Styrene degradation
emerged as the predominant process among polycyclic aromatic hydrocarbon (PAH)
degradation (Dataset 4G).

3.6 Antibiotic resistance rates and genes

Three kinds of antibiotic, macrolides, tetracycline and sulfonamide, were selected for
bacteria resistance test. Macrolides and tetracyclines are among the most widely used
antibiotics globally for both human and agricultural applications (Van Boeckel et al. 2014,
WHO 2019). Sulfonamides were the first synthetically produced antibiotic for widespread
use and extensive resistance has since emerged among clinical isolates (Sköld 2000).
322
All of collected samples in this study showed macrolide, tetracycline and sulfonamide
resistance activity. Fig 6A illustrated the macrolides, tetracycline and sulfamethoxazole
resistance rates on plating count agar. The urban samples antibiotic resistant rate ranged
11.2% - 30.5 % for azithromycin, 13.0-29.0 % for tetracycline, 11.0-26.9 % for sulfonamide.
The natural river samples antibiotic resistant rate ranged 2.4-18.2% for azithromycin,
7.3-18.2 % for tetracycline, 2.4-16.0 % for sulfonamide. The antibiotic resistant rate in urban
samples were higher than those in natural river samples. In the natural river samples, no
significant differences in antibiotic resistance rate were observed, but variability in antibiotic
resistance rates were observed in the urban samples.
332

Five ARGs representing three classes of resistance (two sulfonamide: *sul*1 and *sul2*;
two tetracycline: *tet*O and *tetW*; and one macrolide: *erm*F) were selected for quantification in
all samples to characterize how loads of ARGs change between urban and natural rivers.
The five target ARGs were selected due to their documented prevalence in watersheds
impacted by anthropogenic and agricultural runoff (Garner et al. 2017). All five of these
genes were among those identified as candidate indicators of the extent of the impact of
antibiotic resistance in the environment by the European Cooperation in Science and
Technology Action (Berendonk et al. 2015).
341
The ARGs were found in all samples, and Fig. 6B shows the occurrences and quantities of
ARGs in the sediments of the urban and natural rivers. The concentrations of ARGs varied
greatly, ranging from 1.04×10^3 copies/g (*sul2*, Natural River) to 4.10×10^8 copies/g (*tetO*,
urban river). The quantity of ARGs in the urban river ranged from 1.36×10^4 to 1.63×10^8
copies/g for *sul1*, 2.27×10^4 to 3.00×10^8 copies/g for *sul2*, 5.36×10^3 to 4.10×10^8
copies/g for *tetO,* 2.52×10^3 to 2.17×10^8 copies/g for *tetW,* 1.72×10^3 to 2.02×10^8
copies/g for *ermF*. The quantity of ARGs in the natural river ranged from 1.42×10^3 to
9.49×10^6 copies/g for *sul1*, 1.04×10^3 to 1.9×10^7 copies/g for *sul2*, 1.49×10^3 to 2.5×106
copies/g for *tetO,* 2.34×10^3 to 9.45×106 copies/g for *tetW, 2.12* $\times 10^3$ to 5.52×10^6
copies/g for *ermF*. The quantity of ARG abundances in the sediments of the urban river were
higher than those in the sediments collected from the natural river. A wide concentration
range for each class of resistance genes was observed in the urban river in the tetracycline,
macrolide and sulfonamide resistance genes. In contrast, this trend was not observed in the
natural river.

3.7 Key influencing factors and ecological driving processes

The distance-based redundancy analysis (dbRDA) using environmental parameters as
constraining variables, highlighted the potential significance of these chemical parameters in
elucidating microbial community compositions (Fig. 3F). Ammonium (r sqr= 0.3551,
P=0.001), pH (r sqr= 0.2293, *P*=0.005), Nitrite (r sqr=0.2211, *P*=0.003), Nitrate (r sqr=0.188
P=0.006), TN (r sqr= 0.17, *P*=0.006), TOC (r sqr= 0.1367, *P*=0.032), phosphate (r sqr=
0.0615, *P*=0.010) were more crucial in shaping the microbial community (Fig. 3F).
Ammonium exhibited the strongest influence on bacterial and archaea community
composition within the river network (*P*$\Box$=$\Box$0.001).
365
Across the large river network spanning a geographic distance of ~400 km, the microbial
community increased in dissimilarity with increasing spatial distance, showed a
distance-decay relationships (DDRs) (Fig. 7A). To further assess the underlying factors

governed community dynamics in the river network, the null model (Stegen et al. 2012) and
neutral community model (NCM) (Sloan et al. 2006) were used to quantify the relative
influences of deterministic and stochastic processes. The null model indicated a prevalence
of environmental selection processes, particularly in the form of homogeneous selection
($|\beta NTI| < 2$, RCbary >+0.95) (Fig. 7B, Dataset 3), in governing the community assembly
(details in Materials and Methods). Simultaneously, neutral community model (Fig. 7C)
showed a gradual decline in the contribution of stochastic process with the escalation of
human activity, elucidating 58.5% and 48.3% of the stochastic process within the nature
reserve and urban samples, respectively.

6. Discussion

Urbanization changed the river community diversity, compositions and function
Considerable variation exists within the bacterial and archaeal communities in terms of
abundance, richness, diversity (Fig. 3A-C) and composition (Fig. 3E) between urban and
natural rivers, accompanied by significant niche differentiation (Fig. 3D). Nonmetric
multidimensional scaling (NMDS) (Fig. 3F) and network analysis (Fig. 3I), based on OTU,
demonstrated distinct differences between the microbial communities of urban and those of
the natural river ecosystems. This differentiation persisted even when considering the
exclusion of "rare biosphere" taxa (< 0.1% total sequence).
387
The db-RDA analysis suggested that nitrogen (often in the form of nitrate and ammonium)
and pH contributed significantly to the changes in the urban river sediment microbial
community (Fig. 3G). The high nitrogen loading and lower pH (Fig. 2), are common
consequences of the continuous input of industrial, domestic, and agricultural wastewater,
which was caused by a large population of about 11.06 million in sampled regions (Cheng
and Duan 2021). Excessive nitrogen inputs can lead to eutrophication, a process where the
nutrient enrichment promotes the growth of algae (Chlorophyll *a* content in Fig. 2) (Kuypers
et al. 2018). The proliferation of algae leads to increased organic matter production, which is
subsequently decomposed by bacteria and archaea (TOC content in Fig. 2). The
''productivity/resources hypothesis'' posits that greater resource availability and/or primary
production can support larger population sizes and limit local extinction, promoting species
coexistence (Ibarbalz et al. 2019). Therefore, the influx of organic compounds and nutrients
into the urban river created a more uniform environment, allowing for the coexistence of a
broader range of microbial species. The increased availability of organic matter and the
changed nutrient conditions can profoundly influence the river microbial population.
403
Furthermore, the urban river harbored communities of bacteria and archaea distinct from
those found in the natural river ecosystems (Fig. 3F), with urban river consistently supporting
a significantly greater proportion of Erysipelotrichales, Deinococcales, Methanobacteriales,
Burkholderiales, Chthoniobacterales, Caulobateriales (Fig. S2). The impact of human

activities on river microbial communities was also evident in the predicted metabolic
functions of these communities (Fig. 3I). Therefore, we further investigated key function taxa,
and the results proved the enrichment of nitrogen-transformation microorganisms,
pathogens, PAHs-degrading bacteria, antibiotic-resistant bacteria and the archaea in the
urban river samples.
413
(i) Nitrogen-enriched urbanized rivers exhibited a notably higher abundance of
nitrogen-transformation taxa groups (Dataset 2E-F), including nitrogen-fixing bacteria
(*Anaeromyxobacte*) (Masuda et al. 2020), nitrifying bacteria (such as *Nitrosomonas,*
Nitrospina, Nitrosospira, Nitrospira) and denitrifying bacteria (such as *Paracoccus,*
Hyphomicrobium, and *Gemmobacter*) (Lücker et al. 2010). These taxa are ubiquitous
participants in the nitrogen cycle. The inputs of nitrogen from human activity can lead to
increased abundance of nitrogen-transformation taxa groups (Daims et al. 2015, Henson et
al. 2018), consequently elevating nitrate concentrations in waters (Fig. 2).
422
(ii) Within the urban river, the high abundance of Bacteroidetes and Firmicutes, which are
recognized as good indicators of fecal contamination (Wéry et al. 2008, Wang et al. 2018),
were observed (Dataset 2C). The elevated abundance of Bacteroidetes and Firmicutes in
the urban river areas suggested the presence of untreated wastewater from residential
areas, which may contain fecal pollutants. Additionally, the study also identified a marked
increase in the abundance and diversity of well-known pathogens like toxic *Acinetobacter*,
Arcobacter, *Exiguobacterium*, *Chryseobacterium*, *Escherichia Shigella* and *Staphylococcus*
aureus within urban river environment (Dataset 2E-F). These findings highlighted a distinct
change in microbial structure due to human activities leading to fecal contamination (Drury et
al. 2013, García-Armisen et al. 2014). These genera also have demonstrated a remarkable
capacity to degrade and tolerate a wide array of organic compounds found in sewage water
(Kasana and Pandey 2018, Zhang et al. 2020). This dual capability underscored their
adaptability to environments influenced by human activities and suggested their involvement
in sewage degradation processes (Zhang et al. 2009, VandeWalle et al. 2012).
437
(iii) High abundances of ecological function linked to xenobiotic biodegradation were
detected in urban sites (Dataset 4G). The observed high abundance of genes related to
xenobiotic biodegradation in the urban river is consistent with the findings of a previous
study (Bai et al. 2014). The presence of biodegradation genes could serve as indicators for
the existence of xenobiotics and their corresponding metabolites (Pagé et al. 2015, Kansole
and Lin 2016). Furthermore, the high abundance of PAHs-degrading bacteria such as
Sphingobium, *Rhodococcus*, *Novosphingobium*, *Pseudomonas*, *Sphingomonas*,
Brevibacterium, *Arthrobacter*, *Nocardioides,* and *Mycobacterium* were observed in the urban

river (Datset 2E) (Ghosal et al. 2016), may suggested the accumulation of PAHs in the urban
river sediment. PAHs primarily originating from long-term anthropogenic pollution sources
linked to rapid industrialization and urbanization (Kuppusamy et al. 2017, Mojiri et al. 2019).
These PAHs pollutants, characterized by persistence and low solubility, tend to accumulate
in river sediments due to their hydrophobic nature and strong adsorption onto sediment
particles (Kuppusamy et al. 2017, Patel et al. 2020). Numerous urban rivers in China have
been identified as containing high abundance of PAHs (Cao et al. 2010, Qian et al. 2017),
and significant high biodegradation of PAHs capacity has also been observed in urban areas
(Li et al. 2015, Muangchinda et al. 2017).
455
(iv) Functions linked to drug and antimicrobial metabolism were identified in urban river and
nature reserve river sediment samples (Dataset 4). Notably, higher antibiotic resistance
rates and ARGs were observed in highly urbanized river areas (Fig. 6). The presence of
antibiotics in the river is a widespread concern. Zhang et al. (2015) reported that over 68% of
total antibiotic usage in China in 2013 was excreted by humans and animals, leading to
around 53,800 tons of antibiotics entering the environment (Zhang et al. 2015). Urban water
ecosystems have been identified as potential hotspots for the spread of antibiotic resistant
bacteria in China (Hou et al. 2021, Reddy et al. 2022). The continuous selective pressure
exerted by antibiotic residues, even at sub-inhibitory concentrations, in combination with the
diverse and dense microbial communities sustained by nutrient-rich environments, creates
favorable conditions for the dissemination and proliferation of antibiotic resistance (Rizzo et
al. 2013). Notably, the data showed that significant variation between cities in antibiotic
resistance rates and genes, with no significant differences in natural river samples (Fig. 6).
This correlation was highly significant for antibiotic use and population. This feature
emphasized the importance of the different levels of human activity contribution based on
population size, potentially resulting in varied antibiotic resistance rates among urban
regions. Additionally, pollutants from human activities, such as heavy metals and
disinfectants, further exerted selective pressure on bacteria, contributing significantly to
antibiotic resistance development (Baker-Austin et al. 2006).
475
(v) An increase in the diversity and proportion of the archaea was also identified under the
impact of human activities pollution effluent. In the urban river areas, proportion of the
archaea exhibited 1.58 times higher than those in natural river, and archaeal Chao1 richness
and Shannon diversity exhibited 1.47 and 2.36 times higher than that in the natural river,
respectively. This finding is similar to previous study revealed that in a high-contamination
zone, archaeal Chao1 richness and Shannon diversity exhibited substantial increases,
approximately 1.81 and 1.56 times higher than those in a low-contamination zone,
respectively (Yang et al. 2022). Krzmarzick et al. (2018) proposed that archaea have
demonstrated a notable capacity for enduring harsh environments, leading to the possibility

that numerous archaeal species possessing exceptional adaptability to severe
anthropogenic pollution could flourish in urban river environments (Krzmarzick et al. 2018).
487
(vi) The occurrences of above mentioned taxa observed in the urban river highlighted the
role of human activity influence in urban rivers. This observation is further supported by the
presence of taxa associated with specific functions, such as *Rhodoferax*, *Thiobacillus*,
Methylotenera, *Crenothrix*, *Planktothrix, Microcystis*, *Anabaena*, and *Flavobacteria,* which
exhibited higher abundance in urban areas (Dataset 2E-F). While these taxa are generally
found in freshwater, their prevalence in elevated abundances often links to the presence of
human-impacted pollutants (Beattie et al. 2020). Genera like *Rhodoferax* and *Thiobacillus*
had been identified for their capabilities in iron reduction and sulfur oxidation, respectively
(Finneran et al. 2003, Zhi-Hui et al. 2010). *Methylotenera* and *Crenothrix* play important
roles in freshwater carbon cycling by utilizing reduced carbon sources such as methane as
their primary carbon source (Oshkin et al. 2014, Oshkin et al. 2015, Oswald et al. 2017).
Additionally, previous studies showed that increases in Cyanobacteria, particularly genera
such as *Planktothrix*, *Anabaena*, and *Microcystis* were linked to occurrences of freshwater
eutrophication (Dolman et al. 2012, Domingues et al. 2014). *Flavobacteria also* were
abundant and common heterotrophs in freshwater environments (García-Armisen et al.
2014), and played a crucial contribution in various carbon cycles, such as the degradation of
algal organic matter and other forms of biodegradation (Williams et al. 2013).
505
Overall, our research demonstrated that human population alongside contamination level
serve as the predominant influences shaping urban river microbial communities in the river
network ecosystem. Some bacterial and archaeal communities might have adapted to these
altered conditions over time (Archana et al. 2016, Eckert et al. 2018). Therefore, the
changed environmental conditions can potentially favor specific bacterial and archaeal taxa,
leading to shifts in community composition. These changes in nutrients have cascading
effects on ecosystem functioning and overall microbial processes in the river network. Given
the human activities that carry a myriad of exotic microorganisms and organic/inorganic
contaminants into the urban river, urbanization can profoundly change the riverine
ecosystems (Drury et al. 2013, Xuan et al. 2022).
516
Urbanization non-point source pollution led to increased biotic homogenization
River network sediment communities exhibited high diversity, encompassing thousands of
OTUs across major taxa that inhabit various habitats within the river network (Fig. 3-5).
Intriguingly, the distance-decay relationship (DDR) (Fig. 6) was observed for urban and
natural river samples. Stochastic neutral models postulate that water mass transport must
interact with dispersal limitation to create a distance-decay pattern (Hutchison and
Templeton 1999, Hubbell 2001, Zhang et al. 2014). Therefore, these findings underscored

the coexistence of simultaneous niche-based deterministic and neutral-based stochastic

dispersal processes that jointly govern the community assembly and the biogeographical

signature (Dini-Andreote et al. 2015, Jia et al. 2018).

Previously, the investigation on the impacts of geographic distances and environmental

conditions on microbial community assembly has been a research hotspot in river ecology

(Dini-Andreote et al. 2015, Jia et al. 2018). In large scale watersheds such as Yangtze River

and Tingjiang River, geographic distance was proven key factor governing community

succession (Chen et al. 2019, Gao et al. 2021, Yang et al. 2022). However, in the present

study, we only observed weak DDRs in urban river (Fig. 6). In other words, the geographic

variation in microbial community composition is lower across cities than across natural river

ecosystems. These data indicated that the spatial structures of microbial communities were

less apparent in human-impacted urbanization river ecosystems, emphasizing the role of the

environmental selection. Furthermore, our analyses also show narrower niche

breadths were witnessed for samples in urban river (Fig. 3D), and a greater similarity in the

community composition of archaea, bacteria along the urban river than across the

corresponding natural river ecosystems (Fig. 3F). This finding suggests that the

homogenization effects of human activities and the potential environmental drivers rise from

non-point source pollution created a more uniformity environment in urban river system and

suggesting a homogenization in the urban river microbiome.

Further, the application of the neutral community model (Fig. 7C) in the present study

revealed that deterministic processes predominantly shaped the urban river microbial

communities in the river network. Null model uncovered the homogeneous selection

dominated governs the urban river community assembly, suggested that persistent uniform

non-point source pollution such as high loading nitrogen, PAHs, antibiotic and so on arises

from urbanization human activity exert robust environmental selection deterministic

influences on river system. Therefore, similar urban environmental condition could promote

community fusion, contributing to a more similar community structure (Fig. 3F-G).

According to the United Nations, 68% of the global population will live in cities by 2050,

increasing the environmental and social stresses for the billions of humans living in urban

areas. Previous studies showed that human-induced land-use changes result in the biotic

homogenization of soil microbial communities (Rodrigues et al. 2013). Moreover, land-use

intensification causes multitrophic homogenization of grassland communities have been

previously reported at larger spatial scales (Gossner et al. 2016). Likewise, homogenization

of bacterial communities has also been reported in dust samples collected from urban areas
compared with more rural areas across the United States (Barberán et al. 2015). Our results,
emphasizing a convergence (reduced geographic variation) in the urban rivers microbiome,
are consistent with the effects of urbanization on bird (Murthy et al. 2016), plant, and
invertebrate communities (McKinney 2008). Together, our findings provide novel evidence
that urban river is important hot spots for the local (alpha) diversity of microbes, and allows
for the coexistence of a broader range of microbial species.
567
This result provided an excellent illustration of the concept of biotic homogenization, which
suggested that human modifications of the environment are reducing the biological
differences that exist among natural ecosystems. This process is predicted to result in a
more homogenized biosphere on regional and global scales (McKinney 2006). Further
investigations are needed to explore the universality of biotic homogenization due to human
activity effluents across a range of river ecosystems.

5. Conclusions

Despite the rivers' inherent capacity for dilution and self-purification, the river network was
vulnerable to non-point source pollution from human activities. We found that urban river is
important hot spots of local microbial taxonomic and functional diversity but also support a
more homogenization in the structure and function of the river microbiome. More specifically,
we show that urban river harbor more similar microbiomes than adjacent natural river
ecosystems. The urban river is characterized by higher proportions of fast-growing bacteria,
algae, nitrifiers, PAHs degradation bacteria, and important pathogens, fecal coliform
bacteria and antibiotic resistant bacteria, which were particularly dominant in the nutrient rich,
more population, and more intensively human activity urban river. Our analyses indicate that
anthropogenic pollutants stand out as the principal drivers altering microbial communities,
consequently augmenting the role of homogeneous selection in shaping urban microbial
community assembly. Together, our study represents a comprehensive assessment of the
structure and functional differences in the urban and natural river microbiome.

Acknowledgements

This work was supported by the Youth Program of National Natural Science Foundation of
China (No. 32100081), Central Public-interest Scientific Institution Basal Research Fund,
CAFS (NO.14).

Competing Interest Statement

The authors declare no competing interests.

Declaration of Generative AI and AI-assisted technologies in the writing process

Statement: During the preparation of this work the authors used CHATGPT in order to
improve readability and language. After using this tool/service, the authors reviewed and
edited the content as needed and takes full responsibility for the content of the publication.

Reference

Amend, A. S., T. A. Oliver, L. A. Amaral-Zettler, A. Boetius, J. A. Fuhrman, M. C.
Horner-Devine, S. M. Huse, D. B. M. Welch, A. C. Martiny and A. Ramette (2013).
"Macroecological patterns of marine bacteria on a global scale." Journal of Biogeography
40(4): 800-811.
Archana, A., L. Li, K. Shuh-Ji, B. Thibodeau and D. M. Baker (2016). "Variations in nitrate
isotope composition of wastewater effluents by treatment type in Hong Kong." Marine
Pollution Bulletin **111**(1): 143-152.
Bai, Y., W. Qi, J. Liang and J. Qu (2014). "Using high-throughput sequencing to assess the
impacts of treated and untreated wastewater discharge on prokaryotic communities in an
urban river." Applied Microbiology and Biotechnology **98**(4): 1841-1851.
Baker-Austin, C., M. S. Wright, R. Stepanauskas and J. V. McArthur (2006). "Co-selection of
antibiotic and metal resistance." Trends in Microbiology **14**(4): 176-182.
Barberán, A., J. Ladau, J. W. Leff, K. S. Pollard, H. L. Menninger, R. R. Dunn and N. Fierer
(2015). "Continental-scale distributions of dust-associated bacteria and fungi." Proc Natl
Acad Sci U S A **112**(18): 5756-5761.
Battin, T. J., K. Besemer, M. M. Bengtsson, A. M. Romani and A. I. Packmann (2016). "The
ecology and biogeochemistry of stream biofilms." Nature Reviews Microbiology **14**(4):
251-263.
Beattie, R. E., A. Bandla, S. Swarup and K. R. Hristova (2020). "Freshwater sediment
microbial communities are not resilient to disturbance from agricultural land runoff." Frontiers
in Microbiology **11**: 539921.
Berendonk, T. U., C. M. Manaia, C. Merlin, D. Fatta-Kassinos, E. Cytryn, F. Walsh, H.
Bürgmann, H. Sørum, M. Norström, M.-N. Pons, N. Kreuzinger, P. Huovinen, S. Stefani, T.
Schwartz, V. Kisand, F. Baquero and J. L. Martinez (2015). "Tackling antibiotic resistance:
the environmental framework." Nature Reviews Microbiology **13**(5): 310-317.

Brooks, J., S. Maxwell, C. Rensing, C. Gerba and I. Pepper (2007). "Occurrence of
antibiotic-resistant bacteria and endotoxin associated with the land application of biosolids."
Canadian Journal of Microbiology **53**(5): 616-622.
Cao, Z., J. Liu, Y. Luan, Y. Li, M. Ma, J. Xu and S. Han (2010). "Distribution and ecosystem
risk assessment of polycyclic aromatic hydrocarbons in the Luan River, China."
Ecotoxicology **19**(5): 827-837.
Chen, W., K. Ren, A. Isabwe, H. Chen, M. Liu and J. Yang (2019). "Stochastic processes
shape microeukaryotic community assembly in a subtropical river across wet and dry
seasons." Microbiome **7**(1): 138.
Cheng, M. and C. Duan (2021). "The changing trends of internal migration and urbanization
in China: new evidence from the seventh National Population Census." China Population
and Development Studies **5**: 275-295.
Dai, T., Z. Su, Y. Zeng, Y. Bao, Y. Zheng, H. Guo, Y. Yang and D. Wen (2023). "Wastewater
treatment plant effluent discharge decreases bacterial community diversity and network
complexity in urbanized coastal sediment." Environmental Pollution **322**: 121122.
Daims, H., E. V. Lebedeva, P. Pjevac, P. Han, C. Herbold, M. Albertsen, N. Jehmlich, M.
Palatinszky, J. Vierheilig, A. Bulaev, R. H. Kirkegaard, M. von Bergen, T. Rattei, B.
Bendinger, P. H. Nielsen and M. Wagner (2015). "Complete nitrification by Nitrospira
bacteria." Nature **528**(7583): 504-509.
Ding, X. and L. Liu (2019). "Long-term effects of anthropogenic factors on nonpoint source
pollution in the upper reaches of the Yangtze river." Sustainability **11**(8): 2246.
Dini-Andreote, F., J. C. Stegen, J. D. van Elsas and J. F. Salles (2015). "Disentangling
mechanisms that mediate the balance between stochastic and deterministic processes in
microbial succession." Proc Natl Acad Sci U S A **112**(11): 1326-1332.
Dolman, A. M., J. Rücker, F. R. Pick, J. Fastner, T. Rohrlack, U. Mischke and C. Wiedner
(2012). "Cyanobacteria and cyanotoxins: the influence of nitrogen versus phosphorus." PloS
one **7**(6): e38757.
Domingues, R. B., A. B. Barbosa and H. M. Galvão (2014). "River damming leads to
decreased phytoplankton biomass and disappearance of cyanobacteria blooms." Estuarine,
Coastal and Shelf Science **136**: 129-138.
Douglas, G. M., V. J. Maffei, J. R. Zaneveld, S. N. Yurgel, J. R. Brown, C. M. Taylor, C.
Huttenhower and M. G. I. Langille (2020). "PICRUSt2 for prediction of metagenome
functions." Nature Biotechnology **38**(6): 685-688.
Drury, B., E. Rosi-Marshall and J. J. Kelly (2013). "Wastewater treatment effluent reduces
the abundance and diversity of benthic bacterial communities in urban and suburban rivers."
Appl Environ Microbiol **79**(6): 1897-1905.
Dudley, N. (2008). Guidelines for applying protected area management categories, Iucn.
Eckert, E. M., A. Di Cesare, M. T. Kettner, M. Arias-Andres, D. Fontaneto, H.-P. Grossart
and G. Corno (2018). "Microplastics increase impact of treated wastewater on freshwater
microbial community." Environmental Pollution **234**: 495-502.
EPA. (2022). "Basic Information about Nonpoint Source (NPS) Pollution." from
https://www.epa.gov/nps/basic-information-about-nonpoint-source-nps-pollution.

Finneran, K. T., C. V. Johnsen and D. R. Lovley (2003). "Rhodoferax ferrireducens sp. nov.,
a psychrotolerant, facultatively anaerobic bacterium that oxidizes acetate with the reduction
of Fe(III)." Int J Syst Evol Microbiol **53**(Pt 3): 669-673.
Gao, P., D. Mao, Y. Luo, L. Wang, B. Xu and L. Xu (2012). "Occurrence of sulfonamide and
tetracycline-resistant bacteria and resistance genes in aquaculture environment." Water
Research **46**(7): 2355-2364.
Gao, Y., W. Zhang, Y. Li, H. Wu, N. Yang and C. Hui (2021). "Dams shift microbial
community assembly and imprint nitrogen transformation along the Yangtze River." Water
Research **189**: 116579.
García-Armisen, T., Ö. İnceoğlu, N. K. Ouattara, A. Anzil, M. A. Verbanck, N. Brion and P.
Servais (2014). "Seasonal Variations and Resilience of Bacterial Communities in a Sewage
Polluted Urban River." PLOS ONE **9**(3): e92579.
Garner, E., R. Benitez, E. von Wagoner, R. Sawyer, E. Schaberg, W. C. Hession, L.-A. H.
Krometis, B. D. Badgley and A. Pruden (2017). "Stormwater loadings of antibiotic resistance
genes in an urban stream." Water Research **123**: 144-152.
Ghosal, D., S. Ghosh, T. K. Dutta and Y. Ahn (2016). "Corrigendum: Current State of
Knowledge in Microbial Degradation of Polycyclic Aromatic Hydrocarbons (PAHs): A
Review." Front Microbiol **7**: 1837.
Gibbons, S. M., E. Jones, A. Bearquiver, F. Blackwolf, W. Roundstone, N. Scott, J. Hooker,
R. Madsen, M. L. Coleman and J. A. Gilbert (2014). "Human and Environmental Impacts on
River Sediment Microbial Communities." PLOS ONE **9**(5): e97435.
Gossner, M. M., T. M. Lewinsohn, T. Kahl, F. Grassein, S. Boch, D. Prati, K. Birkhofer, S. C.
Renner, J. Sikorski, T. Wubet, H. Arndt, V. Baumgartner, S. Blaser, N. Blüthgen, C. Börschig,
F. Buscot, T. Diekötter, L. R. Jorge, K. Jung, A. C. Keyel, A. M. Klein, S. Klemmer, J. Krauss,
M. Lange, J. Müller, J. Overmann, E. Pašalić, C. Penone, D. J. Perović, O. Purschke, P.
Schall, S. A. Socher, I. Sonnemann, M. Tschapka, T. Tscharntke, M. Türke, P. C. Venter, C.
N. Weiner, M. Werner, V. Wolters, S. Wurst, C. Westphal, M. Fischer, W. W. Weisser and E.
Allan (2016). "Land-use intensification causes multitrophic homogenization of grassland
communities." Nature **540**(7632): 266-269.
Henson, M. W., J. Hanssen, G. Spooner, P. Fleming, M. Pukonen, F. Stahr and J. C. Thrash
(2018). "Nutrient dynamics and stream order influence microbial community patterns along a
2914 kilometer transect of the Mississippi River." Limnology and Oceanography **63**(5):
1837-1855.
Hou, L., L. Zhang, F. Li, S. Huang, J. Yang, C. Ma, D. Zhang, C.-P. Yu and A. Hu (2021).
"Urban ponds as hotspots of antibiotic resistome in the urban environment." Journal of
Hazardous Materials **403**: 124008.
Hubbell, S. P. (2001). The Unified Neutral Theory of Biodiversity and Biogeography,
Princeton University Press.
Hutchison, D. W. and A. R. Templeton (1999). "Correlation of pairwise genetic and
geographic distance measures: inferring the relative influences of gene flow and drift on the
distribution of genetic variability." Evolution **53**(6): 1898-1914.

Ibarbalz, F. M., N. Henry, M. C. Brandao, S. Martini, G. Busseni, H. Byrne, L. P. Coelho, H. Endo, J. M. Gasol, A. C. Gregory, F. Mahe, J. Rigonato, M. Royo-Llonch, G. Salazar, I. Sanz-Saez, E. Scalco, D. Soviadan, A. A. Zayed, A. Zingone, K. Labadie, J. Ferland, C. Marec, S. Kandels, M. Picheral, C. Dimier, J. Poulain, S. Pisarev, M. Carmichael, S. Pesant, C. Tara Oceans, M. Babin, E. Boss, D. Iudicone, O. Jaillon, S. G. Acinas, H. Ogata, E. Pelletier, L. Stemmann, M. B. Sullivan, S. Sunagawa, L. Bopp, C. de Vargas, L. Karp-Boss, P. Wincker, F. Lombard, C. Bowler and L. Zinger (2019). "Global Trends in Marine Plankton Diversity across Kingdoms of Life." Cell **179**(5): 1084-1097 e1021.

Jia, X., F. Dini-Andreote and J. Falcão Salles (2018). "Community assembly processes of the microbial rare biosphere." Trends in Microbiology **26**(9): 738-747.

Jiao, S., Y. Yang, Y. Xu, J. Zhang and Y. Lu (2020). "Balance between community assembly processes mediates species coexistence in agricultural soil microbiomes across eastern China." The ISME Journal **14**(1): 202-216.

Kansole, M. M. R. and T.-F. Lin (2016). "Microcystin-LR Biodegradation by Bacillus sp.: Reaction Rates and Possible Genes Involved in the Degradation." Water **8**(11): 508.

Kasana, R. C. and C. B. Pandey (2018). "Exiguobacterium: an overview of a versatile genus with potential in industry and agriculture." Critical Reviews in Biotechnology **38**(1): 141-156.

Kiedrzyńska, E., M. Kiedrzyński, M. Urbaniak, A. Magnuszewski, M. Skłodowski, A. Wyrwicka and M. Zalewski (2014). "Point sources of nutrient pollution in the lowland river catchment in the context of the Baltic Sea eutrophication." Ecological Engineering **70**: 337-348.

Krzmarzick, M. J., D. K. Taylor, X. Fu and A. L. McCutchan (2018). "Diversity and Niche of Archaea in Bioremediation." Archaea **2018**: 3194108.

Kuppusamy, S., P. Thavamani, K. Venkateswarlu, Y. B. Lee, R. Naidu and M. Megharaj (2017). "Remediation approaches for polycyclic aromatic hydrocarbons (PAHs) contaminated soils: Technological constraints, emerging trends and future directions." Chemosphere **168**: 944-968.

Kuypers, M. M. M., H. K. Marchant and B. Kartal (2018). "The microbial nitrogen-cycling network." Nature Reviews Microbiology **16**(5): 263-276.

Levins, R. (1968). Evolution in changing environments: some theoretical explorations, Princeton University Press.

Li, H., Q. Yang, J. Li, H. Gao, P. Li and H. Zhou (2015). "The impact of temperature on microbial diversity and AOA activity in the Tengchong Geothermal Field, China." Sci Rep **5**: 17056.

Li, H., Q. Yang and H. Zhou (2020). "Niche differentiation of sulfate-and iron-dependent anaerobic methane oxidation and methylotrophic methanogenesis in deep sea methane seeps." Front Microbiol **11**: 1409.

Li, H., H. Zhou, S. Yang and X. Dai (2023). "Stochastic and Deterministic Assembly Processes in Seamount Microbial Communities." Applied and Environmental Microbiology **0**(0): e00701-00723.

Li, X., L. Hou, M. Liu, Y. Zheng, Y. Li and X. Lin (2015). "Abundance and diversity of polycyclic aromatic hydrocarbon degradation bacteria in urban roadside soils in Shanghai." Applied Microbiology and Biotechnology **99**(8): 3639-3649.

Liu, Y., B. Su, H. Mu, Y. Zhang, L. Chen and B. Wu (2023). "Effects of point and nonpoint source pollution on urban rivers: From the perspective of pollutant composition and toxicity." Journal of Hazardous Materials **460**: 132441.

Louca, S., L. W. Parfrey and M. Doebeli (2016). "Decoupling function and taxonomy in the global ocean microbiome." Science **353**(6305): 1272-1277.

Lücker, S., M. Wagner, F. Maixner, E. Pelletier, H. Koch, B. Vacherie, T. Rattei, J. S. Damsté, E. Spieck, D. Le Paslier and H. Daims (2010). "A Nitrospira metagenome illuminates the physiology and evolution of globally important nitrite-oxidizing bacteria." Proc Natl Acad Sci U S A **107**(30): 13479-13484.

Manyi-Loh, C. E., S. N. Mamphweli, E. L. Meyer, G. Makaka, M. Simon and A. I. Okoh (2016). "An Overview of the Control of Bacterial Pathogens in Cattle Manure." International Journal of Environmental Research and Public Health **13**(9): 843.

Masuda, Y., H. Yamanaka, Z.-X. Xu, Y. Shiratori, T. Aono, S. Amachi, K. Senoo and H. Itoh (2020). "Diazotrophic anaeromyxobacter isolates from soils." Applied and Environmental Microbiology **86**(16): e00956-00920.

McKinney, M. L. (2006). "Urbanization as a major cause of biotic homogenization." Biological conservation **127**(3): 247-260.

McKinney, M. L. (2008). "Effects of urbanization on species richness: a review of plants and animals." Urban ecosystems **11**: 161-176.

Milaković, M., S. Križanović, I. Petrić, A. Šimatović, J. J. González-Plaza, M. Gužvinec, A. T. Andrašević, L. Pole, M. M. Fuka and N. Udiković-Kolić (2020). "Characterization of macrolide resistance in bacteria isolated from macrolide-polluted and unpolluted river sediments and clinical sources in Croatia." Science of The Total Environment **749**: 142357.

Mojiri, A., J. L. Zhou, A. Ohashi, N. Ozaki and T. Kindaichi (2019). "Comprehensive review of polycyclic aromatic hydrocarbons in water sources, their effects and treatments." Science of The Total Environment **696**: 133971.

Muangchinda, C., A. Yamazoe, D. Polrit, H. Thoetkiattikul, W. Mhuantong, V. Champreda and O. Pinyakong (2017). "Biodegradation of high concentrations of mixed polycyclic aromatic hydrocarbons by indigenous bacteria from a river sediment: a microcosm study and bacterial community analysis." Environmental Science and Pollution Research **24**(5): 4591-4602.

Munir, M., K. Wong and I. Xagoraraki (2011). "Release of antibiotic resistant bacteria and genes in the effluent and biosolids of five wastewater utilities in Michigan." Water Research **45**(2): 681-693.

Murthy, A. C., T. S. Fristoe and J. R. Burger (2016). "Homogenizing effects of cities on North American winter bird diversity." Ecosphere **7**(1): e01216.

Oshkin, I. Y., D. A. Beck, A. E. Lamb, V. Tchesnokova, G. Benuska, T. L. McTaggart, M. G. Kalyuzhnaya, S. N. Dedysh, M. E. Lidstrom and L. Chistoserdova (2015). "Methane-fed

microbial microcosms show differential community dynamics and pinpoint taxa involved in communal response." Isme j **9**(5): 1119-1129.

Oshkin, I. Y., C. E. Wegner, C. Luke, M. V. Glagolev, I. V. Filippov, N. V. Pimenov, W. Liesack and S. N. Dedysh (2014). "Gammaproteobacterial methanotrophs dominate cold methane seeps in floodplains of West Siberian rivers." Appl Environ Microbiol **80**(19): 5944-5954.

Oswald, K., J. S. Graf, S. Littmann, D. Tienken, A. Brand, B. Wehrli, M. Albertsen, H. Daims, M. Wagner, M. M. Kuypers, C. J. Schubert and J. Milucka (2017). "Crenothrix are major methane consumers in stratified lakes." Isme j **11**(9): 2124-2140.

Pagé, A. P., É. Yergeau and C. W. Greer (2015). "Salix purpurea Stimulates the Expression of Specific Bacterial Xenobiotic Degradation Genes in a Soil Contaminated with Hydrocarbons." PLOS ONE **10**(7): e0132062.

Pandit, S. N., J. Kolasa and K. Cottenie (2009). "Contrasts between habitat generalists and specialists: an empirical extension to the basic metacommunity framework." Ecology **90**(8): 2253-2262.

Pascual-Benito, M., E. Ballesté, T. Monleón-Getino, J. Urmeneta, A. R. Blanch, C. García-Aljaro and F. Lucena (2020). "Impact of treated sewage effluent on the bacterial community composition in an intermittent mediterranean stream." Environmental Pollution **266**: 115254.

Patel, A. B., S. Shaikh, K. R. Jain, C. Desai and D. Madamwar (2020). "Polycyclic Aromatic Hydrocarbons: Sources, Toxicity, and Remediation Approaches." Front Microbiol **11**: 562813.

Pruden, A., M. Arabi and H. N. Storteboom (2012). "Correlation Between Upstream Human Activities and Riverine Antibiotic Resistance Genes." Environmental Science & Technology **46**(21): 11541-11549.

Qian, X., B. Liang, X. Liu, X. Liu, J. Wang, F. Liu and B. Cui (2017). "Distribution, sources, and ecological risk assessment of polycyclic aromatic hydrocarbons in surface sediments from the Haihe River, a typical polluted urban river in Northern China." Environ Sci Pollut Res Int **24**(20): 17153-17165.

Reddy, S., K. Kaur, P. Barathe, V. Shriram, M. Govarthanan and V. Kumar (2022). "Antimicrobial resistance in urban river ecosystems." Microbiological Research **263**: 127135.

Rizzo, L., C. Manaia, C. Merlin, T. Schwartz, C. Dagot, M. C. Ploy, I. Michael and D. Fatta-Kassinos (2013). "Urban wastewater treatment plants as hotspots for antibiotic resistant bacteria and genes spread into the environment: A review." Science of The Total Environment **447**: 345-360.

Rodrigues, J. L., V. H. Pellizari, R. Mueller, K. Baek, C. Jesus Eda, F. S. Paula, B. Mirza, G. S. Hamaoui, Jr., S. M. Tsai, B. Feigl, J. M. Tiedje, B. J. Bohannan and K. Nüsslein (2013). "Conversion of the Amazon rainforest to agriculture results in biotic homogenization of soil bacterial communities." Proc Natl Acad Sci U S A **110**(3): 988-993.

Ruff, S. E., J. F. Biddle, A. P. Teske, K. Knittel, A. Boetius and A. Ramette (2015). "Global dispersion and local diversification of the methane seep microbiome." Proc Natl Acad Sci U S A **112**(13): 4015-4020.

Salazar, G. (2020). "EcolUtils: Utilities for community ecology analysis." <u>R package</u>.

Singh, R., A. P. Singh, S. Kumar, B. S. Giri and K.-H. Kim (2019). "Antibiotic resistance in
major rivers in the world: A systematic review on occurrence, emergence, and management
strategies." <u>Journal of Cleaner Production</u> **234**: 1484-1505.

Sköld, O. (2000). "Sulfonamide resistance: mechanisms and trends." <u>Drug resistance
updates</u> **3**(3): 155-160.

Sloan, W. T., M. Lunn, S. Woodcock, I. M. Head, S. Nee and T. P. Curtis (2006).
"Quantifying the roles of immigration and chance in shaping prokaryote community
structure." <u>Environ Microbiol</u> **8**(4): 732-740.

Stegen, J. C., X. Lin, A. E. Konopka and J. K. Fredrickson (2012). "Stochastic and
deterministic assembly processes in subsurface microbial communities." <u>ISME J</u> **6**(9):
1653-1664.

Van Boeckel, T. P., S. Gandra, A. Ashok, Q. Caudron, B. T. Grenfell, S. A. Levin and R.
Laxminarayan (2014). "Global antibiotic consumption 2000 to 2010: an analysis of national
pharmaceutical sales data." <u>The Lancet infectious diseases</u> **14**(8): 742-750.

VandeWalle, J. L., G. W. Goetz, S. M. Huse, H. G. Morrison, M. L. Sogin, R. G. Hoffmann, K.
Yan and S. L. McLellan (2012). "Acinetobacter, Aeromonas and Trichococcus populations
dominate the microbial community within urban sewer infrastructure." <u>Environmental
Microbiology</u> **14**(9): 2538-2552.

Wang, L., J. Zhang, H. Li, H. Yang, C. Peng, Z. Peng and L. Lu (2018). "Shift in the microbial
community composition of surface water and sediment along an urban river." <u>Science of The
Total Environment</u> **627**: 600-612.

Wang, Y., H. F. Sheng, Y. He, J. Y. Wu, Y. X. Jiang, N. F. Tam and H. W. Zhou (2012).
"Comparison of the levels of bacterial diversity in freshwater, intertidal wetland, and marine
sediments by using millions of illumina tags." <u>Appl Environ Microbiol</u> **78**(23): 8264-8271.

Wéry, N., C. Lhoutellier, F. Ducray, J.-P. Delgenès and J.-J. Godon (2008). "Behaviour of
pathogenic and indicator bacteria during urban wastewater treatment and sludge
composting, as revealed by quantitative PCR." <u>Water Research</u> **42**(1): 53-62.

WHO (2019). "Critically important antimicrobials for human medicine." <u>6th Revision 2018:
World Health Organization</u>.

Widder, S., K. Besemer, G. A. Singer, S. Ceola, E. Bertuzzo, C. Quince, W. T. Sloan, A.
Rinaldo and T. J. Battin (2014). "Fluvial network organization imprints on microbial
co-occurrence networks." <u>Proceedings of the National Academy of Sciences</u> **111**(35):
12799-12804.

Williams, T. J., D. Wilkins, E. Long, F. Evans, M. Z. DeMaere, M. J. Raftery and R.
Cavicchioli (2013). "The role of planktonic Flavobacteria in processing algal organic matter
in coastal East Antarctica revealed using metagenomics and metaproteomics."
<u>Environmental Microbiology</u> **15**(5): 1302-1317.

Xiao, H., X. Zhang, M. Yan, L. Zhang, H. Wang, Y. Ma and J. Liu (2022). "The
Temporal-Based Forest Disturbance Monitoring Analysis: A Case Study of Nature Reserves
of Hainan Island of China From 1987 to 2020." <u>Frontiers in Environmental Science</u> **10**.

Xie, Y., J. Wang, Y. Wu, C. Ren, C. Song, J. Yang, H. Yu, J. P. Giesy and X. Zhang (2016).
"Using in situ bacterial communities to monitor contaminants in river sediments."
Environmental Pollution **212**: 348-357.
Xu, N., H. Hu, Y. Wang, Z. Zhang, Q. Zhang, M. Ke, T. Lu, J. Penuelas and H. Qian (2023).
"Geographic patterns of microbial traits of river basins in China." Science of The Total
Environment **871**: 162070.
Xuan, Y., Y. Mai, Y. Xu, J. Zheng, Z. He, L. Shu and Y. Cao (2022). "Enhanced microbial
nitrification-denitrification processes in a subtropical metropolitan river network." Water
Research **222**: 118857.
Yang, S., Q. Chen, T. Zheng, Y. Chen, X. Zhao, Y. He, W. Sun, S. Zhong, Z. Li and J. Wang
(2022). "Multiple metal(loid) contamination reshaped the structure and function of soil
archaeal community." Journal of Hazardous Materials **436**: 129186.
Zhang, L., M. Zhong, X. Li, W. Lu and J. Li (2020). "River bacterial community structure and
co-occurrence patterns under the influence of different domestic sewage types." Journal of
Environmental Management **266**: 110590.
Zhang, Q.-Q., G.-G. Ying, C.-G. Pan, Y.-S. Liu and J.-L. Zhao (2015). "Comprehensive
Evaluation of Antibiotics Emission and Fate in the River Basins of China: Source Analysis,
Multimedia Modeling, and Linkage to Bacterial Resistance." Environmental Science &
Technology **49**(11): 6772-6782.
Zhang, Y., C. F. Marrs, C. Simon and C. Xi (2009). "Wastewater treatment contributes to
selective increase of antibiotic resistance among Acinetobacter spp." Science of The Total
Environment **407**(12): 3702-3706.
Zhang, Y., Z. Zhao, M. Dai, N. Jiao and G. J. Herndl (2014). "Drivers shaping the diversity
and biogeography of total and active bacterial communities in the South China Sea." Mol
Ecol **23**(9): 2260-2274.
Zhi-Hui, Y., K. Stöven, S. Haneklaus, B. Singh and E. Schnug (2010). "Elemental sulfur
oxidation by Thiobacillus spp. and aerobic heterotrophic sulfur-oxidizing bacteria."
Pedosphere **20**(1): 71-79.
899
900

901 **Figure**

902
903

904 **Figure 1.** Location of the collected river network sediment samples. Orange circles indicate
905 the urban river sites, and blue circles indicate the nature reserve river sites, respectively.
906 The Inset map shows the extent of the study area relative to China.

907
Figure 2. Comparison of geochemical characterization in human-impacted urban rivers and
undisturbed natural rivers in the river network. The orange circles indicate urban rivers and
blue circles indicate natural rivers, respectively.

Figure 3. The microbiome in the large river network. (A-D) Microbial abundance, Richness

estimates, Shannon index and Niches breadth were investigated in urban (orange circles)

and natural (blue circles) river samples. (E) Venn analysis. (F) Similarity of microbial

communities visualized by two-dimensional (2D) nonmetric multidi-mensional scaling

(NMDS). (F) A distance-based redundancy analysis (db-RDA) showing microbial community

similarity when environmental factors were used as constraining variables and were fitted

onto the ordination as arrows. The length of each arrow indicates the multiple partial

correlation of the variable to RDA axes and can be interpreted as an indication of that

variable's contribution to the explained community similarity. (I) Prediction of the functional

diversity and metabolic pathways between urban and natural river samples.

Figure 4. Relative sequence abundance of bacterial and archaeal. The graph shows the
relative abundance of the most abundant bacterial and archaeal at phylum (A) and classes
(B) level in urban river and natural river according to average relative abundance across all
samples.

Figure 5. Heat map with taxonomic information on the dominant species in all samples.

933
934 **Figure 6.** Antibiotic resistance rates and gene abundance. The orange circles indicate the
935 urban river and blue circles indicate the natural river, respectively. While no significant
936 differences were observed among the natural river, the antibiotic-resistant rates exhibited
937 variability among the samples of the urban river.

Figure 7. General patterns of microbial diversity in the river network system. (A)
Distance-decay curves showing Bray–Curtis dissimilarity against geographic distances
between sampling sites. Solid lines denote the ordinary least-squares linear regressions. (B)
Scatter plot of βNTI values grouped by spatial scales of the river network. (C) Fit of the
neutral community model (NCM) of community assembly. The solid blue lines indicate the
best fit to the NCM as in Sloan et al. (2006), and the dashed blue lines represent 95%
confidence intervals around the model prediction. OTUs that occur more or less frequently
than predicted by the NCM are shown in different colors. Nm indicates the metacommunity
size times immigration, R^2 indicates the fit to this model.