**Urban non-point source pollutants cause microbial community**

**homogenization via increasing deterministic processes**

Haizhou Li[1, *], Xiangyu Fan[2], Zhiwei He[3], Jing Fu[1], Yuekai Wang[1], Jinchen Yu[1], Shanshan

Yang[4], Jiawang Wu[5], Li Wu[6], Jin Zhou[1, *].

Email: lihz.bio@gmail.com, zhou_jin@foxmail.com

1. East China Sea Fisheries Research Institute, Chinese Academy of Fishery Science,
Shanghai, China.
2. Department of Molecular and Cellular Biology, University of California, Davis, United States.
3. State Key Laboratory of Marine Geology, Tongji University, Shanghai, China.
4. College of Marine Science and Technology, China University of Geosciences, Wuhan, Hubei,
China
5. School of Marine Sciences, Sun Yat-Sen University, Guangdong, China.
6. Laboratory for Coastal Ocean Variation and Disaster Prediction, College of Ocean and
Meteorology, Guangdong Ocean University, Guangdong, China.

**Keywords: river network, non-point source pollution, urbanization, biotic**
**homogenization, microbiomes**

## Abstract

Urbanization significantly impacts the community structure of aquatic organisms and poses a

major threat to river biodiversity. However, the extent to which urbanization is linked to the

homogenization of microbial communities and the underlying mechanisms remains poorly

understood. In this study, we investigated the bacterial and archaeal communities from cities

and neighboring natural rivers across river network located in the Qinling Mountains, Northwest

China, and further investigated the alpha and beta diversity patterns and the mechanisms

influenced by urbanization. We found that the influx of urban non-point source pollutants

created a eutrophic condition, and enhanced the urban river microbial populations. Meanwhile,

the rapid urbanization tends to decrease the overall habitat heterogeneity, and imposed

stronger homogeneous selection and caused microbial communities biotic homogenization.

The mechanisms of biotic homogenization can be attributed to modulating generalist/specialist species and invasion of nonnative species. For instance, the urban river had a greater proportion of fast-growing bacteria, algae, nitrifiers, PAH-degrading bacteria, pathogens, fecal bacteria and antibiotic-resistant bacteria than natural river ecosystems. Overall, urbanization leads to a more uniform river biosphere, causing the extinction of unique local species and a subsequent decrease in the regional species pool.

**1. Introduction**

Rivers have been a preferred location for human settlements throughout history (1). However, in recent decades, human activities such as rapid urbanization and industrialization have discharged multiple anthropogenic contaminants and non-native (exotic) microorganisms into the river (2, 3). These contaminants can be either organic (e.g., antibiotics, polycyclic aromatic hydrocarbons, fertilizers, pesticides, manure) (4-7) or inorganic contaminants (e.g., nitrate, phosphorus, heavy metal) (8-10). According to the United Nations, 68% of the global population will live in cities by 2050 (11). Accelerated urbanization and intense human activities substantially influence the community composition and function of the river microbiome, and posing a considerable risk to riverine ecosystems. For example, centralized wastewater treatment plants (WWTPs) represent an important and common urban river pollution source. Evidences showed that effluents from WWTPs exert adverse effects on the functionality of river ecosystems, often leading to the deterioration of water quality, habitat degradation, and a reduction in the richness and diversity of benthic and sedimentary microbial communities (12-15).

WWTPs are a common source of point-source pollution in the river ecosystem and have become a global aquatic environmental issue. However, the immense industrial and domestic sewage are considered as the primary contributors to non-point source pollution (2, 16). Point-source pollution is easy to identify, but non-point source pollution is harder to identify and harder to address. The United States Environmental Protection Agency indicates non-point source pollution remains the primary unresolved issue affecting river water quality, yet there are little data on the impacts of widespread non-point source pollution on microbial communities in river ecosystems (17, 18). The potential ecological and health influences of non-point source pollution in river ecosystems require further research and monitoring.

Here, we focus on a large-scale river network, which flows out of the Qinling Mountains, where

it eventually empties into the Yangtze River (Fig. 1). The Qinling Mountains, located in

Northwest China, serve as a geographically important north-south boundary and natural barrier

for the eastward flow of moisture from the Tibetan Plateau, providing a significant source of

freshwater for downstream human communities. The distinctive feature of the river network is

that it passes numerous cities (a current population of approximately 11.06 million) (19) and

national nature reserves. Establishing nature reserves is an effective measure to avoid human

activity disturbances (20, 21). This river network provides a useful model for understanding the

impact of non-point source pollution on the river ecosystem.

In this study, the collected river surface sediment samples (0–2 cm deep) were in eutrophic

waterbodies (highly urbanized rivers) and oligotrophic or mesotrophic waterbodies (undisturbed

nature reserve rivers), based on trophic state index (TSI) (Fig. 1, Dataset 1B). River sediments

are considered as the major habitat of microbes in the aquatic environment (22, 23). Therefore,

sediment-associated microorganisms drive essential ecological processes and contribute

significantly to river function (24). Here, these collected river surface sediments represent

diverse non-point source pollution gradients, and creating multiple niches in this river

network. We obtained millions of 16S rRNA tags sequences from these sediments by

high-throughput sequencing, aiming to (i) compare the community composition, diversity and

function of bacteria and archaea in urban river with natural river ecosystems; (ii) evaluate the

environmental factors linked to the river microbiome; (iii) identify the river microbial taxa

(bacteria, archaea) community assembly processes; and (iv) assess the microbial functional

attributes characterizing in river ecosystems associated with urbanization, eutrophication,

antibiotic resistance, fecal contamination, pathogenesis, nutrient cycling, and abiotic stress.

**2. Materials and methods**

**2.1 Study area, sampling procedure, and physicochemical analysis**

Sampling was conducted in August-September 2022. From each sampling site on three

sampling occasions, water and sediment samples were collected within approximately 2 square

meters at each site. The geographical data (longitude, latitude, and distance between sampling

sites) were recorded by a GPS device (Dataset 1A). The water quality parameters, including

water temperature, pH, dissolved oxygen (DO), water clarity of Secchi depth, transparency

(SD), turbidity (TUB), total nitrogen (TN), nitrate-nitrogen ($NO_3^-$), ammonium-nitrogen ($NH_4^+$),

total phosphors (TP), total dissolved phosphorus (TDP), phosphate, permanganate index,

volatile phenol, *Chlorophyll*-a concentration were measured according to Environmental Quality

Standards for Surface Water (Ministry of Ecology and Environment of People's Republic of

China, GB3838-2002). Sediment total organic carbon (TOC) was measured by the multi N/C

3100 (Analytik Jena). Each sample was performed in triplicate, and the values were expressed

as the mean (n = 3). To evaluate the trophic status of each site, we calculated a trophic state

index (TSI) based on the concentrations of TN, TP, Chl-a and SD in the water environment (25,

26). Five trophic levels were defined: oligotrophic (TSI < 30), mesotrophic (30 ≤ TSI ≤50),

slightly eutrophic (50 < TSI ≤ 60), moderately eutrophic (60 < TSI ≤70) and highly eutrophic

(TSI >70) (25). For detailed information on the environmental variables of the trophic groups,

see Dataset 1B.

**2.2 Sediment cell counts**

Catalyzed reporter deposition-fluorescent *in situ* hybridization (CARD-FISH) was used to

quantify the cell densities based on our previous studies (27). The probe-labeling peroxidases

were EUB338 (bacteria, 5'-GCW GCC WCC CGT AGG WGT-3') and Arch915 (archaea,

5'-GTG CTC CCC CGC CAA TTC CT-3'). Probe NON338 (5'-ACT CCT ACG GGA GGC

AGC-3') was used as a control. Frozen samples (0.5 g) were fixed with 4% paraformaldehyde

for 24 h at room temperature. The fixed sediments were washed three times by centrifugation

(8,000 × *g* for 10 min) using phosphate buffered saline (PBS) at 4 °C and stored in an

ethanol/PBS buffer (1:1) at -20 °C for further processing. Later, 100 µL of fixed sediment was

diluted with 900 µL ethanol/PBS buffer (1:1) and dispersed using ultrasound. Then, 20 µL of the

dispersed sediment was further diluted in 20 mL of Milli-Q filtered water. The suspended

sediments were filtered onto polycarbonate filters, and 0.1% low melting point agarose was

dripped onto the filters and dried at 46 °C in an incubator. The microbes were permeabilized

using 15 µg/mL proteinase K. Then, 3% $H_2O_2$ was used to inactivate the endogenous

peroxidases.

For hybridization, filters were placed in a tube and mixed with 500 µL hybridization solution (10%

dextran sulfate, 2% blocking reagent (Roche, Germany), 0.1% (w/v) sodium dodecyl sulfate, 20

mM Tris–HCl [pH 8.0], 0.9 M NaCl and formamide) and 1 µL of probe working solution (final

concentration, 0.028 µM) (28). Microorganisms were hybridized for at least 60 min on a rotor at

46 °C, and then the filters were washed twice using washing solution (28) (0.01% SDS, 5 mM

EDTA [pH 8.0], 20 mM Tris–HCl [pH 8.0] and 3 mM NaCl) at 48 °C for 20 min. After washing,

filters were mixed with 1,000 µL of amplification solution (0.0015% $H_2O_2$, 1×PBS [pH 7.4], 0.1%

(w/v) blocking reagent) and 1 µL of Alexa 488-labelled tyramides (Life Technologies[TM], Thermo

Fisher, USA). The probes were incubated at 46 °C in amplification solution for at least 30 min in

the dark.

For second hybridizations, the first probe-labelling peroxidase was inactivated by incubating the

filter sections in 0.01 M HCl for 10 min at room temperature and washing the sections with 50

mL of Milli-Q water. Then, the CARD-FISH protocol was repeated two times with the same filter

sections by using different probes. The second hybridization was performed using Alexa

647-labelled tyramides (Life Technologies, Thermo Fisher, USA). Finally, all microorganisms

were stained using DAPI and mounted with ProLong Gold Antifade reagent (Life Technologies,

Carlsbad, California, USA). Cell counting was performed using ImageJ.

**2.3 High-throughput sequencing**

Frozen sample was aseptically subsampled for DNA extraction. In addition, a negative control

(5 mL of DNase water) was included with each batch of subsamples and subjected to DNA

extraction. DNA was extracted from each subsample using the E.Z.N.A.® soil DNA kit (Omega

Bio-tek, Norcross, Georgia, USA), according to the manufacturer's instructions.  The final DNA

concentration was determined by a NanoDrop 2000 (Thermo Scientific, Waltham, USA), and

DNA quality was checked by 1% agarose gel electrophoresis. The sequencing steps were

conducted by Majorbio Bio-Pharm Technology Co., Ltd. (Shanghai, China). The

bacterial/archaeal universal primers were 515F and 806R (29, 30), targeting the 16S rRNA

gene V4 region. The polymerase chain reaction (PCR) protocol was performed according to

previously described methods (31). The PCR products were extracted from a 2% agarose gel

and further purified using an AxyPrep DNA Gel Extraction Kit (Axygen Biosciences, USA) and

quantified using Quanti Fluor™-ST (Promega, USA). Purified amplicons were pooled in

equimolar amounts and paired-end sequenced (2□×□300) on an Illumina MiSeq platform

(Illumina, San Diego, California, USA). Raw fastq files were demultiplexed, quality filtered by

Trimmomatic and merged by FLASH. OTUs were clustered with a 97% similarity cut-off using

UPARSE, and chimeric sequences were identified and removed using UCHIME. To remove the

effects of sampling intensity (number of reads) on comparisons of richness estimates, we

randomly subsampled a fixed number of reads (n= 10, 000) for each sample. Taxonomic

assignment was performed using the SILVA 16S rRNA gene database (version 138). The

distance-based maximum likelihood was used for phylogenetic analysis. Bootstrap analysis

was performed using 1,000 replications. The Mega software had been used for computing

phylogenetic trees. Alpha-diversity metrics (observed OTU richness, Chao1 richness, Shannon

and Simpson diversity indices, and equitability index) were calculated using Mothur (32).

Here, the "niche breadth" approach (33) was used to quantify habitat specialization. The

formula is described below:

$$B_j = \frac{1}{\sum_{i=1}^{N} P_{ij}^2}$$

where $B_j$ is the habitat niche breadth of OTUj in a metacommunity; and $P_{ij}$ is the proportion of

OTUj in a given site i, and N is the total number of sites. A higher B value of OTUs indicates a

wider habitat niche breadth. In contrast, lower *B* values indicate that the OTUs occurred in

fewer habitats and were unevenly distributed (34). The average of B values from all OTUs in

one community was regarded as the community-level B-value (*Bcom*) (35). We calculated

the *B* value based on species abundance and evenness for NHF and SCS sediment samples

(36).

Furthermore, in order to identify microbial ecological functions at the studied sites, the obtained

OTUs were compared with the PICRUSt2 (37, 38) and FAPROTAX database (39, 40) to predict

metabolic functions of the microbial community.

**2.4 Screening for antibiotic resistant bacteria from river sediments**

The approach that has been used to evaluate the status of antibiotic resistance in the

environment involves the calculation of resistance percentage, which is based on the ratio

between the number of bacteria that are able to grow on culture media that are supplemented

with antibiotics at doses close to the minimal inhibitory concentration (MIC) and the number of

bacteria that grow on antibiotic-free media (41, 42). The approach of heterotrophic plate count

(HPC) method (43) was used to determine the concentration of antibiotic resistant bacteria in

the collected river surface sediments. 0.1 g sediment samples were dissolved in 10 mL of 0.85%

sterilized normal saline, and diluted 10 × fold for immediate processing. 100 μL diluted samples

were cultured onto plate count agar medium. Medium contained four-class plates, 50.4 μg/mL

sulfamethoxazole, 16 μg/mL tetracycline, 15□μg/mL azithromycin and no antibiotics. The

media was added antifungal agent 200 µg/mL cycloheximide according to Gao el al., (2012)

and Brooks et al. (2007) method. Total heterotrophic cultivable bacterial population was

determined by plate count agar medium without antibiotics. The selected concentrations of

antibiotics on the media were determined according to the criteria of the Clinical and Laboratory

Standards Institute (CLSI) guidelines (2015) and the previous study (43-45). Plates were

incubated for 48–72 h at 37 °C (43, 46). Duplicate counting was performed, and colony-forming

units (CFU) were calculated.

**2.5 Real-time quantitative PCR (qPCR)**

Absolute quantification standard curve method was used to quantify the antibiotic resistant

genes in the river sediment samples based on our previous studies (47). The real-time

quantitative PCR (q-PCR) (ABI7500, USA) using SYBR 2×RealUniversal PreMix (Tiangen

Biotech). qPCR was performed to assess the abundance of the macrolide, tetracycline and

sulfonamide antibiotic resistant genes (ARGs). The sequences of the primers and the reaction

conditions used in this study are presented in Table S1 in the Supporting Information (SI)

section.

The PCR reactions were performed in a 20-µL reaction mixture that comprised 10 µL of SYBR

RealUniversal PreMix (2 ×), 0.4 µL of each type of primer (10/20 µM), and 1 µL of a DNA

template (5 ng). The thermocycling protocol consisted of an initial denaturation for 1 min at

95 °C, followed by 40 cycles of denaturation (95 °C, 10 s) and annealing at the temperatures in

Tables S1. The specificity of the PCR products was checked by a melting curve analysis. All

qPCR assays were performed in triplicate for each gene,

The q-PCR standards were synthesized by cloning target genes into *Escherichia coli* DH5a

using a PMD18-T vector Cloning Kit (TaKaRa Biotechnology, China). Vector concentrations

were measured using NanoDrop, and the quantities of target genes per µL of solution were

obtained basing on the length of plasmid and target gene sequence. Six-point calibration curves

were generated by 10-fold serial dilutions of the plasmid carrying ARGs. All PCR reactions were

run in parallel with serially diluted q-PCR standards and DNA-free water as the negative

controls.

**2.6 Statistical analysis**

Null model analysis was carried out using the framework described by Stegen et al. (2012, 2013)
to classify community pairs into underlying drivers of species sorting (or selection), dispersal
limitation, homogeneous dispersal, and drift. The variation of both phylogenetic diversity and
taxonomic diversity was measured using null model-based phylogenetic and
taxonomic $\beta$-diversity metrics, namely $\beta$-nearest taxon index ($\beta$NTI) and Bray–Curtis-based
Raup–Crick ($RC_{Bray}$). A significant deviation (i.e., $|\beta NTI| > 2$) indicates the dominance of
selection processes. $\beta NTI < -2$ indicates significantly less phylogenetic turnover than expected
(i.e., homogeneous selection) while $\beta NTI > +2$ indicates significantly more phylogenetic
turnover than expected (i.e., variable selection). Subsequently, $RC_{Bray}$ was used to further
partition the pairwise comparisons that were not assigned to selection (i.e., $|\beta NTI| < 2$). The
relative influence of homogenizing dispersal was quantified as the fraction of pairwise
comparisons with $|\beta NTI| < 2$ and $RC_{Bray} < -0.95$. Dispersal limitation was quantified as the
fraction of pairwise comparisons with $|\beta NTI| < 2$ and $RC_{Bray} > 0.95$. The fractions of all pairwise
comparisons with $|\beta NTI| < 2$ and $|RC_{Bray}| < 0.95$ were used to estimate influence of
"undominated" assembly, which mostly consists of weak selection, weak dispersal,
diversification, and/or drift.
239
A neutral community model was used to determine the contribution of stochastic processes to
microbial community assembly by predicting the relationship between the frequency with which
taxa occur in a set of local communities and their abundance across the wider metacommunity
(48). The model predicts that abundant taxa are more likely to be dispersed by chance and
widespread across metacommunity, while rare taxa would be lost in different local communities
due to ecological drift. In the model, the estimated migration rate is a parameter for evaluating
the probability that a random loss of an individual in a local community would be replaced by
dispersal from the metacommunity, and, therefore, is a measure of dispersal limitation.
Higher $m$ values indicate that microbial communities are less dispersal limited. The formula is
as follows:

$$Freq_i = 1 - I(1/N \mid N*m*p_i, N*m*(1 - p_i))$$

where $Freq_i$ is the occurrence frequency of taxon $i$ across communities; $N$ is the number of
individuals per community; $m$ is the estimated migration rate; $p_i$ is the average relative
abundance of taxon $i$ across communities; and $I()$ is the probability density function of beta
distribution. $R^2$ indicates the fit of the parameter based on nonlinear least-squares fitting.

**2.7 Data available**

Sequencing data are stored on the National Genomics Data Center (NGDC) Sequence Read
Archive (PRJCA018972).

**3. Results**

**3.1 Environmental heterogeneity between urban and natural river areas**

A total of 28 urban river and 24 undisturbed pristine natural river surface (0-2 cm) sediment
samples were collected (Fig. 1). Samples from all 52 locations covered a wide range of
horizontal distance (0.72-376 km). The environmental chemical profiles of river water showed
remarkable differences between the urban and natural rivers (Fig. 1, Dataset 1A). The
concentration of total nitrogen (TN), nitrite, nitrate, ammonium, total phosphorus (TP), total
dissolved phosphorus (TDP), phosphate, volatile phenol, Chlorophyll *a* permanganate index, in
urban river were obviously higher than natural river (P<0.01). The total organic carbon (TOC)
content in sediment also showed a significant increasing trend from the natural river to urban
river (*P*<0.0001) (Fig. 1). Here, the permanganate index is often used as a comprehensive
indicator of the degree of water pollution caused by reducible organic and inorganic substances
(49).

The sampling locations along the urban river exhibited an average Trophic State Index (TSI) of
53.07, with a range from 41.97 to 69.32 (Dataset 1B), categorizing most of these sites as
eutrophic based on a TSI threshold exceeding 50 (50). Conversely, the sampling sites within
the natural river had an average TSI of 36.23, with range from 20.84 to 53.95 (Dataset 1B),
positioning most sites within the oligotrophic and mesotrophic range (TSI < 50). These
differences indicate substantial environmental heterogeneity between the two habitats.

**3.2 Microbial community abundance, richness, diversity**

Microbial abundance ranged from $1.3 \times 10^7$ to $1.1 \times 10^9$ cells g-1 and $8.0 \times 10^6$ to $7.1 \times 10^8$ cells
g-1 for urban and natural river samples, respectively (Fig. 2A, details in Dataset 2A), and urban
river sites exhibited a significantly higher abundance of microorganisms than those from the
natural river samples (*P*<0.01). The abundance of microorganisms increased with increasing
TSI, indicating that the urban river eutrophication and has a great impact on microbial
population (Fig. 2E).

Using universal primers for bacteria and archaea, 3,309,566 high-quality sequences were

generated throughout the river network (Dataset 2A, Supporting Information Fig. S1). These

high-quality sequences were clustered into 18,038 operational taxonomic units (OTUs, 97%

similarity level) (Dataset 2B). A range of 1,628–6,104 chao1 OTUs and 1.89–6.68 Shannon

index were observed in the urban samples, and a range of 1,321–5,471 chao1 OTUs and

1.88–6.31 Shannon index were observed in natural river samples (Fig. 2B-C). The richness and

diversity were considerably higher in the urban river compared to the natural river samples ($P <$

0.05). The alpha diversity showed predictable patterns along the TSI gradient, and both the

Chao 1 and Shannon increased with increasing TSI (Fig. 2F-G). The natural river communities

exhibit a wider niche breadth (Fig. 2D), which is indicative of greater metabolic flexibility.

74 microbial phyla and 3,599 species were identified in urban samples, and 80 phyla and 3,736

species were identified in natural samples (Dataset 2C-F). At the phylum level, the urban and

natural river samples investigated here shared a high proportion of taxa (Fig. 3A). The ten most

prevalent phyla in all of sites collectively constituted 95.48% of the total sequences, matching

the classical abundance-range relationship (51, 52). Furthermore, an increase in the diversity

and proportion of archaea was also identified under the impact of human activities pollution

effluent. In the urban river areas, proportion of archaea exhibited 1.58 times higher than those

in natural river, and archaeal Chao1 richness and Shannon diversity exhibited 1.47 and 2.36

times higher than that in natural river, respectively.

Although the microbial community of these sites exhibited similarities at the phylum level, 38

orders in samples showed significant differences ($P<0.01$) between urban and natural river

ecosystems (Supporting Information Fig. S2). For example, Bacillales, Clostridiales,

Steroidobacterales were dominant in urban areas, while Erysipelotrichales, Deinococcales,

Alicyclobacillales, Pyrinomonadales, Burkholderiales were dominant in the natural river

(Dataset 2C, Supporting Information Fig. S2). Further, a heatmap was generated to represent

the 50 most prevalent species (Supporting Information Fig. S3). In urban rivers, the dominant

taxa included Unclassified Planococcaceae, Unclassified Steroidobacteraceae, Unclassified

Anaerolineaceae, and Unclassified Rhizobiales. On the other hand, the unclassified

*Acinetobacter*, unclassified Pseudomonas, unclassified Sphingomonadaceae, unclassified

*Deinococcus*, *Pseudomonas parafulva* had high abundances in natural rivers (Dataset 2E). At

OTU level, the NMDS and network analysis showed that the urban samples harbored

significantly dissimilar microbial communities with natural river samples (Fig. 2I-J;

PERMANOVA pseudo F = 2.1455, p.adj = 0.024, 999 permutations; ANOSIM R sqr = 0.0597, P

= 0.03). Along the trophic gradient, we observed taxonomic homogenization for bacteria and

archaea. Such a phenomenon can be supported by the average Bray-Curtis dissimilarity for

each trophic group, which showed lower values in highly urbanization river than in natural rivers

with lower trophic status.

**3.3 The key functional microbial communities**

Microbial community metabolic predictions guided by the FAPROTAX database (39) revealed

distinctions in the metabolic potential related to nitrogen, carbon, and sulfur cycling between

urban and natural rivers (Fig. 4, Dataset 3A-D). For nitrogen transformation, the

microorganisms involved in ammonia oxidation, nitrite oxidation, nitrogen fixation, nitrification in

urban rivers were higher than those in the natural river (ANOVA, $P < 0.001$). High-throughput

sequencing showed a notably higher abundance of nitrogen transformation taxa groups

(Dataset 2E-F, $P < 0.01$) was observed in urbanized rivers, including nitrogen-fixing bacteria

(*Anaeromyxobacte*) (53), nitrifying bacteria (*Nitrosomonas, Nitrospina, Nitrosospira*, *Nitrospira*)

and denitrifying bacteria (*Paracoccus, Hyphomicrobium*, and *Gemmobacter*) (54). The db-RDA

analysis indicated that nitrogen (often in the form of nitrate and ammonium) contributed

significantly to the changes in the urban river microbial community (Fig. 2I).

In all samples, biodegradation pathways encompassing 21 types of xenobiotic compounds

were determined (Dataset 3F-G). The ecological function of xenobiotic biodegradation exhibited

a noticeable increasing trend from the natural river to the human-impacted sites. It is

understood that the detection of biodegradation genes can act as markers for the presence of

xenobiotics and their related metabolites (55, 56). In the urban river locations, a significant

presence of aminobenzoate, benzoate, and naphthalene was observed (Dataset 3G),

coinciding with a rise in the abundance of bacteria capable of degrading PAHs, such as

*Sphingobium*, *Rhodococcus*, *Novosphingobium*, *Pseudomonas*, *Sphingomonas*,

*Brevibacterium*, *Arthrobacter*, *Nocardioides,* and *Mycobacterium* were observed in the urban

river (Dataset 2E, $P < 0.01$) (57). PAHs primarily originating from long-term anthropogenic

pollution sources linked to industrialization and urbanization (58, 59). These PAHs pollutants,

characterized by persistence and low solubility, tend to accumulate in river sediments due to

their hydrophobic nature and strong adsorption onto sediment particles (5, 58). It suggested the

accumulation of PAHs in our collected urban river sediment samples.

Function genes linked to drug and antimicrobial metabolism were observed higher in urban river

sediment samples (Dataset 3D). Therefore, three kinds of antibiotic, macrolides, tetracycline

and sulfonamide, were selected for antibiotic resistance rates (ARRs) test in our study (Fig. 5A).

Tetracyclines and macrolides are the most commonly used antibiotics around the world for both

medical and agricultural purposes (60, 61). Sulfonamides, the first antibiotics to be synthesized

for widespread use, have experienced a significant rise in resistance among clinical samples

(62). The ARRs in urban rivers ranged 11.2% - 29.5 % for azithromycin, 13.0-29.0 % for

tetracycline, 11.0-26.9 % for sulfonamide. The ARRs in natural rivers ranged 2.4-18.2% for

azithromycin, 4.0-16.0 % for tetracycline, 2.4-16.0 % for sulfonamide. The antibiotic resistant

rate in urban samples were higher than those in natural river samples (Fig. 5A). Five antibiotic

resistance genes (ARGs) spanning three categories of resistance mechanisms—specifically,

two for sulfonamides (*sul1* and *sul2*), two for tetracyclines (*tetO* and *tetW*), and one for

macrolides (*ermF*) (Fig. 5B)—were chosen for analysis in all specimens to delineate the

variance in ARG concentrations between urban and natural river settings. The ARGs were

found in all samples, and Fig. 5B shows the occurrences and quantities of ARGs in the

sediments of the urban and natural rivers. The concentrations of ARGs varied greatly, ranging

from $1.04 \times 10^3$ copies/g (*sul2*, Natural River) to $4.10 \times 10^8$ copies/g (*tetO*, urban river). The

quantity of ARGs in the urban river ranged from $1.36 \times 10^4$ to $1.63 \times 10^8$ copies/g for *sul1*,

$2.27 \times 10^4$ to $3.00 \times 10^8$ copies/g for *sul2*, $5.36 \times 10^3$ to $4.10 \times 10^8$ copies/g for *tetO,* 2.52

$\times 10^3$ to $2.17 \times 10^8$ copies/g for *tetW,* $1.72 \times 10^3$ to $2.02 \times 10^8$ copies/g for *ermF*. The quantity of

ARGs in the natural river ranged from $1.42 \times 10^3$ to $9.49 \times 10^6$ copies/g for *sul1*, $1.04 \times 10^3$ to

$1.9 \times 10^7$ copies/g for *sul2*, $1.49 \times 10^3$ to 2.5 × 106 copies/g for *tetO,* $2.34 \times 10^3$ to 9.45 × 106

copies/g for *tetW, 2.12* $\times 10^3$ to $5.52 \times 10^6$ copies/g for *ermF*. The quantity of ARGs

abundances in the sediments of the urban river were higher than those in the sediments

collected from the natural river.

Further, the urban areas displayed a higher abundance of taxa involved in fecal contamination

and pathogens compared to the natural river (ANOVA, $P < 0.05$). For example, high abundance

of Bacteroidetes and Firmicutes were observed in the urban river (Dataset 2C, $P <0.01$).

Firmicutes and Bacteroidetes are two major phyla of bacteria found in the human intestine, and

have been frequently found in WWTPs and wastewater-influenced environments (63), which

384 are recognized as good indicators of fecal contamination (40, 64). The elevated abundance of

385 Bacteroidetes and Firmicutes in our collected urban river sediment samples suggested the

386 presence of untreated wastewater from residential areas in these areas, which may contain

387 fecal pollutants. Additionally, according to Pathogen–Host Interaction (PHI) database (65), our

388 study also identified a marked increase in the abundance and diversity of pathogens like

389 *Acinetobacter*, *Arcobacter*, *Exiguobacterium*, *Chryseobacterium*, *Escherichia Shigella* and

390 *Staphylococcus aureus* within urban river (Dataset 2E-F, P < 0.01). These findings highlighted

391 urban non-point source untreated water pollutants have resulted in fecal and pathogenic

392 bacterium contamination in the river network.

393 **3.6 Key influencing factors and ecological driving processes**

394 Across the large river network spanning a geographic distance of ~400 km, the microbial

395 community increased in dissimilarity with increasing spatial distance, showed a distance-decay

396 relationship (DDR) (Fig. 6A, Dataset 2G). However, we observed microbial communities in the

397 natural rivers had a steeper DDR slope between sampled sites than those in the urban rivers

398 (Fig. 6A), indicating the environmental selection has a stronger effect on the urban river. The

399 distance-based redundancy analysis (db-RDA) using environmental parameters as

400 constraining variables, highlighted the potential significance of these chemical parameters in

401 elucidating microbial community compositions (Fig. 2I). Ammonium (r sqr= 0.3551, *P*=0.001),

402 pH (r sqr= 0.2293, *P*=0.005), Nitrite (r sqr=0.2211, *P*=0.003), Nitrate (r sqr=0.188 *P*=0.006), TN

403 (r sqr= 0.17, *P*=0.006), TOC (r sqr= 0.1367, *P*=0.032), phosphate (r sqr= 0.0615, *P*=0.010)

404 were more crucial in shaping the microbial community.

406 To further assess the underlying factors governed community dynamics in the river network, the

407 null model (66) and neutral community model (NCM) (48) were used to quantify the relative

408 influences of stochastic and deterministic processes in our study. The null model indicated a

409 prevalence of environmental selection processes, particularly in the form of homogeneous

410 selection (|βNTI| < 2, RCbary >+0.95) (Fig. 6B), in governing the urban river community

411 assembly (details in Materials and Methods). Simultaneously, neutral community model (Fig.

412 6C) showed a gradual increase in the contribution of deterministic process with the increase of

413 human activity and eutrophication, elucidating48.3% and 58.5% of the stochastic process within

414 the natural river and urban river samples, respectively. The NCM and null model further

revealed that deterministic process, specifically homogeneous selection, are the primary drivers
of bacterial and archaeal community compositions in urban river sediments.

**4. Discussion**

In this study, we explored the bacterial and archaeal communities in urban and natural rivers
along the trophic gradient and examined the drivers underlying the observed alpha and beta
diversity. Considerable variation exists within the bacterial and archaeal communities in terms
of abundance, richness, diversity (Fig. 2A-C) and composition (Fig. 2H-J) between urban and
natural rivers, and the urbanization enhance microbial population and biodiversity,
accompanied by significant niche differentiation (Fig. 2D). Nonmetric multidimensional scaling
(NMDS) (Fig. 2I) and network analysis (Fig. 2J), based on OTUs, demonstrated distinct
differences between the microbial communities of urban and natural river ecosystems,
indicating that community composition trends became increasingly similar as eutrophication
intensified. The impact of human activities on river microbial communities was also evident in
the predicted metabolic functions of these communities. This differentiation persisted even
when considering the exclusion of "rare biosphere" taxa (< 0.1% total sequence).
430
We inferred that the observed biotic homogenization caused by urbanization eutrophication
result from the habitat heterogeneity and environmental filtering. In highly eutrophic urban river
systems, the rapid urbanization tends to decrease the overall habitat heterogeneity. For
instance, the high nitrogen/phosphorus loading, high organic carbon and lower pH (Fig. 1), are
common consequences of the continuous input of industrial, domestic, and agricultural
wastewater, which was caused by a large population of approximately 11.06 million in sampled
regions (19). The db-RDA analysis (Fig. 2I) indicated that pH and nitrogen (often in the form of
high abundance of nitrate and ammonium) contributed significantly to the changes in the urban
river microbial community. pH is widely recognized as a major environmental driver shaping
microbial communities, influencing the balance and interaction between niche-related and
neutral processes (67). More acidic conditions (Fig. 1) imposed stronger environmental filtering
and largely shifted community composition. Meanwhile, excessive nutrition inputs lead to
eutrophication (Dataset 1B), a process where the nutrient enrichment promotes the growth of
nitrogen-transformation microorganisms and algae (Chlorophyll *a* content in Fig. 1) (68). The
proliferation of nitrogen-transformation microorganisms and algae leads to increased organic
matter production (TOC content in Fig. 1).

Geng et al. (2022) proposed that eutrophication can induce homogenization by modulating interspecies interactions. For instance, organic carbon generated through phytoplankton photosynthesis serves as a crucial carbon source for river microorganisms. Variations in the organic carbon produced by different phytoplankton species select for distinct bacterial communities with varying capacities for organic matter utilization (25). Eutrophication promotes the accumulation of organic matter derived from algae, altering the sources of organic mass and impacting the structure of river microbial communities (69). Additionally, the "productivity/resources hypothesis" suggests that enhanced availability of resources and/or increased primary productivity can sustain larger populations (70). Therefore, the influx of human activity pollutants into the river created a more uniform and homogeneity environment conditions, allowing for the coexistence of a broader range of microbial species. The increased availability of organic matter and the changed nutrient conditions can profoundly influence the river microbial population, and enhanced the microbial population sizes and diversity in urbanized rivers.

Further, higher ARRs and ARGs were also observed in highly urbanized river areas (Fig. 5A-B). Notably, the data showed that significant variation in ARRs and ARGs between cities, with no significant differences in natural river samples. This feature emphasized the importance of the different levels of human activity contribution based on population sizes, potentially resulting in varied ARRs and ARGs among urban rivers. Zhang et al. (2015) reported that over 68% of total antibiotic usage in China in 2013 was excreted by humans and animals, leading to around 53,800 tons of antibiotics entering the environment (71). Urban water ecosystems have been identified as potential hotspots for the spread of antibiotic resistant bacteria in China (72, 73). Hence, the continuous selective pressure exerted by antibiotic residues, even at sub-inhibitory concentrations, in combination with the diverse and dense microbial communities sustained by nutrient-rich environments, creates favorable conditions for the dissemination and proliferation of antibiotic resistance (42).

Meanwhile, nutrient enrichment often coincides with heavy metal pollution in numerous human-impacted aquatic ecosystems (74). In our study, proportion of the archaea in the urban river areas exhibited 1.58 times higher than those in natural river, and archaeal Chao1 richness and Shannon diversity exhibited 1.47 and 2.36 times higher than that in the natural river, respectively. This finding is similar to previous study revealed that in a high heavy metal

contamination zone, archaeal Chao1 richness and Shannon diversity exhibited substantial
increases, approximately 1.81 and 1.56 times higher than those in a low heavy metal
contamination zone, respectively (10). Krzmarzick et al. (2018) proposed that archaea have
demonstrated a notable capacity for enduring harsh environments, leading to the possibility that
numerous archaeal species possessing exceptional adaptability to severe anthropogenic
pollution could flourish in urban river environments (75). Therefore, habitat heterogeneity plays
a crucial role in determining alpha and beta diversity in river microbiomes.
488
Secondly, this study aimed to investigate the assembly mechanisms of both bacterial and
archaeal communities in urban and natural rivers using environmental DNA meta-barcoding
technology. River microbial communities exhibited high diversity, encompassing thousands of
OTUs across major taxa that inhabit various habitats within the river ecosystem (Fig. 2-3). It is
worth noting that the distance-decay relationship (DDR) was observed in urban and natural
river samples (Fig. 6A). Deciphering the relative influence of diverse ecological processes in
microbial community assembly remain an important query in riverine biogeography (76). The
strength of the distance–decay of community similarity is influenced by both stochastic and
deterministic processes (27, 77). Among them, some previous studies had found the
geographic distances (passive dispersal) and environmental filtering are two major ecological
processes impacted on the assembly of microbial communities in connected river ecosystem
(78-80). For example, in large scale watersheds like the Yangtze River and Tingjiang River,
geographic separation and dispersal limitation have been identified as an important factor in the
succession of communities (10, 80, 81).
503
However, in our study, the microbial communities in natural river locations exhibited a more
pronounced DDR slope among sampling sites than those in urban river sites (Fig. 6A). The
variation in microbial community composition tends to be less pronounced across urban areas
compared to the diversity found across natural river ecosystems. Furthermore, our analyses
also show narrower niche breadths were observed for samples in the urban river (Fig. 2D), and
a higher level of homogenization in the community composition along the urban river compared
to the natural river ecosystems (Fig. 2I-J). The application of the neutral community model (Fig.
6C) revealed that deterministic processes predominantly shaped the urban river microbial
communities in the river network. Null model further uncovered the homogeneous selection
dominated governs the urban river community assembly (Fig. 6B). This finding further suggests
that persistent uniform non-point source pollution such as high nutrients, PAHs, antibiotics

pollutions arises from the urbanization created a more uniformity environment in urban river
system, exerting robust homogeneous selection on urban river ecosystem. Homogeneous
selection is likely a potent mechanism for biotic homogenization, particularly in harsh
environments like those affected by eutrophication, where it diminishes the role of stochastic
processes in shaping assemblages (25). Therefore, similar urban river environmental condition
could promote community fusion, contributing to a more similar community structure (Fig. 2I-J)
and function (Fig.4), and indicating that taxonomic and functional homogenization occur
simultaneously in the river network.
523
Our findings, highlighting convergence in the microbiome of urban rivers, align with observed
urbanization impacts on communities of bird (82), fish (83), plant (84), invertebrate (85, 86),
zooplankton (87), dust samples (88). As outlined by Olden and Poff (2003), functional
homogenization occurs when a specific type of environment becomes more uniformly common.
Consequently, the proliferation of analogous species and consistent physical environments in
heavily altered urban river habitats is likely lead to a rise in functional homogenization.
530
On the other hand, differences in the local species pool between urban and natural river areas
were evident, with shared microorganisms accounting for 55% (Fig. 2H). This further
underscore distinct environmental conditions between these study areas result in varied local
species pools and species abundance distributions. The urban river habitats promote the
establishment of 374 "urban-adaptable" species (Fig. 2H), and these "urban-adaptable" species
dominate across urban river sites. Meanwhile, a narrower niche breadth has been found in
urban river areas compared to natural river areas (Fig. 2D), suggesting a lesser metabolic
plasticity of microorganisms. We hypothesize that urban eutrophication favors species adapted
to pollution tolerance, thereby promoting biotic homogenization.
540
In natural ecosystems, microorganisms are typically categorized as habitat generalists and
specialists based on their niche breadth (89). Habitat generalists with broad niche breadth are
less competitive under optimal growth conditions but exhibit greater resistance to changing
environments, whereas specialists have narrower niches and demonstrate specific
environmental fitness (90). We had confirmed that anthropogenic activities decreasing urban
local environmental heterogeneity in interconnected river network on a regional scale. This
decreased environmental heterogeneity not only deteriorates the habitats of indigenous species
that are at a competitive disadvantage but also fosters urban-adaptable species (including the

exotic species or range expansion of native specialist species) acclimatizing to urban river

conditions (Fig. 2D, H) (91). Hence, the mechanisms of biotic homogenization can be attributed

to three aspects: (1) extinction of native generalist species, (2) invasion of widespread

nonnative species, and (3) range expansion of native specialist species.

Non-native (exotic) biota and native specialist species becomes increasingly enriched in urban

rivers, replacing many natural river locally unique, endemic native species from natural rivers,

thereby reducing the unique regional identity of biotas. For example, urbanization-induced biotic

homogenization leads to an increase in local biodiversity within urban areas (Fig. 2B-C), yet it

contributes to a decrease in regional biodiversity in the urban areas (e.g., 3499 species in the

urban areas, 3736 species in natural areas, Fig. 2H). It's worth noting that global diversity

diminishes due to the extinction of unique regional species, which are then absent from the

worldwide pool of species (92). Therefore, the biotic homogenization could potentially have

detrimental effects on crucial ecosystem functions. The finding serves as a compelling example

of biotic homogenization, highlighting how human-induced environmental changes are

diminishing the biological diversity that exists between natural ecosystems. This trend is

expected to lead to a more uniform biosphere at both regional and global levels (92),

consequently diminishing global biodiversity. This contrast between local augmentation and

global diminution of biodiversity represents a critical concern for conservation biology, as it

could potentially divert public attention from the more pressing issue of worldwide species loss.

**5. Conclusion**

Collectively, our research demonstrated that human population and contamination level serve

as the predominant influences shaping urban river microbial communities in the river network.

Changed environmental conditions can potentially favor specific bacterial and archaeal taxa,

alter the alpha and beta diversity of urban river microbiomes, and contribute to the biotic

homogenization of river microbiomes. Biotic homogenization can be attributed to the

modulation of generalist/specialist species and the invasion of nonnative species. These

changes have cascading effects on ecosystem functioning and overall microbial processes in

the river network, potentially profoundly altering riverine ecosystems. Future research is

required to assess the extent of biotic homogenization due to human activities across global

river ecosystems.

**Acknowledgements**

This work was supported by the Youth Program of National Natural Science Foundation of
China (No. 32100081), Central Public-Interest Scientific Institution Basal Research Fund, CAFS
(NO.14).

**Competing Interest Statement**

The authors declare no competing interests.

**Declaration of Generative AI and AI-assisted technologies in the writing process**

Statement: During the preparation of this work the authors used CHATGPT in order to improve
readability and language. After using this tool/service, the authors reviewed and edited the
content as needed and takes full responsibility for the content of the publication.

**Reference**

1.   Widder S, Besemer K, Singer GA, Ceola S, Bertuzzo E, Quince C, et al. Fluvial network
organization imprints on microbial co-occurrence networks. Proceedings of the National
Academy of Sciences. 2014;111(35):12799-804.
2.   Beattie RE, Bandla A, Swarup S, Hristova KR. Freshwater sediment microbial communities
are not resilient to disturbance from agricultural land runoff. Front Microbiol. 2020;11:539921.
3.   Wang Y, Sheng HF, He Y, Wu JY, Jiang YX, Tam NF, et al. Comparison of the levels of
bacterial diversity in freshwater, intertidal wetland, and marine sediments by using millions of
illumina tags. Appl Environ Microbiol. 2012;78(23):8264-71.
4.   Singh R, Singh AP, Kumar S, Giri BS, Kim K-H. Antibiotic resistance in major rivers in the
world: A systematic review on occurrence, emergence, and management strategies. Journal of
Cleaner Production. 2019;234:1484-505.
5.   Patel AB, Shaikh S, Jain KR, Desai C, Madamwar D. Polycyclic Aromatic Hydrocarbons:
Sources, Toxicity, and Remediation Approaches. Front Microbiol. 2020;11:562813.
6.   Xu N, Hu H, Wang Y, Zhang Z, Zhang Q, Ke M, et al. Geographic patterns of microbial traits
of river basins in China. Sci Total Environ. 2023;871:162070.
7.   Liu Y, Su B, Mu H, Zhang Y, Chen L, Wu B. Effects of point and nonpoint source pollution
on urban rivers: From the perspective of pollutant composition and toxicity. J Hazard Mater.
2023;460:132441.

8.   Pruden A, Arabi M, Storteboom HN. Correlation Between Upstream Human Activities and
Riverine Antibiotic Resistance Genes. Environ Sci Technol. 2012;46(21):11541-9.
9.   Manyi-Loh CE, Mamphweli SN, Meyer EL, Makaka G, Simon M, Okoh AI. An Overview of
the Control of Bacterial Pathogens in Cattle Manure. International Journal of Environmental
Research and Public Health. 2016;13(9):843.
10. Yang S, Chen Q, Zheng T, Chen Y, Zhao X, He Y, et al. Multiple metal(loid) contamination
reshaped the structure and function of soil archaeal community. J Hazard Mater.
2022;436:129186.
11. Delgado-Baquerizo M, Eldridge DJ, Liu Y-R, Sokoya B, Wang J-T, Hu H-W, et al. Global
homogenization of the structure and function in the soil microbiome of urban greenspaces.
Science Advances. 2021;7(28):eabg5809.
12. Drury B, Rosi-Marshall E, Kelly JJ. Wastewater treatment effluent reduces the abundance
and diversity of benthic bacterial communities in urban and suburban rivers. Appl Environ
Microbiol. 2013;79(6):1897-905.
13. Pascual-Benito M, Ballesté E, Monleón-Getino T, Urmeneta J, Blanch AR, García-Aljaro C,
et al. Impact of treated sewage effluent on the bacterial community composition in an
intermittent mediterranean stream. Environ Pollut. 2020;266:115254.
14. Dai T, Su Z, Zeng Y, Bao Y, Zheng Y, Guo H, et al. Wastewater treatment plant effluent
discharge decreases bacterial community diversity and network complexity in urbanized coastal
sediment. Environ Pollut. 2023;322:121122.
15. Kiedrzyńska E, Kiedrzyński M, Urbaniak M, Magnuszewski A, Skłodowski M, Wyrwicka A,
et al. Point sources of nutrient pollution in the lowland river catchment in the context of the Baltic
Sea eutrophication. Ecol Eng. 2014;70:337-48.
16. Ding X, Liu L. Long-term effects of anthropogenic factors on nonpoint source pollution in the
upper reaches of the Yangtze river. Sustainability. 2019;11(8):2246.
17. EPA. Basic Information about Nonpoint Source (NPS) Pollution United States
Environmental Protection Agency2022 [Available from:
https://www.epa.gov/nps/basic-information-about-nonpoint-source-nps-pollution.
18. Wang H, Dong Y, Jiang Y, Zhang N, Liu Y, Lu X, et al. Multiple stressors determine the
process of the benthic diatom community assembly and network stability in urban water bodies
in Harbin. Sci Total Environ. 2024;913:169536.
19. Cheng M, Duan C. The changing trends of internal migration and urbanization in China:
new evidence from the seventh National Population Census. China Population and
Development Studies. 2021;5:275-95.
20. Dudley N. Guidelines for applying protected area management categories: Iucn; 2008.
21. Xiao H, Zhang X, Yan M, Zhang L, Wang H, Ma Y, et al. The Temporal-Based Forest
Disturbance Monitoring Analysis: A Case Study of Nature Reserves of Hainan Island of China
From 1987 to 2020. Frontiers in Environmental Science. 2022;10.

22. Gibbons SM, Jones E, Bearquiver A, Blackwolf F, Roundstone W, Scott N, et al. Human
and Environmental Impacts on River Sediment Microbial Communities. PLoS One.
2014;9(5):e97435.
23. Battin TJ, Besemer K, Bengtsson MM, Romani AM, Packmann AI. The ecology and
biogeochemistry of stream biofilms. Nat Rev Microbiol. 2016;14(4):251-63.
24. Xie Y, Wang J, Wu Y, Ren C, Song C, Yang J, et al. Using in situ bacterial communities to
monitor contaminants in river sediments. Environ Pollut. 2016;212:348-57.
25. Geng M, Zhang W, Hu T, Wang R, Cheng X, Wang J. Eutrophication causes microbial
community homogenization via modulating generalist species. Water Res. 2022;210:118003.
26. Carlson RE. A trophic state index for lakes 1. Limnol Oceanogr. 1977;22(2):361-9.
27. Li H, Zhou H, Yang S, Dai X. Stochastic and Deterministic Assembly Processes in
Seamount Microbial Communities. Applied and environmental microbiology. 2023;89(7).
28. Eickhorst T, Tippkötter R. Improved detection of soil microorganisms using fluorescence in
situ hybridization (FISH) and catalyzed reporter deposition (CARD-FISH). Soil Biol Biochem.
2008;40(7):1883-91.
29. Hill JE, Town JR, Hemmingsen SM. Improved template representation in cpn60
polymerase chain reaction (PCR) product libraries generated from complex templates by
application of a specific mixture of PCR primers. Environ Microbiol. 2006;8(4):741-6.
30. Hoshino T, Doi H, Uramoto GI, Wormer L, Adhikari RR, Xiao N, et al. Global diversity of
microbial communities in marine sediment. Proc Natl Acad Sci U S A. 2020;117(44):27587-97.
31. Li H, Yang Q, Zhou H. Niche differentiation of sulfate-and iron-dependent anaerobic
methane oxidation and methylotrophic methanogenesis in deep sea methane seeps. Front
Microbiol. 2020;11:1409.
32. Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, et al. Introducing
mothur: open-source, platform-independent, community-supported software for describing and
comparing microbial communities. Appl Environ Microbiol. 2009;75(23):7537-41.
33. Levins R. Evolution in changing environments: some theoretical explorations: Princeton
University Press; 1968.
34. Pandit SN, Kolasa J, Cottenie K. Contrasts between habitat generalists and specialists: an
empirical extension to the basic metacommunity framework. Ecology. 2009;90(8):2253-62.
35. Jiao S, Yang Y, Xu Y, Zhang J, Lu Y. Balance between community assembly processes
mediates species coexistence in agricultural soil microbiomes across eastern China. The ISME
Journal. 2020;14(1):202-16.
36. Salazar G. EcolUtils: Utilities for community ecology analysis. R package. 2020.
37. Douglas GM, Maffei VJ, Zaneveld JR, Yurgel SN, Brown JR, Taylor CM, et al. PICRUSt2
for prediction of metagenome functions. Nat Biotechnol. 2020;38(6):685-8.
38. Langille MG, Zaneveld J, Caporaso JG, McDonald D, Knights D, Reyes JA, et al. Predictive
functional profiling of microbial communities using 16S rRNA marker gene sequences. Nat
Biotechnol. 2013;31(9):814-21.

39. Louca S, Parfrey LW, Doebeli M. Decoupling function and taxonomy in the global ocean
microbiome. Science. 2016;353(6305):1272-7.
40. Wang L, Zhang J, Li H, Yang H, Peng C, Peng Z, et al. Shift in the microbial community
composition of surface water and sediment along an urban river. Sci Total Environ.
2018;627:600-12.
41. Berendonk TU, Manaia CM, Merlin C, Fatta-Kassinos D, Cytryn E, Walsh F, et al. Tackling
antibiotic resistance: the environmental framework. Nat Rev Microbiol. 2015;13(5):310-7.
42. Rizzo L, Manaia C, Merlin C, Schwartz T, Dagot C, Ploy MC, et al. Urban wastewater
treatment plants as hotspots for antibiotic resistant bacteria and genes spread into the
environment: A review. Sci Total Environ. 2013;447:345-60.
43. Munir M, Wong K, Xagoraraki I. Release of antibiotic resistant bacteria and genes in the
effluent and biosolids of five wastewater utilities in Michigan. Water Res. 2011;45(2):681-93.
44. Gao P, Mao D, Luo Y, Wang L, Xu B, Xu L. Occurrence of sulfonamide and
tetracycline-resistant bacteria and resistance genes in aquaculture environment. Water Res.
2012;46(7):2355-64.
45. Milaković M, Križanović S, Petrić I, Šimatović A, González-Plaza JJ, Gužvinec M, et al.
Characterization of macrolide resistance in bacteria isolated from macrolide-polluted and
unpolluted river sediments and clinical sources in Croatia. Sci Total Environ. 2020;749:142357.
46. Brooks J, Maxwell S, Rensing C, Gerba C, Pepper I. Occurrence of antibiotic-resistant
bacteria and endotoxin associated with the land application of biosolids. Can J Microbiol.
2007;53(5):616-22.
47. Li H, Yang Q, Li J, Gao H, Li P, Zhou H. The impact of temperature on microbial diversity
and AOA activity in the Tengchong Geothermal Field, China. Sci Rep. 2015;5:17056.
48. Sloan WT, Lunn M, Woodcock S, Head IM, Nee S, Curtis TP. Quantifying the roles of
immigration and chance in shaping prokaryote community structure. Environ Microbiol.
2006;8(4):732-40.
49. Xie W-Q, Gong Y-X. Measurement of permanganate index in environmental water via
indirect phase-conversion strategy. Journal of Chromatography A. 2024;1728:464987.
50. Zhang J, Zhang B, Liu Y, Guo Y, Shi P, Wei G. Distinct large-scale biogeographic patterns
of fungal communities in bulk soil and soybean rhizosphere in China. Sci Total Environ.
2018;644:791-800.
51. Ruff SE, Biddle JF, Teske AP, Knittel K, Boetius A, Ramette A. Global dispersion and local
diversification of the methane seep microbiome. Proc Natl Acad Sci U S A
2015;112(13):4015-20.
52. Amend AS, Oliver TA, Amaral-Zettler LA, Boetius A, Fuhrman JA, Horner-Devine MC, et al.
Macroecological patterns of marine bacteria on a global scale. J Biogeogr. 2013;40(4):800-11.
53. Masuda Y, Yamanaka H, Xu Z-X, Shiratori Y, Aono T, Amachi S, et al. Diazotrophic
anaeromyxobacter isolates from soils. Applied and environmental microbiology.
2020;86(16):e00956-20.

54. Lücker S, Wagner M, Maixner F, Pelletier E, Koch H, Vacherie B, et al. A Nitrospira
metagenome illuminates the physiology and evolution of globally important nitrite-oxidizing
bacteria. Proc Natl Acad Sci U S A. 2010;107(30):13479-84.
55. Pagé AP, Yergeau É, Greer CW. Salix purpurea Stimulates the Expression of Specific
Bacterial Xenobiotic Degradation Genes in a Soil Contaminated with Hydrocarbons. PLoS One.
2015;10(7):e0132062.
56. Kansole MMR, Lin T-F. Microcystin-LR Biodegradation by Bacillus sp.: Reaction Rates and
Possible Genes Involved in the Degradation. Water. 2016;8(11):508.
57. Ghosal D, Ghosh S, Dutta TK, Ahn Y. Corrigendum: Current State of Knowledge in
Microbial Degradation of Polycyclic Aromatic Hydrocarbons (PAHs): A Review. Front Microbiol.
2016;7:1837.
58. Kuppusamy S, Thavamani P, Venkateswarlu K, Lee YB, Naidu R, Megharaj M.
Remediation approaches for polycyclic aromatic hydrocarbons (PAHs) contaminated soils:
Technological constraints, emerging trends and future directions. Chemosphere.
2017;168:944-68.
59. Mojiri A, Zhou JL, Ohashi A, Ozaki N, Kindaichi T. Comprehensive review of polycyclic
aromatic hydrocarbons in water sources, their effects and treatments. Sci Total Environ.
2019;696:133971.
60. Van Boeckel TP, Gandra S, Ashok A, Caudron Q, Grenfell BT, Levin SA, et al. Global
antibiotic consumption 2000 to 2010: an analysis of national pharmaceutical sales data. The
Lancet infectious diseases. 2014;14(8):742-50.
61. WHO. Critically important antimicrobials for human medicine. 6th Revision 2018: World
Health Organization. 2019.
62. Sköld O. Sulfonamide resistance: mechanisms and trends. Drug resistance updates.
2000;3(3):155-60.
63. Saarenheimo J, Aalto SL, Rissanen AJ, Tiirola M. Microbial Community Response on
Wastewater Discharge in Boreal Lake Sediments. Front Microbiol. 2017;8:750.
64. Wéry N, Lhoutellier C, Ducray F, Delgenès J-P, Godon J-J. Behaviour of pathogenic and
indicator bacteria during urban wastewater treatment and sludge composting, as revealed by
quantitative PCR. Water Res. 2008;42(1):53-62.
65. Urban M, Cuzick A, Seager J, Wood V, Rutherford K, Venkatesh SY, et al. PHI-base: the
pathogen–host interactions database. Nucleic Acids Res. 2020;48(D1):D613-D20.
66. Stegen JC, Lin X, Konopka AE, Fredrickson JK. Stochastic and deterministic assembly
processes in subsurface microbial communities. ISME J. 2012;6(9):1653-64.
67. Ren L, Jeppesen E, He D, Wang J, Liboriussen L, Xing P, et al. pH Influences the
Importance of Niche-Related and Neutral Processes in Lacustrine Bacterioplankton Assembly.
Applied and environmental microbiology. 2015;81(9):3104-14.
68. Kuypers MMM, Marchant HK, Kartal B. The microbial nitrogen-cycling network. Nat Rev
Microbiol. 2018;16(5):263-76.

69. Han X, Schubert CJ, Fiskal A, Dubois N, Lever MA. Eutrophication as a driver of microbial
community structure in lake sediments. Environ Microbiol. 2020;22(8):3446-62.
70. Ibarbalz FM, Henry N, Brandao MC, Martini S, Busseni G, Byrne H, et al. Global Trends in
Marine Plankton Diversity across Kingdoms of Life. Cell. 2019;179(5):1084-97 e21.
71. Zhang Q-Q, Ying G-G, Pan C-G, Liu Y-S, Zhao J-L. Comprehensive Evaluation of
Antibiotics Emission and Fate in the River Basins of China: Source Analysis, Multimedia
Modeling, and Linkage to Bacterial Resistance. Environ Sci Technol. 2015;49(11):6772-82.
72. Hou L, Zhang L, Li F, Huang S, Yang J, Ma C, et al. Urban ponds as hotspots of antibiotic
resistome in the urban environment. J Hazard Mater. 2021;403:124008.
73. Reddy S, Kaur K, Barathe P, Shriram V, Govarthanan M, Kumar V. Antimicrobial resistance
in urban river ecosystems. Microbiol Res. 2022;263:127135.
74. Jaiswal D, Pandey J. An ecological response index for simultaneous prediction of
eutrophication and metal pollution in large rivers. Water Res. 2019;161:423-38.
75. Krzmarzick MJ, Taylor DK, Fu X, McCutchan AL. Diversity and Niche of Archaea in
Bioremediation. Archaea. 2018;2018:3194108.
76. Martiny JB, Eisen JA, Penn K, Allison SD, Horner-Devine MC. Drivers of bacterial
beta-diversity depend on spatial scale. Proc Natl Acad Sci U S A. 2011;108(19):7850-4.
77. Stegen JC, Lin X, Fredrickson JK, Chen X, Kennedy DW, Murray CJ, et al. Quantifying
community assembly processes and identifying features that impose them. ISME J.
2013;7(11):2069-79.
78. Dini-Andreote F, Stegen JC, van Elsas JD, Salles JF. Disentangling mechanisms that
mediate the balance between stochastic and deterministic processes in microbial succession.
Proc Natl Acad Sci U S A. 2015;112(11):1326-32.
79. Jia X, Dini-Andreote F, Falcão Salles J. Community assembly processes of the microbial
rare biosphere. Trends Microbiol. 2018;26(9):738-47.
80. Chen W, Ren K, Isabwe A, Chen H, Liu M, Yang J. Stochastic processes shape
microeukaryotic community assembly in a subtropical river across wet and dry seasons.
Microbiome. 2019;7(1):138.
81. Gao Y, Zhang W, Li Y, Wu H, Yang N, Hui C. Dams shift microbial community assembly
and imprint nitrogen transformation along the Yangtze River. Water Res. 2021;189:116579.
82. Murthy AC, Fristoe TS, Burger JR. Homogenizing effects of cities on North American winter
bird diversity. Ecosphere. 2016;7(1):e01216.
83. Menezes RF, Borchsenius F, Svenning JC, Davidson TA, Søndergaard M, Lauridsen TL, et
al. Homogenization of fish assemblages in different lake depth strata at local and regional
scales. Freshwater Biology. 2015;60(4):745-57.
84. Salgado J, Sayer CD, Brooks SJ, Davidson TA, Goldsmith B, Patmore IR, et al.
Eutrophication homogenizes shallow lake macrophyte assemblages over space and time.
Ecosphere. 2018;9(9):e02406.

85. McKinney ML. Effects of urbanization on species richness: a review of plants and animals.
Urban ecosystems. 2008;11:161-76.
86. Donohue I, Jackson AL, Pusch MT, Irvine K. Nutrient enrichment homogenizes lake benthic
assemblages at local and regional scales. Ecology. 2009;90(12):3470-7.
87. Liu P, Xu S, Lin J, Li H, Lin Q, Han B-P. Urbanization increases biotic homogenization of
zooplankton communities in tropical reservoirs. Ecological Indicators. 2020;110:105899.
88. Barberán A, Ladau J, Leff JW, Pollard KS, Menninger HL, Dunn RR, et al. Continental-scale
distributions of dust-associated bacteria and fungi. Proc Natl Acad Sci U S A.
2015;112(18):5756-61.
89. Sun J, Zhou H, Cheng H, Chen Z, Wang Y. Distinct strategies of the habitat generalists and
specialists in the Arctic sediments: Assembly processes, co-occurrence patterns, and
environmental implications. Mar Pollut Bull. 2024;205:116603.
90. Wang L, Han M, Li X, Yu B, Wang H, Ginawi A, et al. Mechanisms of niche-neutrality
balancing can drive the assembling of microbial community. Mol Ecol. 2021;30(6):1492-504.
91. Kowarik I. Novel urban ecosystems, biodiversity, and conservation. Environ Pollut.
2011;159(8):1974-83.
92. McKinney ML. Urbanization as a major cause of biotic homogenization. Biol Conserv.
2006;127(3):247-60.
820
821

822    **Figure**

824    **Figure 1.** (A) Location of the collected river network sediment samples. The Inset map shows

825    the extent of the study area relative to China. (B) Differences in environmental variables

826    between the urban and natural river areas. Orange circles indicate the urban river sampling

827    sites, and blue circles indicate the natural river sampling sites, respectively.

Figure 2. The microbiome in the large river network. (A-D) Microbial abundance, Chao1 richness estimates, Shannon index and Niches breadth were investigated in urban (orange circles) and natural (blue circles) river samples. (E-G) Pairwise relationships between the microbial abundance, Chao 1 richness, Shannon index and TSI for observed communities. (H) Venn analysis. (F) A distance-based redundancy analysis (db-RDA) visualized by two-dimensional (2D) nonmetric multidi-mensional scaling (NMDS) showing microbial community similarity when environmental factors were used as constraining variables and were fitted onto the ordination as arrows. Yellow circles indicate the urban river sampling sites, and red circles indicate the natural river sampling sites, respective. The length of each arrow indicates the multiple partial correlation of the variable to RDA axes and can be interpreted as an indication of that variable's contribution to the explained community similarity. (I) Co-occurrence networks of the microbial community between urban and natural river samples.

**Figure 3.** The relative abundance of the phylum (A) and order (B) in urban and natural river sediment samples.

**Figure 4.** The potentially ecological functions based on 16S rRNA genes of microbial community composition.

**Figure 5.** Different of antibiotic resistance rates (A) and antibiotic resistance gene abundance

(B). The orange circles indicate the urban river and blue circles indicate the natural river,

respectively.

**Figure 6.** General patterns of microbial diversity in the river network system. (A)

Distance-decay curves showing Bray–Curtis dissimilarity against geographic distances

between sampling sites. Solid lines denote the ordinary least-squares linear regressions. (B)

Scatter plot of βNTI values grouped by spatial scales of the river network. (C) Fit of the neutral

community model (NCM) of community assembly. The solid blue lines indicate the best fit to the

NCM as in Sloan et al. (2006), and the dashed blue lines represent 95% confidence intervals

around the model prediction. OTUs that occur more or less frequently than predicted by the

NCM are shown in different colors. Nm indicates the metacommunity size times immigration, $R^2$

indicates the fit to this model.