1 Full title: **Starvation resistance in the nematode *Pristionchus pacificus***
2   **requires a conserved supplementary nuclear receptor**
3
4 Short title: **NHR-10 is required for starvation resistance in *P. pacificus***
5
6
7

8   **Tobias Theska, Tess Renahan, and Ralf J. Sommer***
9
10
11
12 Max Planck Institute for Biology, Tübingen,
13 Department for Integrative Evolutionary Biology,
14 Max-Planck-Ring 9,
15 72076 Tübingen, Germany
16
17
18
19 ***Corresponding author**
20 Ralf J. Sommer (ralf.sommer@tuebingen.mpg.de)
21
22
23
24 **ORCIDs**
25 Tobias Theska - https://orcid.org/0000-0002-0920-4427
26 Tess Renahan - https://orcid.org/0000-0002-4326-0254
27 Ralf J. Sommer - https://orcid.org/0000-0003-1503-7749

28
29
30

31 **Keywords**
32 nuclear receptor, phenotypic plasticity, starvation resistance, ortholog conjecture,
33 nematode, *Pristionchus pacificus*

Abstract

Nuclear hormone receptors (NHRs) are a deeply-conserved superfamily of metazoan transcription factors, which fine-tune the expression of their regulatory target genes in response to a plethora of sensory inputs. In nematodes, NHRs underwent an explosive expansion and many species have hundreds of *nhr* genes, most of which remain functionally uncharacterized. However, recent studies elucidated that two sister receptors, *Ppa*-NHR-1 and *Ppa*-NHR-40, are crucial down-stream regulators of feeding plasticity in the diplogastrid model organism *Pristionchus pacificus.* In this study, we functionally characterize *Ppa*-NHR-10, the closest paralog to *Ppa*-NHR-1 and *Ppa*-NHR-40, aiming to demonstrate that it too regulates aspects of feeding plasticity. We used CRISPR/CAS9-mediated mutagenesis to create knock-out mutations of this receptor and applied a combination of geometric morphometrics and unsupervised clustering to characterize potential mutant phenotypes. Surprisingly, we found that *Ppa*-NHR-10 is not involved in the regulation of plasticity in feeding structures. Instead, multiple RNA-seq experiments revealed that many of the target genes of this receptor are crucial for lipid catabolism. We hypothesized that their mis-regulation could affect the survival of mutant worms during starvation, where lipid catabolism is often essential. Using survival assays, we found that mutant worms indeed show drastically decreased starvation resistance, both as young adults and as dauer larvae. We also characterized genome-wide changes to the transcriptional landscape in *P. pacificus* when exposed to 24hrs of acute starvation, and found that *Ppa*-NHR-10 partially regulates some of these responses. Taken together, we were able to demonstrate that *Ppa*-NHR-10 is broadly required for starvation resistance and regulates different biological processes than its closest paralogs *Ppa*-NHR-1 and *Ppa*-NHR-40 and its one-to-one ortholog in *Caenorhabditis elegans.* This shows that, amongst distantly-related model nematodes, even deeply-conserved NHRs can regulate vastly distinct biological processes and that neither paralogs, nor one-to-one orthologs, may be adequate predictors of their biological functions.

Author summary

When predicting the biological function of a gene (or the protein it encodes) we usually use its homologs for reference since it is likely that our gene or protein of interest has homologs in a different organism (orthologs) or in the same organism (paralogs), whose biological roles have already been characterized via molecular genetic interventions. However, predicting biological functions from homologs can have its limitations. In this study, we functionally characterized a conserved nuclear receptor, *Ppa*-NHR-10, in the nematode *Pristionchus pacificus* and found that both dauer larvae (a specialized dispersal stage) and adult worms are less resistant to starvation if this receptor is experimentally rendered non-functional. Not only did these findings reveal that *Ppa*-NHR-10 is crucial for the worms to survive harsh conditions they would frequently encounter in the wild, they also demonstrated that *Ppa*-NHR-10's homologs would not have been adequate predictors of its biological role: its homolog in *Caenorhabditis elegans* regulates a metabolic shunt pathway; its paralogs in *P. pacificus* regulate feeding plasticity. Thus, our data suggests that conserved nuclear receptors can functionally diversify across distantly-related nematodes, and that we cannot categorically infer their biological roles from well-studied homologs.

Introduction

Nematodes are an incredibly abundant and stunningly diverse group of roundworms, which occupy nearly all ecosystems across the globe. Recent estimates predict that they make up approximately 80% of all terrestrial animals [1] and that the number of unnamed species ranks in the millions [2,3]. Historically, nematodes, especially *Caenorhabditis elegans*, have been used by cell biologists to unravel fundamental secrets of animal physiology, metabolism, and cell fate [4-8], and by developmental biologists to dissect the essential processes of animal ontogeny, including heterochrony, developmental systems drift, and phenotypic plasticity [9-14].

Phenotypic plasticity, the ability of a single genotype to produce different phenotypes depending on the environmental cues that the organism receives during development, has re-gained wide-spread attention as a crucial driving force in organismal evolution [15], especially in the context of contemporary debates surrounding a proposed extension of standard evolutionary theory [15-18].

Diplogastrid nematodes such as *Pristionchus pacificus* and its conspecifics are model organisms used to study the molecular mechanisms that govern phenotypic plasticity and the role it plays in phenotypic evolution [12,18-20]. These worms display a morphological novelty in their mouths that renders them unique amongst roundworms: they possess movable cuticular teeth which allow them to predate on other nematodes (Fig 1A). This phenotype is plastic; depending on the environment, one of two alternative adult morphologies can be adopted: the predatory "eurystomatous" morph which has two cuticular teeth, or the strictly bacterial-feeding "stenostomatous" morph that only has a single tooth (Fig 1A) [12]. Recent studies utilizing *P. pacificus* began to elucidate the complex genetics behind this dimorphism [12,21-23] and identified two transcription factors, *Ppa*-NHR-1 and *Ppa*-NHR-40, as crucial down-stream regulators in the gene regulatory network (GRN) that governs mouth-form plasticity [24]. While these NHRs constitute important regulatory elements of the plasticity GRN, new molecular players continue to be found on a regular basis [25-27], expanding our knowledge of this genetic network and how it evolved along with the morphologically novel teeth.

Another plastic phenotype maintained in *P. pacificus*, and most nematodes, is related to their life cycles. Under suboptimal environmental conditions, e.g. overcrowding or food depletion, young larvae can enter an indirect life cycle (Fig. 1B) as dauers, an arrested developmental stage [28,29]. Dauers are specialized non-feeding larvae adapted for stress survival [6,28] and dispersal [30]. Characteristic morphological features of dauers include a thickened body wall cuticle and a closed buccal cavity with a plug clogging the alimentary channel (Fig 1C). Once environmental conditions turn more favorable, the worms can exit dauer and resume development into reproducing adults (Fig 1B). Interestingly, as in mouth-form plasticity, NHRs play important roles in regulating dauer plasticity [29,31].

NHRs - nuclear (hormone) receptors - constitute one of the most important transcription factors classes in animals, in which they integrate cellular metabolism with physiological function [32]. These transcription factors are restricted to metazoans [33] and they generally share an archetypical protein domain organization. Their two most important domains are the DNA-binding domain (DBD), which recognizes specific DNA motifs in the *cis*-regulatory regions of their regulatory target genes, and the ligand-binding domain (LBD), which regulates the activity state of the NHR depending on the presence or absence of a specific ligand [32,33].

Recent advances in genomics revealed that nematodes, in contrast to other animals, often possess hundreds of *nhr* genes [24,34]. The overwhelming majority of these, called "supplementary nuclear receptors" (*supnrs*), originated from an explosive burst of duplications of the nuclear receptor HNF4 [35,36]. Another striking observation among distantly-related nematodes with similarly high numbers of NHRs (e.g. *C. elegans* [n=266] and *P. pacificus* [n=254]) showed that the vast majority of NHRs is completely lineage-specific with no one-to-one orthologs in other species [24]. Additionally, the expression of *supnr* genes in the nervous system is highly biased toward sensory neurons [34], suggesting that the expansion of NHRs, which is particularly blatant in free-living nematodes [34], could be linked to the worm's complex sensory environments filled with entangled microbial foods. Here, *supnrs* might act as sensors of environmental food signals or internal metabolic signals derived from ingested bacterial food, and may adjust the worm's physiological states or behavioral responses according to perceived environmental information [34], an idea that has recently been demonstrated for a *supnr* in *C. elegans* [37].

Intriguingly, *Ppa-nhr-1* and *Ppa-nhr-40*, the aforementioned mouth-form plasticity regulators in *P. pacificus*, are representatives of these nematode-specific *supnrs* [24]. Yet, they are two of the very few *supnrs* which are actually conserved across distantly-related species. A previous study demonstrated that they co-regulate a group of fast-evolving genes encoding enzymes needed for the turnover of the cuticular material that makes up the worm's teeth [24]. The same study also included a phylogeny of the NHRs found in *C. elegans* and *P. pacificus*. Interestingly, this phylogeny indicated that one more NHR, *Ppa-NHR-10*, might be another deeply-conserved *supnr* and the potential sister receptor to *Ppa-NHR-1* and *Ppa-NHR-40*. If this were true, we speculated, *Ppa-NHR-10* could be another important mouth-form plasticity regulator that was hitherto overlooked.

Thus, we set out to investigate the biological function of *Ppa-NHR-10*, using an integrative approach that combines CRISPR-mediated knockouts of this receptor, geometric morphometric analyses of mutant mouth morphologies, and regulatory-target identification via transcriptomics. Surprisingly, our data demonstrated that *Ppa-NHR-10* is not involved in regulating mouth-form plasticity, but that it is required for the survival of acute starvation in young adults and long-term starvation in dauer. We also elucidated genome-wide patterns of transcriptional responses to acute starvation in *P. pacificus* and found that NHR-10's regulatory targets include multiple enzymes involved in lipid catabolism and transport, some of which are starvation-response genes (SRGs).

Results

Ppa-NHR-10 is a deeply-conserved supplementary nuclear receptor and the closest relative of Ppa-NHR-1 and Ppa-NHR-40

We first aimed to investigate whether *Ppa*-NHR-10 is indeed a member of the supplementary nuclear receptor cluster that includes *Ppa*-NHR-1 and *Ppa*-NHR-40, as indicated in the two-species NHR phylogeny in Sieriebriennikov *et al.* [24]. For this, we searched for orthologs of these three receptor proteins across different nematode species, based on which we aimed to reconstruct their phylogeny. We decided to include a total of ten species, representing members of clades III-V, in our analysis. We also expanded the number of representative species in Diplogastridae and Rhabditidae, as this allows a more nuanced reconstruction of the evolution of this small NHR-cluster in these families.

We found that these three NHRs indeed form a cluster of *supnrs* that is deeply conserved across the investigated nematode clades and that, usually, each species has exactly one copy of each of these receptors encoded in their genome (Fig 1D). Notable exceptions to this observation are *Brugia malayi*, which seems to lack an ortholog of NHR-10, and *Haemonchus contortus*, which lacks a copy of NHR-40 but has two copies of NHR-10. We were also able to confirm previous results [24] which suggested that NHR-1 and NHR-10 are sister receptors, and that NHR-40 is their next closest relative. Lastly, our maximum likelihood tree suggests that NHR-68, NHR-101, and NHR-114 form a group of supplementary receptors that evolved via multiple gene duplications in the lineage leading to *Caenorhabditis* nematodes. Thus, these receptors lack any one-to-one orthologs in the other investigated nematode species. In contrast, one-to-one orthology across species could clearly be established for NHR-1, NHR-10, and NHR-40 (Fig 1D).

These results suggest that, although these three NHRs represent nematode-specific *supnrs*, they were conserved across at least 130 million years of evolution [38]. This is stunning, as the vast majority of these receptors is entirely lineage-specific, undergoing rapid evolutionary turnover [24]. Additionally, NHR-10 is deeply nested within this cluster and, in *P. pacificus*, it is the last remaining one among these three receptors for which the biological function remains unknown. We hypothesized that *Ppa*-NHR-10 is likely another regulator of this plasticity and that, probably, the entire NHR-1/-10/-40 cluster was co-opted into this functional context during diplogastrid evolution. To further investigate this possibility, we wanted to study the cellular expression patterns of *Ppa-nhr-10*.

Ppa-nhr-10 is expressed in multiple tissues including the hypodermis, pharyngeal glands, uteri, spermathecae, and rectal glands

Previous studies from our lab demonstrated that *Ppa-nhr-1* and *Ppa-nhr-40* are co-expressed in the pharyngeal muscles, the tissues that produce the cuticular teeth of *P. pacificus* [24]. Additionally, all of their joint regulatory targets show an incredibly consistent expression pattern in a single cell: g1D. This is a gland cell of *P. pacificus*, which is morphologically integrated with the dorsal pharynx, where it runs through the dorsal tooth and opens into the mouth lumen [39]. Thus, we anticipated to find *Ppa*-NHR-10 to be expressed in any of these tissues.

Our transcriptional reporter animals revealed that *Ppa-nhr-10* is expressed in a variety of tissues (Fig 2). These always included the skin of the worms (hypodermis) (Fig 2A, 2B, and 2C), the anterior and posterior uteri and spermathecae (Fig 2A, 2B, and 2D), the

rectal glands (Fig 2A), and, expectedly, the pharyngeal glands (Fig 2A and 2C). While all of our worms consistently showed an expression in all three of the pharyngeal gland cells (g1D, g1VL, g1VR) [39], the relative intensity of the expression signal varied across the two independently acquired reporter lines: in one reporter line the ventral pharyngeal glands (g1V) dominate the signal (Fig 2A), while in the other the dorsal pharyngeal gland (g1D) reports more strongly (Fig 2B and 2C). Additionally, one of the lines only shows a weak expression pattern in the rectal glands (Fig 2A *vs.* 2B). Thus, the reported expression patterns were, at least partially, in line with our expectations.

Taken together, the fact that *Ppa-nhr-10* is the deeply conserved paralog of *Ppa-nhr-1* and *Ppa-nhr-40*, and that it is expressed in the same tissues (and/or their regulatory targets), indicated to us that this *supnr* may indeed be another member of the plasticity-regulating network in *P. pacificus*. Thus, we set out to characterize whether the loss of *Ppa-nhr-10* would lead to any mouth-form plasticity related mutant phenotypes.

Loss of *Ppa-nhr-10* does not affect stoma morphology and only weakly affects mouth-form plasticity

We created multiple independent *loss-of-function* alleles of *Ppa-nhr-10* (genome annotation: El Paco v3, gene_ID: PPA01780, chromosome: ChrIII). We used CRISPR/CAS9 to introduce frameshift mutations into the gene body that would disrupt and truncate the nuclear receptor protein either at the site of the DNA-binding domain (DBD) or the ligand-binding domain (LBD). While we were unable to obtain viable frameshift alleles in the DBD, we succeeded in independently generating four homozygous mutant lines with frameshifts truncating the LBD (crRNA sequence: GGCGCGTGGGTTCGGGCGTA; PAM: CGG). Two of these lines were chosen to establish the *Ppa-nhr-10(lof)* reference strains RS3920(*tu1654*) and RS3921(*tu1655*), which we used for all subsequent experiments in this study (Fig 3A).

Next, we wondered whether our *Ppa-nhr-10* mutants display any morphological phenotypes related to mouth-form plasticity, just as *Ppa-nhr-1* and *Ppa-nhr-40* mutants do [22,24]. We found that both *Ppa-nhr-10* mutant strains show a detectable decrease in the number of predatory morphs when compared to wild-type worms (Fig 3B). While this effect is well supported by our data (wildtype *vs.* RS3920: $Z = 2.1$, $P = 0.0058$; wildtype *vs.* RS3921: $Z = 2.6$, $P = 0.0004$) and consistent across both mutant strains (RS3920 vs. RS3921: $Z = 0.05$, $P = 0.4839$), it is only marginal in size (~ 4-5%). This is in stark contrast to the effects observed for *loss-of-function* and *gain-of-function* mutations in *Ppa-nhr-40*, which lead to a complete loss of the predatory or bacterivorous morphs, respectively. Given the small size of the effect that *Ppa-nhr-10* mutations exert on the worm's morph ratio, and taking into account the vast diversity of factors influencing the mouth-form decision [12,18,27], we concluded that *Ppa-nhr-10* is unlikely to be a direct regulator of mouth-form plasticity. Indeed, it seems more likely that the small observed effects on the morph ratio are indirect consequences of the misregulation of *Ppa-nhr-10*'s true regulatory target pathways.

Additionally, we found that the loss of *Ppa-nhr-10* does not affect the overall morphology of the adult mouth. A geometric morphometric analysis 2-dimensional landmark data from 144 individual worms (48 per strain) showed very homogenous and highly overlapping ranges of morphological variation in mouth form across wild-type worms and *Ppa-nhr-10* mutants (Fig 3C). A principal component analysis (PCA) of our form data revealed that, together, PC1 and PC2 capture the majority of form variation in the data (60.2%). Additionally, we were able to estimate that PC1 and PC2 are the only "meaningful" principal components *sensu* Schlager et al. [40,41] and that the other PCs are likely to contain

noise rather than signal. Subsequent model-based clustering of all individuals in the form dataset (based on the information contained in the "meaningful" PCs) revealed that mutant worms could not be classified as such, or differentiated from wild-type worms, based on their mouth morphologies alone (Fig 3D). Together, our geometric morphometric and model-based clustering analyses support the absence of any discernible mutant phenotypes in the mouth of *Ppa-nhr-10* mutants. Again, these findings are in blatant contrast to the phenotypes caused by the loss *Ppa-nhr-1,* which leads to structural alterations of the mouth and gives rise to worms with an intermediary mouth morphology combining anatomical features of predators and bacterivors [24].

Taken together, the results of our morphological analyses do not support our initial hypothesis that *Ppa*-NHR-10 is another constituent of the gene regulatory network that governs mouth-form plasticity in *P. pacificus*. This also indicates that not the entire NHR-1/-10/-40 cluster was co-opted into said functional context during diplogastrid evolution. Intrigued by this observation, we set out to elucidate the biological function of NHR-10 in *P. pacificus,* starting with exploratory RNA-seq experiments.

Exploratory transcriptomics suggests *Ppa*-NHR-10 might be a regulator of lipid catabolism and transport

Since nuclear receptors are transcription factors, it is possible to identify their potential regulatory target genes, and thus infer their potential biological functions, through comparative transcriptomics. We analyzed the transcriptomes of wild-type worms (PS312), and *Ppa-nhr-10(lof)* mutants (RS3920 and RS3921) at two developmental stages (larvae and adults). We aimed to identify potential "core targets" of *Ppa*-NHR-10 which we considered to be genes that are consistently mis-regulated (i.e., differentially expressed) in both mutant strains, in at least one of the two developmental stages.

In doing so, we were able to identify 32 potential regulatory target genes, of which 25 were down-regulated and seven up-regulated (Fig 4). We found 11 genes consistently differentially expressed in mutant larvae and 12 in mutant adults (compared to their wild-type counterparts). The larval targets include multiple down-regulated collagens and up-regulated proteins which are known to be involved in stress responses, like C-type lectins or the UDP-glycosyltransferases (Fig 4). Interestingly, we also found two proteins, both of them strongly down-regulated, which are predicted to be involved in the binding and transport of lipids: PPA13553 (an ortholog of the *C. elegans* vitellogenin *vit-6*) and PPA05630 (a predicted apolipoprotein). On the other hand, all of the 12 targets we identified in adult mutants were down-regulated and they include a more diverse array of proteins, ranging from a membrane-trafficking protein (PPA08133/*Ppa-scm-1*), to a component of the RNA polymerase II complex (PPA14053/*Ppa-lin-25*), to a chromatin organization modifier (ppa_stranded_DN31598_c1_g4_i2), to one of the TGF-*β* signaling receptors (PPA15180/*Ppa-daf-4*). The latter is particularly interesting, as TGF-*β* signaling in *C. elegans* is involved in both the formation of the dauer larvae and the regulation of body fat [42,43].

Strikingly, we found nine putative target genes which are strongly and constitutively mis-regulated across both developmental stages in *Ppa-nhr-10(lof)* mutants, and the majority of them have predicted functions in lipid catabolism. Briefly, canonical lipid catabolism involves three processes: (I) lipolysis - the cleavage of neutral storage lipids (TAGs) into glycerol and free fatty acids by lipases, (II) activation - the ligation of CoA to free fatty acids by fatty acyl-CoA synthetases (ACSs), and (III) *β*-oxidation - the breakdown of activated fatty acids into acetyl-CoA molecules. The latter process is generally performed in four

steps: dehydrogenation by acyl-CoA dehydrogenases (ACDHs), hydration by enoyl-CoA hydratases (ECHs), dehydrogenation by 3-hydroxyacyl CoA dehydrogenases (3-HACDs), and thiolytic cleavage by thiolases [5,43,44]. Most excitingly, the constitutively mis-regulated putative targets of *Ppa*-NHR-10 include two predicted lipases (PPA07082 and Iso_D.954.1), two predicted ACSs (PPA05769 and PPA13813), as well as a predicted ECH (PPA40903). Furthermore, these genes are among the most strongly affected by the loss of *Ppa-nhr-10.* This is reflected in the fact that they show some of the largest L2FC-estimates among all putative regulatory targets (Fig 4).

Taken together, our exploratory transcriptome and differential expression analyses suggest that *Ppa*-NHR-10 might be an important regulator of body fat in *P. pacificus.* This is evident in the observation that the loss of *Ppa-nhr-10* negatively affects the expression of genes predicted to be involved in all the major processes of lipid catabolism, as well as genes encoding proteins required for the transport of lipids across tissues. Given that lipid metabolism and life span are tightly linked in all animals [45], and knowing that functional interferences with any of the lipid catabolic processes in nematodes can impair their survival capabilities across various environmental conditions [43,44], we decided to quantify the survival capabilities of our *Ppa-nhr-10(lof)* mutants in different conditions, expecting to find that they are unlike those of wild-type worms.

Survival analyses reveal that *Ppa-nhr-10* is required for survival of acute starvation in young adults

In *C. elegans*, interference with the major lipid catabolic pathways can lead to altered life spans and/or rates of survival over time. While some mutations in these pathways constitutively impair the worm's survival capabilities, irrespective of environmental factors, others are known to cause such impairments only in certain conditions like starvation [43-45]. Minding these details, we decided to examine the survival capabilities of wild-type *P. pacificus* and *Ppa-nhr-10(lof)* mutants in two different conditions: well-fed and acutely-starved. To achieve this, we transferred young adult hermaphrodites into small cell culture flasks which either contained a standard liquid culture (LC) medium rich in OP50, or a LC medium that contained no food. To obtain the survival curves, we scored the number of survivors every 24hrs until all worms in the flasks were dead.

First, we found that the obtained time-to-death data from the well-fed condition is compatible with the hypothesis that there are no differences in the overall survival capabilities of *Ppa-nhr-10(lof)* mutants and wild-type worms (Fig 5A and 5C). Neither the global Kaplan-Meier (K-M) survival analysis of well-fed worms (Fig 5C: $\chi^2 = 5.7$, $P = 0.060431$), nor the pairwise comparisons of strain-specific survival curves via a log-rank test, revealed any significant differences in survival (Fig 5A). Yet, we found that hazard ratios were estimated to be slightly elevated in the mutant strains compared to the wild-type level of 1.0 (1.15 for RS3920 and 1.13 for RS3921). However, while these hazard ratios do equate to a ~15% increase in the relative risk of death at any given timepoint in the experiment, these estimates were still found to be compatible with the null hypothesis given their associated uncertainty (Fig 5C).

Remarkably, we found the exact opposite to be true for our starvation experiments: the global K-M analysis indicated stark differences in the overall survival probabilities among starving worms (Fig 5C: $\chi^2 = 175$, $P = 1.51e-38$) and the log-rank pairwise comparisons revealed that the survival curves of both *Ppa-nhr-10(lof)* mutants, while being virtually identical to each other (Fig 5B: $P = 1.0$), differed markedly from the wild-type

survival curve (Fig 5B: $P <$ 2e-16). Additionally, we found highly elevated hazard ratios for the *Ppa-nhr-10(lof)* mutants compared to wild-type *P. pacificus* (Fig 5C: 2.23 for RS3920 and 2.12 for RS3921). Simply, these estimates mean that for every wild-type worm that dies at any given point of time during the starvation experiment, more than two mutant worms will face the same fate. These large effects are incompatible with our null hypothesis (P[RS3920]= 4.79e-30 and P[RS3921]= 3.91e-27) and strongly support decreased survival rates in mutant worms (Fig 5C).

Taken together, our condition-specific survival experiments revealed that acute starvation induces highly elevated rates of premature deaths in young adult worms which lack a functional copy of *Ppa-nhr-10*. While this observation confirmed the hypothesis we formulated based on our exploratory RNA-seq results, one might now want to argue that, given the condition-specificity of the identified phenotype, our initial sequencing approach was, in fact, not ideal. This is because all the worms we used for the exploratory RNA-seq analysis were well-fed rather than starved. This means that, amongst all of the identified putative targets which potentially cause the survival phenotype in starving mutants, the exploratory analysis could have picked up only on those genes which are constitutively mis-regulated (i.e., in well-fed *and* acutely-starved conditions). Therefore, we potentially missed out on a pool of putative target genes, which are mis-regulated in mutants specifically *in response to* starvation. Obviously, such genes would be important to explain the starvation-specific survival phenotypes and, if they exist, this could argue that *Ppa*-NHR-10 itself is a regulator of the starvation-response machinery in these nematodes. In order to expand our analysis as to include genes that are mis-regulated in mutants responding to starvation, and to elucidate whether any of *Ppa*-NHR-10's putative targets are members of the worm's starvation-response machinery, we performed confirmatory RNA-seq experiments which included both, well-fed and acutely-starved, wild-type and mutant worms from our LC set-up.

Starvation induces substantial transcriptional responses in genes predicted to be involved in metabolism, stress responses, and translation

To begin with, we aimed to elucidate genome-wide changes to the transcriptional landscape in wild-type worms which faced 24hrs of acute starvation. Using our differential expression pipeline, we were able to identify a total of 1,007 genes showing a response to starvation (Fig 6A). Of these, 294 genes were notably up-regulated (cluster I in Fig 6) and 713 notably down-regulated (cluster II in Fig 6). Thus, our data shows, for most of these starvation-responsive genes (SRGs), the worms strongly decrease expression levels compared to when food is abundantly available.

Next, we wondered what kinds of genes are up- or down-regulated. Are the SRGs predominately diplogastrid-specific orphan genes, or phylogenetically old? Are they members of any known co-expression modules? What kind of proteins do they encode and are there biases towards specific protein classes responding to starvation? Lastly, do the SRGs fall into any specific biological pathways? To address these questions, we mined a wealth of available data through overrepresentation analyses (ORAs).

First, we found that phylogenetically old genes, rather than diplogastrid-specific orphans, are overrepresented in both SRG clusters (Fig 6B). While these observed effects are well-supported by our data (q[cluster I] = 3.7e-04 and q[cluster II] = 1.8e-05), their sizes are comparatively small (fold enrichment: cluster I = 1.15 and cluster II = 1.12). However, while

they do constitute the minority of SRGs, there are still hundreds of orphan genes functionally responding to starvation (Fig 6B).

Next, we asked whether any of the co-expression modules that were recently identified in a nutrigenomic study [46] were overrepresented among the SRGs. Indeed, genes which belong to the modules "oogenesis 3", "nervous system 2", "nervous system 3", and "nervous system 5" where strongly enriched amongst the up-regulated SRGs. On the other hand, members of the co-expression modules "orphan 3", "cuticle 1", "nervous system 4", "intestine 1", "intestine 3", and "intestine 4" where strongly enriched among the down-regulated SRGs (Fig 6B).

To identify the potential biological functions of the SRGs, we predicted the types of protein domains (PFAM [47]) that they encode and, additionally, we functionally annotated the SRGs with WormCat (ver. 2) terms [48] wherever a single ortholog was available in *C. elegans*. Amongst the up-regulated SRGs, there is a clear bias towards genes that encode proteins which are critical for the translational process and protein synthesis (Fig 6C). This includes large ribosomal subunit proteins (Ribosomal_L6e), elongation factors (GTP_EFTU, GTP_EFTU_D2), and tRNA synthetases (tRNA-synt_1, tRNA-synt_1c). This pattern was further confirmed by our ORA of WormCat terms, which found only a single major category enriched among the up-regulated SRGs: "Ribosome" (Fig 6D). This category contains genes that encode proteins and RNA which constitute, assemble, or regulate the ribosome and translational processes [48]. Within the ribosome category, we further found an overrepresentation of genes belonging to the "biogenesis" subcategory (q = 1.1e-07), which encode proteins acting in ribosome assembly [48] (Fig 6D).

Genes which are down-regulated in response to acute starvation seem to execute entirely different biological functions. We found a clear bias towards genes that encode proteins involved in diverse metabolic pathways (q = 2.4e-11), as well as stress responses (q = 1.7e-05) (Fig 6D). The strongest detected metabolic signal was found for lipid metabolism (q = 9.3e-11), followed by amino acid metabolism (q = 6.7e-07), and the one-carbon cycle (q = 0.0203). We further detected overrepresentation signals for specific aspects of these individual metabolic pathways. This includes genes which encode proteins involved in the elongation and desaturation of fatty acids ("fatty acid"), lipid binding, as well as in the breakdown and synthesis of amino acids, and the conversion of homocysteine to glutathione ("transsulfuration") [48] (Fig 6D). The strong overrepresentation pattern for lipid metabolic pathways, was further corroborated by our ORA of PFAM domains, which revealed strong enrichments for lipases (Lipase_GDSL, Lipase_GDSL_2), fatty acid elongases (ELO), fatty acid desaturases (FA_desaturase), and lipid-binding vitellogenins (Vitellogenin_N) (Fig 6C). The second-most overrepresented functional category among down-regulated SRGs - stress response - is based on the presence of multiple C-type lectins, p450 cytochromes, and lysozymes in cluster II (Fig 6C and 6D). These genes are involved in the worm's immune response, the detoxification of xenobiotic compounds, and the survival of pathogenic stresses, respectively [48].

Taken together, irrespective of the direction of the effect, genes which transcriptionally respond to acute starvation are mostly phylogenetically old (Fig 6B). Interestingly, genes which are up-regulated during starvation are predominantly members of various nervous system and oogenesis related co-expression modules, and they tend to encode proteins that are crucial constituents of the translational machinery. In contrast, genes that are down-regulated in response to starvation are predominantly members of various intestinal co-expression modules and tend to encode proteins that are crucial

enzymes in lipid and amino acids metabolism, as well as important detoxification and immune response elements (Fig 6).

With this reference knowledge of genome-wide patterns of starvation responses in hand, we next set out two analyze the transcriptomes of starved *Ppa-nhr-10(lof)* mutants to see which genes are likely to explain the starvation-induced survival phenotype, and whether any of them are members of the identified SRGs.

Confirmatory transcriptome analyses indicates that *Ppa*-NHR-10 regulates aspects of the wild-type starvation response, lipid metabolism, and TGF-β signaling

Our confirmatory transcriptome analysis revealed a majorly updated list of 37 putative *Ppa*-NHR-10 targets, of which 32 were down-regulated and five up-regulated (Fig 7). To validate that these statistically inferred candidate genes are indeed appropriate to address our questions, we submitted their *rlog*-transformed [49] normalized count values to unsupervised hierarchical clustering. Ideally, these candidate genes should capture the information needed to differentiate well-fed from starved, and wild-type from mutant worms. Thus, the clustering algorithm should be able to recover these experimental groupings from the candidate gene data alone. This was indeed the case (Sup Fig 1), confirming that our improved list of putative regulatory targets is appropriate to address the observed starvation-survival phenotype in mutant worms, and to identify any putative targets that are starvation-responsive (i.e., SRGs).

As expected, our original exploratory RNA-seq experiments missed a majority of putative causal targets (28 out of 37). Unfortunately, many of the newly identified target genes have a wide variety of predicted functions which are not easily grouped into overarching processes (Fig 7). Strikingly, however, we recovered nine of the putative *Ppa*-NHR-10 targets that were already found in the exploratory analysis (Fig 4 and Fig 7), including the TGF-β receptor PPA15180/*Ppa-daf-4*, the p450 cytochrome PPA25752/*Ppa-cyp-33B1*, and the two lipid catabolic enzymes PPA05769/*Ppa-acs-14* and PPA40903 (Fig 4 and Fig 7). This confirms that, while our initial RNA-seq approach was not ideal, it was still able to identify some of the putative phenotype-causing targets of *Ppa*-NHR-10.

More importantly, our confirmatory RNA-seq approach did actually confirm the initially observed pattern: a bias toward regulatory targets that are involved in the transport and break of lipids and fatty acids (Fig 7). PPA33234/*Ppa-strl-1* is a lipid-binding protein predicted to be located in mitochondria and involved in the transport of cholesterol for steroid biosynthesis. PPA27650 is predicted to be involved in intermembrane transport of lipids as part of glycerolipid synthesis. Genes that encode important players in the β-oxidation of fatty acids were recovered, too: PPA05769/*Ppa-acs-14* is a fatty acyl-CoA synthetase, the type of enzyme that "activates" free fatty acids for β-oxidation. Additionally, the gene *ppa_stranded_DN29959_c0_g2_i3* encodes an acyl-CoA dehydrogenase (ACDH) and PPA40903 an enoyl-CoA hydratase (ECH). These two enzymes catalyze the first two of the four canonical steps in mitochondrial β-oxidation [43-45].

Lastly, we wondered what the gene regulatory basis of the starvation-survival phenotype in *Ppa-nhr-10(lof)* mutants looks like and anticipated two different scenarios. One, that NHR-10's regulatory targets would be members of genetic networks which are not part of the starvation response machinery (Fig 6), but might, when mis-regulated, lead to decreased survival capabilities in mutant worms that face starvation. Alternatively, *Ppa*-NHR-10's regulatory targets could actually constitute members of the SRGs. In this latter scenario, the impaired survival capabilities of starving mutants would be directly explained

by malfunctions in the starvation-response machinery itself. Surprisingly, our analysis suggested a combination of both: while the majority of putative *Ppa*-NHR-10 targets are genes that do not respond to starvation, we were still able to identify four SRGs among them (Fig 7). Strikingly, these four SRG targets included the two aforementioned *β*-oxidation enzymes (ACDH and ECH), of which one was already found in our exploratory RNA-seq analysis; an observation that also applies to another SRG target: the p450 cytochrome PPA25752/*Ppa-cyp-33B1* (Fig 7).

Taken together, this new data corroborated our original hypothesis that *Ppa*-NHR-10 controls general aspects of fatty acid breakdown and, additionally, demonstrated that this supplementary nuclear receptor is required for some of the worm's transcriptional responses to acute starvation.

Dauer assays demonstrate that the loss of *Ppa-nhr-10* also affects dauer entry rates, dauer exit speed, and long-term dauer survival

Finally, we wondered whether *Ppa*-NHR-10 might play a more general regulatory role in starvation survival beyond its requirement for starvation resistance in young adult worms. Thus, we investigated the potential effects that the loss of this receptor might have on a different life history stage: dauer larvae (Fig 1B). We hypothesized to see effects on dauers for multiple reasons: First, they represent the worm's natural state of starvation, as their buccal plugs prevent them from taking in any food (Fig 1C) [6,28]. Second, it is well established that dauer metabolism in *C. elegans* is heavily dependent on the generation of energy from stored neutral lipids via lipolysis and fatty acid *β*-oxidation [50-53]. Third, it is known that various NHRs are involved in the regulation of dauer entry, dauer exit, and metabolism [31,42-44]. Taking all these factors together, while integrating our findings that *Ppa*-NHR-10 is required for starvation survival in young adults and that it seems to regulate some of the very aspects of lipid breakdown, which also constitute the foundation of dauer metabolism, we were optimistic to find alterations in dauer larvae of our mutant worms.

We quantified three different aspects of dauer biology: i) the number of worms which entered the dauer stage due to overcrowding, ii) the time it took the worms to exit dauer and resume development when re-introduced to food, and iii) the survival of dauers over extended periods of time (Fig 8A).

Interestingly, for dauer entry, we found an allele-specific phenotype in only one of our *Ppa-nhr-10(lof)* mutants: RS3920(*tu1654*). Across culture conditions, worms belonging to this strain showed a decreased number of dauer larvae in the overall population when compared to the wild-type and the second mutant strain (Fig 8B). However, such allele-specific phenotypes could not be observed in the other dauer assays. In fact, just like in all of our previous experiments, the two mutant strains behaved in an almost indistinguishable manner (Fig 8C and 8D). Here, our data demonstrated that animals lacking a functional copy of *Ppa-nhr-10* needed considerably more time to exit the dauer stage and resume normal development (Fig 8C). Where more than 80% of wild-type worms already exited dauer within the first 24hrs after being re-introduced to food, less than 5% of mutant worms managed to do the same (4.6% in RS3920 and 2.9% in RS3921). Only after another 24hrs on food did the mutants reach somewhat similar numbers of exited worms (70.8% in RS3920 and 75.0% in RS3921), at which point virtually all of the wild-type worms had resumed normal development (99.2% in PS312). While eventually all investigated worms completed the dauer exit (and developed into fully viable adults), it took the mutants twice as long as wildtype to do so (Fig 8C).

Excitingly, we found impaired long-term survival capabilities of mutant dauer larvae (Fig 8D). Initially, after spending another week in dauer following the SDS treatment, the estimated overall survival rates of wild-type and mutant dauers were indistinguishable (53.5% in PS312, 56.5% in RS3920, and 52.5% for RS3921). However, this changed during the second week post-SDS exposure: where wild-type worms showed estimated survival rates which were indistinguishable from those of the previous week (58%), our mutant worms died off rapidly, leaving almost no survivors (1.7% in RS3920 and 6.0% for RS3921) (Fig 8D). This trend continued into the third week, where mutant dauers were practically extinct (0.0% survivors in RS3920 and 0.3% in RS3921), while 19.3% of wild-type dauers were still alive. In summary, these assays revealed that all of the investigated aspects of dauer biology were negatively affected by the loss of *Ppa-nhr-10*. This observation confirmed the hypothesis we conceived based on the combined results from our survival analysis and our confirmatory RNA-seq data, and they demonstrate that *Ppa*-NHR-10 is indeed required in the broader context of starvation resistance throughout the *P. pacificus* life cycle, including the ecologically relevant dauer stage.

Discussion

In this study, we aimed to demonstrate that *Ppa*-NHR-10 is a regulator of mouth-form plasticity in *P. pacificus* and, thus, validate the hypothesis that a small but complete cluster of three closely-related and deeply-conserved supplementary nuclear receptors (*supnrs*) was functionally co-opted into the gene regulatory network that governs plasticity in the feeding structures of *Pristionchus* nematodes. We were able to demonstrate that *Ppa*-NHR-10 is indeed, amongst others, expressed in relevant tissues which are part of the morphological novelties in diplogastrid mouths (e.g., the g1D pharyngeal gland cell associated with the dorsal tooth). This is intriguing, as *Ppa*-NHR-1 and virtually all of the common regulatory targets of *Ppa*-NHR-1 and *Ppa*-NHR-40 are co-expressed in the same cell [24]. Yet, unlike mutations in the latter two receptors, loss of *Ppa*-NHR-10 neither disrupted the worm's ability to develop into predators, nor did it lead to aberrant mouth morphologies. Thus, these findings were incompatible with our initial hypothesis. Instead, our results indicate that it was specifically *Ppa*-NHR-1 and *Ppa*-NHR-40, but not the complete NHR-1/-10/-40 cluster, that were functionally co-opted when the plasticity GRN was assembled during the evolution of mouth morphological novelty in diplogastrids. This also demonstrates that the genetic basis for the evolution of plastic traits and novel structures is more complicated than often presumed, and that paralogs - at least when it comes to nematode *supnrs* - might not necessarily be adequate predictors for the functions of the genes of interest.

Through a combination of exploratory and confirmatory mRNA-seq experiments, we were able to identify *Ppa*-NHR-10's putative regulatory targets. These encompass a broad variety of proteins including many metabolic enzymes and multiple signaling elements (Fig 4 and Fig 7). Interestingly, this pattern is completely different from the regulatory target patterns of its paralogs *Ppa*-NHR-1 and *Ppa*-NHR-40: most of their targets belong to a single class of metalloproteases (astacins) that is crucial for cuticle degradation [24] and the vast majority of them is located on the X-chromosome. This is the same chromosome that also harbors *Ppa-nhr-1* and *Ppa-nhr-40*, as well as the multi-gene locus that regulates the plastic switch decision [23,24]. This pattern, again, does not hold true for the loci harboring *Ppa*-NHR-10's targets, which do not display a bias towards one particular chromosome, let alone chromosome X (Sup Tab 1). In contrast to *Ppa*-NHR-1's and *Ppa*-NHR-40's biases towards regulating a specific class of enzyme, *Ppa*-NHR-10 seems to regulate a defined biological process that involves a variety of different proteins: the activation and β-oxidation of fatty acids (Fig 4 and Fig 7). The differences in the regulatory "behaviors" among these closely-related NHRs demonstrate why, in this particular case, *Ppa*-NHR-10's paralogs were inadequate predictors of function.

Most evolutionary biologists would argue that this is not surprising and reference classical literature in the field, which teems with examples of deeply-conserved "master regulators" of development and physiology, which execute "the same" function in the most distantly related animals [54,55]. After all, they may say, orthologs are more functionally similar to one another than paralogs are [56,57]. Yet, in more recent years, researchers increasingly called into question the notion that orthologs are the better, or maybe even the only viable, homologs to turn to, if we want to predict the function of a gene of interest in a unstudied organism [58-60]. This assumption, the "ortholog conjecture" [58], so say the contenders, is no longer sustainable given the vast amount of bioinformatic evidence [59,60]. In fact, some recent studies demonstrated that paralogs can outperform orthologs in

functional inference tasks, and that the cumulative "big data" evidence suggests orthologs and paralogs should be considered together when predicting biological functions for genes of interest [60]. Thus, our example of *Ppa*-NHR-10 not being involved in mouth-form plasticity, of which both of its closest paralogs are crucial regulators, is indeed surprising.

Stunningly, predicting *Ppa*-NHR-10's function would not have been much more successful if we had looked at its well-studied one-to-one ortholog in *C. elegans* instead. *Cel*-NHR-10 is known to regulate the expression levels of genes which encode enzymes that constitute a nematode-specific metabolic pathway called the "propionate shunt" [61]. This shunt is based on five genes (*acdh-1*, *ech-6*, *hach-1*, *hphd-1*, and *alh-8*), all but one of which have a clear one-to-one ortholog in *P. pacificus*. While four out of five propionate shunt genes in *C. elegans* show a strong mis-regulation upon NHR-10 inactivation [61], not a single one of the propionate shunt orthologs in *P. pacificus* displayed altered expression levels in strains that lacked a functional copy of *Ppa*-NHR-10 (Sup Tab 2). Therefore, neither the closest paralogs of *Ppa*-NHR-10 (*Ppa*-NHR-1 and *Ppa*-NHR-40), nor its one-to-one ortholog in *C. elegans* (*Cel*-NHR-10), were suitable predictors of its starvation-resistance related function in *P. pacificus*. This indicates, when it comes to nematode-specific *supnrs*, that caution must be exerted when assigning biological functions to receptors based on homology, rather than on functional genetic investigations. In fact, the data presented in this study suggest that even the very few *supnrs* which are deeply-conserved across distantly-related species might regulate entirely lineage-specific functions, potentially due to rapid evolutionary turnover in the acquisition of regulatory targets. Future comparative studies in *C. elegans* and *P. pacificus* could directly test this hypothesis for other conserved *supnrs*, e.g. NHR-1 and NHR-40, as their biological functions are already known in one of the species [24].

We also elucidated genome-wide transcriptional responses to 24hrs of acute starvation in young adult hermaphrodites of *P. pacificus*. When focusing on the broad-scale patterns in this data, three interesting observations emerge:

First, during starvation, *P. pacificus* increases the expression of numerous genes which are critical elements of the translational machinery, and it decreases the expression of metabolic pathway genes, especially affecting lipid transport, and the desaturation and elongation of fatty acids for fat storage purposes (Fig 6C and 6D). Thus, the biological pathways affected by acute starvation are consistent between *P. pacificus* and *C. elegans* [6,62-64], indicating that the underlying starvation-response mechanisms are deeply conserved; an observation that is also reflected in the fact that most of the SRGs are phylogenetically old (Fig 6B). Yet differences exist in the details: when *C. elegans* starves, it does not change the expression levels of ribosome-biogenesis genes, but it decreases the translation of their transcripts [64]. We speculate that *P. pacificus* might employ a slightly different strategy, in which it increases the expression of translation-related genes, while it does not increase the translation of their transcripts. This way, the worms would not invest large amounts of energy into the protein synthesis [6], while maintaining the ability to quickly do so if food becomes available again. Once scalable proteomic methods become available for *P. pacificus*, future studies will be able to directly test this hypothesis and study the evolution of these molecular starvation-response strategies.

Second, the majority of up-regulated genes are associated with the nervous system and oogenesis, while the vast majority of the down-regulated genes are associated with the metabolic hub of nematodes, the intestine (Fig 6B). Additionally, we found numerous down-regulated genes to be members of a co-expression module which is associated with the hypodermis (Fig 6B "cuticle 1" [46]), the site of long-term lipid storage [44]. Together with the observation that lipid anabolism is down-regulated during starvation, these data

could suggest that *P. pacificus*, when facing acute starvation as adults, might undergo drastic changes to their neural regulatory landscape which may, in turn, lead the worms to switch strategies from synthesizing and storing lipids to maintain their soma, to investing into egg production and ensuring population survival. This would be in line with recent predictions concerning neural regulation of body fat during starvation in *C. elegans* [44].

Third, a variety of stress responses are down-regulated in the absence of bacterial food. This includes multiple proteins predicted to be involved in the detoxification of xenobiotic compounds (p450 cytochromes), the immune response (C-type lectins), and the defense against microbial pathogens (lysozymes) [48] (Fig 6D). This corroborates an interesting observation: the uracil-auxotrophic OP50 strain of *E. coli*, the standard laboratory food source for nematodes [65], might be slightly toxic to *P. pacificus* [66]. Upon ingesting these bacteria, the worms may have to activate some of these stress response pathways in order to be able to sustainably live off of them as a diet. Taken together, this newly-established reference knowledge of responses to acute starvation complements recent nutrigenomic advances [46] in *P. pacificus*, and it will inform future studies which may aim to evaluate the quality of different diets. If the worm's transcriptional responses to a specific bacterial food source are highly similar to the starvation responses reported in this study, this could demonstrate a poor nutritional value of a given bacterial diet.

The present study also demonstrates that *Ppa*-NHR-10 is broadly required for starvation resistance in *P. pacificus* (Fig 5 and Fig 8) and that some of its regulatory targets are starvation-responsive (Fig 7). This indicates that *Ppa*-NHR-10 is one of the probably many transcriptional regulators that contribute to the wild-type starvation response. Interestingly, the expression levels of the *Ppa-nhr-10* gene itself do not change during acute starvation (l2fc = -0.10894 and *s* = 0.79898), suggesting that *Ppa*-NHR-10 is already present before the starvation signal is sensed. It thus seems likely that, in the context of the adult starvation response, *Ppa*-NHR-10 is activated by a ligand that is present during starvation, or inactivated by the absence of its ligand during starvation. Either way, future biochemical studies may address the activity states of this receptor and identify its ligand. This will lead to a better understanding of *Ppa*-NHR-10's role in the starvation response, and elucidate how a food-depletion signal is detected and converted into a biological response via this *supnr*.

Lastly, we found that the loss of *Ppa*-NHR-10 negatively affected multiple aspects of dauer biology, using a unique method combining several assays. This allowed us to investigate three pertinent dauer characteristics - entry, exit, and survival - from the same initial population of dauers in conditions similar to their ecology. In its natural environment, *Pristionchus* is found primarily as dauers in soil and on its invertebrate hosts, predominantly beetles [20,30]. The ability to enter dauer is vital to ensure success in the wild, as diminishing food resources are a common occurrence. Worms that do not enter dauer under unfavorable conditions risk preserving their populations. In addition to persevering through unpredictable food availability, *Pristionchus* faces interspecific competition [67,68]. The speed of dauer exit may determine the success of a population if cohabitating with another nematode species, as first to the feast may establish, repopulate, and consume all accessible bacteria [68]. Long-term survival in dauer provides protection during extended periods of stressful conditions [30], and may play a role in evading competition while waiting for microbial populations to reestablish [68].

Therefore, starvation resistance is a trait that is crucial for the fitness of these worms in the wild: adults need to adjust to starvation and invest into the production of viable eggs, allowing future generations of worms to persist; dauers need to be able to survive over

extended periods of time, allowing them to disperse and establish new populations of reproducing worms under suitable conditions. Here, we demonstrated that the supplementary nuclear receptor *Ppa*-NHR-10 contributes to the transcriptional regulation of starvation responses in *P. pacificus*, and that it is required for starvation resistance in adults and dauer larvae. Based on our transcriptomic experiments and our survival data, we propose that the underlying mechanism through which this is achieved is based on the activation and β-oxidation of fatty acids. Future molecular genetic investigations will be able to directly test this hypothesis based on the candidate targets we identified in our experiments, ultimately expanding our understanding of the physiological and genetic mechanisms that underlie starvation resistance in these nematodes.

Material & methods

Nematode husbandry

We followed standard protocols for the establishment and maintenance of laboratory cultures of rhabditid and diplogastrid nematodes [65]. Briefly, nematodes of all strains used in this study were cultured on nematode growth medium (NGM) in 6cm Petri dishes, using the uracil-auxotrophic strain "OP50" of *Escherichia coli* as a food source. Bacteria were grown overnight at 37°C in LB medium. We generally added $300\mu l$ of fresh bacterial food solution to each Petri dish and left these plates at room temperature for several days, in order to allow the bacterial lawns to grow. Usually, five gravid adult hermaphrodites were transferred to these fresh plates to maintain cultures. Culture plates were stored in plastic boxes and kept in an incubator set to a constant 20°C.

Light microscopy with differential interference contrast (DIC) and morph ratios

For microscopy, all specimens were mounted on object slides with 5% Noble Agar pads which contained the sedative sodium azide (0.5% NaN_3) and subsequently examined using a Zeiss Axio Imager.Z1 microscope with a Zeiss Plan-Apochromate 100 x 1.4 DIC objective. Image stacks (Grayscale, 16-bit) were taken using a monochromatic Zeiss Axiocam 506 CCD camera. The Zen 2 Pro Software (Carl Zeiss Microscopy GmbH, 2011; version 2.0.14283.302; 64-bit) for digital microscopic analysis and image acquisition.

Relative ratios of eurystomaous (Eu) and stenostomatous (St) morphologies in each strain were scored using the aforementioned microscopy settings. Worms that had a large right ventrosublateral tooth, a hook-shaped dorsal tooth, and a promesostegostom whose anterior tips were clearly posterior to those of the gymnostom were classified as Eu morphs; animals that did not meet these combined characteristics were scored as St morphs. Statistical differences in morph ratios were assessed via a PERMANOVA (see "Dauer assays" section of the methods for details).

Quantification of morphological differences in mutant animals

We quantified morphological differences in the mouths of wild-type and mutant worms, using our recently published protocol [69]. This approach combines 2-dimensional landmark-based geometric morphometrics (GMM) with unsupervised clustering methods.

For this study, we obtained image stacks of the nematode mouths in lateral position and recorded the X and Y coordinates of 18 fixed landmarks which capture the overall form of the mouth using FIJI (ver. 2.1.0) [70]. All steps of the GMM analysis were performed in R (ver. 4.3.0) [71] using a combination of the GEOMORPH (ver. 4.0.5) [72-74], MORPHO (ver. 2.11) [40,41], and MCLUST (ver. 6.0.0.) packages [75,76]. General Procrustes Analysis (GPA) was performed with the *gpagen* function of GEOMORPH, by minimizing Procrustes distances. Procrustes coordinates were projected into a linear tangent space in order to obtain Euclidean distances for downstream statistical analysis. We originally generated landmark data for 150 animals (50 per strain; three strains: PS312, RS3920, RS3921), performed a GPA and subsequently checked for Procrustes distance outliers in our original shape data. Outliers were identified separately for each of the strains of interest, using the *plotOutliers* function of GEOMORPH. Six extreme outliers were identified and removed from the original data to facilitate downstream analysis. The outlier-corrected data set (n=144) thus contained landmark configurations of 48 animals per strain. This dataset was used to quantify

differences in mouth form and to see whether mutant animals can be classified as such, based on the structure of their mouths. In morphometric terms, form refers to all morphological variation in the data, i.e. the variation in both shape and size [77]. As form data takes into account the full spectrum of structural variation between the groups of interest, it generally outperforms pure shape or size datasets for classification tasks such as ours [78].

We generated form data by first performing a GPA on our outlier-corrected landmark dataset to obtain a shape matrix to which subsequently appended a size variable (here, the log-transformed centroid size). We performed a principal component analysis (PCA) on the form data to visualize potential morphological differences among wild-type and mutant worms in a form space *sensu* Mitteroecker *et al.* [79]. We tested whether the form data contains clearly separable groups of specimens with similar morphology, ideally ones which correspond to the different strains. For this, we performed model-based clustering with the *Mclust* function of the MCLUST package. We only used "meaningful" principal components (mPCs) of form variation (identified with MORPHO's *getMeaningfulPCs* function [40,41]) as input variables for unsupervised clustering, in order to avoid overparameterization of our multivariate normal models [69]. Finally, we visualized the clustering results in the form space (i.e., the form PCA plot) by coloring each specimen according to the identified cluster to which it belongs.

CRISPR/CAS9 mutagenesis

We used the CRISPR/CAS9 system to create small "indel" mutations in the coding sequences of *Ppa-nhr-10*, specifically in the part of the gene that encodes the ligand-binding domain (LBD) of *Ppa*-NHR-10. For this we followed established mutagenesis protocols for model nematodes [80,81]. Target-specific CRISPR RNAs (crRNAs) were generated based on the publicly available genome sequence of *P. pacificus* (www.pristionchus.org; El Paco ver. 3) and designed to target 20bp upstream of a 5'-NGG-3'-protospacer adjacent motif (PAM). The CAS9 protein (cat.nr.: 1081058; IDT), the universal trans-acting CRISPR RNA (tracrRNA), and all crRNAs were ordered from IDT (Alt-R product line). Ribonucleoprotein complexes (RNPs) were created by combining $0.5\mu l$ of CAS9 nuclease ($10\mu g/ml$ stock) with $5\mu l$ of tracrRNA ($0.4\mu g/\mu l$ stock) and $2.8\mu l$ of crRNA ($0.4\mu g/\mu l$ stock). This mix was incubated at 37°C for 15min and subsequently cooled down to room temperature. The RNP mix was diluted by adding enough Tris-EDTA buffer to bing the total volume of the injection mix to $20\mu l$. In order to avoid clogging in the injection needles, the final mix was centrifuged at 14,000rpm for 2min. Injections were conducted with an Eppendorf FemtoJet microinjector at 400x magnification under an inverse Zeiss AxioVert microscope that was equipped with a Zeiss Plan-Apochromat 40x DIC objective and coupled to an Eppendorf TransferMan micromanipulator. Injections were aimed at the distal tip of the hermaphrodite gonad in young adults; animals were punctured in a 45° angle to the anterior-posterior body axis and RNP complexes were injected into the gonadal rachis. Injected P0 animals were isolated onto separate NGM plates and allowed to lay eggs for 24h at 20°C. After that, P0s were removed from the plates and F1 animals were left to grow. Upon reaching maturity, F1s were singled out to fresh plates, allowed to lay eggs for 24h and subsequently used for genotyping. DNA of candidate F1 animals was recovered via single worm lysis (SWL). In short, individual worms were placed in $10\mu l$ of SWL buffer (10mM Tris-HCl at pH=8.3, 50mM KCl, 2.5mM $MgCl_2$, 0.45% NP-40, 0.45% Tween 20, $120\mu g/ml$ Proteinase K) and frozen at -80°C for 20min. Afterwards, all samples were

incubated in a thermocycler at 65°C for 1h, followed by a heat inactivation of the proteinase at 95°C for 10min. This lysate was used as a template for subsequent PCR amplification (using the Taq PCR Master Mix produced by Qiagen) of the region of interest. Potential mutations in candidate amplicons were identified using Sanger sequencing. Sanger sequencing was performed by Azenta Life Sciences. If heterozygous F1 mutants were found, their F2 offspring were genotyped in a similar fashion, to identify homozygous mutant carriers. Multiple homozygous mutant lines were isolated, including frameshift mutants with premature stop codons. Backups of the original homozygous mutant strains were frozen and stored at -80°C.

Transgenesis

To visualize which cells actively express *Ppa-nhr-10*, we generated transcriptional reporter lines. We PCR-amplified a 1.371 bp long putative promoter sequence, starting right before the start codon of *Ppa-nhr-10*, and subsequently cloned it into a pUC19 vector that contained a codon-optimized GFP with *P. pacificus* introns and the *Ppa-rpl-23* 3'UTR [82]. We generally used the NEBuilder HiFi DNA Assembly Kit for Gibson assemblies, *E. coli* (NEB 5-alpha) for bacterial transformation, and InVitrogen's PureLink HQ Mini Plasmid kit (including the optional, but essential, wash step with PB buffer) for plasmid extraction from 3ml bacterial overnight cultures. The *Ppa-nhr-10p::GFP* construct was linearized with SphI-HF (New England BioLabs) according to manufacturer's instructions. Following established protocols, we combined 10 ng/µl of our linearized plasmid were injected into the gonad of the wild-type strain of *P. pacificus* (PS312), together with 60 ng/µl of SphI-linearized genomic carrier DNA and 10 ng/µl of SphI-linearized co-injection marker *Ppa-egl-20p::RFP* [82,83]. We confirmed that expression patterns were consistent across two independent reporter lines (Fig 2A and 2B). Fluorescent worms were imaged using a Leica SP8 confocal laser-scanning microscope (cLSM).

Exploratory RNA-seq experiments - sequencing mutant larvae and adults

To facilitate *exploratory* RNA-seq analysis on our *Ppa-nhr-10* mutants comparable to the RNA-seq experiments our lab previously conducted on *Ppa-nhr-1* and *Ppa-nhr-40* mutants, we adopted (with minor modifications) the sample collection and sequencing approaches that were used by Sieriebriennikov *et al.* [24]. We first transferred seven gravid adult *P. pacificus* hermaphrodites to fresh agar plates that were spotted with 300µl of OP50 *E. coli*. These worms were kept at 20°C for 4-5 days, in order to obtain sufficient starting amounts of eggs (F2) laid by their F1 offspring. To isolate the eggs, we suspended all worms and eggs on an individual plate in 1ml water, collected such suspensions form multiple plates in 15ml Falcon tubes, and incubated these samples for 10min in a mix of 0.5M NaOH and household bleach (1:5 dilution). The Falcon tubes were vortexed every other minute to disintegrate the post-hatching life stages. The eggs were centrifuge-pelleted at 500g for 2min, resuspended in 5ml water, and pelleted again; this washing process was repeated three times. In the final step, we resuspended the egg pellets in 1ml water and subsequently pipetted 100µl onto fresh agar plates that were kept at 20°C. Animals were collected for RNA extraction at 24h (mostly J2 and a few J3 larvae), 50h (J3 and J4 larvae), and 70h (young adults) after bleaching; the presence and expected distribution of the desired stages at each time point was verified prior to sample collection by screening 50 specimens on each plate under a dissecting microscope. Worms were collected by adding 1ml water to each of the plates, scraping off the bacterial lawns containing the animals with Lazy L-shaped

spreaders, filtering the worms out of the resulting suspension with a 5μm nylon net filter,
and washing the animals off the filters into 1.5ml Eppendorf tubes with water. Finally, the
eggs were pelleted in these tubes at maximum speed for 2min using a tabletop centrifugate,
and the supernatant was discarded. Total RNA was extracted using the Direct-Zol RNA
Mini prep kit (Zymo Research) according to the instructions provided by the manufacturer.
Following Sieriebriennikov *et al.*'s approach [24], we combined 500ng of RNA that was
extracted at the 24h time point with 500ng of RNA extracted at the 50h time point. This
sample type thus contains a total of 1μg RNA extracted from J2-4 larvae. On the other hand,
we gathered 1μg of RNA extracted at the 70h time point (adult worms) and did not mix it
with RNA extracted from any other time point. Therefore, we ended up with two sample
types (for simplicity referred to as "larval" and "adult"), which we shipped to Novogene for
library preparation and mRNA sequencing. Libraries were sequenced as 150bp paired end
reads on Illumina's NovaSeq6000 platform. We sequenced the mRNA of three strains: the
wild-type *P. pacificus* strain (PS312), and two independently acquired frameshift mutants of
Ppa-nhr-10 (RS3920 and RS3921). Two biological replicates were collected for the larval and
adult stages of each of these three strains.
857
Confirmatory RNA-seq experiments - sequencing starving worms
We also performed *confirmatory* RNA-seq experiments, in order to properly identify
Ppa-NHR-10's regulatory targets in the same conditions in which we identified the survival
phenotype of mutant animals. Additionally, we aimed to investigate the wild-type
starvation response of *P. pacificus*. Therefore, using our starvation assay set-up (see
"Starvation assays and survival analysis"), we sequenced mRNA extracted from young
adult hermaphrodites which have been well-fed or acutely-starved for 24 hours (in liquid
cultures). Just like for the exploratory RNA-seq experiments, we used the wild-type strain
(PS312) of *P. pacificus* and the two reference *Ppa-nhr-10(lof)* strains (RS3920 and RS3921) for
these confirmatory RNA-seq experiments. After the liquid culture treatments, we generally
washed the worms three times with 0.5% Triton X-100/PBS and three times with PBS
through a 20μm nylon filter net, and subsequently preformed total-RNA extractions with
the Direct-Zol RNA Mini prep kit (Zymo Research), following manufacturer instructions.
For each sample, ~1μg of total RNA was extracted and shipped to Novogene for library
preparation and mRNA sequencing. As before, libraries were sequenced as 150bp paired
end reads on Illumina's NovaSeq6000 platform. This time, we sequenced three biological
replicates for each the three strains in each of the two conditions.
875
Quantification of transcripts and differential expression analysis (DEA)
Transcript abundances were estimated from the obtained raw read files relative to the *P.*
pacificus transcriptome using SALMON (ver. 1.5.2) [84,85]. We first build a decoy-aware
transcriptome file, using the entire *P. pacificus* genome as a decoy, following the exact
instructions   of   Salmon's   online   documentation   (see
https://salmon.readthedocs.io/en/latest/salmon.html under "Preparing transcriptome
indices (mapping-based mode)"). The decoy-aware transcriptome was then indexed for
subsequent read-mapping using an auxiliary *k*-mer hash over *k*-mers of the length of 31.
Reads were quantified directly against this index using Salmon's *quant* command, by
running the program in mapping-based mode with selective alignment as mapping strategy
("--validateMappings" flag). We allowed Salmon to automatically infer the library type ("-l

A" flag), and to simultaneously learn and correct for fragment-level GC-biases and
sequence-specific biases ("-- gcBias" and "-- seqBias" flags).

Differential expression analyses were carried out in R using BIOCONDUCTOR (ver.
3.17) [86], TXIMPORT (ver. 1.28.0) [87], and DESEQ2 (ver. 1.40.1) [49]. First, we created a
"transcript-name to gene-ID" cross-reference map ("tx2gene") from the publicly available *P.*
pacificus gene annotations using the GENOMICFEATURES (ver. 1.48.3) [88] and
ANNOTATIONDBI (ver. 1.58.0) [89] packages. After that, we summarized the transcript-level
abundance estimates generated by Salmon into gene-level count matrices using the *tximport*
function together with our tx2gene map. The *DESeqDataSetFromTximport* function of
DESEQ2 was used to specify the statistical designs for the analyses and the dataset was pre-
filtered for transcripts which had at least ten counts across all samples. We generally
performed multifactorial DEA by first combining the two factors of interest (i.e., "stage" and
"strain" in the exploratory RNA-seq experiments; or "condition" and "strain" in confirmatory
RNA-seq) into a single factor called "group" and subsequently modeled the mutation-effects
on transcript abundances in each stage or condition (e.g. "RS3920_larvae" *vs.*
"wildtype_larvae", or "RS3920_starved" *vs.* "wildtype_starved") using the *DESeq* function
[86]. The same logic was followed to extract the effect of starvation on wild-type worms (i.e.,
"wildtype_starved" *vs.* "wildtype_fed").

The obtained maximum likelihood estimates (MLE) of the log2-fold changes (L2FC)
for each gene were shrunken with the *lfcShrink* function using the adaptive shrinkage
estimator from the ASHR package [90]. This approach fits a mixture of normal distributions
to form an empirical Bayesian prior for the shrinkage procedure, in order to overcome the
strong heteroskedasticity in L2FC-estimates for genes with low counts while
simultaneously avoiding over-shrinkage of large L2FC-estimates [49]. The resulting
(shrunken) minimum mean squared error (MMSE) estimates of the L2FC were reported for
this study and used to rank and visualize candidate target genes.

Additionally, following recent advancements in the field of genomics, we abstained
from using the classical hypothesis testing approach for DEA, which assumes a "L2FC = 0"
as the null hypothesis and calculates FDR-corrected *P*-values (aka *q*-values) to claim
discoveries [49]. Briefly, this classical approach computes *P*-values, which tell us only how
confident we can be that there is any observable effect. These *P*-values, unfortunately, do
not at all tell us how confident we can be in either the estimated size of the effect or its
estimated direction. This problem can be addressed by hypothesis tests that (I) include a
minimum effect size in the null hypothesis (e.g. L2FC ≥ 'x'), and (II) compute *s*-values. In
this approach, the *s*-values report what we are actually interested in: whether the evidence
in the data is strong enough to support the belief that an effect can be observed, that the
observed effect is at least of the specified minimum size 'x', and that the direction of the
effect was correctly estimated [49,90]. For this reason, in order to infer differentially
expressed genes from our transcriptome data, we always included a L2FC-threshold of >
0.585 (absolute value) and a *s*-value threshold of < 0.005 into our hypothesis tests. This is
equivalent to asking, whether there is sufficient evidence in the data that a given gene is
indeed differentially expressed, that it is at least 50% up- or down-regulated, and that the
direction of this effect (up *vs.* down) was correctly estimated.

Starvation assays and survival analysis

932   Young adult hermaphrodites (day one of adulthood) were washed off of multiple agar
933   plates with 0.5% Triton-X100/PBS and a lazy-L scraper. The collected worm solutions were

filtered through a 20 μm nylon net filter and washed three times with 1ml 0.5% Triton-X100/PBS and, subsequently, three times with 1ml PBS. The bacteria-free worm pellets were resuspended in 250μl S-medium without cholesterol and added to empty agarose plates. The S-medium was allowed to be absorbed by the agarose and the clean worms were allowed to recover for one hour. During the recovery time, liquid cultures were set up for the experiments. We used small cell culture flasks (50ml) for our assays and added a total amount of 2.5ml liquid culture (LC) to each of them. The LCs for the acute starvation assays consisted only of 2.5ml S-medium (without cholesterol). The control LC with plenty of food were set up as follows: for each 2.5ml of final LC, we centrifuged 10ml of fresh bacterial stock solution (*E. coli* OP50, OD_{600} = 0.5) for 30min at 4250rpm (4°C). The bacterial pellets were resuspended in the according amount (2.5ml per assay flask) of S-medium without liquid culture and added to the cell culture flasks. To prevent unwanted microbial contaminations, we added 1μl of nystatin and 1.2μl of streptomycin to each assay flask, irrespective of whether it contained OP50 or not. Finally, we picked the young hermaphrodites which recovered from the washing step and added them to the assay flasks (~50 worms per flask). All assay flasks were kept in a shaking incubator set to 20°C and 130rpm and the ratio of alive worms was scored every 24hrs until all worms in the flask were dead. Media were replaced every third day in order to limit bacterial contaminations and remove offspring animals produced in the well-fed conditions. Overall differences in the survival curves within each of the two treatments were estimated using the Kaplan-Meier (KM) approach [91]. Pairwise comparisons among individual survival curves were based on the KM estimator and performed using the log-rank test [90]. Additionally, we estimated hazard rates for each of the strains in both experiments via Cox proportional hazard regression modeling [91,92]. Effect sizes were calculated as hazard ratios which compare the hazard rates of *Ppa-nhr-10(lof)* mutants (RS3920 and RS3921) to the wild-type hazard rate. *P*-values were corrected for false discoveries using the Bonferroni method. All survival analyses were carried out in R (ver. 4.3.0) [71] using the SURVIVAL (ver. 3.5-5), GGSURVFIT (ver. 0.3.0), SURVMINER (ver. 0.4.9), and GTSUMMARY (ver. 1.7.1) packages [93-96]. All survival experiments were repeated four times (N=4) independently with all three strains.

Phylogram of the NHR-1/-10/-40 cluster

We searched the genomes of ten nematode species for orthologs of *Ppa*-NHR-1, *Ppa*-NHR-10, and *Ppa*-NHR-40, using BLASTP (e < 0.0001). The list of genomes included *Brugia malayi* (clade III - Spiruromorpha), *Bursaphelenchus xylophilus* (clade IV - Tylenchida), *Panagrellus redivivus* (clade IV - Tylenchida), *Oscheius tipulae* (clade V - Rhabditida), *Caenorhabditis briggsae* (clade V - Rhabditida), *Caenorhabditis elegans* (clade V - Rhabditida; wormbase.org WS289), *Haemonchus contortus* (clade V - Strongylida), *Micoletzkya japonica* (clade V - Diplogasterida), *Parapristionchus giblindavisi* (clade V - Diplogasterida), and *Pristionchus pacificus* (clade V - Diplogasterida) [97-106] (non-diplogastrids: WormBase ParaSite ver. WBPS18; diplogastrids: pristionchus.org). The protein sequences of the obtained orthologs were aligned with MAFFT (ver. 7.48) using default settings [107]. Based on the results of the multiple sequence alignments, we calculated maximum likelihood trees in R (ver. 4.3.0) [71] using the PHANGORN (ver. 2.11.1) package [108] applying a LG substitution model with optimization of topology and base frequencies over 100 bootstrap replicates.

Pfam domain prediction, WormCat annotation, and overrepresentation analyses

Protein domains encoded in the proteome of *P. pacificus* were predicted using HMMER (ver. 3.3.2) [109] in conjunction with the Pfam-A.hmm database (ver. 3.1b2) [47] (hmmscan, e-value cutoff: < 0.001). The ORTHOFINDER software (default mode; ver. 2.5.4) [110] was used to define orthogroups of *P. pacificus* and *C. elegans* proteins; WormCat (ver. 2) annotations [48] for *P. pacificus* were created based on the obtained orthogroups. In a few cases, we found multiple *C. elegans* orthologs with different WormCat terms for an individual *P. pacificus* gene. In such cases, we transferred all unique WormCat terms among the orthologs to the given gene in *P. pacificus*. All overrepresentation analyses (ORAs) were based on Fisher's Exact test; obtained *P*-values were FDR-corrected.

Dauer assays

We performed three different assays regarding dauer biology: dauer entry, dauer exit, and dauer survival. The workflow of these assays is depicted in Fig 8A. First, we added a starting population of 500 young adult worms to 10ml standard liquid cultures (LCs) with OP50 as a food source (S-medium with or without cholesterol, 4μl nystatin and 5μl Streptomycin added). Worms were allowed to propagated for two weeks at 22°C and 180rpm. After two weeks we extracted 10μl from each LC and counted the number of dauers versus the number of feeding stages in the sample to quantify dauer entry (in %). The remaining volume of the LCs was treated with SDS following standard protocols [111] to isolate dauers. Some of the obtained dauers were added to agarose plates spotted with 300μl OP50 (30 worms per plate) and followed for multiple days to quantify the speed of dauer exit. These plates were kept at 20°C and worms were scored every 24 hrs until every individual exited the dauer arrest. The rest of the SDS-isolated dauers was added to LCs for the survival assays. These LCs did not contain food nor cholesterol and were again kept at 22°C and 180rpm for up to three weeks. Every week, we extracted 75μl of worm solution and added 25μl to three OP50-spotted agarose plates. We let the worms recover on these plates for 24 hrs and then scored alive versus dead worms to obtain the ratio of survivors (in %). All experiments were repeated multiple times independently (N=4). For the dauer exit and dauer survival assays we exclusively used dauers isolated from LCs that did not contain cholesterol. To statistically assess whether there are mutant-specific phenotypes in dauer entry, dauer exit, or dauer survival, we applied a two-tailed PERMANOVA [112,113] with null model residual randomization (RRPP) using the *lm.rrpp* function of the RRPP package (ver. 1.3.1) [114,115]. Relative effect sizes (Z-scores) and *P*-values are reported in Fig 8B-8D. We considered our dauer data as incompatible with the null hypothesis (i.e., "statistically significant") if the relative effect size was at least twice as large as the standard deviation (Z ≥ 2.0) and the *P*-value was below a type I error rate of 5% ($P < 0.05$). If at least one of these criteria was not met, we considered the data compatible with the null hypothesis (i.e., "not statistically significant"). Null model residuals were randomized and permuted 10,000 times. Pairwise comparisons of mutant and wild-type strains were performed with the *pairwise* function of RRPP [114,115].

Data processing, data visualization, and illustration

All data was processed, formatted, and wrangled in R (ver. 4.3.0) [71] using the TIDYVERSE [116]. All scientific plots depicted in this study were created using the GGPLOT2 package (ver. 3.4.2) [116]. Heatmaps were created using the PHEATMAP package (ver. 1.0.12) [117]. In order to make our plots accessible to people with color-vision deficiencies or color blindness [118], we exclusively used the scientifically-derived colormaps provided by the SCICO (ver. 1.4.0)

[119] and VIRIDIS (ver. 0.6.3) packages [120]. Microscopic images were edited and adjusted for levels, lightness, and contrast, in Affinity Photo (1.10.5). Scientific illustrations and final figures were created in Affinity Designer (ver. 1.10.5).

Author contributions

Conceptualization: Tobias Theska.
Data curation: Tobias Theska.
Formal analysis: Tobias Theska.
Funding acquisition: Ralf J. Sommer.
Investigation: Tobias Theska, Tess Renahan.
Methodology: Tobias Theska, Tess Renahan.
Project administration: Tobias Theska.
Resources: Ralf J. Sommer.
Software: Tobias Theska.
Supervision: Ralf J. Sommer.
Validation: Tobias Theska, Tess Renahan.
Visualization: Tobias Theska.
Writing - original draft: Tobias Theska.
Writing - review & editing: Tobias Theska, Tess Renahan, Ralf J. Sommer.

Competing interests

The authors declare no competing interests.

Data availability statement

Raw reads from the mRNA-seq experiments have been deposited in the European Nucleotide Archive (ENA) under the accession number PRJEB65081. Primer information, morph ratio data, landmark data for geometric morphometrics, Kaplan-Meier starvation survival data, differential expression analysis results, and dauer assay data are provided in the supplementary material (Sup Data 1-10).

Acknowledgements

We thank the Light Microscopy facility at the Max Planck Institute for Biology, especially Christian Feldhaus and Aurora Panzera, for their assistance with confocal laser-scanning microscopy. We are grateful to Heike Haussmann for freezing samples of all the worm strains produced for this study. Additionally, we thank Christian Rödelsperger, Adrian Streit, and Devansh R. Sharma for helpful discussions of our work.

References

1. Van Den Hoogen J, Geisen S, Routh D, Ferris H, Traunspurger W, Wardle DA, et al. Soil nematode abundance and functional group composition at a global scale. Nature. 2019;572:194–198.

2. Lambshead PJD, Boucher G. Marine nematode deep-sea biodiversity–hyperdiverse or hype? Journal of Biogeography. 2003;30:475–485.

3. Blaxter M. Nematodes: the worm and its relatives. PLOS Biology. 2011;9:e1001050.

4. Corsi AK, Wightman B, Chalfie M. A transparent window into biology: a primer on *Caenorhabditis elegans*. Genetics. 2015;200:387–407.

5. Watts JL, Ristow M. Lipid and carbohydrate metabolism in *Caenorhabditis elegans*. Genetics. 2017;207:413–446.

6. Baugh LR, Hu PJ. Starvation responses throughout the *Caenorhabditis elegans* life cycle. Genetics. 2020;216:837–878.

7. Rothman J, Jarriault S. Developmental plasticity and cellular reprogramming in *Caenorhabditis elegans*. Genetics. 2019;213:723–757.

8. Conradt B, Wu YC, Xue D. Programmed cell death during *Caenorhabditis elegans* development. Genetics. 2016;203:1533–1562.

9. Moss EG. Heterochronic genes and the nature of developmental time. Current Biology. 2007;17:R425–434.

10. Sommer RJ. Evolution of regulatory networks: nematode vulva induction as an example of developmental systems drift. In: Soyer O, editors. Evolutionary Systems Biology. Springer; 2012. p. 79–91.

11. Viney M, Diaz A. Phenotypic plasticity in nematodes: evolutionary and ecological significance. Worm. 2012; 1:98–106.

12. Sommer RJ, Dardiry M, Lenuzzi M, Namdeo S, Renahan T, Sieriebriennikov B, et al. The genetics of phenotypic plasticity in nematode feeding structures. Open biology. 2017;7:160332.

13. Sommer RJ, Ogawa A. Hormone signaling and phenotypic plasticity in nematode development and evolution. Current Biology. 2011;21:R758–766.

14. Haag ES, Fitch DH, Delattre M. From "the worm" to "the worms" and back again: the evolutionary developmental biology of nematodes. Genetics. 2018;210:397–433.

15. West-Eberhard MJ. Developmental plasticity and evolution. Oxford; New York: Oxford University Press; 2003.

16. Laland KN, Uller T, Feldman MW, Sterelny K, Müller GB, Moczek A, et al. The extended evolutionary synthesis: its structure, assumptions and predictions. Proceedings of the Royal Society B: Biological Sciences. 2015;282:20151019.

17. Uller T, Feiner N, Radersma R, Jackson IS, Rago A. Developmental plasticity and evolutionary explanations. Evolution & Development. 2020;22:47–55.

18. Sommer RJ. Phenotypic plasticity: from theory and genetics to current and future challenges. Genetics. 2020;215:1–13.

19. Hong RL, Sommer RJ. *Pristionchus pacificus*: a well-rounded nematode. Bioessays. 2006;28:651–659.

20. Sommer RJ. *Pristionchus pacificus*: a nematode model for comparative and evolutionary biology. Vol. 11. Leiden; The Netherlands: Brill; 2015.

21. Ragsdale EJ, Müller MR, Rödelsperger C, Sommer RJ. A developmental switch coupled to the evolution of plasticity acts through a sulfatase. Cell. 2013;155:922–933.

22. Kieninger MR, Ivers NA, Rödelsperger C, Markov GV, Sommer RJ, Ragsdale EJ. The nuclear hormone receptor NHR-40 acts downstream of the sulfatase EUD-1 as part of a developmental plasticity switch in *Pristionchus*. Current Biology. 2016;26:2174–2179.

23. Sieriebriennikov B, Prabh N, Dardiry M, Witte H, Röseler W, Kieninger MR, et al. A developmental switch generating phenotypic plasticity is part of a conserved multi-gene locus. Cell Reports. 2018;23:2835–2843.

24. Sieriebriennikov B, Sun S, Lightfoot JW, Witte H, Moreno E, Rödelsperger C, et al. Conserved nuclear
hormone receptors controlling a novel plastic trait target fast-evolving genes expressed in a single cell.
PLOS Genetics. 2020;16:e1008687.

25. Sun S, Theska T, Witte H, Ragsdale EJ, Sommer RJ. The oscillating Mucin-type protein DPY-6 has a
conserved role in nematode mouth and cuticle formation. Genetics. 2022;220:iyab233.

26. Sun S, Witte H, Sommer RJ. Chitin contributes to the formation of a feeding structure in a predatory
nematode. Current Biology. 2023;33:15–27.

27. Werner MS, Loschko T, King T, Reich S, Theska T, Franz-Wachtel M, et al. Histone 4 lysine 5/12
acetylation enables developmental plasticity of *Pristionchus* mouth form. Nature Communications.
2023;14:2095.

28. Cassada RC, Russell RL. The dauerlarva, a post-embryonic developmental variant of the nematode
Caenorhabditis elegans. Developmental biology. 1975;46:326–342.

29. Ogawa A, Streit A, Antebi A, Sommer RJ. A conserved endocrine mechanism controls the formation of
dauer and infective larvae in nematodes. Current Biology. 2009;19:67–71.

30. Renahan T, Lo W, Werner MS, Rochat J, Herrmann M, Sommer RJ. Nematode biphasic 'boom and bust'
dynamics are dependent on host bacterial load while linking dauer and mouth-form polyphenisms.
Environmental Microbiology. 2021;23:5102–5113.

31. Motola DL, Cummins CL, Rottiers V, Sharma KK, Li T, Li Y, et al. Identification of ligands for DAF-12
that govern dauer formation and reproduction in *C. elegans*. Cell. 2006;124:1209–1223.

32. Tao LJ, Seo DE, Jackson B, Ivanova NB, Santori FR. Nuclear hormone receptors and their ligands:
metabolites in control of transcription. Cells. 2020;9:2606.

33. Miglioli A, Canesi L, Gomes ID, Schubert M, Dumollard R. Nuclear receptors and development of
marine invertebrates. Genes. 2021;12:83.

34. Sural S, Hobert O. Nematode nuclear receptors as integrators of sensory information. Current Biology.
2021;31:4361–4366.

35. Sluder AE, Mathews SW, Hough D, Yin VP, Maina CV. The nuclear receptor superfamily has undergone
extensive proliferation and diversification in nematodes. Genome Research. 1999;9:103–120.

36. Robinson-Rechavi M, Maina CV, Gissendanner CR, Laudet V, Sluder A. Explosive lineage-specific
expansion of the orphan nuclear receptor HNF4 in nematodes. Journal of Molecular Evolution.
2005;60:577–586.

37. Peterson ND, Samantha YT, Huang QJ, Wani KA, Schiffer CA, Pukkila-Worley R. Non-canonical pattern
recognition of a pathogen-derived metabolite by a nuclear hormone receptor identifies virulent bacteria
in *C. elegans*. Immunity. 2023;56:768–782.

38. Howard RJ, Giacomelli M, Lozano-Fernandez J, Edgecombe GD, Fleming JF, Kristensen RM, et al. The
Ediacaran origin of Ecdysozoa: integrating fossil and phylogenomic data. Journal of the Geological
Society. 2022;179(4):jgs2021-107.

39. Riebesell M, Sommer RJ. Three-dimensional reconstruction of the pharyngeal gland cells in the
predatory nematode *Pristionchus pacificus*. Journal of Morphology. 2017;278:1656–1666.

40. Schlager S. Morpho and Rvcg–Shape Analysis in R: R-Packages for geometric morphometrics, shape
analysis and surface manipulations. In: Zheng G, Li S, Szekely G, editors. Statistical Shape and
Deformation Analysis. Academic Press; 2017. p. 217–256.

41. Schlager S, Jefferis G. Morpho: calculations and visualisations related to geometric morphometrics. R
package version 2.11. 2023. Available from: https://cran.r-
project.org/web/packages/Morpho/index.html

42. Hu PJ. Dauer. In: The *C. elegans* research community, editor. WormBook: The Online Review of *C. elegans*
Biology. 2007.

43. Srinivasan S. Regulation of body fat in *Caenorhabditis elegans*. Annual Review of Physiology. 2015;77:161–
178.

44. Lemieux GA, Ashrafi K. Neural regulatory pathways of feeding and fat in *Caenorhabditis elegans*. Annual
Review of Genetics. 2015;49:413–438.

45. Hansen M, Flatt T, Aguilaniu H. Reproduction, fat metabolism, and life span: what is the connection?
Cell Metabolism. 2013;17:10–19.

46. Athanasouli M, Akduman N, Röseler W, Theam P, Rödelsperger C. Thousands of *Pristionchus pacificus* orphan genes were integrated into developmental networks that respond to diverse environmental microbiota. PLOS Genetics. 2023;19:e1010832.

47. Finn RD, Coggill P, Eberhardt RY, Eddy SR, Mistry J, Mitchell AL, et al. The Pfam protein families database: towards a more sustainable future. Nucleic Acids Research. 2016;44:D279–285.

48. Holdorf AD, Higgins DP, Hart AC, Boag PR, Pazour GJ, Walhout AJ, et al. WormCat: an online tool for annotation and visualization of *Caenorhabditis elegans* genome-scale data. Genetics. 2020;214:279–294.

49. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biology. 2014;15:1–21.

50. O'Riordan VB, Burnell AM. Intermediary metabolism in the dauer larva of the nematode *Caenorhabditis elegans* II. The glyoxylate cycle and fatty-acid oxidation. Comparative Biochemistry and Physiology B, Comparative Biochemistry. 1990;95:125–130.

51. Braeckman B, Houthoofd K, Vanfleteren J. Intermediary metabolism. In: The *C. elegans* research community, editor. WormBook: The Online Review of *C. elegans* Biology. 2009.

52. Erkut C, Gade VR, Laxman S, Kurzchalia TV. The glyoxylate shunt is essential for desiccation tolerance in *C. elegans* and budding yeast. Elife. 2016;5:e13614.

53. Hibshman JD, Doan AE, Moore BT, Kaplan RE, Hung A, Webster AK, et al. daf-16/FoxO promotes gluconeogenesis and trehalose synthesis during starvation to support survival. Elife. 2017;6:e30057.

54. Abouheif E. Developmental genetics and homology: a hierarchical approach. Trends in Ecology & Evolution. 1997;12:405–408.

55. Amundson R. Form and Function in Evo-Devo. In: Nuño de la Rosa L, Müller GB, editors. Evolutionary Developmental Biology: A Reference Guide. Springer, Cham.; 2021, pp. 457–467.

56. Gabaldón T, Koonin EV. Functional and evolutionary implications of gene orthology. Nature Reviews Genetics. 2013;14:360–366.

57. Dolinski K, Botstein D. Orthology and functional conservation in eukaryotes. Annual Review of Genetics. 2007;41:465–507.

58. Nehrt NL, Clark WT, Radivojac P, Hahn MW. Testing the ortholog conjecture with comparative functional genomic data from mammals. PLOS Computational Biology. 2011;7:e1002073.

59. Dunn CW, Zapata F, Munro C, Siebert S, Hejnol A. Pairwise comparisons across species are problematic when analyzing functional genomic data. Proceedings of the National Academy of Sciences. 2018;115:E409–417.

60. Stamboulian M, Guerrero RF, Hahn MW, Radivojac P. The ortholog conjecture revisited: the value of orthologs and paralogs in function prediction. Bioinformatics. 2020;36:i219–i226.

61. Bulcha JT, Giese GE, Ali MZ, Lee YU, Walker MD, Holdorf AD, et al. A persistence detector for metabolic network rewiring in an animal. Cell Reports. 2019;26:460–468.

62. Stadler M, Fire A. Conserved translatome remodeling in nematode species executing a shared developmental transition. PLOS Genetics. 2013;9:e1003739.

63. Larance M, Pourkarimi E, Wang B, Murillo AB, Kent R, Lamond AI, et al. Global proteomics analysis of the response to starvation in *C. elegans*. Molecular & Cellular Proteomics. 2015;14:1989–2001.

64. Harvald EB, Sprenger RR, Dall KB, Ejsing CS, Nielsen R, Mandrup S, et al. Multi-omics analyses of starvation responses reveal a central role for lipoprotein metabolism in acute starvation survival in *C. elegans*. Cell Systems. 2017;5:38–52.

65. Stiernagle T. Maintenance of *C. elegans*. In: The *C. elegans* research community, editor. WormBook: The Online Review of *C. elegans* Biology. 2006.

66. Sanghvi GV, Baskaran P, Röseler W, Sieriebriennikov B, Rödelsperger C, Sommer RJ. Life history responses and gene expression profiles of the nematode *Pristionchus pacificus* cultured on *Cryptococcus* yeasts. PLOS One. 2016;11:e0164881.

67. Renahan T, Sommer RJ. Nematode interactions on beetle hosts indicate a role of mouth-form plasticity in resource competition. Frontiers in Ecology and Evolution. 2021;9:752695.

68. Renahan T, Sommer RJ. Multidimensional competition of nematodes affects plastic traits in a beetle ecosystem. Frontiers in Cell and Developmental Biology. 2022;10:985831.

69. Theska T, Sieriebriennikov B, Wighard SS, Werner MS, Sommer RJ. Geometric morphometrics of
microscopic animals as exemplified by model nematodes. Nature Protocols. 2020;15:2611–2644.

70. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an open-source
platform for biological-image analysis. Nature Methods. 2012;9:676–682.

71. R Core Team R. R: A language and environment for statistical computing. R Foundation for Statistical
Computing. 2023.

72. Adams DC, Otárola-Castillo E. geomorph: an R package for the collection and analysis of geometric
morphometric shape data. Methods in Ecology and Evolution. 2013;4:393–399.

73. Adams, D. C., Collyer, M. & Kaliontzopoulou, A. Geomorph: Software for geometric morphometric
analyses.   R   package   version   4.0.5.   2023.   Available   from: https://cran.r-
project.org/web/packages/geomorph/index.html

74. Baken EK, Collyer ML, Kaliontzopoulou A, Adams DC. geomorph v4. 0 and gmShiny: Enhanced
analytics and a new graphical interface for a comprehensive morphometric experience. Methods in
Ecology and Evolution. 2021;12:2355–2363.

75. Scrucca L, Fop M, Murphy TB, Raftery AE. mclust 5: clustering, classification and density estimation
using Gaussian finite mixture models. The R journal. 2016;8:289.

76. Fraley C, Raftery A, Scrucca L. mclust: Normal mixture modeling for model-based clustering,
classification, and density estimation. R package version 6.0.0. 2023. Available from: https://cran.r-
project.org/web/packages/mclust/index.html

77. Klingenberg CP. Size, shape, and form: concepts of allometry in geometric morphometrics. Development
Genes and Evolution. 2016;226:113–137.

78. Mitteroecker P, Gunz P, Windhager S, Schaefer K. A brief review of shape, form, and allometry in
geometric morphometrics, with applications to human facial morphology. Hystrix, the Italian Journal of
Mammalogy. 2013;24:59–66.

79. Mitteroecker P, Gunz P, Bernhard M, Schaefer K, Bookstein FL. Comparison of cranial ontogenetic
trajectories among great apes and humans. Journal of Human Evolution. 2004;46:679–698.

80. Witte H, Moreno E, Rödelsperger C, Kim J, Kim JS, Streit A, et al. Gene inactivation using the
CRISPR/Cas9 system in the nematode *Pristionchus pacificus*. Development Genes and Evolution.
2015;225:55–62.

81. Ghanta KS, Ishidate T, Mello CC. Microinjection for precision genome editing in *Caenorhabditis elegans*.
STAR Protocols. 2021;2:100748.

82. Han Z, Lo WS, Lightfoot JW, Witte H, Sun S, Sommer RJ. Improving transgenesis efficiency and CRISPR-
associated tools through codon optimization and native intron addition in *Pristionchus* nematodes.
Genetics. 2020;216:947–956.

83. Eren GG, Roca M, Han Z, Lightfoot JW. Genomic integration of transgenes using UV irradiation in
Pristionchus pacificus. microPublication Biology. 2022;29.

84. Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. Salmon provides fast and bias-aware
quantification of transcript expression. Nature Methods. 2017;14:417–419.

85. Srivastava A, Malik L, Sarkar H, Zakeri M, Almodaresi F, Soneson C, et al. Alignment and mapping
methodology influence transcript abundance estimation. Genome Biology. 2020;21:1–29.

86. Gentleman RC, Carey VJ, Bates DM, Bolstad B, Dettling M, Dudoit S, et al. Bioconductor: open software
development for computational biology and bioinformatics. Genome Biology. 2004;5:1–16.

87. Love MI, Anders S, Kim V, Huber W. RNA-Seq workflow: gene-level exploratory analysis and
differential expression. F1000Research. 2015;4:1070.

88. Lawrence M, Huber W, Pagès H, Aboyoun P, Carlson M, Gentleman R, et al. Software for computing
and annotating genomic ranges. PLOS Computational Biology. 2013;9:e1003118.

89. Pagès H, Carlson M, Falcon S, Li N. AnnotationDbi: Manipulation of SQLite-based annotations in
Bioconductor.   R   package   version   1.58.0.   2023.   Available   from:
https://bioconductor.org/packages/release/bioc/html/AnnotationDbi.html

90. Stephens M. False discovery rates: a new deal. Biostatistics. 2017;18:275–294.

91. Rich JT, Neely JG, Paniello RC, Voelker CC, Nussenbaum B, Wang EW. A practical guide to
understanding Kaplan-Meier curves. Otolaryngology—Head and Neck Surgery. 2010;143:331–336.

92. Dey T, Lipsitz SR, Cooper Z, Trinh QD, Krzywinski M, Altman N. Regression modeling of time-to-event
data with censoring. Nature Methods. 2022;19:1513–1515.

93. Therneau T. A package for survival analysis in R. R package version 3.5-5. 2023. Available from:
https://cran.r-project.org/web/packages/survival/index.html

94. Sjoberg D, Baillie M, Haesendonckx S, Treis T. ggsurvfit: Flexible Time-to-Event Figures. R package
version 0.3.0. 2023. Available from: http://www.danieldsjoberg.com/ggsurvfit/

95. Kassambara A, Kosinski M, Biecek P, Fabian S. survminer: Drawing Survival Curves using 'ggplot2.' R
package version 0.4.9. 2023. Available from: https://cran.r-
project.org/web/packages/survminer/index.html

96. Daniel d S, Whiting K, Curry M, Lavery JA, Larmarange J. Reproducible summary tables with the
gtsummary package. The R Journal. 2021;13:570–580.

97. Ghedin E, Wang S, Spiro D, Caler E, Zhao Q, Crabtree J, et al. Draft genome of the filarial nematode
parasite *Brugia malayi*. Science. 2007;317:1756–1760.

98. Dayi M, Sun S, Maeda Y, Tanaka R, Yoshida A, Tsai IJ, et al. Nearly complete genome assembly of the
pinewood nematode *Bursaphelenchus xylophilus* strain Ka4C1. Microbiology Resource Announcements.
2020;9:e01002-20.

99. Srinivasan J, Dillman AR, Macchietto MG, Heikkinen L, Lakso M, Fracchia KM, et al. The draft genome
and transcriptome of *Panagrellus redivivus* are shaped by the harsh demands of a free-living lifestyle.
Genetics. 2013;193:1279–1295.

100. Gonzalez de la Rosa PM, Thomson M, Trivedi U, Tracey A, Tandonnet S, Blaxter M. A telomere-to-
telomere assembly of *Oscheius tipulae* and the evolution of rhabditid nematode chromosomes. G3: Genes,
Genomes, Genetics. 2021;11:jkaa020.

101. Stevens L, Moya ND, Tanny RE, Gibson SB, Tracey A, Na H, et al. Chromosome-level reference genomes
for two strains of *Caenorhabditis briggsae*: an improved platform for comparative genomics. Genome
Biology and Evolution. 2022;14:evac042.

102. Laing R, Kikuchi T, Martinelli A, Tsai IJ, Beech RN, Redman E, et al. The genome and transcriptome of
Haemonchus contortus, a key model parasite for drug and vaccine discovery. Genome Biology. 2013;14:1–
16.

103. Prabh N, Roeseler W, Witte H, Eberhardt G, Sommer RJ, Rödelsperger C. Deep taxon sampling reveals
the evolutionary dynamics of novel gene families in *Pristionchus* nematodes. Genome Research.
2018;28:1664–1674.

104. Röseler W, Collenberg M, Yoshida K, Lanz C, Sommer RJ, Rödelsperger C. The improved genome of the
nematode *Parapristionchus giblindavisi* provides insights into lineage-specific gene family evolution. G3:
Genes, Genomes, Genetics. 2022;12:jkac215.

105. Rödelsperger C, Meyer JM, Prabh N, Lanz C, Bemm F, Sommer RJ. Single-molecule sequencing reveals
the chromosome-scale genomic architecture of the nematode model organism *Pristionchus pacificus*. Cell
Reports. 2017;21:834–844.

106. Athanasouli M, Witte H, Weiler C, Loschko T, Eberhardt G, Sommer RJ, et al. Comparative genomics
and community curation further improve gene annotations in the nematode *Pristionchus pacificus*. BMC
Genomics. 2020;21:1–9.

107. Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7: improvements in
performance and usability. Molecular biology and evolution. 2013;30:772–780.

108. Schliep KP. phangorn: phylogenetic analysis in R. Bioinformatics. 2011;27:592–593.

109. Mistry J, Finn RD, Eddy SR, Bateman A, Punta M. Challenges in homology search: HMMER3 and
convergent evolution of coiled-coil regions. Nucleic Acids Research. 2013;41:e121.

110. Emms DM, Kelly S. OrthoFinder: phylogenetic orthology inference for comparative genomics. Genome
Biology. 2019;20:1–14.

111. Karp X. Working with dauer larvae. In: The *C. elegans* research community, editor. WormBook: The
Online Review of *C. elegans* Biology. 2018.

112. Anderson MJ. A new method for non-parametric multivariate analysis of variance. Austral Ecology.
2001;26:32–46.

113.   Anderson MJ. Permutational multivariate analysis of variance (PERMANOVA). In: Balakrishnan N
Colton T, Everitt B, Piegorsch W, Ruggeri F, Teugels JL, editors. Wiley Statsref: Statistics Reference
Online. 2017. pp. 1–15.

114.   Collyer ML, Adams DC. RRPP: An r package for fitting linear models to high-dimensional data using
residual randomization. Methods in Ecology and Evolution. 2018;9:1772–1779.

115.   Collyer M, Adams D. RRPP: linear model evaluation with randomized residuals in a permutation
procedure.   R   package   version   1.3.1.   2023.   Available   from: https://cran.r-
project.org/web/packages/RRPP/index.html

116.   Wickham H, Averick M, Bryan J, Chang W, McGowan LD, François R, et al. Welcome to the Tidyverse.
Journal of Open Source Software. 2019;4:1686.

117.   Kolde R, Kolde MR. Package 'pheatmap.' R package version 1.0.12. 2023. Available from: https://cran.r-
project.org/web/packages/pheatmap/index.html

118.   Crameri F, Shephard GE, Heron PJ. The misuse of colour in science communication. Nature
Communications. 2020;11:5444.

119.   Pedersen TL, Crameri F. Scico: Colour palettes based on the scientific colour-maps. R package version
1.4.0. 2023. Available from: https://cran.r-project.org/web/packages/scico/index.html

120.   Garnier S, Ross N, Rudis R, Camargo PA, Sciaini M, Scherer C. viridis—colorblind-friendly color maps
for R. R package version 0.6.3. 2023. Available from: https://sjmgarnier.github.io/viridis/

Figures

Fig 1. Phenotypic plasticity throughout the *Pristionchus pacificus* life cycle and the NHR-1/-10/-40 phylogeny. (A) Alternative adult mouth forms in *P. pacificus*. Eu = eurystomatous, St = stenostomatous. (B) *P. pacificus* life cycle, including the alternative "indirect life cycle" via dauer development. (C) Closed mouth morphology of dauer larvae. bp = buccal plug. (D) Maximum-likelihood tree representing the protein phylogeny of NHR-1, NHR-10, and NHR-40 across various nematode species, representing clades III-V. Nodes without dots have a bootstrap support of ≤70%. Bma = *Brugia malayi*, Bxy = *Bursaphelenchus xylophilus*, Cel = *Caenorhabditis elegans*, Cbr = *Caenorhabditis briggsae*, Hco = *Haemonchus contortus*, Mja = *Micoletzkya japonica*, Oti = *Oscheius tipulae*, Pgi = *Parapristionchus giblindavisi*, Ppa = *Pristionchus pacificus*, Pre = *Panagrellus redivivus*.

1355
1356

1357

Fig 2. Patterns of *Ppa-nhr-10* expression across tissues. A 1.5kb long promoter sequence of
Ppa-nhr-10 (starting right before the start codon) was fused with *Ppa*-optimzed *GFP.* Relative
strength of fluorescent-protein expression is indicated via a relative intensity scale. (A) and
(B) each show a representative worm from our two independent reporter lines in different
orientations: (A) right body side facing reader, (B) left body side facing reader. Asterisk (*)
in (A) indicates the position of anterior gonad arm. Note that both, anterior and posterior
gonad arms, are always reporting; absence of the anterior arm in (A) is due to image
stacking only. The crossed double-arrows with letters in (A) and (B) indicate anatomical
orientation of the worm: A = anterior, P = posterior, D = dorsal, V = ventral. (C) Zoom-in
on the head of the worm depicted in (B), merged with DIC image. (D) Zoom-in on the
anterior gonad arm (outlined with white dots) of the worm depicted in (B). Note that neither
the sperm cells nor the eggs are reporting *Ppa-nhr-10* expression; only the somatic gonad
tissues do so. amp = ampulla of the gland cell(s), g1D = dorsal pharyngeal gland, g1V =
ventral pharyngeal glands, sp = sperm.

1372

Fig 3. Loss of *nhr-10* does not affect mouth morphology and only weakly affects mouth-
form plasticity. (A) Gene models of *Ppa-nhr-10* and frameshift alleles which lead to early
truncation of the ligand-binding domain. e = exon. (B) Ratio of predatory (Eu) and
bacterivorous (St) animals in the population. Worms were raised and scored on agar plates
with *E. coli* OP50 as a food source. Three biological replicates were obtained (N=3) and 150
worms of each strain were scored per replicate. Significant differences are indicated in black,
non-significant ones in gray. Z-scoes and *P*-values were derived from a two-tailed
PERMANOVA with null model residual randomization (RRPP). Bars represent mean
values across all replicates; error bars indicate ±1 standard error of the mean (C) Geometric
morphometric analysis of form differences in the nematode mouths. The PCA plot shows
the distribution of mouth morphologies of individual worms in a form space. Note that the
ranges of morphological variation (outlined by convex hulls) in *Ppa-nhr-10* mutant strains
completely overlap with the wild-type range of morphological variation. (D) Unsupervised
(model-based) clustering reveals that all 144 individuals belong to a single morphological
cluster, demonstrating that mutant worms cannot be differentiated from wild-type worms
based on mouth morphology. Only predatory (Eu) animals were used for the analyses in
(C) and (D).

Fig 4. Loss of *nhr-10* affects expression levels of 32 potential regulatory target genes. Bold names along the y-axis correspond to names of the misregulated putative targets (El_Paco_v3). Target genes which have an exact *C. elegans* ortholog were additionally given the public gene name of this ortholog (indicated in brackets). Panels indicate in which life history stage the putative targets genes were found. Labels within the panels indicate function information for the gene in form of WormCat terms (black) and/or predicted PFAM domains (gray). WormCat annotations were adapted from *C. elegans,* based on *Cel-Ppa*-orthogroups identified with OrthoFinder. PFAM domains were only indicated for genes we could not assign a WormCat term to, or genes whose WormCat terms ended in "unassigned". Cutoffs used in the differential gene expression analysis with DESeq2 and for PFAM domain prediction with HMMER are indicated in the lower left. DEGs = differentially expressed genes, MMSE = minimum mean squared error, FSOS = false sign or smaller.

Putative NHR-10 targets found across different life history stages

1408

1409

1410

1411

1412

Fig 5. Survival analyses reveal that the loss of *nhr-10* causes premature death during acute starvation. Comparison of the overall survival curves of wild-type *P. pacififcus* and two *Ppa-nhr-10(lof)* mutants that were well-fed (A) and completely starved (B). Survival curves show the mean survival, across all replicates, over time. Transparent ribbons behind the curves indicate the associated uncertainty in form of ±1 standard errors of the mean. Each of the two experiments (A and B) was repeated four times independently (N=4); the total numbers of worms (n) that were tracked across all replicates are indicated on top of both plots. Indicated *P*-values for pairwise comparisons resulted from a log-rank test that followed a Kaplan-Meier survival probability estimation; bold and black font indicates significant differences in the pairwise comparisons of overall survival. (C) Summary of the global Kaplan-Maier statistics for both experiments (in gray boxes) next to results from Cox proportional hazard regressions (white boxes). Points indicate the estimated hazard ratios for both mutant strains within each experiment (relative to the wild-type reference level of "1.0"). Intervals indicate associated uncertainty in form of 95% confidence intervals. *P*-values in bold black indicate significant differences between the mutant strain and the wildtype.

1429
1430

1431

1432

1433

1434

Fig 6. Analysis of the genome-wide patterns of transcriptional responses to acute starvation in wild-type *P. pacificus*. (A) Hierarchically clustered heat map showing the 1,007 genes which were identified as differentially expressed in response to 24h of acute starvation (SRGs). Trees on the left and the top show clustering results based on similarity in expression levels. Numbers on the gene tree indicate the number of genes in each of the two response clusters. WT = wildtype. (B) Bar chart showing the overrepresented gene-age classes and co-expression networks (according to [46]) identified the two response clusters, relative to the genome. "Phylogenetically old genes" refers to those genes in the genome of *P. pacificus* which are not diplogastrid-specific orphan genes. (C) Lollipop plot showing overrepresented PFAM domains in each response cluster. Y-axis gives PFAM domain names. X-axis gives the percentage of all unique genes encoding this PFAM domain, which were found to be present in the response cluster (e.g 60% means that 60% all unique genes in the genome that encode this domain are present in that cluster). (D) Bubble chart showing overrepresented WormCat terms in each of the response clusters. Enriched categories were broken down hierarchically (see WormCat legend in the upper right). Bubble size scales with the gene count indicated to the right of the bubble. The significance scale (*q*-value) depicted in (C) also applies to the bubbles depicted in (D).

1452

1455

Fig 7. Loss of *nhr-10* disturbs the expression levels of 37 putative regulatory targets in starving worms. Bold names along the y-axis correspond to names of the misregulated putative targets (El Paco v3) which are likely to cause the observed starvation-survival phenotype. Genes which have an exact *C. elegans* ortholog were additionally given the public gene name of this ortholog (indicated in brackets). Panels indicate whether these targets are members of the SRGs or not. Labels within the panels indicate functional information for the gene in form of WormCat terms (black) and/or predicted PFAM domains (gray). WormCat annotations were adapted from *C. elegans*, based on *Cel-Ppa-* orthogroups identified with OrthoFinder. PFAM domains were only indicated for genes we could not assign a WormCat term to, or genes whose WormCat terms ended in "unassigned". Cutoffs used in the differential gene expression analysis with DESeq2 and for PFAM domain prediction with HMMER are indicated in the lower left. Note that, in the current El Paco v3 annotations, PPA25752 (in SRG targets) seems to be a gene model that artificially fuses two separate genes. This is supported by multiple available RNA datasets and our PFAM domain predictions, which revealed that the first half of the hypothetical protein encodes a 7TM_GPCR, while the second half encodes a p450 cytochrome. As it seems unlikely that a single protein would act as both, a GPCR and a cytochrome, we repeated our differential expression analysis after manually splitting PPA25752 into two separate genes. We found no evidence of differential expression for the resulting GPCR-gene, but could perfectly replicate the differential expression signature we originally found for PPA25752, for the cytochrome-encoding gene. FSOS = false sign or smaller, MMSE = minimum mean squared error, SRG = starvation-response gene.

Putative NHR-10 targets that could explain the survival phenotype

Fig 8. Loss of *nhr-10* also causes various dauer-related phenotypes, including impaired long-term survival. (A). Workflow of the three dauer assays. (B-D) Results of the individual phenotypic assays indicated above each plot in the workflow chart. Point ranges indicate the mean estimates across all replicates ±1 standard error of the mean. Note that the dauers used in the exit and survival assays (C-D) were obtained from LCs without cholesterol. Additionally, all of the worms in (C) managed to exit dauer and fully resumed developments into viable adults. Hypothesis tests used a two-tailed PERMANOVA with null model residual randomization (RRPP); relative effect sizes (Z-scores) and *P*-values for pairwise comparisons among the three strains are reported in the upper right of each panel. Incompatibility with the null hypothesis ($Z \geq 2$ and $P < 0.05$) indicates the presence of a strain-specific phenotype; compatibility with the null hypothesis ($Z < 2$ or $P \geq 0.05$) supports the absence of a strain-specific phenotype. LCs = liquid cultures, rpm = rotations per minute, SDS = sodium dodecyl sulfate, SEM = standard error of the mean. All experiments were repeated multiple times independently (N=4). Number of worms in (B): n[PS312, (-)] = 460; n[RS3920, (-)] = 537; n[RS3921, (-)] = 432; n[PS312, (+)] = 750; n[RS3920, (+)] = 877; n[RS3921, (+)] = 879. Number of worms in (C): n[PS312] = 240; n[RS3920] = 240; n[RS3921] = 240. Number of worms in (D): n[PS312] = 395; n[RS3920] = 175; n[RS3921] = 177.

1498

1499

Supplementary Information

Sup Fig 1. Heatmap showing relative expression levels of NHR-10's regulatory targets from the confirmatory RNA-seq experiment. Hierarchical clustering was performed on the relative *rlog*-transformed normalized count values of the 37 genes which were identified as differentially expressed in mutants that have been acutely starved (see Fig 7 in main text). Trees on the left and the top show clustering results based on similarity in expression levels. Hierarchical clustering of all transcriptome samples based on the 37 candidate genes alone, perfectly recovered the grouping structures of the experiment (i.e., wildtype *vs.* mutant and fed *vs.* starved).

Sup Tab 1. Overrepresentation analysis shows now bias towards any given chromosomal location among *Ppa*-NHR-10's targets. The 37 candidate regulatory targets (DEGs) of *Ppa*-NHR-10, which were identified in the confirmatory RNA-seq experiments, were used as input for the analysis. No significant bias (FDR-corrected *P*-value < 0.05) towards any particular chromosome was found for *Ppa*-NHR-10's target genes. DEG = differentially expressed gene, FDR = false discovery rate.

chromosome	number_of_genes	number_of_DEGs	fold_enrichment	FDR_corrected_pvalue
Chr1	6869	5	0.5684764688986556	0.35319808452624085
Chr2	4462	11	1.9253031606236448	0.13589186241137188
Chr3	3736	5	1.045199374963829	0.9702253905888654
Chr4	6594	10	1.1843690824582545	0.8351614949834785
Chr5	4097	5	0.9531034573748757	1
ChrX	3002	1	0.26015089039739275	0.35319808452624085

Sup Tab 2. Loss of *Ppa-nhr-10* does not affect the expression levels of *P. pacificus* orthologs of the *C. elegans* propionate shunt genes. El Paco ver. 3 annotations were used for *P. pacificus*. *s*-values indicate FSOS event probability; significant changes to the expression levels ($s < 0.005$) could not be observed in any case. Strain indicated the *Ppa-nhr-10(lof)* mutant that was compared to wild-type worms. Cel = *C. elegans*, FSOS = false sign or smaller than 50% up- or down-regulated, l2fc = log2-fold change in expression levels, mmse = minimum mean squared error, sd = standard deviation.

Cel_gene	pristionchus_ortholog	l2fc_mmse	l2fc_posterior_sd	s-value	strain	treatment
alh-8	PPA18546	0.3120081894467676	0.183586125982154	0.779593708146979	RS3920	fed
alh-8	PPA18546	0.0700641209645215	0.124433208963777	0.903335595238679	RS3920	starved
alh-8	PPA18546	0.1578326679985086	0.116470311709115	0.870830237769447	RS3921	fed
alh-8	PPA18546	0.1310214980190643	0.111534703830993	0.801497484867020	RS3921	starved
ech-6	ppa_stranded_DN19950_c1_g2_i1	0.2915677450017385	0.121252839902834	0.890590474805354	RS3920	fed
ech-6	ppa_stranded_DN19950_c1_g2_i1	0.0278618218738554	0.058205088196759	0.922220261931968	RS3920	starved
ech-6	ppa_stranded_DN19950_c1_g2_i1	0.302048528212008	0.099893413969880	0.865990199881571	RS3921	fed
ech-6	ppa_stranded_DN19950_c1_g2_i1	0.1002545176406774	0.078147156473995	0.831089635382915	RS3921	starved
hach-1	ppa_stranded_DN24485_c0_g1_i1	0.1503250384307045	0.179955991152806	0.871633632978635	RS3920	fed
hach-1	ppa_stranded_DN24485_c0_g1_i1	-0.003755268928726	0.048962247587748	0.916236852972432	RS3920	starved
hach-1	ppa_stranded_DN24485_c0_g1_i1	0.3403493778772363	0.171220920053186	0.685735755852485	RS3921	fed
hach-1	ppa_stranded_DN24485_c0_g1_i1	-0.029993946051769	0.095674037390754	0.816727072214799	RS3921	starved
hphd-1	PPA05876	0.2710472432776699	0.186482563130857	0.845357698346291	RS3920	fed
hphd-1	PPA05876	-0.021954386415356	0.066963132113470	0.912373249230267	RS3920	starved
hphd-1	PPA05876	0.1715677303340060	0.120611075832775	0.868252620804282	RS3921	fed
hphd-1	PPA05876	-0.388037641153159	0.170339234415574	0.558929874614430	RS3921	starved