The pathogenesis of experimental Emergomycosis in mice.

Running title: A novel mouse model of *Emergomyces africanus* infection

Maxine A. Höft[1,2], Lucian Duvenage[1,2], Sumayah Salie[1,2], Roanne Keeton[2], Alfred Botha[3], Ilan

S. Schwartz[4], Nelesh P. Govender[1,5,6], Gordon D. Brown[1,6], J. Claire Hoving[1,2,6✉].

[1] CMM AFRICA Medical Mycology Research Unit, Institute of Infectious Diseases and

Molecular Medicine (IDM), University of Cape Town, Cape Town, South Africa

[2] Division of Immunology, Department of Pathology, Faculty of Health Sciences, University

of Cape Town, Cape Town, South Africa

[3] Department of Microbiology, Stellenbosch University, Stellenbosch, South Africa

[4] Division of Infectious Diseases, Department of Medicine, Duke University School of

Medicine, Durham, North Carolina, USA

[5] National Institute for Communicable Diseases, Division of the National Health Laboratory

Service, Johannesburg, South Africa; School of Pathology, Faculty of Health Sciences,

University of the Witwatersrand, Johannesburg, South Africa

[6] Medical Research Council Centre for Medical Mycology at the University of Exeter, Geoffrey

Pope Building Stocker Road, Exeter, United Kingdom

Corresponding author: Jennifer Claire Hoving; jennifer.hoving@uct.ac.za

Abstract

Emergomyces africanus is a recently identified thermally-dimorphic fungal pathogen that

causes disseminated infection in people living with advanced HIV disease. Known as

emergomycosis, this disseminated disease is associated with very high case fatality rates. Over

the last decade, improved diagnostics and fungal identification in South Africa resulted in a

dramatic increase in the number of reported cases. Although the true burden of disease is still

unknown, emergomycosis is among the most frequently diagnosed dimorphic fungal infections

in Southern Africa; and additional species in the genus have been identified on four continents.

Little is known about the pathogenesis and the host's immune response to this emerging

pathogen. Therefore, we established a murine model of pulmonary infection using a clinical

isolate, *E. africanus* (CBS 136260). Both conidia and yeast forms caused pulmonary and

disseminated infection in mice with fungus subsequently isolated in culture from lung, spleen,

liver, and kidney tissue. Wild-type C57BL/6 mice demonstrated a drop in body weight at two

weeks post-infection, corresponding to a peak in fungal burden in the lung, spleen, liver, and

kidney. An increase in pro-inflammatory cytokine production was detected in homogenized

lung supernatants including IFN-γ, IL-1β, IL-6, IL12-p40 and IL-17 at three- and four-weeks

post-infection. Rag-1-deficient mice, lacking mature T and B cells, had an increased fungal

burden associated with reduced IFN-γ production. Together our data support a protective T-

helper type-1 immune response to *E. africanus* infection. This may provide a possible

explanation for the susceptibility of only a subset of people living with advanced HIV disease

despite hypothesized widespread environmental exposure. In summary, we have established a

novel murine model of *E. africanus* disease providing critical insights into the host immune

components required for eliminating the infection.

Author summary:

Emergomyces africanus is a thermally-dimorphic fungus that was recently described as the cause of disseminated infections in persons living with advanced HIV disease in South Africa, where emergomycosis is among the most frequently diagnosed dimorphic fungal diseases. Four additional *Emergomyces* species have been described causing serious infections around the world, and the infection has a high case fatality rate. Despite the seriousness of this infection, the pathophysiology, immunology, diagnosis, and optimal management of emergomycosis remain largely unknown, hampered by the absence of an animal model. Here, we established a mouse model of pulmonary and disseminated emergomycosis using a clinical isolate of *E. africanus* with the aim of investigating the host immune response. We found that infection spread from the lungs to other organs within two weeks. We describe the effector immune responses in immune-competent mice aiding in controlled infection. We also highlight the immune-modulating components lacking in immunocompromised mice that promote susceptibilty to infection. Our study provides new insights into the components of the host immune response required for controlling *Emergomyces* spp. infection.

Keywords: *Emergomyces africanus*, thermally-dimorphic fungal pathogens, opportunistic, host immune response fungal pathogens, fungal diseases, endemic mycoses

Introduction

Emergomyces africanus is a thermally-dimorphic fungal pathogen belonging to the Ajellomycetaceae (Onygenales) family (Jiang, Dukik et al. 2018). Emergomycosis is the disease caused by seven dimorphic fungal species belonging to the newly described genus, *Emergomyces* (Dukik, Muñoz et al. 2017, Jiang, Tsui et al. 2020). These include *Emergomyces pasteurianus* (Europe, Asia, Africa), *E. canadensis* (North America), *E. orientalis* (Asia) and *E. europaeus* (Europe) (Dukik, Muñoz et al. 2017). Recently*, E. sola* and *E. crescens* have been included into the *Emergomyce*s genus (Jiang, Tsui et al. 2020). Molecular evidence suggests that *E. africanus* is found in the environment. Using spore traps (for air) and a PCR assay, DNA was detected in air and soil samples (Schwartz, Lerm et al. 2018, Schwartz, McLoud et al 2018). Hyphae produce conidiophores and conidia which are thought to be the infectious agent whereby conidia and hyphal fragments are aerosolised upon soil disruption or wind and inhaled by the host (Schwartz, McLoud et al. 2018). Once inhaled, the change in temperature from environmental to body temperature (37 °C) is believed to drive a morphological switch from conidia to a yeast-like phase, which is responsible for pulmonary and upon dissemination, extrapulmonary disease (Kenyon, Bonorchis et al. 2013).

To date, almost all reported cases of emergomycosis are opportunistic infections, occurring among immunocompromised individuals with the majority of cases in persons with advanced HIV disease (Kenyon, Bonorchis et al 2013, Schwartz, Maphanga et al. 2018). Radiographic findings of pulmonary disease were present in 86% of South African patients at the time of diagnosis (Schwartz, Govender et al. 2015), although most patients are diagnosed late in disease course, after the fungus has disseminated to skin. Cutaneous lesions, appearing as papules, plaques, nodules or ulcers, with widespread distribution, have been described in >96% of patients at the time of diagnosis (Schwartz, Govender et al. 2015). Dissemination to other

tissues such as the gastrointestinal tract, liver, lymph nodes and bone marrow also occurs but are more difficult to diagnose (Schwartz, Govender et al. 2015).

Diagnostic methods include fungal culture from tissue biopsy as well as histopathology. Positive histopathology with special fungal stains results in the visualisation of small yeast-like cells (2-4 µm diameter) with narrow-based budding (Kenyon, Bonorchis et al. 2013). *Emergomyces* spp. can often be misidentified as *Histoplasma capsulatum* as the histopathology findings are very similar (Schwartz, Kenyon et al. 2017). Polymerase chain reaction (PCR) amplification of fungal DNA in infected tissue can also be used as a diagnostic method by amplifying regions of the nuclear ribosomal internal transcribed spacer (ITS) and β-tubulin gene (Dukik, Muñoz et al. 2017, Schwartz, Sanche et al. 2018). There are currently no sensitive and specific serological or antigen tests for emergomycosis and there is an urgent need for a rapid and cost-effective diagnostic that may be available in resource-limited settings.

Emergomycosis is assumed to be universally fatal if not treated, while case-fatality ratios in patients receiving treatment are still high, possibly due to diagnosis occurring at late stages of infection. There are consensus guidelines for the diagnosis and treatment of emergomycosis but many of the recommendations are extrapolated from histoplasmosis treatment approaches due to limited human and absent animal data (Thompson, Le et al. 2021). There are currently no animal models of pulmonary infection with emergomycosis. In a proof-of-principle study, Schwartz et al infected mice via intraperitoneal inoculation. However, this would not represent natural infection believed to be via inhalation and therefore limits the understanding of disease kinetics and immunologic response during infection (Schwartz, Lerm et al. 2018). The development of an animal model of pulmonary emergomycosis to study the pathophysiology and immunology of infection and advance diagnostics and treatment strategies is therefore an urgent priority.

The goal of this study was to establish a mouse model of pulmonary disease and to identify key

immune components during infection. Using a clinical isolate of *E. africanus,* we were able to

induce a pulmonary and disseminated infection in wild-type mice. We showed that following

infection, *E. africanus*, disseminated from the lung to the spleen, liver, and kidneys within two

weeks; and by four weeks, it had been cleared from the lung, liver, and kidney (but not spleen).

Infection triggered a T helper-type 1 immune response supported by the detection of pro-

inflammatory cytokines such as IFN-γ, IL-1β, IL-6, IL-12p40 and IL-17 in the supernatant of

lung homogenates. These observations were further elucidated by conducting studies involving

immunocompromised mice (Rag-1$^{-/-}$), which lack mature T and B cells. These studies revealed

an increased fungal burden coupled with diminished IFN-γ production.

Understanding the kinetics of *E. africanus* in an immunocompetent murine model provides

insight into the vital immune components orchestrating control and clearance of infection. This

model creates the foundation for future mechanistic studies of disease progression and

clearance.

Results

Wild-type mice develop *Emergomyces africanus* infection.

To establish a mouse model of disseminated disease for studying host-pathogen interactions, we first determined whether yeast cells of *E. africanus* could infect different mouse strains. A liquid fungal suspension was prepared from yeast cells of *E. africanus* isolated from a patient with advanced HIV disease. Susceptibility to infection was investigated in wild-type mice from three different genetic backgrounds: BALB/c, C57BL/6 and Sv129. As previously described for other fungi, all mice received a non-invasive intratracheal inoculation of between 0.5 to $1x10^5$ yeast cells/50 µl (Hoving, Munyonho et al. 2023). Survival curves showed that all wild-type mice were susceptible to this high-dose infection (Fig 1A). Progressive weight loss occurred in all wild-type mice from 6-days post-infection (Fig 1B). Yeast infection caused pulmonary as well as disseminated infection in all wild-type mouse strains by two-weeks post infection, shown by organisms cultured at 37 °C on brain heart infusion agar (BHI) plates in lung, spleen, liver, and kidney. Representative plates are shown in Fig 1C and Fig 1D.

Disseminated emergomycosis disease parameters in C57BL/6 wild-type mice.

As many gene-deficient mice have been created on a C57BL/6 background, we focussed most of our subsequent studies on this strain, starting with a time-kinetic study of progression of disease. Due to the higher dose resulting in mortality (described above), we infected mice with a lower dose (i.e., $1x10^3$ *E. africanus* yeast cells/50 µl) via non-invasive intratracheal inoculation and euthanised the mice at weekly intervals. Uninfected, naïve mice were used as a comparison to determine *E. africanus*-specific responses. At the lower dose of $1x10^3$ yeast cells/50 µl, C57BL/6 wild-type mice showed transient weight loss at 14- to 17-days post-infection when compared to uninfected mice (Fig 2A). Fungal organisms could be cultured

from the lung, spleen, liver, and kidney (quantified as CFU per gram of respective tissue, Fig 2B). The fungal burden was detected in the lungs within 7 days post-infection and disseminated to other organs by two weeks post-infection. Except for the spleen, the fungal burden diminished and was undetectable by Day 28 in all organs, suggesting the mice could control infection in these tissues. Infected lung lobes were enlarged with macroscopic evidence of lesion formation, particularly at two- and three-weeks post-infection (Fig 2C). Haematoxylin and eosin (H&E) staining of the left lung lobes showed patchy or localized areas of infection and cellular involvement, resembling granuloma-like formations evident from two-weeks post-infection, waning by four weeks (Fig 2D). High magnification H&E and Grocott's methenamine silver stain (GMS)-stained sections for yeast localisation are shown in Supp Fig S1 and Supp Fig S2. Conidia were also able to cause disseminated disease with detectable organisms in lung, spleen, liver and kidney at day 14 post-infection (Supp Fig S3).

T-helper type 1-mediated immune response to *Emergomyces africanus*.

To further characterise the immune response to infection, we characterised cytokine levels in the lung and spleen during the kinetic study. Cytokine levels were compared with the levels obtained for uninfected mice (UI). Of interest, IFN-γ, IL-1β and IL-6 lung cytokine levels were significantly increased in the lung at both two- and three-weeks post-infection. These levels peaked by Day 21 post-infection and then returned to levels comparable to uninfected animals (see Fig 3 for the statistical description of cytokine production). Lung IL-12p40 was significantly increased at Day 14, Day 21 and Day 28 post-infection while IL-17 production was significantly higher at Day 14 post-infection (Fig 3). There was no notable change in lung TNF, IL-12p70 and IL-10 levels at any time point during infection (Fig 3).

During the kinetic study, blood was collected from mice by cardiac puncture immediately after euthanasia. Total blood serum antibody production over time was evaluated by antibody

ELISA and compared to serum antibody levels in uninfected mice. Serum IgM, IgG1 and IgG2a antibody production was significantly increased at Day 21 and Day 28 post-infection when compared to uninfected mice (Fig 4).

Increased susceptibility of immunocompromised mice to *E. africanus.*

Since the majority of *E. africanus* cases are diagnosed in patients with advanced HIV disease (AIDS) and thus very low $CD4^+$ T-lymphocyte counts, a Rag-1-deficient (devoid of mature T and B lymphocytes) murine model of infection was established to study the infection in an immunocompromised host. $Rag-1^{-/-}$ mice were infected via non-invasive intratracheal inoculation with $1x10^3$ yeast cells/ 50 µl and survival was compared to C57BL/6 wild-type mice. All $Rag-1^{-/-}$ mice succumbed to infection by Day 20 post-infection (Fig 5A) and the increased mortality was associated with a significant increase in fungal burden at Day 14 post-infection in all organs tested (Fig 5B). Histopathology demonstrated that diffuse cellular infiltrates were observed in infected lung tissue of C57BL/6 and $Rag-1^{-/-}$ mice at Day 7 post-infection (Fig 5C). At Day 14, both C57BL/6 and $Rag-1^{-/-}$ mice developed similar pathology with granuloma-like structures. Increased magnification of lung tissue areas confirmed a concentration of yeast cells (positive GMS staining) within the granuloma-like structures when compared to surrounding tissue (Supp Fig S4). These data demonstrate the location of yeast cells, while the CFU data quantifies fungal burden.

Susceptibility of Rag-1-deficient mice associated with reduced cellular recruitment and IFN-γ production.

Since $Rag-1^{-/-}$ mice had significantly increased fungal burden at two-weeks post-infection, we determined if there were differences in cellular recruitment to infected lung tissue in these mice. A myeloid cell flow cytometry panel was established and optimised for the infection model

using C57BL/6 and Rag-1$^{-/-}$ mice infected via non-invasive intratracheal inoculation with

$1x10^3$ yeast cells/ 50 µl. Infected lungs were harvested at Day 14 post-infection and single cell

suspension prepared for cell staining (Supp Fig S5, Supp Fig S6 and Supp Table S1).

Representative scatter plots are shown for each identified cell population in both wild-type and

Rag-1$^{-/-}$ mice (Fig 6A). Despite exacerbated disease and increased CFUs, Rag-1$^{-/-}$ mice showed

a significant increase in the frequency (%) of alveolar macrophages, neutrophils, and NK cells

(Fig 6B and Supp Fig S7). Interestingly there was a significant decrease in eosinophils. Both

inflammatory monocytes and dendritic cells were not different from wild-type mice. Notably,

when measuring cytokine production by ELISA assay from lung tissue, IFN-γ was the only

cytokine affected in Rag-1$^{-/-}$ mice compared to C57BL/6 mice (Fig 7).

Discussion

This study reports the first pulmonary *Emergomyces* infection model in mice. We showed that

following pulmonary infection, *E. africanus* disseminated to the spleen, liver, and kidneys

within two weeks, and was cleared from the lung, liver, and kidney (but not spleen) by four

weeks. Infection triggered a T helper-type 1 immune response supported by the detection of

pro-inflammatory cytokines such as IFN-γ, IL-1β, IL-6, IL-12p40 and IL-17 in the supernatant

of lung homogenates. Next, we used immunocompromised (Rag-1-deficient) mice, to identify

immunologic cascades essential for controlling infection. This work advances our

understanding of the nature and function of the immune system in emergomycosis and sets the

foundation for future development of diagnostics and therapies for the disease.

Our data shows for the first time the susceptibility of experimental mice to the yeast (and

conidia) form of a clinically-isolated *E. africanus* strain. We provide a description of the

disease kinetics and immune responses leading to fungal clearance in the lung and how these

are compromised in Rag-1-deficient mice. Since little is known or published about the host

immune response to *Es. africanus*, we compared our results and observations with other closely

related thermally-dimorphic fungi, specifically, *H. capsulatum* due to its phylogenetic and

genetic similarity to *Emergomyces* spp. To date *E. africanus* infections appear to be limited to

immunocompromised patients with advanced HIV disease; however, with improved

serological and/or antigen testing and therefore surveillance, detection is likely to increase in

other groups of people.

Unlike other dimorphic fungal pathogens, an animal reservoir for *Emergomyces* spp. is yet to

be found, despite efforts in screening 26 different species of small mammals (Cronjé, Schwartz

et al. 2018). Our data has shown that infection with the yeast (and conidia) form is possible in

three distinct mouse strains: C57BL/6, BALB/c and Sv129. During survival studies in which

mice were euthanised once reaching a humane endpoint of 20% weight loss, all mice eventually

succumbed to infection at the higher does. For lower dose experiments, wild-type mice that did

not need to be euthanized were able to clear the infection. Mice of different genetic

backgrounds have been reported to respond differently to certain infection models. This is

genetically determined as mice carry varied immunoregulatory genes within alleles of the H2

region (Matthews, Christiansen et al. 2000). These genes regulate the efficiency of antigen

processing, cytokine production, as well as the expression and functioning of cytokine

receptors. Therefore, C57BL/6 mice are prone to Th1- responses and BALB/c mice Th2-prone

(Matthews, Christiansen et al. 2000). Considering that the mice of interest for future studies

are on a C57BL/6 background, and that *Emergomyces* appears to be an intracellular pathogen

(likely to require Th1 responses for clearance) next we determined the disease progression and

parameters in C57BL/6 mice.

Mice infected with a lower dose over four weeks showed significant weight loss at 14 to 17-

days post-infection when compared to uninfected mice but mice that survived were able to

clear infection. Fungal propagules disseminated to multiple organs from the lung infection and

the fungal burden seemed to peak at two weeks after which it was cleared from most organs.

A similar study comparing disease kinetics of multiple *H. capsulatum* strains in C57BL/6 mice

infected with a similar dose (2.5×10^3 yeast cells per mouse) demonstrated a comparable weight

loss profile, with lung CFU peaking 8- days post-infection for all *H. capsulatum* strains.

Clearance of fungal organisms in the lung followed a similar trend with CFU below the

detection limit by 32-days post-infection (Sepúlveda, Williams et al. 2014).

Organ lesions and granulomas are described as a major interface between the host and fungal

pathogens. H&E staining of lung sections during our kinetic study showed granuloma-like

formations detected from two-weeks post-infection. We speculate that granuloma formation

during emergomycosis is induced by the infection of macrophages by yeast cells within the

lung tissue, causing an infiltration of immune cells, which has also been demonstrated during

histoplasmosis (Heninger, Hogan et al. 2006). Granuloma formation is a beneficial response,

required to localize fungal replication, limit dissemination, and protect the lung from extensive

tissue damage. The reduced lung pathology in these studies coincided with a decrease in lung

fungal burden by four-weeks post-infection.

Interestingly, we saw an increase in cell debris noted in lung histology sections at three- and

four weeks post-infection which may be indicative of cellular apoptosis and killing of yeast

cells with possible necrosis occurring at the resolving granuloma sites. While further

investigation is required to confirm these observations, suppressing apoptosis during

Histoplasma spp. infection has been shown to increase the severity of disease, associated with

increased IL-10 and IL-4 production (Allen 2005). By the end of week four, the granuloma-

like structures had resolved, which may have been facilitated by macrophages clearing

apoptotic and yeast-infected cells.

Studies of the immune response to other thermally-dimorphic fungi show that innate immune

cells, such as macrophages and dendritic cells, cooperate with the adaptive immune cells, such

as $CD4^+$ and $CD8^+$ T cells, to control infection. The studies showed that early in infection,

when alveolar macrophages interact with fungal cells, essential pro-inflammatory cytokines

were produced: IL-12, TNF, IL-1, IFN-γ and GM-CSF. These cytokines were required for the

activation of protective adaptive immune responses (Zhou, Sieve et al. 1995, Deepe, Romani

et al. 2000, Deppe and McGuinness 2006). Similarly, our study showed a significant increase

in lung IFN-γ, IL-1β, IL-6, IL-12p40 and IL-17 cytokine production. However, these cytokines

were only increased later in infection and with no difference in the lung at one-week post-

infection, despite a large fungal burden. This data suggests that *E. africanus* may not elicit a

pronounced innate immune response, unlike other opportunistic fungal pathogens. Our study

showed a steady significant increase in lung IL-12p40 from two- to four-weeks post-infection.

In other dimorphic infection models, IL-12 production has been shown to be produced by

macrophages and dendritic cells, importantly linking both innate and adaptive immunity. IL-

12 production has been shown to be essential in adaptive immune responses to dimorphic fungi

(including *H. capsulatum*), leading to the promotion and generation of Th1 cells and cytokines

(such as IFN-γ production by T cells) and subsequent protection against disseminated infection.

In *Histoplasma* spp., infection, neutralisation of IL-12 led to fatality and was associated with a

decrease in IFN-γ, but not TNF or GM-CSF (Zhou, Sieve et al. 1995). Similar future studies in

our *E. africanus* model, will determine if IL-12 plays a protective role. Interestingly, our data

showed no significant increase in TNF levels in lung supernatant during the *E. africanus* kinetic

study. In contrast, TNF has been shown to play an important role in controlling *H. capsulatum*

infection. Increased production of TNF was detected from granuloma infiltrating cells,

primarily from macrophages (Heninger, Hogan et al. 2006). In addition, studies using anti-TNF

treatment were shown to reactivate latent *H. capsulatum* in some patients, highlighting an

important role for TNF in mediating immune responses, controlling yeast infection and

dissemination (Wood, Hage et al. 2003, Allendoerfer and Deepe 1998). Perhaps the importance

of TNF in response to *E. africanus* occurs earlier and would have been missed in our kinetic

studies starting at one-week post-infection. Therefore, future studies could include cytokine

detection in lung supernatant or bronchoalveolar lavage fluid within the first few days

following infection to determine acute inflammatory immune responses to *E. africanus*

infection. Lastly, increased IL-17 production by CD4[+] T cells and activated macrophages have

been shown to induce macrophage effector function during *H. capsulatum* infection (Heninger,

Hogan et al. 2006). However, experiments in C-type lectin receptor-deficient mice showed a

decrease in lung IL-17 but the mice maintained their ability to clear infection. Therefore, we

hypothesize that although IL-17 was increased at two-weeks post-infection, it may not be

essential and rather that Th1 responses are central in driving clearance.

Antibodies have shown disparate biological effects in fungal infections, but the production of

IFN-γ and opsonizing antibodies promote Th1 cells that activate phagocytes at sites of

infection. Therefore, we measured serum antibody levels in our time-course study. IgM, IgG1

and IgG2a were increased at three- and four-weeks post-infection. Interestingly mice that were

susceptible to *Paracoccidioides* infection have been correlated with preferential secretion of

IgG2b-specific antibodies (Cano, Singer-Vermes et al. 1995) rather than IgG2a and IgG1

which were shown to be correlated with improved disease outcomes. This may suggest a

protective role for significantly increased IgG1 and IgG2a against *E. africanus*. Although, B

cell-deficient mice infected with *H. capsulatum* were not more susceptible to disease

(Chandler, Smith et al. 1969, Allendörfer, Brunner et al. 1999). Based on these findings it may

be prudent to determine if antibodies are playing an essential role in controlling *E. africanus*

infection by performing further studies in B cell-deficient mice.

Given the susceptibility of immunocompromised patients with advanced HIV disease to *E.*

africanus, our next objective was to establish the mouse model using Rag-1-deficient mice.

Rag-1$^{-/-}$ mice lack mature T- and B-cells due to defective V(D)J recombination thus halting T-

and B-cell differentiation at an early stage. We showed Rag-1$^{-/-}$ mice to be more susceptible to

E. africanus compared to C57BL/6 wild-type mice. This was associated with significantly

increased fungal burden at two-weeks post-infection, as well as significantly higher

dissemination to other organs including the spleen, liver, and kidney. Rag-1$^{-/-}$ mice showed

successful "granuloma" formation in the lungs at two-weeks post-infection. This could be

attributed to the fact that macrophages have been shown to be the dominant cell type in *H.*

capsulatum-mediated granuloma formation (Heninger, Hogan et al. 2006). Interestingly, the

only significant difference in lung cytokine production was a decrease in IFN-γ. IFN-γ has been shown to play a major role in all dimorphic fungal immunity. Indeed, IFN-γ-deficient mice succumb to *H. capsulatum* infection with significantly higher lung CFU, associated with significantly higher TNF and GM-CSF but not IL-12 or IL-4 cytokines (Allendoerfer and Deepe 1997, Clemons, Darbonne et al. 2000). Multiple studies show that CD4[+] and CD8[+] T cells play a vital role in *H. capsulatum* pulmonary and systemic infection. Particularly, it is the production of IFN-γ by CD4[+] T cells that drive macrophage activation and fungal clearance. Although granuloma-like formations were observed in the lungs of Rag-1[-/-] mice, these were likely predominantly composed of macrophages. However due to the lack of infiltrating CD4[+] and CD8[+] T cells, macrophages are unable to effectively kill intracellular yeasts which leads to the enhanced and uncontrolled disseminated infection and eventual death in Rag-1[-/-] mice. Although significant, an even greater decrease in Rag-1[-/-] IFN-γ was expected. This observation may be explained by the presence and increase of lymphoid NK cells in Rag-1[-/-] mice which can also produce IFN-γ. However, these levels may not be sufficient to arm all macrophages and inhibit the intracellular growth of yeasts. This together with the absence of CD4[+] and CD8[+] infiltrates surrounding granuloma formation may be the cause of increased dissemination and susceptibility of Rag-1[-/-] mice. IFN-γ immune therapy may be a promising area of research in the context of immunocompromised hosts.

We showed the myeloid cellular infiltrates in infected lung tissue at Day 14 post-infection of both wild-type and Rag-1[-/-] mice. Despite a significant increase in the percentage of neutrophils, alveolar macrophages and NK cells, Rag-1[-/-] mice developed exacerbated disease with increased fungal burden in all organs tested. Perhaps this increased cellular infiltration contributed to pathology and disease progression. While there was no significant difference in dendritic cells and inflammatory monocytes, the percentage of eosinophils was significantly reduced. Although neutrophils have been shown to be vital in the defense against some fungal

infections such as *Candida* spp. and *Aspergillus* spp., dimorphic fungi such as *H. capsulatum,*

Blastomyces dermatitidis and *Coccidioides immitis* do not rely on neutrophils (Romani 2011).

Interestingly, dendritic cells have been shown to reduce the transformation of *H. capsulatum*

from conidia to yeast form, therefore limiting the dissemination (Thind, Taborda et al. 2015).

Dendritic cells have also been shown to kill *H. capsulatum* yeast cells by phagosome-lysosome

fusion causing degradation of internalized yeasts (Thind, Taborda et al. 2015). Dendritic cells

are also known to be an important link between innate and adaptive immunity as they can

recognize and phagocytose fungi as well as present antigens to T cells. Dendritic cells may

play an important role in wild-type mice by recognizing and phagocytosing *E. africanus* yeasts,

resulting in IL-12 production and antigen presentation to T cells. Since there was only a subtle

decrease in dendritic cells recruited to Rag-1$^{-/-}$ mouse lungs, with no decrease in lung IL-12

production; we hypothesize that IL-12 is likely to be produced by macrophages during *E.*

africanus infection in Rag-1$^{-/-}$ mice. One limitation of this immunocompromised mouse model

is that we did not observe the presentation of cutaneous lesions which are observed clinically

in human infection. This may be specific to co-infections with HIV, or perhaps future studies

could look at setting up a dermal infection model, infecting mice by skin abrasions rather than

a pulmonary infection route. Despite this limitation, the animal model of *E. africanus* is an

indispensable step forward in understanding the pathogenesis and immune components

required for clearing infection.

Conclusion

Despite the prevalence and high mortality rates of invasive dimorphic fungal infections, they

continue to be misdiagnosed or underdiagnosed and are largely understudied. This work has

laid the foundation for a deeper understanding of *Emergomyces* spp. infection and provided

valuable insight into disease kinetics and immune components involved in fungal clearance

mechanisms in experimental mice. Case study reports indicate that people living with advanced HIV disease are more likely to develop serious symptomatic and disseminated emergomycosis requiring hospitalization. Late diagnosis leads to high fatality rates despite the initiation of amphotericin B treatment, with surviving patients requiring lengthy maintenance antifungal treatment (1-2 years). Here, we report that *E. africanus* infection drives a Th1 immune response in C57BL/6 mice. Importantly we showed that reduced IFN-γ production was associated with increased disease severity in immunocompromised Rag-1-deficient mice. Our model could be further utilized to explore new therapeutic approaches. Research focusing on immune therapy and possible vaccine development for endemic dimorphic fungal pathogens will have a tremendous impact on the treatment and management of the disease.

Materials and Methods

Mouse strains

Male and female mice between the age of 8-12 weeks (age and sex-matched), were used during *in vivo* experiments. Wild-type mice of different genetic backgrounds: C57BL/6, BALB/c and Sv129 mice from Jackson Laboratories were used. Disease progression and immune responses in wild-type C57BL/6 mice were compared with the immunocompromised Rag-1$^{-/-}$ mouse strain on a C57BL/6 background (Jackson Laboratories). All mice were housed in specific pathogen-free conditions in individually ventilated cages within the University of Cape Town biosafety level 2 (BSL2) animal unit facility.

Ethics statement

All the experiments performed were conducted in accordance with the Animal Research Ethics Committee of South African National Standard (SANS 10386:2008) and UCT, South Africa for the practice of animal procedures. The protocols were approved by the Animal Ethics Committee (permitted ethics protocol numbers AEC: 017/001 and 020/017), Faculty of Health Sciences, UCT, South Africa. Individuals involved in the study were accredited by the South African Veterinary Council for all animal experimental procedures and welfare monitoring.

***Emergomyces africanus* strain and growth conditions**

A clinical isolate of *E. africanus* (CBS 136260; accession no JX398291) was acquired from Nelesh Govender (National Institute for Communicable Diseases, South Africa) and was used in all experiments (Kenyon et al. 2013). Laboratory culture of *E. africanus* was performed as described by Kenyon et al., 2013 within biosafety level 2 facilities. Briefly, *E. africanus* was grown in the filamentous form at 25-29 °C on Difco TM Sabouraud Dextrose Agar (SDA) (Difco Laboratories; Becton Dickinson, and Company (BD)) and in the yeast form at 37 °C on

brain heart infusion (BHI) agar (Sigma-Aldrich, USA). The yeast phase suspension was

propagated by inoculating BHI broth followed by incubation at 37 °C with agitation at 180

RPM and harvested at a mid-log phase. Sterile aliquots of inoculum were stored at -80 °C in

BHI broth with 30 % glycerol, after counting the cells for further use. The stock concentration

of fungal cells was determined by serial dilution and plating of randomly selected vials.

***Emergomyces africanus* murine infection model**

E. africanus yeast inoculum was thawed and washed by centrifugation at 6000 RPM for 10

minutes (min) and the pellet was resuspended in sterile PBS. A serial dilution was performed

to obtain the desired concentration of cells per 50 µl directly before infection. Mice were

anaesthetized by intraperitoneal injection with ketamine and xylazine (80 mg/kg and 16 mg/kg

respectively) and 50 µl of the inoculum was administered via non-invasive intratracheal

inoculation. Inspection for humane endpoints in sick mice was highly regarded and adhered to

during the study. Mice showing weight loss greater than 20% of their baseline weight, or those

that showed signs of severe illness and pain (hunched back, lack of grooming) were euthanized.

Measuring fungal burden in mice

Fungal burden was determined in consistent lung, spleen, liver, and kidney tissue portions that

were harvested and weighed at various time points post-infection. Tissue sections included:

three right lung lobes, namely the superior, middle and lower lobes; a three-quarter section

from the posterior extremity of the spleen; the entire liver except for the left lobe (used for

histology); and the right kidney. After euthanasia, these organ sections were harvested

aseptically and stored in 1-2 ml 0.1 % PBS-Tween80 on ice followed by homogenization. Ten-

fold serial dilutions of the homogenates were made in 0.1 % PBS-Tween80 and 100 µl was

plated in duplicate at various dilutions on BHI agar. Plates were incubated at 37 °C for 10-12

days for colony counting. Colony-forming units (CFU) per gram of tissue were calculated and

presented to represent fungal burden and dissemination. The remaining lung and spleen

homogenates, after plating, were centrifuged at 10 000 RPM for 10 min at 4 °C and the

supernatants were collected, stored in aliquots and frozen at -80 °C for further cytokine

measurement by ELISA.

Tissue histopathology assessment

Consistent tissue sections of the lung (left lobe) were collected and immediately fixed in 4 %

(v/v) formaldehyde in PBS to preserve the tissue before embedding in paraffin wax and cut

into 2-5 μm sections. Staining was done using hematoxylin and eosin (H&E) to view the

alveolar spaces and inflammatory cells and Grocott's Methenamine Silver stain (GMS)

(Sigma-Aldrich, USA) to stain fungal cells in infected tissue. Photomicrographs of stained

tissue sections were captured using a Nikon Eclipse 90i microscope and DS-Ri2 high-

performance camera (Nikon) and Nikon NIS-Elements imaging software (v4.30.01 DU1).

Lung cellular recruitment by flow cytometry

Lung single-cell suspensions, 1 x 10^6 cells/reaction, were stained with myeloid cell markers of

neutrophils, inflammatory monocytes, eosinophils, alveolar macrophages, dendritic cells, and

natural killer cells and blocked with 1% rat serum and 1% anti-FcγRII/III. Cells were identified

as follows: neutrophils (CD11b$^+$, LY6G$^+$, LY6B$^+$), inflammatory monocytes (CD11b$^+$, LY6B$^+$,

SiglecF^{-neg}), eosinophils (CD11b$^+$, SiglecF$^+$, LY6B^{-neg}), alveolar macrophages (CD11c$^+$,

SiglecF$^+$, F4/80$^+$) and CD11b$^+$ dendritic cells (LY6G^{-neg}, CD11c$^+$, CD11b$^+$). Separate staining

for natural killer (NK) cell populations in both mouse strains; gating for NK cells: (CD3^{-neg},

NK1.1$^+$). Stained, and fixed cells were acquired using the LSR Fortessa™ (BD

Immunocytometry Systems, San Jose, CA, USA) and BD FACS Diva software (v6.0). Flowjo

software (v10.0.7) (Treestar, Ashland, OR USA) was used for post-acquisition analysis and

cell population determination. Refer to S1 Fig. for gating strategies and S1 Table for a detailed

list of antibodies used.

Cytokine analysis from lung homogenates

The supernatant from lung and spleen homogenates was used to measure the presence of

various cytokines that were released in infected tissue. The following BD OptEIA™ mouse

ELISA kits were used according to the manufacturer's instructions: IFN-γ, IL-1β, TNF, IL-6,

IL-10, IL-12p40 and IL-12p70. IL-17 (purified anti-mouse IL-17A clone: TC11-18H10.1;

biotin anti-mouse IL-17A clone: TC11-8H4) and GM-CSF (purified anti-mouse GM-CSF;

biotin anti-mouse GM-CSF) ELISA antibodies and standards were purchased separately from

Biolegend.

Serum total antibody ELISAs

Total IgM, IgG1, IgG2a, IgG2b, IgG3 and IgA antibody levels from both naïve and infected

mice were measured in sera by coating Nunc-Immuno Maxisorb 96-well plates with unlabeled

goat anti-mouse capture antibody at 1:500 dilution. Following blocking and incubation with

sera samples, total antibody isotype titers were detected with alkaline phosphatase-conjugated

rat anti-mouse antibodies at 1:1000 dilution. All antibodies used were produced by Southern

Biotech.

Statistical analysis

Two-tailed student's t-tests or One-way analysis of variance (ANOVA) tests were used to

determine statistical significance unless otherwise indicated. Results are graphically

represented as mean $\pm$ SD or SEM, with statistical significance represented as *: $p=0.05$; **:

$p=0.01$; ***: $p=0.001$ and ****: $p=0.0001$. All experiments were independently repeated at

least once unless otherwise indicated.

Acknowledgements: We thank Dr Barbra Topless for her initial training in *Emergomyces* spp.

culture techniques. We are grateful to Faried Abbass and Rodney Lucas for their valuable

technical assistance and the UCT animal unit staff for the running and maintenance of the

facility. We would like to thank Lizette Fick for her excellent technical assistance in histology

services and Prof Dhiren Govender for providing a description of histopathology. **Funding:**

This work was supported by the National Research Foundation (NRF) of South Africa,

Carnegie Corporation, Wellcome Trust (209293, 217163) and the Medical Research Council

(UK) Centre for Medical Mycology (MR/N006364/2). MAH was also supported by the

Oppenheimer Memorial Trust and LD supported by the University Research Council,

University of Cape Town.

Competing Interests: The authors have declared that no competing interests exist. All authors

declare no financial or personal conflict of interest regarding this article.

Authors contributions: JCH conceptualized, initiated, and funded the research. MAH planned

and executed the experiments, analyzed the data, and interpreted the results. MAH, LD and SS

contributed to sample preparation. RK assisted with technical expertise, data analysis and

interpretation of results. JCH and GDB contributed to the interpretation of results and

supervision of the project. NPG provided the *Emergomyces* clinical isolate and provided expert

advice for the design of the project. AB, ISS and NPG advised on establishing the protocols,

experimental design and provided critical expertise for the research. All authors provided

critical feedback on the manuscript.

References

Allen, H. L. (2005). "Apoptosis modulates protective immunity to the pathogenic fungus Histoplasma capsulatum." <u>Journal of Clinical Investigation</u> **115**(10): 2875-2885.

Allendoerfer, R. and G. S. Deepe (1997). "Intrapulmonary response to Histoplasma capsulatum in gamma interferon knockout mice." <u>Infection and Immunity</u> **65**(7): 2564-2569.

Allendoerfer, R. and G. S. Deepe, Jr. (1998). "Blockade of endogenous TNF-alpha exacerbates primary and secondary pulmonary histoplasmosis by differential mechanisms." <u>J Immunol</u> **160**(12): 6072-6082.

Allendörfer, R., G. D. Brunner and G. S. Deepe (1999). "Complex Requirements for Nascent and Memory Immunity in Pulmonary Histoplasmosis." <u>The Journal of Immunology</u> **162**(12): 7389.

Cano, L. E., L. M. Singer-Vermes, C. A. Vaz, M. Russo and V. L. Calich (1995). "Pulmonary paracoccidioidomycosis in resistant and susceptible mice: relationship among progression of infection, bronchoalveolar cell activation, cellular immune response, and specific isotype patterns." <u>Infection and Immunity</u> **63**(5): 1777-1783.

Chandler, J. W., Jr., T. K. Smith, W. M. Newberry, Jr., T. D. Chin and C. H. Kirkpatrick (1969). "Immology of the mycoses. II. Characterization of the immunoglobulin and antibody responses in histoplasmosis." <u>J Infect Dis</u> **119**(3): 247-254.

Clemons, K. V., W. C. Darbonne, J. T. Curnutte, R. A. Sobel and D. A. Stevens (2000). "Experimental histoplasmosis in mice treated with anti-murine interferon-gamma antibody and in interferon-gamma gene knockout mice." <u>Microbes Infect</u> **2**(9): 997-1001.

Cronjé, N., I. S. Schwartz, L. Retief, A. D. S. Bastos, S. Matthee, W. Preiser, N. C. Bennett, T.

Maphanga, N. P. Govender, R. Colebunders and C. Kenyon (2018). "Attempted molecular

detection of the thermally dimorphic human fungal pathogen Emergomyces africanus in

terrestrial small mammals in South Africa." Medical Mycology **56**(4): 510-513.

Deepe, G. S. D., L. Romani, V. L. G. Calich, G. Huffnagle, C. Arruda, E. E. I. W. Molinari-

Madlum and J. R. Perfect (2000). "Knockout mice as experimental models of virulence."

Medical Mycology **38**(s1): 87-98.

Deppe, G., S. and M. McGuinness (2006). "Interleukin-1 and Host Control of Pulmonary

Histoplasmosis." The Journal of Infectious Diseases **194**(6): 855-864.

Dukik, K., J. F. Muñoz, Y. Jiang, P. Feng, L. Sigler, J. B. Stielow, J. Freeke, A. Jamalian, B.

Gerrits Van Den Ende, J. G. McEwen, O. K. Clay, I. S. Schwartz, N. P. Govender, T. G.

Maphanga, C. A. Cuomo, L. F. Moreno, C. Kenyon, A. M. Borman and S. De Hoog (2017).

"Novel taxa of thermally dimorphic systemic pathogens in theAjellomycetaceae(Onygenales)."

Mycoses **60**(5): 296-309.

Heninger, E., L. H. Hogan, J. Karman, S. Macvilay, B. Hill, J. P. Woods and M. Sandor (2006).

"Characterization of theHistoplasma capsulatum-Induced Granuloma." The Journal of

Immunology **177**(5): 3303-3313.

Hoving, J. C., F. T. Munyonho and J. K. Kolls (2023). Genetic Mouse Models of Pneumocystis

Pneumonia. Antifungal Immunity: Methods and Protocols. R. A. Drummond. New York, NY,

Springer US: 169-179.

Jiang, Y., K. Dukik, J. F. Muñoz, L. Sigler, I. S. Schwartz, N. P. Govender, C. Kenyon, P.

Feng, B. G. Van Den Ende, J. B. Stielow, A. M. Stchigel, H. Lu and S. De Hoog (2018).

"Phylogeny, ecology and taxonomy of systemic pathogens and their relatives in

Ajellomycetaceae (Onygenales): Blastomyces, Emergomyces, Emmonsia, Emmonsiellopsis."

<u>Fungal Diversity</u> **90**(1): 245-291.

Jiang, Y., C. K. M. Tsui, S. A. Ahmed, F. Hagen, Z. Shang, A. H. G. Gerrits van den Ende, P.

E. Verweij, H. Lu and G. S. de Hoog (2020). "Intraspecific Diversity and Taxonomy of

Emmonsia crescens." <u>Mycopathologia </u>**185**(4): 613-627.

Kenyon, C., K. Bonorchis, C. Corcoran, G. Meintjes, M. Locketz, R. Lehloenya, H. F. Vismer,

P. Naicker, H. Prozesky, M. Van Wyk, C. Bamford, M. Du Plooy, G. Imrie, S. Dlamini, A. M.

Borman, R. Colebunders, C. P. Yansouni, M. Mendelson and N. P. Govender (2013). "A

Dimorphic Fungus Causing Disseminated Infection in South Africa." <u>New England Journal of

Medicine</u> **369**(15): 1416-1424.

Matthews, V. B., F. T. Christiansen and P. Price (2000). "Lymphocytes fromH2bmice produce

lower levels of several cytokines than congenicH2dorH2kmice." <u>Immunology and Cell

Biology</u> **78**(3): 247-253.

Romani, L. (2011). "Immunity to fungal infections." <u>Nature Reviews Immunology</u> **11**(4): 275-

288.

Schwartz, I. S., N. P. Govender, C. Corcoran, S. Dlamini, H. Prozesky, R. Burton, M.

Mendelson, J. Taljaard, R. Lehloenya, G. Calligaro, R. Colebunders and C. Kenyon (2015).

"Clinical Characteristics, Diagnosis, Management, and Outcomes of Disseminated

Emmonsiosis: A Retrospective Case Series." <u>Clinical Infectious Diseases</u> **61**(6): 1004-1012.

Schwartz, I. S., C. Kenyon, R. Lehloenya, S. Claasens, Z. Spengane, H. Prozesky, R. Burton,

A. Parker, S. Wasserman, G. Meintjes, M. Mendelson, J. Taljaard, J. W. Schneider, N. Beylis,

B. Maloba, N. P. Govender, R. Colebunders and S. Dlamini (2017). "AIDS-Related Endemic

Mycoses in Western Cape, South Africa, and Clinical Mimics: A Cross-Sectional Study of

Adults With Advanced HIV and Recent-Onset, Widespread Skin Lesions." <u>Open Forum</u>

<u>Infectious Diseases</u> **4**(4).

Schwartz, I. S., B. Lerm, J. C. Hoving, C. Kenyon, W. G. Horsnell, W. J. Basson, P. Otieno-

Odhiambo, N. P. Govender, R. Colebunders and A. Botha (2018). "Emergomyces africanus in

Soil, South Africa." <u>Emerging Infectious Diseases</u> **24**(2): 377-380.

Schwartz, I. S., T. G. Maphanga and N. P. Govender (2018). "Emergomyces: a New Genus of

Dimorphic Fungal Pathogens Causing Disseminated Disease among Immunocompromised

Persons Globally." <u>Current Fungal Infection Reports</u> **12**(1): 44-50.

Schwartz, I. S., J. D. McLoud, D. Berman, A. Botha, B. Lerm, R. Colebunders, E. Levetin and

C. Kenyon (2018). "Molecular detection of airborne Emergomyces africanus, a thermally

dimorphic fungal pathogen, in Cape Town, South Africa." <u>PLOS Neglected Tropical Diseases</u>

12(1): e0006174.

Schwartz, I. S., S. Sanche, N. P. Wiederhold, T. F. Patterson and L. Sigler (2018).

"Emergomyces canadensis,a Dimorphic Fungus Causing Fatal Systemic Human Disease in

North America." <u>Emerging Infectious Diseases</u> **24**(4): 758-761.

Sepúlveda, V. E., C. L. Williams and W. E. Goldman (2014). "Comparison of Phylogenetically

Distinct Histoplasma Strains Reveals Evolutionarily Divergent Virulence Strategies." <u>mBio</u>

5(4): e01376-01314-e01376.

Thind, S. K., C. P. Taborda and J. D. Nosanchuk (2015). "Dendritic cell interactions with

Histoplasma and Paracoccidioides." <u>Virulence</u> **6**(5): 424-432.

Thompson, G. R., 3rd, T. Le, A. Chindamporn, C. A. Kauffman, A. Alastruey-Izquierdo, N.

M. Ampel, D. R. Andes, D. Armstrong-James, O. Ayanlowo, J. W. Baddley, B. M. Barker, L.

Lopes Bezerra, M. J. Buitrago, L. Chamani-Tabriz, J. F. W. Chan, M. Chayakulkeeree, O. A.

Cornely, C. Cunwei, J. P. Gangneux, N. P. Govender, F. Hagen, M. T. Hedayati, T. M. Hohl,

G. Jouvion, C. Kenyon, C. C. Kibbler, N. Klimko, D. C. M. Kong, R. Krause, L. Lee Lee, G.

Meintjes, M. H. Miceli, P. M. Rath, A. Spec, F. Queiroz-Telles, E. Variava, P. E. Verweij, I.

S. Schwartz and A. C. Pasqualotto (2021). "Global guideline for the diagnosis and management

of the endemic mycoses: an initiative of the European Confederation of Medical Mycology in

cooperation with the International Society for Human and Animal Mycology." Lancet Infect

Dis **21**(12): e364-e374.

Wood, K. L., C. A. Hage, K. S. Knox, M. B. Kleiman, A. Sannuti, R. B. Day, L. J. Wheat and

H. L. Twigg (2003). "Histoplasmosis after Treatment with Anti–Tumor Necrosis Factor-α

Therapy." American Journal of Respiratory and Critical Care Medicine **167**(9): 1279-1282.

Zhou, P., M. C. Sieve, J. Bennett, K. J. Kwon-Chung, R. P. Tewari, R. T. Gazzinelli, A. Sher

and R. A. Seder (1995). "IL-12 prevents mortality in mice infected with Histoplasma

capsulatum through induction of IFN-gamma." The Journal of Immunology **155**(2): 785.

 Figure Legends

Fig 1. *Emergomyces africanus* **yeast cause disseminated infection in wild-type mice.** Three

wild-type mouse strains C57BL/6, BALB/c and Sv129 were infected with 0.5-1x10^5 yeast cells

in 50 µl PBS via oropharyngeal instillation. (A) A high dose infection resulted in 100%

mortality (determined when mice reached 20% weight loss) by 14 days post infection although

BALB/c mice were significantly more susceptible to infection when compared to C57BL/6 and

Sv129 mice; p<0.0001. (B) Weight loss of mouse strains measured. Data represents two

independent pooled experiments, n=8 /group per experiment, with mean ± SD, ***p<0.001

and ****p<0.0001, (one-way AVOVA). (C) Disseminated fungal infection was confirmed in

the lung, spleen, liver, and kidney by two weeks post-infection and yeast cells grew by plating

homogenized organs on brain heart infusion (BHI) agar and incubated at 37 °C for two-three

weeks. (D) Mold phase organisms were also isolated from tissue by culturing SDA plates at

29°C for 2-3 weeks. WT = wild-type.

Fig 2. Emergomycosis disease profile in wild-type mice. C57BL/6 mice were infected with

1x10^3 yeast cells in 50 µl PBS and euthanized at one-week intervals post-infection for four

weeks. Uninfected/naïve (UI) mice were used as a negative control. (A) C57BL/6 mice showed

a transient weight loss compared to uninfected mice. (B) Infection resulted in the dissemination

of organisms from the lung to the spleen, liver, and kidney shown by CFU per gram tissue. (C)

Macroscopic image of fixed lobes showed lesion/granuloma formation, at Day 14 and Day 21

post-infection. (D) Representative photomicrographs of lung sections stained with H&E with

original magnification: 2X. Data represents two individual experiments, n=5 mice/group per

experiment. UI: uninfected, H&E: hematoxylin and eosin.

Fig 3. Cytokine profile of infected lung tissue over time. C57BL/6 mice were infected with

1x10^3 yeast cells in 50 µl PBS and euthanized at one-week intervals post-infection. Supernatant

from homogenized lung was used to test for cytokine levels using ELISA for, IFN-γ, IL-1β,

IL-6, IL-12p40 and IL-17. Data represents two independent pooled experiments, n=4-5

mice/group per experiment, with mean ± SD, *p<0.05, ***p<0.001 and ****p<0.0001 (one-

way ANOVA using Dunnett's multiple comparisons test), C57BL/6 vs. uninfected mice. UI:

uninfected.

Fig 4. Serum total antibody production over time. Blood was collected during the disease

kinetic studies from Day 7 - Day 28 post-infection. Blood serum antibody levels were

determined by ELISA and compared to antibody levels from uninfected mice. Data represents

two independent pooled experiments, n=4-5 mice/group per experiment, with mean ± SD and

*p<0.05, **p<0.01 and ***p<0.001 (one-way ANOVA using the nonparametric Kruskal-

Wallis test and Dunn's multiple comparisons test), C57BL/6 vs. uninfected mice. UI:

uninfected.

Fig 5. Immunocompromised Rag-1-deficient mice are susceptible to *E. africanus*

infection. C57BL/6 and Rag-1$^{-/-}$ mice were infected with 1x10^3 yeast cells and (A) Survival

was compared. (B) In addition, infected C57BL/6 and Rag-1$^{-/-}$ mice were killed at Day 14 post

infection and fungal burden was determined by CFU/gram of tissue in lung, spleen, liver and

kidney. (C) Representative photomicrographs are shown of lung sections stained with H&E

from C57BL/6 and Rag-1$^{-/-}$ mice at Day 14 post infection with original magnification: 2X. Data

represents two independent pooled experiments, with n=8-10 mice/group for survival and 5-6

mice /group for CFU and histology, mean ± SD, ***p<0.001 and ****p<0.0001 C57BL/6 vs

Rag-1$^{-/-}$ (student t-test or one-way AVOVA).

Fig 6. Myeloid cell recruitment to infected lung tissue after *E. africanus* infection.

C57BL/6 and Rag-1$^{-/-}$ mice were infected with 1x10^3 yeast cells and lungs harvested at Day 14

post infection. (A) Representative myeloid cell population scatter plots from both C57BL/6

and Rag-1-/- mice. (B) Percentage cells shown for myeloid cell populations identified as

follows: neutrophils (CD11b$^+$, LY6G$^+$, LY6B$^+$), inflammatory monocytes (CD11b$^+$, LY6B$^+$,

SiglecF^{-neg}), eosinophils (CD11b$^+$, SiglecF$^+$, LY6B^{-neg}), alveolar macrophages (CD11c$^+$,

Siglecf $^+$, F4/80$^+$) and CD11b$^+$ dendritic cells (LY6G^{-neg}, CD11c$^+$, CD11b$^+$). Separate staining

for NK cell populations in both mouse strains; (CD3^{-neg}, NK1.1$^+$). Data represents two pooled

experiments, n=4-5 mice/group per experiment, mean $\pm$ SD, **p<0.01 and ****p<0.0001

C57BL/6 vs. Rag-1$^{-/-}$ (student t-test). FMO: fluorescence minus one; NK: natural killer; ns =

not significant

Fig 7. Lung cytokine production in Rag-1-deficient mice in response to *E. africanus*

infection. Homogenized lung supernatant was used to test for cytokine levels at Day 14 post

infection using ELISA. Data represents two pooled individual experiments, mean $\pm$ SD, n=5-

6 mice/group per experiment, and *p<0.05 C57BL/6 vs Rag-1$^{-/-}$, student t-test.

Supplementary Figure legends

Supp Fig S1. Histopathology of disease progression in wild-type mice. C57BL/6 mice were infected with 1×10^3 yeast cells in 50 µl PBS and euthanized at weekly intervals post-infection. Sections were stained with H&E for inflammation and represented photomicrographs are shown for 10x and 20x magnification. (B) Representative images of lung sections stained with GMS at 10x and 20x magnification.

Supp Fig S2. Fungal distribution in the lungs of wild-type mice. C57BL/6 mice were infected with 1×10^3 yeast cells in 50 µl PBS and euthanized at weekly intervals post-infection. Sections were stained with GMS for fungal organisms and representative photomicrographs are shown for 10x and 20x magnification. GMS: Grocott methenamine silver.

Supp Fig S3. *Emergomyces africanus* conidia cause disseminated infection in wild-type mice. C57BL/6 mice were infected with 1×10^3 conidia cells in 50 µl PBS and euthanized at Day 14 post-infection. Conidia infection resulted in the dissemination of organisms from the lung to the spleen, liver, and kidney shown by CFU per gram tissue. Data represents two independent pooled experiments, n=6 mice per experiment.

Supp Fig S4. Histopathology and fungal distribution in the lungs of Rag-1-deficient mice. C57BL/6 and Rag-1$^{-/-}$ mice were infected with 1×10^3 yeast cells and euthanized at Day 14 post infection. Representative images of lung sections from C57BL/6 and Rag-1$^{-/-}$ mice stained with H&E and GMS at 10x and 20x magnification. H&E: hematoxylin and eosin; GMS: Grocott methenamine silver.

Supp Fig S5. Myeloid flow cytometry gating strategy for infected lung tissue. C57BL/6 and Rag-1$^{-/-}$ mice were infected with 1×10^3 yeast cells. Lungs were harvested at Day 14 post-infection. The myeloid gating strategy for infected lung tissue was initiated by excluding cell

debris and gating for single cells. Live leukocytes were identified as CD45$^+$ and Zombie Red-

. Myeloid cell populations were gated as follows: neutrophils (CD11b$^+$, LY6G$^+$, LY6B$^+$),

inflammatory monocytes (CD11b$^+$, LY6B$^+$, SiglecF^{-neg}), eosinophils (CD11b$^+$, SiglecF$^+$,

LY6B^{-neg}), alveolar macrophages (CD11c$^+$, SiglecF$^+$, F4/80$^+$) and CD11b$^+$ dendritic cells

(LY6G^{-neg}, CD11c$^+$, CD11b$^+$) (B) (see below).

Supp Fig S6. Natural killer cell flow cytometry gating strategy. C57BL/6 and Rag-1$^{-/-}$ mice

were infected with 1×10^3 yeast cells. Lungs were harvested at Day 14 post-infection. Separate

staining was done to identify NK cell populations in both mouse strains; gating for NK cells:

(CD3^{-neg}, NK1.1$^+$). FMO: fluorescence minus one; NK: natural killer.

Supp Fig S7. Absolute number of myeloid cell recruitment to infected lung tissue after *E.*

***africanus* infection.** C57BL/6 and Rag-1$^{-/-}$ mice were infected with 1×10^3 yeast cells and lungs

harvested at Day 14 post-infection. Absolute cell numbers shown for myeloid cell populations

identified as follows: neutrophils (CD11b$^+$, LY6G$^+$, LY6B$^+$), inflammatory monocytes

(CD11b$^+$, LY6B$^+$, SiglecF^{-neg}), eosinophils (CD11b$^+$, SiglecF$^+$, LY6B^{-neg}), alveolar

macrophages (CD11c$^+$, Siglecf $^+$, F4/80$^+$) and CD11b$^+$ dendritic cells (LY6G^{-neg}, CD11c$^+$,

CD11b$^+$). Separate staining for NK cell populations in both mouse strains; gating for NK cells:

(CD3^{-neg}, NK1.1$^+$). Data represents three-four pooled experiments, n=4-5 mice/group per

experiment, mean $\pm$ SD, *p<0.05 and ***p<0.001 C57BL/6 vs. Rag-1$^{-/-}$ (student t-test). FMO:

fluorescence minus one, NK: natural killer.

Figure 1

A

p<0.0001

Survival (%)

C57BL/6
BALB/c
Sv129

Days post infection

B

Body weight (%)

***

****

***

C57BL6
BALB/c
Sv129

Days post infection

C

| Lung | Spleen | Liver | Kidney |

Yeast

D

| Lung | Spleen | Liver | Kidney |

Conidia

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Supp Figure S1

Supp Figure S2

Supp Figure S3

Supp Figure S4

Lung Day 14 post infection

C57BL/6

Rag-1⁻/⁻

H&E

GMS

Supp Figure S5

Supp Figure S6

Supp Figure S7

Supplementary Table S1: Flow cytometry antibody information.

Target Protein	Species	Conjugation	Clone	Company
Ly6G	rat anti-mouse	APC	1A8	BD Pharmingen
CD11c	hamster anti-mouse	Alexa Fluor® 700	HL3	BD Pharmingen
CD11b	rat anti-mouse	V450	M1/70	BD Horizon
neutrophil (7/4)	rat anti-mouse	FITC	GR288279-11	Abcam
CD45	rat anti-mouse	BV510	30-F11	BD Horizon
F4/80	rat anti-mouse	PE/Cy7	BM8	Biolegend

*Zombie Red™ Fixable Viability Kit (Biolegend, USA). **Blocking agents were added to antibody cocktails to prevent non-specific binding of the antibodies: eBioscience™ normal rat serum (Invitrogen, USA) and mouse BD Fc block™ rat anti-mouse CD16/CD32 antibody (clone:2.4G2, BD Pharmingen).