Plasma Growth Hormone Pulsatility Directly Regulates Male-biased Chromatin Accessibility in Adult Mouse Liver

Andy Rampersaud, Jeannette Connerney, and David J. Waxman

Department of Biology and Bioinformatics Program
Boston University, Boston, MA 02215 USA

Keywords: DNase-I hypersensitive sites, STAT5b tyrosine phosphorylation, hepatocyte open chromatin, sex differences in liver disease, growth hormone-regulated enhancers

Running title: GH pulse regulation of liver chromatin accessibility

Address correspondence to:
Dr. David J. Waxman
Dept. of Biology, Boston University
5 Cummington Mall
Boston, MA 02215 USA
Email: djw@bu.edu
ORCID #: 0000-0001-7982-9206

Summary - Sex-differences in plasma growth hormone (GH) profiles, pulsatile in males and persistent in females, regulate sex differences in hepatic STAT5 activation linked to sex differences in gene expression and liver disease susceptibility, but little is understood about the fundamental underlying, GH pattern-dependent regulatory mechanisms. Here, DNase hypersensitivity site (DHS) analysis of liver chromatin accessibility in a cohort of 21 individual male mice established that the endogenous male rhythm of plasma GH pulse-stimulated liver STAT5 activation induces repeated cycles of chromatin opening and closing at several thousand liver DHS and comprises one of two distinct mechanisms conferring male bias to liver chromatin accessibility. Sex-dependent transcription factor binding patterns and chromatin state analysis identified key factors and epigenetic features distinguishing this dynamic, STAT5-driven mechanism of male-biased chromatin opening from that operative at static male-biased DHS, which are constitutively open in male but not female liver. Notably, dynamic but not static male-biased DHS adopt a bivalent epigenetic state in female liver, as do female-biased DHS in male liver, albeit using distinct repressive histone marks in each sex (H3K27me3 at female-biased DHS in male liver, H3K9me3 at male-biased DHS in female liver). Strikingly, a single physiological replacement dose of GH given to hypophysectomized male mice restored, within 30 min, liver STAT5 activity and chromatin accessibility at 83% of the dynamic male-biased DHS that closed following pituitary hormone ablation. Pulsatile chromatin opening stimulated by endogenous, physiological hormone pulses is thus a novel mechanism for establishing widespread sex differences in chromatin accessibility and transcription factor binding, which are closely linked to sex-biased gene expression and the sexual dimorphism of liver function.

Introduction

Growth hormone (GH) regulates hepatic expression of enzymes and transporters that play critical roles in lipid metabolism and in the detoxification of many drugs and other lipophilic foreign chemicals [1, 2]. Dysregulation of hepatic GH signaling can lead to liver metabolic disorders, including the development of fatty liver disease and non-alcoholic steatohepatitis, with males more susceptible than females, as seen in both mice and humans [3-5]. Correspondingly, GH regulates many liver-expressed genes in a sex-dependent manner, enabling each sex to meet its specific metabolic and hormonal requirements [6]. This program of sex-dependent gene expression is controlled by the sex-dependent temporal patterns of pituitary GH secretion [7, 8], which emerge at puberty but are programmed earlier in life by neonatal exposure to androgen [9-11]. Pituitary GH secretion, and plasma GH profiles, are intermittent (pulsatile) in pubertal and adult males, whereas they are near-continuous (persistent) in pubertal and adult females, as seen in rats, mice and humans [12]. These plasma GH profiles, in turn, regulate sex-specific gene transcription in the liver through both positive regulatory mechanisms (class I sex-biased genes) and negative regulatory mechanisms (class II sex-biased genes) [6].

The sex-dependent hepatic actions of GH require the transcription factor STAT5 [13], which is activated by phosphorylation on a single tyrosine residue catalyzed by the GH receptor-associated tyrosine kinase JAK2 [14]. Each successive male plasma GH pulse induces a cycle of STAT5 tyrosine phosphorylation, dimerization and nuclear translocation, followed by STAT5 tyrosine dephosphorylation and return back to the cytosol in time to reset the overall signaling pathway for the subsequent plasma GH pulse [1]. In male mouse liver, GH pulse-activated STAT5 binds to the DNA motif TTCNNNGGA at open chromatin regions strongly enriched for proximity to male-biased genes, but the relationship between STAT5 binding, chromatin opening and sex differences in chromatin accessibility is unknown. In female liver, persistent activation of STAT5 by the near-continuous presence of circulating GH is associated with a significant enrichment of STAT5 binding nearby female-biased genes [15], however, the underlying mechanisms, including chromatin features that distinguish these STAT5 binding sites from male-biased STAT5 binding sites are poorly understood.

Liver STAT5 is activated by male plasma GH pulses within minutes, enabling STAT5 to rapidly induce the transcription of several male-biased genes [16]; however, a majority of male-biased genes respond slowly to changes in GH secretory patterns [17], likely reflecting the time required for secondary changes, including changes in histone modifications and the underlying chromatin state [18]. Continuous infusion of GH in male mice overrides the endogenous plasma GH pulses and substantially feminizes liver gene expression over a period of days, with individual female-biased genes already in an active chromatin state in male liver generally responding earlier than genes in an inactive chromatin state [17]. These chromatin state changes include changes in chromatin accessibility at genomic regions discovered as DNase-I hypersensitive sites (DHS), a hallmark of epigenetic regulation. DHS can be assayed by DNase-seq, which has identified ~70,000 DHS in male and female mouse liver [19], including thousands of enhancers, promoters, and insulator regions [20]. More than 4,000 of the 70,000 liver DHS show sex differences in chromatin accessibility, as well as sex-biased binding of STAT5 and other GH-regulated transcription factors, which reinforce sex differences in liver gene expression [18, 19] and regulate downstream sex differences in disease susceptibility [21-23]. Importantly, male-biased transcription factor binding is strongly enriched at male-biased DHS nearby male-biased genes, and female-biased transcription factor binding is strongly enriched at female-biased DHS nearby female-biased genes. Sex differences in chromatin structure and accessibility are thus key features of sex-differential liver gene expression [17, 18]. However, little is understood about the mechanisms linking the sex-dependent temporal GH secretion patterns to the robust sex differences in chromatin accessibility and transcription factor binding that regulate liver gene expression.

Here, we elucidate the relationship between the sex-dependent patterns of plasma GH stimulation of hepatocytes and sex differences in liver chromatin accessibility. We identify more than 800 male-biased enhancer DHS regions, where the direct binding of plasma GH pulse-activated STAT5 induces a dynamic cycle of male liver chromatin opening and closing at sites that comprise 31% of all male-biased DHS. Thus, the pulsatility of plasma GH stimulation *per se* confers significant male bias in chromatin accessibility and STAT5 binding at a substantial fraction of the genomic sites linked to sex-biased liver gene expression and liver disease. Furthermore, we establish that a single physiological replacement dose of GH given as a pulse to

hypophysectomized (hypox) mice recapitulates, within 30 min, the pulsatile re-opening of chromatin seen in
pituitary-intact male mouse liver. We elucidate key epigenetic features distinguishing this dynamic, STAT5-
driven mechanism of male-biased chromatin opening from that operative at a second, distinct class comprised of
static male-biased DHS, which are constitutively open in male liver but closed in female liver. Finally, our
analysis of repressive histone marks enriched at each class of sex-biased DHS indicates that distinct epigenetic
mechanisms mediate sex-specific gene repression in each sex. Together, these studies identify pulsatile
chromatin opening as a novel mechanism controlling sex differences in chromatin accessibility and transcription
factor binding closely linked to sex differences in gene expression and liver disease.
107
108   **Results**
109
Sex-biased DHS are primarily distal enhancers that target sex-specific genes – Open (accessible) chromatin
regions identified in mouse liver by DNase-seq analysis have been classified as enhancers, insulators and
promoters based on their chromatin mark patterns and CTCF binding activities (n = 70,211 standard reference
DHS set) [20] (Table S2A). Several thousand of these DHS show greater chromatin accessibility in male than
female liver (n = 2,729 male-biased DHS) or greater accessibility in female than male liver (n = 1,366 female-
biased DHS) [19]. These sex-biased DHS regions were enriched for enhancer marks, with >85% classified as
enhancers or weak enhancers in mouse liver, as compared to 66% for all sex-independent DHS (Fig. 1A).
Assignment of the sex-biased liver DHS to their putative gene targets (closest RefSeq gene or multi-exonic
lncRNA gene transcription start site in the same topologically associating domain (TAD) [20] revealed that sex-
biased DHS were highly enriched for genes of the corresponding sex bias, but not for genes of the opposite sex
bias (Table S2B). Cumulative plots of the distance from each DHS to its target gene revealed that a majority of
sex-biased DHS classified as enhancers or insulators are distal to their target genes. Sex-biased DHS with
H3K4me3 marks, which comprise only 1-2% of all sex-biased DHS, are proximal to their target genes,
validating their classification as promoter DHS (Fig. 1B). Thus, a majority of sex-biased DHS are marked as
positively acting distal enhancers, consistent with our finding that sex-dependent DNA looping at an intra-TAD
scale is common in mouse liver [24].
126
No major differences in chromatin state distributions between male-biased, female-biased and sex-independent
enhancer DHS were seen based on separate chromatin state maps developed for male and female liver [18].
Thus, all three sets of enhancer DHS showed a high frequency (50-66%) of chromatin state E6, whose emission
probabilities (Fig. 1C) indicate a high frequency of DHS in combination with the activating chromatin marks
H3K27ac and H3K4me1, and a lower frequency (12-16%) of chromatin state E5 (high frequency of DHS but
low frequency of H3K27ac and H3K4me1) (Fig. 1D). Sex-biased and sex-independent insulator DHS also
showed similar chromatin state distributions (state E5 ~ state E6), as did sex-biased and sex-independent
promoter DHS, which were primarily in state E7, characterized by a high frequency of H3K4me3 and H3K4me1
marks (Fig. 1D).
136
137   The absence of major differences in overall chromatin state distributions between male-biased and female-biased
138   liver DHS prompted us to investigate other factors that may provide insight into underlying mechanisms
139   regulating sex-differences in hepatic chromatin accessibility and their link to sex differences in gene expression.
140
Impact of endogenous pulses of GH and STAT5 activity in male liver - Sex differences in pituitary GH
secretion – pulsatile in males vs. near continuous (persistent) in females – regulate the sex-dependent expression
of hundreds of genes in adult mouse liver. This regulation requires the GH-activated transcription factor STAT5
[25, 26]. We hypothesize that the pulsatile activation of STAT5 seen in male liver in direct response to plasma
GH stimulation [15, 16, 27] dynamically alters the chromatin accessibility landscape of male mouse liver. More
specifically, we propose that the repeated activation of STAT5 in male mouse liver by plasma GH pulses induces
dynamic cycles of chromatin opening and closing at a subset of liver DHS, and that this response can be
discovered by comparing chromatin accessibility profiles in livers of individual male mice euthanized at a peak
vs. at a trough of hepatic STAT5 activity (Fig. 2A, *top*). To test this hypothesis, we analyzed DNase-seq libraries
prepared from liver nuclei purified from 21 individual adult male mice. In parallel, we determined the STAT5
DNA-binding activity of each liver by EMSA (electrophoretic mobility shift analysis) of whole liver extracts.
Ten of the 21 livers had high STAT5 EMSA activity (STAT5-high activity livers) and 11 livers had very low or

no detectable STAT5 EMSA activity (STAT5-low activity livers) (Fig. 2B, Fig. S1). Next, we performed
DNase-seq analysis on genomic DNA fragments released by light DNase-I digestion of nuclei purified from each
liver, followed by diffReps analysis [28] comparing the DNase-seq-released DNA fragments from each group of
livers. We thus discovered genomic regions showing significant differential chromatin accessibility between
STAT5-high and STAT5-low male livers. Principal component analysis using either the top 200 or the top 600
most significant diffReps-identified differentially accessible regions revealed that 18 of the 21 livers yielded
consistent, STAT5 activity-dependent patterns of DNase-released fragments. Two of the STAT5-high livers and
one of the STAT5-low livers were outliers (Fig. 2C). These outliers were also identified by their discordant
DNase-seq read count distributions (Fig. 2D) and were excluded from all downstream analyses.
162
We re-analyzed the remaining 18 STAT5-high and STAT5-low livers using diffReps then filtered the output list
to retain regions identified as DHS peak regions by MACS2 analysis of the same 18 DNase-seq datasets. We
thus identified n=2,832 genomic sites where chromatin opening is associated with high liver STAT5 activity and
n=123 other sites where chromatin opening is associated with low liver STAT5 activity (Table S4A). As each
liver represents a time point of peak (STAT5-high livers) or trough (STAT5-low livers) levels of GH pulse-
activated liver STAT5 activity (Fig. 2A), the STAT5-high/STAT5-low differential sites identify genomic regions
where chromatin dynamically opens or closes in male mouse liver in response to GH pulse activation of STAT5.
The greater chromatin opening in STAT5-high livers as compared to STAT5-low male livers, and compared to
female livers, was visualized in aggregate plots of normalized DNase-seq cuts across the 2,832 genomic regions
(Fig. 2E, *left*). In contrast, chromatin accessibility was greater in the STAT5-low livers at the 123 STAT5-low
diffReps-identified genomic sites, where STAT5 activation is associated with chromatin closing (Fig. 2E, *right*).
This binary pattern, where individual male mouse livers show either high or low DNase-seq read count
distributions at the top differential genomic sites, was also seen in an independent set of nine male liver DNase-
seq samples from a second mouse strain generated by the ENCODE consortium: 4 of the 9 livers showed read
distributions very similar to the STAT5-high livers, while 5 of the 9 livers were similar to the STAT5-low livers
(Fig. 2D, *green bars*).
179
Dynamic vs static liver DHS – Comparison of the set of 2,832 STAT5-high genomic sites with our standard
reference set of 70,211 liver DHS revealed that 31% (n=834) of the 2,729 male-biased liver DHS described
above are STAT5-high sites, i.e., they respond dynamically to endogenous male plasma GH pulses of liver
STAT5, whereas only 0.5% of female-biased liver DHS and 2% of sex-independent liver DHS showed this
dynamic STAT5 response (Fig. 2F, *green*). This strong enrichment of GH/STAT5 pulse-induced chromatin
opening at male-biased DHS as compared to sex-independent DHS (ES = 12.9; p < 1E-05) identifies intermittent
chromatin opening induced by male plasma GH pulses as a mechanism that can explain the male bias in
chromatin accessibility for a significant subset (31%) of male-biased DHS.
188
To determine whether STAT5 binding *per se* is an important feature of the observed pulsatile changes in
chromatin accessibility, we examined normalized DNase-I cut site aggregate plots for the subset comprised of
n=1,307 male-biased DHS that bind STAT5 in ChIP-seq analyses. STAT5-high livers showed the highest level
of chromatin opening, followed by up to a 2.8-fold lower level of chromatin opening at those same genomic
regions in STAT5-low male livers and in female livers (Fig. 3A, plot 1). In contrast, male-biased DHS that did
not bind STAT5 (n=1,422 DHS) showed nearly equal chromatin accessibility in STAT5-high vs. STAT5-low
male livers but ~2-fold lower accessibility in female livers (Fig. 3A, plot 2). Thus, the male bias in accessibility
at the STAT5-bound but not at the non-STAT5 bound male-biased DHS can be explained by STAT5-induced
pulsatile chromatin opening in male liver. Moreover, it is also apparent that the male bias of the latter DHS set is
associated with chromatin closing in female liver (Fig. 3A, plot 2, *black*). The latter conclusion is supported by
the low mean normalized DNase cutting frequency at the 1,422 male-biased DHS in female liver when compared
to that of the full genome-wide set of 53,404 STAT5-unbound, sex-independent DHS (Fig. 3A, plot 2, *black*
versus plot 4). Chromatin accessibility was much higher at the subset of sex-independent DHS that bound
STAT5 (n=12,712 DHS, plot 3). Strikingly, the high chromatin accessibility at these sex-independent DHS was
seen in livers of both sexes and was largely invariant between STAT5-high and STAT5-low male livers. We
conclude that STAT5 binding is associated with increased chromatin opening, and that dynamic chromatin
opening and closing associated with STAT5 binding is a defining characteristic of a distinct subset of male-
biased DHS but occurs infrequently at female-biased and sex-independent DHS.

207
The above findings allow us to identify two distinct subsets of male-biased DHS, which differ in their
responsiveness to liver STAT5 activation: 1) dynamic male-biased DHS are characterized by a male bias in
chromatin accessibility linked to an increase in chromatin opening following the binding of STAT5 when
activated by a plasma GH pulse in male liver every 3-4 h [1, 16] (n=834 dynamic male-biased DHS, 31%); and
2) static male-biased DHS are open in male liver constitutively, i.e., throughout the pulsatile, on/off cycles of
GH-induced STAT5 activity, and are comparatively closed in female liver (n=1,895 static male-biased DHS,
69%). Consistent with this proposal, mean chromatin opening at the set of 834 dynamic male-biased DHS was
3.8-fold higher in STAT5-high male livers than in STAT5-low male livers or in female livers (Fig. 3A, plot 5).
In contrast, the higher chromatin accessibility in male than female liver at the set of 1,895 static male-biased
DHS was largely independent of the male liver's STAT5 activity status. Of note, the mean level of chromatin
opening at those 1,895 sites was 2.6-fold lower than the peak level of accessibility seen at the dynamic male-
biased DHS regions in STAT5-high male livers (Fig. 3A, plot 6 vs. plot 5) (**Table 1A**).
220
221 Chromatin accessibility at the set of static female-biased DHS (1,359 sites; Fig. 2F) was 2.7 to 2.9-fold higher in
222 female liver than in male liver, where chromatin accessibility was independent of male plasma GH/STAT5
223 pulses (Fig. 3A, plot 7). The aggregate female liver accessibility of these static female-biased DHS was very
224 similar to that of the genome-wide set of 64,584 static sex-independent DHS (Fig. 3A, plot 8 vs. plot 7; peak
225 normalized DNase-I activity value of 699 vs 683, Table S6).
226
STAT5 binding is closely associated with dynamic male-biased DHS - The presence of a canonical STAT5
motif, TTC-NNN-GAA, is a distinguishing feature of dynamic male-biased DHS: 81% of 834 dynamic male-
biased DHS contained a STAT5 motif versus only 38% of 1,895 static male-biased DHS (c.f., 25% background
STAT5 motif frequency for genome-wide set of sex-independent static DHS) (Fig. 3B). In addition, multiple
STAT5 motifs are more frequently found at dynamic DHS than at static DHS (Fig. 3C). Consistently, STAT5
binding, determined experimentally by ChIP-seq, occurs at a greater fraction of dynamic than static male-biased
DHS (85% versus 32%; Fig. S2A), and the level of STAT5 binding (normalized STAT5 ChIP-seq read counts)
was significantly higher at the STAT5-bound subsets of the dynamic vs. static DHS sets (Fig. S2B). *De novo*
motif discovery supported these findings, with top scoring motifs matching the STAT5B motif found in dynamic
DHS (Fig. S3A, Fig. S3B) but not in static DHS sets (Fig. S3C-S3E). A close association between STAT5
binding and chromatin opening is also indicated by the higher chromatin accessibility at the DHS subsets
associated with STAT5 binding (Fig. S2C, *top row* vs. *bottom row*).
239
These findings establish a close link between STAT5 binding, which is pulsatile in male liver [15], and the
repeated opening and closing of chromatin at dynamic DHS. Nevertheless, pulsatile chromatin opening was also
found at 124 male-biased DHS (15% of all dynamic male-biased DHS) that did not bind STAT5 (Fig. S2C, plot
1B). Moreover, STAT5 binding alone is not sufficient to insure dynamic, male-biased chromatin opening and
closing, insofar as 90% of all sex-independent DHS that bind STAT5 do not undergo dynamic chromatin
opening and closing (Fig. S2C, plot 5A (11,473 sites, 90.3%) vs plot 4A (1,239 sites, 9.7%)). Furthermore, only
54% of male-biased DHS that bind STAT5 are dynamic DHS (710 sites), the other 46% (597 sites) being static
male-biased DHS, even though they bind STAT5 (Fig. S2C, plot 1A vs 2A), albeit at a lower level than the
dynamic male-biased DHS and similar to the STAT5-bound static sex-independent DHS (Fig. S2B). Of note, we
observed a 5.6-fold higher frequency of dynamic chromatin opening and closing at the male-biased DHS with
STAT5 bound (54%) than at the sex-independent DHS with STAT5 bound (9.7%). Thus, genomic features other
than pulsatile STAT5 binding *per se* are required for dynamic chromatin opening and closing to occur.
252
Enrichment of other GH-regulated liver transcription factors at dynamic vs static DHS – We investigated
whether dynamic and static male-biased DHS differ with respect to the binding of other, sex-biased transcription
factors (**Table S1B**). We considered two GH/STAT5-regulated transcriptional repressors that reinforce STAT5
regulation of liver sex differences. One factor, BCL6, is a male-biased protein that can compete for STAT5
binding to chromatin and preferentially represses female-biased genes in male liver [15, 18, 22, 29]. The second
factor, CUX2, is a female-specific repressor whose binding sites in female liver are enriched nearby male-biased
genes, which enables CUX2 to repress those genes in female liver [30, 31]. Both repressors showed significant
enrichment for binding to genomic regions defined by male-biased DHS as compared to a background set of

static sex-independent DHS. Thus, male-biased BCL6 binding was significantly enriched at the dynamic male-
biased DHS (ES=2.0, p=E-14) but showed minimal enrichment at static male-biased DHS (ES=1.3, p=1.6E-04).
BCL6 binding was also enriched at dynamic sex-independent DHS (Table S7B). In contrast, CUX2 was most
highly enriched for binding in female liver at genomic sites that correspond to static male-biased DHS (ES=4.6,
p=E-40). This binding of CUX2 may contribute to the greater closure of those DHS in female as compared to
male liver (c.f., Fig. 3A, plot 6). Finally, male-biased STAT5 binding sites showed 2.5-fold greater enrichment at
dynamic than at static male-biased DHS (ES=58, p=E-301 vs ES=24, p=E-190) (Table S7B), consistent with our
findings, above, implicating pulsatile STAT5 binding in dynamic chromatin opening at those sites.
269
We reanalyzed published mouse liver ChIP-seq data for FOXA1 and FOXA2 [32], pioneer factors implicated in
chromatin opening, [33, 34], to identify sex-dependent binding sites for each factor. Male-biased binding sites
discovered for each FOXA factor showed strong enrichment for binding at male-biased DHS, with ~2-fold
higher enrichment seen at the static male-biased DHS (for FOXA1: ES=6.1 (static DHS) vs. ES=3.7 (dynamic
DHS); for FOXA2: ES=44 (static DHS) vs. ES=21 (dynamic DHS) (**Table S1B**). Consistent with this, *de novo*
motif discovery identified a Fox family factor as a close match for one of the top enriched motifs in the set of
static but not in the set of dynamic male-biased DHS (Fig. S3C vs Fig. S3A, p-value 9.2E-06 vs. 2.0E-03).
Finally, female-biased FOXA2 binding sites, but not female-biased FOXA1 binding sites, showed strong
enrichment for static female-biased DHS (ES=18, p=E-75; Table S7B). Taken together, these findings support
the proposal that FOXA2, and to a lesser extent FOXA1, contribute to sex-dependent chromatin opening, in
particular at static DHS.
281
Impact of hypophysectomy on liver chromatin accessibility – Surgical removal of the pituitary gland
(hypophysectomy) ablates pituitary GH secretion and thereby abolishes liver STAT5 activation [16], leading to
widespread loss of sex-specific liver gene expression [6]. We hypothesized that by ablating GH-induced STAT5
activation and DNA binding, hypophysectomy will lead to closure of many of the open chromatin regions that
regulate sex-specific gene expression. Furthermore, we proposed that restoration of a pulsatile GH signal, in the
form of a single exogenous GH pulse given by i.p. injection, will reopen chromatin at many of those sites (Fig.
4A). To test this hypothesis, we used DNase-seq to compare the chromatin accessibility profiles of liver nuclei
from hypox male and female mice to those of pituitary-intact control liver nuclei. Hypophysectomy induced
changes in chromatin accessibility at several thousand sites, including large numbers of sex-independent DHS,
with many more genomic regions undergoing chromatin closing than chromatin opening in male liver, but not in
female liver (Fig. 4B, Table S4). Importantly, male-biased DHS that responded to hypophysectomy were almost
exclusively closed in male liver following hypophysectomy (Fig. 4C, *first two rows*). A much smaller percentage
of static female-biased DHS responded to hypophysectomy, and the responses observed generally involved
chromatin opening in male liver and chromatin closing in female liver (Fig. 4C, Table S4G). Thus, sex-biased
DHS are subject to both positive and negative pituitary hormone regulation and respond differently between the
sexes.
298
DHS responses to pituitary ablation link to class I and class II sex-biased gene regulatory responses -
Hypophysectomy abolishes GH-regulated liver sex differences via two distinct mechanisms, which identify two
classes of sex-biased genes [6]: hypophysectomy leads to down regulation of class I sex-biased genes in the sex
where the gene is more highly expressed, and it leads to up regulation of class II sex-biased genes in the sex
where the gene shows lower expression (Fig. S4, model). Building on this classification, we hypothesized that
DHS that close following hypophysectomy are enhancers mapping to sex-specific genes subject to either positive
regulation by GH (class I sex-biased genes) or negative regulation by GH (class II sex-biased genes). Supporting
this proposal, the DHS that close in male liver following hypophysectomy showed strong, specific enrichment
(ES=5.7, p=1.3E-69) for mapping to class I male-specific genes, which are down regulated in hypox male liver;
whereas the DHS that close in hypox female liver showed strong, specific enrichment (ES=7.8, p=6.6E-46) for
mapping to class I female-specific genes, which are down regulated in hypox female liver (Fig. 4D). In contrast,
the DHS that open in hypox male liver were specifically enriched for mapping to class II female-specific genes
(ES=3.6, p=1.8E-15), which are up regulated (de-repressed) in male liver following hypophysectomy; whereas
the DHS that open in hypox female liver were specifically enriched albeit only moderately (ES=1.9, p=4.8E-04)
for mapping to class II male-specific genes, which are up regulated in female liver following hypophysectomy
(Fig. S4). The enrichment patterns exhibited by these four sets of hypophysectomy-responsive DHS mirror the

corresponding sex-specific gene response patterns (Fig. 4D, *bottom*). Thus, all four DHS sets respond to hypophysectomy in a manner consistent with positively-acting regulatory elements linked to sex-biased genes. DHS that are linked to class I sex-biased genes, and whose chromatin closes following hypophysectomy, require pituitary hormone to maintain open chromatin; when pituitary hormones are ablated by hypophysectomy, chromatin closes and their class I sex-biased target genes are repressed. In contrast, DHS that are linked to class II sex-biased genes of the opposite sex bias, and whose chromatin opens following hypophysectomy, are kept in a closed chromatin state by pituitary hormone; when pituitary hormones are ablated, chromatin opens locally at those DHS and their class II sex-biased target genes are de-repressed: expression of female-biased class II genes increases in hypox male liver and expression of male-biased class II genes increases in hypox female liver.

A single exogenous GH pulse rapidly reopens chromatin at dynamic DHS in hypox male liver – Next, we investigated whether the loss of pituitary GH accounts for the widespread chromatin closing seen in hypox male liver. Fig. 4B shows that exogenous GH treatment induces chromatin opening within 30 min at more than 3,000 DHS, including 71% of all dynamic DHS that closed in male liver following hypophysectomy (Fig. 4E). The fraction of reopening chromatin regions increased to 83% when considering the set of dynamic male-biased DHS that closed following hypophysectomy (Table S2A, column AK). Chromatin reopening was sustained for at least 90 min and then decreased substantially back toward baseline by 240 min (Fig. 4B; Fig. 5A, plot 1), at which time liver STAT5 signaling has terminated [16]. Importantly, the mean level of chromatin re-opening induced by an exogenous GH pulse at dynamic male-biased DHS was very similar to that of the same DHS set in STAT5-high male liver (Fig. 5A, plot 1 vs. Fig. 3A, plot 5; peak DNase-I activity value 1886 vs. 1848). Thus, the rapid chromatin opening induced by a single exogenous pulse of GH recapitulates the effects of an endogenous GH pulse in opening dynamic male-biased DHS.

Hypophysectomy led to closure of many fewer static male-biased DHS than dynamic male-biased DHS (Fig. 4C). Exogenous GH treatment partially reversed chromatin closing at static male-biased DHS within 30 min, with the effect persisting, even after 240 min (Fig. 5A, plot 2). This persistence is consistent with static male-biased DHS remaining constitutively open in male liver when liver STAT5 activity dissipates between endogenous plasma GH pulses. Static female-biased DHS also showed a decrease in mean chromatin opening following hypophysectomy, with further decreases in accessibility seen following GH pulse treatment (Fig. 5A, plot 3). Finally, in an important control, the mean chromatin accessibility at the set of 64,584 static sex-independent DHS was unchanged following hypophysectomy or GH pulse stimulation (Fig. 5A, plot 4).

Stratification of the full set of 2,729 male-biased DHS by the presence or absence of STAT5 binding, as determined by ChIP-seq, highlighted the STAT5 dependence of the rapid increases in chromatin accessibility stimulated by GH treatment (Fig. 5B, plot 1 vs plot 2). GH stimulated chromatin reopening at the static male-biased DHS was also primarily associated with the STAT5-bound DHS subset (Fig. S5, plot 2A vs 2B). In contrast, GH induced a much smaller increase in chromatin opening at the STAT5-bound subset of sex-independent DHS (Fig. 5B, plot 3 vs plot 1), where chromatin is already open and largely independent of plasma GH pulses (Fig. 3A, plot 3). GH decreased chromatin accessibility within 30 min at the 123 genomic sites with lower chromatin accessibility in STAT5-high liver as compared to STAT5-low liver (c.f., Fig. 2E), followed by a return to baseline by 240 min (Fig. 5C); this response pattern is consistent with the repression of chromatin accessibility at these sites by endogenous GH/STAT5 pulses (Fig. 2E). Finally, sex-independent DHS that do not bind STAT5 were unresponsive to both hypophysectomy and GH pulse treatment (Fig. 5B, plot 4).

A subset comprised of 354 dynamic, STAT5-high responsive DHS that close following hypophysectomy did not reopen even 240 min after GH pulse treatment (Fig. 4E, Fig. S6). These 354 DHS showed significantly lower differential chromatin accessibility between STAT5-high and STAT5-low livers than did the dynamic STAT5-high DHS subset that responded to an exogenous GH pulse (Fig. S6C, set 4 vs set 5). These 354 dynamic DHS may require multiple GH pulses to reopen or may be co-dependent on other pituitary-regulated hormones ablated by hypophysectomy.

Enrichment of sex-biased histone marks at static and dynamic male-biased DHS – Our initial analyses revealed no major differences between dynamic and static male-biased DHS regarding enhancer vs insulator vs promoter classifications (Fig. 1A) or overall chromatin state distributions (Fig. 1D, Fig. S7). We therefore

examined the two classes of male-biased DHS for differences in sex-biased histone marks (**Table 1C**). Male-
biased enhancer marks (H3K27ac and H3K4me1) showed strong enrichment at both dynamic and static male-
biased DHS when compared to a background set of 64,584 static, sex-independent DHS (Fig. 6A, Table S7B). In
contrast, male-biased H3K36me3 marks, which are characteristic of transcribed regions, were enriched at static
but not at dynamic male-biased DHS. In addition, static but not at dynamic male-biased DHS were significantly
depleted of female-biased enhancer marks (H3K27ac and H3K4me1) (Fig. 6B). This finding is consistent with
the above finding that static male-biased sites are in a comparatively closed state in female liver (Fig. 3, plot 2;
model in Fig. 6C). Static female-biased DHS showed strong enrichments for female-biased enhancer marks
(H3K27ac and H3K4me1), for female-biased H3K4me3 promoter marks, and for female-biased H3K36me3
transcribed region marks (Fig. 6B), a pattern that was very similar to the enrichments seen at static male-biased
DHS (Fig. 6A).

Examination of two repressive histone marks, H3K27me3 and H3K9me3, revealed their use in a unique way in
each sex to enforce sex differences in chromatin states at sex-biased DHS (Fig. 6C). Male-biased H3K27me3
marks were specifically associated with static female-biased DHS (Fig. 6A), consistent with our prior work
showing deposition of these marks by the Ezh1/Ezh2 enzymatic component of PRC2 as a specific mechanism to
repress many female-biased genes in male liver [35]. In contrast, female-biased H3K9me3 repressive marks were
significantly enriched at both dynamic and static male-biased DHS (Fig. 6B); however, we did not find any
corresponding enrichment of male-biased H3K9me3 marks at female-biased DHS. This novel finding suggests
that H3K9me3 marks are specifically used to repress male-biased genes in female liver.

Chromatin state analysis at sex-biased DHS – To further investigate chromatin state differences between
dynamic and static DHS, we computed enrichment scores and depletion scores at each DHS set for each of 14
chromatin states. These 14 states were defined by a panel of activating and repressive chromatin marks and by
the presence of open chromatin (Fig. 1C) and were determined separately for male and female mouse liver [18].
Enrichments were calculated using a genome-wide background set comprised of all 64,584 static sex-
independent DHS (Fig. 7). In male liver, enhancer state E6 (characterized by the presence of DHS, H3K27ac and
H3K4me1; Fig. 1C) was enriched at both dynamic and static male-biased DHS, and at dynamic sex-independent
DHS. Promoter states E7 and E8 were significantly depleted, consistent with the relative paucity of promoter
states in the full set of sex-biased DHS (Fig. 1A). We also observed strong depletion of the inactive chromatin
states E1 (H3K27me3 marks) and E2 (marks not known) and of the transcribed state E14 (H3K36me3 marks),
specifically from the dynamic but not static male-biased DHS in male liver (Fig. 7A). Finally, dynamic and static
male-biased DHS were both strongly enriched for the inactive state E2 in female liver (Fig. 7B).

Dynamic male-biased DHS were enriched for the enhancer state E9 in female liver (ES=2.42, p=1.6E-69; Fig.
7B) but not male liver (Fig. 7A). The emission parameters of state E9 (Fig. 1C) are very similar to those of state
E6, namely, it shows a high frequency of the activating chromatin marks H3K27ac and H3K4me1 but lacks the
open chromatin (DHS) feature characteristic of state E6. Thus, in female liver, dynamic male-biased DHS
contain histone marks that typify an active enhancer but are inactive due to the comparatively closed state of
their chromatin. Indeed, the extent of chromatin opening at these DHS in female liver is equivalent to the level of
chromatin opening seen at the same set of sites in male liver between plasma GH/liver STAT5 pulses (Fig. 3A,
plot 5; model in Fig. 6C). Dynamic male-biased DHS were also enriched for chromatin state E10 in female liver
(ES=2.89, p=1.9E-18) but not male liver. State E10 shows a high frequency of H3K27ac marks, but not
H3K4me1 marks, and lacks DHS. This pattern of chromatin state E9 and E10 enrichment at dynamic male-
biased DHS in female liver, combined with the strong enrichment of H3K9me3 repressive marks (Fig. 6B),
indicates that the genomic regions encompassing dynamic male-biased DHS are in a bivalent chromatin state in
female liver (Fig. 6C). Conceivably, such a bivalent state may facilitate chromatin opening under certain
biological conditions, e.g., chemical exposure or biological stress.

Static female-biased DHS showed little or no enrichment of specific chromatin states in female liver when
compared to the genome-wide background set of static sex-independent DHS (Fig. 7B). This supports a model
whereby their female bias in accessibility is largely due to active suppression in male liver (Fig. 6C). Indeed, the
genomic regions encompassing static female-biased DHS were strongly enriched in male liver for inactive
chromatin states E1 (characterized by repressive mark H3K27me3), E2 and E4 (Fig. 7A). Static female-biased

DHS showed weak enrichment for enhancer states E10 and E11 in male liver, which are respectively characterized by the active enhancer marks H3K27ac and H3K4me1 but devoid of DHS. This is analogous to the finding, above, that male-biased DHS are enriched for chromatin states replete with active histone marks but deficient in DHS in female liver. Static female-biased DHS also showed modest enrichment in male liver for bivalent state E12, which is characterized by a mixture of activating histone marks and the presence of repressive H3K27me3 marks, consistent with the strong enrichment of the latter marks seen in Fig. 6A, above. Finally, promoter-like states E7 and E8 were significantly depleted from static female-biased DHS (Fig. 7), consistent with the female-biased DHS largely being gene distal sex-biased enhancers.

Distinct gene targets and enriched biological processes of dynamic and static sex-biased DHS – Dynamic and static male-biased DHS both showed strong enrichment for mapping to male-biased gene targets (8-fold and 11.7-fold enrichments, respectively), as did static female-biased DHS for female-biased gene targets (12.5-fold enrichment) when DHS were mapped to the single nearest transcription start site in the same TAD (Table S2B). Further, mapping of DHS to putative target genes using GREAT, which typically maps each DHS to two genes, revealed how many male-biased genes may be regulated by multiple male-biased DHS. In some cases, all of the associated male-biased DHS were static male-biased DHS (e.g., *Cyp4a12a*, with 7 static male-biased DHS), while in other cases they were a mixture of dynamic and static male-biased DHS (e.g., *Cyp7b1*, with 7 dynamic and 8 static male-biased DHS) (Table S9A). Top female-biased genes enriched for nearby static female-biased DHS included *Cux2* and three *Cyp3a* genes, with 10-15 female-biased DHS each (Table S10C). Sex-independent genes that are well-established direct targets of STAT5 [36] include *Igf1*, a target of 12 dynamic sex-independent DHS and 3 dynamic male-biased DHS, and *Socs2*, a target of 8 dynamic sex-independent DHS (Table S10D, Table S10A). Other analyses revealed that all three sets of sex-biased DHS were significantly enriched for mapping to hypophysectomy class I-responsive sex-biased genes as compared to hypophysectomy class II-responsive sex-biased genes (Table S2B). This finding is consistent with the sex-biased DHS sets being enriched for positively-acting enhancers that close following hypophysectomy, which leads to decreased expression of their class-I (i.e., hypophysectomy repressed) gene targets.

The gene targets of each sex-biased DHS set were enriched for distinct but partially overlapping Gene Ontology (GO) Biological Processes, as determined by GREAT analysis. Top enriched GO terms common to both dynamic and static male-biased DHS included lipid metabolic process and steroid metabolic process, consistent with the major role that GH plays in the male-prevalence of fatty liver development and liver metabolic disease [3-5]. Unique enriched terms for dynamic male-biased DHS included cell adhesion, gland development and hepaticobiliary development, whereas gene targets of static male-biased DHS were uniquely enriched for cellular response to glucocorticoid stimulus and EGF receptor signaling, among others (Fig. S8, Table S10, bottom). Static female-biased DHS gene targets were uniquely enriched for long chain fatty acid metabolic pathway and related terms. Finally, the set of dynamic sex-independent DHS mapped to gene targets enriched for terms related to glucose transport and metabolism, IGF receptor signaling and fibroblast proliferation (Fig. S8). This finding is consistent with the widespread metabolic effects that GH has in both sexes, including complex effects on glucose uptake and glucose oxidation, and on hepatic gluconeogenesis and glycogenolysis [37].

Discussion

Sex differences in chromatin accessibility are a central epigenetic feature that enables regulatory proteins, such as the GH-activated transcription factor STAT5, to bind chromatin and regulate hepatocyte gene transcription in a sex-specific manner. However, the underlying mechanisms controlling sex differences in liver chromatin accessibility, which occur at more than 4,000 distinct sites across the mouse genome, are poorly understood. GH, working through its sex-dependent temporal patterns of secretion by the pituitary gland, is the major hormonal factor controlling STAT5-dependent sex differences in liver gene transcription, but little is known about the potential of GH to directly regulate sex differences in the epigenetic landscape required for sex-dependent transcriptional outputs. To address this question, we elucidated the impact of male plasma GH pulses on global patterns of liver chromatin accessibility by analyzing a population of 21 individual male mouse livers euthanized at either a peak or a trough of plasma GH pulse-stimulated hepatic STAT5 DNA-binding activity. Our findings establish that the naturally occurring, endogenous plasma GH pulses characteristic of males induce dynamic cycles of chromatin opening and closing at several thousand DNase-I hypersensitive sites (DHS) in male mouse

liver and that these events comprise one of two major mechanisms regulating the male bias in liver chromatin accessibility. Analysis of sex-dependent transcription factor binding patterns and chromatin states elucidated key features distinguishing this dynamic mechanism of male-biased enhancer activation from that of static, GH pulse-unresponsive male-biased DHS and from female-biased DHS (see model, Fig. 6C). GH thus acts at three distinct steps to regulate sex-dependent hepatocyte gene expression, all three involving the GH-activated transcription factor STAT5 (Fig. 8): GH pulse-induced chromatin opening at dynamic male-biased DHS driven by pulsatile GH activation of STAT5, as discussed further, below; direct transcriptional activation of sex-specific genes by GH-activated STAT5; and GH/STAT5-dependent transcriptional regulation of downstream repressors, such as the female-specific CUX2, which binds to male-biased enhancers in female liver to reinforce sex differences in gene transcription.

GH pulse activation of dynamic male-biased DHS - GH-induced chromatin opening in male mouse liver is shown to be directly controlled by the endogenous male rhythm of plasma GH stimulation of hepatocytes, the major liver cell type contributing to sex-biased liver gene expression [38]. The GH pulse-regulated chromatin regions identified here responded to changes in plasma GH levels in a rapid, dynamic manner, with extensive GH-induced chromatin opening occurring within 30 min, as seen when hypophysectomized male mice were given a physiological replacement dose of GH pulse by intraperitoneal injection (Fig. 4A). This time course is consistent with the rapid activation of GH receptor signaling to STAT5, which occurs within 5 min in cell culture [39] and within 15 min *in vivo* in a hypophysectomized rat model [40]. Importantly, 85% of the dynamic male-biased DHS identified here (710 of 834 dynamic male-biased DHS) were bound by STAT5, which appears to be a key driver of these GH pulse-induced chromatin opening events. Chromatin closing followed the termination of nuclear STAT5 signaling, which is complete within 4 h [16] and occurs in sufficient time to reset the GH receptor-JAK2 signaling complex and resensitize hepatocytes before the next plasma GH pulse [39] (Fig. 2A). STAT5 binding and chromatin opening at dynamic male-biased DHS were both significantly lower in livers of female mice, where plasma GH levels and liver STAT5 activity are persistent (near-continuous) yet ineffective at maintaining an open chromatin state. Indeed, the extent of chromatin opening at these male-biased DHS in female liver is very similar to the level in livers of male mice euthanized between plasma GH pulses (STAT5-low male livers; Fig. 3A, panel 5). Pulsatile chromatin opening stimulated by endogenous plasma GH pulses is thus a unique mechanism for establishing and maintaining male-biased chromatin accessibility and transcription factor binding at 834 male-biased DHS, corresponding to 31% of all male-biased DHS in the liver.

GH pulse-responsive DHS were discovered by comparing global chromatin accessibility patterns in a set of 8 livers collected from mice euthanized at a peak of GH-activated STAT5 DNA-binding activity (STAT5-high livers) to those of 10 other livers from mice euthanized between pulses of STAT5 DNA-binding activity, i.e., when liver STAT5 activity is very low or undetectable (STAT5-low livers). Three other male livers gave DNase-seq profiles inconsistent with their EMSA-determined STAT5 DNA-binding activity. These outlier livers may have come from mice euthanized just after the onset of a STAT5 pulse [27], when more time is needed to induce chromatin opening (2 outliers), or shortly after STAT5 is deactivated by tyrosine dephosphorylation [41] but prior to the reversal of chromatin opening (1 outlier). Importantly, the top 200 genomic regions showing differential chromatin opening between STAT5-high and STAT5-low livers also separated a group of nine C57Bl/6 male mouse liver DNase-seq samples from the ENCODE consortium into two distinct classes, corresponding to STAT5-high activity livers (n=4) and STAT5-low activity livers (n=5), respectively. Thus, the dynamic pattern of chromatin opening and closing at these genomic regions is robust across studies and mouse strains.

STAT5-high male livers showed the highest levels of chromatin accessibility among the male-biased DHS that bind STAT5. Similarly, a higher level of chromatin opening was seen at sex-independent DHS that bind STAT5 compared to those that do not bind STAT5 (Fig. 3A, panel 3 vs 4). There is thus a close linkage between STAT5 binding and chromatin opening. This finding is consistent with the proposal that STAT5 acts as a pioneer factor to enable chromatin opening, as was reported for STAT5 action at the *Il9* gene locus in Th9 T-cells [42]. It should be noted, however, that pulsatile chromatin opening also occurred at the small subset (15%) of dynamic male-biased DHS that did not bind STAT5, suggesting a role for other GH receptor signaling pathways [43] in male-biased chromatin opening at those sites. Furthermore, pulsatile STAT5 activation and pulsatile STAT5 DNA binding alone are not sufficient to insure pulsatile chromatin opening, insofar as many sex-independent

DHS do bind STAT5, yet are constitutively open in both male and female liver (Fig. S2C, plot 5A). Finally, plasma GH pulse stimulation decreased chromatin accessibility at a small number of liver DHS, most of which were sex-independent. The mechanism for this decrease and its physiological significance are unknown.

Regulation of static male-biased DHS - We identified a second, but less well-defined mechanism that controls the male bias in chromatin accessibility at a distinct DHS set, comprised of 1,895 static male-biased DHS. These DHS represent 69% of all male-biased DHS and mapped to a set of target genes and enriched biological processes distinct from but overlapping those of the dynamic male-biased DHS. Static male-biased DHS are constitutively open in male liver, with chromatin accessibility largely unchanged across the peaks and valleys of GH-induced liver STAT5 activity. Moreover, in contrast to dynamic male-biased DHS, the male bias in chromatin accessibility at static male-biased DHS is primarily a reflection of their being in a closed state in female liver (Fig. 6C). Static male-biased DHS are relatively deficient in STAT5 binding but showed significant enrichment for binding CUX2, a female-specific repressor protein [30], which we infer contributes to the closure of static male-biased DHS in female liver through its established transcriptional repressor activity [44]. Female-biased DHS showed even greater depletion of STAT5 binding (comparable to genome-wide background of sex-independent DHS), and correspondingly, the vast majority (99%) are static i.e., unresponsive to plasma pulse stimulation in male liver.

Our findings support the following proposal to help explain the widespread closure of male-biased liver DHS, most notably dynamic male-biased DHS (95% closing; Table S2C) when male mice are given GH as a continuous infusion, which mimics the female plasma GH pattern and substantially feminizes liver gene expression [17, 45] (Fig. 6C). First, dynamic male-biased DHS close due to loss of the pulsatile plasma GH pattern and its intrinsic ability to stimulate chromatin opening. Second, static male-biased DHS close due to the strong induction of CUX2 [17], whose binding is enriched at those sites and is associated with chromatin closing, as noted above. The pioneer factors FOXA1 and FOXA2 showed strongest enrichment of their male-biased binding sites at static male-biased DHS and may help maintain these DHS in their constitutively open, male-biased chromatin state. Correspondingly, static female-biased DHS showed strongest enrichment for female-biased FOXA2 binding sites, which likely contributes to their greater accessibility in female compared to male liver (Fig. 6C).

Novel insights from chromatin state analysis - Dynamic and static male-biased DHS are both largely (~95%) comprised of promoter-distal enhancer DHS, as indicated by their flanking histone modifications. This supports our prior finding that sex-dependent intra-TAD DNA looping mechanisms are common in mouse liver [24] and indicates such looping mechanisms likely bring both subsets of male-biased DHS into closer proximity with their sex-biased promoter targets. Dynamic male-biased DHS were enriched for the active enhancer state E6 in male liver but were enriched for enhancer states E9 and E10 in female liver. States E9 and E10 are characterized by a high frequency of the activating chromatin marks H3K27ac and H3K4me1 (E9) or H3K27ac alone (E10), similar to state E6, but unlike E6, they are both deficient in open chromatin (DHS) (Fig. 1C). Thus, in female liver, dynamic male-biased DHS regions contain active histone marks but are in a comparatively closed chromatin state. This may in part be due to their enrichment for H3K9me3 (repressive) histone marks in female liver, which gives them bivalent character (Fig. 6C). This bivalent chromatin state may give dynamic male-biased DHS the potential for chromatin opening and activation of their latent enhancer activity in female liver, e.g., in response to chemical exposure or biological stress [46]. In contrast, static male-biased DHS are depleted of female-biased enhancer marks (H3K27ac and H3K4me1) and are in an inactive chromatin state in female liver (Fig. 6C). Of note, the bivalent chromatin state of dynamic male-biased DHS in female liver is distinct from the poised state these genomic regions apparently adopt in male liver between plasma GH pulses (i.e., in STAT5-low male liver) and in hypophysectomized male liver, where a single GH pulse is all that is required to induce rapid chromatin opening, even after several weeks of pituitary hormone ablation.

Static female-biased DHS are deficient in DHS and are strongly enriched for the repressive histone mark H3K27me3 in male liver, which is thought to contribute directly to their closed, inactive chromatin state in male liver (Fig. 6C). Prior studies have revealed heterogeneity in these female-biased DHS regarding the time-dependent loss of H3K27me3 marks and chromatin opening in livers of male mice exposed to GH as a continuous infusion [17], and in the extent of female-biased target gene de-repression following the loss of

H3K27me3 marks in livers of male mice deficient in the H3K27-trimethylase complex PRC2 [35]. Novel insight into the fundamental underlying epigenetic mechanisms of sex-biased gene regulation comes from our striking finding that H3K27me3 and H3K9me3, two widely used repressive histone marks, are used in a unique way in each sex to enforce sex differences in chromatin states at sex-biased DHS. Thus, whereas male-biased H3K27me3 repressive marks were highly enriched at, and specifically associated with, static female-biased DHS, consistent with our prior work [18, 35], we show here that female-biased H3K9me3 repressive marks are highly enriched at both dynamic and static male-biased DHS, but without a corresponding enrichment of male-biased H3K9me3 marks at static female-biased DHS. This novel discovery supports the proposal that sex-specific gene repression is mediated by two distinct mechanisms, with H3K9me3 marks specifically used to repress male-biased genes in female liver, and H3K27me3 marks specifically used to repress female-biased genes in male liver. Further study will be required to elucidate the underlying mechanisms whereby these repressive histone marks are respectively deposited in a sex-dependent and site-specific manner and the role of GH in regulating these molecular events.

Regulatory elements associated with class I and class II sex-biased genes - DNase-seq profiling identified more than 5,000 liver DHS that open or close following hypophysectomy. Strikingly, a single physiological replacement dose of GH restored, within 30 min, liver STAT5 activity and chromatin accessibility at 83% of dynamic male-biased DHS that closed following hypophysectomy. Remarkably, these chromatin sites are maintained in a primed (poised) state in hypox male liver over a period of several weeks, despite the prolonged deficiency of GH and other pituitary hormones and the dysregulation of several thousand liver-expressed genes [6, 16]. DHS that close in male liver following hypophysectomy were enriched for proximity to class I male-specific genes, which are down regulated in hypox male liver, whereas DHS that close in hypox female liver showed the strongest enrichment for class I female-specific gene targets, which are down regulated in hypox female liver (model, Fig. S4). Further, DHS that open in hypox male liver were enriched for proximity to class II female-specific genes and DHS that open in female liver were enriched for proximity to class II male-specific genes. These enrichments are consistent with the definition of class II sex-biased genes, i.e., genes that are up regulated by hypophysectomy in the sex where they show lower expression in intact mice. Together, these findings lend strong support for the functional role of each class of sex-biased and pituitary hormone-dependent DHS as regulatory elements with intrinsic, positively acting enhancer potential, keeping gene expression on in intact liver by maintaining open chromatin, e.g., in the case of class I male-biased DHS and class I male-biased genes in male liver, or keeping gene expression off by maintaining a closed chromatin state, e.g., in the case of class II male-biased DHS and class II female-biased genes in male liver. Further study will be required to elucidate the underlying regulatory factors and molecular mechanisms governing these gene regulatory circuits.

Methods

Animal treatments and DNase-seq analysis - All mouse work was carried out in accordance with ARRIVE Essential guidelines 2.0 [47] for study design, sample size, randomization, experimental animals and procedures, and statistical methods, and with approval of the Boston University Institutional Animal Care and Use Committee. Male and female CD1 mice (ICR strain, strain code # 022), 7-weeks old and purchased from Charles River Laboratories (Wilmington, MA), were kept on a 12-hour light/dark cycle with food and water without restriction. Livers were collected from individual untreated mice between 8 and 10 weeks of age. Where indicated, male and female mice were hypophysectomized (hypox) by the supplier at ~7-8 weeks of age. Completeness of hypophysectomy was verified by the absence of weight gain over a 2-3 week period and by the lack of Mup protein in urine samples (SDS-PAGE analysis) [16]. Hypox male mice were treated with recombinant rat GH (purchased from Prof. Arieh Gertler, Protein Laboratories Rehovot Ltd, Rehovot, Israel), given as a single intraperitoneal injection at 125 ng of GH per gram body weight [6], or vehicle (control), and were euthanized by cervical dislocation 30, 90 or 240 min later [16]. A protein extract prepared from a small piece of each liver to be used for DNase-seq analysis (see below) was assayed for liver STAT5 DNA-binding activity by EMSA, as described [16]. Results of this assay allowed us to classify each individual male mouse as STAT5-high activity (n=10) or STAT5-low activity (n=11) at the time of euthanasia and liver collection. EMSA was also used to verify the ablation of liver STAT5 activity following hypophysectomy and the effectiveness of exogenous GH administration at restoring endogenous STAT5-high liver levels of STAT5 EMSA activity within 30 min.

DNase-seq libraries, Illumina sequencing, and data processing - Livers used for these analyses were obtained from intact male and female mice, from hypox male and hypox female mice, and from hypox male mice given a single injection of GH and euthanized 30, 90 or 240 min later (n=6-12 livers per group). Nuclei were purified from individual fresh livers as described [19] and stored at -80 ºC. Nuclei were subsequently treated with DNase I to release DNA fragments from hypersensitive genomic regions to identify liver DNase hypersensitive sites (DHS) [16]. The DNA fragments released from each liver were combined (3 livers per pool) to give n=2 to n=4 independent pooled samples, which were used to prepare DNase-seq libraries for each mouse group (Table S1). In the case of STAT5-high and STAT5-low liver DHS analysis, however, DNase-seq libraries were prepared directly from DNase-I digested fragments released from each individual liver without pooling, as described below. DNase-seq libraries were prepared using the NEBNext Ultra Library Prep Kit for Illumina (New England Biolabs). Illumina sequencing was performed on an Illumina HiSeq instrument to a depth of 12 to 33 million mapped 40-50 bp single-end sequence reads per sample. Detailed sequencing statistics are shown in Table S1. Raw and processed data are available at www.ncbi.nlm.nih.gov/gds under accession numbers GSE131848 and GSE131852 (SuperSeries GSE131853).

Sequencing data was analyzed using a custom in-house DNase-seq pipeline [46]. Briefly, the pipeline processes raw FASTQ files and outputs various control metrics, including FASTQC reports (FastQC v0.11.7), confirmation of read length, verification of the absence of read strand bias, and identification of contaminating adaptor sequences (Trim_galore v0.4.2). Reads were mapped to mouse genome mm9 using Bowtie2 (v2.2.6) [48]. Regions of DNase hypersensitivity (DHS) were discovered as peaks identified by MACS2 (v2.1.0.20150731) [49] using the option (-nomodel –shift -100 –extsize 200) to inhibit read shifting, and the option (-keep-dup) to retain all reads that contribute to the peak signal. Peaks were discovered for each individual DNase-seq library then filtered to remove peaks that overlap ENCODE blacklisted regions [50] as well as peaks comprised of 5 or more identical reads that do not overlap any other read ("straight peaks"). Nine additional individual male mouse liver DNase-seq datasets generated by the ENCODE consortium (FASTQ files downloaded at http://genome.ucsc.edu/cgi-bin/hgFileUi?db=mm9&g=wgEncodeUwDnase; GEO accession: GSM1014195, liver replicates #5 to #13) were analyzed using the same DNase-seq pipeline (cell line, liver; strain, C57BL/6; age, adult 8 weeks; sex, male). These DNase-seq datasets were single end sequencing reads, 36 bp in length, with a sequencing depth of 19 to 37 million total mapped reads per sample.

Liver DHS classification and DHS gene target expression data - A set of 72,862 mouse liver DHS regions previously identified by DNase-seq analysis of male and female mouse liver [19] was filtered to remove 515 DHS regions that overlapped ENCODE blacklisted regions. 2,136 DHS regions that could not be classified according to a 5-class DHS model defined previously [20] were also removed to obtain a standard reference set of 70,211 mouse liver DHS. Each DHS was designated as a promoter, weak promoter, enhancer, weak enhancer, or insulator DHS based on the 5-class model, which is primarily based on ChIP-seq signals for the H3 histone marks H3K4me1and H3K4me3, combined with CTCF ChIP-seq binding in adult male mouse liver [20]. Weak enhancers were identified by their low levels of H3K27ac ChIP-seq signal compared to DHS classified as enhancers, combined with a distance from RefSeq gene transcription start sites inconsistent with a weak promoter designation [20]. Further, individual DHS were designated male-biased (n=2,729), female-biased (n=1,366), or sex-independent (n= 66,116) based on significant differences in chromatin accessibility between male and female mouse liver [19]. Except where noted, each DHS was assigned a single putative gene target (RefSeq gene or multi-exonic lncRNA gene) corresponding to the closest transcription start site within the same TAD [20]. Table S2A lists DHS target genes, their overlap with STAT5 binding sites determined by ChIP-seq [15], their DNase-seq activity (chromatin accessibility) in male and female mouse liver, and their responses to hypophysectomy and to hypophysectomy + GH treatment. Where indicated, DHS target genes were defined using GREAT (v.4.0.4) [51, 52] (http://great.stanford.edu/public/html/index.php) and the default settings for the Basal + extension gene association rule, which typically assigns two genes to each DHS (see output in Table S9 and Table S10, below).

Differential DHS between STAT5-high and STAT5-low activity livers – Nuclei purified from each individual STAT5-high activity (n=10) and STAT5 low-activity mouse liver (n=11) (Table S1), classified based on EMSA as described above, were analyzed by DNase-seq to discover genomic regions (i.e., DHS) that responded

dynamically to endogenous plasma GH pulses and the associated changes in liver STAT5 DNA-binding activity, as follows. diffReps analysis [28] was used to discover genomic sites that were more open or were more closed (DNase-seq normalized intensity |fold-change| > 2 and FDR < 0.05 [Benjamini-Hochberg adjusted p-value]) for the set of EMSA-identified STAT5-high activity livers as compared to the set of EMSA-identified STAT5-low activity livers. The diffReps nucleosome option (200 bp window size) was used and the option (-frag) was set to zero for all comparisons. The differential sites that diffReps identified as significant were further analyzed by two methods, principal component analysis and boxplot analysis, to determine the distributions of normalized sequence read counts (reads per kilobase per million mapped reads) for each individual DNase-seq library. Three outlier DNase-seq samples were thus identified: two from livers designated STAT5-high activity based on EMSA (samples G74A_M1 and G74A_M2), which gave DNase-seq read count distributions (top 200 and top 600 diffReps-identified differential sites) more similar to STAT5-low activity livers; and one from a STAT5-low EMSA activity liver (sample G92_M5), which gave a DNase-seq read count distribution more similar to STAT5-high activity livers (Fig. 2C and Fig. 2D, below). These three outlier liver DNase-seq libraries were excluded from all downstream analysis. Next, we implemented diffReps analysis using the same cutoffs described above to compare the DNase-seq activity profiles of the remaining EMSA-identified STAT5-high activity (n=8) and STAT5-low activity (n=10) DNase-seq samples. The resultant set of diffReps differential sites was overlapped with a MACS2 DHS peak union list, which was obtained by merging the MACS2 DHS peak calls from all 18 male mouse liver DNase-seq samples using the BEDTools "merge" command. This overlap analysis yielded a final set of 70,767 MACS2 DHS peak union sites (see Table S4A), of which 2,832 MACS2-identified DHS that were more open (more accessible chromatin state) and 123 DHS were in a more closed chromatin state based on their overlap with the diffReps-identified STAT5-high vs. STAT5-low livers differential regions (see Fig. 2E, below). The other 67,812 DHS regions were designated static DHS, as they did not overlap a diffReps-identified STAT5-high vs. STAT5-low differential region.

Comparison of these 70,767 MACS2-identified DHS peak union sites with our standard reference set of 70,211 mouse liver DHS identified n=2,373 reference set DHS that overlapped a STAT5-high differential site (see Fig. 2F, below), n=98 reference set DHS that overlapped a STAT5-low differential site, and n=45,754 liver DHS that overlapped a static DHS site (Table S2A, columns H and I). The remaining 21,986 standard reference set DHS did not overlap the 70,767 DHS peak union list and were also labeled static DHS. We applied the designation of sex bias determined previously [19], namely, male-biased, female-biased, or sex-independent, to each DHS that overlapped the reference set 70,211 liver DHS (Table S2A, column F). The standard reference set liver DHS were further designated dynamic DHS if they overlapped a STAT5-high > STAT5-low differential DHS (i.e., a GH/STAT5 pulse-opened DHS). Liver DHS not identified as dynamic (including the 98 STAT5-low > STAT5-high DHS peak union sites that overlap a standard reference set DHS) were designated static with respect to chromatin accessibility changes induced by endogenous GH-induced STAT5 pulses. Sex-specific DHS were thus designated dynamic male-biased DHS, static male-biased DHS, dynamic female-biased DHS, or static female-biased DHS with respect to GH/STAT5-induced chromatin opening (Table S2A, column I). Sex-independent DHS were similarly designated dynamic or static sex-independent DHS (Fig. 2F, below).

DNase-I cut site aggregate plots - Normalized DNase-I cut site aggregate plots were generated using an input DNase-seq dataset and the set of input genomic regions (DHS sequences) to be analyzed by sequence read counting. First, FASTQ files from DNase-seq biological replicates were concatenated to obtain a single combined replicates file for each treatment group. For example, for male liver, we generated a single combined STAT5-high DNase-seq FASTQ file by merging the n=8 biological replicate STAT5-high liver FASTQ files, and separately, we generated a single combined STAT5-low sample by merging the n=10 biological replicate STAT5-low DNase-seq FASTQ files (Table S1). Combined replicate FASTQ files were generated for intact female, hypox female, hypox male, and each of the hypox male + GH treatment time point DNase-seq samples in the same manner. Second, for each combined replicate file, a BED file comprised of positive and negative strand reads was processed to determine the DNase-I cut site, which corresponds to the 5'-end of each sequence read. Third, the set of input genomic regions was processed to generate a list of 2 kb regions centered at the midpoint of each DHS. The BEDTools "coverage" command using the (-d) option was used to count the number of DNase-I cut sites at each nt position of the 2 kb midpoint-centered regions, thus producing a read count matrix composed of 2,000 read counts for each input genomic region. Fourth, a custom R script was used to load the read count matrix, calculate the sum of read counts at each nt position, normalize the raw read counts by the

number of reads in the subset of 70,211 standard reference DHS to be analyzed, normalize by the number of
input genomic regions, and then generate a plot of the DNase-I signal across the input genomic regions. The
resultant normalized DNase-I cutting profiles were then smoothed using a LOWESS smoother as implemented in
R (package: gplots v3.0.1.1). An offset was applied to the profile by subtracting an average read count
(calculated from the first 200 nt positions) from the normalized read count intensity to standardize the baseline of
each profile. Profiles for other, related DNase-seq datasets were processed in the same manner and were plotted
on the same set of axes to enable direct comparisons across all such plots in Figs. 2, 3, 5 and Figs. S1, S4 and S5.
754
Impact of hypophysectomy and GH injection on chromatin opening and closing - MACS2 was used to
discover differential DHS peaks in DNase-seq samples prepared from intact male and intact female mouse liver,
hypox male and hypox female mouse liver, and from livers of hypox male mice given a single injection of GH
and euthanized either 30, 90, or 240 min later (Table S1). The effects of hypophysectomy on chromatin
accessibility were determined by comparing hypox liver DNase-seq samples to the corresponding samples
prepared from intact liver (control) samples in male liver and separately in female liver. The effects of a single
injection of GH were determined by comparing DNase-seq samples from livers of hypox male mice treated with
GH to the untreated hypox male liver controls at each time point. For each comparison, a DHS peak union list
was generated by merging the MACS2 DHS peak calls from all individual biological replicates for the
corresponding treatment group. For example, a single peak union list for the hypox male compared to intact male
liver samples was generated by merging all the MACS2 peaks from each of the n=8 individual male mouse liver
DNase-seq samples (n=6 intact male control livers and n=2 hypox male livers; Table S1). Genomic regions that
showed significantly differential DNase-seq signal between hypox and intact control livers, or between hypox
male + GH and hypox male livers were discovered separately for each comparison using diffReps, using the
nucleosome option (200 bp window). Significance was based on |fold-change| > 2 and FDR < 0.05 (Benjamini-
Hochberg adjusted p-value) for diffReps-normalized signal intensity values. The diffReps-identified differential
sites were then filtered by their overlap with the peak union list for all 8 samples to obtain the sets of differential
DHS (e.g., 2,142 DHS that open and 4,856 DHS that close in male liver following hypophysectomy; Table S4,
Summary sheet). MACS2 peak union DHS that did not overlap a diffReps region were annotated as static DHS
peaks (50,055 sites). Concatenation of the diffReps-identified differential DHS with this set of static DHS
yielded the full set of 57,053 DHS peak union sites for this dataset. Each of the 70,211 standard reference set
liver DHS (see above) was then labeled based on whether it overlapped a diffReps-identified differential DHS
that opened following hypophysectomy, a diffReps differential DHS that closed following hypophysectomy, a
static DHS, or "none", for those reference set DHS that did not overlap any of the set of 57,053 hypophysectomy
study DHS peak union sites. Corresponding analyses were performed for the intact and hypox female liver
samples, and for the hypox male + GH time-course comparisons mentioned above. Table S4 summarizes the
numbers of hypophysectomy and hypophysectomy + GH responsive differential DHS for each of these
comparisons, and the subsets that overlap the reference set of 70,211 liver DHS.
783
Sex-specific genes and hypophysectomy response classification - RNA-seq data from intact male and intact
female mouse liver was previously used to identify sex-specific genes based on a gene list comprised of 24,197
RefSeq genes [16]. This list was expanded to include both RefSeq and multi-exonic sex-specific lncRNA genes,
as follows. Sex-specific RefSeq genes were identified from polyA-selected total liver RNA-seq samples from
intact male and intact female liver using the "genebody" method of sequence read counting [16] at a threshold of
|fold-change| > 1.5, adjusted p-value < 0.05 and FPKM > 0.25 for the sex with a greater signal intensity, which
yielded 387 male-specific and 517 female-specific RefSeq genes. Sex-specific multi-exonic lncRNA genes were
identified from polyA-selected liver nuclear RNA-seq samples in male and female mouse liver using the
"ExonCollapsed" read counting method [53] at a threshold of |fold-change| > 2, adjusted p-value < 0.05 and
FPKM > 0.25 for the sex with a higher signal intensity, which yielded 121 male-specific and 102 female-specific
multi-exonic lncRNA genes, for a total of 508 male-specific and 619 female-specific genes (Table S3). A
stringent set of sex-independent genes was defined by FPKM > 1 in both male and female liver, |fold-change| <
1.2, and adjusted p-value > 0.1, which yielded a total of 7,253 stringently sex-independent genes, including
lncRNA genes (Table S3, column N). Sex-biased genes responsive to hypophysectomy were defined by |fold-
change| > 2 and adjusted p-value < 0.05 for hypox mouse liver versus intact mouse liver, determined separately
for male liver and for female liver. Using these thresholds, RNA-seq data from intact and hypox male and female
liver samples were used to group sex-specific genes into classes I and II and their corresponding subclasses (IA,

IB, IC, IIA, and IIB) [6, 16]. Class I sex-biased genes are those sex-biased that are down regulated by
hypophysectomy in the sex where they show higher expression in intact mice. Class II sex-biased genes are those
that are up regulated by hypophysectomy in the sex where they show lower expression in intact mice. Subclasses
A, B, and C indicate the response to hypophysectomy in the dominant sex (class II genes) or in the opposite sex
(class I genes), as defined in Table 3 of [6]. The number of sex-biased genes in each hypophysectomy response
class is shown in Table S3.

STAT5 binding and motif analysis - ChIP-seq analysis of STAT5 binding in male and female mouse liver
identified 15,094 merged peaks comprised of male-enriched and female-enriched, and male-female common
STAT5 binding sites [15]. A small subset of these STAT5 binding sites, 75 peaks, overlapped ENCODE
blacklisted regions and was excluded from downstream analysis. BEDTools overlap analysis of these STAT5
binding sites allowed us to designate each of the 70,211 reference DHS as "STAT5-bound", if a STAT5 binding
site was present, or "not bound". Table S2A provides full details, including the STAT5 binding site, the sex-
specificity of STAT5 binding (male-enriched, female-enriched, or common to both sexes), the normalized ChIP-
seq read counts for STAT5 binding in STAT5-high activity male livers (MH) and in STAT5-high activity female
livers (FH), and the average of these two sets of sequence read counts. The STAT5B motif M00459 from the
TRANSFAC motif database (Release 2011.1) [54] was used to determine the frequency of STAT5 motif
occurrence in each set of DHS sequences. Motifs found in DHS sequences were identified using FIMO (v4.12.0)
[55] using the option (--thres 0.0005) to improve detection of short length motifs. The number of STAT5B motif
occurrences in each of the 70,211 reference set DHS sequences is shown in Table S2A.

Enrichment analysis - For all enrichment calculations described below, the significance of the enrichment was
determined by a Benjamini and Hochberg adjusted Fisher's Exact test p-value as implemented in R. Enrichments
with adjusted p-value < 1E-03 were considered statistically significant. Enrichments of sex-biased DHS for
being enhancer, promoter, or insulator regions (e.g. male-biased DHS for being enhancers compared to the
background set of sex-independent DHS) were calculated, as follows: Enrichment score = (ratio A) / (ratio B),
where: ratio A = the number of sex-biased DHS that are enhancers, divided by the number of sex-biased DHS
that are not enhancers; and ratio B = the number of sex-independent DHS that are enhancers, divided by the
number of sex-independent DHS that are not enhancers. For example, 2,551 male-biased DHS were classified as
enhancers, and 178 other male-biased DHS were not enhancers (2,551/178=14.3), whereas 43,591 sex-
independent DHS were classified as enhancers, and 22,525 sex-independent DHS were not enhancers
(43,591/22,525=1.93), which gives an enrichment score A/B=14.3/1.93=7.41. The set of 66,116 sex-independent
DHS was used as the background for these enrichment calculations.

Enrichments of sex-biased DHS subsets (enhancer, insulator, or promoter) for mapping to sex-specific genes
were calculated (e.g. male-biased enhancer DHS mapping to male-specific genes) as follows (Table S2B):
Enrichment score = (ratio A) / (ratio B), where: ratio A = the number of male-biased enhancer DHS that map to
male-specific genes, divided by the number of male-biased enhancer DHS that map to sex-independent genes;
and ratio B = the number sex-independent enhancer DHS that map to male-specific genes, divided by the number
of sex-independent enhancer DHS that map to sex-independent genes. For example, among the male-biased DHS
classified as enhancers, 404 male-biased enhancer DHS map to male-specific genes, and 525 male-biased
enhancer DHS map to sex-independent genes (404/525 = 0.77), whereas 1,495 sex-independent enhancer DHS
map to male-specific genes, and 15,457 sex-independent enhancer DHS map to sex-independent genes
(1,495/15,457 = 0.097), which gives an enrichment score A/B = 0.77/0.097 = 8.0. The set of 66,116 sex-
independent DHS was used as the background for these enrichment calculations.

Enrichments of hypophysectomy-responsive DHS for mapping to class I or class II sex-specific genes (e.g. DHS
that close in response to hypophysectomy in male liver and that map to male class I genes) were calculated for
male and female liver as follows: Enrichment score = ratio A/ ratio B, where: ratio A = the number of DHS that
open (or that close) following hypophysectomy and that map to class I (or to class II) sex-specific genes, divided
by the number of DHS that open (or that close) following hypophysectomy and that map to sex-independent
genes; and ratio B = the number of hypophysectomy-unresponsive DHS (static DHS) that map to class I or class
II sex-specific genes, divided by the number of static DHS that map to sex-independent genes. For example, in
male liver, 217 DHS that close following hypophysectomy map to a male class I sex-specific gene, and 1,203

other DHS that close map to sex-independent genes (217/1,203=0.1803), whereas 444 static DHS map to a male class I sex-specific gene, and 14,053 static DHS map to a sex-independent gene (444/14,053 = 0.0316), which gives an enrichment score A/B = 0.1803/0.0316 = 5.7. The static DHS used for these enrichment calculations correspond to the set of 35,562 static DHS in male liver and 30,394 static DHS in female liver.

Enrichments for overlap with genomic regions containing biologically relevant sets of transcription factor binding sites, chromatin marks and combinations of epigenetic features (chromatin states) were calculated for each of the following 4 sets of DHS: (1) dynamic male-biased DHS (834 sites), (2) static male-biased DHS (1,895 sites), (3) dynamic sex-independent DHS (1,532 sites), and (4) static female-biased DHS (1,359 sites) relative to a background set of static sex-independent DHS (64,584 sites) (see Fig. 2F). Briefly, these sets of DHS were identified based on their sex-specificity [19] and were classified as "dynamic" if they overlapped a STAT5-high vs. STAT5-low differential DHS, otherwise they were classified as "static". Characterization of the 70,211 reference set DHS members as static or dynamic and their sex-specificity designations are listed in Table S2A. Biologically relevant regions annotated include sex-biased transcription factor binding sites for STAT5 and BCL6 [15], CUX2 [30], and FOXA1/2 [32]. We also examined DHS enrichment for sex-biased chromatin marks and chromatin states defined for male and female liver [18]. FOXA1 and FOXA2 ChIP-seq data [32] was processed and reanalyzed with MAnorm [56] in analyses performed by Gracia Bonilla of this laboratory, which identified sex-biased FOXA1 and sex-biased FOXA2 binding sites in mouse liver. Data from the MAnorm analysis of FOXA1 and FOXA2 ChIP-seq peaks (Table S8) was filtered by log2(fold-change) value (i.e., M-value) to define male-biased and female-biased binding sites. Genomic coordinates for each of the sex-biased transcription factor binding sites, chromatin marks, and chromatin states are provided (Table S5, Table S7). Enrichments of DHS for overlapping biologically relevant regions were calculated (e.g., dynamic male-biased DHS for containing STAT5-High (Male) binding sites compared to the background set of static sex-independent DHS) as follows: Enrichment score = ratio A/ ratio B, where: ratio A = the percentage of dynamic male-biased DHS with a STAT5 binding site; and ratio B = the percentage of static sex-independent DHS with the same type of binding site. For example, 235 dynamic male-biased DHS contain a STAT5-High (Male) binding site (235/834 = 0.2818), whereas 311 static sex-independent DHS contain a binding site (311/64,584 = 0.0048), which gives an enrichment score A/B = 0.2818/0.0048 = 58.7. The set of 64,584 static sex-independent DHS (Fig. 2F) was used as the background for these enrichment calculations.

Chromatin state map analysis - Chromatin state maps (14 state model), developed for male mouse liver, and separately for female mouse liver, are based on epigenetic data from a panel of six histone marks and DHS data in each sex [18]. These maps were used to identify sex differences in chromatin state and chromatin structure at each DHS region and their relationships to sex-biased gene expression, as follows. BEDTools was used to assign one of the 14 chromatin states to each of the 70,211 reference DHS based on the overlap of the DHS with the male liver chromatin state map, and separately, based on its overlap with the female liver chromatin state map (Table S2A). DHS whose genomic coordinates span two or more different chromatin states were assigned to the state with the largest number of overlapping base pairs, or in case of equal numbers of base pairs, the chromatin state with the smaller genomic coordinates. Enrichments of sets of static and dynamic male-biased DHS for being in one of the 14 chromatin states in male liver (e.g. dynamic male-biased DHS for being in chromatin state E6, which is an enhancer-like state) were calculated as follows: Enrichment score = (ratio A) / (ratio B), where: ratio A = the number of male-biased DHS that are in a particular chromatin state, divided by the number of male-biased DHS not in that chromatin state; and ratio B = the number of static sex-independent DHS in that chromatin state, divided by the number of static sex-independent DHS not in that chromatin state. For example, 604 dynamic male-biased DHS are classified as chromatin state E6, and 230 other dynamic male-biased DHS are not classified as chromatin state E6 (604/230 = 2.626), whereas 24,082 static sex-independent DHS are classified as chromatin state E6, and 40,502 static sex-independent DHS are not classified as chromatin state E6 (24,082/40,502 = 0.5946), which gives an enrichment score A/B = 2.626/0.5946 = 4.4. The set of 64,584 static sex-independent DHS was used as the background set for these enrichment calculations.

DHS peak normalization - DNase-seq data to be visualized in the UCSC Genome browser (https://genome.ucsc.edu/) was normalized using the number of sequence reads in each DHS peak region per million mapped sequence reads (reads-in-peaks-per-million, RiPPM) as a scaling-factor [46]. Normalization was carried out using a comprehensive list of DHS peak regions merged across each dataset (i.e., the peak union list),

obtained by concatenating FASTQ files for biological replicates of each control and treatment group, as described above. DHS peak regions identified in both individual liver samples and by analysis of the combined samples were concatenated into a single list, and then the BEDTools *merge* command was used to combine overlapping features to generate a single list of non-overlapping DHS peaks. The fraction of reads in peaks for each sample was then calculated to obtain a scaling factor. Raw sequence read counts were divided by this per-million scaling factor to obtain RiPPM normalized read counts.

Acknowledgements – Supported in part by NIH grant DK121998 (to DJW).

Authors' contributions – AR and DJW conceived of the study. All animal studies and wet lab analyses including STAT5 EMSA gels, isolation of liver nuclei, DNase-seq analysis and library preparation were performed by JC. Data processing, primary computational analysis, preparation of analysis scripts, and data visualization were carried out by AR. All other data analysis and preparation of datasets and figures for publication were carried out by AR and DJW. AR and DJW jointly prepared a preliminary draft of the manuscript. The final manuscript was written by DJW and was reviewed and approved by all the authors. DJW supervised the overall project and edited the final manuscript for publication.

Table 1: Sex-biased DHS	Dynamic male-biased DHS (n=834, 31%)	Static male-biased DHS (n=1895, 69%)	Static female-biased DHS (n=1359)
A. DHS activity (chromatin opening)			
STAT5-high male liver	+++	+	-
STAT5-low male liver	+	+	-
Female liver	+	-	+
B. Sex-biased TF binding sites			
STAT5 (binding in male liver)	+++	+	-
CUX2 (binding in female liver)	+	++	-
FOXA1 (male-biased sites)	+	++	-
FOXA2 (male-biased sites)	+++	+++++	-
FOXA2 (female-biased sites)	-	-	+++
C. Enriched H3 histone marks and chromatin states			
K27ac, K4me1, with DHS (male liver) [State E6]	++	++	-
K27ac and/or K4me1 DHS (male-liver) [States E10, E11]	-	-	++
K36me3 (male-biased)	-	++	-
K36me3 (female-biased)	-	-	++
K27me3 (male-biased)	-	-	+++
K9me3 (female-biased)	++	++	-

Figure legends

Fig. 1. Classification of liver DHS and mapping to liver-expressed genes. A. Shown are the percentages of each indicated liver DHS subset (x-axis) that are classified as weak enhancer, enhancer, insulator or promoter regions [20]. Enrichments of sex-biased DHS for being enhancer, insulator or promoter regions were determined by comparing to a background set comprised of sex-independent DHS (66,116 sites), with the significance determined by Fisher Exact Test with Benjamini-Hochberg p-value adjustment: *, P<0.01; **, P<1e-10; ***, P<1e-50. Black asterisks, enrichment; red asterisks, depletion as compared to the background DHS set. **B.** Cumulative frequency distribution of the distance to the nearest transcription start site in the same TAD for male-biased and female-biased enhancer (e), insulator (i), and promoter (p) DHS. **C.** ChromHMM emission probabilities for each of the 14 chromatin states developed for male and female mouse liver [18], which serves as a reference for data shown in panel D and in Fig. 7. Summary descriptions of the characteristics of each state are shown at the left. **D.** Chromatin state distributions for each of the indicated sex-independent and sex-biased DHS sets. Chromatin state data for male liver was used for male-biased DHS; data for female liver was used for female-biased DHS. See Table S2 for full datasets.

Fig. 2. Discovery and characterization of dynamic and static male-biased DHS. A. Model showing pulsatile, male plasma GH pattern, with mice sampled between GH pulses, when STAT5 is inactive and cytoplasmic (STAT5-low, *blue*), or at a peak of GH, when liver STAT5 is activated to its homodimeric, nuclear DNA-binding form, which enables STAT5 to open chromatin and bind to its consensus motif, TTCNNNGAA (STAT5-high, *red*). This intermittent (pulsatile) activation STAT5 leads to chromatin opening and closing at dynamic DHS (*top*). Static DHS also bind STAT5 intermittently but remain open between plasma GH pulses (*bottom*). **B.** EMSA of STAT5 DNA-binding activity in liver extracts prepared from individual male mice. These data represent liver extracts from 1 of 3 separate cohorts of mice; the other 2 cohorts are shown in Fig. S1. Labels at top indicate the DNase-seq library prepared from each sample. Red labels at top indicate STAT5-high activity based on the EMSA patterns displayed, and blue labels indicate STAT5-low activity livers. Numbers at bottom: number of each mouse whose liver was analyzed. Samples were all run on the same gel. **C.** Principal component (PC) analysis of the distributions of DNase-seq reads per kilobase per million for the set of diffReps differential sites that are more open in STAT5-high as compared to STAT5-low livers. Eigenvector values for principal component 1 are shown for the individual STAT5-high and STAT5-low liver DNase-seq libraries, calculated for the top 600 (*left*) or the top 200 (*right*) diffReps differential sites ordered by decreasing fold-change in chromatin opening. Dotted red line: empirical cutoff separating STAT5-high from STAT5-low liver DNase-seq samples; dotted black circles: outlier samples in each dataset. **D.** Boxplots of DNase-seq activity, in log2(reads per kilobase per million mapped reads), across the top 200 diffReps differential sites that open for the STAT5-high liver DNase-seq libraries (*red* bars), for the STAT5-low liver DNase-seq libraries (*blue* bars), and for DNase-seq libraries for 9 individual male mouse liver samples generated by the ENCODE consortium (*green* bars, replicates 5 to 13, marked on x-axis). Thick dashed black line: arbitrary cutoff used to separate STAT5-high and STAT5-low liver samples. Samples from each group that did not pass the cutoff (*red asterisks at bottom*) are the same outlier livers circled in panel B. **E.** Normalized DNase-I cut site aggregate plots for STAT5-high (*red*) and STAT5-low male livers (*blue*), and for female livers (*black*). Peak values are as shown. Cut sites were aggregated across the sets of diffReps-identified DHS that open (*left*) or close (*right*) in response to endogenous STAT5 pulses in male liver, i.e., sites that show greater diffReps normalized DNase-seq signal intensity in STAT5-high compared to STAT5-low male livers. **F.** Venn diagram indicating overlap between endogenous STAT5 pulse-opened DHS sets identified by diffReps (2,832 sites that respond to liver STAT5 activity in a dynamic manner, which map to a total of 2,373 of the 70,211 DHS comprising the standard reference DHS set) and the indicated sets of sex-biased and sex-independent DHS that do not respond to a change in liver STAT5 activity, i.e., are static DHS. See Table S2A, columns H and I for full listing.

Fig. 3. Dynamic and static male-biased DHS. A. Normalized DNase-I cut site aggregate plots for STAT5-high (*red*) and STAT5-low male livers (*blue*) and female livers (*black*) across the genomic regions included in the indicated sets of male-biased DHS (plots 1, 2) and sex-independent DHS (plots 3, 4), separated into subsets of DHS that either do (plots 1, 3) or do not bind STAT5 by ChIP-seq analysis (plots 2, 4). Plots 5-8 show corresponding plots for each of the indicated dynamic and static DHS sets. See Table S6 for peak normalized DNase-I site values. **B.** Bar plot showing the number (values above bars) and percent of static and dynamic DHS

with one or more occurrences of the indicated STAT5 motif. Based on FIMO scan of the 70,211 standard reference DHS sequences for the occurrence of a TRANSFAC STAT5 motif (box). Dashed horizontal line: background, genome-wide level of STAT5 motifs at static sex-independent DHS. **C.** Percentage of dynamic (*left*) and static (*right*) DHS with zero or more occurrences of the STAT5 motif. X-axis: number of motif occurrences in each DHS (n =0-9), y-axis: % DHS in each group with the corresponding number of motifs. Patterns for dynamic female-biased DHS represent a total of only 7 DHS and are thus not reliable.

Fig. 4. DHS responsive to hypophysectomy, their enrichment for class I and II sex-biased gene targets and responses to GH pulse replacement. A. Model for effects of hypophysectomy, which ablates pulsatile pituitary secretion of GH, on STAT5-induced chromatin opening. DHS that close following hypophysectomy due to the loss of STAT5 binding reopen within 30 min of exogenous GH treatment, which rapidly reactivates STAT5. **B.** Numbers of liver DHS that open or close following hypophysectomy (Hx) and in response to a single injection of GH given to hypox male (M) or female (F) mice and euthanized 30, 90 or 240 min later. DHS opening and closing were determined compared to the indicated controls. **C.** Distributions of sex-biased and sex-independent DHS that open, close or are unchanged (static) hypophysectomized (Hx) male and female mice. See Table S4G for full details. **D.** Enrichment of hypophysectomy-responsive DHS as compared to hypophysectomy-unresponsive (static) DHS for mapping to the four indicated classes of sex-specific genes (see Fig. S4). Class I and II sex-specific genes were identified from RNA-seq gene expression data collected from intact and hypox male and female liver samples (Table S3). The bottom section of the table shows the total number and percentage of sex-specific genes in each of the four indicated sex-biased gene classes that respond to hypophysectomy, as indicated (Table S4H). **E.** Subset of all dynamic DHS (Fig. 2F) that close following hypophysectomy in male mouse liver (n=1,487) and then respond to GH pulse replacement at the indicated time points. Total number of dynamic DHS shown here is lower than the full set of 2,373 dynamic DHS (Fig. 2F), as 70 of these DHS were not identified as DHS in the hypophysectomy study (Table S4G).

Fig. 5. DHS activity aggregate plots for hypophysectomy and GH replacement pulse time course study. **A.** Normalized DNase-I cut site aggregate plots showing the effects of hypophysectomy of male (MHx) and female mice (FHx) and of GH pulse treatment (MHx+GH) after 30, 90 and 240 min compared to intact females for each of the indicated sets of static and dynamic DHS (c.f., Fig. 3A). Reference values for normalized DNase-I cut site activity in intact male liver are from Fig. 3 and Table S6. **B.** Plots are shown as in **A** for the indicated subsets of 2,729 male-biased DHS (plots 1 and 2), and for the indicated subsets of the set of 66,116 sex-independent DHS (plots 3 and 4), i.e., DHS subsets with STAT5 bound (plots 1 and 3) or without STAT5 bound (plots 2 and 4), based on ChIP-seq data for STAT5 in male mouse liver from [15].

Fig. 6. Enrichments of sex-biased histone marks at sex-biased and sex-independent dynamic and static DHS. A. Enrichments of male-biased histone marks, and **B.** enrichments of female-biased histone marks. Data is presented as bar graphs showing significant enrichments, and significant depletions, of the 6 indicated sets of liver histone marks for the four indicated sets of DHS (c.f., Fig. 2F). Enrichment data are graphed as 6 sets of four bars each, separated by vertical blue lines and ordered from 1 to 4, as shown at the right and as marked above select bars. The set of 64,584 static sex-independent DHS was used as the background for the enrichment calculations. Enrichment scores (and depletion scores) are indicated by bar height (Y-axis values), and Fisher Exact test significance values (log p-values, indicated by bar color) are shown for all values that are significant at p<E-03. Values that did not meet this significance threshold are graphed as ES = 1 (horizontal dashed green line); thus, all bars shown, except those graphed at ES = 1.0, represent statistically significant enrichment or depletion. Full details on the numbers of sites, the source publications used to identify these genomic regions, and corresponding BED files are shown in Table S7. **C.** Proposed model for chromatin states adopted by dynamic male-biased DHS, by static male-biased DHS and by static female-biased DHS in male liver (*left*) and in female liver (*right*) in response to the stimulatory and/or repressive actions of plasma GH pulses (in male liver) and near-continuous GH exposure (in female liver). Histone H3 marks are shown by small colored ovals attached to histone tails (box). H3K27me3 is specifically used to repress chromatin in at female-biased DHS in male liver, and H3K9me3 is specifically used to repress chromatin in at male-biased DHS in female liver. The degree of chromatin opening is indicated by the relative distance between nucleosomes. Black arrow indicates the DHS region is active in stimulating gene transcription upon interaction with a nearby or distal gene promoter.

Fig. 7. Enrichments of chromatin states at sex-biased and sex-independent dynamic and static DHS. A. Enrichments of male liver chromatin states at each of the 4 indicated DHS, and **B.** enrichments of female liver chromatin states at each of the 4 indicated DHS sets. Data are presented as described in Fig. 6 and used the set of 64,584 static sex-independent DHS as background for the enrichment calculations. Full details on the numbers of sites, the source publications used to identify these genomic regions, and corresponding BED files are shown in Table S7.

Fig. 8. STAT5 regulates sex-dependent hepatocyte gene expression at three distinct steps. Sex-biased chromatin opening (step 1): GH pulse-induced chromatin opening at dynamic male-biased DHS is driven by pulsatile GH activation of STAT5 in male liver, whereas persistent activation of STAT5 in female liver is associated with static female-biased chromatin opening. Sex-biased transcriptional activation (step 2): Sex differences in chromatin accessibility enable GH-activated STAT5 to bind chromatin in a sex-biased manner and induce the transcriptional activation of sex-biased genes. Sex-based transcriptional repression (step 3): The sex-biased regulatory genes include the GH/STAT5-dependent repressor proteins BCL6 (male-biased) and CUX2 (female-specific), which reinforce sex differences in transcription by preferentially suppressing the expression of female-biased and male-biased genes, respectively, as indicated.

References

[1] D.J. Waxman, M.G. Holloway, Sex differences in the expression of hepatic drug metabolizing enzymes, Mol Pharmacol, 76 (2009) 215-228.

[2] M.C. Vazquez-Borrego, M. Del Rio-Moreno, R.D. Kineman, Towards Understanding the Direct and Indirect Actions of Growth Hormone in Controlling Hepatocyte Carbohydrate and Lipid Metabolism, Cells, 10 (2021).

[3] L.E. Dichtel, J. Cordoba-Chacon, R.D. Kineman, Growth Hormone and Insulin-Like Growth Factor 1 Regulation of Nonalcoholic Fatty Liver Disease, J Clin Endocrinol Metab, 107 (2022) 1812-1824.

[4] D. Kaltenecker, M. Themanns, K.M. Mueller, K. Spirk, T. Suske, O. Merkel, L. Kenner, A. Luis, A. Kozlov, J. Haybaeck, M. Muller, X. Han, R. Moriggl, Hepatic growth hormone - JAK2 - STAT5 signalling: Metabolic function, non-alcoholic fatty liver disease and hepatocellular carcinoma progression, Cytokine, 124 (2019) 154569.

[5] M. Oxley, H. Francis, K. Sato, Growth Hormone Signaling in Liver Diseases: Therapeutic Potentials and Controversies, Semin Liver Dis, 43 (2023) 24-30.

[6] V. Wauthier, A. Sugathan, R.D. Meyer, A.A. Dombkowski, D.J. Waxman, Intrinsic sex differences in the early growth hormone responsiveness of sex-specific genes in mouse liver, Mol Endocrinol, 24 (2010) 667-678.

[7] B. Brie, M.C. Ramirez, C. De Winne, F. Lopez Vicchi, L. Villarruel, E. Sorianello, P. Catalano, A.M. Ornstein, D. Becu-Villalobos, Brain Control of Sexually Dimorphic Liver Function and Disease: The Endocrine Connection, Cell Mol Neurobiol, 39 (2019) 169-180.

[8] L.S. Farhy, C.Y. Bowers, J.D. Veldhuis, Model-projected mechanistic bases for sex differences in growth hormone regulation in humans, Am J Physiol Regul Integr Comp Physiol, 292 (2007) R1577-1593.

[9] J.A. Chowen, L.M. Garcia-Segura, S. Gonzalez-Parra, J. Argente, Sex steroid effects on the development and functioning of the growth hormone axis, Cell Mol Neurobiol, 16 (1996) 297-310.

[10] M.C. Ramirez, G.M. Luque, A.M. Ornstein, D. Becu-Villalobos, Differential neonatal testosterone imprinting of GH-dependent liver proteins and genes in female mice, J Endocrinol, 207 (2010) 301-308.

[11] D.J. Waxman, G.A. Dannan, F.P. Guengerich, Regulation of rat hepatic cytochrome P-450: age-dependent expression, hormonal imprinting, and xenobiotic inducibility of sex-specific isoenzymes, Biochemistry, 24 (1985) 4409-4417.

[12] D.J. Waxman, C. O'Connor, Growth hormone regulation of sex-dependent liver gene expression, Mol Endocrinol, 20 (2006) 2613-2629.

[13] K.H. Clodfelter, M.G. Holloway, P. Hodor, S.H. Park, W.J. Ray, D.J. Waxman, Sex-dependent liver gene expression is extensive and largely dependent upon signal transducer and activator of transcription 5b (STAT5b): STAT5b-dependent activation of male genes and repression of female genes revealed by microarray analysis, Mol Endocrinol, 20 (2006) 1333-1351.

[14] M.J. Waters, The growth hormone receptor, Growth Horm IGF Res, 28 (2016) 6-10.

[15] Y. Zhang, E.V. Laz, D.J. Waxman, Dynamic, sex-differential STAT5 and BCL6 binding to sex-biased, growth hormone-regulated genes in adult mouse liver, Mol Cell Biol, 32 (2012) 880-896.

[16] J. Connerney, D. Lau-Corona, A. Rampersaud, D.J. Waxman, Activation of Male Liver Chromatin Accessibility and STAT5-Dependent Gene Transcription by Plasma Growth Hormone Pulses, Endocrinology, 158 (2017) 1386-1405.

[17] D. Lau-Corona, A. Suvorov, D.J. Waxman, Feminization of Male Mouse Liver by Persistent Growth Hormone Stimulation: Activation of Sex-Biased Transcriptional Networks and Dynamic Changes in Chromatin States, Mol Cell Biol, 37 (2017).

[18] A. Sugathan, D.J. Waxman, Genome-wide analysis of chromatin states reveals distinct mechanisms of sex-dependent gene regulation in male and female mouse liver, Mol Cell Biol, 33 (2013) 3594-3610.

[19] G. Ling, A. Sugathan, T. Mazor, E. Fraenkel, D.J. Waxman, Unbiased, genome-wide in vivo mapping of transcriptional regulatory elements reveals sex differences in chromatin structure associated with sex-specific liver gene expression, Mol Cell Biol, 30 (2010) 5531-5544.

[20] B.J. Matthews, D.J. Waxman, Computational prediction of CTCF/cohesin-based intra-TAD loops that insulate chromatin contacts and gene expression in mouse liver, Elife, 7 (2018).

[21] D.J. Waxman, R.D. Kineman, Sex matters in liver fat regulation, Science, 378 (2022) 252-253.

[22] J. Nikkanen, Y.A. Leong, W.C. Krause, D. Dermadi, J.A. Maschek, T. Van Ry, J.E. Cox, E.J. Weiss, O. Gokcumen, A. Chawla, H.A. Ingraham, An evolutionary trade-off between host immunity and metabolism drives fatty liver in male mice, Science, 378 (2022) 290-295.

[23] M. Baik, J.H. Yu, L. Hennighausen, Growth hormone-STAT5 regulation of growth, hepatocellular carcinoma, and liver metabolism, Ann N Y Acad Sci, 1229 (2011) 29-37.

[24] B.J. Matthews, D.J. Waxman, Impact of 3D genome organization, guided by cohesin and CTCF looping, on sex-biased chromatin interactions and gene expression in mouse liver, Epigenetics Chromatin, 13 (2020) 30.

[25] P. Hao, D.J. Waxman, STAT5 Regulation of Sex-Dependent Hepatic CpG Methylation at Distal Regulatory Elements Mapping to Sex-Biased Genes, Mol Cell Biol, 41 (2021).

[26] M.G. Holloway, Y. Cui, E.V. Laz, A. Hosui, L. Hennighausen, D.J. Waxman, Loss of sexually dimorphic liver gene expression upon hepatocyte-specific deletion of Stat5a-Stat5b locus, Endocrinology, 148 (2007) 1977-1986.

[27] G.S. Tannenbaum, H.K. Choi, W. Gurd, D.J. Waxman, Temporal relationship between the sexually dimorphic spontaneous GH secretory profiles and hepatic STAT5 activity, Endocrinology, 142 (2001) 4599-4606.

[28] L. Shen, N.Y. Shao, X. Liu, I. Maze, J. Feng, E.J. Nestler, diffReps: detecting differential chromatin modification sites from ChIP-seq data with biological replicates, PLoS One, 8 (2013) e65598.

[29] R.D. Meyer, E.V. Laz, T. Su, D.J. Waxman, Male-specific hepatic Bcl6: growth hormone-induced block of transcription elongation in females and binding to target genes inversely coordinated with STAT5, Mol Endocrinol, 23 (2009) 1914-1926.

[30] T.L. Conforto, Y. Zhang, J. Sherman, D.J. Waxman, Impact of CUX2 on the female mouse liver transcriptome: activation of female-biased genes and repression of male-biased genes, Mol Cell Biol, 32 (2012) 4611-4627.

[31] T.L. Conforto, G.F.t. Steinhardt, D.J. Waxman, Cross Talk Between GH-Regulated Transcription Factors HNF6 and
CUX2 in Adult Mouse Liver, Mol Endocrinol, 29 (2015) 1286-1302.

[32] Z. Li, G. Tuteja, J. Schug, K.H. Kaestner, Foxa1 and Foxa2 are essential for sexual dimorphism in liver cancer, Cell,
148 (2012) 72-83.

[33] A. Balsalobre, J. Drouin, Pioneer factors as master regulators of the epigenome and cell fate, Nat Rev Mol Cell Biol, 23
(2022) 449-464.

[34] K.S. Zaret, Pioneer Transcription Factors Initiating Gene Network Changes, Annu Rev Genet, 54 (2020) 367-385.

[35] D. Lau-Corona, W.K. Bae, L. Hennighausen, D.J. Waxman, Sex-biased genetic programs in liver metabolism and liver
fibrosis are controlled by EZH1 and EZH2, PLoS Genet, 16 (2020) e1008796.

[36] P. Rotwein, Regulation of gene expression by growth hormone, Mol Cell Endocrinol, 507 (2020) 110788.

[37] S.H. Kim, M.J. Park, Effects of growth hormone on glucose metabolism and insulin resistance in human, Ann Pediatr
Endocrinol Metab, 22 (2017) 145-152.

[38] C.N. Goldfarb, K. Karri, M. Pyatkov, D.J. Waxman, Interplay Between GH-regulated, Sex-biased Liver Transcriptome
and Hepatic Zonation Revealed by Single-Nucleus RNA Sequencing, Endocrinology, 163 (2022).

[39] C.A. Gebert, S.H. Park, D.J. Waxman, Regulation of signal transducer and activator of transcription (STAT) 5b
activation by the temporal pattern of growth hormone stimulation, Mol Endocrinol, 11 (1997) 400-414.

[40] D.J. Waxman, P.A. Ram, S.H. Park, H.K. Choi, Intermittent plasma growth hormone triggers tyrosine phosphorylation
and nuclear translocation of a liver-expressed, Stat 5-related DNA binding protein. Proposed role as an intracellular
regulator of male-specific liver gene transcription, J Biol Chem, 270 (1995) 13262-13270.

[41] A.A. Able, J.A. Burrell, J.M. Stephens, STAT5-Interacting Proteins: A Synopsis of Proteins that Regulate STAT5
Activity, Biology (Basel), 6 (2017).

[42] Y. Fu, J. Wang, G. Panangipalli, B.J. Ulrich, B. Koh, C. Xu, R. Kharwadkar, X. Chu, Y. Wang, H. Gao, W. Wu, J. Sun,
R.S. Tepper, B. Zhou, S.C. Janga, K. Yang, M.H. Kaplan, STAT5 promotes accessibility and is required for BATF-
mediated plasticity at the Il9 locus, Nat Commun, 11 (2020) 4882.

[43] S.J. Frank, Classical and novel GH receptor signaling pathways, Mol Cell Endocrinol, 518 (2020) 110999.

[44] H. Gingras, O. Cases, M. Krasilnikova, G. Bérubé, A. Nepveu, Biochemical characterization of the mammalian Cux2
protein, Gene, 344 (2005) 273-285.

[45] D. Lau-Corona, H. Ma, C. Vergato, A. Sarmento-Cabral, M. Del Rio-Moreno, R.D. Kineman, D.J. Waxman,
Constitutively Active STAT5b Feminizes Mouse Liver Gene Expression, Endocrinology, 163 (2022).

[46] N.J. Lodato, A. Rampersaud, D.J. Waxman, Impact of CAR Agonist Ligand TCPOBOP on Mouse Liver Chromatin
Accessibility, Toxicol Sci, 164 (2018) 115-128.

[47] N. Percie du Sert, A. Ahluwalia, S. Alam, M.T. Avey, M. Baker, W.J. Browne, A. Clark, I.C. Cuthill, U. Dirnagl, M.
Emerson, P. Garner, S.T. Holgate, D.W. Howells, V. Hurst, N.A. Karp, S.E. Lazic, K. Lidster, C.J. MacCallum, M.
Macleod, E.J. Pearl, O.H. Petersen, F. Rawle, P. Reynolds, K. Rooney, E.S. Sena, S.D. Silberberg, T. Steckler, H. Würbel,
Reporting animal research: Explanation and elaboration for the ARRIVE guidelines 2.0, PLoS Biol, 18 (2020) e3000411.

[48] B. Langmead, S.L. Salzberg, Fast gapped-read alignment with Bowtie 2, Nat Methods, 9 (2012) 357-359.

[49] J. Feng, T. Liu, B. Qin, Y. Zhang, X.S. Liu, Identifying ChIP-seq enrichment using MACS, Nat Protoc, 7 (2012) 1728-
1740.

[50] S.G. Landt, G.K. Marinov, A. Kundaje, P. Kheradpour, F. Pauli, S. Batzoglou, B.E. Bernstein, P. Bickel, J.B. Brown,
P. Cayting, Y. Chen, G. DeSalvo, C. Epstein, K.I. Fisher-Aylor, G. Euskirchen, M. Gerstein, J. Gertz, A.J. Hartemink, M.M.
Hoffman, V.R. Iyer, Y.L. Jung, S. Karmakar, M. Kellis, P.V. Kharchenko, Q. Li, T. Liu, X.S. Liu, L. Ma, A. Milosavljevic,
R.M. Myers, P.J. Park, M.J. Pazin, M.D. Perry, D. Raha, T.E. Reddy, J. Rozowsky, N. Shoresh, A. Sidow, M. Slattery, J.A.
Stamatoyannopoulos, M.Y. Tolstorukov, K.P. White, S. Xi, P.J. Farnham, J.D. Lieb, B.J. Wold, M. Snyder, ChIP-seq
guidelines and practices of the ENCODE and modENCODE consortia, Genome Res, 22 (2012) 1813-1831.

[51] C.Y. McLean, D. Bristor, M. Hiller, S.L. Clarke, B.T. Schaar, C.B. Lowe, A.M. Wenger, G. Bejerano, GREAT
improves functional interpretation of cis-regulatory regions, Nat Biotechnol, 28 (2010) 495-501.

[52] Y. Tanigawa, E.S. Dyer, G. Bejerano, WhichTF is functionally important in your open chromatin data?, PLoS Comput
Biol, 18 (2022) e1010378.

[53] T. Melia, D.J. Waxman, Sex-Biased lncRNAs Inversely Correlate With Sex-Opposite Gene Coexpression Networks in
Diversity Outbred Mouse Liver, Endocrinology, 160 (2019) 989-1007.

[54] V. Matys, O.V. Kel-Margoulis, E. Fricke, I. Liebich, S. Land, A. Barre-Dirrie, I. Reuter, D. Chekmenev, M. Krull, K.
Hornischer, N. Voss, P. Stegmaier, B. Lewicki-Potapov, H. Saxel, A.E. Kel, E. Wingender, TRANSFAC and its module
TRANSCompel: transcriptional gene regulation in eukaryotes, Nucleic Acids Res, 34 (2006) D108-110.

[55] C.E. Grant, T.L. Bailey, W.S. Noble, FIMO: scanning for occurrences of a given motif, Bioinformatics, 27 (2011)
1017-1018.

[56] Z. Shao, Y. Zhang, G.C. Yuan, S.H. Orkin, D.J. Waxman, MAnorm: a robust model for quantitative comparison of
ChIP-Seq data sets, Genome Biol, 13 (2012) R16.

1178

Fig. 1

Fig. 2

Fig. 4

A — Hypox / GH / STAT5 STAT5 / TTCN₃GAA / + GH (30 min)

B
- dDHS_open (green)
- dDHS_close (dark red)

Y-axis: Number of DHS

Categories: MHx vs M, FHx vs F, MHx GH30 vs MHx, MHx GH90 vs MHx, MHx GH240 vs MHx

C

DHS class	% Opening (Male Hx / Female HX)	% Closing	% Static
Dynamic Male-biased	0	76	24
Static Male-biased	1	32	67
Dynamic Male-biased	5	13	82
Static Male-biased	9	7	84
Static Female-biased	17	2	81
Static Female-biased	2	12	86
Dynamic Sex-indep	0	58	41
Static Sex-indep	4	6	90
Dynamic Sex-indep	6	16	78
Static Sex-indep	8	6	86

E

Dynamic, STAT5-High responsive DHS — *Male Hypox*
- Opening DHS 0.2%
- Static DHS 35%
- Closing DHS 65% (n=1,487)
- +GH pulse → DHS reopening
  - n=1,063 (71% at 30 min)
  - n=1,119 (75% at 90 min)
  - n=477 (32% at 240 min) } 1,133 DHS
  - + Unresponsive DHS (n=354)

D

DHS –> Hypox response class	DHS closing in Male Hx liver (Male Hypox/Male control)		DHS closing in Female Hx liver (Female Hypox/Female control)		DHS opening in Male Hx liver (Male Hypox/Male control)		DHS opening in Female Hx liver (Female Hypox/Female control)	
	Enrichment score and adj P-value for *chromatin closing*				Enrichment score and adj P-value for *chromatin opening*			
Male class I	5.7	1.3E-69	3.9	1.69E-21	0.3	9.55E-04	0.3	6.03E-08
Female class I	2.9	4.1E-11	7.8	6.64E-46	0.9	8.83E-01	0.2	3.38E-06
Male class II	1.3	2.2E-01	0.7	4.35E-01	1.7	5.62E-02	1.9	4.83E-04
Female class II	1.3	1.7E-01	1.3	1.45E-01	3.6	1.80E-15	0.6	2.03E-02

Hypox response class of gene	# genes **Down**: Male Hypox	% genes in sex-biased class	# genes **Down**: Female Hypox	% genes in sex-biased class	# genes **Up**: Male Hypox	% genes in sex-biased class	# genes **Up**: Female Hypox	% genes in sex-biased class
Male class I	202	100.0%	89	44.1%	0	0.0%	3	1.5%
Female class I	35	26.1%	134	100.0%	9	6.7%	0	0.0%
Male class II	0	0.0%	0	0.0%	40	44.4%	90	100.0%
Female class II	0	0.0%	0	0.0%	178	100.0%	70	39.3%

Fig. 6

Fig. 7

A Enrichment of **male** liver chromatin states

1. Dynamic male (n=834)
2. Static male (n=1,895)
3. Dynamic sex-independ (n=1,532)
4. Static female (n=1,359)

B Enrichment of **female** liver chromatin states

1. Dynamic male (n=834)
2. Static male (n=1,895)
3. Dynamic sex-independ (n=1,532)
4. Static female (n=1,359)

Fig. 8

Supplemental figures

Fig. S1. EMSA analysis of STAT5 DNA-binding activity in liver extracts prepared from individual male mice. Shown here are EMSA data for 2 of 3 separate cohorts of mice; the 3rd cohort is shown in Fig. 2B. Livers whose nuclei were used for DNase-seq analysis are marked at the top (Table S1); red labels at top indicate STAT5-high activity based on the EMSA patterns displayed, and blue labels indicate STAT5-low activity livers. Also shown here are EMSA activity for 8 other livers, which have low to intermediate STAT5 EMSA activity and were not included in downstream analysis. Numbers at bottom: liver sample numbers from each mouse cohort.

Fig. S2. STAT5 binding is associated with chromatin opening. A. Bar graph of the number and percentage of dynamic and static DHS that have one or more STAT5 binding sites, based on STAT5 ChIP-seq data for mouse liver. BEDTools was used to determine the overlap between the merged list of 15,094 STAT5 binding sites [1] and the standard reference set of 70,211 liver DHS used in this study. **B.** Distribution of STAT5 ChIP-seq signal intensity values for the sets of dynamic and static DHS. For each DHS that contains a STAT5 binding site (numbers shown in **A**), the corresponding normalized STAT5 ChIP-seq read count (average of STAT5 male-high samples and STAT5 female-high liver samples; [1]) was obtained and used to calculate the indicated distributions. **C.** Normalized DNase-I cut site aggregate plots for male livers at STAT5-high, STAT5-low, and female liver for the indicated sets of male-biased, female-biased, and sex-independent DHS, analyzed separately for dynamic DHS subsets (plots 1A, 1B, 4A, 4B) and static DHS subsets (plots 2A, 2B, 3A, 3B, 5A. 5B). Top row: STAT5-bound subsets; bottom row: non-STAT5-bound subsets. See Table S6 for DNase-seq aggregate plot peak values.

Fig. S3. TF motifs discovered de novo from sets of DHS sequences. De-novo motif discovery was carried out on the sets of dynamic and static DHS sequences using the MEME and DREME algorithms of MEME-ChIP [2]. Listed are the de-novo discovered motifs and their associated p-values for: **A**, dynamic male-biased DHS (834 sites); **B**, dynamic sex-independent DHS (1,532 sites); **C**, static male-biased DHS (1,895 sites); **D**, static female-biased DHS (1,359 sites); and **E**, static sex-independent DHS (64,584 sites). Shown are the top most significant motifs. No motifs were discovered from the very small number of dynamic female-biased DHS (7 sites). The presence of the STAT5B motif is a distinguishing characteristic that separates the dynamic from static DHS, which supports our hypothesis that direct binding of STAT5 plays a key role in chromatin opening in response to GH pulses at these dynamic DHS.

Fig. S4. **Sex-biased genes can be classified based on their pituitary hormone-dependence. A**. Model for plasma GH profile dependence and responses of class I and class II male-biased and female-biased genes to pituitary hormone ablation in each sex by hypophysectomy ('hypox'). Figure from [3]. **B**. Sex-specific genes were identified from RNA-seq data from intact male and female mouse liver based on a gene list comprised of 24,197 RefSeq and 3,152 multi-exonic lncRNA genes. First, sex-specific genes were identified with |fold-change| > 1.5, adjusted p-value < 0.05 (for RefSeq genes), and |fold-change| > 2.0, adjusted p-value < 0.05 (for lncRNA genes), with FPKM > 0.25 for the sex with greater signal intensity for both RefSeq and lncRNA datasets (see Methods). Sex-specific genes that were responsive to hypophysectomy (|fold-change| > 2 and an adjusted p-value < 0.05) were further classified into class I, II and corresponding subclasses (IA, IB, IC, IIA, and IIB) based on RNA-seq data from intact and hypox

male and female liver samples. Shown are the full list of RefSeq and lncRNA genes. See Table S3 for full gene listings.

Fig. S5. DNase cut site aggregate plots for the GH time course DHS data. Shown are normalized DNase-I cut site aggregate plots for livers from hypophysectomized male mice (MHx), hypophysectomized male mice treated with GH (MHx+GH) then euthanized after 30, 90 or 240 min, and intact female and hypophysectomized female mice (FHx) across various sets of dynamic and static male-biased DHS, static female-biased DHS, and static sex-independent DHS (see Fig. S1C). Each DHS set was separated into subsets based on STAT5 binding, as determined by Chip-seq (top row: STAT5-bound DHS subsets; bottom row: corresponding non-STAT5-bound DHS subsets). Key results: MHx mice treated with GH for either 30 or 90 min show the largest degree of chromatin opening in the set of 710 dynamic male-biased DHS that bind STAT5. Chromatin opening decreased substantially after 240 min (plot 1A), at which time the activation of liver STAT5 DNA binding activity is fully reversed [4]. Smaller increases in chromatin opening were observed with GH pulse treatment at the subset of 124 dynamic male-biased DHS that did not bind STAT5 (plot 1B), consistent with their dynamic responses to endogenous GH pulsation, and suggesting that chromatin opening at these sites proceeds by a distinct mechanism than at the STAT5-bound dynamic male-biased sites. Static male-biased DHS with STAT5 bound (597 sites) showed a more modest increase in chromatin opening with GH pulse treatment (c.f., higher basal level in MHx control and lower induced level with GH pulse; plot 2A), while static male-biased DHS without STAT5 binding (1,298 sites) showed little or no GH pulse responsiveness (plot 2B). Importantly, the increase in chromatin opening at the STAT5-bound static male-biased DHS largely persists at 240 min (plot 2A), in contrast to the more substantial decline in chromatin opening seen for at STAT5-bound dynamic male-biased DHS (plot 1A). This suggests that, while STAT5 can open chromatin at the static male-biased DHS, it is not required to maintain chromatin accessibility between the naturally occurring endogenous plasma GH pulses. Chromatin opening at female-biased DHS was decreased by hypophysectomy, both at the 258 sites bound by STAT5 and at the 1101 sites that did not show STAT5 binding (plots 3A and 3B). Further, GH pulse treatment of MHx mice stimulated a modest decrease in liver chromatin accessibility at both sets of female-biased DHS. Finally, the dynamic, but not the static sex-independent DHS showed large increases in chromatin opening following GH pulse treatment (plot 4A and plot 5A), similar to the dynamic male-biased DHS. Overall, DHS that bind STAT5 showed higher levels of chromatin opening than DHS that do not bind STAT5 (top row vs. bottom row). Notably, this difference in chromatin accessibility is preserved in hypophysectomized male and female liver, even though liver STAT5 is inactive and cannot bind DHS under these conditions due to the absence of GH stimulation, indicating a role for other pituitary-determined factors in chromatin opening.

Fig. S6. Impact of hypophysectomy and GH pulse replacement on set of 2,373 dynamic DHS.

S6A. Four-oval Venn diagram showing the number of STAT5-high DHS that were close following hypophysectomy and/or open in response to a single exogenous GH pulse given to hypophysectomized male mice. The set of STAT5-high (dynamic) DHS (n = 2,373 sites; Fig. 2E) was analyzed to determine their overlap with the set of DHS that close following hypophysectomy (n = 1,487 sites) or DHS that open when a single exogenous GH pulse is given to hypophysectomized male mice and liver tissue collected after 30 min (n = 1, 475 sites), 90 min (n = 1,393 sites) or 240 min (n= 547 sites) (see Fig. 4 and Table S2: column I = "dynamic" combined with columns AK-AM = "dDHS_open").

S6B. Flowchart of STAT5-high DHS (i.e., n = 2,373 dynamic DHS) identifying hypox-responsive and exogenous GH pulse-responsive DHS in male mouse liver. Subsections of the 4-oval Venn diagram (shown in A) are used to illustrate the DHS subsets defined by the flowchart (7 subsets), including a set of 399 DHS that do not undergo chromatin closing following hypophysectomy and also do not undergo chromatin opening following an exogenous GH pulse.

S6C. Boxplot analysis showing the magnitude of chromatin opening between STAT5-high and STAT5-low male livers for the 7 DHS subsets. The x-axis shows the DHS subsets numbered 1-7 with the corresponding number of DHS sites (as defined in B). RiPPM normalized DNase-seq read counts were obtained from the STAT5-high and STAT5-low male liver samples for the standard reference set of 70,211 DHS (Table S2). The relative magnitude of chromatin opening (i.e., fold-change value) was calculated from the ratio of STAT5-high/STAT5-low RiPPM-normalized DNase-seq read counts in the DHS region. Shown are the distributions of fold-change values for DHS subsets numbered 1-7 defined from the flowchart in B. A Wilcoxon rank-sum test with Benjamini–Hochberg p-value adjustment was used to determine significant differences between distributions of fold-change values (*P < 0.05, **P < 1e-03, ***P < 1e-10). See Table S6 for DNase-seq aggregate plot peak values.

S6D. DNase-I cut site aggregate plots for the 7 DHS subsets defined in B. See Table S6 for DNase-seq aggregate plot peak values. Key findings shown in B, C and D: Another dynamic DHS subset, comprised of 399 DHS, was unresponsive to hypophysectomy and to GH pulse treatment (**B**). These DHS showed weaker mean chromatin accessibility and lower differential accessibility between STAT5-high and STAT5-low livers than the hypophysectomy and exogenous GH responsive sites (**C, D**). Of the STAT5-high DHS (n=2,373 sites), the subset that responded to hypox and/or a single exogenous GH pulse (set 3, n=1,974) showed significantly greater chromatin opening induced by an endogenous GH/STAT5 pulse than the non-responsive subset (n=399) in pituitary-intact male livers (set 3 vs. set 2). Data in **D** validate the large increase in chromatin accessibility in the set of responsive DHS compared to non-responsive DHS (set 3 vs. set 2). Of the responsive DHS (n=1,974 sites), the subset that was opened by an exogenous GH pulse (set 5, n=1,620 DHS) showed greater chromatin opening induced by an endogenous GH/STAT5 pulse than the DHS subset not opened by a single exogenous GH pulse (n=354) (set 4), a finding that was confirmed by the DNase-I aggregate cutting profiles shown in **D**. Of the GH pulse opened DHS (n=1,620 sites), the subset that was closed following hypophysectomy (n=1,133 sites) showed greater chromatin opening due to an endogenous GH/STAT5 pulse than the subset that was not closed following hypophysectomy (n=487 sites) (set 7 vs. set 6), as was also confirmed by DNase-I aggregate cutting profiles in D. Further, the DHS that closed after hypophysectomy and then reopen following an exogenous GH pulse at any of the three time points (30, 90, or 240 min) (n=1,133 sites; see Fig. 4D) showed the largest magnitude (**C**) and relative levels of chromatin opening (**D**) due an endogenous GH/STAT5 pulse when compared to the other DHS subsets (set 7 vs. sets 1-6).

Fig. S7. Chromatin state analysis of static and dynamic male-biased DHS. A. Distribution of static and dynamic DHS in the defined classes of enhancer DHS, weak enhancer DHS, insulator DHS and promoter DHS, based on histone mark patterns [5]. **B.** Chromatin state distributions in male mouse liver of dynamic and static male-biased DHS, based on the 14 chromatin state model developed from the combination of six active and repressive histone marks and DHS, which segment the mouse genome into inactive, bivalent, enhancer-like, promoter-like, or transcribed-like states [6]. Chromatin state data are shown in Table S5. See Fig. 1C for emission probabilities for the six histone marks and DHS for each of the 14 chromatin states. Key results: Overall, 92-96% of dynamic and static male-biased DHS were classified as enhancers, with a larger fraction being weak enhancers [5] in the case of static male-biased DHS. Promoter DHS and insulator DHS comprised the balance of each male-biased DHS set (2-5% each) (**A**). Similarly, 86% of female-biased DHS and 90% of dynamic sex-independent DHS were classified as enhancers or weak enhancers, unlike static sex-independent DHS, where insulator and promoter DHS designations were much more common (15-20% each versus 3-8% for dynamic sex-independent DHS (**A**; also see Fig. 1A). We also did not observe major differences in chromatin state distributions between dynamic and static male-biased DHS based on a 14-state map developed for male and female mouse liver [6]. Thus, dynamic and static male-biased DHS both showed a high frequency (59-72%) of chromatin state E6, whose emission parameters indicate a high frequency of DHS and of the activating chromatin marks H3K27ac and H3K4me1 (**B**). Much smaller percentages of both male-biased DHS subsets were in other chromatin states, primarily enhancer states E5 and E11, promoter state E7, and bivalent state E12,

which is characterized by the presence of both activating marks (H3K4me1) and repressive marks (H2K27me3).

Fig. S8. Top enriched Gene Ontology (GO) terms identified by GREAT analysis of the predicted gene targets of each of the indicated 4 DHS sets. Mouse mm9 genomic coordinates for each DHS set were entered into the web-based tool GREAT (v.4.0.4) (http://great.stanford.edu/public/html/) and target genes identified under default conditions (Association rule: Basal+extension: 5000 bp upstream, 1000 bp downstream, 1 million bp max extension). Shown are the top enriched GO Biological Process terms for each DHS set using the whole genome background setting, with –log10 Binomial p-values shown. Complete data output from GREAT, including listings of gene targets and the DHS associated with each DHS are shown in Table S8.

References

[1] Y. Zhang, E.V. Laz, D.J. Waxman, Dynamic, sex-differential STAT5 and BCL6 binding to sex-biased, growth hormone-regulated genes in adult mouse liver, Mol Cell Biol, 32 (2012) 880-896.

[2] W. Ma, W.S. Noble, T.L. Bailey, Motif-based analysis of large nucleotide data sets using MEME-ChIP, Nat Protoc, 9 (2014) 1428-1450.

[3] T. Melia, D.J. Waxman, Sex-Biased lncRNAs Inversely Correlate With Sex-Opposite Gene Coexpression Networks in Diversity Outbred Mouse Liver, Endocrinology, 160 (2019) 989-1007.

[4] J. Connerney, D. Lau-Corona, A. Rampersaud, D.J. Waxman, Activation of Male Liver Chromatin Accessibility and STAT5-Dependent Gene Transcription by Plasma Growth Hormone Pulses, Endocrinology, 158 (2017) 1386-1405.

[5] B.J. Matthews, D.J. Waxman, Computational prediction of CTCF/cohesin-based intra-TAD loops that insulate chromatin contacts and gene expression in mouse liver, Elife, 7 (2018).

[6] A. Sugathan, D.J. Waxman, Genome-wide analysis of chromatin states reveals distinct mechanisms of sex-dependent gene regulation in male and female mouse liver, Mol Cell Biol, 33 (2013) 3594-3610.

C Comparison of chromatin accessibility differences between STAT5-bound (top row) and non-STAT5-bound DHS subsets (bottom row)

DNase-seq sample:
male STAT5-high (red)
male STAT5-low (blue)
female control (black)

Male biased dynamic DHS (834 sites)

Motif Logo	E-value for motif discovery (MEME/DREME)	Closest matching motif (Tomtom p-value)
	2.1e-30 (MEME)	motif72 (IRF2) (5.44e-04) motif45 (FOXI1) (2.00e-03) motif78 (FOXD1) (2.08e-03)
	4.1e-23 (MEME)	motif29 (STAT5B) (2.84e-06)
	3.7e-12 (MEME)	motif29 (STAT5B) (6.63e-03)
	4.3e-12 (MEME)	motif15 (Pax4) (1.05e-05) motif39 (MZF1_5-13) (5.14e-04) motif47 (SP1:SP3) (6.03e-04)
	2.2e-12 (MEME)	motif48 (SF1) (1.39e-03) motif68 (Hand1-Tcfe2) (4.38e-03)
	1.2e-15 (DREME)	motif29 (STAT5B) (1.20e-04) motif80 (c-Ets-2) (7.50e-03) motif79 (ELF5) (9.13e-03)

Sex independent dynamic DHS (1,532 sites)

Motif Logo	E-value for motif discovery (MEME/DREME)	Closest matching motif (Tomtom p-value)
	2.4e-32 (MEME)	motif29 (STAT5B) (2.76e-09) motif38 (DEAF1) (3.69e-03)
	3.5e-10 (MEME)	No motif match reported
	7.8e-10 (MEME)	motif45 (FOXI1) (8.73e-04) motif78 (FOXD1) (4.69e-03) motif88 (Elf-1) (6.39e-03)
	1.3e-12 (MEME)	motif2 (NR1H2-RXRA) (2.01e-06) motif71 (NR2F1) (3.10e-06)
	1.1e-27 (DREME)	motif29 (STAT5B) (1.52e-04)
	1.4e-14 (DREME)	motif45 (FOXI1) (3.34e-04) motif78 (FOXD1) (7.76e-03)

Male biased static DHS (1,895 sites)

Motif Logo	E-value for motif discovery (MEME/DREME)	Closest matching motif (Tomtom p-value)
	6.3e-21 (MEME)	motif92 (SP1) (1.28e-05) motif39 (MZF1_5-13) (1.72e-04) motif47 (SP1:SP3) (8.89e-04)
	7.2e-13 (MEME)	No motif match reported
	1.4e-12 (MEME)	motif2 (NR1H2-RXRA) (1.26e-06) motif71 (NR2F1) (3.25e-05) motif48 (SF1) (1.08e-03)
	2.7e-11 (MEME)	motif81 (MEIS1B:HOXA9) (8.98e-03)
	9.4e-28 (DREME)	motif19 (PBX1) (7.33e-04) motif24 (Clox) (1.04e-03) motif31 (CDP) (3.37e-03)
	3.9e-17 (DREME)	motif71 (NR2F1) (7.43e-06) motif2 (NR1H2-RXRA) (5.52e-05) motif48 (SF1) (8.75e-03)
	4.0e-12 (DREME)	motif45 (FOXI1) (9.20e-06) motif78 (FOXD1) (3.28e-03) motif97 (TEF-1) (5.15e-03)

Female biased static DHS (1,359 sites)		
Motif Logo	**E-value for motif discovery (MEME/DREME)**	**Closest matching motif (Tomtom p-value)**
	4.3e-17 (MEME)	motif39 (MZF1_5-13) (1.28e-03) motif15 (Pax4) (2.79e-03) motif92 (SP1) (5.47e-03)
	2.6e-10 (MEME)	No motif match reported
	2.8e-013 (MEME)	motif13 (TP53) (7.26e-03) motif48 (SF1) (7.26e-03) motif51 (NRSF) (7.27e-03)
	3.8e-017 (DREME)	No motif match reported

Sex independent static DHS (64,584 sites)

Motif Logo	E-value for motif discovery (MEME/DREME)	Closest matching motif (Tomtom p-value)
	1.8e-35 (MEME)	No motif match reported
	1.4e-12 (MEME)	motif78 (FOXD1) (4.22e-05) motif45 (FOXI1) (2.20e-03) motif72 (IRF2) (2.81e-03)
	3.3e-458 (DREME)	motif88 (Elf-1) (3.97e-05) motif79 (ELF5) (5.49e-05) motif80 (c-Ets-2) (3.61e-04)
	4.7e-286 (DREME)	motif78 (FOXD1) (3.27e-05) motif45 (FOXI1) (3.72e-04) motif76 (Lhx3) (7.75e-03)
	1.3e-281 (DREME)	motif48 (SF1) (2.15e-05) motif2 (NR1H2-RXRA) (2.90e-04) motif18 (RORA_1) (3.10e-04)
	2.5e-212 (DREME)	motif92 (SP1) (3.95e-06) motif39 (MZF1_5-13) (3.18e-03) motif87 (Egr) (6.29e-03)
	4.2e-172 (DREME)	motif62 (TLX1-NFIC) (1.24e-03) motif68 (Hand1-Tcfe2) (4.94e-03)
	1.0e-142 (DREME)	motif72 (IRF2) (3.46e-03) motif45 (FOXI1) (4.29e-03) motif78 (FOXD1) (7.79e-03)
	7.1e-108 (DREME)	motif17 (Myc) (1.09e-07) motif6 (MYC-MAX) (1.34e-07) motif42 (Myf) (2.73e-04)
	2.7e-104 (DREME)	motif34 (LUN-1) (6.09e-04) motif29 (STAT5B) (3.56e-03)

A. Hypophysectomy-response classes of sex-biased genes

Fig. S4

B. Pituitary hormone-dependent sex-biased genes

Class	Number of Genes (%)	Response to MHx	Response to FHx	Gene Subclass	Number of Genes
Male class I	202 (69)	Down	-	IA	110
		Down	Down	IB	89
		Down	Up	IC	3
Male class II	90 (31)	-	Up	IIA	50
		Up	Up	IIB	40
Female class I	134 (43)	-	Down	IA	90
		Down	Down	IB	35
		Up	Down	IC	9
Female class II	178 (57)	Up	-	IIA	108
		Up	Up	IIB	70

Comparison of chromatin accessibility changes between STAT5-bound (top row) and non-STAT5-bound DHS subsets (bottom row)

A

[MHX/Male Control]
DHS that close
(1,487 sites)

[MHX+GH 30 min/MHx]
DHS that open (1,475 sites)

[MHX+GH 90 min/MHx]
DHS that open (1,393 sites)

[MHX+GH 240
min/MHx]
DHS that open
(547 sites)

Note: Total sum of genes in the above Venn diagram = 1,974. In addition, 399 other DHS did not undergo chromatin closing in response to hypox (salmon oval) and lacked chromatin opening in any of the GH time-course data sets (green and blue ovals). The sum of the above 1,974 DHS plus these 399 other DHS = the full set of 2,373 DHS that are more open in STAT5-high than in STAT5-low livers (i.e., dynamic DHS).

B

1. DHS that open due to an endogenous GH pulse (2,373 sites)

3. DHS that close due to hypox and/or are opened by an exogenous GH pulse (1,974 sites: 83%)

2. DHS that do not close due to hypox and are not opened by an exogenous GH pulse (399 sites: 17%)

4. DHS that are closed by hypox but are not opened by an exogenous GH pulse (354 sites: 18%)

5. DHS opened by an exogenous GH pulse at one or more time points (1,620 sites: 82%)

6. DHS that not closed by hypox but are opened by an exogenous GH pulse (487 sites: 30%)

7. DHS that are closed by hypox and are opened by an exogenous GH pulse (1,133 sites: 70%)

C Relative magnitude of chromatin opening between STAT5 high and STAT5 low male livers

DHS sets 1-7 are defined in panel B

A Static and dynamic male-biased DHS

B Chromatin state distribution of male biased DHS

A. 834 dynamic male-biased DHS

B. 1895 static male-biased DHS

C. 1359 static female-biased DHS

D. 1532 dynamic sex-independent DHS