Improved Estimation of Bird Abundance and Energy Consumption by Combining Point and Transect Observations

Maria Jose Ayala Bolagay[1], Aaron Brothers[2], Marta Quitián[3], Eike Lena Neuschulz[4], Matthias Schleuning[4], Katrin Böhning-Gaese[4], Orlando Alvarez[1], Vinicio Santillán[1], and Christopher Thron[*2]

[1]Universidad Católica de Cuenca, Ecuador
[2]Texas A&M University-Central Texas, Killeen TX USA
[3]Instituto Mediterráneo de Estudios Avanzados (CSIC-UIB)
[4]Senckenberg Biodiversity and Climate Research Centre, Frankfurt Germany

April 16, 2023

Abstract

Point counts and transect surveys are both common methods for estimating bird abundance (by species, genus, or family) in a given environment. Sometimes both methods are used in the same study. Point counts reflect abundance directly, while transect observations of bird feeding instances reflect both the bird abundance and the mean energy consumption rate per bird.

In this paper we exploit the relationship between these two types of observations to draw inferences about bird energy consumption; to develop an improved estimator of bird abundance; and to derive error bars for abundance estimates. The mathematical methods developed are applied to a dataset consisting of point counts and transect surveys of Ecuadorian frugivore bird species in six different types of environment (three elevations and two degrees of forestation). By comparing transect-based and point count-based estimates of energy consumption across species, we show that mean fruit weight consumed does not have a significant effect on number of feeding instances. We also derive a procedure for forming an optimal linear combination of two relative abundance estimators based on point counts and transect observations respectively. The improvement in accuracy achieved by this combined estimator is verified by simulations applied to bird family data from the above-mentioned survey. Finally, we

[*]Corresponding author: thron@tamuct.e du

use simulations to obtain 90% confidence intervals for the bird family relative abundance estimates. These confidence intervals were also verified using simulations.

Keywords: Energy consumption, Abundance, Estimation, Survey, Point counts, Transect sampling, Bird, Ecosystem, Frugivore, Species, Family, Simulation

1   Introduction

Two common methods for estimating abundance of bird types (whether species, genera, or families) in a given area are point counts and transect surveys. Point counts are obtained when the observer chooses a set of fixed observation points within the area of interest, and counts bird detections (by sight or by sound) during multiple sessions over an extended period of several days or months. Transect surveys are conducted by the observer specifying a transect (i.e. narrow section) within the area and repeatedly walking through and recording bird observations at the same time of day over an extended period. Transect measurements can target bird feedings, so that the frequency of feeding on each different type of plant is recorded. These census methods have been extensively discussed in the literature, from several points of view [6], [7].

The energy consumption of different bird types can be expressed in two different ways. The total energy consumption of a bird type can be expressed as the number of individuals of that type times the mean consumption per individual. On the other hand, the same quantity may be expressed as the total number of feedings by birds of that type times the amount consumed per feeding. Point counts can be used to estimate bird abundance required for the first expression, while transect surveys can provide feeding data for the second expression.

We may make use of these two alternative estimates of energy consumption in several different ways. First, by comparing the two estimates, we may gain information about bird consumption rates. Second, we may combine the two estimates to obtain a more accurate estimate of energy consumption per bird type, and hence of bird type relative abundance. Finally, given that count measurements are Poisson distributed (which is commonly recognized as valid for count measurements), we can use the results obtained and simulations to obtain confidence intervals for the relative abundance of different bird types.

The methodologies that we have developed may be applied to other surveys where multiple forms of data related to species abundance are present. Our results can be used both to obtain more accurate abundance estimates, and to rigorously specify accuracies of the estimates thus obtained. We have also demonstrated a technique for estimating an important biological parameter, namely the exponent in the functional dependence of bird energy consumption on bird weight. Using this technique, other datasets may provide similar estimates.

The remainder of this paper is organized as follows. Section 2 consists of two parts. In the first part, we present the theoretical derivations for the statistical methods used for the data analysis. These methods include the construction of point count-based and transect observation-based estimators for energy consumption and for bird abundance; and the combination of bird abundance estimators and log abundance estimators into more accurate estimators. In the second part, the methods used to evaluate the theoretical results are described. These methods include the use of an empirical dataset obtained from several regions in southern Ecuador; this dataset is used both to estimate model parameters from bird species data, and as a basis for simulations to test the accuracy of the different theoretical estimators using bird family data. In Section 3 the results of the evaluation are presented; all figures displaying these results are placed at the end of the paper, after the bibliography. Finally, in Section 4 our conclusions from this study are summarized.

2 Methodology

This section consists of two subsections dealing with theoretical and empirical methods, respectively. The first section derives statistical estimators that make use of point counts and transect observations to estimate model parameters and demographics. The second section describes the application of these theoretical results to a specific ecological survey.

2.1 Theoretical equations

2.1.1 Transect observation-based expression for energy consumption per bird type

The energy consumption per bird type over a given period of time at a given location may be expressed as the total number of feedings times the mean energy transfer per feeding. We may rewrite this statement in the form of an equation:

$$\mathcal{E}_n = \sum_{m=1}^{N} \mathcal{F}_{nm} \mathcal{T}_{nm} \qquad (n = 1 \ldots N, m = 1 \ldots M), \tag{1}$$

where

- M, N: Number of fruit and bird types, respectively.

- $\mathcal{E}_n$: Total energy consumption of all birds of type n at the specified location during the specified period.

- $\mathcal{F}_{nm}$: Total number of feedings of bird type n on fruit type m; $\mathcal{T}_{nm}$: Mean energy transfer per feeding for bird type n on fruit type m.

Under the assumption that the transect observations are representative and capture the same proportion of all bird-fruit interactions, we have:

$$X_{nm} = K_{\mathcal{F}} \mathcal{F}_{nm} - u_{nm} \qquad (n = 1 \ldots N, m = 1 \ldots M), \tag{2}$$

where

- $K_\mathcal{F}$: Constant of proportionality, assumed to be independent of bird and fruit types;

- X_{nm}: Number of transect observations of bird type n consuming fruit type m;

- u_{nm}: Observational uncertainty for bird type n consuming fruit type m.

We furthermore assume initially that energy transfer per feeding depends on both the mean fruit weight and the mean bird weight. This assumption presumes that larger fruit may require less foraging, and larger birds may be able to consume more per feeding. This relationship is summarized in the equation:

$$\mathcal{T}_{mn} = K_\mathcal{T} f_m^{p_f} w_n^{p_{b1}} \qquad (n = 1 \dots N), \tag{3}$$

where

- $K_\mathcal{T}$: Constant of proportionality, assumed to be independent of bird type.

- f_m: mean fruit weight for type m.

- w_n: Mean weight of bird type n;

- p_f, p_{b1}: Exponents of fruit weight and bird weight respectively, in power model.

Substituting $\mathcal{F}_{nm}$ and $\mathcal{T}_{nm}$ in (1) with their expressions from (2) and (3) respectively, we have finally:

$$\mathcal{E}_n = \kappa_1 \sum_{m=1}^{M} (X_{nm} + u_{nm}) f_m^{p_f} w_n^{p_{b1}} \qquad (n = 1 \dots N), \tag{4}$$

where $\kappa_1 \equiv (K_\mathcal{F} K_\mathcal{T})^{-1}$ is a constant of proportionality that is independent of bird and fruit types. In order to simplify the model and make it amenable to analysis, we replace f_m in (4) with $\bar{f}_n$, which is defined as the mean fruit weight consumed by bird species n. This simplification will be accurate when each bird species mostly consumes fruit species with similar average weights. An additional simplification concerns the magnitude of the measurement errors u_{nm}. We assume that these errors are nearly constant in percentage across species: that is to say, the error for species n will be roughly proportional to the number of observations X_n.

With the two simplifications described above, the final simplified model is given by the equation:

$$\mathcal{E}_n = \kappa_1 X_n (1 + u_n)(\bar{f}_n)^{p_f} (w_n)^{p_{b1}}, \tag{5}$$

where

- X_n: Sum of transect observations for bird type n. The formula for X_n is $X_n = \sum_{m=1}^{M} X_{nm}, n = 1 \dots N$.

- u_n: Combined proportionate observational uncertainty for bird type n. The variables $\{u_n\}, n = 1, \dots N$ are modeled as independent, identically distributed random variables with mean 0 and variance $\sigma_{X,n}^2$;

- $\bar{f}_n$: Average weight of fruit consumed by bird type n, given by the formula:

$$\bar{f}_n \equiv \frac{\sum_{m=1}^{M} X_{nm} f_m}{\sum_{m=1}^{M} X_{nm}}. \tag{6}$$

2.1.2 Point count-based estimate of energy consumption per bird type

A second expression for a bird type's energy consumption at a given location during a prescribed period is the product of the number of birds times the mean energy intake per bird, as expressed by the following formula:

$$\mathcal{E}_n = \mathcal{N}_n \mathcal{M}_n \qquad n = 1 \dots N, \tag{7}$$

where $\mathcal{N}_n$ and $\mathcal{M}_n$ represent respectively the total number and mean energy consumption of birds of type n.

As with transect observations, we assume that point counts are representative samples across all bird types, so we have:

$$Z_n = K_{\mathcal{N}} \mathcal{N}_n - \tilde{v}_n \qquad n = 1 \dots N, \tag{8}$$

where

- $K_{\mathcal{N}}$: Constant of proportionality, assumed to be independent of bird types;

- Z_n: Number of point count observations of bird type n, $n = 1 \dots N$;

- $\tilde{v}_n$: Uncertainty in point counts for bird type n.

As with transect observations, we assume that the uncertainty is proportional to the number of observations:

$$\tilde{v}_n = Z_n v_n \qquad n = 1 \dots N, \tag{9}$$

where $\{v_n, n = 1 \dots N\}$ are independent, identically distributed random variables with mean 0 and variance $\sigma_{Z,n}^2$.

In the literature, bird type mean energy consumption is modeled as proportional to mean bird weight raised to a power as follows [3]:

$$\mathcal{M}_n \equiv K_{\mathcal{M}} w_n^{p_{b2}}, \tag{10}$$

where

- $K_{\mathcal{M}}$: Constant of proportionality, independent of bird type n;

- w_n: Mean weight of bird type n, $n = 1, \dots N$;

- p_{b2}: Weight exponent in energy model.

By substituting (8)-(10) into (7) we obtain finally:

$$\mathcal{E}_n = \kappa_2 Z_n (1 + v_n)(w_n)^{p_{b2}}, \tag{11}$$

where $\kappa_2 \equiv K_{\mathcal{M}}/K_{\mathcal{N}}$ is independent of n.

2.1.3 Parameter estimation based on energy consumption equations

In practice, equations (5) and (11) developed in the previous two sections cannot be used to estimate total energy consumption of bird types, because the constants κ_1 and κ_2 are impossible to obtain. However, by setting the two expressions equal to each other obtain a linear relationship that can be used to estimate the model exponents p_{b1}, p_{b2} and p_f based on point count and transect data. Equating $\mathcal{E}_n$ in (5) and (11) gives:

$$\kappa_1 X_n (1 + u_n)(\bar{f}_n)^{p_f}(w_n)^{p_{b1}} = \kappa_2 Z_n (1 + v_n)(w_n)^{p_{b2}}. \tag{12}$$

Taking logarithms and rearranging, we find:

$$\log(X_n) = \log(Z_n) + C_0 + p_b \log(w_n) - p_f \log(\bar{f}_n) + e_n, \tag{13}$$

where:

$$
\begin{aligned}
C_0 &\equiv \log\left(\frac{\kappa_2}{\kappa_1}\right); \\
p_b &\equiv p_{b2} - p_{b1} \\
e_n &\equiv \log\left(\frac{1 + u_n}{1 + v_n}\right) \text{(combined error term)}
\end{aligned}
\tag{14}
$$

Note that if $u_n, v_n \ll 1$ then $e_n \approx u_n - v_n$. We will assume that $\{e_n, n = 1, \dots N\}$ are independent, identically distributed random errors. Then if we have transect data $\{X_1, \dots, X_N\}$ and point count data $\{Z_1, \dots, Z_M N\}$, we may use (13) to perform ordinary unweighted least squares regression and find values for C_0, p_b, and p_f that give the best fit to the data.

2.1.4 Estimates of relative abundance

In this section we make use of the results from previous sections to develop a pair of estimates for relative abundances of bird types, where the relative abundance a_n of species n is defined as type abundance divided by total bird abundance:

$$a_n = \frac{\mathcal{N}_n}{\sum_{k=1}^{N} \mathcal{N}_k}. \tag{15}$$

The two estimates are based on point counts and transect observations, respectively. We also show that in the case where the estimators are unbiased and uncorrelated, a linear combination of the two estimators can be formed that is unbiased and has smaller variance than either of the two individual estimators.

First we develop an estimator for relative abundance based on point counts. From (8) and (9) and the definition (15) we have:

$$
\begin{aligned}
a_n &= \frac{Z_n(1 + v_n)}{\sum_{k=1}^{N} Z_k(1 + v_k)} \\
&= \frac{Z_n}{\sum_{k=1}^{N} Z_k}(1 + v_n) \cdot \left(1 + \frac{\sum_{k=1}^{N} v_k Z_k}{\sum_{k=1}^{N} Z_k}\right)^{-1}.
\end{aligned}
\tag{16}
$$

The final multiplicative term in (16) is independent of n, and involves the sum of independent random errors, which will tend to cancel each other. We may therefore approximate this term as:

$$
\left(1 + \frac{\sum_{k=1}^{N} v_k Z_k}{\sum_{k=1}^{N} Z_k}\right)^{-1} \approx (1 - \nu)^{-1} \approx 1 + \nu,
\tag{17}
$$

where $\nu \ll 1$ is a mean-zero random variable. This gives finally

$$
\begin{aligned}
a_n &\approx \frac{Z_n}{\sum_{k=1}^{N} Z_k}(1 + v_n)(1 + \nu) \\
&\approx \frac{Z_n}{\sum_{k=1}^{N} Z_k}(1 + \nu + v_n'),
\end{aligned}
\tag{18}
$$

where $\{v_n' = v_n(1 + \nu), n = 1, \ldots N\}$ are statistically independent errors. It follows that an estimator $\widehat{a}_n$ for a_n that is based on point counts $\{Z_n\}$ can be obtained from (18) with the random error terms removed:

$$
(\widehat{a}_n)_Z \equiv \frac{Z_n}{\sum_{k=1}^{N} Z_k}.
\tag{19}
$$

Next, an alternative expression for a_n based on transect observations can be obtained. Combining (8), (9) and rearranging,, we have:

$$
Z_n(1 + v_n) \equiv K_{\mathcal{N}} \mathcal{N}_n \qquad n = 1 \ldots N,
\tag{20}
$$

and substituting into (12) and rearranging gives:

$$
\mathcal{N}_n = \frac{\kappa_1}{\kappa_2 K_{\mathcal{N}}} X_n(1 + u_n)(\bar{f}_n)^{p_{f,n}}(w_n)^{-p_{b,n}}.
\tag{21}
$$

(Note that we are generalizing (12) to the case where the fruit weight and bird weight exponents $p_{f,n}$ and $p_{b,n}$ depend on bird type.) This enables us to give an expression for relative abundance in terms of X_n instead of Z_n:

$$
a_n = \frac{\mathcal{N}_n}{\sum_{k=1}^{N} \mathcal{N}_k} = \frac{X_n(1 + u_n)(\bar{f}_n)^{p_{f,n}}(w_n)^{-p_{b,n}}}{\sum_{k=1}^{N} X_k(1 + u_k)(\bar{f}_k)^{p_{f,n}}(w_k)^{-p_{b,n}}}.
\tag{22}
$$

By a similar argument as before, we obtain the following X_n-based estimator for relative abundance a_n:

$$(\widehat{a}_n)_X \equiv \frac{(\bar{f}_n)^{\widehat{p}_{f,n}}(w_n)^{-\widehat{p}_{b,n}}X_n}{\sum_k (\bar{f}_k)^{\widehat{p}_{f,n}}(b_k)^{-\widehat{p}_{b,n}}X_k} \tag{23}$$

where $\widehat{p}_{f,n}, \widehat{p}_{b,n}$ are estimates for the fruit and bird weight coefficients $p_{f,n}$ and $p_{b,n}$, respectively.

2.1.5 Minimum variance combination of relative abundance estimators

Now consider the linear combination of estimators:

$$(\widehat{a}_n)_{\beta_n,\gamma_n} \equiv \beta_n(\widehat{a}_n)_Z + \gamma_n(\widehat{a}_n)_X, \tag{24}$$

where $\{\beta_k, \gamma_k, k = 1 \ldots N\}$ are positive coefficients. It follows by the linearity of distribution mean that:

$$\text{mean}\left[(\widehat{a}_n)_{\beta_n,\gamma_n}\right] = \beta_n \text{mean}\left[(\widehat{a}_n)_Z\right] + \gamma_n \text{mean}\left[(\widehat{a}_n)_X\right]; \tag{25}$$

In the case where the estimators $(\widehat{a}_n)_Z$ and $(\widehat{a}_n)_X$ are unbiased, we have:

$$\text{mean}\left[(\widehat{a}_n)_Z\right] = \text{mean}\left[(\widehat{a}_n)_X\right] = a_n; \tag{26}$$

The requirement that $(\widehat{a}_n)_{\beta_n,\gamma_n}$ be an unbiased estimator of a_n is expressed mathematically as:

$$\text{mean}\left[(\widehat{a}_n)_{\beta_n,\gamma_n}\right] = a_n. \tag{27}$$

Combining (25), (26), and (27) gives the requirement:

$$\gamma_n = 1 - \beta_n. \tag{28}$$

Given that $(\widehat{a}_n)_Z$ and $(\widehat{a}_n)_X$ are based on independent observations and are thus uncorrelated, it follows:

$$\text{Var}\left[(\widehat{a}_n)_{\beta_n,\gamma_n}\right] = \beta_n^2 \text{Var}\left[(\widehat{a}_n)_Z\right] + \gamma_n^2 \text{Var}\left[(\widehat{a}_n)_X\right], \tag{29}$$

and a standard calculation shows that the unbiased minimum-variance estimators for a_n is obtained by choosing:

$$\gamma_n \equiv (1 - \beta_n); \quad \beta_n \equiv \frac{\text{Var}[(\widehat{a}_n)_X]}{\text{Var}[(\widehat{a}_n)_Z] + \text{Var}[(\widehat{a}_n)_X]}, \tag{30}$$

which has variance

$$\min_{\beta_n} \text{Var}\left[(\widehat{a}_n)_{\beta_n,1-\beta_n}\right] = \frac{\text{Var}[(\widehat{a}_n)_X]\text{Var}[(\widehat{a}_n)_Z]}{\text{Var}[(\widehat{a}_n)_Z] + \text{Var}[(\widehat{a}_n)_X]}. \tag{31}$$

2.1.6 Combined estimates of log relative abundance

Sometimes it is more advantageous to consider the logarithms of relative abundances of bird species, genera, or families, instead of relative abundances themselves. Typically the logarithms have a more symmetrical distribution which is closer to normal. By taking natural logarithms of $(\widehat{a}_n)_Z$ and $(\widehat{a}_n)_X$, we obtain estimators for the logarithm of relative abundance, which we denote as $(\widehat{\ell}_n)_Z$ and $(\widehat{\ell}_n)_X$ respectively:

$$
(\widehat{\ell}_n)_Z \equiv \log\left((\widehat{a}_n)_Z\right) = \log Z_n - \log\left(\sum_k Z_k\right);
$$

$$
(\widehat{\ell}_n)_X \equiv \log\left((\widehat{a}_n)_X\right)
$$

$$
= \log X_n - \widehat{p}_{b,n} \log w_n - \widehat{p}_{f,n} \log f_n - \log\left(\sum_k (f_k)^{-\widehat{p}_{f,n}}(b_k)^{-\widehat{p}_{b,n}} X_k\right).
\tag{32}
$$

As with the relative abundance estimators, we may form linear combinations of the log relative abundance estimators:

$$
(\widehat{\ell}_n)_{\widetilde{\beta}_n, \widetilde{\gamma}_n} \equiv \widetilde{\beta}_n (\widehat{\ell}_n)_Z + \widetilde{\gamma}_n (\widehat{\ell}_n)_X,
\tag{33}
$$

where $\{\widetilde{\beta}_k, \widetilde{\gamma}_k, k = 1 \ldots N\}$ are positive coefficients. In the case where $(\widehat{\ell}_n)_Z$ and $(\widehat{\ell}_n)_X$ are unbiased and uncorrelated, it follows as before that:

$$
\mathrm{Var}\left[(\widehat{\ell}_n)_{\widetilde{\beta}_n, \widetilde{\gamma}_n}\right] = (\widetilde{\beta}_n)^2 \mathrm{Var}\left[(\widehat{\ell}_n)_Z\right] + (\widetilde{\gamma}_n)^2 \mathrm{Var}\left[(\widehat{\ell}_n)_X\right],
\tag{34}
$$

and an unbiased minimum-variance estimator for ℓ_n is obtained by choosing:

$$
\widetilde{\gamma}_n \equiv (1 - \widetilde{\beta}_n); \quad \widetilde{\beta}_n \equiv \frac{\mathrm{Var}[(\widehat{\ell}_n)_X]}{\mathrm{Var}[(\widehat{\ell}_n)_Z] + \mathrm{Var}[(\widehat{\ell}_n)_X]}.
\tag{35}
$$

2.1.7 Calculation of coefficients for combined estimators

The combined estimators of relative abundance and log relative abundance defined in Sections 2.1.5 and 2.1.6 respectively require the specification of coefficients $\{\beta_n, \beta'_n, n = 1 \ldots N\}$. These coefficients are not determined a priori, but rather must be supplied by the user. In this section, we describe a simulation-based procedure for determining coefficients which will give estimators that have reduced variance.

The proposed estimation procedure for β_n is based on a bootstrapping apporach, and runs as follows. Suppose the user has recorded point counts $z'_1, \ldots z'_N$ and transect observations $x'_1, \ldots x'_N$ for N different species.

1. Compute abundance estimates based on $\{z'_1 \ldots z'_N\}$, according to (19):

$$
(\widehat{a}_n)_{z'} \equiv \frac{z'_n}{\sum_{k=1}^N z'_k}.
\tag{36}
$$

2. Generate M simulated sets of point counts and transect observations $(z''_{1m} \ldots z''_{Nm}, x''_{1m} \ldots x''_{Nm}), m = 1 \ldots M$ based on the user's observed data. Simulation values are independently generated according to the following distributions:

$$z''_{nm} \sim Pois(z'_n); \qquad x''_{nm} \sim Pois \left(\frac{z'_n (\bar{f}_n)^{p_{f,n,m}} w_n^{p_{b,n,m}}}{\sum_{k=1}^{N} z'_k (\bar{f}_k)^{p_{f,n,m}} w_j^{p_{b,n,m}}} \sum_{k=1}^{N} x'_k \right), \tag{37}$$

where $Pois(\lambda)$ denotes a Poisson random variable with mean event rate λ; and $p'_{f,n,m}$ and $p'_{b,n,m}$ are randomly generated according to probability distributions reflecting the user's knowledge of fruit and weight exponents in the model (3). (Note that Poisson distributions are commonly used to model survey counts [2].) This expression generates simulated point counts with expected values equal to the observed point counts $z'_1 \ldots z'_N$, and simulated transect observations whose expected values yield energy consumption rates that agree with those calculated from the point counts, and whose sum agrees with the sum of the user's observed transect observations.

3. For each of the M simulated sets of observations, the user calculates abundance estimators according to (19):

$$(\widehat{a}_{nm})_{z''} \equiv \frac{z''_{nm}}{\sum_k z''_{km}}; \qquad (\widehat{a}_{nm})_{x''} \equiv \frac{(\bar{f}_n)^{\bar{p}_{f,n}} (w_n)^{-\bar{p}_{b,n}} x''_{nm}}{\sum_k (\bar{f}_k)^{\bar{p}_{f,n}} (b_k)^{-\bar{p}_{b,n}} x''_{km}}. \tag{38}$$

4. The user calculates values of $\beta_n, n = 1 \ldots N$ that minimize root mean squared error (RMSE) of the combined estimator $(\widehat{a}_n)_\beta \equiv \beta_n (\widehat{a}_n)_Z + (1 - \beta_n)(\widehat{a}_n)_X$:

$$\beta_n = \mathrm{argmin}_\beta \sqrt{\frac{1}{N} \sum_{m=1}^{M} \left(\beta(\widehat{a}_{nm})_{z''} + (1-\beta)(\widehat{a}_{nm})_{x''} - (\widehat{a}_n)_{z'} \right)^2}. \tag{39}$$

The minimized RMSE values in (39) can also be used to estimate confidence intervals for $\widehat{a}_n)_\beta$, since the RMSE value should approximate one standard deviation. So if the RMSE value obtained is denoted as $\Delta_{\beta,n}$, the $(\widehat{a}_n)_\beta \pm 1.65\Delta_{\beta,n}$ should give a 90% confidence interval for the true value of a_n.

A similar procedure can be to determine values of $\widetilde{\beta}_n, n = 1 \ldots N$ and confidence intervals for an optimal linear combination of log abundance estimators, defined as $(\widehat{\ell}_n)_{\widetilde{\beta}} \equiv \widetilde{\beta}_n (\widehat{\ell}_n)_Z + (1 - \widetilde{\beta}_n)(\widehat{\ell}_n)_X$.

2.2 Empirical evaluation and verification of theoretical equations

In this section, we describe how the equations derived in Section 2.1 are evaluated and verified using data from an actual survey of frugivore birds conducted

Location	Approx. altitude (m)	Environment type
Bombuscaro	1000	Natural
Copalinga	1000	Fragmented
ECSF	2000	Natural
Finca	2000	Fragmented
Cajanuma	3000	Natural
Bellavista	3000	Fragmented

Table 1: Summary of characteristics for six locations studied

in several locations in Ecuador. Both the data collected and the verification methods applied are described.

2.2.1 Dataset used in methodology evaluation

Data was collected from 18 1-hectare plots spread across three elevations (1000 m above sea level (asl) (970–1092 $\pm$ 47.2 m), 4° 6′ S, 78° 58′ W; 2000 m asl (2025–2119 $\pm$ 32.4 m), 3° 58′ S, 79° 4′ W; and 3000 m asl (2874–2966 $\pm$ 32.9 m], 4° 6′ S, 79° 10′ W. These plots represented two types of habitat (natural forest and fragmented forest), within and around Podocarpus National Park (PNP) and San Francisco reserve (BRSF) on the southeastern slope of the Andes in Ecuador. The six sites are characterized in Table 1. Study plots were selected jointly by the German Platform for Biodiversity and Ecosystem Monitoring and Research in South Ecuador, to ensure typical representation of each forest type. The area exhibits a humid tropical montane climate [5] with a bimodal rainfall pattern (highest precipitation season from May through June, lowest precipitation season from October through November) [4].

To assess bird diversity and abundance, nine observation points were established within each 1-hectare plot. For a duration of 10 minutes at each point, we recorded and identified all bird species heard or seen within a 20-meter radius [1]. This sampling was repeated eight times per plot over a span of two years (1/6 hours $\times$ 9 observation points $\times$ 8 replicates $\times$ 18 plots = 216 total sampling hours). The total number of bird species (species richness) and the overall number of individual birds (abundance) were quantified by aggregating the records from all observation points and replicates per plot. Bird species were classified as frugivores if their diet consisted of 40% or more fruit [8].

2.2.2 Parameter estimation using energy models

Best fits for the parameters p_f and p_b were found by applying linear regression to (13), for several different datasets. Fits were found for species data from each of the six regions separately; for species data grouped by altitude; for species data grouped by environment type (natural versus fragmented); and for species data for all six regions combined. For each fit, scatter plots were made for

actual versus predicted transect observations, based on (13). R^2 and Akaike information criterion (AIC) values and 95 percent confidence intervals for p_f and p_b were also computed. Scatter plots, R^2 and AIC values and 95 percent confidence intervals for p_b were also generated for the same datasets, but with a simplified single-parameter model that assumes $p_f = 0$ in (13). The 2-parameter and 1-parameter models are compared for goodness of fit. The purpose is to determine whether p_f has a significant effect on mean energy consumption per feeding.

2.2.3 Performance evaluation of abundance estimation using empirically-based simulations

Statistical characteristics of the abundance and log abundance estimators defined in Sections 2.1.4 and 2.1.6 were evaluated using simulations as follows. In this evaluation, the purpose was not to estimate true relative abundances of the regions from which the dataset was taken–since we don't know these true relative abundances, the accuracy of such estimates cannot be ascertained. Rather, the regional data is used to supply realistic scenarios; the regional data from the six regions is used as "ground truth" for six hypothetical regions, and simulations are randomly generated based on the data, and used to predict this "ground truth". In this way we can evaluate the accuracy of estimators using simulations with realistic characteristics. We base simulations on per-family data instead of per-species data, because per-species point counts are too small to establish differences in accuracies between estimators.

Testing for estimator accuracy and precision

Simulations were conducted to determine the accuracy and precision of point count-based, transect observation-based, and combined estimators as follows.

10000 simulated scenarios were generated based on the bird family data from each region. The data from the six actual regions was used to generate "ground truth" for six hypothetical regions as follows. If $PC_1, PC_2, \ldots PC_N$ are the observed point counts in a given actual region, then the true relative abundance of family n in the corresponding hypothetical region is given by:

$$a_n = \frac{PC_n}{\sum_{j=1}^{N} PC_j}. \tag{40}$$

Next, we simulated 10,000 surveys in each hypothetical region, where each survey includes point counts and transect observations. For a given simulation, the point count for each bird family n is generated by a Poisson random variable whose mean is the point count from the observed data. We denote these Poisson random variables as $z_n, n = 1, \ldots N$, where

$$z_n \sim Pois(PC_n). \tag{41}$$

Transect observations for each simulation are randomly generated as follows. First, for each bird family n we generate a weight exponent $p_{b,n}$ as a Gaussian

random variable with 95% confidence interval [-0.6,-0.1]:

$$p_{b,n} \sim Gauss(\mu = -0.35, \sigma^2 = 0.125^2). \tag{42}$$

The choice of distribution (42) will be motivated in Section 3.1 (see the final line of Table 3): this represents the uncertainty in the parameter p_b which is estimated as described in Section 2.2.2. The transect observation for birds in family n (denoted as x_n) is then simulated as follows:

$$x_n \sim Pois\left(PC_n w_n^{p_{b,n}} \frac{\sum_{j=1}^N TO_j}{\sum_{j=1}^N PC_j w_j^{p_{b,j}}}\right), \tag{43}$$

where TO_j denotes the number of transect observations of bird family j in the original data ($j = 1 \ldots N$). The choice of Poission parameter in (43) was made so that the expected sum of simulated observations over bird families at a given region is equal to the sum of transect observations in the original data. The Poisson parameter also reflects a 1-parameter model of energy consumption that presumes $p_f = 0$ in (4). This choice will be justified by the results in Section 3.1, which show there is no statistical evidence of energy consumption's dependence on mean fruit weight.

Given these simulated point counts $\{z_n\}$ and transect observations $\{x_n\}$, We then compute estimates $(\widehat{a}_n)_z$ and $(\widehat{a}_n)_x$ for a_n based on $\{z_n\}$ and $\{x_n\}$ respectively according to (19) and (23):

$$(\widehat{a}_n)_z \equiv \frac{z_n}{\sum_k z_k}; \quad (\widehat{a}_n)_x \equiv \frac{(w_n)^{-\widehat{p}_{b,n}} x_n}{\sum_k (b_k)^{-\widehat{p}_{b,n}} x_k}, \tag{44}$$

where $\widehat{p}_{b,n}$ is the investigator's best determination of the fruit weight exponent $p_{b,n}$. Two cases were considered: (a) the investigator does not have detailed knowledge of the fruit weight exponent, and sets $\widehat{p}_{b,n}$ equal to the mean value $\bar{p}_b$ of the distribution in (42); and (b) the investigator knows exactly the values of $p_{b,n}$ for each n, and uses $\widehat{p}_{b,n} = p_{b,n}$.

Besides the estimators $\widehat{a}_n)_z$ and $\widehat{a}_n)_x$, we also computed the combined estimate $(\widehat{a}_n)_\beta = \beta(\widehat{a}_n)_z + (1 - \beta)(\widehat{a}_n)_x$ as a function of β for $0 \le \beta \le 1$ for each simulation. Taking these results over all simulations, we were able to compute the means and variances of $\widehat{a}_n)_z$, $\widehat{a}_n)_x$, and $\widehat{a}_n)_\beta$. This enables us to compute the accuracy and precision of all estimators for each family in each of the six hypothetical regions, for the two cases (a) and (b) definedin the previous paragraph.

A similar procedure was used to gauge the accuracy and precision of log abundance estimators $(\widehat{\ell}_n)_z, (\widehat{\ell}_n)_x$, and $(\widehat{\ell}_n)_\beta$ defined as follows:

$$(\widehat{\ell}_n)_z \equiv \log(\widehat{a}_n)_z; \qquad (\widehat{\ell}_n)_x \equiv \log(\widehat{a}_n)_x; (\widehat{\ell}_n)_\beta \equiv \beta(\widehat{\ell}_n)_z + (1 - \beta)(\widehat{\ell}_n)_x. \tag{45}$$

Combined estimator parameter determination and confidence interval estimation

The second set of simulations is intended to test the procedure for determining variance-reducing linear combinations of estimators described in Section 2.1.7.

- First, for each of the six different regions observed in the study, 2500 simulated scenarios for point count and transect observation values $\{z_1 \ldots z_n, x_1, \ldots x_n\}_m$, $m = 1 \ldots 2500$ were generated according to (41). Each observation vector $z_1 \ldots z_N$ was to create a relative abundance vectors $a_n = z_n / \sum_{k=1}^{n} z_k$, $n = 1 \ldots N$. These relative abundances are taken as "ground truth" for the scenario (these are generated so as to increase the variety in the "ground truth" scenarios, making the test more robust.)

- Next, for each scenario, point counts and transect observations (simulating the dataset collected by a single hypothetical "ecologist" studying the scenario) were randomly generated as z'_n and x'_n respectively according to (compare (41)-(43))

$$
\begin{aligned}
z'_n &\sim Pois(z_n), n = 1 \ldots N; \\
p_{b,n} &\sim Gauss(\mu = -0.35, \sigma^2 = 0.125^2), n = 1 \ldots N; \\
x'_n &\sim Pois\left(z_n w_n^{p_{b,n}} \frac{\sum_{k=1}^{N} x_k}{\sum_{k=1}^{N} z_k w_k^{p_{b,k}}} \right), n = 1 \ldots N.
\end{aligned}
\tag{46}
$$

- Each simulated ecologist uses her own point counts and transect observations to compute estimators $(\widehat{a}_n)_{z'}, (\widehat{a}_n)_{x'}, (\widehat{\ell}_n)_{z'}$ and $(\widehat{\ell}_n)_{x'}$ according to (38). Two alternative situations are modeled: (a) the ecologist does not have detailed knowledge of the coefficients $p_{b,n}$, and uses the mean value $\bar{p}_b = 0.35$ to estimate $(\widehat{a}_n)_{x'}$ and $(\widehat{\ell}_n)_{x'}$; (b) the ecologist has precise knowledge of $p_{b,n}$, and uses the same values that are used in (46) for these estimates.

- Each simulated ecologist then uses the prescription in Section 2.1.7 with $M = 2500$ to estimate the linear combination parameters β and $\widetilde{\beta}$ for the estimators $(\widehat{a}_n)_\beta$ and $(\widehat{\ell}_n)_\beta$, as well as 90% confidence intervals for these estimates. As in the previous step, two scenarios apply: (a) the ecologist uses the mean value $\bar{p}_b = 0.35$ to estimate $(\widehat{a}_n)_\beta$ and $(\widehat{\ell}_n)_\beta$; (b) the ecologist uses the same values of $p_{b,n}$, that are used in (46) for these estimates.

To compare the accuracies of the point count-based, transect-based, and combined estimators, 95% confidence intervals for the relative error in relative abundance (estimate − ground truth relative abundance)/(relative abundance) are plotted. In order to verify the confidence intervals computed in the last bullet above, we also recorded the proportion of time that the ground truth relative abundances were within the 90% confidence intervals.

3   Results

3.1   Application of energy consumption estimation methodology to dataset

The six panels in Figure 1 contain scatter plots showing actual transect observations of bird species in log scale (y axis) versus transect observations predicted by the model (13) in log scale, for each of the six locations studied. The x and y axes in these plots represent the right and left sides of (13) respectively, and the parameters and constant C_0 are obtained by unweighted linear regression. An exact fit to the theoretical model corresponds to a line of slope 1. We may visually appraise goodness of fit by seeing how close the data points lie to this theoretical line. The dots' colors represent the mean weight of the corresponding bird species. For most regions there does seem to be a generally positive trend, albeit with numerous errant values particularly on the low side. There is no discernible influence of bird weight on the accuracy of the predictions.

Figures 2, 3, and 4 similarly show predicted versus actual transect observations grouped by elevation, fragmentation, and all data respectively. These figures show the same general positive trend and loose fits as were seen in the plots for individual regions.

Results of the best fit for the two parameter model (13) for the six separate regions are summarized in Table 2. The table includes the model parameter estimates and 95% confidence intervals, as well as the R^2 values and Akaike information criterion (AIC) values for the different regions, altitudes, environment types, and total for all families for all regions. R^2 values range from 0.03 to 0.40, indicating relatively poor fits.

Figure 5 shows 95% confidence intervals (± 2 standard deviations) for the model coefficients p_b (*left*) and p_f (*right*) as estimated from different data subsets. Confidence intervals are shown for the coefficient estimates using per-species data from six separate locations (yellow bars); using data grouped by altitude (purple bars); using data grouped by fragmentation (blue bars); and using all data combined (green bar). In 11 of the 12 cases (including the overall data model), the confidence intervals for p_f contain 0, indicating a lack of statistical evidence for a nonzero value of p_f. On the other hand, in 6 of the 12 cases (including the model for overall data), the confidence interval for p_b lies entirely below 0, indicating that p_b is negative at 5% significance.

The fact that p_f is not significantly different from 0 in virtually all cases motivates replacing $p_f = 0$ in (13). The model then has only the coefficient p_b plus the constant C_0 that are fitted to the transect and point count data:

$$\log(X_n) = \log(Z_n) + C_0 + p_b \log(w_n) + e_n. \qquad (47)$$

Figures 6–9 show the transect observation prediction accuracy for the one-coefficient model (47) by region, elevation, environment type and combined,

Location	C_0 est.	C_0 CI	p_b est.	p_b CI	p_f est.	p_f CI	R^2	AIC
By region								
Bombuscaro	3.6	[2.4,4.9]	-0.6	[-0.9,-0.3]	0.2	[-0.2,0.5]	0.30	92
ECSF	3.7	[0.4,7.1]	-0.6	[-1.4,0.2]	0.2	[-0.8,1.0]	0.19	64
Cajanuma	3.6	[-7.2,14.5]	-0.7	[-4.0,2.1]	0.3	[-1.0,1.6]	0.12	24
Copalinga	2.0	[0.3,3.7]	-0.2	[-0.5,0.2]	0.2	[-0.2,0.5]	0.03	141
Finca	3.0	[0.6,5.3]	-0.7	[-1.3,-0.04]	0.02	[-0.4,0.5]	0.25	60
Bellavista	3.7	[-4.0,11.4]	-0.7	[-2.4,0.9]	0.7	[-0.3,1.8]	0.40	23
By altitude								
Altitude 1000	3.7	[-4.0,11.4]	-0.7	[-2.4,0.9]	0.7	[-0.3,1.8]	0.40	23
Altitude 2000	3.2	[1.4,5.1]	-0.7	[-1.2,0.9]	-0.2	[-0.6,0.2]	0.3	86
Altitude 3000	4.2	[-0.8,9.1]	-0.8	[-2.0,0.3]	-0.7	[-1.4,-0.0]	0.40	33
By environment								
Natural	3.4	[2.2,4.5]	-0.6	[-0.8,-0.3]	-0.1	[-0.4,0.1]	0.40	166
Fragmented	1.9	[0.6,3.1]	-0.3	[-0.6,-0.0]	0.0	[-0.2,0.2]	0.08	208
All birds	2.2	[1.2,3.1]	-0.4	[-0.6,-0.1]	-0.1	[-0.3,0.1]	0.11	245

Table 2: Summary of parameter estimations from regression for the two-coefficient model (13), which includes both bird weight and fruit weight

Location	C_0 est.	C_0 CI	p_b est.	p_b CI	R^2	AIC
By region						
Bombuscaro	3.4	[2.2,4.5]	-0.6	[-0.9,-0.2]	0.27	91
ECSF	3.4	[0.4,7.1]	-0.6	[-1.3,0.2]	0.18	63
Cajanuma	1.7	[-3.9,7.3]	-0.3	[-1.9,1.4]	0.03	22
Copalinga	1.5	[0.28,2.7]	-0.14	[-0.5,0.2]	0.01	140
Finca	2.9	[0.8,5.0]	-0.7	[-1.3,-0.0]	0.25	58
Bellavista	-0.9	[-5.4,3.6]	0.09	[-1.15,1.33]	0.00	25
By altitude						
Altitude 1000	1.9	[0.9,2.9]	-0.3	[-0.5,0.0]	0.06	156
Altitude 2000	2.9	[1.2,4.6]	-0.6	[-1.1,-0.2]	0.27	85
Altitude 3000	0.2	[-3.7,4.1]	-0.1	[-1.1,1.0]	0.00	37
By environment type						
Natural	3.1	[2.1,4.0]	-0.5	[-0.8,-0.25]	0.22	166
Fragmented	1.9	[0.9,2.9]	-0.3	[-0.6,0.0]	0.08	206
All	2.0	[1.2,2.8]	-0.3	[-0.6,-0.1]	0.1	244

Table 3: Summary of parameter estimations from regression for the single coefficient model with $p_f = 0$ in (13)

respectively (compare Figures 1–4, which show corresponding accuracies for the two-coefficient model). Visually, the one-coefficient model appears to have accuracies comparable to the two-coefficient model.

Table 3 summarizes the fit parameters for the single-coefficient model that includes only bird weight and not fruit weight. The AIC values for all fits are reduced, thus justifying the exclusion of fruit weight as a significant parameter.

Figure 10 summarizes confidence intervals for the bird weight coefficient in the single coefficient model of the form (47). Confidence intervals are consistent across all regions, and overlap in the range of -0.3 to -0.5 As with the two-coefficient model, the bird weight coefficients tend to be negative, and some are negative with 95% significance (including the overall estimate). Confidence intervals for high altitudes are smaller, which may reflect the fact that bird species are more variable at low altitudes. Taken together, the data gives a 95% confidence interval for p_b as $[-0.6, -0.1]$. Recall that p_b is defined in (14) as the difference of the two exponents p_{b1} and p_{b2}, where p_{b1} is related to bird weight's influence on energy consumption per bird, while p_{b2} is related to bird weight's influence on energy consumption per observed feeding. Estimates of p_{b1} in [3] for most families are close to 0.7 (hummingbirds are an exception, with $p_{b1} > 1$). Our finding of $p_b \approx -0.4$ is consistent with a value of $p_{b1} \approx 1$. This

would imply that bird energy consumption per feeding is roughly proportional to bird weight. This is not unreasonable, because heavier birds proportionately have larger anatomical features, including larger stomachs.

Figure 11 shows a scatter plot of the ratio of feeding observation/transect observations versus transect observations frequency. The figure indicates a negative correlation between the feeding/observation ratio and the number of observations. This is an indication of regression to the mean: overpredictions (or underpredictions) based on transect observations will tend to be revised downwards (or upwards) based on feeding observations—and vice versa. The presence of regression to the mean indicates that point counts and transect observations tend to counterbalance each other, and estimators based on both factors may achieve greater accuracy. (This indication is confirmed in the following sections.)

3.2 Peformance evaluation of estimators

Figures 12–27 show the results of the performance evaluation tests described in Section 2.2.3. The figures comprise two subsets: Figures 12–19 show results for estimators that were computed using the mean value $\bar{p}_b$ of the bird weight exponent, while Figures 20–27 use the exact values of $p_{b,n}$ in the calculations. The two subsets corresponds to cases (a) and (b) respectively described in Section 2.2.3, in which the investigator has imprecise or precise information respectively about the dependence of energy consumption on weight.

Case (a): estimators computed using mean value of bird weight exponent $\bar{p}_b$

Figure 12 shows the mean over the 10,000 simulations for the estimators $(\widehat{a}_n)_z$ and $(\widehat{a}_n)_x$ for different bird families for different regions, compared to the true value. The figure shows that there is very little bias in most estimators, where the estimators for birds with lower relative abundance tending have the most bias. Figure 13 shows similar results for the estimators of log abundance $(\widehat{\ell}_n)_z$ and $(\widehat{\ell}_n)_x$.

Figure 14 shows the relative standard deviations for $(\widehat{a}_n)_\beta$ for values of β between 0 and 1, for all different bird families in the six different regions. The standard deviations are based on estimates of 10,000 simulated ecologists. The triangles on the horizontal axis mark the optimal values of β as calculated by (30). The figure shows that (30) gives a very accurate estimate of the β that minimizes standard deviation. All of the optimal β values are between 0.5 and 1, reflecting the fact that $(\widehat{a}_n)_z$ (corresponding to $\beta = 1$) is consistently smaller than $(\widehat{a}_n)_x$ (corresponding to $\beta = 0$). Figure 15 shows similar results for the standard deviations for $(\widehat{\ell}_n)_\beta$ for the different bird families in different regions.

Figure 16 shows the relative root mean squared error (RMSE) of $(\widehat{a}_n)_\beta$ as a function of β for different bird families per region. The graph is almost identical to Figure 14, showing that bias error is negligible compared to standard deviation. Comparison of Figure 17 with Figure 15 similarly shows that bias is negligible for $(\widehat{\ell}_n)_\beta$ for the different bird families in different regions.

Figure 18 shows RMSE-minimizing β values for $(\widehat{a}_n)_\beta$ versus relative abundance of bird family for the six regions. It is clear that lower β values (between

0.4 and 0.9) only occur for birds with relative abundance less than about 0.1, with only two exceptions from the Cajanuma dataset (which had only 2 families). Figure 19 shows that similar results hold for RMSE-minimizing β values for $(\widehat{\ell}_n)_\beta$.

Case (b): Estimators computed using exact values of bird weight exponent $p_(bn)$

Figures 20–27 correspond to Figures 12–19, except they show results obtained when the investigator has accurate information about $p_{b,n}$. It is evident that the estimates $(\widehat{a}_n)_X$ and $(\widehat{\ell}_n)_X$ (corresponding to $\beta = 0$ in Figures 22–25) are more accurate than the corresponding estimates when $p_{b,n}$ is not known exactly (as depicted in Figures 22–25). As a result the optimal values of β tend to be lower when $p_{b,n}$ is known exactly, as shown in Figures 26 and 27 (compare Figures 18 and 19 for the inexact case).

3.3   Verification of accuracy of combined estimators

Figures 28–29 show 95% confidence intervals for the relative errors for point count-based, transect observation-based, and combined estimators of relative abundance obtained from the simulations described in Section 2.1.7; results are also summarized numerically in Tables 4 and 5. Figures 28–29 both show that relative abundance estimates using the combined estimator $(\widehat{a}_n)_\beta$ are always at least as accurate as the best of the two estimators $(\widehat{a}_n)_Z$ and $(\widehat{a}_n)_X$. Comparing between the two figures shows that combined estimates using the exact values for $p_{b,n}$ are mjuch more accurate than those using $\overline{p_b}$.

Figures 30 and 31 show similar results for relative errors of different estimators of log relative abundance. Taken together, the results indicate that more accurate knowledge about individual bird species' energy consumption habits and requirements can enable improved estimates of relative abundance and log relative abundence. The degree of improvement is displayed quantitatively in Table 4 (for relative abundance estimators) and in Table 5 (for log relative abundance estimators).

4   Conclusions

The principal conclusions of this study are as follows:

- For the dataset drawn from six habitats in Ecuador described in Section 2.2.1, there is no statistical evidence that bird species' energy consumption depends on the weight of fruit consumed.

- On the other hand, there is statistical evidence from this dataset supporting existing literature that energy consumption of bird species depends on bird weight.

	$\widehat{a}_Z$ rel. err.		$\widehat{a}_X$ rel. err.		$\widehat{a}_\beta$ rel. err.	
	(mean $\pm$ std)		mean $\bar{p}_b$	exact p_b	mean $\bar{p}_b$	exact p_b
Bombuscaro	0.28 ± 0.21		0.55 ± 0.18	0.16 ± 0.11	0.22 ± 0.12	0.14 ± 0.10
Copalinga	0.30 ± 0.19		0.64 ± 0.27	0.23 ± 0.17	0.25 ± 0.15	0.18 ± 0.13
ECSF	0.23 ± 0.10		0.66 ± 0.29	0.21 ± 0.12	0.21 ± 0.15	0.15 ± 0.08
Finca	0.16 ± 0.10		0.54 ± 0.28	0.16 ± 0.14	0.15 ± 0.09	0.11 ± 0.08
Cajanuma	0.18 ± 0.11		0.33 ± 0.21	0.14 ± 0.09	0.16 ± 0.10	0.10 ± 0.07
Bellavista	0.20 ± 0.16		$0.6\ pm\ 0.34$	0.49 ± 0.50	0.2 ± 0.16	0.18 ± 0.15
Altitude 1	0.29 ± 0.20		0.61 ± 0.24	0.20 ± 0.15	0.24 ± 0.14	0.16 ± 0.12
Altitude 2	0.20 ± 0.10		0.61 ± 0.29	0.19 ± 0.13	0.18 ± 0.09	0.13 ± 0.08
Altitude 3	0.19 ± 0.15		0.51 ± 0.33	0.37 ± 0.44	0.18 ± 0.14	0.16 ± 0.13
Natural	0.25 ± 0.18		0.57 ± 0.26	0.26 ± 0.27	0.21 ± 0.11	0.16 ± 0.13
Fragmented	0.25 ± 0.18		0.61 ± 0.29	0.26 ± 0.27	0.22 ± 0.15	0.16 ± 0.13
TOTAL	0.25 ± 0.18		0.59 ± 0.27	0.22 ± 0.22	0.21 ± 0.13	0.15 ± 0.11

Table 4: Means and standard deviations of RMSE for relative error of point count-based, transect-based, and combined relative abundance estimators, for individual regions and grouped by altitude, environment type, and total.

	$\widehat{\ell}_Z$ rel. err.		$\widehat{\ell}_X$ rel. err.		$\widehat{\ell}_\beta$ rel. err.	
	(mean $\pm$ std)		mean $\bar{p}_b$	exact p_b	mean $\bar{p}_b$	exact p_b
Bombuscaro	0.26 ± 0.15		0.53 ± 0.15	0.18 ± 0.13	0.22 ± 0.11	0.14 ± 0.09
Copalinga	0.29 ± 0.17		0.58 ± 0.18	0.23 ± 0.16	$0.25/0.18$	$0.13/0.11$
ECSF	0.24 ± 0.12		0.63 ± 0.20	0.22 ± 0.13	0.22 ± 0.10	0.16 ± 0.09
Finca	0.17 ± 0.11		0.52 ± 0.18	0.16 ± 0.14	0.16 ± 0.10	0.12 ± 0.09
Cajanuma	0.2 ± 0.14		0.33 ± 0.20	0.15 ± 0.10	0.17 ± 0.11	0.12 ± 0.08
Bellavista	0.21 ± 0.18		0.53 ± 0.11	0.34 ± 0.2	0.18 ± 0.12	0.17 ± 0.12
Altitude 1	0.28 ± 0.16		0.56 ± 0.17	0.21 ± 0.15	0.24 ± 0.12	0.16 ± 0.11
Altitude 2	0.21 ± 0.12		0.58 ± 0.20	0.19 ± 0.13	0.20 ± 0.10	0.14 ± 0.09
Altitude 3	0.21 ± 0.16		0.46 ± 0.17	$0.17/\pm 0.19$	0.18 ± 0.12	0.15 ± 0.11
Natural	0.25 ± 0.14		0.55 ± 0.20	0.19 ± 0.13	0.22 ± 0.11	0.15 ± 0.09
Fragmented	0.25 ± 0.17		0.56 ± 0.17	0.23 ± 0.17	0.22 ± 0.13	0.16 ± 0.11
Total	0.25 ± 0.15		0.57 ± 0.22	0.23 ± 0.22	0.22 ± 0.12	0.15 ± 0.10

Table 5: Means and standard deviations of RMSE for error of point count-based, transect-based, and combined log relative abundance estimators, for individual regions and grouped by altitude, environment type, and total.

- There is also statistical evidence from this same dataset that energy consumption per transect observation is larger for heavier birds than for lighter birds.

- Simulations based on the Ecuador data demonstrate that point counts and transect observations can be used to obtain independent estimates of relative abundance and log of relative abundance. Typically, the point

	$\widehat{a}_\beta$ 90% C.I. proportions		$\widehat{\ell}_\beta$ 90% C.I. proportions	
	using $\overline{p}_b$	using exact p_b	using $\overline{p}_b$	using exact p_b
Bombuscaro	0.906 ± 0.007	0.900 ± 0.004	$0.907 \pm .007$	0.901 ± 0.004
Copalinga	0.905 ± 0.008	0.901 ± 0.004	0.906 ± 0.001	0.901 ± 0.004
ECSF	0.905 ± 0.007	0.903 ± 0.002	0.905 ± 0.007	0.904 ± 0.002
Finca	0.903 ± 0.005	0.905 ± 0.008	0.903 ± 0.005	0.906 ± 0.008
Cajanuma	0.902 ± 0	0.907 ± 0	0.902 ± 0	0.908 ± 0
Bellavista	0.904 ± 0.005	0.901 ± 0.009	0.904 ± 0.005	0.902 ± 0.009
Altitude 1.	0.906 ± 0.008	0.901 ± 0.004	0.906 ± 0.008	0.901 ± 0.004
Altitude 2	0.904 ± 0.006	0.904 ± 0.005	0.904 ± 0.006	0.904 ± 0.005
Altitude 3	0.903 ± 0.004	0.903 ± 0.008	0.9034 ± 0.004	0.904 ± 0.008
Natural	0.905 ± 0.007	0.902 ± 0.004	0.905 ± 0.007	0.9028 ± 0.004
Fragmented	0.904 ± 0.007	0.902 ± 0.006	0.905 ± 0.007	0.902 ± 0.006
TOTAL	0.905 ± 0.007	0.902 ± 0.005	0.905 ± 0.007	0.903 ± 0.005

Table 6: Proportion of instances where the true relative abundance (respectively log relative abundance) lay within the 90% confidence intervals computed by the technique described in this section. Estimates using the mean value of the bird weight exponent $\overline{p}_b$ and the exact value of p_b are shown. The error bars are for one standard deviation calculated from the proportions obtained for different families in the region.

count-based estimates are more accurate.

- The two estimates mentioned in the previous item may be linearly combined to form another estimate which has lower RMSE. The user may run simulations to estimate the parameter β needed to form a linear combination of point count based and transect observation based abundance estimators with smaller root mean squared error.

- Error bars may be computed for the linearly combined relative abundance estimates that accurately reflect the uncertainty in these estimates.

- With more accurate information concerning the dependence of bird type energy consumption as a function of weight, a better estimate of β can be made that leads to combined estimators of abundance with lower root mean squared error.

CRediT statement

Conceptualization: V.S., M.J.A.B., A.B., C.T.; **Data curation**: V.S., M.Q., E.L.N., M.S.; **Formal analysis**:M.J.A.B., A.B., C.T.; **Funding acquisition**: K.B.S.,O.A.; **Investigation**: V.S., M.Q., E.L.N., M.S.; **Methodology**: M.J.A.B.,A.B.,C.T.; **Project administration**: V.S., K.B.S., O.A.; **Resources**: K.B.S., O.A.; **Software**: M.J.A.B., A.B., C.T.; **Supervision**: K.B.S., V.S., C.T.; **Validation**: M.J.A.B., C.T.; **Visualization**: M.J.A.B., A.B., C.T.; **Writing** −

original draft: M.J.A.B., A.B., C.T.; **Writing – review & editing**: V.S., E.L.N., C.T.

Acknowledgements

We are grateful to the Fulbright Program administered by the US Department of State, the Catholic University of Cuenca and to Texas A & M University-Central Texas for providing travel funding and logistical support during the conduct of this research.

References

[1] Colin J Bibby, Neil D Burgess, David M Hillis, David A Hill, and Simon Mustoe. *Bird census techniques*. Elsevier, 2000.

[2] Stefany Coxe, Stephen G West, and Leona S Aiken. The analysis of count data: A gentle introduction to poisson regression and its alternatives. *Journal of personality assessment*, 91(2):121–136, 2009.

[3] J. Gurney D. Crocker, A. Hart and C. McCoy. Project pn0908: Methods for estimating daily food intake of wild birds and mammals. Technical report, Central Science Laboratory, 2002.

[4] Paul Emck. *A climatology of south Ecuador-with special focus on the major Andean ridge as Atlantic-Pacific climate divide*. PhD thesis, Friedrich-Alexander-Universität Erlangen-Nürnberg (FAU), 2007.

[5] Markus Kottek, Jürgen Grieser, Christoph Beck, Bruno Rudolf, and Franz Rubel. World map of the köppen-geiger climate classification updated. *Meteorologische Zeitschrift*, 15(3):259–263, 07 2006.

[6] C. John Ralph, John R. Sauer, and Sam Droege. Monitoring bird populations by point counts. Technical report, 1995.

[7] William J Sutherland. *Ecological census techniques: a handbook*. Cambridge university press, 2006.

[8] Hamish Wilman, Jonathan Belmaker, Jennifer Simpson, Carolina de la Rosa, Marcelo M Rivadeneira, and Walter Jetz. Eltontraits 1.0: Species-level foraging attributes of the world's birds and mammals: Ecological archives e095-178. *Ecology*, 95(7):2027–2027, 2014.

Figures

Figure 1: Transect observation prediction accuracy by location for two-coefficient model that includes both bird and fruit weights. The 45 degree line corresponds to zero prediction error.

Figure 2: Transect observation prediction accuracy of two-coefficient model, by elevation (species data from the two regions at each elevation are combined).

Figure 3: Transect observation prediction accuracy of two-coefficient model, by fragmentation (species data from natural and fragmented regions are combined separately).

Figure 4: Transect observation prediction accuracy for two-coefficient model, all observations (species data from all regions are combined).

Figure 5: 95% confidence intervals for bird weight coefficient (*left*) and fruit weight coefficient (*right*) in two-coefficient model (bird_weight +fruit_weight) by location (yellow), by elevation (purple), by fragmentation (blue) and overall(green)

Figure 6: Transect observation prediction accuracy for one-coefficient model, by region.

Figure 7: Transect observation prediction accuracy for one-coefficient model, by elevation.

Figure 8: Transect observation prediction accuracy for one-coefficient model, by fragmentation.

Figure 9: Transect observation prediction accuracy for one-coefficient model, all observations.

Figure 10: 95% confidence intervals for one-coefficient model (bird weight only) model coefficients by location, by elevation, by fragmentation and overall

Figure 11: Per-family (transect observation/point count) ratio versus point count for combined data.

Figure 12: Mean $(\widehat{a}_n)_z$ and $(\widehat{a}_n)_x$ compared to true relative abundance for different bird families in different regions. Bird families in each region are sorted in increasing order of relative abundance.

Figure 13: Mean $(\widehat{\ell}_n)_z$ and $(\widehat{\ell}_n)_x$ compared to true log relative abundance for different bird families in different regions. Bird families in each region are sorted in increasing order of relative abundance.

Figure 14: Relative standard deviation (standard deviation divided by true value) of $(\widehat{a}_n)_\beta$ for $0 \le \beta \le 1$, for different bird families in different regions. $\beta = 0$ and $\beta = 1$ correspond to $(\widehat{a}_n)_X$ and $(\widehat{a}_n)_Z$, respectively. The stars locate the minima of the different standard deviation curves, while the corresponding triangles on the x axis indicate the value of β corresponding to the minimum value.

Figure 15: Standard deviation of $(\widehat{\ell}_n)_\beta$ for $0 \leq \beta \leq 1$, for different bird families in different regions. $\beta = 0$ and $\beta = 1$ correspond to $(\widehat{\ell}_n)_X$ and $(\widehat{\ell}_n)_Z$, respectively. The stars locate the minima of the different standard deviation curves, while the corresponding triangles on the x axis indicate the value of β corresponding to the minimum value.

Figure 16: Relative root mean squared error (RMSE) (RMSE / true value) of $(\widehat{a}_n)_\beta$ for $0 \le \beta \le 1$, for different bird families in different regions. $\beta = 0$ and $\beta = 1$ correspond to $(\widehat{a}_n)_X$ and $(\widehat{a}_n)_Z$, respectively. The stars locate the minima of the different standard deviation curves, while the corresponding triangles on the x axis indicate the value of β corresponding to the minimum value.

Figure 17: Root mean squared error (RMSE) of $(\widehat{\ell}_n)_\beta$ for $0 \leq \beta \leq 1$, for different bird families in different regions. $\beta = 0$ and $\beta = 1$ correspond to $(\widehat{\ell}_n)_X$ and $(\widehat{\ell}_n)_Z$, respectively. The stars locate the minima of the different standard deviation curves, while the corresponding triangles on the x axis indicate the value of β corresponding to the minimum value.

Figure 18: Optimal value of β (i.e. value of β that minimizes RMSE error) for $(\widehat{a}_n)_\beta$ for $0 \leq \beta \leq 1$ as a function of bird family true relative abundance.

Figure 19: Optimal value of β (i.e. value of β that minimizes RMSE error) for $(\widehat{\ell}_n)_\beta$ for $0 \leq \beta \leq 1$ as a function of bird family true relative abundance.

Figure 20: Mean $(\widehat{a}_n)_z$ and $(\widehat{a}_n)_x$ compared to true relative abundance for different bird families in different regions, for ecologists with exact information about p_b. Bird families in each region are sorted in increasing order of relative abundance.

Figure 21: Mean $\widehat{(\ell_n)}_z$ and $\widehat{(\ell_n)}_x$ compared to true log relative abundance for different bird families in different regions, for ecologists with exact information about p_b. Bird families in each region are sorted in increasing order of relative abundance.

Figure 22: Relative standard deviation (standard deviation divided by true value) of $(\widehat{a}_n)_\beta$ for $0 \leq \beta \leq 1$, for different bird families in different regions. $\beta = 0$ and $\beta = 1$ correspond to $(\widehat{a}_n)_X$ and $(\widehat{a}_n)_Z$, respectively, for ecologists with exact information about p_b. The stars locate the minima of the different standard deviation curves, while the corresponding triangles on the x axis indicate the value of β corresponding to the minimum value.

Figure 23: Standard deviation of $(\widehat{\ell}_n)_\beta$ for $0 \leq \beta \leq 1$, for different bird families in different regions. $\beta = 0$ and $\beta = 1$ correspond to $(\widehat{\ell}_n)_X$ and $(\widehat{\ell}_n)_Z$, respectively, for ecologists with exact information about $p_{b,n}$. The stars locate the minima of the different standard deviation curves, while the corresponding triangles on the x axis indicate the value of β corresponding to the minimum value.

Figure 24: Relative root mean squared error (RMSE) (RMSE / true value) of $(\hat{a}_n)_\beta$ for $0 \leq \beta \leq 1$, for different bird families in different regions. $\beta = 0$ and $\beta = 1$ correspond to $(\hat{a}_n)_X$ and $(\hat{a}_n)_Z$, respectively, for ecologists with exact information about p_b. The stars locate the minima of the different standard deviation curves, while the corresponding triangles on the x axis indicate the value of β corresponding to the minimum value.

Figure 25: Root mean squared error (RMSE) of $(\widehat{\ell}_n)_\beta$ for $0 \leq \beta \leq 1$, for different bird families in different regions. $\beta = 0$ and $\beta = 1$ correspond to $(\widehat{\ell}_n)_X$ and $(\widehat{\ell}_n)_Z$, respectively, for ecologists with exact information about p_b. The stars locate the minima of the different standard deviation curves, while the corresponding triangles on the x axis indicate the value of β corresponding to the minimum value.

Figure 26: Optimal value of β (i.e. value of β that minimizes RMSE error) for $(\widehat{a}_n)_\beta$ for $0 \leq \beta \leq 1$ as a function of bird family true relative abundance, for ecologists with exact information about p_b.

Figure 27: Optimal value of β (i.e. value of β that minimizes RMSE error) for $(\widehat{\ell}_n)_\beta$ for $0 \leq \beta \leq 1$ as a function of bird family true relative abundance, for ecologists with exact information about p_b.

Figure 28: 95% confidence intervals for relative errors (error divided by true value) for three estimators of relative abundance (AZ = point-count based, AX = transect count based, and combined) for different bird families for scenarios based on data from six different regions. The AX and combined estimators are calculated using $\overline{p}_b$ as in (38). Families are ordered in order of increasing abundance in each region.

Figure 29: 95% confidence intervals for relative errors (error divided by true value) for three estimators of relative abundance (AZ = point-count based, AX = transect count based, and combined) for different bird families for scenarios based on data from six different regions. The AX and combined estimators are calculated using the exact value of $p_{b,n}$. Families are ordered in order of increasing abundance in each region.

Figure 30: 95% confidence intervals for three estimators of logarithm of relative abundance (AZ = point-count based, AX = transect count based, and combined) for different bird families for scenarios based on data from six different regions. The AX and combined estimators are calculated using $\overline{p}_b$ as in (38). Families are ordered in order of increasing abundance in each region.

Figure 31: 95% confidence intervals for three estimators of logarithm of relative abundance (AZ = point-count based, AX = transect count based, and combined) for different bird families for scenarios based on data from six different regions. The AX and combined estimators are calculated using the exact value of $p'_{b,n}$ Families are ordered in order of increasing abundance in each region.