A Spatiotemporal Molecular Atlas of the Ovulating Mouse Ovary

AUTHORS: Madhav Mantri[1], Hanxue Hannah Zhang[2], Emmanuel Spanos[1], Yi A Ren[2,#,*], Iwijn De Vlaminck[1,#,*]

AFFILIATIONS:

[1]Nancy E. and Peter C. Meinig School of Biomedical Engineering, Cornell University, Ithaca, New York, [2]Department of Animal Science, Cornell University, Ithaca, New York

[#]These authors contributed equally

*To whom correspondence should be addressed: vlaminck@cornell.edu and yar9@cornell.edu

ABSTRACT

Ovulation is essential for reproductive success, yet the underlying cellular and molecular mechanisms are far from clear. Here, we applied high-resolution spatiotemporal transcriptomics to map out cell-type- and ovulation-stage-specific molecular programs as function of time during follicle maturation and ovulation in mice. Our analysis revealed dynamic molecular transitions within granulosa cell types that occur in tight coordination with mesenchymal cell proliferation. We identified new molecular markers for the emerging cumulus cell fate during the preantral-to-antral transition. We describe transcriptional programs that respond rapidly to ovulation stimulation and those associated with follicle rupture, highlighting the prominent roles of apoptotic and metabolic pathways during the final stages of follicle maturation. We further report stage-specific oocyte-cumulus cell interactions and diverging molecular differentiation in follicles approaching ovulation. Collectively, this study provides insights into the cellular and molecular processes that regulate mouse ovarian follicle maturation and ovulation with important implications for advancing therapeutic strategies in reproductive medicine.

INTRODUCTION

Ovulation is a complex process that involves extensive molecular, cellular and structural changes in distinct compartments in ovarian follicles, which culminate in follicle wall rupture and release of oocytes for fertilization. Successful ovulation is predicated on precise coordination and interaction between cell types within the ovary, both temporally and spatially. Temporally, the gene expression processes that are important for ovulation are regulated dynamically on a time scale of minutes to hours[1]. Spatially, the formation of follicle rupture sites requires spatially restricted tissue remodeling between granulosa and mesenchymal cells in the surrounding stromal tissues[2]. The cellular and molecular mechanisms responsible for controlling this temporal and spatial specificity of follicle rupture, however, remain poorly understood.

Single-cell RNA-sequencing has been used extensively to characterize the diversity of cell types and phenotypes in the ovary[3–6]. Yet, single-cell RNA sequencing requires tissue dissociation and therefore leads to the loss of spatial information regarding cellular niches, cell-cell interactions and the interplay between morphological changes and cellular changes in tissues[7,8]. Spatial RNA sequencing has been applied recently to study the cellular organization within the ovary, but at limited spatial resolution and without incorporating temporal analyses[9,10]. Here, we address these challenges by implementing spatial RNA sequencing at 10 µm spatial resolution and by profiling gene expression at eight well defined stages preceding ovulation in mice. To induce ovulation, we used exogenous gonadotropin stimulation in immature mice, a widely used model to study the molecular and cellular mechanisms that regulate ovulation. To support our analysis, we developed new analytical approaches to study cell-cell interactions in specific spatial niches within ovarian follicles. Altogether, our work leads to a highly detailed spatiotemporal atlas of ovulation in mice and provides insights into how ovarian follicle maturation and rupture are regulated in time and space.

RESULTS

High-resolution spatial transcriptomics of murine ovaries undergoing ovulation

To investigate the spatiotemporal regulation of ovarian follicle maturation and ovulation in mice, we implemented high-resolution spatial gene expression profiling on ten ovaries, collected at eight well-defined time points between ovulation induction and follicle rupture (Seeker, Curio Biosciences, **Methods, Fig. 1A & 1B**). Following an initial quality control — removing bead transcriptomes with fewer than 100 unique molecules detected and those not located directly under the tissue —we analyzed 121,536 spatial transcriptomes (**Supp. Fig. 1A-1C**). We assigned the most likely cell type to each spatial transcriptome via deconvolution using a scRNA-seq reference (**Methods**). The spatial transcriptomes represented seven broad cell types: granulosa cells, mesenchymal cells, oocytes, endothelial cells, epithelial cells, immune cells, and erythrocytes (**Fig. 1C, Supp. Fig. 1D**). As each bead used in spatial transcriptomics

measures 10 µm, this dataset provides the first near single-cell resolution, whole-genome spatiotemporal transcriptomic atlas of immature and ovulating ovaries.

To refine the classification of granulosa cell types, we used additional clustering and analysis of expression of canonical cell-type specific markers in conjunction with observed spatial location within the tissue (**Supp. Table 1, Supp. Fig. 1C & 2A, Methods**). This allowed us to identify mural cells located along the follicular walls, and cumulus cells surrounding the oocytes at each stage during ovulation (**Fig. 1D, Supp. Fig. 2A & 2B**). We further identified four types of mural granulosa cells with distinct transcriptomic profiles that are linked to follicle state (preantral, antral, atretic, and lytic follicles, **Fig. 1D, Supp. Fig. 2A & 2B**). Additionally, we refined cell type labels for beads within the mesenchyme via deconvolution using a scRNA-seq reference[3], and identified smooth muscle cells, fibroblast-like cells, theca cells, dividing mesenchyme cells, and pericytes (**Fig. 1C, 1D, Supp. Fig. 2C & 2D**). Mesenchymal smooth muscle and fibroblast-like cells were visible in some of the ovarian tissue sections both in the medullary stroma (**Supp Fig. 2C & 2D**) and associated with growing follicles (**Supp Fig. 2C**). To enable the characterization of cell-cell interactions and molecular changes specific to defined stages of follicle maturation, we manually segmented the 335 follicles represented in the ten ovary tissue sections (**Supp. Fig. 3A-C**). Spatiotemporal maps for the transcripts of *Pgr*, *Ptgs2* and *Ihh* demonstrated patterns of expression consistent with previous reports[11–14] (**Fig. 1E, 1F and 1G**). Collectively, we generated a spatially resolved hierarchical map of the cell types and their phenotypes in immature and preovulatory mouse ovaries during hormone-induced ovulation.

Figure 1 | Spatial transcriptomics profiling of murine ovaries undergoing hormone-induced ovulation.
A) Illustration depicting the internal components of the mouse ovary. **B)** Workflow schematic outlining the steps followed in the experiment. PMSG = pregnant mare serum gonadotropin; hCG = human chorionic gonadotropin. Spatial transcriptomics was conducted on tissue sections from three untreated immature ovaries and seven ovaries undergoing hormone-controlled ovulation. **C)** Tree diagram showing multi-level hierarchy of different cell types within the ovary. The first level of the tree represents the broad cell type labels predicted for Slide-seq beads using scRNA-seq data from cycling mouse ovaries. The leaves of the tree represent the fine cell type labels assigned using canonical markers after unsupervised clustering of bead transcriptomes in individual samples. **D)** Spatial transcriptomics maps of three immature ovaries and seven representative ovaries undergoing hormone-controlled ovulation colored by fine-grained cell type labels. **E-G)** Spatial transcriptomics maps showing expression of *Pgr*, *Ptgs2*, and *Ihh* over eight time points before and after ovulation stimulation.

Spatial delineation and novel molecular markers of granulosa cells associated with preantral-to-antral transition

We first analyzed granulosa cells within the immature ovaries to map their molecular changes during folliculogenesis prior to ovulation induction (**Fig. 2A**). Unsupervised clustering identified four distinct granulosa cell types: *i)* granulosa with preantral phenotype expressing markers *Kctd14* and *Gas6*, *ii)* granulosa with early antral phenotype expressing markers *Inhba* and *Inhbb*, *iii)* granulosa with atretic phenotype expressing markers *Ctgf* and *Cfh*, *iv)* and granulosa cells with mitotic phenotype expressing *Top2a* and *Cdk1* (**Fig. 2B & 2C**). We used the phenotype of granulosa cells to classify follicles as preantral (*Kctd14*+ granulosa cells), antral (*Inhbb*+ granulosa cells) or atretic (*Cfh*+ granulosa cells, **Fig. 2D**). The majority of follicles in immature ovaries were atretic, and most antral follicles were located along the edge of the ovary. Consistent with the absence of antrum observed in H&E staining, preantral follicles mapped by spatial transcriptomics were completely filled with granulosa cells (**Supp. Fig. 1C**). Notably, mitotic granulosa cells in G2M phase were more prevalent in antral follicles, consistent with the observation of *Top2a* and *Cdk1* expression in these follicles, and in line with the expected growth of antral follicles (**Fig. 2E**). Interestingly, in atretic follicles, while the outer layers of granulosa cells expressed markers of degeneration, the inner layers of granulosa cells expressed mitotic markers (**Supp. Fig. 4A & 4B**). Last, we used the follicle segmentation and classification to identify novel markers of granulosa cells associated with follicle maturation stage. *Gas6*, *Gstm2*, *Isyna1* were enriched in preantral follicles; *Nap1l5* in antral follicles; and *Gm26917*, *Itih2* in atretic follicles (**Fig. 2F**).

The transition between preantral and antral follicle is a crucial stage of follicle maturation. We overlaid markers of preantral (*Kctd14*) and antral (*Inhbb*) follicle phenotypes, which revealed intricate spatial patterns of granulosa cell differentiation markers during the preantral-to-antral follicle transition (**Fig. 2G**). Preantral and early antral follicles showed an enrichment of a *Inhbb*+*Kctd14*+ cell population in the innermost layers of granulosa cells (boxed follicles: upper, preantral; lower, early antral. Histology staining of these follicles is shown in **Supp. Fig. 1C**. In the same follicles, the *Inhbb*+*Kctd14*- granulosa cells surrounded the double-positive cells. As antral follicles grew bigger, the innermost granulosa cells retained the expression of both *Inhbb* and *Kctd14* while the population of *Inhbb*+*Kctd14*- granulosa cells expanded. In preovulatory follicles, we found that mural granulosa cells expressed only *Inhbb*, while cumulus cells expressed both *Inhbb* and *Kctd14* (**Supp. Fig. 4C-4E**). Together, these observations strongly suggest that *Inhbb*+*Ktcd14*- cells along the follicle wall are precursors of mural granulosa cells, and that the *Inhbb*+*Kctd14*+ cells in the center of small antral follicles are precursors of cumulus cells, which corroborates with a previous report on the expression of *Kctd14* in preantral follicles and cumulus cells[3]. In summary, these observations revealed distinct layers of granulosa cells emerging and expanding during preantral-to-antral transition.

To delve deeper into the molecular signatures underlying the transition between
preantral to antral follicles, we performed differential gene expression analysis and found *Ank1*,
Aldh3b1, *Irx3*, and *Cacna1a* upregulated in *Inhbb+Kctd14+* double-positive granulosa cells (**Fig.
2H, Supp. Fig. 4F**). We performed multiplexed Fluorescence In Situ Hybridization (FISH) to
visualize *Ank1* expression in ovaries from untreated mice, and found that *Ank1* (ankyrin 1 that
regulates cell membrane stability and shape) was indeed expressed in the *Inhbb+Kctd14+* double-
positive granulosa cells surrounding oocytes (**Fig. 2I**). Intriguingly, in preantral follicles where all
granulosa cells expressed *Kctd14* but not *Inhbb*, *Ank1* was enriched in granulosa cells surrounding
the oocytes towards the center of the follicles; while in preovulatory follicles, *Ank1* expression is
retained in cumulus cells but not mural granulosa cells (**Supp. Fig. 4G & 4H**). We suggest that
Ank1, *Aldh3b1*, *Irx3* and *Cacna1a* are early molecular markers for further differentiation of the
cumulus cell fate, and that their induction of expression is associated with the preantral-to-antral
follicle transition. Taken together, our findings delineate molecular and spatial changes in distinct
granulosa cell layers and identified novel molecular regulators during the crucial preantral-to-
antral transition.

**Figure 2 | Spatial and molecular differentiation of distinct granulosa cell layers emerging during
prenatal-to-antral transition. A)** Illustration showing the structural transition from preantral to antral
ovarian follicles. **B)** Spatial transcriptomics maps of three untreated immature ovaries colored by
granulosa cell type. **C)** Spatial expression of three immature untreated ovaries showing the expression of
previously reported and novel gene markers for preantral, antral, atretic, and mitotic follicles. **D)** Spatial
transcriptomics maps of immature ovaries colored by follicle type as determined by gene expression of
mural granulosa cells within the follicles. **E)** Bar plot showing proportion of cells in various cell cycle phases
across different follicle types. **F)** Heatmap showing differentially expressed genes in mural granulosa cells
across different follicle types. **G)** Spatial transcriptomics maps showing spatial organization for the
expression of preantral follicle marker, *Kctd14*; and antral follicle marker, *Inhbb*. **H)** Spatial transcriptomics
maps showing the expression of genes specific to granulosa cells in follicles transitioning from preantral
to antral stage: *Ank1*, *Aldh3b1*, *Irx3*, and *Cacna1a*. **I)** Multiplexed RNA FISH staining for preantral marker

172   *Kctd14* (yellow), antral marker *Inhbb* (magenta), and *Ank1* gene (cyan) on untreated immature ovary. The
173   dotted boxes show zoomed-in images of a representative antral follicle. Representative images are from
174   five biological replicates.

175   ───
176

177   **Spatiotemporal response of mural granulosa cells to ovulation stimulation**

178   To delineate molecular changes driving follicle maturation and ultimately ovulation, we
first analyzed the rapid transcriptomic response of mural granulosa cells to hCG treatment that
mimics the preovulatory luteinizing hormone surge. Differential gene expression analysis
identified 77 genes with significantly upregulated transcript levels (log fold change > 1.0, adjusted
p-value < 0.01) as early as one-hour post-hCG treatment (0h post-hCG vs. 1h post-hCG, **Supp. Fig.**
5A & Fig. 3A), including *Serpin3a*, *Lars2*, and *Enc1.* As expected, they were predominantly
expressed in mural granulosa cells of antral follicles (**Fig. 3A**). Gene ontology analysis of genes
upregulated in mural granulosa cells one-hour post-hCG revealed an enrichment of ontology
terms representing negative regulation of apoptosis and chromatin remodeling (**Fig. 3B**).
Consistent with previous reports that hCG/LH inhibits granulosa cell apoptosis[15–17], our analysis
revealed an acute induction of transcripts that negatively regulate apoptotic processes and of
cell cycle regulator *Ccnd2* in preovulatory follicles at 0-1h post-hCG[18,19] (**Supp. Fig. 5B & 5C**).
Regulatory network analysis on spatial transcriptomes of mural granulosa cells further revealed
regulons that are responsive to hCG treatment (**Methods**), including *Mtf1*(+), *Tfap2e*(+), *Sp1*(+),
Klf4(+), and *Mecp2*(+) (**Fig. 3C**). *Tfap2e*, *Sp1* and *Klf4* regulons have been implicated in
transcriptional regulation of steroidogenesis[20]. Consistent with previous work[21], we also
observed in mural granulosa cells of preovulatory follicles a sharp increase in transcript levels of
Cyp19a1 between 0-1h post-hCG, before they decreased to base levels after 4h post-hCG (**Supp.**
Fig. 5D). Interestingly, expression of *Cyp17a1* in theca cells was also initially upregulated and then
decreased to base levels with similar temporal dynamics as *Cyp19a1* (**Supp. Fig. 5E**). These
observations support a temporally coordinated increase in steroidogenic activities between
mural granulosa cells and theca cells as a rapid response to hCG stimulation.
200

201   We further clustered genes into groups based on their temporal expression patterns in
mural granulosa cells (**Methods**, **Supp. Fig. 5F**). Genes that exhibited a gross trend of gradual
decrease in expression following hormone treatment included *Nppc*, an important regulator of
oocyte meiotic arrest[22], and those involved in apoptosis regulation, *Grem1*, *Grem2*[23], and *Comp*
(**Fig. 3D & 3E, Supp. Fig. 5G**). *Comp* (also known as *thrombospondin-5*, a non-collagenous
extracellular matrix protein that suppresses apoptosis), was expressed in mural granulosa cells
and cumulus cells of small antral follicles in the immature ovaries, and in large antral follicles of
preovulatory ovaries, with its expression increasing abruptly at 0-1h post-hCG treatment but
gradually declining after 1h post-hCG treatment (**Fig. 3E**). We calculated and visualized a module

score for a panel of 12 genes associated with negative regulation of apoptosis and found this score first increased between 0-1h post-hCG, then sharply decreased in mural granulosa cells of the preovulatory follicles in contrast to preantral and atretic follicles (**Fig. 3F, Supp. Fig. 5H**). The initial increase of this gene module score is consistent with an upregulation of genes that negatively regulate apoptosis (**Fig. 3B**). We also found that, between 4h and 8h post-hCG, while this module score gradually decreased in mural granulosa cells of preovulatory follicles, it remained at appreciable levels in cumulus cells until immediately before ovulation (**Supp. Fig. 5I**). These observations support an important role of apoptosis regulation in the final stage of follicle maturation and preparation for ovulation.

Among genes that exhibit a continuous increase in expression in response to hormone treatment, *Edn2* emerged as one of the most variable genes. *Edn2* was expressed in mural granulosa cells of the preovulatory follicles immediately prior to ovulation (**Fig. 3G and 3H**). *Edn2* is known for its crucial role in ovulation[24,25]. To uncover potential novel regulators of follicle rupture and the transition to luteal phase, we searched for genes that are spatially and temporally associated with *Edn2* in preovulatory follicles (**Methods**). *Adamts1* was among these genes, consistent with previous reports of its important role in ovulation[12,26,27] (**Fig. 3I**). We also identified a gradual increase in the expression of *Apln* in mouse mural granulosa cells, consistent with the role of *Apln* in ovarian steroidogenesis as reported in several species other than mice[28,29] (**Fig. 3I**). *Cyr61* (also known as *Ccn1*) promotes adhesion of endothelial cells and upregulates angiogenic genes[30], both important processes for the formation of corpus luteum. *Gpx3* codes for glutathione peroxidase, where glutathione protects cells from oxidative stress and is associated with fertility[31]. Notably, among genes that shared a similar expression pattern as *Edn2*, we found an enrichment of genes regulating cell metabolism, including *Akr1c18*[32] (more commonly known as *20alpha-HSD*; metabolizes progesterone to its inactive form), *Rora* (an orphan nuclear receptor regulating cell differentiation, circadian gene expression and cell metabolism), *Ldhd* (lactate dehydrogenase D; catabolizes D-lactate to pyruvate), and *Fabp4*[33] (regulates long-chain fatty transport and metabolism), which is consistent with the results from gene ontology enrichment analysis (**Supp. Fig. 5K & 5L**). Multiplexed FISH demonstrated that *Ldhd* was not expressed in the ovary at 0h post-hCG but co-expressed with *Rhox8* and *Edn2* in mural granulosa cells shortly before ovulation (**Fig 3J and Supp. Fig. 5J**). Together, these observations reveal temporally coordinated changes of metabolic activities and follicle rupture in mural granulosa cells during hormone-induced ovulation.

Figure 3 | Spatiotemporal response of mural granulosa cells to ovulation induction. A) Spatial maps
showing genes upregulated in mural granulosa cells one hour after PMSG and hCG hormone treatment.
B) Bar plot showing the top six gene ontology (GO) terms for genes upregulated one hour after hCG
treatment. **C)** Rank for gene regulatory networks, referred to as regulons, in mural granulosa cells based
on regulon specificity score (RSS) one-hour post-hCG treatment. Top five most specific regulons at 0h and
1h post-hCG is labeled in red. **D)** Line plot showing temporal trends of genes that monotonously decreased
with time post-hCG in mural granulosa cells of preovulatory follicles. **E)** Spatial transcriptomics map
showing the temporal expression of *Comp* in immature and preovulatory ovaries. **F)** Spatial
transcriptomics map showing the temporal gene module score for negative regulation of apoptosis in
untreated immature and preovulatory ovaries. **G)** Line plot showing temporal trends of genes that
monotonously increased with time post-hCG in mural granulosa cells from preovulatory follicles. **H)**
Spatial transcriptomics map showing the temporal expression of *Edn2* in untreated immature and
preovulatory ovaries. **I)** Scatter plots showing expression of four genes that correlate with the expression
of *Edn2* in mural granulosa cells. Each point in the scatter plot represents a spatial transcriptome. **J)**

Multiplexed RNA FISH staining for granulosa cell marker *Rhox8* (yellow), rupture-associated marker *Edn2*
(magenta), and metabolic activity gene *Ldhd* (cyan) in preovulatory follicles at 11h post-hCG. The dotted
boxes show zoomed-in images of a representative preovulatory follicle. Representative images are from
three biological replicates.

───

Dynamic and temporally controlled oocyte-cumulus cell interactions in preovulatory follicles

The rupture of ovarian follicles at ovulation must be closely coordinated with meiotic
maturation of oocytes to ensure successful fertilization[34] (**Fig. 4A**). To gain insights into the
molecular changes associated with oocyte maturation, we analyzed the transcriptomes of
individual oocytes represented in the spatial transcriptomic data (**Fig. 4B**). We implemented
stringent quality control (**Methods**) and produced pseudo-bulk transcriptomes specific to 83
whole oocytes (average number of beads per cell = 16, **Supp. Fig. 6A & 6B** and **Fig. 4B & 4C**).
Visualization of the total number of detected genes within oocytes in response to hormone
treatments revealed an initial increase in transcriptional diversity up to 6h post-hCG treatment,
followed by a subsequent decrease (**Fig. 4D**). Clustering analysis of oocyte transcriptomes
showed only subtle variations among oocytes from different follicle types and stages, with one
notable exception: a well-defined cluster of oocytes obtained from ovaries at 11h and 12h post-
hCG (**Supp. Fig. 6B & 6C**). We observed an upregulation of *Btg4*, *C86187*, *Riok1*, and *Ccdc186* in
these mature oocytes (**Fig. 4E**). Of functional interest to oocyte maturation and early
embryogenesis, we also found in this cluster an enrichment of genes that regulate RNA activities
(*Dnajc21*, *Zcchc8*, *Unc50* and *Rbm36*), and DNA demethylation (*Riok1* and *Tet3*). These results
provide valuable insights into the coordination of ovulation and oocyte maturation.

Cumulus cells provide the oocyte with crucial nutrients and undergo expansion in
response to ovulation induction. We computed trends in the expression of highly variable genes
in cumulus cells and clustered the genes into groups based on their temporal expression patterns
(**Methods, Supp. Fig. 6D-6F**). In line with existing knowledge, we observed upregulation of *Ptx3*,
which encodes pentraxin-3, a major component of the extracellular matrix of cumulus cells
during expansion[35] (**Supp. Fig. 6E & 6I**). Among genes that exhibited a sharp increase in
expression between 0h and 4h post-hCG, *Star* and *Cyp11a1* emerged as the most variable gene
(**Supp. Fig. 6G & 6H**), supporting previous findings on steroidogenic activities of cumulus cells
after LH/hCG stimulation[31]. Among genes that display a more gradual increase in expression were
Nupr1, which encodes a nuclear protein transcriptional regulator, and *Spp1*, which encodes
secreted phosphoprotein that binds to the extracellular hydroxyapatite and is responsible for
extracellular matrix organization (**Fig. 4F-G, Supp. Fig. 6J**).

The interaction between cumulus cells and oocytes is highly dynamic and context
dependent. We performed bi-directional cell-cell signaling analysis for cumulus and oocyte cells
within individual follicles at different time points between hCG stimulation and ovulation
(**Methods**). We found time-specific cellular interactions from cumulus to oocyte post-hCG:
INHBB->ACVR1B at 1h; BTC->MAPK3 and VCAN->SELP at 4h; NTS->PRKD1 at 4h and 6h; NPY-
>MAPK3 at 8h, FGF2->MAPK3 at 11h, and IL18->MAPK3 at 12h (**Fig. 4I**). We further identified
novel ligand-receptor interactions, including SDC1->MAPK3 from 6h to 12h post-hCG and
S100A6->ESR1 specifically enriched immediately before ovulation at 11h and 12h post-hCG (**Fig.**
4I). Similarly, we found novel communication pathways from oocytes to cumulus cells post-hCG:
such as PTPN6->CDK1/JAK1 at 1h, FYN->PTGS2 at 4h, JAG2->NOTCH2 at 8h, and NTN1->PIK3R1
at 12h post-hCG (**Fig. 4J**). Collectively, these observations reveal new cellular interactions in the
dynamic and temporally controlled cellular niche of cumulus cells and oocytes during the
preovulatory stage.
309

Figure 4 | Dynamic and temporally controlled oocyte-cumulus cell interaction in preovulatory follicles.
A) Illustration showing the structure of a preovulatory follicle with oocyte and surrounding cumulus cells.
B) Analysis workflow for oocyte segmentation and characterization of oocyte heterogeneity across follicle types. **C)** Bar plot showing the counts of segmented oocytes in preovulatory ovaries at different stages across different follicle types. **D)** Box plot showing the number of genes detected per oocyte across time points. **E)** Spatial transcriptomics maps of immature and ovulating ovaries showing the expression of a pan-oocyte marker (*H1foo*) and four genes enriched in mature oocytes: *C86187*, *Riok1*, *Ppig*, and *Dnajc21*. **F)** Line plot showing temporal trends of genes that show a gradual increase in expression over time post-hCG in cumulus cells from preovulatory follicles. **G)** Spatial transcriptomics map showing the temporal expression of *Nupr1* in immature and preovulatory ovaries. **H)** Spatial transcriptomics map showing the temporal expression of *Spp1* in immature and preovulatory ovaries. **I)** Heatmap showing the mean expression of top-five differentially expressed ligand-receptor genes expressed in cumulus cells and

323   oocytes from preovulatory follicles, respectively. Each row represents one cumulus-oocyte-complex
324   (COC). **K)** Heatmap showing the mean expression of top-five differentially expressed ligand-receptor
325   genes expressed in oocytes and cumulus cells from preovulatory follicles, respectively. Each row
326   represents one COC.

327
328

Spatiotemporally restricted cellular interactions and molecular programs during follicle
maturation and rupture

Maturation and ovulation of ovarian follicles requires significant remodeling of the
surrounding stromal tissues and acute formation of rupture sites on the ovary's surface,
facilitating the release of oocytes for fertilization. We quantified the proportions of different
mesenchymal cell types and observed enrichment in the proportion of dividing mesenchymal
cells, first in immature ovaries and then at 6h post-hCG (**Supp Fig. 7A**). Visual examination of
dividing mesenchymal cells in the immature ovaries revealed their specific localization around
antral follicles enriched in proliferating granulosa cells, which suggested coordinated
proliferation of the mesenchyme and granulosa cells (**Fig. 5A and Supp Fig. 4B**). To understand
the molecular basis of this coordination, we explored cell-cell communication between the
dividing mesenchyme and mural granulosa cells of antral follicles in the immature ovaries
(**Methods**). This analysis revealed INHA/INHBA->TGFBR3, INHA/INHBB->ACVR1, and LAMA1/2-
>RPSA among the most significantly enriched ligand-receptor pairs (**Fig. 5B**). The enrichment in
the proportion of dividing mesenchyme at 6h post-hCG can be attributed to increased
vascularization of stromal cell layers surrounding the preovulatory follicles as suggested by an
increase in the proportion of endothelial cells as well as pericytes between 4h and 6h post-hCG
(**Supp Fig. 7**). Overall, these observations pinpoint tight coordination between mesenchyme and
mural granulosa cells during follicular maturation.

We identified three distinct time-dependent, mesenchyme-specific gene expression
signatures: ***i)*** a sharp initial increase in the expression of genes associated with neurotrophic
signaling from 0h to 4h post-hCG treatment (*Ntrk2*, *Bdnf*, and *Pdgfra); **ii)*** an intermediate
increase in the expression of genes associated with extracellular structure organization between
4h and 8h post-hCG treatment (*Postn*, *Mmp2*, *Eln*, *Col5a2*, and *Tgfbi); **iii)*** and a sharp, late-onset
increase in the expression of genes associated with myofibril assembly and muscle contraction
at 8h post-hCG treatment (*Myom1*, *Lmod1*, *Pdgfrb*, **Fig. 5C and 5D, Supp Fig. 7**). We visualized a
module score for genes associated with muscle contraction and found that this score is increased
in the mesenchyme of preovulatory follicles after hCG treatment (**Fig. 5E**). This observation is in
line with the reported role of muscle gene-expressing cells, in particular vascular smooth muscle
cells, for ovulation[36–38]. Collectively, these results revealed distinct activities of the ovarian
stroma during defined time windows preceding ovulation.

To gain new insight into follicle rupture and the transition to the luteal phase, we analyzed

the granulosa cell transcriptomes at 11h and 12h post-hCG. While follicles appeared similar on

histology immediately before follicle rupture at 12h post-hCG (**Supp. Fig. 1C**), we identified two

distinct types of large ovulating follicles based on transcriptomic signatures of their granulosa

cells (**Fig. 5F & 5G**). We asked whether these transcriptomic signatures are associated with the

fate of the follicle to ovulate. And indeed, granulosa cells in one type of periovulatory follicles

were enriched with transcripts of genes known to be crucial for ovulation, including *Edn2*, *Ptgs1*,

Junb, and *Adamts1*[39–41] (**Fig. 5H**). In contrast, transcript levels of these genes were low in the

other type of follicles, which share a similar transcriptomic signature with granulosa cells in

prematurely luteinized structures (lytic-like) (**Fig. 5G**). Multiplexed FISH of *Rhox8*, *Edn2*, and *Ltbp1*

(latent transforming growth factor binding protein 1; a novel gene associated with of the lytic-

like periovulatory follicles) confirmed that periovulatory follicles that express *Ltbp1* do not

express *Edn2* (**Fig. 5I**) Hence, the molecular heterogeneity of periovulatory follicles may

correspond to divergent follicle fates of ovulation versus premature luteinization. Interestingly,

luteinization marker *Lhcgr* increased at 12h post-hCG in both types of periovulatory follicles,

suggesting a transition stage right before ovulation (**Supp. Fig. 8A**). These findings highlight the

rapid and intricate processes governing follicular fate determination immediately preceding

ovulation.

Figure 5 | Spatially restricted molecular programs in ovarian stroma and periovulatory follicles. **A)**
Spatial transcriptomics map of immature ovaries with mesenchymal transcriptomes colored by predicted
mesenchymal cell types (left). Spatial transcriptomics map of immature ovaries colored by follicle type
predicted based on the granulosa cell phenotypes (right). Arrows point to dividing mesenchyme cells in
the untreated immature ovaries and asterisks on the follicle type map label the corresponding antral
follicles. **B)** Dot plot showing ligand-receptor interaction between dividing mesenchymal cells and
granulosa cells of antral follicles in immature ovaries. **C)** Line plot showing temporal trends of genes that
display a sharp, late onset increase in expression post-hCG in mesenchymal cells. **D)** Spatial
transcriptomics map showing the temporal expression of *Myom1* in immature and preovulatory ovaries.
E) Spatial transcriptomics map showing the temporal gene module score for muscle contraction in
immature and preovulatory ovaries. **F)** Spatial transcriptomics map of a preovulatory ovary at 12h post-
hCG, with granulosa cell types colored by their transcriptomes. **G)** Spatial transcriptomics map of a

preovulatory ovary at 12h post-hCG showing the expression of gene markers of mural granulosa cells (top) and lytic-like granulosa cells (bottom). **H)** Spatial transcriptomics map of a preovulatory ovary at 12h post-hCG showing the expression of follicle rupture-associated gene markers such as *Edn2*, *Ptgs1*, *Junb*, and *Adamts1*. **I)** Multiplexed RNA FISH staining for antral follicle associated granulosa cell marker *Rhox8* (yellow), lytic-like granulosa cell marker *Ltbp1* (cyan), and rupture-associated marker *Edn2* (magenta) in preovulatory follicles at 12h post-hCG. Representative images are from four biological replicates.

Discussion

We produced a spatiotemporal transcriptomic atlas using spatially resolved RNA-sequencing at near single-cell resolution to understand cell-type- and folliculogenesis-stage-specific mechanisms underlying mouse ovarian follicle maturation and ovulation. Our analysis revealed intricate spatial and molecular differentiation of distinct granulosa cell layers during the crucial preantral-to-antral transition[42,43]. We find that the transition toward the antral follicle state requires the coordinated expansion of *Inhbb+Kctd14+* cells in the center layers surrounding the oocytes, and of *Inhbb+Kctd14-* mural granulosa cells surrounding the double-positive cells. We identified *Ank1* as a potential early marker for cumulus cell fate at the preantral stage. While *Kctd14* is ubiquitous in preantral follicles, we found that its expression is suppressed in the outer layers of granulosa cells of antral follicles. In contrast, we observed that *Ank1* is induced in inner layers of granulosa cells. This sharp spatial demarcation in molecular marker expression suggests that interactions between oocyte-derived factors and the spatial location of granulosa cells relative to the basement membrane dictates their differentiation and fate.

We further delineated cell-type-specific interaction mechanisms of ovarian cells. The changes that occur in the ovarian stroma and mesenchymal cells are important for follicle growth and rupture but challenging to study *in vivo*. Using spatial transcriptomics, we observed a surge in the proliferative activity of mesenchymal cells in early antral follicles which also had high proliferative activity of granulosa cells. These coordinated proliferative activities are possibly mediated by the ovarian local effect of inhibins and laminins. We observed a subsequent burst of mesenchymal cell proliferation at around 6h post-hCG in parallel with an enrichment of genes regulating extracellular matrix remodeling and an increase in the number of endothelial cells and pericytes. Transcriptomic changes in cumulus cells and oocytes in response to ovulation stimulation have been studied extensively[44–46], but rarely without separating the two cell types *in vitro*, or at time points immediately preceding ovulation. We profiled oocyte-cumulus cell interaction mechanisms specific to defined post-hCG time points. Close to ovulation, we observed increased expression of *Nupr1*, *Spp1* and *S100a6* in cumulus cells, and an enrichment in the expression of genes that regulate RNA activity and DNA demethylation in oocytes.

In addition, we report gene regulatory programs at the follicle level both in rapid response
to ovulation stimulation, and in association with the process of follicle rupture and transition to
the luteal phase. Limited information was previously available on gene regulation during the first
hour after hCG simulation and the last hour before ovulation. We found that during the first hour
post-hCG, genes related to chromatin remodeling are upregulated, the expression of
steroidogenic genes is increased in both mural granulosa cells and theca cells, and an anti-
apoptotic program is induced, supporting the notion that coordinated cell survival and
steroidogenic programs respond rapidly to hormone stimulation. In the hour leading up to follicle
rupture, we observed genes such as *Akr1c18* and *Ldhd*, which modulate cell metabolism and
steroidogenesis, show the same trend of increase as a gene module score of muscle contraction
and genes associated with follicle rupture, including *Edn2* and *Adamts1*[12,24]. Immediately before
follicle rupture, we find that periovulatory follicles exhibit acute luteinization and divergent
heterogeneity, with a subpopulation of follicles poised to rupture. These observations highlight
the dynamic biological processes that occur at the final stage of follicle maturation immediately
before ovulation.

Given that regulators and pathways that are important for ovarian follicle maturation and
ovulation are highly conserved between humans and mice, our findings, and data promise to
translate into valuable insights for infertility management in humans.

METHODS

Ethical approval for animal experiments

All animal work was conducted ethically, conforming to the U.S. Public Health Service policy, and
was approved by the Institutional Animal Care and Use Committee at Cornell University (IACUC
approved protocol number 2019-0006). C57BL/6J mice were bred in-house for all experiments in
the study. Mice were housed in 11.5" x 7.5" IVC Polycarb Shoebox Cage for the duration of the
experiment. Temperature 68-77°F and humidity between 30 to 70% were maintained in the
rodent room. Lights were turned ON from 5AM and OFF at 7PM in the rodent room.

Hormone treatment for induction of ovulation

21-23 days old female mice with body weight of 10~11 grams were treated with pregnant mare
serum gonadotropin (PMSG) and human chorionic gonadotropin (hCG) to induce ovulation.
Untreated mice of the same age were used as immature controls. Mice were first administered
5 IU PMSG (BioVendor, RP1782725000) followed by 5 IU hCG 48h later (EMD Millipore
Corporation, #230734-1MG) in 100µl 0.1% BSA-PBS per mouse via intraperitoneal injection using
a 27G insulin syringe. Three immature and seven preovulatory mice across eight different time
points, namely untreated immature, 0h, 1h, 4h, 6h, 8h, 11h, and 12h post-hCG treatment, were
used for spatial transcriptomics experiments. To evaluate the response of mice to hormone
treatment, RNA extraction followed by reverse transcription (RT) and quantitative polymerase
chain reaction (qPCR) was performed on one ovary from every mouse for a panel of selected
genes. The other ovaries from the mice selected based on RT-qPCR results were used for spatial
transcriptomics experiments.

Sample preparation for Curio spatial transcriptomics

Whole ovaries were isolated using aseptic technique and placed in ice cold and sterile 1X
Phosphate Buffered Saline (1X PBS, without calcium and magnesium chloride; Gibco 10010023).
Blood and other contamination were carefully removed by washing the tissues with fresh 1X PBS.
Fresh tissues were immediately embedded in Optimal Cutting Compound (OCT) media (SAKURA
25608-930) and frozen directly on dry ice for spatial transcriptomics experiments. The tissue
blocks were cut into 10 µm sections using Thermo Scientific CryoStar NX50 cryostat and mounted
on a Seeker Tiles (Curio Bioscience) for spatial transcriptomics experiments.

Slide-seq spatial transcriptomics library preparation

Slide-seq spatial transcriptomics[47,48] experiment was performed using the Curio Seeker Kit (Curio
Bioscience) according to manufacturer's instructions. Briefly, tissue sections from fresh-frozen
murine ovaries were mounted on a 3mm x 3mm spatially indexed bead surface (Curio Seeker Kit,
Curio Bioscience). After RNA hybridization and reverse transcription, the tissue sections were
digested, and the beads were removed from the glass tile and resuspended. Second strand
synthesis was then performed by semi-random priming followed by cDNA amplification.

Sequencing libraries were then prepared using the Nextera XT DNA sample preparation kit. The
libraries were pooled and sequenced on an Illumina NextSeq 2K (P3 flow cell) using the 100-cycle
kit (Read 1 = 50 bp, Read 2 = 72 bp, Index 1 = 8 bp, and Index 2 = 8 bp). The sequencing data was
aligned to the mouse genome (assembly: GRCm38) using the STAR Solo[49] (version=2.7.9a)
pipeline to derive a feature x bead barcode expression matrix. Bead barcode location files for the
corresponding tiles were provided by Curio Bioscience.

Slide-seq data preprocessing for quality control and smear removal
Slide-seq count matrix and the position information for every bead barcode were loaded into an
AnnData object using scanpy[50] (v1.9.1). After filtering out the beads with less than 100 transcripts
detected, we log-normalized the Slide-seq expression data and computed principal components
using highly variable genes. The transcriptomes were then clustered using the Leiden clustering
algorithm and visualized on spatial maps. Cluster maps were then compared to Hematoxylin and
Eosin (H&E) stained images of consecutive tissue sections, and clusters representing beads not
located under the tissue were manually removed from individual Slide-seq datasets. Spatial
distances between all pairs of beads within individual samples were then used to filter out beads
with less than 10 beads within 100 µm distance. Individual samples were then reprocessed, and
an additional round of filtering was performed to remove clusters of beads not located under the
tissue. After filtering, individual datasets were processed and used for cell type labeling.

Multi-level cell type label assignment for spatial transcriptomics datasets
scRNA-seq dataset from cycling murine ovaries[3] was used as a reference to perform a multi-level
deconvolution of the spatial transcriptomics datasets. First, cell2location[51] (v0.1.3) was used to
deconvolve spatial transcriptomes and broad cell type labels were assigned to each bead. Genes
in the reference were filtered with cell_count_cutoff=5, cell_percent_cutoff=0.03, and
nonz_mean_cutoff=1.12 to select for highly expressed markers of rare cell types while removing
most uninformative genes. Cell type signatures were determined using NB regression and used
for spatial mapping of scRNA-seq broad cell type labels on Slide-seq data with hyperparameters
N_cells_per_location=1 and detection_alpha=20. Additional round of clustering was performed
on beads labeled as granulosa cells within individual datasets and differential gene expression
analysis in conjunction with observed spatial location within the tissue was used to label cumulus
cells and four distinct types of mural granulosa cells with distinct transcriptomic profiles that are
linked to follicle state (preantral, antral, atretic, and lytic follicles). Additionally, another round of
deconvolution using a scRNA-seq reference was performed to refine cell type labels for beads
classified as mesenchyme. Cell type signatures were determined using NB regression and used
for spatial mapping of scRNA-seq fine cell type labels on Slide-seq data with hyperparameters
N_cells_per_location=1 and detection_alpha=20. Fine cell type labels were assigned to

mesenchymal cells located in both cortical and medullary stroma of the ovary tissue sections. All
celltype labels were integrated together for visualization and analysis in the scanpy package.
525
Follicle segmentation and follicle state assignment in Slide-seq datasets
Individual Slide-seq datasets were read using squidpy[52] package (v1.2.3) and visualized in the
interactive Napari image viewer[53] (v0.4.15). Broad cell type labels assigned to individual beads
were visualized on spatial maps and polygon shapes were drawn in the mesenchyme separating
the follicles to manually segment 335 follicles across eight Slide-seq datasets. The beads within
individual segmented follicles were assigned a unique follicle-ID for follicle level analysis and
visualization. Each follicle was assigned a follicle state (preantral, antral, atretic, or lytic follicle)
based on the transcriptional phenotype of mural granulosa cell within that phenotype as labeled
during the multi-level cell type label assignment for spatial transcriptomics datasets.
535
Single-cell regulatory network inference for Slide-seq datasets
pySCENIC implementation (v0.12.1) of the SCENIC pipeline[54,55] (Single-Cell rEgulatory Network
Inference and Clustering) was used to infer transcription factors, gene regulatory networks for
Slide-seq data. Raw spatial gene expression counts for combined Slide-seq datasets were saved
in loom file format using the loompy package (v2.0.17). These counts were used for gene
regulatory network inference and generation of co-expression modules using pySCENIC package.
Inferred modules were used for regulon prediction aka cisTarget and cellular enrichment of
regulons (AUCell) was estimated for every bead in Slide-seq data. AUCell regolon enrichment
scores were compared between representing different stages of luteinizing hormone treatment
to estimate Regulon Specificity Scores (RSS) for specific stages/ conditions.
546
Temporal gene expression analysis and clustering for Slide-seq datasets
Custom scripts were used to study temporal trends in gene expression across stages of
preovulatory ovaries. In short, Generalized Additive Models (GAMs) were trained on expression
values of highly variable genes across discrete time points using the pyGAM package[56] (v0.8.0).
LinearGAM function as implemented in pyGAM package was used with n_splines=4 and
spine_order=2 to derive the GAM models for individual genes which were then used to predict
gene expression trends on the same eight discrete time points. Predicted gene expression values
for all genes were then scaled and clustered using K-means clustering with k from 4-6 as
implemented in the scikit-learn[57] package (v1.2.0). Scaled gene expression trends were then
visualized for temporally variable genes within gene groups. Genes of interest within individual
gene groups were visualized on spatial maps using scanpy. Pseudo-bulk granulosa mural cell
transcriptomes at a follicle resolution were used to study temporal changes in mural granulosa
cells. Raw gene expression counts for mural granulosa cells within each follicle were added to
construct pseudo-bulk granulosa mural transcriptomes.

Construction and analysis of pseudo-bulk oocyte transcriptomes

Bulk oocyte transcriptomes were constructed from spatial transcriptomics beads under individual oocyte cells. Beads classified as oocytes during deconvolution were extracted and filtered to remove beads with oocyte prediction score <= 0.5 and select the beads most likely to be in oocytes. Spatial distances between all pairs of beads under oocytes were then used to filter out beads with less than five beads within 100 μm distance. Individual samples were then reprocessed, and an additional round of filtering was performed to remove prediction noise. Raw gene expression counts for oocyte beads within each follicle were added to construct pseudo-bulk oocyte transcriptomes.

Cell-cell interactions analysis on spatial transcriptomics data

Cell-cell signaling analysis was performed using the ligand-receptor interaction analysis module of squidpy package. For oocytes and cumulus cells in preovulatory follicles, mean expression of annotated ligand-receptor interaction pairs from Omnipath database was calculated at follicle resolution using the squidpy implementation of the method CellPhoneDB[58]. Mean expression values of ligand receptor paris across follicles were then compared between different stages of ovulation to identify stage specific cell-cell interactions. For interactions between dividing mesenchyme and granulosa cells in the immature untreated ovaries, ligand receptor interaction analysis on all annotated pairs was performed on beads classified as granulosa and dividing mesenchyme.

Enrichment analysis for Gene Ontology (GO) for spatial transcriptomics

GO term enrichment analysis was performed on differentially expressed genes using gseapy[59,60] (version v1.0.4) wrapper package. Differential gene expression analysis results were used to select genes which were used for GO term enrichment analysis using GO_Biological_Processes_2021 gene sets in enrichr command. The enriched GO terms of interest were selected and visualized on a bar plot. The differentially expressed genes that were associated with GO terms of interest were used to calculate module scores using the score_genes command in scanpy. Module scores for individual beads then compared across different follicle types and across stages of ovulation.

Sample preparation for RNA fluorescence in-situ hybridization (FISH), immunofluorescence, and histology

Whole ovaries were isolated using aseptic technique and placed in ice cold sterile 1X PBS. Fresh tissues were immediately embedded in Optimal Cutting Compound (OCT) media and frozen in liquid nitrogen cooled isopentane, cut into 10 μm sections using a Thermo Scientific Microm 550

598 cryostat, and mounted on -20°C cooled histological glass slides which were then stored at -80°C
599 until used.
600
RNA fluorescence in-situ hybridization (FISH) split probe design and Signal Amplification using
Hybridization Chain Reaction HCR-V3
Two-step hybridization strategy with split probe design and Hybridization Chain Reaction (HCR)-
V3[61] was used to label up to three transcripts in a single tissue section. Split probes for individual
gene targets were designed using NCBI primer-blast[62]. The probes were checked for binding
specificity against the RefSeq mRNA database for *Mus musculus*. We designed 20-21 bp primer
pairs for an amplicon length of 40-42 bp (2 x primer length), primer melting temperature between
57°C and 63°C, and primer GC content between 35% and 65%. 5-10 pairs of reverse
complemented forward primers and reverse primers were then concatenated to the flanking
initiator sequence for HCR, ordered from Integrated DNA Technologies (IDT) with standard
desalting purification. Split probes for each gene target, mixed and diluted in nuclease-free water
to create a split probe pool stock solution at 10µM total probe concentration for every target.
Hairpin pairs labeled with three different fluorophores namely Alexa-488, Alexa-546, and Alexa-
647 (Molecular Instruments) were used for HCR-V3.
615
RNA fluorescence in-situ hybridization (FISH) experiments
Slides with tissue sections were then brought to room temperature until the OCT melts and were
then immediately fixed in 4% paraformaldehyde for 12 minutes at room temperature. Post
fixation, the sections were washed for 5 mins in 1x PBS twice, incubated for 1 hour in 70% ethanol
for tissue permeabilization, washed again for 5 mins in 1x PBS, and then used for primary
hybridization. Hybridization Buffer (HB) mix was prepared with 2x SSC, 5x of Denhart Solution,
10% Ethylene Carbonate, 10% Dextran Sulfate, 0.01% SDS, 1µM of probe pool mix per target for
the hybridization reaction. 20-30 µl of HB mix (with probes) per section was then put on each
slide to cover the tissue section, covered with parafilm, and incubated overnight at 37°C inside a
humidifying chamber for primary hybridization. After primary hybridization, parafilm was
removed and slides were washed in Hybridization Wash Buffer-1 (0.215M NaCl, 0.02M Tris HCl
pH 7.5, and 0.005M EDTA) for 20-30 minutes at 48°C. Amplification Buffer (AB) mix was prepared
with 2x SSC, 5x of Denhart Solution, 10% Dextran Sulfate, 0.01% SDS, 0.06µM of HCR hairpins for
the amplification reaction. 2µl of each fluorophore labeled hairpins at 3µM corresponding to the
target genes were mixed, incubated at 95°C for 1.5 minutes, covered in aluminum foil, and left
to cool down at room temperature for 30 minutes to form hairpins before adding it to AB mix.
20 µl of AB mix per section was then put on each slide to cover the tissue section, covered with
parafilm, and incubated overnight at room temperature in the dark for signal amplification. After
signal amplification, parafilm was removed, and slides were washed in the 5x SSCT buffer twice
for 30-40 minutes and then twice for 10 mins. The slides were then carefully cleaned with

Kimwipe and treated with Ready Probes Auto-fluorescence Quenching Reagent Mix (Thermo
Fisher, R37630) for 5 minutes and washed for 5 minutes in 1X PBS twice. Last, tissue sections
were counterstained with DAPI for 5 minutes at room temperature, washed for 5 minutes in 1x
PBS twice, excess PBS cleaned using Kimwipe, immediately mounted on coverslips using Slowfade
antifade media, left overnight for treatment, and imaged the next day on a Zeiss Axio Observer
Z1 Microscope using a Hamamatsu ORCA Fusion Gen III Scientific CMOS camera. smFISH images
were shading corrected, stitched, rotated, thresholded, and exported as TIFF files using Zen 3.1
software (Blue edition).

REFERENCES

1. Espey, L. L. & Richards, J. S. Temporal and spatial patterns of ovarian gene transcription following an ovulatory dose of gonadotropin in the rat. *Biol. Reprod.* **67**, 1662–1670 (2002).

2. Duffy, D. M., Ko, C., Jo, M., Brannstrom, M. & Curry, T. E. Ovulation: Parallels With Inflammatory Processes. *Endocr. Rev.* **40**, 369–416 (2019).

3. Morris, M. E. *et al.* A single-cell atlas of the cycling murine ovary. *eLife* **11**, e77239 (2022).

4. Park, C. J. *et al.* Progesterone Receptor Serves the Ovary as a Trigger of Ovulation and a Terminator of Inflammation. *Cell Rep.* **31**, 107496 (2020).

5. Li, S. *et al.* Single-cell RNA sequencing analysis of mouse follicular somatic cells†. *Biol. Reprod.* **105**, 1234–1245 (2021).

6. Stévant, I. *et al.* Dissecting Cell Lineage Specification and Sex Fate Determination in Gonadal Somatic Cells Using Single-Cell Transcriptomics. *Cell Rep.* **26**, 3272-3283.e3 (2019).

7. Mantri, M. *et al.* Spatiotemporal single-cell RNA sequencing of developing chicken hearts identifies interplay between cellular differentiation and morphogenesis. *Nat. Commun.* **12**, 1771 (2021).

8. Mantri, M. *et al.* Spatiotemporal transcriptomics reveals pathogenesis of viral myocarditis. *Nat. Cardiovasc. Res.* **1**, 946–960 (2022).

9. Shi, Y. *et al.* A spatiotemporal gene expression and cell atlases of the developing rat ovary. *Cell Prolif.* cpr.13516 (2023) doi:10.1111/cpr.13516.

10. Russ, J. E., Haywood, M. E., Lane, S. L., Schoolcraft, W. B. & Katz-Jaffe, M. G. Spatially resolved transcriptomic profiling of ovarian aging in mice. *iScience* **25**, 104819 (2022).

11. Ismail, P. M., Li, J., DeMayo, F. J., O'Malley, B. W. & Lydon, J. P. A novel LacZ reporter mouse reveals complex regulation of the progesterone receptor promoter during mammary gland development. *Mol. Endocrinol. Baltim. Md* **16**, 2475–2489 (2002).

12. Robker, R. L. *et al.* Progesterone-regulated genes in the ovulation process: ADAMTS-1 and cathepsin L proteases. *Proc. Natl. Acad. Sci. U. S. A.* **97**, 4689–4694 (2000).

13. Joyce, I. M., Pendola, F. L., O'Brien, M. & Eppig, J. J. Regulation of prostaglandin-endoperoxide synthase 2 messenger ribonucleic acid expression in mouse granulosa cells during ovulation. *Endocrinology* **142**, 3187–3197 (2001).

14. Wijgerde, M., Ooms, M., Hoogerbrugge, J. W. & Grootegoed, J. A. Hedgehog signaling in mouse ovary: Indian hedgehog and desert hedgehog from granulosa cells induce target gene expression in developing theca cells. *Endocrinology* **146**, 3558–3566 (2005).

15. Hirata, R., Hojo, T., Sano, M., Hayashi, N. & Okuda, K. Potential role of hCG in apoptosis of human luteinized granulosa cells. *J. Reprod. Dev.* **61**, 67–73 (2015).

16.   Svensson, E. C., Markström, E., Andersson, M. & Billig, H. Progesterone receptor-mediated inhibition of apoptosis in granulosa cells isolated from rats treated with human chorionic gonadotropin. *Biol. Reprod.* **63**, 1457–1464 (2000).

17.   Carletti, M. Z. & Christenson, L. K. Rapid effects of LH on gene expression in the mural granulosa cells of mouse periovulatory follicles. *Reprod. Camb. Engl.* **137**, 843–855 (2009).

18.   Li, Y. *et al.* Down-regulation of long non-coding RNA MALAT1 inhibits granulosa cell proliferation in endometriosis by up-regulating P21 via activation of the ERK/MAPK pathway. *Mol. Hum. Reprod.* **25**, 17–29 (2019).

19.   Sun, L., Zhang, P. & Lu, W. lncRNA MALAT1 Regulates Mouse Granulosa Cell Apoptosis and 17β-Estradiol Synthesis via Regulating miR-205/CREB1 Axis. *BioMed Res. Int.* **2021**, 6671814 (2021).

20.   LaVoie, H. A. Transcriptional control of genes mediating ovarian follicular growth, differentiation, and steroidogenesis in pigs. *Mol. Reprod. Dev.* **84**, 788–801 (2017).

21.   Carletti, M. Z. & Christenson, L. K. Rapid effects of LH on gene expression in the mural granulosa cells of mouse periovulatory follicles. *REPRODUCTION* **137**, 843–855 (2009).

22.   Zhang, M., Su, Y.-Q., Sugiura, K., Xia, G. & Eppig, J. J. Granulosa cell ligand NPPC and its receptor NPR2 maintain meiotic arrest in mouse oocytes. *Science* **330**, 366–369 (2010).

23.   Pangas, S. A., Jorgez, C. J. & Matzuk, M. M. Growth differentiation factor 9 regulates expression of the bone morphogenetic protein antagonist gremlin. *J. Biol. Chem.* **279**, 32281–32286 (2004).

24.   Ko, C. *et al.* Endothelin-2 in ovarian follicle rupture. *Endocrinology* **147**, 1770–1779 (2006).

25.   Cacioppo, J. A. *et al.* Granulosa cell endothelin-2 expression is fundamental for ovulatory follicle rupture. *Sci. Rep.* **7**, 817 (2017).

26.   Brown, H. M. *et al.* ADAMTS1 cleavage of versican mediates essential structural remodeling of the ovarian follicle and cumulus-oocyte matrix during ovulation in mice. *Biol. Reprod.* **83**, 549–557 (2010).

27.   Roche, J. *et al.* Apelin (APLN) and Apelin Receptor (APLNR) in Human Ovary: Expression, Signaling, and Regulation of Steroidogenesis in Primary Human Luteinized Granulosa Cells. *Biol. Reprod.* **95**, 104 (2016).

28.   Shokrollahi, B. *et al.* Apelin and Apelin Receptor in Follicular Granulosa Cells of Buffalo Ovaries: Expression and Regulation of Steroidogenesis. *Front. Endocrinol.* **13**, 844360 (2022).

29.   Roche, J. *et al.* Apelin (APLN) regulates progesterone secretion and oocyte maturation in bovine ovarian cells. *Reprod. Camb. Engl.* **153**, 589–603 (2017).

30.   Leu, S.-J., Lam, S. C.-T. & Lau, L. F. Pro-angiogenic Activities of CYR61 (CCN1) Mediated through Integrins αvβ3 and α6β1 in Human Umbilical Vein Endothelial Cells *. *J. Biol. Chem.* **277**, 46248–46255 (2002).

31.   Su, Y.-Q., Nyegaard, M., Overgaard, M. T., Qiao, J. & Giudice, L. C. Participation of mitogen-activated protein kinase in luteinizing hormone-induced differential regulation of steroidogenesis and steroidogenic gene expression in mural and cumulus granulosa cells of mouse preovulatory follicles. *Biol. Reprod.* **75**, 859–867 (2006).

32.   Stocco, C. O., Chedrese, J. & Deis, R. P. Luteal expression of cytochrome P450 side-chain cleavage, steroidogenic acute regulatory protein, 3beta-hydroxysteroid dehydrogenase, and 20alpha-hydroxysteroid dehydrogenase genes in late pregnant rats: effect of luteinizing hormone and RU486. *Biol. Reprod.* **65**, 1114–1119 (2001).

33.   Fan, R. *et al.* Effects of the FABP4 gene on steroid hormone secretion in goose ovarian granulosa cells. *Br. Poult. Sci.* **62**, 81–91 (2021).

34.   Russell, D. L. & Robker, R. L. Molecular mechanisms of ovulation: co-ordination through the cumulus complex. *Hum. Reprod. Update* **13**, 289–312 (2007).

35.   Varani, S. *et al.* Knockout of pentraxin 3, a downstream target of growth differentiation factor-9, causes female subfertility. *Mol. Endocrinol. Baltim. Md* **16**, 1154–1167 (2002).

36.   Migone, F. F. *et al.* In vivo imaging reveals an essential role of vasoconstriction in rupture of the ovarian follicle at ovulation. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 2294–2299 (2016).

37.   Ren, Y., Cowan, R. G., Harman, R. M. & Quirk, S. M. Dominant activation of the hedgehog signaling pathway in the ovary alters theca development and prevents ovulation. *Mol. Endocrinol. Baltim. Md* **23**, 711–723 (2009).

38.   Ren, Y., Cowan, R. G., Migone, F. F. & Quirk, S. M. Overactivation of hedgehog signaling alters development of the ovarian vasculature in mice. *Biol. Reprod.* **86**, 174 (2012).

39.   Lee-Thacker, S. *et al.* Core Binding Factor β Expression in Ovarian Granulosa Cells Is Essential for Female Fertility. *Endocrinology* **159**, 2094–2109 (2018).

40.   Choi, Y. *et al.* FOS, a Critical Downstream Mediator of PGR and EGF Signaling Necessary for Ovulatory Prostaglandins in the Human Ovary. *J. Clin. Endocrinol. Metab.* **103**, 4241–4252 (2018).

41.   Choi, Y., Jeon, H., Akin, J. W., Curry, T. E. & Jo, M. The FOS/AP-1 Regulates Metabolic Changes and Cholesterol Synthesis in Human Periovulatory Granulosa Cells. *Endocrinology* **162**, bqab127 (2021).

42.   Diaz, F. J., Wigglesworth, K. & Eppig, J. J. Oocytes are required for the preantral granulosa cell to cumulus cell transition in mice. *Dev. Biol.* **305**, 300–311 (2007).

43.   Wigglesworth, K., Lee, K.-B., Emori, C., Sugiura, K. & Eppig, J. J. Transcriptomic diversification of developing cumulus and mural granulosa cells in mouse ovarian follicles. *Biol. Reprod.* **92**, 23 (2015).

44.   Su, Y.-Q. *et al.* Selective degradation of transcripts during meiotic maturation of mouse oocytes. *Dev. Biol.* **302**, 104–117 (2007).

45.   Pan, H., O'brien, M. J., Wigglesworth, K., Eppig, J. J. & Schultz, R. M. Transcript profiling during mouse oocyte development and the effect of gonadotropin priming and development in vitro. *Dev. Biol.* **286**, 493–506 (2005).

46.   Hernandez-Gonzalez, I. *et al.* Gene expression profiles of cumulus cell oocyte complexes during ovulation reveal cumulus cells express neuronal and immune-related genes: does this expand their role in the ovulation process? *Mol. Endocrinol. Baltim. Md* **20**, 1300–1321 (2006).

47.   Rodriques, S. G. *et al.* Slide-seq: A scalable technology for measuring genome-wide expression at high spatial resolution. *Science* **363**, 1463–1467 (2019).

48.   Stickels, R. R. *et al.* Highly sensitive spatial transcriptomics at near-cellular resolution with Slide-seqV2. *Nat. Biotechnol.* **39**, 313–319 (2021).

49.   Kaminow, B., Yunusov, D. & Dobin, A. STARsolo: accurate, fast and versatile mapping/quantification of single-cell and single-nucleus RNA-seq data. 2021.05.05.442755 Preprint at https://doi.org/10.1101/2021.05.05.442755 (2021).

50.   Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene expression data analysis. *Genome Biol.* **19**, 15 (2018).

51.   Kleshchevnikov, V. *et al.* Cell2location maps fine-grained cell types in spatial transcriptomics. *Nat. Biotechnol.* **40**, 661–671 (2022).

52.   Palla, G. *et al.* Squidpy: a scalable framework for spatial omics analysis. *Nat. Methods* **19**, 171–178 (2022).

53.   Ahlers, J. *et al.* napari: a multi-dimensional image viewer for Python. (2023) doi:10.5281/ZENODO.3555620.

54.   Aibar, S. *et al.* SCENIC: single-cell regulatory network inference and clustering. *Nat. Methods* **14**, 1083–1086 (2017).

55.   Van De Sande, B. *et al.* A scalable SCENIC workflow for single-cell gene regulatory network analysis. *Nat. Protoc.* **15**, 2247–2276 (2020).

56.   Servén, D., Brummitt, C., Abedi, H., & Hlink. dswah/pyGAM: v0.8.0. (2018) doi:10.5281/ZENODO.1208723.

57.   Pedregosa, F. *et al.* Scikit-learn: Machine Learning in Python. *J. Mach. Learn. Res.* **12**, 2825–2830 (2011).

58.   Efremova, M., Vento-Tormo, M., Teichmann, S. A. & Vento-Tormo, R. CellPhoneDB: inferring cell–cell communication from combined expression of multi-subunit ligand–receptor complexes. *Nat. Protoc.* **15**, 1484–1506 (2020).

59.   Fang, Z., Liu, X. & Peltz, G. GSEApy: a comprehensive package for performing gene set enrichment analysis in Python. *Bioinformatics* **39**, btac757 (2023).

60.   Chen, E. Y. *et al.* Enrichr: interactive and collaborative HTML5 gene list enrichment analysis tool. *BMC Bioinformatics* **14**, 128 (2013).

61.   Choi, H. M. T. *et al.* Third-generation *in situ* hybridization chain reaction: multiplexed, quantitative, sensitive, versatile, robust. *Development* **145**, dev165753 (2018).

62.   Ye, J. *et al.* Primer-BLAST: a tool to design target-specific primers for polymerase chain reaction. *BMC Bioinformatics* **13**, 134 (2012).

DATA AVAILABILITY

The authors declare that all sequencing data supporting the findings of this study have been deposited in NCBI's Gene Expression Omnibus (GEO) with GEO series accession number GSE240271. Bead barcode location files matched to spatial transcriptomics datasets have been made publicly available on GitHub (https://github.com/madhavmantri/mouse_ovulation). All other data supporting the findings in this study are included in the main article and associated files. Source data for individual figures are provided with this paper.

CODE AVAILABILITY

Scripts to reproduce the analysis presented in this study have been deposited on GitHub (https://github.com/madhavmantri/mouse_ovulation).

ACKNOWLEDGEMENTS

We would like to thank Dr. Peter Schweitzer and the Cornell Genomics Center for help with sequencing assays and the Cornell Bioinformatics facility for assistance with bioinformatics. We also thank the members of the Ren and De Vlaminck labs for many valuable discussions. This work was supported by a seed grant provided by Cornell Center of Vertebrate Genomics (to Y.A.R. and I.D.V.) and R01HD109392 from NICHD (to Y.A.R).

AUTHOR CONTRIBUTIONS

M.M., Y.A.R., and I.D.V. designed the study. M.M. and H.H.Z. performed the animal experiments. M.M. and E.S. performed the spatial transcriptomics experiments. M.M. and E.S. analyzed the data. M.M., H.H.Z., and E.S. performed histology, RNA FISH, and analyzed the images. M.M., H.H.Z., E.S, Y.A.R., and I.D.V. wrote the manuscript. All authors provided feedback and comments.

COMPETING INTERESTS

The authors declare no competing interests.