DEBBIES to compare life history strategies across ectotherms

Authors
Isabel M. Smallegange, Sol Lucas

Affiliations
School of Natural and Environmental Sciences, Newcastle University, Newcastle upon Tyne, NE1 7RU, UK

corresponding author(s): Isabel Smallegange (Isabel.smallegange@newcastle.ac.uk)

Abstract
Demographic models are used to explore how life history traits structure life history strategies across species. This study presents the DEBBIES dataset that contains estimates of eight life history traits (length at birth, puberty and maximum length, maximum reproduction rate, fraction energy allocated to respiration versus reproduction, von Bertalanffy growth rate, mortality rates) for 185 ectotherm species. The dataset can be used to parameterise dynamic energy budget integral projection models (DEB-IPMs) to calculate key demographic quantities like population growth rate and demographic resilience, but also link to conservation status or biogeographical characteristics. Our technical validation shows a satisfactory agreement between observed and predicted longevity, generation time, age at maturity across all species. Compared to existing datasets, DEBBIES accommodates (i) easy cross-taxonomical comparisons, (ii) many data-deficient species, and (iii) population forecasts to novel conditions because DEB-IPMs include a mechanistic description of the trade-off between growth and reproduction. This dataset has the potential for biologists to unlock general predictions on ectotherm population responses from only a few key life history traits.

Background & Summary

Matrix population models (MPMs) and integral projection models (IPMs) provide the basis for exploring how variation in the demographic rates of survival, growth and reproduction fuels variation across species in life history traits (like the timing, intensity, frequency and duration of key demographic processes, such as longevity, generation time or degree of iteroparity) and in combinations of life history traits that form life history strategies (including pace of life and reproductive strategy)[1-3]. Life history traits and strategies calculated from these structured population models predict not only key demographic properties such as population growth rate and demographic resilience, but also have important connections to other disciplines like biogeography, evolutionary biology and conservation biology[4-6].

Important datasets exist that collate MPMs (COMPADRE[7]; COMADRE[8]) and IPMs (PADRINO[9]) for plants and animals, and even algae, fungi, bacteria, and viruses. Recent efforts have furthermore linked the latter datasets in a centralised meta-database of trait data (MOSAIC[10]) so they can be interrogated at the same time. While these datasets are valuable in improving both data access and data usability[11] and are used globally in networks like the Open Trait Network (https://opentraits.org/), the parameterisation of the structured population models that are within these datasets requires long-term individual-level data that are scored from birth till death. Yet, there are many organisms for which such data are not available, for example, because it is difficult to track individuals over their lifetime (e.g. micro-organisms, small (soil-dwelling) animals). Because such MPM and IPM datasets can form part of pipelines to develop, for example, essential biodiversity variables to observe and report global biodiversity change[12], it is pertinent to avoid any unwilling species bias. We thus critically need to include datasets of structured population models that can also accommodate more data-deficient species to have a taxonomically most balanced representation as possible.

One type of structured population model that does not require many long-term individual-level life history data is the dynamic energy budget (DEB) IPM[13]. To parameterise a DEB-IPM for a species, one requires eight life history traits to be estimated (traits include length measures, rates of growth, reproduction and mortality [Fig. 1]) to be able to predict survival, growth and reproduction for a simple life cycle (more complex life cycles would require more parameters)[13]. These traits are assumed to be fixed for a life cycle (Fig. 1); that is, length at birth and at puberty represent specific moments in the life cycle and are time-invariant; maximum length and maximum reproduction rate, in turn, are maximum values that are achieved under the most favourable circumstances a life cycle is in, and thus are also time-invariant. We assume that kappa remains constant over a life cycle because empirical evidence suggests so[14]. To estimate the von Bertalanffy growth rate and mortality parameters (Fig. 1) one does require repeated observations on individual growth or survival of individuals within a population. So far, only two small datasets of DEB-IPM parameters have been generated, comprising estimates of the eight life history traits for 13 species of marine megafauna[3] and 13 microorganisms[15]. Here, we introduce a much larger and taxonomically diverse dataset that we refer to as DEBBIES in which we compiled estimates of the eight life history traits for 185 ectotherm species. Ectotherms are taxonomically diverse and their growth and reproduction can be captured in simple energy budget models, like the one incorporated into the DEB-IPM. Also, more than 99% of species are ectotherms[16]; consequently, no biological prediction can be considered universal if it is does not include these organisms. We find in our technical validation that our model outputs exhibit good agreement with observations on key life history traits (age at maturity, longevity, generation time). The dataset can be used for a variety of different applications of eco-evolutionary studies (e.g.[15,17]) (Fig. 1). Because DEB-IPMs (unlike MPMs and other IPMs) across species are built from the same life history traits, they can readily be used for comparative studies of life

81   histories and population dynamics across a wide range of species, for which DEBBIES
82   provides all the necessary input data. Finally, because DEB-IPMs include a mechanistic
83   description of the trade-off between growth and reproduction, they are particularly suited to
84   create population forecasts to novel conditions (like those created by climate change)[13] for
85   the species currently listed in the dataset.
86
Methods
88
89   Because the main purpose of DEBBIES is to parameterise DEB-IPMs for further analysis (Fig.
90   1), we first give a brief description of what a DEB-IPM is, and how a species' eight life history
91   traits feed into a DEB-IPM, before explaining how we collated the life history data presented
92   in DEBBIES.
93
Brief description of a DEB-IPM
The three demographic rates of survival, growth, reproduction and the relationship between
parent and offspring size are captured in the DEB-IPM by four fundamental functions, which
describe the dynamics of a population comprising cohorts of females of different sizes[3,13]: (1)
the survival function, $S(L(t))$, describing the probability of surviving from time t to time t+1;
(2) the growth function, $G(L', L(t))$, describing the probability that an individual of body
length L at time t grows to length L' at $t + 1$, conditional on survival; (3) the reproduction
function, $R(L(t))$, giving the number of offspring produced between time t and $t + 1$ by an
individual of length L at time t; and (4) the parent-offspring function, $D(L', L(t))$, the latter
which describes the association between the body length of the parent L and offspring
length L' (i.e. to what extent does offspring size depend on parental size). Denoting the
number of females at time t by $N(L, t)$ means that the dynamics of the body length number
distribution from time t to t+1 can be written as:
107
108   $$N(L', t + 1) = \int_{\Omega} \left[D(L', L(t))R(L(t)) + G(L', L(t))S(L(t)) \right] N(L, t)dL \qquad (1)$$
109
110   where the closed interval Ω denotes the length domain.
111   The survival function $S(L(t))$ in equation (1) is the probability that an individual of
112   length L survives from time t to $t + 1$:
113
114   $$S(L(t)) = \begin{cases} e^{-\mu_j} & \text{for } L_b \leq L < L_p \ \& \ L \leq L_m E(Y)/\kappa \,, \\ e^{-\mu_a} & \text{for } L_p \leq L \leq L_m \ \& \ L \leq L_m E(Y)/\kappa \,, \\ 0 & \text{otherwise} \end{cases} \qquad (2)$$
115
where $E(Y)$ can range from zero (empty gut) to one (full gut). Individuals die from starvation
at a body length at which maintenance requirements exceed the total amount of assimilated
energy, which occurs when $L > L_m \cdot E(Y)/\kappa$ and hence, then, $S(L(t)) = 0$ (e.g., an
individual of size L_m will die of starvation if $E(Y) < \kappa$). Juveniles and adults often have different
mortality rates, and, thus, juveniles ($L_b \leq L < L_p$) that do not die of starvation (i.e. $L \leq L_m \cdot$
$E(Y)/\kappa$) have a mortality rate of μ_j, and adults ($L_p \leq L \leq L_m$) that do not die of starvation
(i.e. $L \leq L_m \cdot E(Y)/\kappa$) have a mortality rate of μ_a.
123   The demographic functions that describe growth and reproduction in the DEB-IPM
are derived from the Kooijman-Metz model[3,13,18]. This is is a simple version of the standard
model of Kooijman's DEB theory, but one that still fulfils the criteria for general explanatory
models for the energetics of individuals[19]. The Kooijman-Metz model assumes that individual
organisms are isomorphic (which means that body surface area and volume are proportional
to squared and cubed length, respectively). The rate at which an individual ingests food, I, is
assumed to be proportional to the maximum ingestion rate I_{max}, the current feeding level Y

and body surface area, and hence to the squared length of an organism: $I = I_{max}YL^2$.
Ingested food is assimilated with a constant efficiency ε. A constant fraction κ of assimilated
energy is allocated to respiration; this respiration energy equals $\kappa\varepsilon I_{max}YL^2$ and is used to
first cover maintenance costs, which are proportional to body volume following ξL^3 (ξ is the
proportionality constant relating maintenance energy requirements to cubed length), while
the remainder is allocated to somatic growth. The remaining fraction $1 - \kappa$ of assimilated
energy, the reproduction energy, is allocated to reproduction in case of adults and to the
development of reproductive organs in case of juveniles, and equals $(1 - \kappa)\varepsilon I_{max}YL^2$. This
means that, if an individual survives from time *t* to time *t*+1, it grows from length *L* to length
L' following a von Bertalanffy growth curve, $\frac{dL}{dt} = \dot{r}_B(L_m \cdot Y - L)$, where $\dot{r}_B$ is the von
Bertalanffy growth rate (here assumed to be constant across feeding levels, but can be
adjusted if necessary[13]) and $L_m = \kappa\varepsilon I_{max}/\xi$ is the maximum length under conditions of
unlimited resource. Both κ and I_{max} are assumed to be constant across experienced feeding
levels, and therefore L_m is also assumed constant.
Implicitly underlying the population-level model of equation (1), like in any IPM, is a
stochastic, individual-based model, in which individuals follow Markovian growth trajectories
that depend on an individual's current state[20]. This individual variability is in standard IPMs
modelled in the functions describing growth, $G(L', L(t))$, and the parent-offspring
association, $D(L', L(t))$ (see below), using a probability density distribution, typically
Gaussian[21]. In the DEB-IPM, this individual variability arises from how individuals experience
the environment; specifically, the experienced feeding level *Y* follows a Gaussian distribution
with mean *E(Y)* and standard deviation $\sigma(Y)$. It means that individuals within a cohort of
length *L* do not necessarily experience the same feeding level due to demographic
stochasticity (e.g. individuals, independently of each other, have good or bad luck in their
feeding experience). Taken together, this means that the function $G(L', L(t))$ is the
probability that an individual of body length *L* at time *t* grows to length *L'* at *t* + 1, conditional
on survival, and, following common practice[20-22], follows a Gaussian distribution:
157
158 $$G(L', L(t)) = \frac{1}{\sqrt{2\pi\sigma_L^2(L(t+1))}} e^{\frac{-(L'-E(L(t+1)))^2}{2\sigma_L^2(L(t+1))}} \tag{3}$$
159
160 with the growth realized by a cohort of individuals with length *L(t)* equalling
161
162 $$E\big(L(t+1)\big) = \begin{cases} L(t)e^{-\dot{r}_B} + \left(1 - e^{-\dot{r}_B}\right)L_m E(Y) & \text{for } L \leq L_m E(Y) \\ L(t) & \text{otherwise} \end{cases}, \tag{4}$$
163
164 and the variance in length at time *t* + 1 for a cohort of individuals of length *L* as
165
166 $$\sigma^2(L(t+1)) = \begin{cases} (1 - e^{-\dot{r}_B})^2 L_m^2 \sigma^2(Y) & \text{for } L \leq L_m E(Y) \\ 0 & \text{otherwise} \end{cases} \tag{5}$$
167
where $\sigma(Y)$ is the standard deviation of the expected feeding level.
If a surviving female is an adult, she also produces offspring. According to the
Kooijman-Metz model[18], reproduction, i.e. the number of offspring produced by an
individual of length *L* between time *t* and *t* + 1, equals $Y \cdot R_m \cdot L^2/L_m^2$. The parameter R_m is
the maximum reproduction rate of an individual of maximum length L_m. Note that R_m is
proportional to $(1 - \kappa)$[18], whereas L_m is proportional to κ; κ thus controls the trade-off
between energy allocation to reproduction versus growth. However, the role of κ in the DEB-
IPM is mostly implicit, as κ is used as input parameter only in the starvation condition (see
below), whereas R_m and L_m are measured directly from data. Like L_m, R_m is also proportional

to I_{max}; since both κ and I_{max} are assumed to be constant across experienced feeding levels,
R_m is also assumed constant. The reproduction function $R(L(t))$ gives the number of
offspring produced between time t and $t + 1$ by an individual of length L at time t:
180

$$R(L(t)) = \begin{cases} 0 & \text{for } L_b \leq L < L_p \\ E(Y)R_mL(t)^2/L_m^2 & \text{for } L_p \leq L \leq L_mE(Y) \\ \dfrac{R_m}{1-\kappa}\left[E(Y)L(t)^2 - \dfrac{\kappa L(t)^3}{L_m}\right] & \text{for } L_mE(Y) < L \leq L_mE(Y)/\kappa \end{cases} \quad (6)$$

Individuals are mature when they reach puberty at body length L_p and only surviving adults
reproduce (equation (1)); thus, only individuals within a cohort of length $L_p \leq L \leq L_mY/\kappa$
reproduce.
The probability density function $D(L', L(t))$ gives the probability that the offspring of an
individual of body length L are of length L' at time $t + 1$, and hence describes the association
between parent and offspring character values:
188

189     $$D\big(L', L(t)\big) = \begin{cases} 0 & \text{for } L < L_p \\ \dfrac{1}{\sqrt{2\pi\sigma_{L_b}^2(L(t))}}e^{\frac{-(L'-E_{L_b}(L(t))^2}{2\sigma_{L_b}^2(L(t))}} & \text{otherwise} \end{cases} \quad (7)$$

190
where $E_{L_b}\big(L(t)\big)$ is the expected size of offspring produced by a cohort of individuals with
length $L(t)$, and $\sigma_{L_b}^2(L(t))$ the associated variance. For simplicity, we set $E_{L_b}\big(L(t)\big)$ constant
and assumed its associated variance, $\sigma_{L_b}^2\big(L(t)\big)$, to be very small.
Finally, note that the DEB-IPM assumes no effect of temperature on fundamental
functions. Temperature effects on fundamental functions, however, can be approximated by
varying experienced feeding level. Alternatively, one could resort to a more detailed and
more parameter-rich DEB-IPM that links individuals' size- and temperature-dependent
consumption and maintenance via somatic growth, reproduction, and size-dependent
energy allocation to emergent population responses[23], but this is not linked to DEBBIES.
200
Collection of life history trait data required to parameterise a DEB-IPM
Running a DEB-IPM for a species requires estimates for eight life history traits: the fraction
respiration energy κ, length at birth L_b, length at puberty L_p, maximum length L_m, juvenile
mortality rate μ_j, adult mortality rate μ_a, von Bertalanffy growth rate $\dot{r}_B$ and maximum
reproduction rate R_m (Fig. 1).
For the elasmobranchs, we obtained von Bertalanffy growth rate values using the
following search, in order of priority: (i) primary literature, using female growth curve,
measured empirically using data from animals, (ii) from Froese[24] supplementary material, or
(iii) using the equation $\dot{r}_B = -\log(1 - 0.95)/t_{max}$, where t_{max} is a species' longevity
(years), sourced from Fishbase[25]. If multiple values were available, only those in the highest
priority group were kept. If there were still multiple values, their median value was used. Any
values listed as 'Questionable' on Fishbase[25] were not used. Body lengths for 152 species
were sourced from Sharks of the World[26] or Rays of the World[27]. Total lengths were
converted to fork lengths using scalar values on Fishbase[26]. Length at birth was sourced from
the IUCN red list[28] for five species (*M. ambigua, M. birostris, A. parmifera, D. trachyderma*
and *R. australiae*). Length at puberty for *Maculabatis ambigua* was sourced from primary
literature[29]. We calculated maximum reproduction rate R_m as $R_m = (c \times n)/i$, where c is
the mean clutch size, n is the mean number of litters produced per year, and i is the
remigration interval, which is the minimal number of years between reproductive seasons.

Minimum and maximum pup numbers were sourced from Sharks of the World[26], Rays of the
World[27], IUCN red list[28], Fishbase[25], or Barrowclift et al.[30]. The sources for maximum and
minimum pup numbers were not prioritised, and therefore some maximums and minimums
were obtained from different sources. Breeding intervals were found in the same way,
except for *C. granulosus*, *S. californica*, *F. macki* and *R. porosus,* which data we sourced from
primary literature[31-34]. If the number of pups or breeding interval were not found, we took
those data from the next closest species within the same genus. Mode of reproduction was
sourced from the Sharks of the World[26] and Rays of the World[27]. Adult mortality rate was
calculated[35] by taking the inverse of the mean of longevity and age at maturity (*a*) ($\mu_a =$

$1/[(t_{max} + a)/2]$. Juvenile mortality rate[36] was calculated as $\mu_j = -\log{(l_\alpha^{\frac{1}{a}})}$, where
survival to maturity[37], l_α, equals $l_\alpha = e^{-\mu_a(t_{max}-a)}$.
Values for all other species were taken directly from the primary literature. Finally,
for most species, we were able to take values for κ from the Add My Pet database[38]. If no
values for κ were available for a species, we assumed κ = 0.8 as is common practice[39]. The
parameter kappa is explicit in the starvation condition[13], which states that individuals of
length *L* die of starvation when they can no longer cover their maintenance costs, which
occurs when $L = L_m E(Y)\kappa = L_{\text{starvation}}$. At any feeding level, the ultimate length that
individuals can grow to equals $L_\infty = L_m E(Y)$. Substituting L_∞ into the starvation condition
returns: $L_{\text{starvation}} = L_\infty/\kappa$. In our case, $\kappa = 0.8$ so that $L_{\text{starvation}} = 1.25 L_\infty$. This means that
at any constant feeding level, individuals will never reach the length at which they starve
because that is 25% larger than the ultimate length they can achieve at that feeding level.
Only when feeding level varies over time, like in a stochastic time series, can κ affect
population responses if, for example, individuals that were large in a good feeding
environment suddenly find themselves in a poor feeding environment.
244
Data Records
246
The DEBBIES dataset (Version 5) includes eight life history trait records for 185 ectotherm
species that were sourced from the scientific literature and are stored in one csv file with
accompanying metadata text file on FigShare[40]. Each row in the file represents one species
record. For most species, we sourced life history traits from different studies as rarely only
one single study provided estimates for all traits for one species. Each data record therefore
describes a species' general life history and is not specific to a particular population. On the
one hand, this approach is in line with the assumption of DEB theory that individual-level
differences are sufficiently small to take mean values to represent species-specific
parameters[41]. On the other hand, recent work on different populations of Trinidadian
guppies cautions against combining life history data from different studies, because
systematic bias during parameter estimation can generate substantial variation, and similar
patterns of growth and reproduction can be produced with very different parameter sets[42].
In contrast, focusing only on parameter sets specific to a particular population of Trinidadian
guppies returns life history variation predictions that are in line with observations[42]. Users
should thus carefully interpret their results, especially when predictions are not in line with
expectations. One could, for example, explore the robustness of a DEB-IPM to uncertainty or
perturbation of the input data to assess how much the data can be changed before any
desired property of the model, like predicted population growth rate, is lost[43,44]. The columns
of the data file are described in Table 1.
266
Technical Validation
268
Two model performance tests on a subset of DEBBIES that have validated the reliability of
the dataset are presented elsewhere[3,13]. Here, we conducted a similar model performance

test on the full dataset by exploring the distribution of predicted population growth rates,
and by comparing predicted and observed generation time (years), age at maturity (years),
and longevity (the sum of age at maturity and mature life expectancy; years) for three
different feeding levels. Because not all species were represented at all feeding levels (less
than half of all species in the dataset can persist at low feeding levels), any effect of feeding
level cannot be estimated independently. Therefore, we ran the model performance tests
separately for each feeding level. Predicted values of the latter quantities are calculated as
explained below in the usage notes (see also Fig. 1: derived life history traits). For ray-finned
fish, observed values were obtained from Fishbase[25] and, if unavailable, we obtained
generation times from the IUCN Red List[28] and age at maturity values from the Animal
Diversity Web[45]. For the cartilaginous fish, we obtained generation times (for all species) and
longevities (18 species) from the IUCN Red List[28], and longevity and age at maturity values
from the Sharks of the World book[26], the Rays of the World book[27] or from Barrowclift et
al.[30]. For one cartilaginous fish, we obtained the longevity value directly from the scientific
literature (silvertip shark *Carcharhinus albimarginatus*)[47] and for the Galapagos shark
Carcharhinus galapagensis from Fishbase[25]. For all other species, we first consulted the
Animal Diversity Web[45], then the IUCN Red List[28] for observations on generation time and
AnAge[47] for observations on age at maturity and/or longevity if these were unavailable in the
Animal Diversity Web[45]. If, for a particular quantity, a value range was given, we took the
median; if a series of observations was given, we took the mean.
Our validation shows that the predicted population growth rate λ (calculated as the
dominant eigenvalue of the matrix approximation of equation (1), see also Table 3) most
species are slightly higher than $\lambda = 1$ (denoting population increase) at high experienced
feeding level ($E(Y) = 0.9$), centred around $\lambda = 1$ (denoting stability) at an intermediate feeding
level ($E(Y) = 0.7$), and mostly lower than $\lambda = 1$ (denoting population decline) at low feeding
level ($E(Y) = 0.5$) (Fig. 2a). This is in line with the general expectation in ecology that
populations increase under favourable conditions but decline when conditions deteriorate.
Predicted generation times were higher than observed generation times at the lower
experienced feeding levels but not significantly different from observed generation times at
the highest experienced feeding level (Table 2; Fig. 2b). Using the root mean square error
(RMSE) (Table 2), we can quantify the overall deviation. Specifically, we estimate that 95% of
the observed generation time values fall within a range that extends 19 − 23 years from the
predicted generation times across feeding levels (assuming the residuals follow a Normal
distribution, 95% of observed values fall within $\pm 2 \times$ RMSE from the predicted values, and we
took the lowest and highest RMSE value from across the range of feeding levels to estimate
this 95% confidence interval). The average RMSE value for generation time across the three
feeding levels equals 10 years (Table 2). Given the fact that the highest observed generation
time is 53 years (*Carcharodon carcharias*), the average RMSE of 10 years indicates that the
model predictions have an average error rate of 19% (10 ÷ 53 = 0.19). We surmise that one
reason for this relatively large error rate is that predicted generation time is, following
convention [1], calculated as $T = \log(R_0)\,/\log(\lambda)$, where R_0 is the net reproductive rate and
λ the population growth rate. Calculated this way, generation time represents the time it
takes a population to increase by a factor R_0, which might not always be a good
approximation of how generation time is measured in the field.
Predicted longevity was lower than observed longevity at the lower experienced
feeding levels but not significantly different from observed values at the highest experienced
feeding level (Table 2; Fig. 2c). Using the root mean square error (RMSE) (Table 2), we
estimate that 95% of the observed longevity values fall within a range that extends 12 − 43
years from the predicted longevities across feeding levels. The average RMSE value for
longevity across the three feeding levels equals 13 years (Table 2). Given the fact that the
highest observed longevity is 100 years (*Squalus suckleyi*), the average RMSE of 13 years
indicates that the model predictions have an average error rate of 13% (13 ÷ 100 = 0.13).

Finally, predicted age at maturity was higher than observed age at maturity and the
best fit with observed values was at the lowest experienced feeding level (Table 2; Fig. 2d).
We are unsure why this is the case, but it most likely indicates a mismatch between the
functional biology of maturation, and the assumptions underlying the calculations of age at
maturity[48]. Using the root mean square error (RMSE) (Table 2), we estimate that 95% of the
observed age at maturity values fall within a range that extends 11 – 13 years from the
predicted ages at maturity across feeding levels. The average RMSE value for longevity
across the three feeding levels equals 6 years (Table 2). Given the fact that the highest
observed longevity is 39 years (*Chelonia mydas*), the average RMSE of 13 years indicates that
the model predictions have an average error rate of 15% ($6 \div 39 = 0.15$).

In summary, the model predicts generation times and longevity values accurately at
high experienced feeding levels, but predicted generation times showed the highest average
error rate. Predicted age at maturity values were significantly higher than observed values.
However, their error rate was overall lower than the error rate of predicted generation times
and only slightly higher than the error rate of predicted longevity values. Depending on the
specific question a user is interested in, these error rates can be acceptable or not. What
gives us confidence in the technical quality of the dataset and its potential applications, is
the fact that predicted population growth rates are within the range that we would expect
biologically (Fig. 1a).

Usage Notes

This data descriptor was peer reviewed in 2023 based on version 5 of the DEBBIES dataset[40].
All versions are available online[40] as described as in Table 1. Version 1 contains data on 47
species; some of these entries were incorrect and removed in version 2. In version 3, 157
species of elasmobranchs were added, with three more elasmobranchs added in version 4. In
the current version 5, we changed the symbols in the dataset that denote κ and $\dot{r}_B$ to match
the symbols used in this data descriptor. We anticipate future versions to include more
species.

The main purpose of the DEBBIES dataset is to be able to parameterise a DEB-IPM
for each of the species in the dataset, but the dataset on its own can also feed into other
databases (Fig. 1). Once parameterised, running a DEB-IPM requires setting a value for the
experienced feeding level $E(Y)$, which value can range from zero (empty gut) to one (full gut).
We find that for many species, $E(Y)$ should be higher than about 0.7 for the model to run,
although it can run for some species at much lower values, down to 0.4[13]. A feeding level of
around 0.7 can be considered to represent a gut that is 'just filled', on a scale between
empty and bursting[49]. In that sense, we should perhaps not be surprised DEB-IPMs require a
minimum experienced feeding level of around 0.7.

Many different quantities can be calculated from a parameterised DEB-IPM. For
example, like current MPM and IPM datasets[7-9], DEB-IPMs can be used to calculate key
demographic quantities such as population growth rate and demographic resilience, but also
nine key derived life history traits that inform on a species' turnover rate, longevity, growth
and reproduction (Fig. 1) (Table 3)[1]. Additionally, because all DEB-IPMs have the same
structure, one can run perturbation analyses to estimate for each species the proportional
change in the population growth rate for a proportional change in each of the input life
history traits (κ, L_b, L_p, L_m, μ_j, μ_a, $\dot{r}_B$, R_m). These results can be used to pinpoint those parts of
an organism's life history that should be the focus of monitoring or research to inform
conservation management, or those life history traits that contribute most to fitness. We
provide MatLab code to calculate the aforementioned quantities (see Code Availability
below). Some users, however, might be more interested in exploring the dynamics of single
species, rather than performing cross-taxonomical analyses. For those, we have included

code to explore how the aforementioned quantities vary with experienced feeding level $E(Y)$
for a particular species (see Code Availability below).
The nine derived life history traits[1] (Fig. 1, Table 3) can be used for more in-depth
cross-taxonomical analyses (Fig. 1). For example, statistical analyses (e.g., a phylogenetically
corrected principal component analysis[50] can be used to explore how life history variation
can be structured and summarised into a few life history strategies[51]. Previous studies using
MPM databases have revealed that plant and animal life history variation is predominantly
structured along a fast-slow life history speed axis (rapid growth, high reproduction and
short lifespan versus slow growth, low reproduction but long lifespan) and a reproductive
strategy axis (concentrated versus more dispersed reproduction events across the adult
lifespan)[1-2]. An interesting exercise would be to see if the same structuring holds when
individual growth and reproduction rates are described mechanistically using an energy
budget model (like in DEB-IPMs) instead of being estimated from observational data (like in
MPMs). What is more, because a DEB-IPM is run for a specific feeding level set by the user,
one could also investigate if the same structuring applies across a range of feeding levels (but
note that our validation only covers feeding levels between 0.5-0.9). Finally, if species can be
ranked along one or two life history strategy axes, their position along these axes can be
linked to quantities from other disciplines (Fig. 1). This opens op possibilities for users to
explore the potential for life history strategies to, for example, predict the likelihood of
extinction or invasion (e.g.[4]) investigate to what extent such strategies map onto
phylogenetic trees (traitgrams) to answer questions about their evolution (e.g.[6]).
Finally, users should be aware that, for the species currently included in DEBBIES, the
projection interval between time t and time $t + 1$ is set at 1 year; all rates are therefore
expressed per year and all durations are expressed in years. Likewise, all length
measurements in DEBBIES are currently expressed in centimetres. Future versions, however,
could include species which demographic rates are better projected over shorter time
intervals and which lengths are better expressed in millimetres or even micrometres (e.g.,
microorganisms[6]).
402

Code Availability

MatLab code and instructions in a README text file for cross-taxonomical analysis can be
downloaded from FigShare (https://doi.org/10.6084/m9.figshare.13241972; folder:
'Derived life history traits across ectotherms'[40]) to calculate (i) generation time, survivorship
curve, age at sexual maturity, progressive growth, retrogressive growth, mean sexual
reproduction, degree of iteroparity, net reproductive rate, and mature life expectancy (Table
3), as well as population growth rate and demographic resilience (damping ratio), and (ii) run
an elasticity analysis for each species listed in DEBBIES for a predefined, experienced feeding
level.
For users interested in exploring the dynamics of single species, MatLab code and
instructions in a README text file can be downloaded from FigShare
(https://doi.org/10.6084/m9.figshare.13241972; folder: 'Derived life history traits single
ecotherm'[40]) to calculate (i) generation time, survivorship curve, age at sexual maturity,
progressive growth, retrogressive growth, mean sexual reproduction, degree of iteroparity,
net reproductive rate, and mature life expectancy (Table 3), as well as population growth
rate and demographic resilience (damping ratio), and (ii) run an elasticity analysis for a single
species for a range of experienced feeding levels defined by the user.
420

Acknowledgements

422   The authors would like to acknowledge Gemma Crawford, Naomi Eeltink, Tom Hopman and
423   Iris van Rijn for their help in collecting some of the data as part of their studies, and thank
424   Amy Attle and Burkeleigh Boyd for testing the code as part of their studies. For the purpose

425 of Open Access, the author has applied a CC BY public copyright licence to any Author
426 Accepted Manuscript (AAM) version arising from this submission.
427

Author contributions
429 Isabel Smallegange - project conception – data acquisition – data validation – writing.
430 Sol Lucas - data acquisition – data validation – proof reading.
431

Competing interests
433 The authors declare no competing interests.
434

References
436

1. Salguero-Gómez, R., et al. Fast–slow continuum and reproductive strategies structure
plant life-history variation worldwide. *Proc. Natl Acad. Sci. U.S.A.* **113**, 230–235
(2016).
2. Paniw, M., Ozgul, A., & Salguero-Gómez, R. Interactive life-history traits predict
sensitivity of plants and animals to temporal autocorrelation. *Ecol. Lett.* **21**, 275–286
(2018).
3. Smallegange, I. M., Flotats Avilés, M., Eustache, K. Unusually paced life history
strategies of marine megafauna drive atypical sensitivities to environmental
variability. *Front. Mar. Sci.* **7**, 597492 (2020).
4. Salguero-Gómez, R. Applications of the fast–slow continuum and reproductive
strategy framework of plant life histories. *New Phytol.* **213**, 1618–1624 (2017).
5. Capdevila, P., et al. Longevity, body dimension and reproductive mode drive
differences in aquatic versus terrestrial life history strategies. *Funct. Ecol.* **34**, 1613–
1625 (2020).
6. Romeijn, J., Smallegange, I. M. Exploring how the fast-slow pace of life continuum and
reproductive strategies structure microorganism life history variation. Preprint at
https://doi.org/10.1101/2022.11.28.517963 (2022).
7. Salguero-Gómez, R., et al. The COMPADRE Plant Matrix Database: an online
repository for plant population dynamics. *J. Ecol.* **103**, 202-218 (2014).
8. Salguero-Gómez, R., et al. COMADRE: a global database of animal demography. *J.
Anim. Ecol.* **85**, 371-384 (2016).
9. Levin, S., et al. PADRINO. *Zenodo* v0.0.1. https://doi.org/10.5281/zenodo.6573870
(2022).
10. Bernard, C., et al. MOSAIC: a unified trait database to complement structured
population models. *Sci. Data* **10**, 335 (2023).
11. Gallagher, R. V., et al. Open Science principles for accelerating trait-based science
across the Tree of Life. *Nature Ecol. Evol.* **4**, 294–303 (2020).
12. Kissling, W. D., et al.. Towards global data products of Essential Biodiversity Variables
on species traits. *Nature Ecol. Evol.* **2**, 1531-1540 (2018).
13. Smallegange, I. M., Caswell, H., Toorians, M. E. M., & de Roos, A. M. Mechanistic
description of population dynamics using dynamic energy budget theory incorporated
into integral projection models. *Methods Ecol. Evol.* **8**, 146-154 (2017).
14. Augustine, S., & Kooijman, S. A. L. M. A new phase in DEB research. *J Sea Res.* **143**, 1-7
(2019).
15. Romeijn, J., & Smallegange, I. M. Exploring how the fast-slow pace of life continuum
and reproductive strategies structure microorganism life history variation. Preprint at
https://doi.org/10.1101/2022.11.28.517963 (2022).
16. Atkinson, D., & Sibly, R. M. Why are organisms usually bigger in colder environments?
Making sense of a life history puzzle. *Trends Ecol. Evol.* **12**, 235-239 (1997).

17. Rademaker, M., van Leeuwen, A., & Smallegange, I. M. Why we should not necessarily
expect life history strategies to inform on sensitivity to environmental change.
Preprint at https://doi.org/10.22541/au.164848872.26565315/v1 (2022).
18. Kooijman, S. A. L. M., & Metz, J. A. J. On the dynamics of chemically stressed
populations: The deduction of population consequences from effects on individuals.
Ecotoxicol. Environ. Saf. **8**, 254–274 (1984).
19. Sousa, T., Domingos, T., Poggiale, J. C., & Kooijman, S. A. L. M. Dynamic energy budget
theory restores coherence in biology. *Philos. Trans. R. Soc. Lond. B* **365**, 3413–3428
(2010).
20. Easterling, M. R., Ellner, S. P., & Dixon, P. M. Size-specific sensitivity: Applying a new
structured population model. *Ecology* **81**, 694–708 (2000).
21. Coulson, T. Integral projections models, their construction and use in posing
hypotheses in ecology. *Oikos* **121**, 1337–1350 (2012).
22. Merow, C., et al. Advancing population ecology with integral projection models: a
practical guide. *Methods Ecol. Evol*. **5**, 99-110 (2014).
23. Thunell V., Gårdmark, A., Vindenes, Y. Optimal energy allocation trade-off driven by
size-dependent physiological and demographic responses to warming. *Ecology* **4**,
e3967 (2023).
24. Froese, R. Estimating somatic growth of fishes from maximum age or maturity. *Acta
Ichthyol. Piscat.* **52**, 125-133 (2022).
25. Froese, R., & Pauly, D. Editors. FishBase. World Wide Web electronic publication.
www.fishbase.org, (02/2023).
26. Ebert, D. A., Dando, M., & Fowler, S. *Sharks of the World. A complete guide* (Princeton
University Press, 2021).
27. Last, P., et al. *Rays of the World* (CSIRO Publishing, 2016).
28. IUCN. The IUCN Red List of Threatened Species. Version 2022-2.
https://www.iucnredlist.org. Accessed on [19 May 2023].
29. Temple, A. J., et al. Life-history, exploitation and extinction risk of the data-poor Baraka's
whipray (*Maculabatis ambigua*) in small-scale tropical fisheries. *J. Fish Biol.* **97**, 708-719
(2020).
30. Barrowclift, E., et al. Tropical rays are intrinsically more sensitive to overfishing than
the temperate skates. *Biol. Cons.* **281**, 110003 (2023).
31. Guallart, J., & Vicent, J. J. Changes in composition during embryo development of the
gulper shark, *Centrophorus granulosus* (Elasmobranchii, Centrophoridae): an
assessment of maternal-embryonic nutritional relationships. *Environ. Biol. Fishes* **61**,
135-150 (2001).
32. Natanson, L. J., & Cailliet, G. M. Reproduction and development of the Pacific angel
shark, *Squatina californica*, off Santa Barbara, California. *Copeia* **4**, 987-994 (1986).
33. Simpfendorfer, C. A., & Unsworth, P. Reproductive biology of the whiskery shark,
Furgaleus macki, off south-western Australia. *Mar. Freshwater Res*. **49**, 687-793
(1998).
34. Mattos, S. M., Broadhurst, M., Hazin, F. H., & Jones, D. M. Reproductive biology of the
Caribbean sharpnose shark, *Rhizoprionodon porosus*, from northern Brazil. *Mar.
Freshwater Res.* **52**, 745-752 (2001).
35. Pardo, S. A., Kindsvater, H. K., Reynolds, J. D., & Dulvy, N. K. Maximum intrinsic rate of
population increase in sharks, rays, and chimaeras: the importance of survival to
maturity. *Can. J. Fish. Aquat. Sci.*, **73**, 1159-1163 (2016).
36. Smith, S. E., Au, D. W., & Show, C. Intrinsic rebound potentials of 26 species of Pacific
sharks. *Mar. Freshwater Res.* **49**, 663-678 (1998).
37. Dulvy, N. K., et al. Methods of assessing extinction risk in marine fishes. *Fish Fish.* **5**,
255-276 (2004).

38. Add-my-pet. *Database of code, data and DEB model parameters* (www.debtheory.org)
(2023)
39. Kooijman, S. A. L. M. *Dynamic energy budget theory for metabolic organization*
(Cambridge, UK: Cambridge University Press 2010).
40. Smallegange, I. M. DEBBIES. A database to compare life history strategies across
ectotherms. *Figshare*. Dataset. https://doi.org/10.6084/m9.figshare.13241972 (2020).
41. Marques, G. M., et al. The AmP project: Comparing species on the basis of dynamic
energy budget parameters. PLoS Computational Biology, 14(5), e1006100 (2018).
42. Pottor, T., Reznick, D. N., & Coulson, T. Substantial intraspecific variation in energy
budgets: Biology or artifact? *Funct. Ecology* **35**, 1693-1707 (2021).
43. Hodgson, D. J., & Townley, S. Linking management changes to population dynamic
responses: the transfer function of a projection matrix perturbation. *J Applied Ecol.*
41, 1155-1161 (2004).
44. Deines A., et al. Robust population management under uncertainty for structured
population models. *Oikos* **17**, 2175-2183 (2007).
45. Myers, P., et al. *The Animal Diversity Web* (online). Accessed at
https://animaldiversity.org (2023).
46. Smart, J. J., et al. Stochastic demographic analyses of the silvertip shark (*Carcharhinus*
albimarginatus) and the common blacktip shark (*Carcharhinus limbatus*) from the
Indo-Pacific. *Fish. Res.* **191**, 95-107 (2017).
47. de Magalhaes, J. P., & Costa, J. A database of vertebrate longevity records and their
relation to other life-history traits. *J. Evol. Biol.* **22**, 1770-1774 (2009).
48. Caswell, H. Sensitivity Analysis: Matrix Methods in Demography and Ecology.
Demographic Research Monographs. Springer Open, chapter 10 (2019).
49. Piet, G. J., & Guruge, W. A. H. P. Diel variation in feeding and vertical distribution of
ten co-occurring fish species: consequences for resource partitioning. *Env. Biol. Fishes*
50, 293-307 (1997).
50. Revell, L. J. phytools: An R package for phylogenetic comparative biology (and other
things). *Methods Ecol. Evol.* **3**, 217–223 (2012).
51. Stearns, S. C. The influence of size and phylogeny on patterns of covariation among
life-history traits in the mammals. *Oikos* **41**, 173–187 (1983).
52. Caswell, H. Matrix Population Models. (Sunderland, MA: Sinauer Associates 2001)
53. Tuljapurkar, S., Gaillard, J. M. & Coulson, T. From stochastic environments to life
histories and back. *Phil. Trans R. Soc. B.* **364**, 1499–1509 (2009).
54. Keyfitz N., & Caswell H. Applied mathematical demography. (Springer, New York 2005).

Figure 1. Workflow of parameterising a DEB-IPM and example applications, including what databases DEBBIES can feed into (Essential Biodiversity Variables[12]; MOSAIC[10]). DEBBIES currently contains 185 ectotherms of 18 different orders. Eight life history trait values are required to parameterise a DEB-IPM (top-right box). Once parameterised, it can be used to calculate a further nine derived life history traits (Table 3) that can in turn be summarised into life history strategies[1]. The resulting 'fast–slow continuum and reproductive strategy framework' can be linked to quantities from other disciplines[4].

572 **Figure 2.** Technical validation. Shown are for three levels of experienced feeding level (blue:
573 $E(Y)$ = 0.9; red: $E(Y)$ = 0.7; yellow: $E(Y)$ = 0.5) the frequency distributions of the population
574 growth rate λ (**a**), observed versus predicted generation time (years) (**b**); observed versus
575 predicted longevity (years) (**c**); and observed versus predicted age at maturity (years) (**d**).
576 The black lines in (B-D) denote the $x = y$ line of equal values.

Table 1. Description of the columns in the data records file; each row in the file is a species.

Name	Column	Description
RecordID	1	Record ID
Class	2	Species class
Order	3	Species order
Family	4	Species family
Species	5	Species Latin name
Common_name	6	Species common name
κ	7	Fraction of energy allocated to respiration (κ) (as opposed to reproduction)
L_b	8	Length at birth (cm)
L_p	9	Length at puberty (cm)
L_m	10	Maximum length (cm)
μ_j	11	Juvenile mortality rate (y^{-1})
μ_a	12	Adult mortality rate (y^{-1})
r_B	13	von Bertalanffy growth rate (y^{-1})
R_m	14	Maximum reproduction rate (y^{-1})
Contributor	15	Name of the person who collected the trait data
kappa_REF	16	Reference for κ
L_b_REF	17	Reference for L_b
L_p_REF	18	Reference for L_p
L_m_REF	19	Reference for L_m
μ_j_REF	20	Reference for μ_j
μ_a_REF	21	Reference for μ_a
r_B_REF	22	Reference for r_B
R_m_REF	23	Reference for R_m

Table 2. Technical validation that fitted linear regression models without an intercept ($y \sim x$) on observed (x) and predicted (y) values of generation time (G), longevity (L) and age at maturity (A) for three experienced feeding levels $E(Y)$ = 0.5, $E(Y)$ = 0.7, and $E(Y)$ = 0.9. The 95% confidence intervals (CIs) of the regression coefficient are approximated as the regression coefficient $\pm$ twice its standard error. If a 95% CI overlaps with 1, predicted values do not significantly differ from observed values. Also given is the root mean square error (RMSE) and the coefficient of determination (R_2) of each model. See also Fig. 2.

$E(Y)$	model $y \sim x$	regression coefficient	95% CI	RMSE	R^2
0.5	$G_{predicted} \sim G_{observed}$	1.24	1.10 – 1.38	9.34	0.27
0.7	$G_{predicted} \sim G_{observed}$	1.23	1.16 – 1.30	11.40	0.28
0.9	$G_{predicted} \sim G_{observed}$	1.02	0.97 – 1.07	9.92	0.38
0.5	$L_{predicted} \sim L_{observed}$	0.59	0.57 – 0.61	6.14	0.55
0.7	$L_{predicted} \sim L_{observed}$	0.84	0.81 – 0.87	11.20	0.40
0.9	$L_{predicted} \sim L_{observed}$	0.99	0.93 – 1.05	21.50	0.16
0.5	$A_{predicted} \sim A_{observed}$	1.25	1.21 – 1.29	5.70	0.49
0.7	$A_{predicted} \sim A_{observed}$	1.30	1.26 – 1.34	5.51	0.57
0.9	$A_{predicted} \sim A_{observed}$	1.45	1.40 – 1.50	6.60	0.57

Table 3. Nine key derived life history traits that inform on a species' turnover rate, longevity, growth and reproduction, for which we provide MatLab code to calculate them (see Code Availability)[40]. To calculate life history traits, one needs to discretise the IPM (equation (1)) by dividing the length domain Ω into very small-width discrete bins (we chose 200 bins), resulting in a matrix **A** of size $p \times q$, where $p = q = 200$, and which dominant eigenvalue equals λ, the population growth rate. Mean lifetime reproductive success R_0 is the dominant eigenvalue of the matrix $\mathbf{F} = \mathbf{V}(\mathbf{I} - \mathbf{GS})^{-1}$, where **I** is the identity matrix and $\mathbf{V} = \mathbf{DR}$, with **D** as the parent-offspring association, **R** the reproduction, **G** the growth and **S** the survival matrix[52]; this gives generation time[52] $T = \log(R_0)/\log(\lambda)$. The mean life expectancy, η_e, is calculated as $\eta_e = \mathbf{1}^T\mathbf{N}$, where **1** is a vector of ones of length m and **N** is the fundamental matrix $\mathbf{N} = (\mathbf{I} - \mathbf{S})^{-1}$. The longevity of an individual of length L is η_L, which means we can calculate age at sexual maturity $L_\alpha = \eta_{L_b}$ and mature life expectancy $L_\varpi = \eta_{L_p}$ so that $\eta_e = L_\alpha + L_\varpi$ [48]. l_x is the probability of surviving to age at least x, and m_x is the average fertility of age class x (cf.[53]). $\overline{\mathbf{G}}$ is the mean of **G**, $\overline{\mathbf{V}}$ is the mean of **V,** and i and j are the row and column entries of the matrix, respectively. The vital rates included in progressive growth γ, retrogressive growth ρ, and sexual reproduction φ, were averaged across the columns j (the length bins), weighted by the relative contributions of each stage at stationary equilibrium. For example, to calculate mean sexual reproduction φ, we summed the values in the columns j of the **V** matrix and multiplied each φ_{ij} by the corresponding jth element w_j of the stable stage distribution **w**, calculated as the right eigenvector of **A**. Finally, the degree of iteroparity $S = \sqrt{V_\mathbf{f}}/\overline{a}_\mathbf{f}$, with $V_\mathbf{f} = \overline{a_\mathbf{f}^2} - (\overline{a}_\mathbf{f})^2$, where $\overline{a_\mathbf{f}^2} = [(\mathbf{c} \circ \mathbf{c})^T \otimes \sum_{x=0}^{x=\eta_e} m_x]/[\mathbf{1}^T \otimes \sum_{x=0}^{x=\eta_e} m_x]$ and $\overline{a}_\mathbf{f} = [\mathbf{c}^T \otimes \sum_{x=0}^{x=\eta_e} m_x]/[\mathbf{1}^T \otimes \sum_{x=0}^{x=\eta_e} m_x]$, with $\mathbf{c}^T = (1\ 2\ \cdots\ p)$ [48].

Life history trait	Symbol	Definition	Equation
Generation time	T	Number of days required for the individuals of a population to be fully replaced by new ones	$T = \dfrac{\log(R_0)}{\log(\lambda)}$
Survivorship curve	H	Keyfitz' entropy ($H < 1$ denotes increasing mortality rate with age, $H > 1$ denotes decreasing mortality rate with age, $H = 1$ denotes a constant mortality rate with age)[54].	$H = -\dfrac{\sum_{x=0}^{x=\eta_e} \log(l_x)l_x}{\sum_{x=0}^{x=\eta_e} l_x}$
Age at maturity	L_α	Number of days that it takes an average individual in the population to become reproductive	$L_\alpha = \eta_{L_b}$
Progressive growth	γ	Mean probability of growing to a larger length across the length domain Ω.	$\gamma = \sum_{i}^{m} \overline{\mathbf{G}}_{i,j}\vert_{i<j}$
Retrogressive growth	ρ	Mean probability of growing to a smaller length across the length domain Ω.	$\rho = \sum_{i}^{m} \overline{\mathbf{G}}_{i,j}\vert_{i>j}$
Mean recruitment success	φ	Mean per-capita number of recruits across the length domain Ω.	$\varphi = \sum_{i}^{m} \overline{\mathbf{V}}_{i,j}$
Degree of iteroparity	S	Coefficient of variation in age at reproduction	$S = \sqrt{V_\mathbf{f}}/\overline{a}_\mathbf{f}$
Net reproductive rate	R_0	Mean number of recruits produced during the mean life expectancy of an individual in the population.	$R_0 = \sum_{x=0}^{x=\eta_e} l_x m_x$
Mature life expectancy	L_ω	Number of days from the mean age at maturity (L_α) until the mean life expectancy (η_e) of an individual in the population.	$L_\varpi = \eta_{L_p}$