The three-dimensionally articulated oral apparatus of a Devonian
heterostracan sheds light on feeding in Palaeozoic jawless fishes

Richard Dearden[1,2*], Andy Jones[1], Sam Giles[1,3], Agnese Lanzetti[1,3], Madleen Grohganz[4], Zerina
Johanson[3], Stephan Lautenschlager[1], Emma Randle[1], Philip C. J. Donoghue[4], Ivan J. Sansom[1]

[1] School of Geography, Earth & Environmental Sciences, University of Birmingham, Edgbaston,
Birmingham, B15 2TT,

[2] Naturalis Biodiversity Center, 2 Darwinweg, Leiden, 2333 CR, The Netherlands.

[3] Natural History Museum, Cromwell Road, London SW7 5BD, UK

[4] Bristol Palaeobiology Group, School of Earth Sciences, University of Bristol, Bristol BS8 1TQ, UK.

*Corresponding author: r.dearden@bham.ac.uk

Attempts to explain the origin and diversification of vertebrates have commonly invoked the
evolution of feeding ecology, contrasting the passive suspension feeding of invertebrate chordates
and larval lampreys with active predation in living jawed vertebrates. Of the extinct jawless
vertebrates that phylogenetically intercalate these living groups, the feeding apparatus is preserved
only in the early diverging stem-gnathostome heterostracans and its anatomy remains poorly
understood. Here we use X-ray microtomography to characterise the feeding apparatus of the
pteraspid heterostracan *Rhinopteraspis dunensis* (Roemer, 1855). We show that the apparatus is
composed of thirteen plates arranged approximately bilaterally, the majority of which articulate
from the postoral plate. Our reconstruction of the apparatus shows that the oral plates would have
been capable of movement within the dorso-ventral plane, but their degree of movement was
limited. The functional morphology of the apparatus in *Rhinopteraspis* precludes all proposed
interpretations of feeding except for suspension/deposit feeding and we interpret the apparatus as
having served primarily to moderate the oral gape. This is consistent with evidence that at least
some early jawless gnathostomes were suspension feeders and runs contrary to macroecological
scenarios that envisage early vertebrate evolution as characterised by a directional trend towards
increasingly active food acquisition.

Keywords: Heterostracan, Pteraspid, Feeding, Devonian, Palaeozoic, Ostracoderm

1. Introduction

Feeding figures prominently in attempts to understand the evolutionary origins of vertebrates (Janvier 1996, Anderson *et al.* 2011). In contrast to invertebrate chordates, which exclusively suspension feed with either a ciliated pharynx or a mucus net (1), the dorso-ventrally closing jaws of living jawed vertebrates (crown-gnathostomes plus 'placoderms') (2–4) or antero-posteriorly moving system of cartilages (in cyclostomes, hagfishes and lampreys) (5–8) allow for a far broader range of feeding strategies. The evolution of these unique vertebrate feeding modes plays a major role in attempts to explain the evolution of vertebrate anatomy and the origins of its modern diversity (9,10). Prominently, the New Head Hypothesis (11–15), argues that the shift from suspension feeding to predation accompanied the emergence of neural crest, neurogenic placodes, and the accompanying evolution of a prechordal head. This hypothesis predicts the earliest vertebrates with a prechordal head, i.e. parts formed from trabecular elements of the neurocranium anterior to the notochord(16), had a predatory feeding mode.

The fossil record of Palaeozoic vertebrates provides a means of testing such hypotheses, but current interpretations of that record are far from decisive. In particular, heterostracans, an extinct group of jawless stem-gnathostomes, have been the focus of much of the debate over feeding in early vertebrates. This is because their oral region is more commonly and completely preserved than in any other such group, and they are often interpreted as one of the earliest diverging lineages of stem-gnathostomes. As such, heterostracans have the potential to inform on the feeding ecology of the earliest members of the gnathostome lineage (17). The heterostracan feeding apparatus is best known in pteraspids, where the oral region is characterised by distinctive macromeric dermal plates (18–22). The function of these plates has been much debated, variously interpreted as biting (19) or slicing (23) 'jaws', a cyclostome-like feeding apparatus (24–27), a sediment scoop (22), or a filtering structure (28–32). Equally varied are the inferred ecologies, with heterostracans interpreted as active predators (19), macrophagous selective predators (12,33) , hagfish-like scavengers (24–27), herbivores (23), detritivores (22,34), or suspension feeders (including filter feeding) (10,20,35–38). The most recent investigation suggests that heterostracans were suspension feeders because the analysed oral plates exhibited no evidence of the wear anticipated of a 'tooth-like' function (20).

The difficulty in studying the articulated heterostracan oral apparatus *in situ* contributes to this lack of consensus. In the rare cases where they are preserved, articulated heterostracan oral apparatuses consist of small plates suspended in encasing sediment. As a result, previous reconstructions of the oral apparatus have focused either on the gross arrangement of the apparatus, in which the morphology and detailed arrangement of the individual plates are not characterised (22,24,26,27), or describe isolated elements with little or no reference to articulated apparatuses (22,39,40). Evidently, the feeding ecology of heterostracans remains in its infancy and so we sought to advance understanding through a detailed characterization and reconstruction of the heterostracan oral apparatus. We used X-Ray microtomography to characterise the three-dimensionally (3D) articulated oral apparatus of an exceptionally well-preserved specimen of the pteraspid heterostracan *Rhinopteraspis dunensis* (Roemer, 1855). Using computed tomography, we generated volumetric models of the components of the oral apparatus and used these models to reconstruct their three dimensional arrangement in vivo. We use this reconstruction to assess competing hypotheses of heterostracan feeding.

2. Materials & Methods

(a) Specimens and locality

Rhinopteraspis dunensis (Roemer, 1855) NHMUK PV P 73217 is housed in the collections of the Natural History Museum, London (NHMUK). In the museum catalogue the specimen is listed as being collected from "Odenspiel Quarry", likely corresponding to Jaeger Steinbruch, a quarry near the village of Odenspiel, Reichshof, North-Rhine Westphalia, Germany or, possibly, outcrops in the local area (41,42). The Jaeger Quarry and surrounding outcrops expose sandstones and mudstones belonging to the Siegenian (?upper Pragian or lower Emsian, Lower Devonian) Odenspiel Formation (43,44), deposited on the northern margin of the Rhenohercynian Basin, which was a marginal transgressive and regressive delta-dominated setting (45). The Odenspiel Formation falls within the 'Pararhenotypics subfacies' of Jansen (46), representing a marginal marine, intertidal lagoonal setting preserving a restricted fauna of fish, bivalve molluscs, lingulid and terebratulid brachiopods, and eurypterids (41,42,47–49).

(b) Terminology

Various terminologies have been applied to the pteraspid oral region. Here we follow the terminology of Randle and Sansom (50) and, where that is not possible, that of Blieck (51). We extend existing terminology to describe the nature and arrangement of the oral plates (Fig. 1G). For the anatomical axes of the animal as a whole we use dorsal/ventral in the dorsoventral axis, rostral/caudal for the sagittal axis, and dextral/sinistral for the transverse axis. When describing the oral apparatus itself we also use lateral/medial to describe lateral positions relative to the sagittal axis, ad/aboral to describe the surfaces of plates relative to the mouth (i.e. adoral is the surface of a plate facing the mouth, aboral the surface facing away), and proximal/distal to refer to positions on the oral plates relative to their articulation with the postoral plate with oral plate tips being distal.

(c) Computed tomography

Rhinopteraspis dunensis NHMUK PV P 73217 (Figs 1,S1) was scanned using a Nikon Metrology XTH 225ST X-ray tomography instrument based in Bristol Palaeobiology, University of Bristol. Two scans were undertaken, each composed of two stacked scans. The first scan included the whole headshield at a voltage of 223 kV and a current of 193 µA, with 3141 projections, and with a 1 mm Sn filter, obtaining a dataset with 63.41 µm voxel resolution (Fig. 1A,B). The second scan targeted the oral region specifically, including the posterior third of the rostrum, the orbital, oral, and pineal parts of the headshield, and the anterior quarter of the dorsal and ventral discs, at a voltage of 180 kV and a current of 178 µA, with 3141 projections of 8 frames and 708 ms each, and with a 0.5 mm Cu filter, achieving a voxel resolution of 22.91 µm (Figs 1D-H,2).The resulting tomographic datasets were segmented in Mimics v.25 (materialise) to create 3D models. All 3D models were visualised in Blender 3.5 (blender.org).

(d) Reconstruction and animation

3D models of the higher resolution scan set of *Rhinopteraspis* were imported into Blender for retrodeformation (Fig. 3). Use of Blender tools for retrodeformation followed the recommendations and techniques set out by Herbst et al. (52). Initial retrodeformation focused on the best preserved plates and those with clearly delineated articulations (53), including the rostral, orbital, supraoral and paraoral plates. After rearticulating and repairing deformation (see Supplementary Information),

these elements provided a framework to delimit the dorsal and lateral extent of the oral plate array.
The oral plates were aligned, maintaining their preserved order, within this delimited area by placing
their dorsal tips close to the margin of the mouth and rotating the plates caudoventrally to match
the angle of the surrounding plates. The postoral plates were then fitted to the proximal ends of the
oral plates. The proximal ends of the oral plates were then readjusted into articulation with the
sulcus preserved in the postoral plates (Fig. 2C), while maintaining alignment with the closely
articulated lateral oral- and paraoral plates and each other. Finally the ventral disc was
retrodeformed and articulated with the postoral plate. The reconstructed specimen was animated in
Blender to simulate the movement of the oral plates Fig. S3). Empties (single geometry-less points
that act as handles for object transformation without interfering in the render process) were placed
between each oral plate and the postoral plate below, so that the local x-axis of each was aligned
with the approximate outer boundary of the postoral plate. Each oral plate was then parented
(=linked) to the empty below it. These empties were then animated to rotate around their local x-
axes, causing the parented oral plate to also rotate around that axis.
134
3. Description
NHMUK PV P 73217 comprises an almost-complete three-dimensional specimen of *Rhinopteraspis*
dunensis, preserving the entire headshield and articulated body scales (Fig. 1A,B,S1). The specimen
has been crushed laterally, with the oral region and ventral shield displaced rostro-dorsally (Fig. 1).
Otherwise, the specimen is complete and individual oral elements appear to have maintained their
original shape and relative location, as evidenced by their approximately symmetrical arrangement.
The dermal skeletal anatomy of pteraspid heterostracans is well-characterised in numerous taxa
(18,54) to which that of *Rhinopteraspis dunensis* NHMUK PV P 73217 conforms (32,55,56). The
headshield is composed of large dorsal and ventral shields separated by paired cornual and branchial
plates, with a dorsal spine set into the posterior margin of the dorsal shield (Fig. 1A,B,S1). Anterior to
the dorsal shield is an elongate rostrum that is separated from the dorsal shield by paired orbital
plates (Figs 1A,B,S1). The anterior length of the rostrum is broken off from the rest of the specimen
(Figs 1A,B,S1). The unpaired pineal plate is indistinguishable from the top of the orbital plates in the
scan data.
149
The oral region is bordered dorsally by the supraoral plate, laterally by paired paraoral plates, and
caudally by the postoral plate (Fig. 1). In previous descriptions of pteraspids, including
Rhinopteraspis, the subrostral lamella and ascending postrostral lamella have been characterised as
part of the rostral plate (51,57), although Friman & Bardenheuer described paired plates in this
position they termed the 'subrostral plates' (32). In NHMUK PV P 73217 they comprise a separate
structure, broken into three parts (postmortem), which we term the supraoral plate (Figs 1D,G,H,
2A,B). The supraoral plate is trapezoidal in shape, narrowing rostrally. A pronounced furrow runs
around its lateral and rostral margins, which is overlapped by the paraoral,

Figure 1. *Rhinopteraspis dunensis* NHMUK PV P 73217. A-C, Rendering of head shield based
on computed tomographic data in (A) sinistral, (B) dextral view, and (C) transparent with
scheme of anatomical axes. D-H, renders based on higher resolution data showing the oral
apparatus in more detail in D, ventral view; E, sinistral view; F, dextral view; G rostral view;
H, Rostro-ventral view. Green and blue parts of 3D renders represent oral region.
Abbreviations: S, sinistral (left), D, dextral (right). Scale bars represent 5 cm in panels A,B, 1
cm in panels D-H.

orbital, and rostral plates; the margins of the plate sweep ventro-caudally to form two latero-ventral
arms (Fig. 2B). The rostral half of the ventral surface of the supraoral plate (the subrostral lamella) is
surmounted by a superficial layer of ornament and made convex by a prominent median crest,
aligned rostro-caudally (Fig. 2B). The caudalmost half (the ascending postrostral lamella) curves
upwards into the mouth and lacks dentine ornament (as in *Rhinopteraspis cornubica* Tarlo, 1961, p.
373), the rostral border of this area presumably represents the position of the oral opening. This
border is marked by a row of tubercles, comprising one large medial tubercle and two pairs of
smaller tubercles on either side (Fig. 2B). The paired paraoral plates are elongate, taper rostrally, and
overlie large rostro-lateral overlap surfaces on the orbital plates, rostral to the orbits (Figs 1D-F, 2A).
We interpret the 'olfactory grooves' identified in *Rhinopteraspis cornubica* by Tarlo (1961, fig. 1) as
the overlap surfaces between the paraoral plates and supraoral plate (Fig. 2A) (32).

The postoral plate is bow-shaped and originally symmetrical, although the right process is damaged
(Fig. 2A,C). The ventral surface is smoothly convex. The inner surface has a caudal overlap surface
that is concave to curve around the anterior rim of the ventral shield, and a dorsal surface that forms
the ventral margin of the oral opening, bearing paired sulci for the oral plates that are interrupted
medially by a raised area (Fig. 2C). Paired lateral plates and lateral postoral plates lie between the
postoral plate and the orbital plates (Figs 1,2). The extensive overlap between the plates
surrounding the mouth and the larger head shield plates strongly suggests that they comprised an
integrated structural unit with little or no movement relative to each other.

The oral apparatus itself is composed of thirteen imbricated oral plates and one pair of lateral oral
plates (Figs 1,2,S2). The aboral, mediolateral surfaces of each oral plate, as well at the lateral
surfaces of the hook, are all faced with tuberculated ornament (Fig. 2F-I). However, the lateral and
adoral surfaces of the oral plates, as well as the ventral side of the hook and the proximal end of the
plates, are all unornamented, instead exhibiting a porous surface texture reflecting open vascular
canals. The plates are arranged bilaterally about the midline into sinistral and dextral series.
Although plate pairs that occur in equivalent positions on either side of the midline are similar, they
do not exhibit mirror-image symmetry. In particular, the unpaired medial plate is not symmetrical
but, rather, is continuous with the sinistral series (Fig. 2E,S2). Each oral plate has the same general
morphology of a main limb with a rhomboidal cross-section, a distal hook (except for the most
laterally placed plates), and a proximal articulation surface for the postoral plate (Figs 2,S2). There
are six oral plates in the sinistral series (S1-6) and six in the dextral series (D1-6), each preserved
inclined at varying degrees (maximum about 45 deg) to each other along their coronal axis. The
more medially placed the plate, the more inclined it is along its long axis to provide a fit with the
adjacent oral plate; the lateral and medial faces of the plates overlap and imbricate, inclined at
increasing angles relative to the sagittal plane, from medial to lateral. The distal hooks of the oral
plates curve adorally, while the proximal ends of the plates are notched, reflecting the ventral limit
of the external dermal ornament, serving as articulation with the postoral plate (except for the
medial plate M).

Figure 2. *Rhinopteraspis dunensis* NHMUK PV P 73217 oral region. A, oral apparatus and
surrounding plates as preserved, in ventral view; B, ventral view of supraoral plate with
three fragments rearticulated; C, postoral plate in dorsal view; D, oral apparatus as
preserved with ventral plates removed; E, dorsal view of oral plates as preserved; F-I, oral
plate R4 in aboral (F), lateral (G), adoral (H), and medial (I) views, alongside drawings
depicting the inferred extent of dermal ornament in grey, based on comparison with
isolated plates of *Loricopteraspis dairydinglensis*(20,40,58). Abbreviations: S, sinistral (left),
D, dextral (right). Scale bars represent 1 cm in panels A-E, 0.5 cm in panels F-I.
214

Although the oral plates have similar morphologies, individually they vary in relative proportion
depending on their position within the apparatus (Figs 2,S2). The hook of the unpaired medial oral
plate M is as long as the main limb. This plate is preserved overlying the lateral two oral plates D1
and S1. The lateral side of its proximal end is notched to fit with the left medial face of D1. Plate M
curves dextrally such that it fits the curvature of the dextral face of the adjacent oral plate S1. The
main limb of oral plates D1 and S1 are twice as long that of M, with a narrow base; similarly, it may
not have articulated with the postoral plate, but the hooks in D1 and S1 are as large and very similar
in shape. These plates fit around oral plates S2 and D2, which fit around S3 and D3, etc.; these pairs
have narrow bases and slightly smaller hooks than S1 and D1. Plates S4 and D4 have a slightly
broader base and proportionally shorter hook. This trend continues laterally, with increasingly
shorter hooks and wider bases in S5 and D5. Finally S6 and D6 have no perceptible hook and a very
broad base. The lateral edges of plates S6 and D6 is concave, fitting the medial edge of the lateral
oral plates.

The lateral oral plates are morphologically distinct from the oral plates (Fig. 2,S2), approximately
triangular, tapering rostrally with a curved medial margin that matches the lateral profile of the
lateral-most oral plates (Figs 2A,S2). The better preserved sinistral lateral oral plate appears to have
a distinct notch in its posterior side (Fig. S2A,C), although this is difficult to corroborate from the
dextral lateral oral plate.

4. Reconstruction

The combined width of the bases of the oral plates, when aligned perpendicular to a sagittal plane,
matches the length of the sulcus on the postoral plate (Fig. 3) and the complementary symmetry
exhibited by adjacent plates indicates that they were capable of almost completely filling the width
of the oral opening. In this position, the lateral-facing surface of each oral plate overlaps the adoral
surface of its outer neighbour (Fig. 3). The lateral surfaces of the outermost oral plates fit closely
with the lateral oral plates (Figs. 3C,D). M, S1 and D1 appear to lie on top of the neighbouring oral
plates rather than contacting the postoral plate. When viewed caudally, this brings the hooks of the
plates into alignment, forming a plane above the posterior unornamented surface of the plates (Fig.
3d). This reconstruction also suggests that the oral apparatus fits together such that the tops of the
oral plates extend to the top of the mouth to almost meet the supraoral plate. The medial crest of
the supraoral plate creates a convex dorsal margin of the mouth, and the increasing length of the
oral plates in lateral positions means that their dorsal tips also follow this convex line. Thus, when
fully closed, there would have remained a narrow "letterbox-shaped" opening (cf. (59)) into which
projected the dorsal tips of the oral plates and the medial crest and associated tubercles of the
dorsal oral plate. Indeed, the medial crest of the supraoral plate would have effectively divided the
residual opening into two. There is no evidence that the plates intercalated with the tubercles
demarcating the dorsal margin of the mouth at the rostral margin of the ascending postrostral
lamella, Fig. 2B). When modelled to open around the axis of the sulcus on the postoral plate
synchronously, the oral plates do not overlap as they rotate (Fig. S3). Instead, their placement along
the curved axis of the postoral plate sulcus causes them to splay outwards (Fig. S3).

Figure 3. Reconstruction of *Rhinopteraspis dunensis* based on NHMUK PV P 73217, anterior

section of rostrum not shown. A, rostrolateral view; B, ventral view; C, rostral view; D,

caudal view of articulated oral plates; E, sagittal cross-section through centre of the head; F,

close up cross-section lateral to E, showing junction between base of oral plate and the

postoral plate.

5. Discussion

Based upon our 3D reconstruction of the oral region in *Rhinopteraspis*, we are able to consider the oral plates as an integrated apparatus and test hypotheses of its form and function. The endoskeleton of heterostracans is completely unknown beyond what can be inferred from the dermal skeleton (60). Hence, we attempt to consider the constraints imposed by the dermal skeleton, without speculating as to endoskeletal structure.

(i) Interpretation of the oral apparatus of *Rhinopteraspis*

Most previous hypotheses of function in heterostracan feeding assume significant movement and a degree of rotation of the oral apparatus, often following a cyclostome-like mode (19,22,24). In many of these the postoral plate(s) is assumed to move significantly, either inwards (24) or outwards (22,59). The large, curved overlap surface at a 45° angle between the postoral plate and the ventral plate in *Rhinopteraspis* (Fig. 2C) suggests that this movement would not have been possible and that the postoral plate was static during feeding. In contrast, the imbricated nature of the oral plates strongly suggests that they were mobile relative to the postoral plate. Any movement requires rotation around the point of their attachment to the postoral plate. The rhomboid cross-section and the complementary symmetry of the oral plates would have prevented them either moving independently, or moving in an entirely sagittal plane (Fig. S3). Rather, they must have moved as an integrated unit, splaying ventro-laterally as the plates rotated aborally on the sulcus of the postoral plate. The medial plates (M, S1, D1) that do not articulate with the postoral plate are the exceptions: their cross-sectional shape and ab/adoral overlap would have precluded their movement relative to the other plates.

The morphologies of individual oral plates are closely comparable to those observed in acid-prepared isolated oral plates of *Loricopteraspis dairydinglensis* recovered through acid digestion of limestone (20,40,58). The open vasculature on the lateral surfaces and adoral surfaces, as well as the ventral side of the hook and the proximal end of each plate, is significant. This indicates that these parts of the plates were embedded in soft tissue, the upper limit of which would have been the proximal surface of the large projecting hooks on the adoral side of the oral plates. The medial plates would have been supported entirely by this soft tissue which must have provided the basis of any movement of the plates and so may have included unmineralized cartilage, muscles or tendons.

Our reconstruction indicates that, even when the mouth was fully closed by occlusion of the oral plates, there would have remained a short but wide gape that was effectively divided in two by the median crest of the supraoral plate. The associated tubercles of the supraoral plate and tips of the oral plate would have projected into this space, further occluding it. This oral morphology precludes interpretation as a hagfish-like prenasal sinus (26,27) because the opening is too small. There is also no evidence for separate upper oral plates as inferred by Stensio (26,27). Assuming the presence of forward pointing denticles on the lateral surfaces of the oral plate hooks, as in *Loricopteraspis* and *Pteraspis* (20,58), these structures would not have bordered the oral opening but, rather, the junction between each plate and its neighbour, linked by soft tissue. Rather than being involved in food capture or processing, this ornamentation may have helped to prevent particles from becoming lodged in the spaces between the plates (58) and their associated soft tissue, preventing fouling of the oral apparatus (cf. Hamann and Blanke (1)). Any movement seems unlikely to have been great in

310 magnitude due to the weak joint between the oral plates and the postoral plate, as well as the
311 suspension of the small median oral plates in soft tissue. A structural analogue in a living vertebrate
312 for heterostracan oral plates might be the branchiostegal plates in osteichthyans, which support the
313 branchiostegal membrane, and make limited, coordinated movements to aid the suction pump (61).
314
The oral apparatus of *Rhinopteraspis* could only have been moved from the adoral side with rostro-
caudal movements, as the aboral surfaces of the oral plates lack any kind of attachment surfaces.
This is inconsistent with a gnathostome-like organisation of paired mandibular adductor muscles.
However, living cyclostomes both operate the oral apparatus by moving cartilages rostrally and
caudally along the floor of the pharynx. In hagfishes, keratinous toothlets are mounted on a
cartilaginous dental plate that is pulled anteriorly along a basal plate to evert the lingual apparatus,
and posteriorly to return it to resting position (5,6,8). In lampreys, a medial piston cartilage is
protracted rostrally, moving a medial apical cartilage, the action of which brings keratinous toothlets
in front of the apical cartilage into contact with other toothlets in a rasping action (7,62,63). A
medial groove along the visceral surface of the ventral shield in some pteraspids has been cited as
evidence for a cyclostome-like medial structure of the oral mechanism (60).
326
327 **(ii) Comparison of oral anatomy with other heterostracans**
328
The anatomy of the oral apparatus of *Rhinopteraspis* is comparable to that of other pteraspids.
Individually, the shape of the oral plates conforms closely to the morphologies of isolated plates acid
prepared from an articulated specimen of *Loricopteraspis dairydinglensis* ((40), pl. 37; (58)). The
arrangement of these plates is similar to articulated specimens of *Protopteraspis* (19) and *Errivaspis*
(22). Importantly, the posterior alignment of the hooks in *Rhinopteraspis* (Fig. 3) can also be seen in
other pteraspids where the adoral side of an articulated oral apparatus is visible: *Mylopteraspidella*
(Stensiö, (27), Fig. 44, p. 197) and *Protopteraspis* (19), as well as in *Errivaspis* (White, (22)figs 41-44).
Together, these indicate that our interpretation of the feeding apparatus of *Rhinopteraspis* is more
broadly applicable within pteraspids. However, varied body shapes and positions of the oral opening
indicate some diversity in feeding ecology (e.g. *Doryaspis* (64), *Drepanaspis* (65)).
339
Athenaegis, the oldest articulated heterostracan (66), is often assumed to be an outgroup to all
other macromeric heterostracans (21,50,67,68), and its oral apparatus appears broadly
anatomically consistent with *Rhinopteraspis*. By contrast, cyathaspids, a likely paraphyletic grade of
heterostracans (21), have arrangements of plates that are difficult to rationalise with the oral
apparatuses of *Athenaegis* and pteraspids (e.g. *Anglaspis* (37), *Poraspis* (69), *Capitaspis* (70), and
Allocryptaspis (29)). In amphiaspids, the oral aperture occurs at the end of a tube in a single, fused
headshield (71). Meanwhile the oral apparatuses of the "tessellate heterostracan" taxa are
completely unknown (34) and what is known of their anatomy is difficult to reconcile with the
pteraspids, including *Rhinopteraspis*. Thus, while the oral apparatus of *Rhinopteraspis* may be
representative of pteraspids, *Athenaegis* and, by implication, macromeric heterostracans primitively,
it may not be representative of heterostracans more generally, which may have exhibited greater
diversity in terms of feeding ecology. Further investigation is required to assess the ubiquity of
suspension-feeding within the group, and the position in the water column at which they fed.
353
(iii) Implications for feeding in pteraspids and ancestral heterostracans

Our articulated oral apparatus model can be used to consider the various feeding strategies that
have been proposed for pteraspid heterostracans. These strategies can be roughly divided into
macrophagy, predation, deposit/detritus feeding, and microphagy or suspension feeding (72).

Hypotheses of macrophagous and predatory heterostracan feeding are based on analogy between
heterostracan oral plates and the gnathostome mandible (e.g. (19)), or more rarely the hagfish oral
apparatus (e.g. (24)). However, the oral apparatus in Rhinopteraspis is poorly constructed for biting
or grasping. Although the ascending lamella bears tubercles that have been interpreted as the upper
'jaw' analogue in such a scenario (57), they do not occlude with the tips of the oral plates. Moreover,
the angle of approach of the plates to the preoral plate during closing is oblique, with a low
mechanical advantage, and would have been poorly suited to generating force. Finally, the oral
plates themselves would not have formed a firm biting surface, with a poorly developed joint with
the postoral plate even in the broad based lateral oral plates (e.g. plates S/D4-6), and the medial oral
plates (e.g. plates M, S1, D1) suspended in soft tissue or with very narrow-based articulations with
the postoral plate. Scenarios that envisage see the oral plates as elements analogous to hagfish-like
toothlets (24) can be similarly rejected because the junction between the proximal tips of the oral
plates and the sulcus in the postoral plate, and the fixed position of the postoral plate, would all
have acted to dramatically restrict the movement of the oral plates.

Deposit or detritus feeding interpretations typically envisages the oral apparatus as a 'scoop' that
would have been used to acquire sediment from the bottom of the water body (22). This requires
the plates to evert substantially from their resting position, with some reconstructions also require
movement of the postoral plate (22 figs 49,50). We can reject this based on our reconstruction of
Rhinopteraspis, where significant movement of the oral plates is limited by the joint between the
postoral and ventral plates. However, we cannot rule out the burial of the snout/mouth in the
substrate as a means of deposit feeding. The presence or absence of wear, which is often used as a
line of evidence in discussion of deposit feeding (20,73), are not visible in Rhinopteraspis at the
resolution of our scan data. We note though that the long-snout in pteraspids such as
Rhinopteraspis, along with the strongly convex ventral abdomen, would have restricted the ability of
the animal to place its mouth in contact with the substrates.

In either suspension feeding or deposit feeding modes, rostral rotation of the oral plates would have
resulted in a greater area of gape, increasing intake. The occluded oral apparatus leaves a restricted
opening with digitate margins defined by the tips of the oral plates and the tubercles of the dorsal
oral plate that would have served to prevent fouling (cf. (1,58)) while facilitating inflow in a manner
analogous to straining functions seen in animals as diverse as flamingos, brachiopods, oysters and
gastropods (74). It is possible that the exposed tips of the oral plates in Rhinopteraspis acted as an
analogous structure along with the tubercles on the supraoral plate. The limited movement acted to
control the entry of larger particles into the mouth and may also have served to provide a means to
expel larger particles trapped between the plates (1).

(iv) Implications for early vertebrate evolution
The New Head and New Mouth hypotheses (9,12) have argued for a long term shift towards
increasingly active food acquisition, from the filter-feeding of invertebrate chordates to

macrophagous predation in living jawed vertebrates. The feeding ecology of the extinct
heterostracans is key since this group represents one of the earliest diverging members of the
gnathostome stem-lineage. A diversity of feeding ecologies have been attributed to heterostracans,
from macrophagous predation, through scavenging and herbivory, to deposit and filter feeding. Our
three-dimensional reconstruction of the feeding apparatus of *Rhinopteraspis dunensis* suggests its
finger-like oral plates were limited to only a small degree of concerted rotation, precluding all
proposed feeding strategies bar suspension feeding or deposit feeding by burial of the mouth in the
substrate. Comparison to other heterostracans suggests that while this condition may be
unrepresentative of the clade generally, it is representative of pteraspids and the oldest and earliest
well-known heterostracan; as such it may be plesiomorphic for the clade. Given macrophagy in
earlier diverging lampreys, hagfish and conodonts (72), it is clear that early vertebrates and stem-
gnathostomes had already established a diversity of feeding ecologies, long before the origin of
jaws. This finding is consistent with recent challenges to the predictions of New Head Hypothesis,
demonstrating that vertebrate innovations and elaborations cannot be characterised by a directional
trend towards increasingly active food acquisition (75–77), but, rather, increasing ecological
diversity.
416
6. Conclusion
A lack of knowledge of the 3D anatomy of early vertebrate feeding apparatuses has obscured their
feeding ecology, and in turn hindered testing macroecological scenarios seeking to explain early
vertebrate evolution. Using X-Ray microtomography and computed tomography we have
reconstructed the oral apparatus of the pteraspid heterostracan *Rhinopteraspis dunensis*. The oral
apparatus in *Rhinopteraspis* is composed of one medial and six pairs of bilaterally arranged oral
plates, plus a pair of lateral oral plates. Inferred articulation of these plates indicates that range of
motion was limited; the oral plates could only move in concert and could not rotate far. When
occluded, the oral plates left a short wide gape to the mouth, closed partially by a convex crest
extending from the supraoral plate and the distal tips of the oral plates. The reconstructed anatomy
precludes all proposed feeding modes bar suspension or deposit feeding in *Rhinopteraspis*.
Heterostracans more generally show a far wider range of oral anatomies and body shapes than in
pteraspids (29), and a diversity of feeding ecologies was established early in vertebrate evolution,
long before the origin of jaws. Given the existence of contemporary macrophagous vertebrate
predators and scavengers, the presence of suspension feeding heterostracans is incompatible with
the directional trend towards increasingly active food acquisition envisaged by the New Head
hypotheses.
434

435   **Acknowledgements** Many thanks to Liz Martin-Silverstone (University of Bristol) for help
436   with CT scanning. Thanks also to Emma Bernard (NHMUK) for specimen access.
437
Funding This work was supported by the Leverhulme Trust [RPG-2021-271 "Feeding without
jaws: innovations in early vertebrates"]. RPD is currently supported by Marie Skłodowska-
Curie Action DEADSharks (Grant Agreement ID 101062426). SG was also supported by a
Royal Society Dorothy Hodgkin Research Fellowship (DH160098). MG is funded by the
Natural Environment Research Council (GW4+ DTP studentship) while PD is funded by the

Leverhulme Trust (RF-2022-167) and the John Templeton Foundation (JTF 62220; JTF

62574).

Data Availability

On publication all CT data and 3D models upon which this study is based will be made

publicly and freely available.

References

References

1.   Hamann L, Blanke A. Suspension feeders: Diversity, principles of particle separation
and biomimetic potential. J R Soc Interface. 2022;19(186).
2.   Anderson PSL, Westneat MW. Feeding mechanics and bite force modelling of the
skull of Dunkleosteus terrelli, an ancient apex predator. Biol Lett. 2007;3(1):77–9.
3.   DeLaurier A. Evolution and development of the fish jaw skeleton. Wiley Interdiscip
Rev Dev Biol. 2019;8(2):1–36.
4.   Smithwick FM. Feeding ecology of the deep-bodied fish Dapedium (Actinopterygii,
Neopterygii) from the Sinemurian of Dorset, England. Palaeontology. 2015;58(2):293–
311.
5.   Clark AJ, Summers AP. Morphology and kinematics of feeding in hagfish: Possible
functional advantages of jaws. Journal of Experimental Biology. 2007;210(22):3897–
909.
6.   Clubb BL, Clark AJ, Uyeno TA. Powering the hagfish "bite": The functional
morphology of the retractor complex of two hagfish feeding apparatuses. J Morphol.
2019;280(6):827–40.
7.   Hilliard RW, Potter IC, Macey DJ. The Dentition and Feeding Mechanism in Adults of
the Southern Hemisphere Lamprey Geotria australis Gray. Acta Zoologica.
1985;66(3):159–70.
8.   Dawson JA. The oral cavity, the "jaws" and the horny teeth of Myxine glutinosa. In:
Brodal AB, Fänge R, editors. The Biology of Myxine. Oslo: Universitetsforlaget; 1963.
p. 231–55.
9.   Mallatt J. Ventilation and the origin of jawed vertebrates: a new mouth. Zool J Linn
Soc. 1996;117:329–404.
10.   Mallatt J. Feeding ecology of the earliest vertebrates. Zool J Linn Soc. 1984;82:261–
72.
11.   Northcutt RG. The new head hypothesis revisited. J Exp Zool B Mol Dev Evol.
2005;304(4):274–97.
12.   Gans C, Northcutt RG. Neural crest and the origin of vertebrates: a new head.
Science (1979). 1983;220(4594):268–73.
13.   Northcutt RG. The origin of craniates: neural crest, neurogenic placodes, and
homeobox genes. Israeli Journal of Zoology. 1996;42:273–313.
14.   Gans C. Stages in the origin of vertebrates: analysis by means of scenarios.
Biological Reviews. 1989;64:221–68.
15.   Gans C. Evolutionary origin of the vertebrate skull. In: The Skull: Patterns of Structural
and Systematic Diversity vol 2. Chicago: University of Chicago Press; 1993. p. 1–35.
16.   Kuratani S, Ahlberg PE. Evolution of the vertebrate neurocranium: Problems of the
premandibular domain and the origin of the trabecula. Zoological Lett. 2018;4(1):1–
10.
17.   Keating JN, Marquart CL, Marone F, Donoghue PCJ. The nature of aspidin and the
evolutionary origin of bone. Nat Ecol Evol [Internet]. 2018; Available from:
http://www.nature.com/articles/s41559-018-0624-1
18.   Blieck ARM. Hétérostracés, Pteraspidiformes: systemastique,phylogenie,
biostratigraphie et biogeographie. Paris: Editions du CNRS; 1984. 1–25 p. (Cahiers
du Palaéontologie (section Vertébrés)).
19.   Kiaer. The structure of the mouth of the oldest known vertebrates, pteraspids and
cephalaspids. Palaeobiologica. 1928;1.

20.   Purnell MA. Feeding in Extinct Jawless Heterostracan Fishes and Testing Scenarios
497   of Early Vertebrate Evolution. Proc R Soc Lond B Biol Sci. 2002;269(1486):83–8.
498   21.   Randle E, Sansom RS. Phylogenetic relationships of the higher heterostracanś
499   (heterostraci: Pteraspidiformes and cyathaspididae), extinct jawless vertebrates. Zool
500   J Linn Soc. 2017;181(4):910–26.
501   22.   White EI. The ostracoderm Pteraspis KNER and the relationships of the agnathous
502   vertebrates. Philosophical Transactions of the Royal Society of London B.
503   1935;225(527).
504   23.   Bendix-Almgreen SE. Silurian ostracoderms from Washington Land (North
505   Greenland), with comments on cyathaspid structure, systematics and phyletic
506   position. Rapport Grønlands Geologiske Undersøgelse. 1986;132(1981):89–123.
507   24.   Janvier P. The Structure of the naso-hypophysial Complex and the Mouth in Fossil
508   and Extant Cyclostomes, with Remarks on Amphiaspiforms. Zool Scr. 1974;3(4):193–
509   200.
510   25.   Jarvik E. Basic structure and evolution of vertebrates. London: Academic Press; 1980.
511   26.   Stensiö E. Les Cyclostomes fossiles ou Ostracodermes. In: Grassé PP, editor. Traité
512   de Zoologie. Paris: Masson; 1958. p. 173–425.
513   27.   Stensiö E. Les Cyclostomes fossiles ou Ostracodermes. In: Traité de paleontologie.
514   Paris: Masson; 1964.
515   28.   Janvier P. Facts and fancies about early fossil chordates and vertebrates. Nature
516   [Internet]. 2015;520(7548):483–9. Available from:
517   http://www.nature.com/doifinder/10.1038/nature14437
518   29.   Elliott DK, Reed RC, Loeffler EJ. A new species of Allocryptaspis (Heterostraci) from
519   the Early Devonian, with comments on the structure of the oral area in cyathaspids.
520   In: Recent Advances in the Origin and Early Radiation of Vertebrates. 2004. p. 455–
521   72.
522   30.   Lingham-Soliar T. The vertebrate integument volume 1: Origin and evolution. The
523   Vertebrate Integument Volume 1: Origin and Evolution. 2014;1:1–268.
524   31.   Stiefel KM. Evolutionary trends in large pelagic filter-feeders. Hist Biol [Internet].
525   2021;33(9):1477–88. Available from: https://doi.org/10.1080/08912963.2019.1711072
526   32.   Friman L, Bardenheuer P. A new reconstruction of Rhinopteraspis dunensis (Agnatha,
527   Heterostraci) with some remarks on pteraspid ontogeny. Munster Forsch Geol
528   Paläont. 1984;61:147–73.
529   33.   Denison RH. Feeding mechanisms of agnatha and early gnathostomes. Am Zool.
530   1961;1(2):177–81.
531   34.   Dineley DL, Loeffler EJ. Ostracoderm faunas of the Delorme and associated Siluro-
532   Devonian Formations, North West Territories, Canada. Palaeontological Association
533   Special Papers in Palaeontology. 1976;18:1–214.
534   35.   Romer AS. The Vertebrate Story. Chicago: University of Chicago Press\; 1959.
535   36.   Mallatt J. Reconstructing the life cycle and the feeding of ancestral vertebrates. In:
536   Foreman RE, Gorbman A, Dodd JM, Olson R, editors. Evolutionary biology of
537   primitive fishes. New York: Plemen Press; 1985.
538   37.   Blieck A, Heintz N. The cyathaspids of the Red Bay Group (Lower Devonian) of
539   Spitsbergen. Polar Res. 1983;1(1):49–74.
540   38.   Halstead LB. The Heterostracan Fishes. Biological Reviews [Internet].
541   1973;48(3):279–332. Available from: http://doi.wiley.com/10.1111/j.1469-
542   185X.1973.tb01005.x

39. Ball HW, Dineley DL. The Old Red Sandstone of Brown Clee Hill and the adjacent
area. I. Stratigraphy. Bulletin of the British Museum (Natural History), Geology.
1961;5(7):175–242.

40. White EI. The Old Red Sandstone of Brown Clee Hill and the adjacent area. II.
Palaeontology. Bulletin of the British Museum (Natural History), Geology.
1961;5(7):243–310.

41. Conzen T, Gerlach R. Fische aus dem Unterdevon. Archäologie im Rheinland. 1993;

42. Hartkopf-Fröder C, Knaak M, Weber HM, Wrede V. Der Steinbruch Jaeger bei
Odenspiel – ein Unterdevon-Fossilfundpunkt mit großem Potenzial. Archäologie im
Rheinland. 2015;

43. Becker RT, Marshall JEA, Da Silva A, Agterberg FP, Gradstein FM, Ogg JG. Chapter
22 – The Devonian Period. In: Geologic Time Scale 2020. 2020. p. 733–810.

44. Poschmann M, Schoenemann B, Mccoy VE. Telltale eyes: The lateral visual systems
of Rhenish Lower Devonian eurypterids (Arthropoda, Chelicerata) and their
palaeobiological implications. Palaeontology. 2016;59(2):295–304.

45. Stets J, Schäfer A. The Siegenian delta: Land-sea transitions at the northern margin
of the Rhenohercynian Basin. Geol Soc Spec Publ. 2009;314(September):37–72.

46. Jansen U. Brachiopod faunas, facies and biostratigraphy of the Pridolian to lower
Eifelian succession in the Rhenish Massif (Rheinisches Schiefergebirge, Germany).
Becker, R. T., Ko¨nigshof, P. & Brett, C. E. (eds) 2016. Geological Society, London,
Special Publicat. In: Becker RT, Königshof P, Brett CE, editors. Devonian Climate,
Sea Level and Evolutionary Events. Geological Society, London,; 2016. p. 45–122.
(Special Publications).

47. Weber HM, Hartkopf-Fröder C. Reste eines seltenen Seeskorpions aus dem
Unterdevon von Odenspiel. Archäologie im Rheinland. 2016;52–3.

48. Hartkopf-Fröder C, Weber HM. From Emsian coastal to Famennian marine
environments: palaeogeographic evolution and biofacies in the Bergisch Gladbach-
Paffrath Syncline area (Rhenish Massif, Germany). Münstersche Forschungen zur
Geologie und Palaeontologie. 2016;108:46–75.

49. Schmidt W. Grundlagen einer Pteraspiden-Stratigraphie im Unterdevon der
Rheinische Geosynklinale. Fortschritte in der Geologie von Rheinland und Westfalen.
1959;5.

50. Randle E, Sansom RS. Exploring phylogenetic relationships of Pteraspidiformes
heterostracans (stem-gnathostomes) using continuous and discrete characters. J Syst
Palaeontol [Internet]. 2016;2019(August):1–17. Available from:
http://www.tandfonline.com/doi/full/10.1080/14772019.2016.1208293

51. Blieck A. Le genre Rhinopteraspis Jaekel (Vertébrés, Hétérostracés) du Dévonien
Inférieur. Bull Mus natl Hist nat, Paris, 4è sér. 1980;2(1):25–47.

52. Herbst EC, Meade LE, Lautenschlager S, Fioritti N, Scheyer TM. A toolbox for the
retrodeformation and muscle reconstruction of fossil specimens in Blender. R Soc
Open Sci. 2022;9(8).

53. Lautenschlager S. Reconstructing the past: methods and techniques for the digital
restoration of fossils. R Soc Open Sci. 2016;3(160342).

54. Keating JN, Marquart CL, Donoghue PCJ. Histology of the heterostracan dermal
skeleton: Insight into the origin of the vertebrate mineralised skeleton. J Morphol
[Internet]. 2015;276:657–80. Available from: http://doi.wiley.com/10.1002/jmor.20370

55. Fahlbusch K. Pteraspis dunensis ROEMER eine neuarbeitung der pteraspidenfunde
(agnathen) von Overath. Palaeontographica, Abteilung A. 1957;108:1–56.

56.   Fahlbusch K. Eine Pteraspiden-Fauna aus dem Unterdevon von Alken an der Mosel.
592   Senckenbergiana lethaea. 1966;47(2):165–91.
593   57.   Tarlo B. Rhinopteraspis (McCOY), with notes on the classification and evolution of the
594   pteraspids. Acta Palaeontol Pol. 1961;VI(4).
595   58.   Grohganz M, Ballell A, Ferron HG, Johanson Z, Rayfield EJ, Donoghue PCJ. Testing
596   hypotheses on heterostracan feeding. In Review.
597   59.   Janvier P. The phylogeny of the Craniata, with particular reference to the significance
598   of fossil "agnathans." J Vertebr Paleontol. 1981;1(2):121–59.
599   60.   Janvier P. Early Vertebrates. Clarendon; 1996.
600   61.   Farina SC, Near TJ, Bemis WE. Evolution of the branchiostegal membrane and
601   restricted gill openings in Actinopterygian fishes. J Morphol. 2015;276(6):681–94.
602   62.   Dawson J. the Breathing and Feeding Mechanism of the Lampreys. Biol Bull.
603   1905;9(1):1–21.
604   63.   Dawson J. the Breathing and Feeding Mechanism of the Lampreys. Biol Bull.
605   1905;9(1):1–21.
606   64.   Pernègre VN. The genus Doryaspis white (Heterostraci) from the Lower Devonian of
607   Vestspitsbergen, Svalbard. J Vertebr Paleontol. 2003;22(4):735–46.
608   65.   Gross W. Drepanaspis gemuendenensis Schluter neuuntersuchung. 1963;34.
609   66.   Soehn KL, Wilson MVH. A complete, articulated heterostracan from Wenlockian
610   (Silurian) beds of the Delorme Group, Mackenzie Mountains, Northwest Territories,
611   Canada. J Vertebr Paleontol. 1990;10(4):405–19.
612   67.   Lundgren M, Blom H. Phylogenetic relationships of the cyathaspidids (Heterostraci).
613   GFF [Internet]. 2013 Mar [cited 2014 Nov 18];135(1):74–84. Available from:
614   http://www.tandfonline.com/doi/abs/10.1080/11035897.2013.770792
615   68.   Elliott DK, Lassiter L, Blieck A. A phylogenetic analysis of the Cyathaspididae
616   (Vertebrata: Pteraspidomorphi: Heterostraci). Acta Palaeontol Pol. 2021;66(X).
617   69.   Elliott DK, Petriello MA. New poraspids (Agnatha, Heterostraci) from the Early
618   Devonian of the western United States. J Vertebr Paleontol. 2011;31(3):518–30.
619   70.   Elliott DK, Petriello MA. A new cyathaspid (Agnatha, Heterostraci) with an articulated
620   oral cover from the Late Silurian of the Canadian Arctic. J Vertebr Paleontol.
621   2013;33(1):29–34.
622   71.   Novitskaya LI. Les Amphiaspides (Heterostraci) du Devonien de la Sibérie. Cahiers
623   de Paléontologie, Section Vertebres, Centre National de la Recherche Scientifique.
624   1971;1–130.
625   72.   Purnell MA. Scenarios, selection, and the ecology of early vertebrates. In: Ahlberg PE,
626   editor. Major events in early vertebrate evolution. London: Taylor and Francis; 2001.
627   p. 187–208. (Systematics Association special volume series).
628   73.   Denison RH. Growth and wear of the shield in Pteraspididae (Agnatha).
629   Palaeontographica, Abteilung A. 1973;143:1–10.
630   74.   Rudwick MJS. The function of zigzag deflexions in the commissures of fossil
631   brachiopods. Palaeontology. 1964;7(1):135–71.
632   75.   Ferrón HG, Donoghue PCJ. Evolutionary analysis of swimming speed in early
633   vertebrates challenges the 'New Head Hypothesis.' Commun Biol. 2022;5(1).
634   76.   Gai Z, Li Q, Ferrón HG, Keating JN, Wang J, Donoghue PCJ, et al. Galeaspid
635   anatomy and the origin of vertebrate paired appendages. Nature.
636   2022;609(7929):959–63.

637   77.   Gai Z, Lin X, Shan X, Ferrón HG, Donoghue PCJ. Postcranial disparity of galeaspids
638   and the evolution of swimming speeds in stem-gnathostomes. Natl Sci Rev.
639   2023;10(7).

Supplementary Material Captions

Supplementary Figure 1. *Rhinopteraspis dunensis* NHMUK PV P 73217. A-D, photographs of
complete specimen in (A) left lateral view with full length of preserved rostrum, (B) right
lateral view, (C) close-up of front of head in left lateral view, and (D) close-up of oral region
in ventral view; E-G, Abbreviations: S, sinistral (left), D, dextral (right). Scale bars represent
5 cm in panels A,B, 2.5 cm in panel C, 1 cm in panel D.

648
649

655 **Supplementary Figure 3.** Reconstruction of *Rhinopteraspis dunensis* based on NHMUK PV P
656 73217 with oral plates animated to open to 30 degrees. A-D, lateral view; E-H , anterior
657 view; I-L, dorsal view (with surrounding plates removed). A,E,I, resting position; B,F,J, open
658 to 10 degrees; C,G,K, open to 20 degrees; D,H,L, open to 30 degrees.