A Continuous Extension of Gene Set Enrichment Analysis using the Likelihood Ratio

Test Statistics Identifies VEGF as a Candidate Pathway for Alzheimer's disease

Ali Mahzarnia[1*], Michael W. Lutz[2], Alexandra Badea[1,2,3, 4]

[1] Department of Radiology, Duke University School of Medicine, Durham, NC, USA

[2] Department of Neurology, Duke University School of Medicine, Durham, NC, USA

[3] Biomedical Engineering, Duke University, Durham, NC, USA

[4] Brain Imaging and Analysis Center, Duke University School of Medicine, Durham, NC, USA

*Corresponding Author:

Ali Mahzarnia, PhD

Rm139, Joseph and Kathleen Bryan Research Building

Duke University Medical Center, Durham, NC, 27710

Ali.mahzarnia@duke.edu

919-684-7654

Abstract

Background

17   Alzheimer's disease involves brain pathologies such as amyloid plaque depositions and

18   hyperphosphorylated tau tangles and is accompanied by cognitive decline. Identifying

19   the biological mechanisms underlying disease onset and progression based on

20   quantifiable phenotypes will help understand the disease etiology and devise therapies.

Objective

21   Our objective was to identify molecular pathways associated with AD biomarkers

22   (Amyloid-β and tau) and cognitive status (MMSE) accounting for variables such as age,

23   sex, education, and APOE genotype.

Methods

We introduce a novel pathway-based statistical approach, extending the gene set

likelihood ratio test to continuous phenotypes. We first analyzed independently each of

the three phenotypes (Amyloid-β, tau, cognition), using continuous gene set likelihood

ratio tests to account for covariates, including age, sex, education, and APOE genotype.

The analysis involved a large sample size with data available for all three phenotypes,

allowing for the identification of common pathways.

Results

30   We identified 14 pathways significantly associated with Amyloid-β, 5 associated with tau,

31   and 174 associated with MMSE. Surprisingly, the MMSE outcome showed a larger

32   number of significant pathways compared to biomarkers. A single pathway, vascular

33   endothelial growth factor receptor binding (VEGF-RB), exhibited significant associations

34   with all three phenotypes.

Conclusions.

35   The study's findings highlight the importance of the VEGF signaling pathway in aging in

36   AD. The complex interactions within the VEGF signaling family offer valuable insights for

37   future therapeutic interventions.

38   **Keywords:** Continuous GSLRT, Likelihood Ratio-Derived Test Statistics, Alzheimer's Disease,

39   RNA-Seq

Introduction

One strategy to better understand and accurately model late-onset Alzheimer's disease, is to

incorporate various genetic, clinical, and environmental factors into a cohesive model. This

model should establish connections between measurable biomarkers and risk factors. Several

key factors play a significant role in shaping the risk of Alzheimer's disease (AD). These factors

encompass both genetic influences, such as APOE genotype and sex, as well as environmental

elements, including education level, diet, and age. In recent years, the identification and

characterization of AD have been facilitated by the use of biomarkers like Amyloid-β (Aβ),

phosphorylated tau (tau), and measures of neurodegeneration. Furthermore, AD is

distinguished by memory impairment, often evaluated using the Minimental State Evaluation

(MMSE).

AD has a deleterious impact on American lives, as over 6 million individuals are currently

afflicted with Alzheimer's, with a projected twofold increase by 2050. The disease's high

mortality rate claims 1 in 3 seniors while imposing significant economic burdens, costing the

nation billions [1]. However, the biological background conducive to developing AD remains

unknown. The aim of this study was to address the existing knowledge gap by pinpointing

molecular pathways that play a crucial role in modulating levels of hallmark AD pathologies, as

well as memory function.

Recent publications have revealed a role for multiple pathways in AD, based on brain

proteomics and transcriptomic analyses [2, 3]. These studies have revealed pathways that are

relevant not only to neurons but also to cells regulating response to inflammation [4], such as

microglia [5]. Interestingly endothelial cells, astrocytes and neurons that control neurovascular

functions have been shown to play an important role in AD [6]. Other cell types and subcellular

components such as mitochondria may be involved [7]. [8] identifies a novel brain-enriched

RING finger E3 ligase, RNF182, which shows elevated expression in AD brains and may play a

role in controlling neurotransmitter release. Pathways involved with filament-based processes,

cellular detoxification, and wound healing have also been involved [2, 3]. A decline in sensory

function, including taste has also been reported with aging and AD [9]. Importantly, the vascular

endothelial growth factor has been associated with AD (VEGF) [10, 11], and while its role in

neurodegeneration is yet to be understood, it presents a druggable target for therapies.

However, most studies focused on comparisons of two or three groups of subjects using

discrete classification variables, such as case/control, without accounting for the relationships

between multiple hallmark biomarkers. Here we propose an approach to detect common gene

pathways based on RNA-Seq changes associated with continuous-scale changes in multiple

biomarkers and clinical phenotypes.

Our study involves developing a statistical approach centered around gene pathways. This

method incorporates estimations of Amyloid-β and tau tangle brain levels, along with memory

scores from the MMSE, and integrates them into a comprehensive statistical model.

Additionally, we include disease-relevant traits such as age, sex, education, and APOE

genotype in this model. The primary goal is to rank the pathways that undergo alterations in AD,

considering the influence of each of the biomarkers. By doing so, we can effectively identify the

shared pathways across the three domains: amyloid, tau, and MMSE scores.

Moreover, our approach factors in the unique characteristics of human subjects, including age-

specific, sex-specific, education-specific, and APOE genotype-specific differences. It also

evaluates the significance of the relationship between pathway-level interactions and the

presence of AD in relation to each of these factors. We conduct analyses using human

transcriptomic data on Alzheimer's disease progression and explore interactions with the APOE

genotype.

To identify relevant gene sets [12] introduced Gene Set Enrichment Analysis (GSEA), a robust

analytical method for interpreting gene expression data from genome-wide RNA analysis. GSEA

focuses on gene sets—groups of genes with shared biological function, chromosomal location,

or regulation—and demonstrates effectiveness in identifying common biological pathways e.g.

for cancer-related data sets, such as leukemia and lung cancer, where single-gene analysis falls

short. This method using the Kolmogorov-Smirnov statistics has a limitation in that it does not

account for gene-gene interactions. Still, it has spurred the development of many other methods

[13]. Other statistics have been proposed as well, e.g. the gene set likelihood ratio test (gsLRT),

which uses a logistic regression model [14]. However, this particular model has its limitations as

it is only applicable to binary outcome variables. In contrast, our study deals with continuous-

scale variables, specifically amyloid, tau, and MMSE scores. As a result, we require a different

model that can accommodate such continuous data. Additionally, we incorporated

transcriptomic data, encompassing more than 20 thousand genes, as opposed to the mere 142

genes used in the referenced gsLRT study involving proteomics.

Methodology

102   Our proposition introduces a novel approach utilizing all three phenotypes and biomarkers

103   (amyloid, tau, and MMSE scores) to identify shared and meaningful pathways associated with

104   them. Our methodology harnesses gene-level data, encompassing transcriptomics. By utilizing

105   this data, we generate significance scores at the biological pathway level, serving as indicators

106   of the novelty of observed experimental outcomes.

Data and preprocessing

The data sample was taken from a subset of the Religious Orders Study and Rush Memory and

Aging Project (ROSMAP) dataset [15-17] that had RNAseq data available from the dorsolateral

frontal cortex. ROS has enlisted nuns and brothers since 1994. MAP recruited individuals from

the northern Illinois region since 1997. Both studies were run by the same investigators using

similar data collection techniques. Thus, the results from both were comparable. For the

analyses reported in this paper, the clinical consensus diagnoses of Alzheimer's disease or mild

cognitive impairment were used to define a case while the diagnosis of no cognitive

impairment/no impaired domains defined controls. Additional covariates for the statistical

models were age, sex, education, and APOE genotype. The total sample with both gene

expression and clinical data contained 634 subjects, with 433 cases and 201 controls.

Demographic information for the sample is summarized in Table 1.

Group	N	age (mean)	age (SD)
All	634	87.93	6.70
Female	406	89.02	6.57
Male	228	85.99	6.51

Table 1- Demographic Information of Participants by Gender. This table presents the

demographic information for a sample of 634 participants, categorized by gender. It includes the

mean age and standard deviation (SD) for each group.

In the context of [18], the amyloid and tangles definitions and measurements are as follows.

The overall amyloid level is determined as the mean percentage of cortex occupied by Amyloid-

β protein in eight specific brain regions. This measurement is obtained through molecular-

specific immunohistochemistry, where the Amyloid-β protein is targeted and quantified using

image analysis techniques. The Amyloid-β score is calculated in eight brain regions, namely the

hippocampus, entorhinal cortex, midfrontal cortex, inferior temporal cortex, angular gyrus,

calcarine cortex, anterior cingulate cortex, and superior frontal cortex. At least four of these

regions are required to calculate the mean Amyloid-βscore.

Tangle density is determined as the mean density of neuronal neurofibrillary tangles in eight

specific brain regions. These tangles are identified using molecular-specific

immunohistochemistry, employing antibodies specific to abnormally phosphorylated Tau protein,

known as AT8. The cortical density of tangles is measured per square millimeter using

systematic sampling. The tangle score is calculated as the mean density in eight brain regions,

including the hippocampus, entorhinal cortex, midfrontal cortex, inferior temporal cortex, angular

gyrus, calcarine cortex, anterior cingulate cortex, and superior frontal cortex. A minimum of

four regions is necessary to compute the mean tangle density.

Clinical phenotypic information for the sample is summarized in Table 2.

Cognitive state/Phenotypes	N Rows	MMSE (mean)	MMSE (SD)	Amyloid (mean)	Amyloid (SD)	log(tangles) (mean)	log(tangles) (SD)
No cognitive impairment	201	28.21	1.6	2.78	3.44	0.32	1.58
MCI: Mild cognitive impairment (one impaired domain) and NO other cause of CI	158	25.89	3.38	4.03	4.63	1.07	1.28
MCI: Mild cognitive impairment (One impaired domain) AND another cause of CI	10	25.19	3.73	1.35	1.33	-0.03	1.3
AD: Alzheimer's dementia and NO other cause of CI (NINCDS PROB AD)	220	13.49	8.22	5.72	4.36	1.83	1.25
AD: Alzheimer's dementia AND another cause of CI (NINCDS POSS AD)	33	15.49	7.94	3.62	3.69	1.41	0.97

Other dementia: Other primary cause of dementia	12	15.17	5.47	2.59	3.6	0.5	1.31

Table 2-Clinical Phenotypic Data for Different Cognitive States. This table presents key clinical phenotypic data for different cognitive states in a diverse sample population. The study includes individuals with no cognitive impairment, mild cognitive impairment (MCI), Alzheimer's dementia (AD), and other primary causes of dementia. The table provides mean scores and standard deviations for Mini-Mental State Examination (MMSE), amyloid levels, and log(tangles) for each cognitive state/phenotype. The findings highlight distinct cognitive profiles and potential biomarkers associated with various cognitive conditions, contributing to better understanding and targeted interventions for cognitive disorders.

The RNAseq data was obtained from the Accelerating Medicines Partnership Program for Alzheimer's Disease Data Knowledge Portal (https://adknowledgeportal.synapse.org/), specifically, the RNA-seq Harmonization study (https://www.synapse.org/#!Synapse:syn9702085). The ROSMAP data from this study was used to create a combined dataset of RNA-seq data in combination with the three clinical phenotypes of MMSE, amyloid burden, and tangles [19]. The RNA-seq Harmonization study has the goal of creating an RNA-seq database based on a consensus set of analytical tools. The methodological details of the RNA-seq processing are given in Wan et al [20] and at the RNA-Seq reprocessing study website for the ROSMAP project (https://www.synapse.org/#!Synapse:syn8456629). In brief, RNA was extracted from samples consisting of approximately 100 mg of gray matter tissue from the dorsolateral prefrontal cortex. The RNA samples were prepared and sequenced as described in [19]. The reprocessing of the RNAseq data was done using a consensus set of tools with only library type-specific parameters varying between pipelines. Picard (https://broadinstitute.github.io/picard/) was used to generate FASTQs from source BAMs. Generated FASTQ reads were aligned to the GENCODE24

(GRCh38) reference genome using STAR [21] and gene counts were computed for each

sample. To evaluate the quality of individual samples and to identify potentially important

covariates for expression modeling, two sets of metrics were computed using the

CollectAlignmentSummaryMetrics and CollectRnaSeqMetrics functions in Picard. To account for

differences between samples, studies, experimental batch effects, and unwanted RNA-Seq

specific technical variations library normalization and covariate adjustments for each study

separately using fixed/mixed effects modeling. The workflow consists of the following steps: (i)

gene filtering: Genes that are expressed more than 1 CPM (read Counts Per Million total reads)

in at least 50% of samples in each tissue and diagnosis category were used for further analysis,

(ii) conditional quantile normalization was applied to account for variations in gene length and

GC content, (iii) sample outlier detection using principal component analysis and clustering, (iv)

Covariates identification and adjustment, where confidence of sampling abundance were

estimated using a weighted linear model using voom-limma package in Bioconductor [22]. For

the differential expression analysis, fixed/mixed effect linear regression was used with the

following models: gene expression ~Diagnosis + Sex + covariates + (1| Donor) or gene

expression ~Diagnosis x Sex + covariates + (1|Donor), where each gene is linearly regressed

independently with Diagnosis, a variable explaining the AD status of an individual, identified

covariates, and donor information as a random effect. Observation weights (if any) were

calculated using the voom-limma [22] pipeline such that observations with higher presumed

precision will be up-weighted in the linear model fitting process.

The data used for the gsLRT analysis included the phenotypes MMSE, amyloid burden and

log(tangles). For gene expression, normalized and filtered counts were used as per the

description   on   the   RNA-Seq   reprocessing   study   website

(https://www.synapse.org/#!Synapse:syn8456629). The specific datasets used for this study

are: RNAseq   data   is   obtained   from   (https://www.synapse.org/#!Synapse:syn8456638),

covariates are recorded in (https://www.synapse.org/#!Synapse:syn11024258), differential

expression results are from (https://www.synapse.org/#!Synapse:syn8456721) and clinical

phenotypic data was obtained from the ROSMAP data sharing resource

(https://www.radc.rush.edu/). These datasets were organized into the specific matrices for gene

expression results, covariates, and phenotypes required by the gsLRT program.

Data and Code Availability

192   ROSMAP resources can be requested at https://www.radc.rush.edu. The Code is available at

193   https://github.com/Ali-Mahzarnia/gslrt2.

194
195

Statistical Model

To derive the pathways associated with each AD related phenotypes (amyloid, tau, or MMSE

scores), we define the following hypothesis tests. In our models, we utilize a matrix $G \in \mathbb{R}^{n \times m}$ to

represent measurement values (e.g., RNA expression) for m genes and n samples. Additionally,

we employ a matrix, $X \in \mathbb{R}^{n \times d}$, which contains d=4 covariates such as "sex," years of

education", "APOE Genotype" (APOE2,3,4), and "age at death" associated with the samples.

Moreover, we utilize a continuous vector $y \in \mathbb{R}^{n \times 1}$ for the n samples that is the phenotype

measurements (amyloid, tau, or MMSE scores). To conduct data analysis, we employ nested

models for each gene g_j, $j = 1, \ldots, m$ and all samples $i = 1, \ldots, n$ as outlined below:

$$H_0 : y_i = x_i \beta^j + g_i^j \alpha^j + e_i$$

$$H_a : y_i = x_i \beta^j + g_i^j \alpha^j + Genotype_i * g_i^j \gamma^j + e_i$$

where $e_i \sim N(0, \sigma)$. We denote the maximum likelihood estimator of the two above models with

$\widehat{\omega}_0 = (\widehat{\beta_0}, \widehat{\alpha_0})$, $\widehat{\omega}_a = (\widehat{\beta_a}, \widehat{\alpha_a}, \widehat{\gamma^j})$.

In order to assess the enhanced explanatory capacity of the interaction term between the gene

measurement profile (g_j) and genotype (APOE2,3,4), in contrast to the simpler model that

merely includes the covariate matrix X and (g_j), we introduce statistical measures that evaluate

the disparity in the models via log-likelihoods at the gene level. These statistics act as our metric

for quantifying the supplementary explanatory power.

$$\Lambda^j = 2\,[\,l\,(\hat{\omega}_a) - l(\hat{\omega}_0)]$$

211   where $l(.)$, is the log-likelihood. We define the enrichment score for gene set G_j:

$$ES = \sum_{j \in G_j} \Lambda^j$$

We calculate p-values for enrichment scores through nperm = 10,000 permutation sampling,

which provides more conservative estimates but is computationally intensive. Since gene set

tests share overlapping membership and exhibit interdependence, the presented p-values in

this context and subsequent tables have not been adjusted for multiple testing. Consequently,

they do not possess theoretical guarantees for controlling the False Discovery Rate (FDR).

Once the pathways are sorted by their significance or ES, identifying the shared pathways

among the three model runs becomes possible.

The model utilized in this study closely resembles that developed by Bryan et al [14], based on

a logistic model, whereas our model accommodates a continuous outcome variable.

Results

Using public resources from RNA-seq analyses of prefrontal cortex in ROSMAP participants, we

have identified pathways associated for Amyloid-β, tau tangles and MMSE, accounting for

covariates i.e. sex, genotype, education, age, and the interactions between RNA expression

levels and APOE genotype.

We have identified significant pathways when examining the outcome of Amyloid-β. We have

chosen a p-value threshold of 0.1 instead of the conventional 0.05 for hypothesis decision-

making in order to capture a broader range of potentially significant pathways. This slightly

relaxed threshold allows for a more inclusive analysis, potentially uncovering additional

pathways that may contribute to our understanding of the relationship between Amyloid-β and

the identified pathways. Table 3 and Figure 1 present the 14 significant pathways associated

with Amyloid-β, such as TRANSLOCASE ACTIVITY, C3HC4 TYPE RING FINGER DOMAIN

BINDING, INSULIN RECEPTOR BINDING, and VASCULAR ENDOTHELIAL GROWTH

FACTOR RECEPTOR 2 BINDING.

Gene Pathways	p-value	ES
GOMF DNA TRANSLOCASE ACTIVITY	0.032	3.761
GOMF C3HC4 TYPE RING FINGER DOMAIN BINDING	0.040	3.047
GOMF VASCULAR ENDOTHELIAL GROWTH FACTOR RECEPTOR 2 BINDING	0.043	2.445
GOMF SUMO POLYMER BINDING	0.059	2.637
GOMF ATP DEPENDENT PROTEIN DISAGGREGASE ACTIVITY	0.064	2.468
GOMF ACETYLESTERASE ACTIVITY	0.067	2.315
GOMF DOPAMINE NEUROTRANSMITTER RECEPTOR ACTIVITY	0.082	2.383
GOMF BH DOMAIN BINDING	0.084	2.074
GOMF INSULIN RECEPTOR BINDING	0.086	2.217
GOMF SMALL RIBOSOMAL SUBUNIT RRNA BINDING	0.087	2.002
GOMF CLASS I DNA APURINIC OR APYRIMIDINIC SITE ENDONUCLEASE ACTIVITY	0.091	2.298
GOMF SHORT CHAIN CARBOXYLESTERASE ACTIVITY	0.094	2.001
GOMF VASCULAR ENDOTHELIAL GROWTH FACTOR RECEPTOR BINDING	0.098	1.787
GOMF ALPHA N ACETYLGALACTOSAMINIDE ALPHA 2 6 SIALYLTRANSFERASE ACTIVITY	0.099	1.869

Table 3- Significant Pathways Associated with Amyloid-β Using a P-value Threshold of 0.1. This

table presents the pathways that show significance (p-values smaller than 0.1) in relation to

Amyloid-β,. The pathways listed provide insights into potential biological mechanisms and

molecular processes associated with Amyloid-β.

Figure 1-Pathway Analysis of Amyloid-β. Each pathway is represented on the y-axis, while the

corresponding -log10(P-value) is represented by bars parallel to the x-axis, positioned in front of

each pathway. The length of the bars reflects the statistical significance of the pathways with

amyloid burden; longer bars indicating greater significance. This visualization allows for a quick

assessment of the significance levels for each pathway, aiding in the identification of key

pathways associated with Amyloid-β.

The 5 significant pathways associated with log(tangles) are presented in Table 4 and illustrated

in Figure 2 such as BITTER TASTE RECEPTOR ACTIVITY, PROTEIN GLUTAMINE GAMMA

GLUTAMYLTRANSFERASE ACTIVITY, TASTE RECEPTOR ACTIVITY, and VASCULAR

ENDOTHELIAL GROWTH FACTOR RECEPTOR 2 BINDING.

Gene Pathways	p-value	ES
GOMF BITTER TASTE RECEPTOR ACTIVITY	0.035	3.547
GOMF PROTEIN GLUTAMINE GAMMA GLUTAMYLTRANSFERASE ACTIVITY	0.055	2.658
GOMF TASTE RECEPTOR ACTIVITY	0.066	2.486
GOMF VASCULAR ENDOTHELIAL GROWTH FACTOR RECEPTOR 2 BINDING	0.097	1.910
GOMF HYALURONONGLUCOSAMINIDASE ACTIVITY	0.098	2.074

250   *Table 4- Significant Pathways Associated with log(tangles). This table shows the pathways that*

251   *exhibited statistical significance (p-values smaller than 0.1) in relation to log(tangles).*

Figure 2- Pathway Analysis of log(tangles). This figure presents the results of the pathway significance analysis for log(tangles). Each pathway is displayed on the y-axis, while bars parallel to the x-axis represent the corresponding -log10(P-value). The length of each bar reflects the statistical significance of the pathway association with log(tangles). Longer bars indicate greater significance/smaller p values.

In addition to the pathways depicted in Figure 3, Table 5 includes the top candidate of such pathways, which represent a subset of significant associations (with p-values smaller than 0.1) with the Mini-Mental State Examination (MMSE) outcome. These results highlight specific pathways that exhibit statistical significance in association with MMSE. Figure 3 provides a partial representation of 174 top pathways, including TRAIL BINDING, VASCULAR ENDOTHELIAL GROWTH FACTOR RECEPTOR BINDING, VASCULAR ENDOTHELIAL GROWTH FACTOR RECEPTOR 2 BINDING.

Gene Pathways	p-value	ES
GOMF TRAIL BINDING	0.0010	7.071
GOMF VASCULAR ENDOTHELIAL GROWTH FACTOR RECEPTOR BINDING	0.0018	4.114
GOMF VASCULAR ENDOTHELIAL GROWTH FACTOR RECEPTOR 2 BINDING	0.0021	4.747
GOMF DEATH RECEPTOR ACTIVITY	0.0023	4.800
GOMF MECHANOSENSITIVE MONOATOMIC ION CHANNEL ACTIVITY	0.0027	4.061
GOMF HEXOKINASE ACTIVITY	0.0031	4.691
GOMF TUMOR NECROSIS FACTOR BINDING	0.0032	5.289
GOMF CC CHEMOKINE BINDING	0.0049	4.886
GOMF NERVE GROWTH FACTOR BINDING	0.0049	4.603
GOMF RAGE RECEPTOR BINDING	0.0060	3.650

Table 5- Significant Gene Pathways Associated with MMSE Outcome. It highlights the top

candidate pathways, including TRAIL BINDING and VASCULAR ENDOTHELIAL GROWTH

FACTOR RECEPTOR BINDING, among others.

Figure 3- Pathway Analysis of MMSE. This figure shows a portion of the pathway significance

analysis results for MMSE. Each pathway is plotted on the y-axis, accompanied by bars parallel

to the x-axis that represent the corresponding -log10(P-value). The length of each bar reflects

the statistical significance of the pathway, with longer bars indicating higher significance. This

visual representation enables a quick evaluation of the significance levels associated with each

pathway. It assists in identifying key pathways that are linked to MMSE, contributing to a better

understanding of the underlying mechanisms influencing MMSE.

Finally, we investigated the shared pathways among all 3 studies involving Amyloid-β,

log(tangles), and MMSE, shown in Figure 4. This analysis allows us to gain insights into the

shared biological mechanisms and molecular processes that may contribute to the interplay

between Amyloid-β, log(tangles), and MMSE.

Figure 4- Common Significant Pathways Between for Amyloid-β, log(tangles), and MMSE. This

Venn diagram illustrates the common pathways identified among the studies involving Amyloid-

β, log(tangles), and MMSE. The diagram consists of overlapping circles that represent each

study, with labeled sections indicating the shared pathways among them. The shared pathways

are listed within the diagram, providing a concise overview of the biological processes and

molecular mechanisms that are consistently implicated across these phenotypes. This analysis

highlights the interconnectedness of these factors and underlying pathways that contribute to

the associations between Amyloid-β, log(tangles), and MMSE.

Results for the comparisons of expression differences between control and AD samples are

shown   in   Table   6   as   reported   in   the   AMP-AD   analysis   of   the   dataset

(https://www.synapse.org/#!Synapse:syn8456721).   Higher expression in AD samples relative

to normal cognition is observed for VEGFA, VEGFB, VEGFD, PGF and FLT1 while lower

expression in AD samples relative to controls is observed for FLT4, KDR, NRP1 and NRP2.

The only significant difference between the two groups after adjusting for multiple comparisons

is for VEGFB.

Gene	Comparison	logFC	CI.L	CI.R	AveExpr	t	P.Value	adj.P.Val
VEGFA	AD-CONTROL	0.16	-0.08	0.40	1.86	1.28	0.20	0.43
VEGFB	AD-CONTROL	0.10	0.03	0.16	5.67	2.81	0.01	0.05
VEGFD	AD-CONTROL	0.02	-0.07	0.12	1.10	0.50	0.62	0.79
PGF	AD-CONTROL	0.15	-0.02	0.32	0.17	1.69	0.09	0.26
FLT1	AD-CONTROL	0.14	-0.05	0.32	3.86	1.44	0.15	0.36
FLT4	AD-CONTROL	-0.03	-0.15	0.09	0.15	-0.54	0.59	0.78
KDR	AD-CONTROL	-0.07	-0.20	0.07	1.05	-1.00	0.32	0.55
NRP1	AD-CONTROL	-0.07	-0.18	0.03	2.14	-1.41	0.16	0.37
NRP2	AD-CONTROL	-0.02	-0.10	0.06	1.37	-0.58	0.56	0.75

299   *Table 6- Differential Expression of VEGF family related Genes in Alzheimer's Disease (AD)*

300   *Samples Compared to Controls. This table presents the results of a comparative analysis*

301   *between control and Alzheimer's disease (AD) samples, focusing on the expression differences*

302   *of VEGF-family--related genes.*

Discussion

The progression of Alzheimer's disease (AD) manifests through changes in biomarkers that

reflect abnormal protein expression, such as Amyloid-β, phosphorylated tau, as well as clinically

measurable symptoms including memory decline. In our study, we developed new methods to

reveal pathways related to changes in brain RNA-Seq for each of these two neuropathological

biomarkers, as well as for MMSE. Surprisingly we observed a larger number of significant

pathways for gene expression association with MMSE (174 pathways) than those associated

with the two hallmark biomarkers for AD, Amyloid-β (14 pathways), and tau tangles (5

pathways). More importantly, we identified a single pathway, vascular endothelial grown factor

receptor binding (VEGF-RB) that was associated with differences reflective of all three

phenotypes analyzed by the continuous gene set likelihood ratio test (gsLRT).

The methodological advance of this study was the extension of gsLRT from binary (e.g.

disease/control status) to continuous value phenotypes. Our study extends prior work by

analyzing three continuous scale phenotypes (Amyloid-β , tau tangles, MMSE) in a pathway

analysis that accounts for the covariates age, sex and APOE genotype (gsLRT for continuous

phenotypes).

Among the pathways identified as significant for Amyloid-β we noted several candidate

pathways that support changes in DNA repair ability (GOMF DNA TRANSLOCASE ACTIVITY)

[23], cell mediated immunity (GOMF C3HC4 TYPE RING FINGER DOMAIN BINDING) [24],

apoptosis (GOMF BH DOMAIN BINDING) [25], protein synthesis (GOMF SMALL RIBOSOMAL

SUBUNIT RRNA BINDING) [26] and disaggregation (GOMF ATP DEPENDENT PROTEIN

DISAGGREGASE ACTIVITY), as well as insulin signaling (GOMF INSULIN RECEPTOR

BINDING) [27, 28], which have all been connected to AD.

Among the pathways identified as significant for tau tangles we noted two that support

alterations in sensory processing, pointing to taste (GOMF BITTER TASTE RECEPTOR

ACTIVITY; GOMF TASTE RECEPTOR ACTIVITY). Recent studies have shown alterations in

the sour taste [29], while here we identified changes in pathways associated with bitter taste,

and taste in general. Since taste and olfaction are closely linked, these results suggest possible

changes in olfactory function. Among sensory changes in AD, olfaction has been proposed as

one of the more promising biomarkers for early detection [30]. We also noted pathways pointing

to differences in the transfer of amino acids across the membrane, cell survival during oxidative

stress (GOMF PROTEIN GLUTAMINE GAMMA GLUTAMYLTRANSFERASE ACTIVITY) [31],

and glutathione homeostasis, relevant to several neurodegenerative diseases, such as AD, PD

and ALS [31, 32]. Finally, the role of extracellular matrix was suggested by the presence of the

GOMF HYALURONONGLUCOSAMINIDASE ACTIVITY pathway, which has a less understood

but complex role in aging and disease [33, 34].

Among the large number of pathways related to MMSE, we noted top candidates involved in

apoptosis through TRAIL which binds to death receptors, suggesting a relation with immune

related mechanisms [35].

Amyloid-β and tau biomarkers in combination with measures of neurodegeneration have been

associated with the progression of AD neuropathology and memory loss [36-41]. We note that

neuropsychological measures such as MMSE can reflect other causes of cognitive impairment

including damage to the cerebrovascular system although individuals with moderate to severe

Alzheimer's disease tend to have MMSE scores less than 15 [42].

The role of the VEGF signaling family in neurodegeneration and Alzheimer's disease has been

extensively studied including with multiomic approaches that analyzed bulk RNA sequencing

data, single nucleus sequencing data, and mass spectrometry proteomics data [10]. As a

consequence of the numerous pathways that intersect with VEGF receptor signaling, it has

been difficult to identify the specific receptors and molecules that associate with disease

endophenotypes or covariates including age and sex.

VEGF includes a family of five ligands (VEGFA, VEGFB, VEGFC, VEGFD, and PGF), three

tyrosine kinase receptors (FLT1, FLT4, and KDR), and two modulating receptors (NRP1 and

NRP2). The pathway identified as significantly associated with the three phenotypes (GOMF

VASCULAR ENDOTHELIAL GROWTH FACTOR RECEPTOR BINDING (GO:0005172)

includes the ligands and several other genes.   The pathway GOMF VASCULAR

ENDOTHELIAL GROWTH FACTOR RECEPTOR ACTIVITY (GO:0005021) includes the

tyrosine kinase receptors and modulating receptors).

The presence of VEGF alongside Amyloid-β plaques in AD brains and its strong binding to

Amyloid-β suggest that VEGF may contribute to neurodegeneration and vascular dysfunction

[43]. Additionally, Amyloid-β inhibits VEGF receptor signaling, impairing angiogenesis [44].

VEGF accumulation around amyloid plaques interacts directly with Amyloid-β, rescuing synaptic

dysfunction caused by the toxic Amyloid-β oligomers [45].

VEGF also interacted with tau and amyloid-β_{42}, predicting hippocampal atrophy and memory

decline. Another study [46] revealed that VEGF genes, particularly FLT4 and FLT1, were

associated with AD neuropathology and cognition. Higher levels of VEGF were associated with

slower hippocampal atrophy and better cognitive function [47]. These findings emphasize the

importance of understanding the relationship between VEGF, tau pathology, and AD for

potential therapeutic interventions.

Several studies have investigated the role of vascular endothelial growth factor (VEGF) in

cognitive impairment. For example one study found that VEGF AA genotype is associated with

an increased risk of developing AD and MCI, while higher VEGF levels are observed in AD

patients [48]. Another study found lower VEGF levels in AD patients and amnestic MCI patients

compared to controls, correlating with cognitive decline [49]. Additionally, [50] finds that higher

serum VEGF levels in ischemic stroke patients are associated with post-stroke cognitive

impairment. Conversely, [51] shows that VEGF signaling is crucial for maintaining cognition and

neurogenesis, cautioning against inhibiting VEGF signaling. Interestingly, [52] demonstrates that

VEGF levels increase during the early stage of AD but decrease as the disease progresses,

suggesting a link between VEGF levels and cognitive decline. VEGF produced by macrophages

plays a role in preserving cognitive function in obesity, which can be a risk factor during aging

and AD [53] However the literature is still controversial and more work is needed to understand

the role of various VEGF isoforms role in modulating cognition [54] .

The ensemble of molecules in the VEGF pathways and their interactions have been reported to

have varied effects on AD phenotypes. The members of the VEGF measured in the brain and

blood have been characterized with respect to cognitive performance, neural and

cerebrovascular pathology, and CSF biomarkers [11, 55]. Blood and brain VEGFA has been

reported to be protective against memory impairment and brain atrophy in AD [10]. Higher

expression levels of VEGFB, PGF, FLT1 and FLT4 are reported to be associated with faster

cognitive decline and greater neuropathological lesion development [10, 55] and our data

supported this direction of difference in expression levels. The VEGF family is involved in

multiple signaling pathways, leading to potentially different effects on AD-related phenotypes;

for example, VEGFA can signal through KDR or FLT1 where the receptors can elicit effects in

opposing directions [10]. In a study of microglial control of astrocytes in response to microbial

metabolites, microglial VEGFB was shown to trigger FLT1 signaling in astrocytes and promote

CNS inflammation [56]. Neutropilin expression (NRP1 and NRP2) was decreased AD samples

relative to samples from individuals with normal cognition in agreement with prior evidence [10].

NRP1 and NRP2 have well-established roles in angiogenesis[10, 57]. Moreover, interactions

between VEGF family proteins with APOE have also been reported. Higher levels of VEGFA

were reported to be associated with worse outcomes among APOE E4 carriers and better

outcomes among non-E4 carriers [10].

Our study has several strengths. First, the extension of the gsLRT to continuous phenotypes

allowed a higher statistical level (continuous vs. nominal) of data for the pathway analysis. The

relatively large sample size where data was available for all three phenotypes allowed for

comparisons at the gene and pathway level. For the pathway/signature analysis, well-

established databases including gene ontology were used to enable replication studies and

other future work.

A limitation of the analysis is that the phenotypes are measured at a single time point.

Longitudinal data for the Amyloid-β and tau biomarkers are not available from post-mortem

brains, though nuclear imaging or fluid biomarkers present a great promise for the future of such

longitudinal studies. MMSE is measured over time, so another possible phenotype to consider

would be the decline in MMSE from baseline to death. Studies that examine changes in CSF

and blood biomarkers for VEGF over time will provide information on temporal relationships

between VEGF family mRNA and protein concentrations and biomarker changes and can open

new avenues for exploiting its therapeutic potential [10].

Our study proposes a method for pathway identification using continuous phenotypes and public

data bases on RNA-Seq, but can potentially be used for proteomic analyses, or extended to

multinetwork omic studies and make use of extensive public data base resources or de novo

analyses to better understand the mechanistic substrates for neurodegenerative diseases.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or

financial relationships that could be construed as a potential conflict of interest.

Funding

This work was supported by RF1 AG057895, R01 AG066184, U24 CA220245, RF1 AG070149,

P30 AG072958.

Acknowledgment

425   We thank all the participants of ROS and MAP studies. These studies were funded by the

426   National Institute on Aging: P30AG010161 ADCC, P30AG072975 ADRC, R01AG015819 RISK,

427   R01AG017917 MAP, U01AG46152 AMP-AD Pipeline I, U01AG61356 AMP-AD Pipeline II.

428   Furthermore, we are grateful to Dr. Jordan Bryan for helpful discussions.

References

1. Association, A.s., *2019 Alzheimer's disease facts and figures.* Alzheimer's & dementia, 2019. **15**(3): p. 321-387.

2. Seyfried, N.T., et al., *A Multi-network Approach Identifies Protein-Specific Co-expression in Asymptomatic and Symptomatic Alzheimer's Disease.* Cell Systems, 2017. **4**(1): p. 60-72.e4.

3. Huang, Z., et al., *Brain proteomic analysis implicates actin filament processes and injury response in resilience to Alzheimer's disease.* Nature Communications, 2023. **14**(1).

4. McGeer, E.G. and P.L. McGeer, *The importance of inflammatory mechanisms in Alzheimer disease.* Experimental Gerontology, 1998. **33**(5): p. 371-378.

5. Ofengeim, D., et al., *RIPK1 mediates a disease-associated microglial response in Alzheimer's disease.* Proceedings of the National Academy of Sciences of the United States of America, 2017. **114**(41): p. E8788-E8797.

6. Sweeney, M.D., S. Ayyadurai, and B.V. Zlokovic, *Pericytes of the neurovascular unit: Key functions and signaling pathways.* Nature Neuroscience, 2016. **19**(6): p. 771-783.

7. Zhang, Y., et al., *Identification of mitochondrial related signature associated with immune microenvironment in Alzheimer's disease.* Journal of Translational Medicine, 2023. **21**(1).

8. Liu, Q.Y., et al., *A novel brain-enriched E3 ubiquitin ligase RNF182 is up regulated in the brains of Alzheimer's patients and targets ATP6V0C for degradation.* Molecular neurodegeneration, 2008. **3**: p. 1-16.

9. Sakai, M., et al., *Decline of gustatory sensitivity with the progression of Alzheimer's disease.* International Psychogeriatrics, 2016. **28**(3): p. 511-517.

10.   Seto, M., et al., *Multi-omic characterization of brain changes in the vascular endothelial*

growth factor family during aging and Alzheimer's disease. Neurobiol Aging, 2023. **126**:

p. 25-33.

11.   Libby, J.B., et al., *Whole blood transcript and protein abundance of the vascular*

endothelial growth factor family relate to cognitive performance. Neurobiol Aging, 2023.

124: p. 11-17.

12.   Subramanian, A., et al., *Gene set enrichment analysis: a knowledge-based approach for*

interpreting genome-wide expression profiles. Proceedings of the National Academy of

Sciences, 2005. **102**(43): p. 15545-15550.

13.   Tamayo, P., et al., *The limitations of simple gene set enrichment analysis assuming*

gene independence. Stat Methods Med Res, 2016. **25**(1): p. 472-87.

14.   Bryan, J., et al., *Likelihood ratio statistics for gene set enrichment in Alzheimer's disease*

pathways. Alzheimer's & Dementia, 2021. **17**(4): p. 561-573.

15.   Bennett, D.A., et al., *Overview and findings from the Religious Orders Study.* Curr

Alzheimer Res, 2012. **9**(6): p. 628-45.

16.   Bennett, D.A., et al., *Overview and findings from the Rush Memory and Aging Project.*

Curr Alzheimer Res, 2012. **9**(6): p. 646-63.

17.   De Jager, P.L., et al., *Alzheimer's disease: early alterations in brain DNA methylation at*

ANK1, BIN1, RHBDF2 and other loci. Nat Neurosci, 2014. **17**(9): p. 1156-63.

18.   Bennett, D.A., et al., *Religious orders study and rush memory and aging project.* Journal

of Alzheimer's disease, 2018. **64**(s1): p. S161-S189.

19.   De Jager, P.L., et al., *A multi-omic atlas of the human frontal cortex for aging and*

Alzheimer's disease research. Sci Data, 2018. **5**: p. 180142.

20.   Wan, Y.W., et al., *Meta-Analysis of the Alzheimer's Disease Human Brain Transcriptome*

and Functional Dissection in Mouse Models. Cell Rep, 2020. **32**(2): p. 107908.

21. Dobin, A., et al., *STAR: ultrafast universal RNA-seq aligner.* Bioinformatics, 2013. **29**(1):

p. 15-21.

22. Ritchie, M.E., et al., *limma powers differential expression analyses for RNA-sequencing*

and microarray studies. Nucleic Acids Res, 2015. **43**(7): p. e47.

23. Ciccia, A. and S.J. Elledge, *The DNA damage response: making it safe to play with*

knives. Mol Cell, 2010. **40**(2): p. 179-204.

24. Lundegaard, C., et al., *Major histocompatibility complex class I binding predictions as a*

tool in epitope discovery. Immunology, 2010. **130**(3): p. 309-18.

25. Yuan, J. and B.A. Yankner, *Apoptosis in the nervous system.* Nature, 2000. **407**(6805):

p. 802-809.

26. Hernández-Ortega, K., et al., *Altered Machinery of Protein Synthesis in Alzheimer's:*

From the Nucleolus to the Ribosome. Brain Pathol, 2016. **26**(5): p. 593-605.

27. Xie, L., et al., *Alzheimer's beta-amyloid peptides compete for insulin binding to the*

insulin receptor. J Neurosci, 2002. **22**(10): p. RC221.

28. Matioli, M. and R. Nitrini, *Mechanisms linking brain insulin resistance to Alzheimer's*

disease. Dement Neuropsychol, 2015. **9**(2): p. 96-102.

29. Ferraris, C.A.-O., et al., *Association between Sour Taste SNP KCNJ2-rs236514, Diet*

Quality and Mild Cognitive Impairment in an Elderly Cohort. LID - 10.3390/nu13030719

[doi] LID - 719. (2072-6643 (Electronic)).

30. Murphy, C., *Olfactory and other sensory impairments in Alzheimer disease.* Nature

Reviews Neurology, 2019. **15**(1): p. 11-24.

31. Zhang, H., H.J. Forman, and J. Choi, *Gamma-glutamyl transpeptidase in glutathione*

biosynthesis. Methods Enzymol, 2005. **401**: p. 468-83.

32. Fujii, J., et al. *Critical Roles of the Cysteine–Glutathione Axis in the Production of*

γ-Glutamyl Peptides in the Nervous System. International Journal of Molecular

Sciences, 2023. **24**, DOI: 10.3390/ijms24098044.

33.   Zakusilo, F.T., et al., *Matters of size: Roles of hyaluronan in CNS aging and disease.*

505   Ageing Research Reviews, 2021. **72**: p. 101485.

506   34.   Sun, Y., et al., *Role of the Extracellular Matrix in Alzheimer's Disease.* Frontiers in Aging

507   Neuroscience, 2021. **13**.

508   35.   Pimentel, J.M., J.Y. Zhou, and G.S. Wu, *The Role of TRAIL in Apoptosis and*

509   *Immunosurveillance in Cancer.* Cancers (Basel), 2023. **15**(10).

510   36.   Dubois, B., et al., *Preclinical Alzheimer's disease: Definition, natural history, and*

511   *diagnostic criteria.* Alzheimers Dement, 2016. **12**(3): p. 292-323.

512   37.   Jack, C.R., Jr., et al., *NIA-AA Research Framework: Toward a biological definition of*

513   *Alzheimer's disease.* Alzheimers Dement, 2018. **14**(4): p. 535-562.

514   38.   Jack, C.R., Jr., et al., *A/T/N: An unbiased descriptive classification scheme for Alzheimer*

515   *disease biomarkers.* Neurology, 2016. **87**(5): p. 539-47.

516   39.   Jack, C.R., Jr., et al., *Tracking pathophysiological processes in Alzheimer's disease: an*

517   *updated hypothetical model of dynamic biomarkers.* Lancet Neurol, 2013. **12**(2): p. 207-

518   16.

519   40.   Jack, C.R., Jr., et al., *Hypothetical model of dynamic biomarkers of the Alzheimer's*

520   *pathological cascade.* Lancet Neurol, 2010. **9**(1): p. 119-28.

521   41.   Knopman, D.S., et al., *The National Institute on Aging and the Alzheimer's Association*

522   *Research Framework for Alzheimer's disease: Perspectives from the Research*

523   *Roundtable.* Alzheimers Dement, 2018. **14**(4): p. 563-575.

524   42.   Wimo, A., et al., *The GERAS Study: a prospective observational study of costs and*

525   *resource use in community dwellers with Alzheimer's disease in three European*

526   *countries--study design and baseline findings.* J Alzheimers Dis, 2013. **36**(2): p. 385-99.

527   43.   Yang, S., *Bae DG, Kang HJ, Gwag BJ, Gho YS, Chae CB.* Co-accumulation of vascular

528   endothelial growth factor with beta-amyloid in the brain of patients with Alzheimer's

529   disease. Neurobiol Aging, 2004. **25**: p. 283-290.

44.   Patel, N.S., et al., *Alzheimer's β-amyloid peptide blocks vascular endothelial growth*

factor mediated signaling via direct interaction with VEGFR-2. Journal of

neurochemistry, 2010. **112**(1): p. 66-76.

45.   Martin, L., et al., *VEGF counteracts amyloid-β-induced synaptic dysfunction.* Cell

reports, 2021. **35**(6).

46.   Mahoney, E.R., et al., *Brain expression of the vascular endothelial growth factor gene*

family in cognitive aging and alzheimer's disease. Molecular psychiatry, 2021. **26**(3): p.

888-896.

47.   Hohman, T.J., et al., *The role of vascular endothelial growth factor in neurodegeneration*

and cognitive decline: exploring interactions with biomarkers of Alzheimer disease.

JAMA neurology, 2015. **72**(5): p. 520-529.

48.   Chiappelli, M., et al., *VEGF gene and phenotype relation with Alzheimer's disease and*

mild cognitive impairment. Rejuvenation research, 2006. **9**(4): p. 485-493.

49.   Huang, L., J. Jia, and R. Liu, *Decreased serum levels of the angiogenic factors VEGF*

and TGF-β1 in Alzheimer's disease and amnestic mild cognitive impairment.

Neuroscience letters, 2013. **550**: p. 60-63.

50.   Prodjohardjono, A., et al., *Higher level of acute serum VEGF and larger infarct volume*

are more frequently associated with post-stroke cognitive impairment. PloS one, 2020.

15(10): p. e0239370.

51.   Koester-Hegmann, C., et al., *High-altitude cognitive impairment is prevented by enriched*

environment including exercise via VEGF signaling. Frontiers in cellular neuroscience,

2019. **12**: p. 532.

52.   Tang, H., et al., *Expression level of vascular endothelial growth factor in hippocampus is*

associated with cognitive impairment in patients with Alzheimer's disease. Neurobiology

of aging, 2013. **34**(5): p. 1412-1415.

53.   Jais, A., et al., *Myeloid-cell-derived VEGF maintains brain glucose uptake and limits*

556   *cognitive impairment in obesity.* Cell, 2016. **165**(4): p. 882-895.

557   54.   Ng, T., et al., *Vascular endothelial growth factor inhibitors and cognitive impairment:*

558   *evidence and controversies.* Expert opinion on drug safety, 2014. **13**(1): p. 83-92.

559   55.   Mahoney, E.R., et al., *Brain expression of the vascular endothelial growth factor gene*

560   *family in cognitive aging and alzheimer's disease.* Mol Psychiatry, 2021. **26**(3): p. 888-

561   896.

562   56.   Rothhammer, V., et al., *Microglial control of astrocytes in response to microbial*

563   *metabolites.* Nature, 2018. **557**(7707): p. 724-728.

564   57.   Lampropoulou, A. and C. Ruhrberg, *Neuropilin regulation of angiogenesis.* Biochem Soc

565   Trans, 2014. **42**(6): p. 1623-8.