The resting state of the human T-cell receptor–CD3 complex

Ryan Q. Notti[1,2,*], Fei Yi[3,4], Søren Heissel[5], Martin W. Bush[1], Henrik Molina[5], Zaki Molvi[6],

Christopher A. Klebanoff[2,3,4], and Thomas Walz[1]

[1] Laboratory of Molecular Electron Microscopy, The Rockefeller University, New York, NY

[2] Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, NY

[3] Human Oncology Pathogenesis Program, Memorial Sloan Kettering Cancer Center, New

York, NY

[4] Parker Institute for Cancer Immunotherapy, Memorial Sloan Kettering Cancer Center, New

York, NY

[5] Proteomics Resource Center, The Rockefeller University, New York, NY

[6] Immunology Program, Memorial Sloan Kettering Cancer Center, New York, NY

* To whom correspondence should be addressed: rnotti@rockefeller.edu

Summary

The T-cell receptor (TCR) is central to the ligand-dependent activation of T lymphocytes and as

such orchestrates both adaptive and pathologic immune processes[1]. However, major questions

remain regarding the structure and function of the human TCR[2–4]. Here, we present cryogenic

electron microscopy structures for the unliganded human TCR–CD3 complex in a native-like

lipid bilayer, revealing two related conformations that are distinct from its structure in detergent.

These new "closed and compacted" conformations afford insights into the interactions between

the TCR–CD3 and the membrane, including conserved surface patches that make extensive

outer leaflet contact, and suggest novel conformational regulation by glycans. We show that the

closed/compacted conformations, not the extended one previously reported in detergent[5–8],

represent the unliganded resting state for the TCR–CD3 *in vivo*, underscoring the importance of

structural interrogation of membrane proteins in native-like environments. We use conformation-

locking disulfide mutants to show that ectodomain opening is necessary for maximal ligand-

dependent TCR–CD3 activation, demonstrating that TCR-intrinsic conformational change is

necessary for full TCR–CD3 activation and opening numerous avenues for immunoreceptor

engineering.

Main

T cells are central to a wide array of both adaptive and pathologic immune responses. The T-

cell receptor (TCR), in association with members of the CD3 protein family, allows T cells to

recognize and respond to antigenic peptides presented on human leukocyte antigen (HLA)

complexes of other cells. While the identities of the TCR–CD3 complex constituents and its

downstream signaling elements have been known for some time, questions remain regarding its

activation mechanism.

The TCR–CD3 is a hetero-octameric membrane protein complex[5–8]. The $\alpha\beta$-type TCR is

comprised in part by two variable chains, TCRα and TCRβ, which directly interface with the

HLA-antigen complex. Antigen specificity is generated through somatic recombination of the

variable segments in the TCRα and TCRβ genes. The remainder of the core complex is

comprised of six invariant CD3 family chains: a CD3γ-CD3ϵ heterodimer, a CD3δ-CD3ϵ

heterodimer, and a CD3ζ homodimer. Herein, we differentiate the CD3ϵ protomers by the CD3

component with which each pairs (CD3ϵ_D pairs with CD3δ, and CD3ϵ_G pairs with CD3γ) and the

CD3ζ protomers as CD3ζ and CD3ζ'.

While the TCRα and TCRβ chains lack intracellular signaling domains, the CD3 proteins

possess cytoplasmic immunoreceptor tyrosine activation motifs (ITAMs) that become

phosphorylated upon TCR-HLA interaction and then bind and activate downstream signaling

kinases. Dynamic interactions between ITAMs and the membrane regulate signaling[9,10], but the

mechanism by which ligand binding to the extracellular TCRα-TCRβ domains results in

phosphorylation of cytoplasmic CD3 ITAMs remains a matter of debate. Specifically, it is unclear

whether ligand binding and activation involves conformational change in the TCR–CD3

complex. While reconstitution of TCR–CD3 signaling *in vivo* with minimal constructs has argued

against transmembrane conformational change[11], biochemical evidence in support of

conformational change[12–15] and specifically for reversible structural extension of the TCR under

mechanical force suggests the opposite[16], although the structural details of such conformational

transitions remain to be determined.

Structures of the hetero-octameric $\alpha\beta$-type TCR–CD3 have been determined by

cryogenic electron microscopy (cryo-EM)[5–8]. These studies revealed an antenna-like structure,

with the TCRα and TCRβ variable domains projecting over a ring formed by the CD3

ectodomains and the TCRα and TCRβ constant domains, with numerous interactions between

their respective transmembrane (TM) helices (Extended Data Figure 1a). Addition of HLA-

antigen to this complex did not meaningfully alter the conformation of the TCR–CD3[7,8], arguing

against models of intrinsic conformational change in the ligand-dependent TCR–CD3 activation

process (Extended Data Figure 1a,b).

However, the TCR–CD3 complexes used in these studies were solubilized in detergent.

Biochemical data have previously shown important roles for the lipid bilayer in the regulation of

TCR–CD3 signaling specifically[6,17,18], raising questions as to the physiologic relevance of the

determined structures. Membrane protein structure determination is generally preferable in a

native-like membrane environment that increases the likelihood that important protein–lipid

interactions are captured in the obtained molecular models[19].

Here, we report cryo-EM structures of a human TCR–CD3 stabilized both in detergent

micelles and in a native-like lipid environment. The structures reveal novel conformations for the

TCR–CD3 in a lipid bilayer, distinct from those previously reported, and cross-linking data

support the notion that these conformations represent the resting state of an unstimulated TCR–

CD3 on T cells. We hypothesize that the previously published structures of the TCR–CD3 in

detergent approximate a ligand-activated conformation and show that exit from the resting state

conformation is required for maximal TCR–CD3 signaling.

Results

Structure of the NY-ESO-1-specific 1G4 TCR–CD3 in detergent micelles. In this study, we

investigated the structure of the 1G4 (LY-variant) TCR–CD3. We chose the 1G4 TCR, which

binds to an HLA-A*02:01-restricted 9 amino acid epitope derived from the NY-ESO-1 cancer-

testis antigen, because it is both a model immunoreceptor and being developed as an anti-

cancer therapeutic[20]. To confirm that 1G4 has a similar structure to the other TCR–CD3

complexes analyzed in detergent, we purified this complex in glyco-diosgenin (GDN) and

determined its structure by cryo-EM to a nominal resolution of 3.3 Å (Figure 1a, Extended Data

Figure 1c-f, Extended Data Table 1, Supplementary Methods Figures 1, 2, 5-8). Herein, this

model and the conformation it represents will be abbreviated as "GDN." Overall, GDN was very

similar to the structures of three other TCR–CD3 constructs, including those with and without

HLA bound (Extended Data Figure 1g). This is not surprising, given the nearly identical CD3

and TCR constant region sequences shared by these receptors (Extended Data Figure 1h).

Structure of the TCR–CD3 in a native-like lipid environment. To determine the structure of the

1G4 TCR–CD3 in a native-like lipid environment, we reconstituted the TCR–CD3 complex into

proteolipid nanodiscs loaded with mammalian polar lipids and cholesteryl hemisuccinate

(Supplementary Methods Figure 1). Computational processing of cryo-EM data yielded two

distinct maps determined to nominal resolutions of 3.3 to 3.1 Å (for locally refined maps used for

model building and refinement; Supplementary Methods Figures 3-5, Extended Data Figure 2).

We denote these as nanodisc (ND)-I and ND-II. These maps were clearly different from GDN

(Figure 1) and the previously published TCR structures. Attempts to bias the particle search

towards the published conformations with projections of the structure in detergent or that same

structure docked into a lipid nanodisc model did not yield any maps approximating the

previously published conformation (Supplementary Methods Figure 3).

Despite equal or higher nominal overall resolutions than GDN, ND-I and ND-II were not

as high quality (compare local resolution estimations in Extended Data Figure 2d,h with

Extended Data Figure 1f). This was likely due to a high degree of conformational variability, but

also due in part to map anisotropy (compare Extended Data Figure 2c,g to Extended Data

Figure 1e). Efforts to improve the orientation distribution of the particles with graphene oxide

grids and fluorinated detergents did not improve reconstruction quality (data not shown).

Nonetheless, these maps were of sufficient quality to build models into the density, with much of

the density having estimated local resolution of better than 4 Å.

Molecular models were built for ND-I and ND-II by docking the GDN structure into the

maps and refining them by standard approaches (Extended Data Table 1, Supplementary

Methods Figures 6-8; see Supplementary Methods for details). These models reveal two distinct

but related conformations of the TCR–CD3 (Figure 1a). The ND structures show two large

conformational changes relative to GDN: a "closure" of the TCR ectodomains against the CD3

ectodomains (Figure 1a,b) and a "compaction" of the juxtamembrane (JM) linker peptides

(Figure 1a,c,d). Collectively, these changes shorten the TCR projection into the extracellular

milieu by approximately 35 Å. Closure of the TCR ectodomains (Figure 1b) results in part from

rotation of the variable domains towards their respective constant domains about the

interdomain hinge region (Extended Data Figure 3a,b). Relative to the "extended" JM linker

conformations seen in GDN, the CD3 JM linkers are compacted, with closer approximation of

the CD3 ectodomains to the amino-terminal ends of their TM helices (Figure 1c,d).

The conformational changes in the TCR ectodomains are accompanied by rotations of

TM helices. While the in-plane arrangement of the TM helices is largely similar when viewed

along the membrane normal (Figure 1e), examination from orthogonal axes reveals differences

in TM helix conformation. Compaction of the CD3 JM linkers requires opening of the angle

between the TM helices of the CD3$\varepsilon\delta$ and CD3$\varepsilon\gamma$ pairs by 23° (from 31° to 54°) and 28° (from

39° to 67°), respectively, when comparing ND-II with GDN (Figure 1c,d). For ND-I, the TM

helices of the CD3$\varepsilon\delta$ are bent, with the first helical turn displaying a wide splay (76°, Extended

Data Figure 3c), but the overall angle between the modeled helices (46°) is intermediate to ND-

II and GDN. The CD3ε_G TM helix was not modeled, as its register could not be confirmed, and

thus the CD3$\varepsilon\gamma$ TM angle was not measured. Additionally, there is splaying of the TCRβ and

distal CD3ζ TM helices away from the membrane normal (Figure 1f). In GDN, the TCRα TM

helix extends outward past the amino-terminal ends of the CD3 TM helices, where it is

ensconced by the extended CD3 JM linkers. Compaction of the JM linkers in nanodisc

necessitates reorganization of the first turn of the TCRα TM helix, which is unstructured and

displaced towards the lipid bilayer (Figure 1f). Taken together, these cryo-EM maps reveal

marked conformational differences between the TCR–CD3 in detergent micelles and a lipid

bilayer.

Additionally, the transmembrane helices of ND-I and ND-II appear shorter than in GDN.

Specifically, these structures lack clear density for the cluster of basic and aromatic amino acids

visualized in GDN that extend the TM helices (Extended Data Figure 3d). In a membrane, these

extensions would likely be interfacial or extend into the cytoplasm[21]. Although there is additional

density beyond the modeled carboxy-termini of some of these shortened helices in ND-I and

ND-II, it is not of sufficient quality to model with confidence (Extended Data Figure 3e,f). This

suggests the TCR–CD3 TM helices are less rigid in the inner leaflet of a lipid bilayer than in a

micelle.

Lipid interactions with the TCR–CD3. The TCR–CD3 is known to interface directly with

membrane cholesterol[22], and the binding sites have been visualized in previously published

structures[6]. In GDN, ND-I, and ND-II, density corresponding to two cholesterols in the previously

published structures was again apparent (Extended Data Figure 3g). The two cholesterol

molecules in ND-I and ND-II are oriented tail-to-tail in a groove between CD3ζ', TCRα, TCRβ,

and CD3γ, as in GDN (Figure 2a). However, the compaction-associated changes in the TCR–

CD3 TM helices results in a sliding of the cholesterol molecules towards each other, with

deflection of their branched aliphatic tails to prevent clashing (Figure 2a, insets). We refer to the

cholesterol molecules as $Chol_A$ and $Chol_B$, corresponding to the TCR–CD3 TM helix with which

each is approximated. The deflected tail of $Chol_B$ is interposed between adjacent TM helices,

accommodated by their increased splay in the compact conformation (Figure 2b, inset). The

cholesterol-binding groove is more open to the membrane bilayer in nanodiscs, creating a

possible path for cholesterol loading into the assembled TCR–CD3 and/or exchange with the

membrane environment (asterisks in Figure 2b, bottom left).

Imaging of the TCR–CD3 in a lipid bilayer revealed additional sites of protein–lipid

interaction. Unmodeled tubular densities consistent with phospholipid acyl chains are present at

several sites adjacent to the TCR–CD3 TM helices (yellow densities, Extended Data Figure 3h-

k). Structured lipid is seen parallel to the tail-to-tail cholesterol dimer, filling the aforementioned

channel between the cholesterol-binding groove and the bulk membrane lipids (Figure 2b,

bottom left; Extended Data Figure 3g). Splay of the CD3ζ TM helix away from that of CD3ε_D

enlarges a cavity, which is filled by several presumptive lipid acyl chains in the nanodisc

(Extended Data Figure 3j). Additional probable sites of structured lipid include a shroud of

density over the CD3ζ TM helices (Extended Data Figure 3h) and in the cleft over the TCR$\alpha\beta$

TM helices between CD3δ and CD3ε_G (Extended Data Figure 3i).

The ND-I and ND-II maps show clear density for the proteolipid nanodisc that enshrouds

not only the TM helices, but also the membrane proximal faces of the TCR–CD3 ectodomains

(Figure 3a, left). The lipid contacts with the CD3 ectodomains are apparent even in maps

contoured at a higher threshold, consistent with high occupancy (Figure 3a, right). While the

CD3 ectodomain surfaces are generally poorly evolutionarily conserved in comparison to the

TM helices, patches on the membrane-proximal surfaces of the CD3 ectodomains do show high

sequence conservation (Figure 3a,c). These highly conserved ectodomain surface patches

(black triangles in Figure 3b-e) interface with the extracellular membrane leaflet and are neutral

to electropositive (Figure 3e), facilitating phospholipid head group interactions. Of note, we

observe a similar conserved electropositive patch at the interface of CD79$\alpha\beta$[23], the analogous

ITAM-bearing chains in the B-cell receptor, raising the prospect that this interface may also

favor membrane interactions and a different conformation in the lipid bilayer from that reported

in detergent micelles (Extended Data Figure 4).

Probing TCR conformation in situ. Which conformation of the TCR–CD3 is the physiologic

resting state on the cell membrane: the open/extended conformation seen in detergent micelles,

or the closed/compacted conformations seen in the lipid bilayer of the nanodisc? Given that the

lipid bilayer of a nanodisc more closely approximates the physicochemical milieu of the plasma

membrane, we hypothesized that the TCR–CD3 is closed and compacted in the absence of

activating ligand *in vivo.* To test this notion, we engineered a series of disulfide bonds that would

only be geometrically favorable in ND-I and/or ND-II but not GDN (or vice versa), and we read

out their presence or absence by one of the following approaches: (1) non-reducing sodium

dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and Western blot of whole cell

lysates for interchain disulfides, or (2) mass spectrometry on affinity-purified protein for

intrachain disulfides. Analysis of disulfide bonds in the Protein Databank have shown that

disulfide bonds almost never have a Cys-Cys Cα-Cα distance greater than 7 Å, with most being

between 5 and 6.5 Å[24]. Accordingly, we sought residue pairs in the TCR–CD3 ectodomains with

Cα-Cα distances below 7 Å in one conformation and above 7 Å in the other as candidates for

disulfide engineering.

The interface between CD3δ and TCRα contains several pairs of residues with geometry

favorable for engineered disulfide formation in ND-I but not GDN (Figure 4a). As TCRα and

TCRβ are linked by an interchain disulfide natively (Extended Data Figure 5a), formation of a

TCRα–CD3δ disulfide would result in an upward gel shift of the TCR$\alpha\beta$ heterodimer in

nonreducing SDS-PAGE. For all three tested engineered disulfide pairs meeting the 7 Å Cα-Cα

distance constraint, a gel shift consistent with TCRα–CD3δ disulfide formation was observed

(Figure 4b, asterisks; all full-length gels in Supplementary Methods Figure 9). Confirming that

this mass shift was due to disulfide formation, no gel shift was apparent under reducing

conditions (Supplementary Methods Figure 9b). Moreover, the mass shift was due to the

formation of the engineered disulfide and not an off-target bond, as adduct formation was only

seen when both residues in a given target pair were mutated to cysteines (Figure 4b). As an

internal negative control, a permutation of cysteine mutants at this interface not meeting

geometric criteria for disulfide formation in ND-I (TCRα S78C – CD3δ N38C, Figure 4a, right,

orange labels) was produced. It did not form a disulfide *in vivo* (Figure 4b, orange) despite S78C

and N38C successfully pairing with cognate cysteine mutants that did meet the inter-Cα

threshold. This control shows the structural precision and specificity of the engineered disulfide

bond approach. These experiments were performed using whole-cell lysates, but we obtained

similar results with NiNTA-affinity-purified protein (Extended Data Figure 5b), and we were able

to reconstitute the two G79C-crosslinked mutants into nanodiscs (Extended Data Figure 5c,d),

suggesting proper folding and assembly of the holo TCR–CD3.

The region around the TCRα and TCRβ interdomain hinges is also notably different in

the closed/compacted conformation compared to the open/extended one, with the variable

domains coming into closer apposition with their respective constant domains, allowing the

engineering of conformation-specific disulfide pairs in this region (Figure 4c,d). We probed

formation of these intrachain disulfides by mass spectrometry. Proteolytic digestion of the TCR–

CD3 with trypsin and chymotrypsin yielded distinct peptides, several of which were linked by

disulfide-bonds. Electron-transfer/higher-energy collision dissociation (EThcD) fragmentation

was used to fragment the disulfide-bonded peptides providing *in vacuo* reduction to the linear

peptide constituents along with peptide backbone fragmentation, thereby allowing unambiguous

identification of the engineered disulfides[25]. All native TCR–CD3 disulfides spanning two

proteolytic fragments (Extended Data Figure 5a) were observed in the wild-type 1G4 TCR–CD3

by this approach (Extended Data Table 2). For both ND conformation-compatible engineered

disulfides tested (Figure 4c,d), we observed formation of the engineered disulfide (Figure 4e,f

and Extended Data Table 2). In contrast, an engineered disulfide that would only be present in

the open/extended conformation (Extended Data Figure 5e) did not form *in vivo* (Extended Data

Table 2). Taken together, these data support the conclusion that a TCR–CD3 in the unliganded,

resting state adopts the closed/compact conformation *in vivo*.

Conformational variability of the TCR–CD3 in a lipid bilayer. To allow direct comparison of the

two distinct, but related, closed/compacted structures (ND-I and ND-II), they were aligned by

their TCRα TM helices. TCRα was chosen because it is relatively perpendicular to the

membrane plane (alongside CD3ζ') and thus might be least affected by any differences in TM

helix splay. Indeed, this alignment strategy brought the nanodisc density of their maps into

alignment as well (Extended Data Figure 6a), giving both models the same orientation relative

to the membrane bilayer. In contrast to the strong overall alignment of the TCR–CD3 TM helices

by this approach (Extended Data Figure 6b,c), the ectodomains of ND-I and ND-II has marked

conformational differences, as seen in the aligned cryo-EM densities (Extended Data Figure

6a,b) and the resulting protein models (Figure 5a). Comparison of the structures reveals the

motions allowing exchange between these conformations (Figure 5b,c, Supplementary Movie

1). ND-I is the most closed of the three structures in this study, and transition to ND-II requires

several motions: the depression of the CD3ε_D (Figure 5b) and CD3γ ectodomains (Figure 5c)

away from the TCR variable domains and towards the bilayer outer leaflet; rotation of the of the

TCRα variable domain about the interdomain hinge (Figure 5b); and retraction of the TCRβ

variable domain (Figure 5c). This is accompanied by sliding of CD3ζ along its helical axis

towards the ectodomains (Figure 5b). While the TCRα and CD3ζ' TM helices remain relatively

fixed in this frame of reference, the TCRβ and CD3γ TM helices shift away from the cholesterol-

binding groove (Figure 5d, Extended Data Figure 6c). To accommodate this movement, there is

rotation of Chol$_A$ and elevation of Chol$_B$ (Figure 5d), showing cholesterol to be an integral and

dynamic component of the TCR–CD3 structure.

The GDN map displays densities for many of the known asparagine (N)-linked glycans in

the TCR–CD3 (Extended Data Figure 6d). There were fewer densities modellable as N-linked

glycans in the ND-I and ND-II maps (Extended Data Figure 6d). However, all three maps show

an array of densities lateral to the CD3δ and TCRα ectodomains, corresponding to sites of

known N-linked glycosylation. In ND-II, the TCRα (linkage unclear) and CD3δ N74-linked

glycans are visible, emerging radially from the TCR–CD3, as in GDN, but are not well structured

(Figure 5e). By comparison, ND-I reveals clear reorganization of the glycans (Figure 5e), with

increased stacking of adjacent glycans brought into close proximity by tight ectodomain closure.

This arraying of N-linked glycans is reminiscent of that seen on HIV Env, where interacting

adjacent glycans form a structured arcade of glycan cores (Extended Data Figure 6e). Two

densities in the array are attributable to CD3δ N74- and TCRα N82-linked glycans, with a third

density emerging between them without definitive attribution (purple question mark in Figure 5e,

right inset). In GDN, as in other structures of TCR–CD3 in detergent, the CD3δ N38-linked

glycan is oriented inward towards the cleft between the TCR and CD3 ectodomains (Figure 5f,

top). In ND-II, CD3δ N38 is unstructured (Figure 5f, middle), but is grossly positioned at the

entrance to the cleft, and the linked glycan could conceivably orient either into or away from the

cleft. In ND-I, CD3δ N38 is structured (Figure 5f, bottom), but its glycan cannot be confidently

modeled. However, density protrudes from N38 into the open cleft (Figure 5f, asterisk) and

additional unmodeled density is seen within the cleft, consistent with the glycan remaining within

the cleft in the ND-I conformation.

The structure of the TCR variable domains in GDN is grossly unchanged from that seen

in previous crystal structures of the 1G4 TCR bound to HLA-β2M-antigen[26] (herein, TCR-HLA;

Extended Data Figure 6g). Despite the unstructuring of their amino-termini and some surface

loops in the ND-I and ND-II, as well as the domain-scale movements described above, there

was little change in the orientations of the TCR variable domains relative to one another when

compared to GDN and the TCR-HLA crystal structure (Extended Data Figure 6g). However,

differences in ectodomain closure seen between ND-I and ND-II could have consequences for

the accessibility of the TCR variable domains to HLA-antigen complexes. To assess the impact

of ectodomain cleft opening and closure on TCR–HLA interactions, we modeled the binding

geometry of HLA onto GDN, ND-I, and ND-II by alignment of the TCR-HLA crystal structure to

the TCR variable domains in our models (Figure 5g). There appears to be adequate clearance

for the TCR variable domains to interact with HLA in GDN and ND-II. For ND-I, the modeled fit

is tighter, with the CD3ε_D ectodomain predicted to be in close contact with HLA. However, on

assessment of space-filling model cross-sections, no overt steric clashes are evident (Figure

5h). Because a portion of a potentially clashing surface loop in the CD3ε_D ectodomain is not

modeled in ND-I, the TCR-aligned HLA model was docked into the ND-I cryo-EM density

(Figure 5i), but no unmodeled density appears to clash with HLA. The CD3ε_D ectodomain from

GDN is completely modeled and aligns closely with that from ND-I (Extended Data Figure 6f,

inset). As an additional sensitivity analysis, the GDN CD3ε_D ectodomain was modeled into ND-I

and the HLA interaction again assessed. While there is a small overlap in their space-filling

models (Extended Data Figure 6f), no large clashes are observed. Thus, while it remains

unclear what quantitative effect ectodomain cleft closure might have on HLA affinity, particularly

in the physiologic context of a glycosylated HLA molecule, it is at least plausible that both ND-I

and ND-II may be receptive to HLA binding.

A model for TCR–CD3 conformational change in receptor activation. Taking the above

structural, biochemical, and modeling data together and in context with the literature, we

developed a model for ligand-dependent activation of the TCR–CD3. Our data support a model

in which the unliganded "resting" TCR–CD3 explores a thermally accessible spectrum of

closed/compacted conformations from the most closed ND-I to the more open ND-II. Previous

data have shown HLA monomer binding to the TCR is insufficient to drive T-cell activation, but

that subsequent force application induces activation[27]. Force-dependent activation was

accompanied by an increase in TCR length normal to the membrane[16], raising the possibility

that the open/extended structure of the TCR reported in detergent micelles is actually an

approximation of a physiologically active state. We do note that other data support the

possibility of TCR–CD3 conformational change upon ligand binding, even without external force

application, suggesting an allosteric mechanism[12,28], which would also be consistent this model.

Moreover, TCR–CD3 opening and/or extension may be necessary for coreceptor interaction

with HLA bound to the TCR–CD3. Modeling the HLA class II-CD4 complex onto GDN, ND-I, and

ND-II reveals an unfavorable geometry for CD4 interaction with a class II HLA when bound to a

TCR–CD3 in the ND-I or ND-II conformation because of their proximity to the membrane

(Extended Data Figure 6h), at least for the CD4 conformation reported in the CD4-HLA-TCR

ectodomains ternary complex crystal structure[29]. In contrast, CD4-HLA can be accommodated

in the GDN conformation (Extended Data Figure 6h). The full ectodomain structure of the CD8

coreceptor has not been described, precluding a similar analysis applicable to TCRs

recognizing HLA class I-restricted epitopes, like the one used in this study.

We hypothesize that binding of HLA-antigen to the TCR–CD3 results in a structural

transition from the closed/compacted resting conformation to an open/extended active

conformation (Figure 6a). We favor this transition occurring via the ND-II conformation, which is

more open than ND-I and thus closer to the active state along this trajectory. Morphing the

closed/compacted conformation to the open/extended one (Supplementary Movie 2) highlights

the straightening of the TM helices towards the membrane normal (that is, decreased splay,

Figure 6b). These changes in TM conformation add to the plausibility of the model, as they

provide a means to transmit conformational changes in the ectodomains to the cytoplasmic

ITAMs.

Thus, transition from the closed/compacted to the open/extended conformation would

seem to require further ectodomain opening from the ND-II state (Figure 6a). We hypothesized

that the intra-TCRα disulfide-forming double mutant S104C–V182C (characterized above,

Figure 4c) would prevent ectodomain opening to the more open active conformation, and thus

impair ligand-dependent TCR–CD3 activation. To investigate the impact of restrained TCR–CD3

opening on ligand-dependent activation of T cells, we introduced wild-type and TCRα S104C–

V182C mutant TCR constructs into TCR$\alpha\beta$-negative Jurkat T cells bearing an NFAT-responsive

fluorescent reporter. TCR–CD3 assembly and surface localization (as assessed by anti-CD3ϵ

antibody binding) and HLA-binding competence were confirmed by flow cytometry for the

mutant and were comparable to cells transduced with wild-type 1G4 (Figure 6c). In the absence

of HLA-antigen tetramers, there was no detectable reporter fluorescence in these cells, but

upon exposure to HLA-antigen tetramers, cells expressing wild-type TCR–CD3 became

activated, with reporter fluorescence peaking 16 h after HLA-tetramer addition (Figure 6d). By

contrast, peak reporter fluorescence was significantly decreased in the TCRα S104C–V182C

mutant, with peak activation attenuated by 26% ($P < 0.05$, Figure 6d, source data in

Supplementary File 1) relative to wild-type on exposure to 8 μg/mL of tetramers. The TCRα

S104C–V182C mutant and wild-type controls were equally responsive to phorbol 12-myristate

13-acetate (PMA) and ionomycin, which bypass TCR–CD3 signaling to activate T cells.

Discussion

Here, we report structures of a human TCR–CD3 both in a lipid bilayer and in detergent

micelles. We find that the TCR–CD3 hetero-octamer assumes a spectrum of more closed and

compacted conformations when embedded in a lipid bilayer relative to that seen in detergent

micelles, that the closed/compacted conformation is indeed the native unliganded state of the

TCR–CD3, and that exit from this conformation by ectodomain opening is necessary for

maximal ligand-dependent activation of the TCR–CD3. While this latter finding does not rule out

roles for TCR–CD3-extrinsic phenomena (e.g., higher-order clustering, steric exclusion of

inhibitory phosphatases) in the activation of the TCR–CD3 or regulation thereof, it argues that

intrinsic conformational change in the TCR–CD3 is necessary for early TCR–CD3 activation and

argues against previous models that have excluded this requirement[11,30].

Broadly, these findings reinforce the importance of structural interrogation of membrane

proteins in lipid environments that closely resemble their physiologic environment. Comparison

of TCR–CD3s in detergent micelles in the absence and presence of activating ligand had

revealed no change in TCR–CD3 structure[7,8], consistent with models excluding TCR–CD3

conformational change in the activation process. Although these studies were technically well

performed, our results now reveal that they did not identify the true resting state of the TCR–

CD3, calling into question the conclusion that there is no intrinsic conformational change in the

TCR–CD3 during ligand-dependent activation.

Additionally, the structures in detergent did not capture the full scope of TCR–CD3

interaction with the membrane. Not only are membrane lipids closely interwoven with the TM

helices, but large conserved surface patches on the ectodomains mediate lipid–protein

interactions, likely partially counteracting the thermodynamic costs of JM linker compaction and

"spring loading" the TCR with potential energy for conformational change during activation. In

this model, ligand binding and/or force application provides the activation energy to release the

membrane-binding surfaces of the CD3 ectodomains and allow JM extension. This would be

consistent with existing models of TCR–CD3 response to force, which invoke conformational

changes with free energy barriers rendering them inaccessible to thermal motion alone[31].

Detergent extraction would overcome this thermodynamic barrier by removing the lipid bilayer

all together. During reconstitution into nanodiscs, we first extracted the TCR–CD3 complex in

detergent, opening and extending the receptor, which suggests that the addition of a lipid bilayer

is sufficient to reset the TCR–CD3 to its resting conformation without the need for extrinsic

chaperones. Lastly, we note that a similar conserved surface patch is present on the CD79

heterodimer in the B-cell receptor[23], raising the possibility that other hetero-multimeric ITAM-

containing immunoreceptors exploit protein surface interactions with the lipid bilayer to constrain

conformation.

As with the TCR–CD3 structures in detergent, the CD3 ITAMs are not visible in our

structures. The ITAMs are membrane-associated in the unliganded state[9,10], but are likely to be

radially disordered along the intracellular membrane face. However, the TM helix movements

seen between our closed/compacted and the open/extended conformations suggest a means

for communication of extracellular conformational changes in the TCR–CD3 complex to the

ITAMs. Within the cytoplasmic domains of CD3 proteins, the basic residues proximal to the CD3

TM helix are critical for ITAM association with the membrane[9,10]. Upon transition from the

receptive to the hypothesized active conformation, the decreased splay in the CD3 TM helices

could alter basic patch interaction with the membrane and favor ITAM release.

Analysis of the basal conformational heterogeneity of the TCR–CD3 suggests novel

means of TCR–CD3 regulation. The TCRα and CD3δ glycan ordering observed in the ND-I

conformation is notably different from the glycan arrangement in the more open ND-II

conformation or the TCR–CD3 in detergent. While molecular modeling of the branched glycan

sugars was not possible with sufficient confidence, the stacked arrangement of glycan core

densities raises the possibility of extensive hydrogen bonding between the glycan branches

distally and perhaps also with TCR–CD3 surface residues. While glycan–glycan and glycan–

protein interactions are well described *in trans*, there are fewer examples of structured glycan–

glycan interactions within a given protein or stable complex. The human immunodeficiency

virus-1 Env protein is an example of the latter[32,33] and the arrangement of the radial glycan array

in ND-I is reminiscent of that seen in crystallographic studies of Env. However, the stacking of

glycans observed in the TCR–CD3 is unique in being protein conformation-dependent. Breaking

an extensive glycan hydrogen-bonding network to transition out of ND-I would be energetically

costly, and thus modification of TCR–CD3 glycosylation could serve as a means of

posttranslational regulation of the TCR–CD3 activation threshold by increasing or decreasing

the free energy barrier to ND-I exit.

Previous studies have explored the role of cholesterol in the regulation of TCR–CD3

activation[6,18,22]. The substantive movements of the two cholesterol molecules seen between

TCR–CD3 conformations raise the possibility that changes in the sterol composition of the T-cell

membrane (either global or local) could alter TCR–CD3 activity even prior to ligand binding by

shaping the basal TCR–CD3 conformational landscape. Moreover, our structures reveal

channels for cholesterol loading or exchange with the lipid bilayer not previously observed,

making it plausible that already assembled TCR–CD3 could respond to changes in membrane

sterol content. Non-sterol lipid species are also known regulators of T-cell signaling[17], including

phosphoinositol lipids[34,35] and oxidized lipids in the tumor microenvironment[36]. This work

delineates multiple sites of structured interaction between the TCR–CD3 and membrane lipids

beyond the sterol-binding site, which may allow specific lipids to modulate TCR–CD3

conformation and activity.

The extensive contacts between TCR variable domain residues and those of the TCR

constant domains or CD3 ectodomains in the closed/compacted conformations raises questions

about how variation at these sites might confer specific baseline sensitivities to given TCR

alleles. Whether use of certain V or J alleles changes the occupancy of different segments of

the TCR–CD3 conformational landscape will need to be determined and has broad implications

for vaccine design and immunotherapy engineering. Likewise, whether individually rare germline

coding variants in the TCR constant domain *loci* or CD3 proteins affects an individual's basal T-

cell reactivity is unclear outside of known deleterious loss-of-function alleles[37]. Such information

might predict individual propensity for autoimmunity or responsiveness to immune-oncology

agents and will provide insights for engineering TCRs with favorable activity profiles[38].

Methods

Full methods are detailed in the Supplementary Methods.

Data Availability Statement

Cryo-EM maps and atomic coordinates for GDN, ND-I, and ND-II have been deposited in the

Protein Data Bank and Electron Microscopy Data Bank with the following accession numbers:

GDN, PDB 9BBC, EMD-44417; ND-I, PDB 8TW4, EMD-41658; ND-II, PDB 8TW6, EMD-41660.

Acknowledgements

The authors thank the members of the Walz, T. de Lange (RU), and R. K. Hite (MSKCC) labs

and Drs. E. L. Reinherz (HMS, DFCI) and W. D. Tap (MSKCC) for insightful conversations. We

thank M. Ebrahim, J. Sotiris, and H. Ng at the Evelyn Gruss Lipper Cryo-EM Resource Center of

The Rockefeller University for assistance with cryo-EM data collection. We thank Pujita Das for

excellent technical assistance. R.Q.N. was supported by National Institutes of Health (NIH)

grants T32CA9207, KL2TR001865, and UL1TR001866; the Black Family Metastasis Center at

Rockefeller University; the Shapiro-Silverberg Fund for the Advancement of Translational

Research; and the Mark Foundation for Cancer Research (R.Q.N. and T.W.). C.A.K. was

supported, in part, by NIH grants R37 CA259177, R01 CA269733, R01 CA286507, P30

CA008748, and P50 CA217694; Cycle for Survival and the Shteinbuk and Mead Family; the

Parker Institute for Cancer Immunotherapy; and The Sarcoma Center at MSKCC. Mass

spectrometry data was generated by the Proteomics Resource Center at The Rockefeller

University (RRID:SCR_017797) using instrumentation funded by the Sohn Conferences

Foundation and the Leona M. and Harry B. Helmsley Charitable Trust.

Author Contributions

R.Q.N conceived of the project, designed and performed experiments, analyzed data, built

models, and wrote and revised the manuscript; F.Y. designed and performed experiments and

analyzed data; S.H. designed and performed experiments and analyzed data; M.B. performed

experiments and analyzed data; H.M. designed and performed experiments and analyzed data;

C.A.K. designed experiments, analyzed data, and revised the manuscript; T.W. designed

experiments, analyzed data, and revised the manuscript.

Competing Interests

C.A.K. is an inventor on patents related to TCR discovery and public neoantigen-specific TCRs

and is recipient of licensing revenue shared according to MSKCC institutional policies. C.A.K.

has consulted for or is on the scientific advisory boards for Achilles Therapeutics, Affini-T

Therapeutics, Aleta BioTherapeutics, Bellicum Pharmaceuticals, Bristol Myers Squibb,

Catamaran Bio, Cell Design Labs, Decheng Capital, G1 Therapeutics, Klus Pharma, Obsidian

Therapeutics, PACT Pharma, Roche/Genentech, Royalty Pharma, and T-knife. C.A.K. is a

scientific co-founder and equity holder in Affini-T Therapeutics.

References

1. Janeway, C. A. *et al. Immunobiology*. (Garland Science, 2001).

2. Reinherz, E. L. The structure of a T-cell mechanosensor. *Nature* **573**, 502–504 (2019).

3. Courtney, A. H., Lo, W.-L. & Weiss, A. TCR Signaling: Mechanisms of Initiation and

Propagation. *Trends Biochem. Sci.* **43**, 108–123 (2018).

4. Mariuzza, R. A., Agnihotri, P. & Orban, J. The structural basis of T-cell receptor (TCR)

activation: An enduring enigma. *J. Biol. Chem.* **295**, 914–925 (2020).

5. Dong, D. *et al.* Structural basis of assembly of the human T cell receptor–CD3 complex.

Nature **573**, 546–552 (2019).

6. Chen, Y. *et al.* Cholesterol inhibits TCR signaling by directly restricting TCR–CD3 core tunnel

motility. *Mol. Cell* **82**, 1278-1287.e5 (2022).

7. Sušac, L. *et al.* Structure of a fully assembled tumor-specific T cell receptor ligated by pMHC.

Cell **185**, 3201-3213.e19 (2022).

8. Saotome, K. *et al.* Structural analysis of cancer-relevant TCR–CD3 and peptide-MHC

complexes by cryoEM. *Nat. Commun.* **14**, 2401 (2023).

9. Xu, C. *et al.* Regulation of T Cell Receptor Activation by Dynamic Membrane Binding of the

CD3ε Cytoplasmic Tyrosine-Based Motif. *Cell* **135**, 702–713 (2008).

10.   Zhang, H., Cordoba, S.-P., Dushek, O. & Anton van der Merwe, P. Basic residues in the

T-cell receptor ζ cytoplasmic domain mediate membrane association and modulate signaling.

Proc. Natl. Acad. Sci. **108**, 19323–19328 (2011).

11.   James, J. R. & Vale, R. D. Biophysical Mechanism of T Cell Receptor Triggering in a

Reconstituted System. *Nature* **487**, 64–69 (2012).

12.   Minguet, S., Swamy, M., Alarcón, B., Luescher, I. F. & Schamel, W. W. A. Full activation

of the T cell receptor requires both clustering and conformational changes at CD3. *Immunity*

26, 43–54 (2007).

13.   Gil, D., Schamel, W. W. A., Montoya, M., Sánchez-Madrid, F. & Alarcón, B. Recruitment

of Nck by CD3 epsilon reveals a ligand-induced conformational change essential for T cell

receptor signaling and synapse formation. *Cell* **109**, 901–912 (2002).

14.   Natarajan, K. *et al.* An allosteric site in the T-cell receptor Cβ domain plays a critical

signalling role. *Nat. Commun.* **8**, 15260 (2017).

15.   Rangarajan, S. *et al.* Peptide-MHC (pMHC) binding to a human antiviral T cell receptor

induces long-range allosteric communication between pMHC- and CD3-binding sites. *J. Biol.*

Chem. **293**, 15991–16005 (2018).

16.   Das, D. K. *et al.* Force-dependent transition in the T-cell receptor β-subunit allosterically

regulates peptide discrimination and pMHC bond lifetime. *Proc. Natl. Acad. Sci.* **112**, 1517–

1522 (2015).

17.   Wu, W., Shi, X. & Xu, C. Regulation of T cell signalling by membrane lipids. *Nat. Rev.*

Immunol. **16**, 690–701 (2016).

18.   Wang, F., Beck-García, K., Zorzin, C., Schamel, W. W. A. & Davis, M. M. Inhibition of T

cell receptor signaling by cholesterol sulfate, a naturally occurring derivative of membrane

cholesterol. *Nat. Immunol.* **17**, 844–850 (2016).

19.   Notti, R. Q. & Walz, T. Native-like environments afford novel mechanistic insights into

membrane proteins. *Trends Biochem. Sci.* (2022).

20.   Robbins, P. F. *et al.* A Pilot Trial Using Lymphocytes Genetically Engineered with an NY-

ESO-1–Reactive T-cell Receptor: Long-term Follow-up and Correlates with Response. *Clin.*

Cancer Res. **21**, 1019–1027 (2015).

21.   Ulmschneider, M. B. & Sansom, M. S. P. Amino acid distributions in integral membrane

protein structures. *Biochim. Biophys. Acta BBA - Biomembr.* **1512**, 1–14 (2001).

22.   Swamy, M. *et al.* A Cholesterol-Based Allostery Model of T Cell Receptor

Phosphorylation. *Immunity* **44**, 1091–1101 (2016).

23.   Su, Q. *et al.* Cryo-EM structure of the human IgM B cell receptor. *Science* **377**, 875–880

(2022).

24.   Gao, X., Dong, X., Li, X., Liu, Z. & Liu, H. Prediction of disulfide bond engineering sites

using a machine learning method. *Sci. Rep.* **10**, 10330 (2020).

25.   Liu, F., van Breukelen, B. & Heck, A. J. R. Facilitating Protein Disulfide Mapping by a

Combination of Pepsin Digestion, Electron Transfer Higher Energy Dissociation (EThcD), and

a Dedicated Search Algorithm SlinkS*. *Mol. Cell. Proteomics* **13**, 2776–2786 (2014).

26.   Chen, J.-L. *et al.* Structural and kinetic basis for heightened immunogenicity of T cell

vaccines. *J. Exp. Med.* **201**, 1243–1255 (2005).

27.   Kim, S. T. *et al.* The αβ T Cell Receptor Is an Anisotropic Mechanosensor *. *J. Biol.*

Chem. **284**, 31028–31037 (2009).

28.   Lanz, A.-L. *et al.* Allosteric activation of T cell antigen receptor signaling by quaternary

structure relaxation. *Cell Rep.* **36**, 109375 (2021).

29.   Yin, Y., Wang, X. X. & Mariuzza, R. A. Crystal structure of a complete ternary complex of

T-cell receptor, peptide-MHC, and CD4. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 5405–5410

(2012).

30.   Fernandes, R. A. *et al.* T Cell Receptors are Structures Capable of Initiating Signaling in

the Absence of Large Conformational Rearrangements*. *J. Biol. Chem.* **287**, 13324–13335

(2012).

31.   Reinherz, E. L., Hwang, W. & Lang, M. J. Harnessing αβ T cell receptor mechanobiology

to achieve the promise of immuno-oncology. *Proc. Natl. Acad. Sci.* **120**, e2215694120

(2023).

32.   Stewart-Jones, G. B. E. *et al.* Trimeric HIV-1-Env Structures Define Glycan Shields from

Clades A, B, and G. *Cell* **165**, 813–826 (2016).

33.   Lee, J. H., de Val, N., Lyumkis, D. & Ward, A. B. Model Building and Refinement of a

Natively Glycosylated HIV-1 Env Protein by High-Resolution Cryoelectron Microscopy.

Structure **23**, 1943–1951 (2015).

34.   Chouaki Benmansour, N. *et al.* Phosphoinositides regulate the TCR/CD3 complex

membrane dynamics and activation. *Sci. Rep.* **8**, 4966 (2018).

35.   Costello, P. S., Gallagher, M. & Cantrell, D. A. Sustained and dynamic inositol lipid

563   metabolism inside and outside the immunological synapse. *Nat. Immunol.* **3**, 1082–1089

564   (2002).

565   36.   Xu, S. *et al.* Uptake of oxidized lipids by the scavenger receptor CD36 promotes lipid

566   peroxidation and dysfunction in CD8+ T cells in tumors. *Immunity* **54**, 1561-1577.e7 (2021).

567   37.   Kent, A., Longino, N. V., Christians, A. & Davila, E. Naturally Occurring Genetic

568   Alterations in Proximal TCR Signaling and Implications for Cancer Immunotherapy. *Front.*

569   *Immunol.* **12**, (2021).

570   38.   Klebanoff, C. A., Chandran, S. S., Baker, B. M., Quezada, S. A. & Ribas, A. T cell

571   receptor therapeutics: immunological targeting of the intracellular cancer proteome. *Nat. Rev.*

572   *Drug Discov.* **22**, 996–1017 (2023).

575 **Figures and Legends**

576 **Figure 1**
577

Figure 1: Architecture of the TCR–CD3 in a native-like lipid environment. (a) Cryo-EM

maps (top, middle) and space-filling solvent-excluded surfaces for the models (bottom; herein,

model surfaces) of the TCR–CD3 in a detergent micelle (GDN) and in nanodiscs (conformations

ND-I and ND-II). Top row shows cryo-EM density (without local refinement) after B-factor

sharpening, contoured to a low threshold to highlight micelle and nanodisc positioning. Red

dashed box indicates approximate plasma membrane span as inferred from the micelle and

nanodisc densities (red arrows and labels). Extracellular and intracellular faces indicated as e.c.

and i.c., respectively. Middle row shows DeepEMhancer-sharpened maps after local refinement,

accentuating the protein and glycan densities. Protein regions indicated at far right abbreviated

as follows: var, variable domains; ecto, ectodomains; const, constant domains; JM,

juxtamembrane linkers; TM, transmembrane helices. (b) Comparison of GDN and ND-I models,

highlighting ectodomain closure in the context of a lipid bilayer. Model backbone cartoon inside

transparent model surfaces shown. (c,d) Comparison of GDN and ND-II models, highlighting

juxtamembrane linker compaction in the direction normal to the lipid bilayer (green dashed box)

and coincident splay of the TM helices (cyan lines and angular measures). The CD3$\varepsilon\delta$ and

CD3$\varepsilon\gamma$ heterodimers are shown in (c) and (d), respectively. (e) Overlay of the TM helices in the

detergent micelle and lipid bilayer (both ND-I and ND-II), viewed from the extracellular space

showing similar overall arrangement. (f) Comparison of GDN (transparent grey) and ND-II (solid,

color-coded) transmembrane helices aligned to TCRα residues 245-260 and viewed parallel to

the plasma membrane. Green and orange arrows highlight increased splay of the TCRβ

transmembrane helix and the carboxy-terminal end of the CD3ζ TM helix, respectively. Blue

arrow highlights the unstructuring and deflection of the first helical turn at the amino-terminus of

the TCRα transmembrane helix in the lipid bilayer.

Figure 2

605
606
607

Figure 2: TCR–CD3–lipid interactions. (a) Comparison of the cholesterol-binding sites of the

TCR in detergent (top) and in nanodiscs (bottom). Overview shown at left and insets show the

cholesterol-binding sites in detail (left) and with protein removed (right) to highlight the deflection

of the branched aliphatic tails. $Chol_B$ denotes the cholesterol proximal to the ectodomains and

the TCRβ transmembrane helix; $Chol_A$ denotes the cholesterol proximal to the cytoplasmic face

and the TCRα transmembrane helix. (b) Sections parallel to the membrane plane highlighting

the opening of the cholesterol-binding groove to the lipid environment in nanodiscs relative to

that in detergent. Sections are shown at the levels of $Chol_A$ and $Chol_B$, as indicated. Density

from locally refined and B-factor-sharpened cryo-EM maps is shown as black mesh,

cholesterols in lime green. Dashed circle highlights structured lipid density in the nanodisc

bilayer adjacent to the opened cholesterol-binding groove. Insets show the cholesterol binding

groove from the same angle as the main panels, but with both cholesterols shown in the slice,

highlighting intercalation of the deflected $Chol_B$ aliphatic chain between TM helices in nanodiscs

(black dashed oval) as compared to the structure in GDN.

Figure 3

626
627
628

Figure 3: Conserved CD3 surface patches form extensive membrane contacts. (a) ND-II

model surface colored by sequence conservation (Consurf score 1 through 9) and yellow

unmodeled map density from the nanodisc enshrouding the TCR–CD3 TM helices and

membrane-proximal ectodomain surfaces. When visualized with a higher contouring threshold

(right), structured lipid density adjacent to the CD3 ectodomains persisted (red dotted box). Note

also the relative sequence conservation in the CD3 TM helices and membrane-adjacent

ectodomain surfaces, but not solvent-exposed surfaces. (b) Top, ND-II space-filling model for

orientation. Arrows and eyes indicate the viewing direction for inspection of the CD3$\varepsilon\gamma$ and

CD3$\varepsilon\delta$ interfaces examined in the lower panels. Bottom, overviews orienting the close-up views

in panels c through e. Red dashed lines denote the border between TM and ectodomain (Ecto)

surfaces. (c) Close-up views of the membrane-proximal face of the CD3 heterodimers, with

black triangles highlighting the conserved surface patch at the subunit interface. (d) Lipid

density (yellow) is most robust over this conserved patch, visualized at high contouring

threshold (as in panel a, right). (e) The conserved ectodomain patch is relatively electropositive

compared to the surrounding ectodomain surfaces colored by electrostatic potential (red to blue,

-10 to 10 kcal/(mol*e)).

Figure 4
648

Figure 4: Confirmation of the TCR–CD3 resting conformation *in situ.* (a) Comparison of

$C\alpha$-$C\alpha$ distances between apposing faces of $CD3\delta$ and $TCR\alpha$ in detergent and in a lipid bilayer,

highlighting plausible engineered disulfide positions in ND-I (distances in blue shades) that are

not geometrically plausible in GDN (distances in yellow, orange, red, magenta). (b) Non-

reducing SDS-PAGE gel of whole-cell lysates from HEK293T cells infected with the indicated

TCR–CD3 virus genotypes Western blotted for EGFP-tagged $TCR\beta$. Engineered disulfides

yielding a $TCR\alpha\beta$–$CD3\delta$ covalent adduct resulted in an upward gel shift (asterisks). Color

coding and distances as in panel a. (c,d) Design of intrachain disulfides in the closed

conformation of $TCR\alpha$ (c) and $TCR\beta$ (d). Distances color coded as above. (e,f) Mass

spectrometric detection of intrachain disulfides in $TCR\alpha$ (e) and $TCR\beta$ (f). EThcD fragment

spectra of disulfide-linked dipeptides are shown. Gas-phase reduced linear peptides are shown

in red (peptide A) and blue (peptide B) with the reduced Cys highlighted in bold. Fragment ions

of the two linear peptides are shown in magenta (fragments of peptide A) and cyan (fragments

of peptide B). Charge-reduced precursor ions are shown in gray.

Figure 5

Figure 5: Basal conformational variability of the TCR–CD3. (a) Comparison of ND-I

(transparent cyan) and ND-II (red), shown here as model surfaces, aligned on the TCRα

transmembrane helix. (b,c) Gradient arrows highlight the domain movements underlying the

transition between ND-I (cyan) and ND-II (red). (d) TCRβ and CD3γ TM helix movement away

from the cholesterol-binding site during the transition from ND-I to ND-II, with arrows indicating

the accompanying movements of the cholesterols in the sterol-binding groove. (e) Comparison

of the TCRα and CD3δ lateral glycosylation sites (red and cyan dashed lines, respectively) in

the density maps showing increasing ordering of the glycan array with ectodomain closure

(purple dashed line). Insets highlight modeled asparagine-linked glycan cores (or partial cores)

modeled into these densities in GDN and ND-I. (f) Comparison of the CD3δ N38 residue in

GDN, ND-II and ND-I. Grey surface is the locally refined, DeepEMhancer-sharpened cryo-EM

density. (g) Modeling HLA-β2M-antigen (grey) binding to the TCR in the indicated

conformations. (h) Cross section through the HLA-TCR interface in ND-I shows close

association of HLA and CD3ε_D, but no large clashes. (i) Similarly, modeling HLA onto the locally

refined, DeepEMhancer-sharpened ND-I density does not show any large clashes between HLA

and unmodelled density.

Figure 6
691

692
693
694
695

Figure 6: Model of TCR–CD3 activation. (a) Proposed model of TCR–CD3 conformational

dynamics. The unliganded TCR–CD3 explores resting subconformations (between ND-I and

ND-II). Ligand-induced activation is mediated by a conformational transition from ND-II to an

open, extended conformation, like in GDN. Magenta arrows and text indicate movements

required to transition to the next conformation in the series. (b) Changes in the TM helices

between hypothesized "resting" (ND-I shown) and "active" (GDN shown) conformations. (c)

Flow cytometry histogram from J8Zb2m-α-β- cells post tetramer-APC enrichment and stained

as indicated (n = 3 replicates). Cells were untransduced (top) or transduced with a virus

expressing the TCR$\alpha\beta$ 1G4 genotype indicated (middle, bottom). Vertical axis is cell count

normalized to peak; horizontal axis is fluorescence in arbitrary units. Red line indicates

fluorescence intensity mode (histogram peak) for wild-type 1G4. (d) Time-dependent T-cell

activation quantified using an NFAT-ZsGreen reporter by live cell imaging. Top shows ZsGreen

fluorescence over time after addition of indicated concentrations of HLA-tetramer or

PMA/ionomycin for 1G4 wild-type (red) or TCRα S104C/V182C mutant-transduced cells (cyan).

Data shown are mean +/- standard deviation using biologic triplicates. Peak CD3-TCR

activation amplitudes are compared for each condition at the 8 μg/mL HLA-tetramer peak

(t=16h) for all HLA-tetramer concentrations or at the PMA/Ionomycin peak (t=10h) specifically

for that condition (bottom). P values calculated by unpaired, two-tailed t-test shown for each

comparison. $P < 0.05$ highlighted in red.

Extended Data Figures and Legends
717
718 **Extended Data Figure 1**
719

720
721
722
723

Extended Data Figure 1: The TCR–CD3 in detergent. (a) Cartoon representations of

previously published structures of the TCR–CD3 by itself and in complex with HLA-β2M-antigen

determined in detergent micelles by cryo-EM. Models are colored by chain as indicated. (b)

Superposition of published structures of a TCR–CD3 by itself (red) or in complex with HLA-β2M-

antigen (cyan), showing no meaningful structural change throughout the TCR structure upon

ligand binding. (c) Gold-standard Fourier shell correlation (FSC) curves for the map of the TCR–

CD3 in GDN (herein, simply "GDN") determined in this study after non-uniform refinement. (d)

FSC curve for the locally refined GDN map (mask excluded micelle density). While resolution

was not nominally improved by masking out the micelle, map interpretability was qualitatively

improved, and this map was used for further analysis. (e) Three-dimensional (3D) FSC analysis

for the locally refined GDN map, showing relatively little map anisotropy. (f) Central slices of the

locally refined GDN map, colored by local resolution estimate, as indicated in the panel. (g)

Backbone representation of the GDN structure compared with those of TCR–CD3 sequences

previously determined in other studies. Models containing HLA-β2M-antigen have had those

atoms removed for this analysis. (h) GDN model colored according to sequence identity among

the four TCR–CD3 sequences (this study and three priors) with cryo-EM structures determined

in detergent micelles.

Extended Data Figure 2

745
746
747
748

Extended Data Figure 2: Map assessment for ND-I and ND-II. (a) Gold-standard FSC curves

for the ND-I map after non-uniform refinement. (b) FSC curves for the locally refined ND-I map

(mask excluded nanodisc density). While resolution was not nominally improved by masking out

the nanodisc, map interpretability was qualitatively improved, and this map was used for further

analysis. (c) 3D FSC analysis for locally refined ND-I. Note the more extensive anisotropy than

was seen for the TCR–CD3 in detergent micelles. (d) Central slice of locally refined ND-I,

colored by local resolution estimate, as indicated in the panel. (e) Gold-standard FSC curves for

the ND-II map after non-uniform refinement. (f) FSC curves for the locally refined ND-II map

(mask excluded nanodisc density). As with ND-I, nominal resolution was not improved by

masking out the nanodisc, but map interpretability was qualitatively improved, and this map was

used for further analysis. (g) 3D FSC analysis for locally refined ND-II. Note again the more

extensive anisotropy than was seen for the TCR in detergent micelles. (h) Central slice of locally

refined ND-II, colored by local resolution estimate, as indicated in the panel.

Extended Data Figure 3

766
767
768
769

Extended Data Figure 3: Structural detail of the TCR–CD3 in nanodiscs. (a,b) Cartoon

representations of ND-I (solid) and GDN (transparent gray) TCRα (panel a) and TCRβ (panel b),

aligned to the respective TCR constant domains. Arrows indicate the movements of the variable

domains relative to the constant domains that comprise the "closure" of the variable domains in

ND-I/II relative to GDN. (c) The bent TM helices of the ND-I CD3ε_D heterodimer. Splay

measured at the level of the first helical turn (cyan) is greater than that of the helices overall

(orange). (d) Cartoon representation of the indicated models with basic and aromatic residues

highlighted as space-filling spheres. Blue dashed box highlights the clusters of basic and

aromatic residues modeled at the carboxy-termini of the TM helices in GDN, but unstructured in

ND-I and ND-II. Color legend applies to all figure panels. (e,f) Density projecting from the

carboxy-termini of the TM helices modeled in ND-II that could not be confidently modeled, but

likely corresponding to continuation of the polypeptide chains. Grey, DeepEMhancer-sharpened

maps. Dashed lines encircle the densities in question. (g) Density corresponding to the

cholesterols modeled in ND-I, ND-II, and GDN. The transparent green surface is the indicated

locally refined, B-factor-sharpened cryo-EM map, cropped to a zone of 2.2 Å around the

cholesterols (black). (h-k) Additional densities likely attributable to lipids in the ND-II map.

Proteins are shown as model surfaces, and cholesterol in stick representation. Yellow surface is

the ND-II B-factor-sharpened map after local refinement (h) or non-uniform refinement (i-k) with

the modeled densities subtracted to highlight unmodeled densities.

Extended Data Figure 4
792

793
794
795
796

Extended Data Figure 4: Insights into BCR structure and membrane interaction. (a,b)

Surface representation of the human B-cell receptor (BCR) model, as determined in a detergent

micelle by cryo-EM (PDB 7XQ8)[23]. The model is colored by chain (a) and by sequence

conservation (b, scale applies to remainder of figure). Red arrows and eye icons labeled

CD79$\alpha\beta$ and heavy-heavy' indicate the direction for the magnified views in panels (c) and (d),

respectively. (c) Membrane-proximal face of the CD79$\alpha\beta$ interface, colored by conservation

(left, as in panel a) and electrostatic potential (right). Black triangle highlights region of high

sequence conservation and corresponding surface electropositivity. However, this patch is

oriented away from the micelle surface in the structure in detergent (dashed arrow to panel e).

(d) Membrane-proximal face of the IgM heavy chains, with black box highlighting broad

sequence conservation and largely electronegative conserved face. Dashed arrow indicates

contact between this surface patch and the detergent micelle density in panel e. Electrostatic

potential scale applies to panels c and d. (e) Surface model of the BCR with the unmodeled

map density shown (transparent gray). Note the detergent micelle enshrouding the TM helices

and making contact with the electronegative heavy-heavy' interface (red dashed arrow) but not

the electropositive CD79$\alpha\beta$ interface (blue dashed arrow), which is the opposite of what might

be expected in a phospholipid membrane.

Extended Data Figure 5

818
819
820
821

Extended Data Figure 5: Engineered disulfide crosslinking. (a) Location of all native

disulfides (solid cyan and magenta ovals) and unpaired cysteines (dashed black ovals) in this

TCR construct. Disulfides indicated by cyan ovals are amenable to mass spectrometric

detection as peptide pairs after EThcD, whereas disulfides indicated by magenta ovals are not

unambiguously detectable by this approach, as they are C-X-X-C lariats or symmetric between

CD3ζ homodimers. Subunits colored as indicated to the right. (b) Non-reducing SDS-PAGE gel

of NiNTA-affinity-purified TCR complex from HEK293T cells infected with the indicated TCR

virus genotypes and Western blotted for EGFP-tagged TCRβ. Engineered disulfides yielding a

TCR$\alpha\beta$–CD3δ covalent adduct results in an upward gel shift (asterisks). Color coding and

distances as in Figure 4a,b. (c,d) Acquastain-stained SDS-PAGE gels of NiNTA-affinity-purified

TCR complex reconstituted into nanodiscs and purified by size-exclusion chromatography

(SEC). Virus genotypes indicated above gels. The presence (+) or absence (-) of reducing

agent dithiothreitol (DTT) in the sample-loading buffer is indicated. In panel c, dashed line

markers 1, 2, and 3 indicate the inferred electrophoretic mobility of the TCR$\alpha\beta$::EGFP–

CD3δ adduct, TCR$\alpha\beta$::EGFP, and TCRβ::EGFP, respectively. Note the mobility shift between

the nonreduced wild-type TCR complex and the non-reduced TCRα(G79C)–CD3δ(D77C)

mutant complex. Molecular weight ladder (L) is color-coded with magenta, orange, green, and

white asterisks denoting 250 kDa, 150 kDa, 100 kDa, and 75 kDa, respectively. In panel d, the

non-reduced TCRα(G79C)–CD3δ(D38C) mutant complex displays an electrophoretic mobility

similar to that of the non-reduced TCRα(G79C)–CD3δ(D77C) mutant complex, consistent with

engineered disulfide formation. (e) Design of an engineered disulfide bond plausible in the GDN

conformation but not in the closed ND-I conformation.

845
846

847   **Extended Data Figure 6**
848

Chain	Residue	GDN	ND-I	ND-II
TCRα	N41			
TCRα	N82			
TCRα	N166			
TCRα	N200			
TCRα	N211			
TCRα	N247			
TCRβ	N84			
TCRβ	N202			
CD3δ	N38			
CD3δ	N74			
CD3γ	N52			
CD3γ	N92			

849
850
851

Extended Data Figure 6: Basal conformational variation and glycosylation of the TCR–

CD3. (a,b) Cryo-EM maps for ND-I (transparent cyan) and ND-II (solid red) aligned to the TCRα

transmembrane helix. The B-factor-sharpened map without local refinement and

DeepEMhancer-sharpened, locally refined maps are shown (a and b, respectively). (c) Relative

positions of the TM helices for ND-I and ND-II (colored as in panel a) viewed along the axis

perpendicular to the membrane plane (left) and parallel to the membrane (right). Dashed circle

indicates cholesterol-binding groove. (d) Graphical representation of partially modeled N-linked

glycans in GDN, ND-I, and ND-II. All asparagine residues within an N-X-S/T motif in the

structure are included in this list for completeness; however, TCRα N247 is located in the TM

domain and is unlikely to undergo glycosylation. Green, modeled; magenta, not modeled. (e)

Glycans (grey sticks) in a structured array on the surface of HIV Env as determined by x-ray

crystallography (PDB 5FYJ)[32]. Protein shown as cartoon backbone. 2Fo-Fc map contoured to

1σ and clipped to within 5 Å of the model. (f) Space-filling model of HLA docked onto ND-I with

the CD3ϵ_D ectodomain from GDN docked in place of that from ND-I, showing minimal, if any

steric clashing. Cartoon representation of the remainder of ND-I shown for orientation. Inset

shows alignment of CD3ϵ_D from GDN and ND-I by their ectodomains. Red dashed circle

highlights surface loop (residues 107-109) not modeled in ND-I (purple dashed line) but

modeled in GDN. (g) Alignment of TCR variable domains in this study to PDB 2BNR, a crystal

structure of the 1G4 TCR bound to HLA[26]. (h) Alignment of the CD4-Class II HLA-TCR ternary

complex structure (PDB 3T0E)[29] to the Class I HLA docked into the our structures as in panel F

and Figure 5. The Class II HLA from 3T0E is shown in yellow, and CD4 in cyan. Other color

coding as indicated in figure. Dashed black line represents the approximate boundary of the

lipid bilayer outer leaflet.

Extended Data Tables

Extended Data Table 1: Model building and refinement statistics.

	GDN	ND-I	ND-II
Model			
Composition (#)			
Chains	10	12	10
Atoms	8440 (Hydrogens: 0)	6460 (Hydrogens: 0)	6242 (Hydrogens: 0)
Residues	Protein: 1052 Nucleotide: 0	Protein: 836 Nucleotide: 0	Protein: 821 Nucleotide: 0
Water	0	0	0
Ligands	BMA: 3	BMA: 1	BMA: 1
	NAG: 15	NAG: 9	NAG: 3
	CLR: 2	CLR: 2	CLR: 2
Bonds (RMSD)			
Length (Å) (# > 4s)	0.003 (0)	0.004 (0)	0.003 (0)
Angles (degrees) (# > 4s)	0.923 (6)	0.855 (6)	0.906 (3)
MolProbity score	1.9	2.71	2.67
Clash score	13.57	21.05	18.43
Ramachandran plot (%)			
Outliers	0.1	0.26	0.4
Allowed	3.67	7.14	9.49
Favored	96.24	92.6	90.12
Rama-Z (Ramachandran plot Z-score, RMSD)			
whole (N = 1036)	0.51 (0.26)	-2.15 (0.27)	-1.91 (0.29)
helix (N = 214)	0.63 (0.34)	-1.96 (0.42)	-1.14 (0.41)
sheet (N = 286)	1.85 (0.32)	-0.40 (0.37)	0.54 (0.37)
loop (N = 536)	-0.64 (0.26)	-1.77 (0.26)	-2.35 (0.28)
Rotamer outliers (%)	0.56	3.72	2.99
Peptide plane (%)			
Cis proline/general	11.6/0.2	6.1/0.0	3.3/0.0
Twisted proline/general	0.0/0.1	0.0/0.0	0.0/0.3
CaBLAM outliers (%)	3.04	5.6	7.32
ADP (B-factors)			
Iso/Aniso (#)	8440/0	6460/0	6242/0
min/max/mean			
Protein	78.97/260.88/146.95	46.79/362.28/144.81	51.96/387.42/144.45
Nucleotide	---	---	---

Ligand	92.44/292.30/180.12	78.92/388.12/221.92	124.98/195.38/160.46
Water	---	---	---
Occupancy			
Mean	1	1	1
occ = 1 (%)	100	100	100
0 < occ < 1 (%)	0	0	0
occ > 1 (%)	0	0	0
Data			
Box			
Lengths (Å)	(88.56, 101.52, 138.24)	(95.04, 87.48, 110.16)	(92.88, 92.88, 101.52)
Supplied Resolution (Å)	3.3	3.3	3.1
Resolution Estimates (Å), Masked			
d FSC (half maps; 0.143)	3.4	3.2	3.1
d 99 (full/half1/half2)	4.3/3.3/3.3	4.5/3.7/3.7	4.2/3.4/3.4
d model	3.7	3.3	3.1
d FSC model (0/0.143/0.5)	3.2/3.3/3.8	3.2/3.4/7.4	3.1/3.3/7.2
Resolution Estimates (Å), Unmasked			
d FSC (half maps; 0.143)	3.5	3.3	3.1
d 99 (full/half1/half2)	4.3/2.5/2.5	4.5/3.5/3.4	4.2/3.2/3.2
d model	3.8	3.3	3.1
d FSC model (0/0.143/0.5)	3.3/3.4/3.8	3.2/3.5/7.6	3.1/3.4/7.3
Map min/max/mean	-0.60/1.36/0.06	-0.65/1.68/0.09	-0.67/1.77/0.10
Model vs. Data			
CC (mask)	0.82	0.57	0.59
CC (box)	0.81	0.65	0.68
CC (peaks)	0.7	0.41	0.48
CC (volume)	0.82	0.58	0.59
Mean CC for ligands	0.72	0.56	0.58

Extended Data Table 2: Disulfide-linked peptides identified by cross-linking mass

spectrometry. Disulfide-linked peptides identified for each TCR genotype. Peptides with an

XLinkX score greater than 30 are shown, as no native peptides were observed below this

threshold. Color coding as follows: green, expected native disulfide; cyan, expected engineered

disulfide; red, non-native disulfide or contaminant; magenta, off-target disulfide involving the

engineered cysteines.

XLINKX SCORE	SEQUENCE 1	CHAIN 1	CYS 1	SEQUENCE 2	CHAIN 2	CYS 2
TCRα(WT), TCRβ(WT)						
120.57	V[C]LATGFYPDHVEL	B	163	[C]QVQFY	B	228
106.91	LT[C]DAEAK	G	46	GMYQ[C]K	G	87
95.96	Q[C]AQDMNHEY	B	42	F[C]ASSY	B	110
90.84	T[C]PQYPGSEILW	E	49	VN[C]NTSITW	D	37
73.16	FPSPESS[C]DVK	A	228	AD[C]GFTSESY	B	263
70.28	SVPEGENLVLN[C]SFTDSAIY	A	42	L[C]AVRPLY	A	109
69.67	SNKSDFA[C]ANAF	A	206	SV[C]LF	A	156
58.32	T[C]PQYPGSEILW	E	49	[C]QVQFY	B	228
53.59	VFVN[C]NTSITW	D	37	GIYR[C]NGTDIY	D	73
40.68	NNSIIPEDTFFPSPESS[C]DVK	A	228	ATLV[C]L	B	163
30.56	T[C]PQYPGSEILW	E	49	YV[C]YPR	E	98
TCRα(S104C-V182C), TCRβ(WT)						
120.8	LT[C]DAEAK	G	46	GMYQ[C]K	G	87
106.26	V[C]LATGFYPDHVEL	B	163	R[C]QVQFY	B	228
104.38	Q[C]AQDMNHEY	B	42	F[C]ASSY	B	110
92.7	GIYR[C]NGTDIYK	D	73	VFVN[C]NTSITW	D	37
80.51	SVPEGENLVLN[C]SFTDSAIY	A	42	L[C]AVRPLY	A	109
78.49	SNKSDFA[C]ANAF	A	206	SV[C]LF	A	156
72.7	T[C]PQYPGSEILW	E	49	[C]QVQFY	B	228
72.62	IAASQPGD[C]ATY	A	104	ITDKT[C]LDMR	A	182
70.28	SVPEGENLVLN[C]SFTDSAIY	A	42	[C]NGTDIY	D	73
65.27	V[C]LATGFYPDHVEL	B	163	T[C]PQYPGSEILW	E	49
60.55	SDFA[C]ANAF	A	206	ITDKT[C]L	A	182
56.37	FPSPESS[C]DVK	A	228	AD[C]GFTSESY	B	263
49.26	ITDKT[C]L	A	182	SV[C]LF	A	156
48.23	T[C]PQYPGSEILW	E	49	VN[C]NTSITW	D	37
45.66	T[C]PQYPGSEILW	E	49	YV[C]YPR	E	98
TCRα(WT), TCRβ(T130C-H172C)						
99.32	QEDGSVLLT[C]DAEAK	G	46	GMYQ[C]KGSQNK	G	87
96.82	Q[C]AQDMNHEY	B	42	F[C]ASSY	B	110
88.95	FGEGSRL[C]VLEDLK	B	130	ATGFYPD[C]VELSW	B	172
79.2	SNKSDFA[C]ANAF	A	206	SV[C]LF	A	156
72.18	SVPEGENLVLN[C]SFTDSAIY	A	42	L[C]AVRPLY	A	109
71.14	GIYR[C]NGTDIYK	D	73	VFVN[C]NTSITW	D	37
63.42	L[C]VLEDLK	B	130	LT[C]DAEAK	G	46
62.17	L[C]VLEDLK	B	130	SV[C]LF	A	156
54.2	L[C]VLEDLK	B	130	F[C]ASSY	B	110
53.77	T[C]PQYPGSEILW	E	49	L[C]VLEDLK	B	130

53.5	FPSPESS[C]DVK	A	228	AD[C]GFTSESY	B	263
47.92	T[C]PQYPGSEILW	E	49	[C]QVQFY	B	228
41.91	L[C]VLEDLK	B	130	[C]QVQFY	B	228
41.51	AD[C]GFTSESY	B	263	L[C]VLEDLK	B	130
38.79	T[C]PQYPGSEILW	E	49	YV[C]YPR	E	98
36.32	ATGFYPD[C]VEL	B	172	[C]LSSR	B	207
34.92	VFVN[C]NTSITW	D	37	GMYQ[C]K	G	87
30.54	LT[C]DAEAK	G	46	[C]QVQFY	B	228
TCRα(WT), TCRβ(K33C-L135C)						
96.43	GIYR[C]NGTDIYK	D	73	VFVN[C]NTSITW	D	37
95.43	LT[C]DAEAK	G	46	GMYQ[C]K	G	87
90.79	ATLV[C]LATGFYPDHVEL	B	163	R[C]QVQFY	B	228
81.71	LTVLED[C]K	B	135	ATLV[C]L	B	163
81.57	Q[C]AQDMNHEY	B	42	F[C]ASSY	B	110
81.57	LTVLED[C]K	B	135	[C]QVQFY	B	228
79.2	SNKSDFA[C]ANAF	A	206	SV[C]LF	A	156
76.07	T[C]PQYPGSEILW	E	49	[C]QVQFY	B	228
72.9	SVPEGENLVLN[C]SFTDSAIY	A	42	L[C]AVRPLY	A	109
69.49	LTVLED[C]K	B	135	F[C]ASSY	B	110
65.27	V[C]LATGFYPDHVEL	B	163	T[C]PQYPGSEILW	E	49
64.22	ATLV[C]LATGFYPDHVEL	B	163	ATLV[C]LATGFYPDHVEL	B	163
53.91	T[C]PQYPGSEILW	E	49	VN[C]NTSITW	D	37
48.05	V[C]LATGFYPDHVEL	B	163	SDFA[C]ANAF	A	206
47.05	LTVLED[C]K	B	135	SV[C]LF	A	156
45.66	V[C]LATGFYPDHVEL	B	163	SV[C]LF	A	156
43.9	T[C]PQYPGSEILW	E	49	V[C]YPR	E	98
36.08	FPSPESS[C]DVK	A	228	AD[C]GFTSESY	B	263
35.13	T[C]PQYPGSEILW	E	49	LTVLED[C]K	B	135
35.04	T[C]PQYPGSEILW	E	49	[C]AVRPL	A	109
34.06	V[C]LATGFYPDHVEL	B	163	VLN[C]SFTDSAIY	A	42
30.82	SDFA[C]ANAF	A	206	LTVLED[C]K	B	135

886