DNA polymerase diversity reveals multiple incursions of Polintons during nematode evolution

Dae-Eun Jeong[1,7], Sameer Sundrani[2,6], Richard Nelson Hall[2], Mart Krupovic[3], Eugene V. Koonin[4], and Andrew Z. Fire[1,5,7]

[1]Department of Pathology, Stanford University School of Medicine, Stanford, California, USA

[2]Department of Bioengineering, Stanford University, Stanford, California, USA

[3]Institut Pasteur, Université Paris Cité, CNRS UMR6047, Archaeal Virology Unit, Paris, France

[4]National Center for Biotechnology Information, National Library of Medicine, National Institutes of Health, Bethesda, Maryland, USA

[5]Department of Genetics, Stanford University School of Medicine, Stanford, California, USA

[6]Current address: Vanderbilt University School of Medicine, Nashville, Tennessee, USA

[7]Corresponding (dejeong@stanford.edu, afire@stanford.edu)

Abstract

Polintons are dsDNA, virus-like self-synthesizing transposons widely found in eukaryotic genomes. Recent metagenomic discoveries of Polinton-like viruses are consistent with the hypothesis that Polintons invade eukaryotic host genomes through infectious viral particles. Nematode genomes contain multiple copies of Polintons and provide an opportunity to explore the natural distribution and evolution of Polintons during this process. We performed an extensive search of Polintons across nematode genomes, identifying multiple full-length Polinton copies in several species. We provide evidence of both ancient Polinton integrations and recent mobility in strains of the same nematode species. In addition to the major nematode Polinton family, we identified a group of Polintons that are overall closely related to the major family, but encode a distinct protein-primed B family DNA polymerase (pPolB) that is related to homologs from a different group of Polintons present outside of the *Nematoda*. Phylogenetic analyses on the pPolBs support the evolutionary scenarios in which these extrinsic pPolBs that seem to derive from Polinton families present in oomycetes and molluscs replaced the canonical pPolB in subsets of Polintons found in terrestrial and marine nematodes, respectively, suggesting inter-phylum horizontal gene transfers. The pPolBs of the terrestrial nematode and oomycete Polintons share a unique feature, an insertion of a HNH nuclease domain, whereas the pPolBs in the marine nematode Polintons share an insertion of a VSR nuclease domain with marine mollusc pPolBs. We hypothesize that horizontal gene transfer occurs among Polintons from widely different but cohabiting hosts.

Polintons/Mavericks (hereafter referred to as Polintons) were discovered as double-
stranded DNA (dsDNA) transposons that encode a self-synthesizing, protein-primed DNA
polymerase of the B family (pPolB) and a retroviral-element-like integrase (INT) (hence the
name Polintons)[1,2]. Thus far primarily identified and characterized *in silico*, Polintons are among
the larger known DNA transposons found widely across unicellular and multicellular eukaryotes,
ranging from 13-25 kilobase pair (kbp) with 100-1500 base pair (bp) terminal inverted repeats
(TIR) and with 5-8 bp target site duplications (TSD)[1-5]. In addition to pPolB and INT, Polintons
typically encode a core set of conserved genes encoding homologs of dsDNA virus proteins
involved in virion morphogenesis, such as adenovirus-like maturation protease (PRO), genome
packaging ATPase, and major and minor capsid proteins (MCPs and mCPs, respectively)[5-11].
Polintons generally occupy a low fraction of their host genome; however, there are genomes
with much higher occurrence, such as the excavate *Trichomonas vaginalis* where Polintons
expanded to occupy more than 30% of the genome[3,12-18].
While Polintons were first thought to be transposable elements, recent experimental and
metagenomic discoveries of Polinton-like virophages and Polinton-like viruses suggest that
Polintons found in multicellular eukaryotic genomes are capable of transmission between host
genomes as infectious viral particles[19-25]. A combination of inferred protein characteristics (the
presence of genes encoding homologs of the major and minor capsid protein and the ATPase
and protease involved in capsid maturation), strongly supports this possibility[5,7]. A dual
capability as viruses and transposons[5,7,18] would afford Polintons an ability to propagate both
within and between genomes.
Nematode genomes provide an excellent opportunity to observe transmissible genetic
elements[26,27] from diverse environments. The nematode phylum includes both parasitic and
free-living species, with the latter having successfully adapted to diverse environmental niches
including soil, rotten fruits, freshwater and seawater[28-33]. Because nematodes interact with
diverse types of organisms (viruses, bacteria, protists, animals, and plants)[34-38], the phylum is
likely to have been exposed to a broad cross section of potentially infectious or transmissible
elements. Thus, nematode genomes serve as a potentially-broad detector for elements capable
of interspecies transfer and acquisition. Although several RNA viruses have been reported to
infect or to be associated with nematodes[35,39-46], only limited information about possible dsDNA
virus infection and evolution in nematodes is available[47,48].
In silico analyses have identified a handful of Polintons from nematode species[1,2,25,49]
including *C. briggsae*. These nematode Polintons encode a core set of Polinton proteins
including pPolB, INT, ATPase, PRO, mCP and MCP, and nematode Polinton-specific protein
PC1[1], which was recently proposed to function as a fusogen[49]. The presence of a fusogen
suggests that nematode Polintons may form an enveloped viral particle[50]. However, the natural
distribution and evolution of Polintons across nematode genomes have not been extensively
studied.
In this study, we use the diversity of available nematode genome sequences to
investigate the natural variation and history of Polintons in this phylum. Of particular note, we
observed an apparently modular history of the elements, with two distinct classes of pPolBs in
otherwise-similar nematode Polintons. Additional analysis of these two classes outside of
Nematoda suggests that Polintons have acquired and exchanged pPolB genes during evolution.

The distribution of pPolB homologs in the Earth biome suggests that Polintons act as gene
transfer vehicles.
3
4
Results and Discussion
6
Two groups of Polintons encoding distinct DNA polymerases in *Caenorhabditis*
nematodes
To explore the distribution and evolution of Polintons in nematodes, we searched for full-
length Polintons across 354 nematode genomes from 180 species and identified 266 Polintons
in 66 genomes of 29 species (Extended Data Table 1; see Methods). While surveying the
identified Polinton sequences, we noticed that a group of nematode Polintons encompassed a
pPolB that was only weakly similar to the typical pPolB encoded by other nematode Polintons,
in contrast to all other core Polinton genes and terminal inverted repeats (TIRs) that were highly
conserved. We denoted the distant pPolB group 'pPolB2' to distinguish it from the typical
nematode Polinton pPolBs (hereafter 'pPolB1') (Fig 1A).
The unexpected identification of two distinct groups of pPolBs in the nematode Polintons
prompted us to examine the molecular diversification of Polinton proteins in nematodes in
greater detail. To compare the extent of protein sequence similarity among different Polinton
proteins, we constructed the corresponding multiple sequence alignments and calculated the
percentage amino acid identity for each pair of homologous proteins from Polintons of *C.*
briggsae (*C. briggsae* was chosen because this species was found to encompass the greatest
number of both pPolB1- and pPolB2-class Polintons). The resulting comparisons showed 92%
median identity for MCP, 85% for Fusogen, 81% for ATPase, 80% for INT, 66% for PRO, 78%
for puORF, 68% for mCP, 77% for pPolB1 and 91% for pPolB2 (Fig 1B), indicating that these
proteins are highly conserved among *C. briggsae* Polintons.
In a sharp contrast, pairwise comparisons between pPolB1 and pPolB2 showed a
median of 16% amino acid sequence identity. The Polinton pPolBs belong the protein-primed
family B DNA polymerases that contain conserved exonuclease motifs (Exo I, II, and III)
required for proofreading activity and polymerase motifs (Pol A, B, and C) required for DNA
polymerization activity[51-54]. Examination of the multiple sequence alignment of representative
pPolB1 and pPolB2 proteins from Polintons of 19 nematode species with the prototypical Phi29
phage pPolB indicated that all three exonuclease and all three polymerase motifs are conserved
in both pPolB1 and pPolB2 (Extended Data Fig 1 and Extended Data Fig 2A), suggesting that
both groups of the nematode Polintons possess fully functional DNA polymerases.
Given the low sequence similarity between pPolB1 and pPolB2, we also compared the
structural models of the two proteins obtained using AlphaFold2[55] through ColabFold[56].
Structural alignment and superposition of *C. briggsae* pPolB1 and pPolB2 structures show
evidence of a close structural similarity between the two groups of pPolBs, in spite of the low
sequence identity, in line with the prediction that both are active DNA polymerases (Extended
Data Fig 2B-D and Fig 2A). In addition to the conserved structural core of pPolB1 and pPolB2,
the structural alignment also revealed potential pPolB1- and pPolB2-specific domains (Fig 2A).
In particular, we identified a pPolB2-specific domain that is conserved across all nematode
pPolB2 proteins except the pPolB2 from three Polintons in a marine species, *Trissonchulus*

latispiculum (Fig 2A and Extended Data Fig 1). A protein domain search using HHpred[57]
identified this domain as a putative HNH endonuclease[58] (Fig 2B-E). An additional structural
alignment between terrestrial *C. briggsae* pPolB2 and marine *T. latispiculum* (*Tlat*) pPolB2
revealed a *Tlat* pPolB2-specific domain, which was subsequently identified as a VSR
endonuclease domain (Fig 2G-J). Apparently, this pPolB2 family acquired an additional
endonuclease domain during its evolution.
Taken together, these observations suggest an ancient separation between pPolB1 and
pPolB2 polymerase families. In the following sections, we investigate the potential origins and
evolution of the two groups of nematode Polintons.
10
Inter- and intra-species copy number variations of pPolB1 and pPolB2 classes of
Polintons in nematode genomes
An extensive search for Polintons in diverse nematode genomes revealed variations in
the inter- and intra-species distributions and copy numbers of the pPolB1 and pPolB2 groups of
Polintons. The pPolB1-class and pPolB2-class Polintons were detected in 25 and 12 nematode
species, respectively (out of a total of 180 species) (Fig 3 and Extended Data Table 1).
Polintons of both classes were detected in several nematode genera (*Pristionchus*, *Oscheius*,
and *Caenorhabditis*), with varying copy numbers; notable copy number variations were also
observed between species within a genus, in particular, *Caenorhabditis*, for which multiple
genomes are available (Fig 3). Generally, the pPolB1-class Polintons were substantially more
abundant than the pPolB2-class ones, with several species carrying only pPolB1 Polintons and
only four possessing pPolB2 Polintons (Fig 3 and Extended Data Table 1). These observations
appear to be compatible with independent spread of the pPolB1 and pPolB2 classes of
Polintons among nematodes.
We also observed distinct patterns and copy numbers of Polintons within each species,
most clearly, in a set of *Caenorhabditis* species for which genome sequences are available for
multiple wild-type strains (Extended Data Fig 3A-D). Because genomes for three wild-type
strains of *C. briggsae* have been assembled at chromosome level[59,60] with the largest copy
numbers of both pPoB1- and pPolB2-class Polintons in nematodes (Fig 3 and Extended Data
Fig 3D), we further analyzed the genomic neighborhoods of the Polintons, to test whether
transposition occurred before or after the diversification of the wild-isolates. Only two pPolB1-
class Polintons shared the same positions in all three *C. briggsae* strains and 6 pPolB1-class
Polintons shared the positions in AF16 and QX1410 strains (Extended Data Fig 3E), suggesting
that these Polintons already occupied the same positions in the genomes of the respective *C.*
briggsae ancestors. In contrast, the remaining pPolB1-class Polintons and all pPolB2-class
Polintons were found at unique positions (Extended Data Fig 3E), suggesting transposition
events that occurred after the strain divergence. These intra-species variations imply that
Polintons were actively integrating into *Caenorhabditis* genomes both before and after the
diversification of the characterized strains, with each genome reflecting a distinct natural history
of exposure to the two classes of Polintons.
41
Transfer of pPolB genes among Polintons
The discovery of two distinct groups of pPolBs in nematode Polintons could be explained
through one of two alternative scenarios. Under the first scenario, there are two distinct classes

of nematode Polintons with only an ancient relationship, in which case, all proteins would show
low sequence conservation, in the same range as the pPolBs. The second scenario involves a
relatively recent pPolB swap, in which case, the other proteins would be far more similar than
the pPolBs.
The bulk protein sequence comparisons described above favor the second scenario, but
to investigate the evolution of the two groups of Polintons further, we examined the relationships
among the Polintons within a single species (*C. briggsae*) by calculating sequence similarities
for all possible pairwise combinations of Polintons. This analysis showed strong positive
correlations between the pairwise similarities of all proteins other than pPolB, regardless of
whether the compared Polintons encoded pPolB1 or pPolB2 (Fig 4A-B and Extended Data Fig
4A-B). In within-class comparisons, strong correlations between the similarities among pPolBs
and those among other Polinton proteins (Fig 4C) suggest that pPolB1 and pPolB2 proteins can
be maintained within Polintons through substantial evolutionary spans. By contrast, in between-
class comparisons, the similarities among pPolB1 and pPolB2 did not correlate with those
among other proteins. (Fig 4C). These observations strongly suggest that in one of the
nematode Polinton classes pPolB was relatively recently replaced by a distantly related
homolog.
We further searched for examples of pPolB1 and pPolB2 genes being embedded in
otherwise closely similar Polinton sequences. One such example, providing a useful illustration,
was observed in *C. briggsae* strain QX1410 (Fig 4D-E), where Polintons Plnt_Cbri_41 and
Plnt_Cbri_35 both showed high similarity to the canonical Plnt_Cbri_43 sequence in all non-
pPolB proteins (Fig 4D), whereas the pPolBs showed dramatic divergence (only 16% sequence
identity) (Fig 4E). A more general view of these relationships is apparent from the phylogenetic
trees of the pPolBs and the rest of the Polinton proteins (Extended Data Fig 5A-H). In the pPolB
tree, the two groups of pPolBs formed two well separated clades whereas the other Polinton
proteins associated with either pPolB1 or pPolB2 were interspersed in the respective trees (Fig
4F, and Extended Data Fig 4C and 5A-H). In addition to *C. briggsae*, we also observed a
possible inter-Polinton exchange of pPolB genes in an *Oscheius* species (DF5120) as indicated
by the observation that an MCP protein from pPolB2-class Polinton clustered with a pPolB1-
class Polinton MCP in the phylogenetic tree (Extended Data Fig 6A-B). These observations are
compatible with the proposed exchange of distinct pPolB genes between Polintons.
32
Distribution of pPolB1 and pPolB2 homologs across eukaryotes suggests distinct
evolutionary trajectories of pPolB1 and pPolB2
To characterize the evolutionary trajectories of the two pPolB types on a larger
evolutionary scale, we explored the distribution of pPolB1 and pPolB2 homologs across the
available eukaryotic genome assemblies (7,897 assemblies from 64 phyla; 1.63 Terabases
available through NCBI genome database as of late April, 2023). To focus on Polintons, we
restricted our analysis to pPolB genes linked to INT homologs (see Methods). We identified
20,702 INT-associated pPolB1 and 1,331 pPolB2 homologs distributed across 960 and 164
genomes of diverse eukaryotes, respectively (Extended Data Table 2 and 3). These pPolBs
found from all taxon groups contained conserved exonuclease and polymerase motifs for
pPolB1 and pPolB2 groups (Extended Data Fig 7), suggesting that these pPolBs are functional
Polinton polymerases.

1   Distinct patterns of pPolB1 and pPolB2 distribution at the phylum level were identified
2 (Fig 5A). Mimicking the case of nematodes, pPolB1-containing Polintons showed a wide spread
3 across 30 phyla. pPolB1 were found to be most abundant in invertebrates, with multiple copies
4 found in 17 of the 23 invertebrate phyla. By contrast, pPolB2 showed a sparse distribution,
5 being represented mostly in two unicellular eukaryote phyla, *Metamonada* and *Oomycota*, and
6 five invertebrate phyla, *Annelida*, *Brachiopoda*, *Echinodermata*, *Mollusca*, and *Nematoda*. The
7 highest pPolB2 copy numbers were detected in *Metamonada*, particularly, in *Trichomonas*
8 *vaginalis* (388 copies) and *Tritrichomonas foetus* (259 copies), consistent with a previous report
9 on the high abundance of Polintons in *T. vaginalis*[3].
10   A broader phylogenetic analysis that included a set of representatives from the main
11 groups of pPolBs encoded by viruses of prokaryotes and eukaryotes as well as Polintons and
12 DNA plasmids further illuminated the evolutionary trajectories of pPolB1 and pPolB2 (Fig 5B).
13 Each of the two groups of pPolBs was found to be monophyletic (Fig 5B and Extended Data Fig
14 8). pPolB1 and pPolB2 formed distinct clades in the tree, and their relative positions suggested
15 that these two DNA polymerase forms diverged early on. Given the greater diversity and
16 broader distribution among the nematodes of pPolB1 compared to pPolB2 (Fig 3 and Extended
17 Data Fig 3A-E), it appears most likely that the ancestral nematode Polintons encoded pPolB1
18 which was replaced in one nematode branch. That replacement might have occurred via
19 horizontal gene transfer from oomycete Polintons, given the tree topology and the highest
20 sequence similarities between oomycete pPolBs and nematode pPolB2s (Fig 5B and Extended
21 Data Fig 8). Furthermore, protein domain search showed that pPolB2 of oomycete Polintons
22 shares the HNH endonuclease domain with the homolog from the nematode Polintons (Fig 6,
23 and Extended Data Fig 9 and 10A), to the exclusion of other pPolB2s. Thus, the HNH domain is
24 a unique synapomorphy (shared derived character) that links the oomycete and nematode
25 Polintons encoding pPolB2, strongly supporting the horizontal gene transfer scenario. The HNH
26 domain could potentially have been captured by the oomycete pPolB2 from a Group I self-
27 splicing intron, before finding its way into the nematode genomes[61].
28   While analyzing protein profiles of oomycete Polintons, we noted that oomycete
29 Polintons encode a conserved protein showing similarity to beta-helix-fold tail fiber proteins of
30 tailed bacteriophages (Fig 6), raising a possibility that these proteins mediate recognition and
31 binding of oomycetes Polintons and oomycetes hosts. Notably, unlike nematode Polintons,
32 oomycete Polintons do not encode membrane fusion proteins and hence are unlikely to be
33 enveloped. Whether potential envelopment is an exclusive property of the nematode Polintons
34 compared to Polintons from other organisms remains to be investigated.
35   Given the apparent relatively recent incursion of pPolB2 into nematode Polintons, direct
36 inter-phyla horizontal transfer of the pPolB2 gene between oomycete and nematode Polintons,
37 perhaps, via virus particles formed by Polintons appears likely. Under this scenario, a nematode
38 would have been infected by the oomycete Polinton virus particles, with subsequent
39 recombination with a resident nematode Polinton such that only the pPolB gene was
40 transferred. Alternatively, the immediate source of the pPolB2 homologs in nematodes might
41 have been an intermediate species. In either case, the pPolB2-encoding Polinton family was
42 fixed in the nematodes and spread to a variety of nematode species, again, possibly, through
43 Polinton virions.
44

A distinct group of pPolB2 Polintons in a marine nematode species

A recent study of marine nematode genomes[62] allowed us to further examine Polintons in an ecological niche for which genome data were initially limited. In addition to land-dwelling nematodes (both free and parasitic), there are many nematode species that dwell in aquatic habitats[28,32,33]. Notably, analysis of the recently sequenced marine nematode genomes identified a Polinton-containing genome (*Trissonchulus latispiculum*[62]) that encompasses no pPolB1-class Polintons but carries a distinct group of three pPolB2-class Polintons (Fig 1A and Extended Data Table 1). Sequence and structural comparison of the terrestrial *C. briggsae* pPolB2 and a marine *T. latispiculum* pPolB2 showed that *T. latispiculum* pPolB2 lacked the HNH nuclease domain but instead contained a VSR (very short patch repair) endonuclease domain (Fig 2F-G and Extended Data Fig 9 and 10B), which was not found in any other nematode pPolB2s (Extended Data Fig 1).

The pPolB2s of the marine nematode Polinton are highly diverged from the pPolB2s of the terrestrial nematode species. Among the known Polintons, the marine nematode pPolB2s showed the highest similarity to pPolB2s of mollusc Polintons as indicated by the phylogenetic tree topology (Fig 5B, Fig 6, and Extended Data Fig 8). Polintons of a mud snail (*B. attramentaria*) and a sea slug (*E. chlorotica*) encode the closest relatives of the marine nematode pPolB2 with the shared VSR endonuclease domain, an apparent synapomorphy of this subgroup of pPolB2 (Fig 6 and Extended Data Fig 8). The presence of the VSR domain along with the phylogenetic tree topology support an additional, independent inter-phylum horizontal gene transfer, in this case between Polintons of nematodes and molluscs Polintons that inhabit the same marine environments.

Conclusions

In this work, we present the unexpected finding that nematode Polintons form two distinct groups that differ by encoding highly diverged pPolBs. The pPolB1 group is by far more widely spread and more abundant among the nematodes than the pPolB2 group suggesting that pPolB2 comparatively recently displaced pPolB1 in two distinct lineages of nematode Polintons at least one of which subsequently spread across nematode species. Phylogenetic analysis of a broad selection of protein-primed B family DNA polymerases suggested that pPolB2s originated from oomycete and mollusc Polintons independently of pPolB1. The DNA polymerase displacement then might have occurred by horizontal transfer of the pPolB2 gene, possibly, via infection of nematodes by oomycete and mollusc Polinton virus particles, with subsequent recombination with a resident nematode Polinton, resulting in the displacement of the DNA polymerase gene alone. The two independent incursions of pPolB2 subfamilies into the terrestrial and marine nematodes, from the oomycete and marine mollusc Polinton families respectively, indicate that the transfers are consistent with the native ecosystems of different groups of nematodes. In this context, we note that terrestrial nematodes can serve as hosts for natural oomycete infection, which could have facilitated the transfer[63,64], whereas marine nematodes most likely shared environments, such as sediment, with prevalent marine molluscs. Although Polinton virions so far have not been observed, the conservation of the major and minor capsid proteins, the packaging ATPase and the maturation protease across the entire diversity of Polintons strongly suggests that such particles exist. Most likely, these virions

mediate the Polinton spread, with the precise characteristics of genomic Polinton elements in a species group reflecting the historical environmental niche rather than the phylogeny. The findings presented in this paper are compatible with this prediction.

Methods

Search of Polintons through *Nematoda* genomes

Assembled genomes of *Nematoda* were downloaded from the NCBI genome assembly database by using the NCBI datasets (https://www.ncbi.nlm.nih.gov/datasets/) command-line tools. The taxon keyword "*Nematoda*" was used to download fasta format genome data. Genome sequences were subjected to a tBLASTn search (with 0.001 as an *e*-value cutoff and 50 as a minimum bitscore). Query sequences (pPolB, INT, MCP, ATPase and Pro) for tBLASTn search were obtained from open reading frame (ORF) prediction by ORFfinder (https://www.ncbi.nlm.nih.gov/orffinder/) on *C. briggsae* Polinton-1 (Polinton-1_CB, WBTransposon00000832)[1] that was downloaded from the WormBase (https://wormbase.org/). We filtered out candidate genetic blocks that had all five query proteins within 20 kb and gathered all of these segments with upstream and downstream buffers of 10 kb. Next, the candidates were further filtered out by the presence of terminal inverted repeats (TIR) that were detected using Inverted Repeat Finder (https://tandem-test.bu.edu/irf/irf.download.html)[65]. The 6 bases upstream and downstream of each candidate TIR were used to test whether a target site duplication (TSD) was present. Polinton candidate sequences were further validated by the presence of at least 5 Polinton proteins through ORF prediction on candidate sequences and protein-protein BLAST (BLASTp, bitscore cut-off: 100) of the ORFs against query Polinton proteins. A total of 266 Polintons from 66 nematode genomes (29 species) were identified and summarized in Extended Data Table 1.

Multiple sequence alignment and phylogenetic tree analysis

Multiple sequence alignment for nematode pPolB proteins, *C. briggsae* Polinton proteins, and pPolB proteins found from each taxon group was performed using ClustalO[66,67]. Phylogenetic trees were generated with Maximum likelihood (ML) method by using IQ-TREE[68-70], and mid-point rooted and visualized by ETE3 Python package[71]. Percent identity between protein sequences were calculated based on the pairwise comparison of sequences within the ClustalX environment[72]. Conserved motifs were searched from the multiple sequence alignment results and enrichment of amino acids were visualized by using LogoMaker Python package[73].

For phylogenetic analyses of pPolBs from multiple taxa, pPolB protein sequences in each taxon group were clustered and a representative sequence was selected for each cluster by using CD-HIT[74]. Then, pPolBs were subjected to multiple sequence alignment by using ClustalO[66,67]. Gap regions in the alignment data were trimmed off with trimAl gappyout option[75] and then phylogenetic trees were reconstructed by using ML method inferred by IQ-TREE v2.0.3 with ultrafast bootstrapping (1000 iterations)[68-70] and with LG+F+R10 model as the best fitting model. The nodes of the phylogenetic tree were collapsed if all the pPolB leaves of the node were from same taxon groups. The phylogenetic trees were mid-point rooted and visualized by using ETE3 Python package[71].

For the more global phylogenetic analysis, a dataset of pPolB sequences was retrieved

from Krupovic and Koonin (2014)[10]. The dataset included pPolB sequences from bacterial

tectiviruses, linear mitochondrial and cytoplasmic plasmids, adenoviruses and a limited number

of bidnaviruses and Polintons. This dataset was supplemented with a greatly expanded set of

pPolB sequences of Polintons, bidnaviruses and adintoviruses[25]. The dataset was then

clustered to 50% identity over 80% of alignment length using MMseqs2[76,77]. The sequences

were aligned using MAFFT with the G-INS-1 option[78]. Low-information content positions were

removed using trimAl with the gappyout option[75]. The maximum likelihood phylogenetic tree was

calculated using IQ-TREE v2.0.6[68] with the best fitting substitution model determined for the

data, which was LG+F+R10.

Protein structure prediction and alignment

Structure prediction steps of *C. briggsae* pPolB1 or pPolB2 proteins were executed

using MMSeqs2[76,77] and AlphaFold2[55] on ColabFold v1.5.2[56]

(https://colab.research.google.com/github/sokrypton/ColabFold/blob/main/AlphaFold2.ipynb).

Pairwise structural alignment of predicted pPolB1 and pPolB2 structures was performed with

FATCAT[79,80] (https://www.rcsb.org/alignment) with default parameters. The structure was

visualized using the PyMOL Molecular Graphics System, Version 2.5 Schrödinger, LLC.

Polinton pPolB search through NCBI genome assemblies

We downloaded 7,897 genomes from 64 phyla of eukaryotes (except Viridiplantae or

Chordata) available through the NCBI genome database as of late 2023 April (assembly source:

genbank). To explore the distribution of nematode pPolB1 and pPolB2 homologs in the

genomes, each fasta format genome file was indexed for tBLASTn search, and Polinton pPolB1

and pPolB2 protein sequences were searched through the genome using tBLASTn. To provide

a stringent search, we filtered out small matches by setting a match length cutoff (650 amino

acids for both pPolB1 and pPolB2). Then, the matched sequences with 20 kb both upstream

and downstream were used to find potential ORFs for pPolB and other Polinton proteins. We

further filtered candidates based on the length of predicted pPolB ORFs (a minimum ORF

length: 900 amino acids). Subsequently, hmmsearch[81] (http://hmmer.org/) was performed to

detect homology between predicted ORFs and Polinton proteins. Profiles for the Polinton

proteins (INT, MCP, and ATPase) were built from the multiple sequence alignments of each

Polinton protein obtained from RepBase Polintons (https://www.girinst.org/repbase)[82,83]. To

ensure that pPolB gene candidates are indeed Polinton polymerases, we selected pPolBs that

co-detected with an INT protein nearby (see Extended Data Table 3 for pPolB-Polinton

information). For each taxon group, copy numbers of pPolB1 and pPolB2 were normalized by

the number of genomes that had at least a single copy of pPolB and visualized by using Python

ETE3 package[71].

Fig 1

A

B

Fig 1. **Two distinct DNA polymerase B families in nematode Polintons.**
A. A phylogenetic tree (left) was constructed by applying a maximum likelihood-based method
(IQ-TREE) to a multiple sequence alignment of pPolB proteins from Polintons of 19 nematode
species. The genetic architectures of these elements are shown on the right. These Polintons
were chosen for initial sequence alignment based on uninterrupted DNA polymerase and other
Polinton protein coding regions. Navy and magenta colors of tree nodes indicate pPolB-1- and
pPolB2-class Polintons, respectively. Bootstrap supporting values are shown on the top position
of tree branches. Colors in right panels denote conserved genes encoding Polinton proteins;
orange for an adenovirus-like maturation protease (PRO), navy and magenta for protein-primed
DNA polymerase B genes (pPolB1 and pPolB2, respectively), beige for Polinton-
uncharacterized ORF (puORF), purple for a packaging ATPase, lawngreen and cyan for major
and minor capsid proteins (MCP and mCP, respectively), dark red for a fusogen, green for a
retorviral-like-element integrase (INT), and white for novel (potentially either spurious or
functional) ORFs that have not been previously described in other Polintons. Labels in
parentheses denote unique identifiers (UI) for each Polinton (see Extended Data Table 1).
B. A dot plot and a background box plot show percent (%) amino acid identities between each
Polinton protein found in 49 *C. briggsae* Polintons.

Fig 2

A

pLDDT (pPolB1)

pPolB1

pPolB2

pLDDT (pPolB2)

HNH Endo

Aligned
Non-aligned
Gap

B *C. briggsae* pPolB2

C *C. briggsae* pPolB2 HNH domain

D *P. alcaligenes* PacI

E Superimposed

F

pLDDT for pPolB2 (*C. briggsae*)

C. briggsae pPolB2 — HNH Endo

T. latispiculum pPolB2 — VSR Endo

pLDDT for pPolB2 (*T. latispiculum*)

Aligned
Non-aligned
Gap

G *T. latispiculum* pPolB2

H *T. latispiculum* pPolB2 VSR domain

I *E. coli* VSR protein

J Superimposed

Fig 2. HNH and VSR endonuclease domains are found in terrestrial and marine nematode pPolB2s, respectively, but not in pPolB1.

A. Aligned (blue), non-aligned (skyblue) and gap (white) parts of pPolB1 and pPolB2 (middle panel) from sequence-free structural alignment are shown with pLDDT (predicted local distance difference test) score predicted at the single amino acid level by AlphaFold2. Overall aligned parts for both pPolB1 and pPolB2 showed high pLDDT score, suggesting confident prediction of the structures. **B.** The predicted structure of *C. briggsae* pPolB2 (green) with the HNH endonuclease domain (red). **C-E.** The predicted structure of pPolB2-specific HNH domain (blue, **C**) and the crystal structure of *P. alcaligenes* PacI HNH endonuclease (orange, **D**) (PDBid: 3M7K)[58] are superimposed (**E**) (RMSD: 3.03, TM-score: 0.55) to show potentially conserved structural folds. **F.** Aligned (blue) and non-aligned (skyblue) parts of *C. briggsae* pPolB2 and *T. latispiculum* pPolB2 (middle panel) from sequence-free structural alignment are shown with pLDDT score. **G.** The predicted of *T. latispiculum* pPolB2 (cyan) with the VSR domain (yellow). **H-J.** The predicted structure of *T. latispiculum* pPolB2 VSR endonuclease domain (blue, **H**) and the crystal structure of *E. coli* VSR endonuclease (orange, **I**) (PDBid: 1VSR)[84] are superimposed (**J**) (RMSD: 3.06, TM-score: 0.63) to show potentially conserved structural folds. Dark and light colors for both orange and blue (**C-E** and **H-J**) indicate aligned and non-aligned positions, respectively.

Fig 3

1   **Fig 3. Inter-species copy number variations of nematode Polintons.**

2   Bar graphs (right panel) showing average copy numbers per genome of full-length pPolB1-class

3   (navy) and pPolB2-class (magenta) Polintons in *Nematoda* species are shown with

4   a nematode species phylogenetic tree (left panel) that was obtained from Ahmed et al. (2022)[85].

Fig 4

A

mCP identity (%) vs INT identity (%)

● Within-class (pPolB1-class vs pPolB1-class) or (pPolB2-class vs pPolB2-class)
● Between-class (pPolB1-class vs pPolB2-class)

$r^2 = 0.9$
$r^2 = 0.86$

B

puORF identity (%) vs ATPase identity (%)

$r^2 = 0.9$
$r^2 = 0.8$

C

pPolB1/2 identity (%) vs ATPase identity (%)

$r^2 = 0.82$
$r^2 = 0.19$

D

Amino acid identity (%) to query Polinton proteins

pPolB1-class
pPolB2-class

type, PRO, pPolB1/2, puORF, ATPase, mCP, Fusogen, MCP, INT

E

C. briggsae (QX1410) Polintons

pPolB1-class (Plnt_Cbri_38)
68 78 82 81 68 83 94 78

pPolB1-class (Plnt_Cbri_43)
79 90 97 96 95 95 98 98

pPolB1-class (Plnt_Cbri_41)
87 16 95 95 94 95 99 92

pPolB2-class (Plnt_Cbri_35)

PRO
pPolB1
pPolB2
puORF
ATPase
mCP
Fusogen
MCP
INT
TIR

F

pPolB — Fusogen

● pPolB1-class Polinton
● pPolB2-class Polinton

1.44891

0.301992

1	**Fig 4**. **Two distantly related groups of pPolBs in *C. briggsae* Polintons.**
2	**A-C**. Scatter plots showing percent amino acid identity of one protein for each axis from *C.*
3	*briggsae* Polinton species; mCP and INT (**A**), puORF and ATPase (**B**), and pPolB1/2 and
4	ATPase (**C**). Each dot indicates a pair of two *C. briggsae* Polintons, whose proteins indicated at
5	the X and Y axes were aligned, respectively, for calculating percent identity. Dot colors indicate
6	whether two Polintons in a pair for comparison are within the same pPolB group (blue) or from
7	different pPolB groups (orange). **D.** Heatmap showing percentage amino acid identity of
8	proteins. Each row shows an individual *C. briggsae* Polinton (navy for pPolB1-class and
9	magenta for pPolB2-class), while each column represents one of the conserved Polinton
10	proteins. All comparisons were performed against a query *C. briggsae* Polinton (Plnt_Cbri_18).
11	**E.** Genetic architectures of four *C. briggsae* Polintons (QX1410 strain, a wild isolate) shown
12	with % amino acid identities for each Polinton protein. Plnt_Cbri_41 and Plnt_Cbri_43 are
13	relatively close to each other, showing > 90% identity, while Plnt_Cbri_43 and Plnt_Cbri_38 are
14	more distant (evidenced by lower % identities). The continuous nature of the homology outside
15	of the pPolB region provides evidence for gradual divergence in these regions during the
16	ancestry of the current elements. By contrast, the Plnt_Cbri_35 Polinton showed strong
17	similarity in all proteins except pPolB (particularly to PInt_Cbri_41), with dramatic divergence for
18	pPolB, suggesting a swapping event of the pPolB gene during the derivation of this Polinton. **F.**
19	A tanglegram showing the relationship between two phylogenetic trees constructed from
20	multiple sequence alignment of *C. briggsae* pPolB1/2 (left panel) and Fusogen (right panel)
21	proteins. The colors of the nodes indicate pPolB1-class (navy) and pPolB2-class (magenta).
22	Dashed lines between two trees denote the correspondence of pPolB and Fusogen proteins
23	found from the same Polinton. The tree exemplifies that close relationships between non-pPolB
24	proteins can be accompanied by very distant relationships between the corresponding pPolBs.

Fig 5

A

Fig 5. **Distinct distribution and evolutionary trajectories of pPolB1 and pPolB2 groups in**
the biosphere.
A. A NCBI taxonomy tree with Eukaryotic taxa. The numbers in parentheses indicate the
number of genomes having at least one copy of pPolB and the number of total genomes
subjected to the Polinton pPolB search (see Methods for details). The sizes and colors in the pie
charts indicate the copy number of Polinton pPolBs per pPolB-containing genomes and the
classes of pPolB; navy for pPolB1 and magenta for pPolB2, respectively. **B.** A phylogenetic tree
was constructed from multiple sequence alignment of pPolB proteins found in diverse taxa
including viruses of prokaryotes, eukaryotes, DNA plasmids, and Polintons using the IQ-TREE-
inferred maximum likelihood method. Bootstrap supporting values are shown on the bottom
positions of tree branches. All pPolBs identified in this study were found within Group 2 Polinton
pPolBs where two big monophyletic groups for pPolB1 and pPolB2 classes are shown with navy
and magenta branch colors, respectively. Red asterisks indicated the *Nematoda* pPolB1 and
pPolB2 proteins identified in this study.

Fig 6

HNH endo in pPolB

VSR endo In pPolB

pPolB2-class Polinton *Oscheius sp.(DF5120)* (Plnt_Osp._1)

pPolB2-class Polinton *Auanema rhodensis* (Plnt_Arho_1)

pPolB2-class Polinton *Caenorhabditis remanei* (Plnt_Crem_7)

pPolB2-class Polinton *Caenorhabditis tropicalis* (Plnt_Ctro_6)

pPolB2-class Polinton *Caenorhabditis briggsae* (Plnt_Cbri_57)

pPolB2-class Polinton *Caenorhabditis latens* (Plnt_Clat_4)

Nematoda (Terrestrial)

pPolB2-class Polinton *Pythium brassicum* (Plnt_Pbra_2)

pPolB2-class Polinton *Pythium guiyangense* (Plnt_Pgui_3)

pPolB2-class Polinton *Phytophthora capsici* (Plnt_Pcap_1)

pPolB2-class Polinton *Phytophthora palmivora* (Plnt_Ppal_2)

pPolB2-class Polinton *Phytophthora sojae* (Plnt_Psoj_3)

pPolB2-class Polinton *Phytophthora cactorum* (Plnt_Pcac_26)

pPolB2-class Polinton *Phytophthora vignae* (Plnt_Pvig_7)

pPolB2-class Polinton *Phytophthora tubulina* (Plnt_Ptub_2)

Oomycota

pPolB2-class Polinton *Batillaria attramentaria* (Plnt_Batt_12)

pPolB2-class Polinton *Elysia chlorotica* (Plnt_Echl_1)

Mollusca

pPolB2-class Polinton *Trissonchulus latispiculum* (Plnt_Tlat_2)

Nematoda (Marine)

pPolB2-class Polinton *Haliotis rufescens* (Plnt_Hruf_3)

pPolB2-class Polinton *Tegillarca granosa* (Plnt_Tgra_2)

pPolB2-class Polinton *Crassostrea virginica* (Plnt_Cvir_1)

pPolB2-class Polinton *Pecten maximus* (Plnt_Pmax_3)

Mollusca

PRO
pPolB1
pPolB2
puORF
ATPase
mCP
Fusogen
Potential fiber protein
MCP
INT
Endonuclease
TFIIB
TIR
Uncharacterized

Fig 6. Phylogenetic relationships of Polinton pPolB2 proteins from three distinct phyla.
A Phylogenetic tree (left) has been built by using IQ-TREE-inferred maximum likelihood on a
multiple sequence alignment of pPolB2 protein sequences found from *Nematoda*, *Oomycota*
and *Mollusca* (See Extended Data Fig 9 for full alignment data.). Bootstrap supporting values
are shown on the top position of tree branches. Colors in right panels denote conserved genes
encoding Polinton proteins; orange for an adenovirus-like maturation protease (PRO), navy and
magenta for protein-primed DNA polymerase B genes (pPolB1 and pPolB2, respectively), beige
for Polinton-uncharacterized ORF (puORF), purple for a packaging ATPase, lawngreen and
cyan for major and minor capsid proteins (MCP and mCP, respectively), dark red for a fusogen,
green for a retorviral-like-element integrase (INT), olive for an endonuclease, ivory for potential
fiber protein, black for a potential transcription initiation factor (TFIIB) and white for novel
(potentially either spurious or functional) ORFs.

Extended Data Fig 1

Extended Data Fig 1 *(continued)*

Extended Data Fig 2

A

Motifs	Exo_I	Exo_II	Exo_III	Pol_A	Pol_B	Pol_C

		* *	*	*	*	* *
pPolB	*Phi29*	DFET	HNLK-FD	YIKNDIQ	DVNSLYPAQM	KLMLNSLYGKFASN	YCDTDS
pPolB1	*Heterorhabditis bacteriophora*	DFES	HNLKGYD	YMINDVL	DANNLYGWAM	KLMNNSVFGKTMEN	YQDTDS
pPolB1	*Micoletzkya japonica*	DFED	HNLKGYD	YMINDVL	DANNLYGWAM	KLMNNSVFGKTMEN	YQDTDS
pPolB1	*Pristionchus pacificus*	DFES	HNLKGYD	YMSNDVL	DANKLYGWAM	KLMNNSVFGKTMEN	YQDTDS
pPolB1	*Oscheius sp. (DF5120)*	DFES	HNLRGYD	YMINDVL	DANNLYGWAM	KLMNNSVFGKTMEN	YQDTDS
pPolB1	*Caenorhabditis elegans*	DSES	HNLKGYD	YMINDVL	DANNLYGWAM	KLMNNSVFGKTMEN	YQDTDS
pPolB1	*Caenorhabditis remanei*	DFES	HNLKGYD	YMINDVL	DANNLYGWAM	KLMNNSVFGKTMEN	YQDTDS
pPolB1	*Caenorhabditis tropicalis*	DFES	HNLKGYD	YMINDVL	DANNLYGWAM	KLMNNSVFGKTMEN	YQDTDS
pPolB1	*Caenorhabditis briggsae*	DFES	HNLKGYD	YMINDVL	DANNLYGWAM	KLMNNSVFGKTMEN	YQDTDS
pPolB1	*Caenorhabditis latens*	DFES	HNLKGYD	YMINDVL	DANNLYGWAM	KLMNNSVFGKTMEN	YQDTDS
pPolB1	*Halicephalobus mephisto*	DFES	HNLKGYD	YMINDVL	DANNLYGWSM	KLMNNAVFGKTMEN	YQDTDS
pPolB1	*Halicephalobus sp. (NKZ332)*	DFES	HNLKGYD	YMINDVL	DANNLYGWAM	KLMNNAVFGKTMEN	YQDTDS
pPolB1	*Diploscapter coronatus*	DFEA	HNLTSYD	YMKTDVL	DVNSLYPTAM	KLAMNIVFGKTIEN	YMDTDS
pPolB1	*Bursaphelenchus mucronatus*	DFEA	HNLTSYD	YVKTDVL	DVNSLYPTAM	KLAMNIVFGKTIEN	YMDTDS
pPolB2	*Trissonchulus latispiculum*	DFEC	FNSGKFD	YNNRDVE	DGNALYLWCL	KLLGNSYYGRTITN	IQMDTDS
pPolB2	*Oscheius sp. (JU1382)*	DFEV	FNSGGYD	YNNLDVR	DANALYLWCL	KLVGNSAFGVTGMD	EMDTDS
pPolB2	*Auanema rhodensis*	DFEA	FNSGGYD	YNNLDVR	DANALYLWCL	KLVGNSAFGVTGMD	EMDTDS
pPolB2	*Caenorhabditis remanei*	DMES	FNSSFYD	YNILDVK	DANALYLHSL	KLVGNSAFGRTGMD	EMDTDS
pPolB2	*Caenorhabditis tropicalis*	DMES	FNSSFYD	YNILDVR	DANALYLHSL	KLVGNSAFGRTGMD	EMDTDS
pPolB2	*Caenorhabditis briggsae*	DMES	FNSSFYD	YNILDVR	DANALYLHSL	KLVGNSAFGRTGMD	EMDTDS
pPolB2	*Caenorhabditis latens*	DMES	FNSSFYD	YNILDVK	DANALYLHSL	KLVGKLAFGRTGMD	EMDTDS

B *C. briggsae* pPolB1

C *C. briggsae* pPolB2

D Superimposed

Extended Data Fig 2. **pPolB1 and pPolB2 proteins have closely similar structures with**
conserved catalytic motifs.
A. Multiple sequence alignment (right panels) of pPolB1 and pPolB2 proteins from 19 nematode
species. The phylogenetic tree (left panels) is equivalent to that in Fig 1A with the minor addition
of the outgroup Phi29 phage DNA polymerase). Six conserved pPolB protein motifs (Exo_I,
Exo_II, Pol_A, Pol_B, and Pol_C) are selectively shown (See Extended Data Fig 1 for a full
alignment result). **B-D**. show AlphaFold2 predictions of *C. briggsae* pPolB1 (orange, **B**), pPolB2
(blue, **C**) and a superimposed structure (RMSD: 4.22, TM-score: 0.53) (**D**) of pPolB1 and
pPolB2 constructed by structural alignment. Dark and light colors indicate aligned and non-
aligned positions within these putative structures, respectively.

Extended Data Fig 3

A *Caenorhabditis elegans*

Legend: pPolB1-class (navy), pPolB2-class (magenta)

Rows (top to bottom): JU1400, MY2147, MY2693, N2, NIC526, DL226, JU1395, EG4725, JU2600, NIC2, CB4856, DL238, ECA36, ECA396, XZ1516, JU2526, QX1794, JU310

X-axis: Copy numbers in genome (0–16)

B *Caenorhabditis remanei*

Rows: PX439, EM464, PB4641, PX356, PX506

C *Caenorhabditis tropicalis*

Rows: JU1373, NIC203

D *Caenorhabditis briggsae*

Rows: QX1410, AF16, VX34

X-axis: Copy numbers in genome (0–16)

E

Legend: pPolB1-class (navy), pPolB2-class (magenta); Shared (square), Unique (triangle)

Rows: VX34, AF16, QX1410 — ChrI, ChrII, ChrIII

Rows: VX34, AF16, QX1410 — ChrIV, ChrV, ChrX

Extended Data Fig 3. intra-species copy number and position variations of *Caenorhabditis* **Polintons.**

A-D. Bar graphs showing average copy numbers per genome of pPolB1-class (navy) and pPolB2-class (magenta) full-length Polintons in wild-type strains of *C. elegans* (A), *C. remanei* (**B**), *C. tropicalis* (**C**) and *C. briggsae* (**D**). **E.** Insertion positions of pPolB1- (navy) and pPolB2-class (magenta) Polintons from three *C. briggsae* strains (VX34, AF16, and QX1410) at the chromosome scale. Dot shapes indicate whether the positions were found as unique (inversed triangle) or shared (square) in at least two strains.

Extended Data Fig 4

A

● Within-class (pPolB1-class vs pPolB1-class) or (pPolB2-class vs pPolB2-class)
● Between-class (pPolB1-class vs pPolB2-class)

$r^2 = 0.82$
$r^2 = 0.69$

B

$r^2 = 0.71$
$r^2 = 0.7$

C

● pPolB1-class Polinton
● pPolB2-class Polinton

pPolB

	MCP

Plnt_Cbri_59
Plnt_Cbri_49
Plnt_Cbri_55
Plnt_Cbri_28
Plnt_Cbri_42
Plnt_Cbri_27
Plnt_Cbri_37
Plnt_Cbri_18
Plnt_Cbri_16
Plnt_Cbri_14
Plnt_Cbri_38
Plnt_Cbri_29
Plnt_Cbri_51
Plnt_Cbri_54
Plnt_Cbri_23
Plnt_Cbri_26
Plnt_Cbri_39
Plnt_Cbri_45
Plnt_Cbri_30
Plnt_Cbri_43
Plnt_Cbri_13
Plnt_Cbri_33
Plnt_Cbri_41
Plnt_Cbri_24
Plnt_Cbri_56
Plnt_Cbri_47
Plnt_Cbri_57
Plnt_Cbri_50
Plnt_Cbri_52
Plnt_Cbri_17
Plnt_Cbri_35
Plnt_Cbri_58
Plnt_Cbri_22
Plnt_Cbri_25

Plnt_Cbri_59
Plnt_Cbri_49
Plnt_Cbri_55
Plnt_Cbri_37
Plnt_Cbri_28
Plnt_Cbri_42
Plnt_Cbri_27
Plnt_Cbri_16
Plnt_Cbri_14
Plnt_Cbri_38
Plnt_Cbri_29
Plnt_Cbri_18
Plnt_Cbri_51
Plnt_Cbri_54
Plnt_Cbri_23
Plnt_Cbri_47
Plnt_Cbri_57
Plnt_Cbri_30
Plnt_Cbri_43
Plnt_Cbri_26
Plnt_Cbri_39
Plnt_Cbri_56
Plnt_Cbri_45
Plnt_Cbri_50
Plnt_Cbri_13
Plnt_Cbri_33
Plnt_Cbri_41
Plnt_Cbri_24
Plnt_Cbri_17
Plnt_Cbri_22
Plnt_Cbri_25
Plnt_Cbri_52
Plnt_Cbri_35
Plnt_Cbri_58

1.43364

0.0241324

Extended Data Fig 4. **Polinton genes diverge together.**

A-B. Scatter plots showing percent (%) amino acid identity of a protein at each axis from *C.*

briggsae Polinton species; ATPase and PRO (**A**), INT and MCP (**B**). Each dot indicates a pair of

two *C. briggsae* Polintons, whose proteins indicated at the X and Y axes were aligned,

respectively, for calculating % identity. Dot colors indicate whether two Polintons in a pair for

comparison are within the same groups of Polintons (within pPolB-class, blue) or from different

groups (between pPolB-class, orange). **C.** A tanglegram showing the relationship between two

phylogenetic trees constructed from multiple sequence alignment of *C. briggsae* pPolB1/2 (left

panel) and MCP (right panel) proteins. The colors of the nodes indicate pPolB1-class (navy) and

pPolB2-class (magenta). Dashed lines between two trees denote the correspondence of pPolB

and MCP proteins found from the same Polinton.

Extended Data Fig 5

Extended Data Fig 5. **Phylogenetic analysis suggests acquisition of two classes of pPolB**
in *C. briggsae* Polintons during evolution.
A-H. Phylogenetic trees were constructed by maximum likelihood method on multiple sequence
alignment of pPolB (**A**), Fusogen (**B**), ATPase (**C**), mCP (**D**), puORF (**E**), INT (**F**), MCP (**G**) and
PRO (**H**) proteins found in *C. briggsae* Polintons. Navy and magenta colors of tree nodes
indicates pPolB1- and pPolB2-class Polintons, respectively, where the proteins are encoded.
The numbers on each branch indicates bootstrap supporting values.

Extended Data Fig 6

A

B

pPolB1-class Polinton
pPolB2-class Polinton

A

Nematode

- *Bursaphelenchus mucronatus*
- *Halicephalobus mephisto*
- *Diploscapter coronatus*
- *Diploscapter pachys*
- *Caenorhabditis elegans*
- *Caenorhabditis tropicalis*
- *Caenorhabditis remanei*
- *Caenorhabditis latens*
- *Caenorhabditis briggsae*
- *Heterorhabditis bacteriophora*
- *Oscheius sp.DF5120*
- *Auanema rhodensis*
- *Pristionchus entomophagus*
- *Pristionchus mayeri*
- *Pristionchus pacificus*
- *Pristionchus fissidentatus*
- *Micoletzkya japonica*
- *Trissonchulus latispiculum*

pPolB

B

Nematode

- *Caenorhabditis elegans*
- *Caenorhabditis tropicalis*
- *Caenorhabditis briggsae*
- *Caenorhabditis remanei*
- *Caenorhabditis latens*
- *Diploscapter pachys*
- *Diploscapter coronatus*
- *Heterorhabditis bacteriophora*
- *Oscheius sp.DF5120*
- *Oscheius sp.JU1382*
- *Auanema rhodensis*
- *Allodiplogaster sudhausi*
- *Micoletzkya japonica*
- *Pristionchus pacificus*
- *Pristionchus mayeri*
- *Pristionchus entomophagus*
- *Pristionchus fissidentatus*
- *Bursaphelenchus mucronatus*
- *Halicephalobus mephisto*
- *Trissonchulus latispiculum*

MCP

 Extended Data Fig 6. Evidence for horizontal gene transfer mediated by Polintons across

nematode species.

A-B. Tanglegrams that show the link between phylogenetic trees of nematode species (left) and

Polinton pPolB (**A**) and MCP (**B**) proteins. The colors of tree nodes indicate pPolB1-class (navy)

and pPolB2-class (magenta). Dashed lines between two trees denote the correspondence of

pPolB (**A**) or MCP (**B**) proteins found from the nematode species (navy and magenta for the

links from species to pPolB1- and pPolB2-class Polintons, respectively). Subsets of closely

related PolBs and/or MCPs were found across distant nematode species, suggesting either that

these Polintons originate from a common Polinton ancestor or that horizontal gene transfer

event occurred across different nematode species.

Extended Data Fig 7

Extended Data Fig 7. Polinton pPolBs have conserved exonuclease and polymerase
domains.
Conserved exonuclease (Exo_I, Exo_II, and Exo_III) and polymerase (Pol_A, Pol_B, and
Pol_C) motifs for pPolBs found from each taxon groups were searched through multiple
sequence alignments and are shown with graphical representation of the conserved amino
acids.

Extended Data Fig 8

Extended Data Fig 8. Monophyletic groups of Polinton pPolB1 and pPolB2 families.
A phylogenetic tree of representative pPolB1 and pPolB2 proteins from multiple taxon groups
was built by using IQ-TREE-inferred maximum likelihood method on multiple sequence
alignment of the proteins (see Method for details). Bootstrap supporting values for each branch
were shown on the top position of the branches. Navy and magenta colors indicate pPolB1 and
pPolB2, respectively. The numbers in parentheses denote the number of pPolB proteins in each
node of taxon group used for the multiple sequence alignment. Nematode pPolB2 groups found
from the terrestrial and marine species are marked with red arrows.

Extended Data Fig 9

Extended Data Fig 9. Terrestrial nematode and oomycete, and marine nematode and
mollusc pPolB2 proteins share unique synapomorphies, an HNH and a VSR domain,
respectively.
Full-length multiple sequence alignment data of representative pPolB2 proteins from three
distinct phyla; *Nematoda*, *Oomycota*, and *Mollusca*. Red boxes indicate conserved catalytic
motifs for exonuclease (Exo_I, Exo_II, Exo_III) and polymerase (Pol_A, Pol_B, Pol_C) activity.
The conserved HNH endonuclease domain marked with a blue box is found in terrestrial
nematode and oomycete pPolB2s but not in mollusc pPolB2s. In contrast, The VSR
endonuclease domain marked with a purple box was conserved in a marine nematode pPolB2
(nema_Tlat_2) and mollusc pPolB2 proteins. Labels at the most left column show pPolB2
unique identifier that includes phylum and species information; moll for *Mollusca*, nema fo*r*
Nematoda, oomy for *Oomycota*, Cvir for *Crassorstrea virginica*, Batt for *Batillaria attramentaria*,
Hruf for *Haliotis rufescens*, Echl for *Elysia chlorotica*, Pmax for *Pecten maximus*, Tgra for
Tegillarca granosa, Tlat for *Trissonchulus latispiculum*, Osp. for *Oscheius* species, Arho for
Auanema rhodensis, Crem for *Caenorhabditis remanei*, Clat for *Caenorhabditis latens*, Ctro for
Caenorhabditis tropicalis, Cbri for *Caenorhabditis briggsae*, Aeut for Aphanomyces euteiches,
Lgig for Lottia gigantea, Pcac for *Phytophthora cactorum*, Pcap for *Phytophthora capsici*, Pfra
for *Phytophthora fragariae*, Plat for *Phytophthora lateralis*, Ppal for *Phytophthora palmivora*,
Psoj for *Phytophthora sojae*, Ptub for *Phytophthora tubulina*, Pvig for *Phytophthora vignae*, Pbra
for *Pythium brassicum*, Pgui for *Pythium guiyangense*.

Extended Data Fig 10

A

HNH Endonuclease domain in pPolB2

B

VSR Endonuclease domain in pPolB2

Extended Data Fig 10. HNH and VSR domains of pPolB2s appear functional.
A. A phylogenetic tree (left) and multiple sequence alignment (right) of HNH endonuclease
domains found in Polinton pPolB2 proteins of nematode and oomycete species with PacI
bacterial HNH endonuclease found in *Pseudomonas alcaligenes*. Red and blue asterisks
indicate zinc-binding Cys residues and catalytic residues for PacI[58], both of which are conserved
across nematode and oomycete pPolB2 proteins except a Tyr residue. **B.** A phylogenetic tree
(left) and multiple sequence alignment (right) of VSR endonuclease domains found in Polinton
pPolB2 proteins of marine nematode and mollusc species with *E. coli* VSR protein. Blue
asterisks indicate catalytically important residues for *E. coli* VSR endonuclease activity[86].

Acknowledgements

We thank Karen Lynn Artiles, Lamia Wahba, Massa Shoura, Matthew James McCoy, Orkan Ilbay, Jingxun Chen, Usman Enam, Emily Greenwald, Ivan Nikolay Zheludev, Drew Galls, and Janie Soo-hyun Kim for helpful discussion and their critical reading on the manuscript. This study was supported by the NIGMS grant (R35GM130366 to A.Z.F.), and by the Long-term Postdoctoral Fellowship from the Human Frontier Science Program (LT000329/2019-L) and the Bernard Cohen Postdoctoral Fellowship (to D-E.J.). S.S. is supported by the Stanford Cardiovascular Institute Summer Undergraduate Fellowship. E.V.K. is supported by the Intramural Research Program of the National Institutes of Health of the USA (National Library of Medicine).

References

Kapitonov, V. V. & Jurka, J. Self-synthesizing DNA transposons in eukaryotes. *Proc Natl Acad Sci U S A* **103**, 4540-4545, doi:10.1073/pnas.0600833103 (2006).

Pritham, E. J., Putliwala, T. & Feschotte, C. Mavericks, a novel class of giant transposable elements widespread in eukaryotes and related to DNA viruses. *Gene* **390**, 3-17, doi:10.1016/j.gene.2006.08.008 (2007).

Feschotte, C. & Pritham, E. J. DNA transposons and the evolution of eukaryotic genomes. *Annu Rev Genet* **41**, 331-368, doi:10.1146/annurev.genet.40.110405.090448 (2007).

Jurka, J., Kapitonov, V. V., Kohany, O. & Jurka, M. V. Repetitive sequences in complex genomes: structure and evolution. *Annu Rev Genomics Hum Genet* **8**, 241-259, doi:10.1146/annurev.genom.8.080706.092416 (2007).

Krupovic, M. & Koonin, E. V. Polintons: a hotbed of eukaryotic virus, transposon and plasmid evolution. *Nat Rev Microbiol* **13**, 105-115, doi:10.1038/nrmicro3389 (2015).

Feschotte, C. & Pritham, E. J. Non-mammalian c-integrases are encoded by giant transposable elements. *Trends Genet* **21**, 551-552, doi:10.1016/j.tig.2005.07.007 (2005).

Krupovic, M., Bamford, D. H. & Koonin, E. V. Conservation of major and minor jelly-roll capsid proteins in Polinton (Maverick) transposons suggests that they are bona fide viruses. *Biol Direct* **9**, 6, doi:10.1186/1745-6150-9-6 (2014).

Gao, X. & Voytas, D. F. A eukaryotic gene family related to retroelement integrases. *Trends Genet* **21**, 133-137, doi:10.1016/j.tig.2005.01.006 (2005).

Krupovic, M. & Koonin, E. V. Self-synthesizing transposons: unexpected key players in the evolution of viruses and defense systems. *Curr Opin Microbiol* **31**, 25-33, doi:10.1016/j.mib.2016.01.006 (2016).

Krupovic, M. & Koonin, E. V. Evolution of eukaryotic single-stranded DNA viruses of the Bidnaviridae family from genes of four other groups of widely different viruses. *Sci Rep* **4**, 5347, doi:10.1038/srep05347 (2014).

Koonin, E. V., Krupovic, M. & Yutin, N. Evolution of double-stranded DNA viruses of eukaryotes: from bacteriophages to transposons to giant viruses. *Ann N Y Acad Sci* **1341**, 10-24, doi:10.1111/nyas.12728 (2015).

Haapa-Paananen, S., Wahlberg, N. & Savilahti, H. Phylogenetic analysis of Maverick/Polinton giant transposons across organisms. *Mol Phylogenet Evol* **78**, 271-274, doi:10.1016/j.ympev.2014.05.024 (2014).

Novick, P. A., Smith, J. D., Floumanhaft, M., Ray, D. A. & Boissinot, S. The evolution and diversity of DNA transposons in the genome of the Lizard Anolis carolinensis. *Genome Biol Evol* **3**, 1-14, doi:10.1093/gbe/evq080 (2011).

da Silva, A. F., Dezordi, F. Z., Loreto, E. L. S. & Wallau, G. L. Drosophila parasitoid wasps bears a distinct DNA transposon profile. *Mob DNA* **9**, 23, doi:10.1186/s13100-018-0127-2 (2018).

Shao, F., Han, M. & Peng, Z. Evolution and diversity of transposable elements in fish genomes. *Sci Rep* **9**, 15399, doi:10.1038/s41598-019-51888-1 (2019).

Klai, K. *et al.* Screening of Helicoverpa armigera Mobilome Revealed Transposable Element Insertions in Insecticide Resistance Genes. *Insects* **11**, doi:10.3390/insects11120879 (2020).

Bellas, C. *et al.* Large-scale invasion of unicellular eukaryotic genomes by integrating DNA viruses. *Proc Natl Acad Sci U S A* **120**, e2300465120, doi:10.1073/pnas.2300465120 (2023).

Chase, E. E., Desnues, C. & Blanc, G. Integrated viral elements suggest the dual lifestyle of Tetraselmis spp. Polinton-like viruses. *Virus Evol* **8**, veac068, doi:10.1093/ve/veac068 (2022).

Fischer, M. G. & Suttle, C. A. A virophage at the origin of large DNA transposons. *Science* **332**, 231-234, doi:10.1126/science.1199412 (2011).

Fischer, M. G. & Hackl, T. Host genome integration and giant virus-induced reactivation of the virophage mavirus. *Nature* **540**, 288-291, doi:10.1038/nature20593 (2016).

Roux, S. *et al.* Updated Virophage Taxonomy and Distinction from Polinton-like Viruses. *Biomolecules* **13**, doi:10.3390/biom13020204 (2023).

Yutin, N., Shevchenko, S., Kapitonov, V., Krupovic, M. & Koonin, E. V. A novel group of diverse Polinton-like viruses discovered by metagenome analysis. *BMC Biol* **13**, 95, doi:10.1186/s12915-015-0207-4 (2015).

Bellas, C. M. & Sommaruga, R. Polinton-like viruses are abundant in aquatic ecosystems. *Microbiome* **9**, 13, doi:10.1186/s40168-020-00956-0 (2021).

Roitman, S. *et al.* Isolation and infection cycle of a polinton-like virus virophage in an abundant marine alga. *Nat Microbiol* **8**, 332-346, doi:10.1038/s41564-022-01305-7 (2023).

Starrett, G. J. *et al.* Adintoviruses: a proposed animal-tropic family of midsize eukaryotic linear dsDNA (MELD) viruses. *Virus Evol* **7**, veaa055, doi:10.1093/ve/veaa055 (2021).

Bessereau, J. L. Transposons in C. elegans. *WormBook*, 1-13, doi:10.1895/wormbook.1.70.1 (2006).

Laricchia, K. M., Zdraljevic, S., Cook, D. E. & Andersen, E. C. Natural Variation in the Distribution and Abundance of Transposable Elements Across the Caenorhabditis elegans Species. *Mol Biol Evol* **34**, 2187-2202, doi:10.1093/molbev/msx155 (2017).

De Ley, P. A quick tour of nematode diversity and the backbone of nematode phylogeny. *WormBook*, 1-8, doi:10.1895/wormbook.1.41.1 (2006).

Vanreusel, A., De Groote, A., Gollner, S. & Bright, M. Ecology and biogeography of free-living nematodes associated with chemosynthetic environments in the deep sea: a review. *PLoS One* **5**, e12449, doi:10.1371/journal.pone.0012449 (2010).

Nigon, V. M. & Felix, M. A. History of research on C. elegans and other free-living nematodes as model organisms. *WormBook* **2017**, 1-84, doi:10.1895/wormbook.1.181.1 (2017).

Frezal, L. & Felix, M. A. C. elegans outside the Petri dish. *Elife* **4**, doi:10.7554/eLife.05849 (2015).

Hodda, M. Phylum Nematoda: a classification, catalogue and index of valid genera, with a census of valid species. *Zootaxa* **5114**, 1-289, doi:10.11646/zootaxa.5114.1.1 (2022).

Majdi, N. & Traunspurger, W. Free-living nematodes in the freshwater food web: a review. *J Nematol* **47**, 28-44 (2015).

Schulenburg, H. & Felix, M. A. The Natural Biotic Environment of Caenorhabditis elegans. *Genetics* **206**, 55-86, doi:10.1534/genetics.116.195511 (2017).

Felix, M. A. *et al.* Natural and experimental infection of Caenorhabditis nematodes by novel viruses related to nodaviruses. *PLoS Biol* **9**, e1000586, doi:10.1371/journal.pbio.1000586 (2011).

Zhang, F. *et al.* Caenorhabditis elegans as a Model for Microbiome Research. *Front Microbiol* **8**, 485, doi:10.3389/fmicb.2017.00485 (2017).

Felix, M. A. & Wang, D. Natural Viruses of Caenorhabditis Nematodes. *Annu Rev Genet* **53**, 313-326, doi:10.1146/annurev-genet-112618-043756 (2019).

Pulavarty, A., Egan, A., Karpinska, A., Horgan, K. & Kakouli-Duarte, T. Plant Parasitic Nematodes: A Review on Their Behaviour, Host Interaction, Management Approaches and Their Occurrence in Two Sites in the Republic of Ireland. *Plants (Basel)* **10**, doi:10.3390/plants10112352 (2021).

Bekal, S., Domier, L. L., Niblack, T. L. & Lambert, K. N. Discovery and initial analysis of novel viral genomes in the soybean cyst nematode. *J Gen Virol* **92**, 1870-1879, doi:10.1099/vir.0.030585-0 (2011).

Bekal, S. *et al.* A novel flavivirus in the soybean cyst nematode. *J Gen Virol* **95**, 1272-1280, doi:10.1099/vir.0.060889-0 (2014).

Lin, J. *et al.* A novel picornavirus-like genome from transcriptome sequencing of sugar beet cyst nematode represents a new putative genus. *J Gen Virol* **99**, 1418-1424, doi:10.1099/jgv.0.001139 (2018).

Ruark, C. L. *et al.* Soybean cyst nematode culture collections and field populations from North Carolina and Missouri reveal high incidences of infection by viruses. *PLoS One* **12**, e0171514, doi:10.1371/journal.pone.0171514 (2017).

Vieira, P. & Nemchinov, L. G. A novel species of RNA virus associated with root lesion nematode Pratylenchus penetrans. *J Gen Virol* **100**, 704-708, doi:10.1099/jgv.0.001246 (2019).

Franz, C. J., Zhao, G., Felix, M. A. & Wang, D. Complete genome sequence of Le Blanc virus, a third Caenorhabditis nematode-infecting virus. *J Virol* **86**, 11940, doi:10.1128/JVI.02025-12 (2012).

Williams, S. H. *et al.* Discovery of two highly divergent negative-sense RNA viruses associated with the parasitic nematode, Capillaria hepatica, in wild Mus musculus from New York City. *J Gen Virol* **100**, 1350-1362, doi:10.1099/jgv.0.001315 (2019).

Brown, D. J. F., Robertson, W. M. & Trudgill, D. L. Transmission of Viruses by Plant Nematodes. *Annual Review of Phytopathology* **33**, 223-249, doi:DOI 10.1146/annurev.py.33.090195.001255 (1995).

Hess, R. T. & Poinar, G. O., Jr. Iridoviruses infecting terrestrial isopods and nematodes. *Curr Top Microbiol Immunol* **116**, 49-76, doi:10.1007/978-3-642-70280-8_4 (1985).

Poinar, G. O., Jr., Hess, R. T. & Cole, A. Replication of an Iridovirus in a nematode (Mermithidae). *Intervirology* **14**, 316-320, doi:10.1159/000149202 (1980).

Widen, S. A. *et al.* Virus-like transposons cross the species barrier and drive the evolution of genetic incompatibilities. *Science* **380**, eade0705, doi:10.1126/science.ade0705 (2023).

Widen, S. A., Bes, I. C., Koreshova, A., Krogull, D. & Burga, A., doi:10.1101/2022.07.12.499685 (2022).

Vance, T. D. R. & Lee, J. E. Virus and eukaryote fusogen superfamilies. *Curr Biol* **30**, R750-R754, doi:10.1016/j.cub.2020.05.029 (2020).

Blanco, L. & Salas, M. Mutational analysis of bacteriophage phi 29 DNA polymerase. *Methods Enzymol* **262**, 283-294, doi:10.1016/0076-6879(95)62024-9 (1995).

Blanco, L. & Salas, M. Relating structure to function in phi29 DNA polymerase. *J Biol Chem* **271**, 8509-8512, doi:10.1074/jbc.271.15.8509 (1996).

Redrejo-Rodriguez, M. *et al.* Primer-Independent DNA Synthesis by a Family B DNA Polymerase from Self-Replicating Mobile Genetic Elements. *Cell Rep* **21**, 1574-1587, doi:10.1016/j.celrep.2017.10.039 (2017).

Kazlauskas, D., Krupovic, M., Guglielmini, J., Forterre, P. & Venclovas, C. Diversity and evolution of B-family DNA polymerases. *Nucleic Acids Res* **48**, 10142-10156, doi:10.1093/nar/gkaa760 (2020).

Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583-589, doi:10.1038/s41586-021-03819-2 (2021).

Mirdita, M. *et al.* ColabFold: making protein folding accessible to all. *Nat Methods* **19**, 679-682, doi:10.1038/s41592-022-01488-1 (2022).

Gabler, F. *et al.* Protein Sequence Analysis Using the MPI Bioinformatics Toolkit. *Curr Protoc Bioinformatics* **72**, e108, doi:10.1002/cpbi.108 (2020).

Shen, B. W. *et al.* Unusual target site disruption by the rare-cutting HNH restriction endonuclease PacI. *Structure* **18**, 734-743, doi:10.1016/j.str.2010.03.009 (2010).

Stein, L. D. *et al.* The genome sequence of Caenorhabditis briggsae: a platform for comparative genomics. *PLoS Biol* **1**, E45, doi:10.1371/journal.pbio.0000045 (2003).

Stevens, L. *et al.* Chromosome-Level Reference Genomes for Two Strains of Caenorhabditis briggsae: An Improved Platform for Comparative Genomics. *Genome Biol Evol* **14**, doi:10.1093/gbe/evac042 (2022).

Hausner, G., Hafez, M. & Edgell, D. R. Bacterial group I introns: mobile RNA catalysts. *Mob DNA* **5**, 8, doi:10.1186/1759-8753-5-8 (2014).

Lee, Y. C. *et al.* Single-worm long-read sequencing reveals genome diversity in free-living nematodes. *Nucleic Acids Res*, doi:10.1093/nar/gkad647 (2023).

Osman, G. A. *et al.* Natural Infection of C. elegans by an Oomycete Reveals a New Pathogen-Specific Immune Response. *Curr Biol* **28**, 640-648 e645, doi:10.1016/j.cub.2018.01.029 (2018).

Grover, M. & Barkoulas, M. C. elegans as a new tractable host to study infections by animal pathogenic oomycetes. *PLoS Pathog* **17**, e1009316, doi:10.1371/journal.ppat.1009316 (2021).

Warburton, P. E., Giordano, J., Cheung, F., Gelfand, Y. & Benson, G. Inverted repeat structure of the human genome: the X-chromosome contains a preponderance of large, highly homologous inverted repeats that contain testes genes. *Genome Res* **14**, 1861-1869, doi:10.1101/gr.2542904 (2004).

Sievers, F. *et al.* Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Mol Syst Biol* **7**, 539, doi:10.1038/msb.2011.75 (2011).

Sievers, F. & Higgins, D. G. Clustal Omega for making accurate alignments of many protein sequences. *Protein Sci* **27**, 135-145, doi:10.1002/pro.3290 (2018).

Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. *Mol Biol Evol* **37**, 1530-1534, doi:10.1093/molbev/msaa015 (2020).

Hoang, D. T., Chernomor, O., von Haeseler, A., Minh, B. Q. & Vinh, L. S. UFBoot2: Improving the Ultrafast Bootstrap Approximation. *Mol Biol Evol* **35**, 518-522, doi:10.1093/molbev/msx281 (2018).

Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A. & Jermiin, L. S. ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat Methods* **14**, 587-589, doi:10.1038/nmeth.4285 (2017).

Huerta-Cepas, J., Serra, F. & Bork, P. ETE 3: Reconstruction, Analysis, and Visualization of Phylogenomic Data. *Mol Biol Evol* **33**, 1635-1638, doi:10.1093/molbev/msw046 (2016).

Larkin, M. A. *et al.* Clustal W and Clustal X version 2.0. *Bioinformatics* **23**, 2947-2948, doi:10.1093/bioinformatics/btm404 (2007).

Tareen, A. & Kinney, J. B. Logomaker: beautiful sequence logos in Python. *Bioinformatics* **36**, 2272-2274, doi:10.1093/bioinformatics/btz921 (2020).

Fu, L., Niu, B., Zhu, Z., Wu, S. & Li, W. CD-HIT: accelerated for clustering the next-generation sequencing data. *Bioinformatics* **28**, 3150-3152, doi:10.1093/bioinformatics/bts565 (2012).

Capella-Gutierrez, S., Silla-Martinez, J. M. & Gabaldon, T. trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972-1973, doi:10.1093/bioinformatics/btp348 (2009).

Mirdita, M., Steinegger, M. & Soding, J. MMseqs2 desktop and local web server app for fast, interactive sequence searches. *Bioinformatics* **35**, 2856-2858, doi:10.1093/bioinformatics/bty1057 (2019).

Steinegger, M. & Soding, J. MMseqs2 enables sensitive protein sequence searching for the analysis of massive data sets. *Nat Biotechnol* **35**, 1026-1028, doi:10.1038/nbt.3988 (2017).

Katoh, K., Rozewicki, J. & Yamada, K. D. MAFFT online service: multiple sequence alignment, interactive sequence choice and visualization. *Brief Bioinform* **20**, 1160-1166, doi:10.1093/bib/bbx108 (2019).

Ye, Y. & Godzik, A. FATCAT: a web server for flexible structure comparison and structure similarity searching. *Nucleic Acids Res* **32**, W582-585, doi:10.1093/nar/gkh430 (2004).

Li, Z., Jaroszewski, L., Iyer, M., Sedova, M. & Godzik, A. FATCAT 2.0: towards a better understanding of the structural diversity of proteins. *Nucleic Acids Res* **48**, W60-W64, doi:10.1093/nar/gkaa443 (2020).

Eddy, S. R. Accelerated Profile HMM Searches. *PLoS Comput Biol* **7**, e1002195, doi:10.1371/journal.pcbi.1002195 (2011).

Jurka, J. Repbase update: a database and an electronic journal of repetitive elements. *Trends Genet* **16**, 418-420, doi:10.1016/s0168-9525(00)02093-x (2000).

Jurka, J. *et al.* Repbase Update, a database of eukaryotic repetitive elements. *Cytogenet Genome Res* **110**, 462-467, doi:10.1159/000084979 (2005).

Tsutakawa, S. E. *et al.* Crystallographic and functional studies of very short patch repair endonuclease. *Mol Cell* **3**, 621-628, doi:10.1016/s1097-2765(00)80355-x (1999).

Ahmed, M. *et al.* Phylogenomic Analysis of the Phylum Nematoda: Conflicts and Congruences With Morphology, 18S rRNA, and Mitogenomes. *Frontiers in Ecology and Evolution* **9**, doi:10.3389/fevo.2021.769565 (2022).

Tsutakawa, S. E. & Morikawa, K. The structural basis of damaged DNA recognition and endonucleolytic cleavage for very short patch repair endonuclease. *Nucleic Acids Res* **29**, 3775-3783, doi:10.1093/nar/29.18.3775 (2001).