# The bacterial microbiome of symbiotic and menthol-bleached polyps of long-term aquarium-reared *Galaxea fascicularis*

Puntin Giulia[1]*, Wong Jane C. Y.[2,3], Röthig Till[4,5], Baker David M.[2,3], Sweet Michael[5], Ziegler Maren[1]*

[1] Marine Holobiomics Lab, Department of Animal Ecology & Systematics, Justus Liebig University Giessen, Heinrich-Buff-Ring 26-32 IFZ, 35392 Giessen, Germany
[2] The Swire Institute of Marine Science, University of Hong Kong, Cape d'Aguilar Road, Shek O, Hong Kong SAR, China
[3] School of Biological Sciences, Kadoorie Biological Science Building, University of Hong Kong, Pok Fu Lam Road, Hong Kong SAR, China
[4] Branch of Bioresources, Fraunhofer Institute for Molecular Biology and Applied Ecology, Giessen, Germany
[5] Aquatic Research Facility, Nature-based Solutions Research Centre, University of Derby, UK

       *Corresponding authors

Correspondence: maren.ziegler@bio.uni-giessen.de; puntin.giulia@gmail.com

**ABSTRACT**

Coral reefs support the livelihood of half a billion people but are at high risk of collapse due to the vulnerability of corals to climate change and local anthropogenic stressors. While understanding coral functioning is essential to guide conservation efforts, research is challenged by the complex nature of corals. They exist as metaorganisms (holobionts), constituted by the association between the (coral) animal host, its obligate endosymbiotic algae (Symbiodiniaceae), and other microorganisms comprising bacteria, viruses, archaea, fungi and other protists. Researchers therefore increasingly turn to model organisms to unravel holobiont complexity, dynamics, and how these determine the health and fitness of corals. The coral *Galaxea fascicularis* is an emerging model organism for coral symbiosis research with demonstrated suitability to aquarium rearing and reproduction, and to manipulation of the host-Symbiodiniaceae symbiosis. However, little is known about the response of the *G. fascicularis* microbiome to menthol bleaching—the experimental removal of the Symbiodiniaceae which represents the first step in coral-algal symbiosis manipulation. For this, we characterized the bacterial microbiome of symbiotic and menthol-bleached *G. fascicularis* originating from the Red Sea and South China Sea (Hong Kong) that were long-term aquarium-reared in separate facilities. We found that the coral-associated microbiomes were composed of relatively few bacterial taxa, when compared to literature reports from wild corals. Symbiotic polyps (clonal replicates) from the same colony had similar microbiomes, which were distinct from those of other colonies despite co-culturing in shared aquaria. The response of the bacterial microbiome to menthol bleaching varied between the two facilities, warranting further investigation into the role of rearing conditions. Nevertheless, the changes in community composition observed in both facilities appeared to be stochastic and indicative of a dysbiotic state. Considering the importance of bleaching treatment of captive corals for symbiosis research, our results—although preliminary—contribute fundamental knowledge for the development of the Galaxea model for coral symbiosis research.

*Keywords:* coral, model organism, 16S, holobiont, symbiosis, aquarium, ex situ

**Introduction**

By the end of this century, the livelihood of more than half a billion people will largely depend on the ability
of corals to cope with changing ocean conditions (Hughes et al. 2017). Corals are vulnerable to climate change,
pollution, and overfishing, which have caused the loss of half of the world's coral cover since the 1950s (Eddy
et al. 2021; Eakin et al. 2022). As such, the reef structures they build and the multi-billion dollar ecosystem
services they provide are at high risk of collapse (Costanza et al. 2014; van Hooidonk et al. 2016). Considering
the high stakes, understanding coral functioning is essential to predict future scenarios and guide management
and conservation efforts.
Corals exist as metaorganisms, or so-called holobionts (Rohwer et al. 2002), where complexity hinders our
ability to unravel coral functioning and how it ultimately affects coral physiology and ecology (Rosenberg et al.
2007; Jaspers et al. 2019). Indeed, the coral holobiont comprises the animal host, obligate intracellular algal
symbionts (Symbiodiniaceae), a rich and diverse bacterial community, together with other microorganisms
such as archaea, fungi, viruses, and protists (Bourne et al. 2016; Pogoreutz et al. 2020). Even richer than their
taxonomy is the potential diversity of relationships and interactions among members, and how these could
contribute to holobiont health and resilience (Thompson et al. 2015; Pogoreutz et al. 2020).
Besides the coral host, the best understood member of the coral holobiont is the algal symbiont, as its
photosynthesis-derived energy fuels the calcification process that builds reefs and allows corals to thrive in
nutrient-poor waters (Muscatine and Porter 1977; Muscatine 1990). Bacteria are also involved in nutrient
cycling and metabolism (Robbins et al. 2019; Tandon et al. 2020), as well as other essential physiological
processes such as development (Webster et al. 2004; Tebben et al. 2015) and immunity (Certner and Vollmer
2018; Miura et al. 2019). Interestingly, bacteria also mediate host-Symbiodiniaceae dynamics through the
mitigation of thermal and light stress on the algal symbiont (Motone et al. 2020; Connelly et al. 2022), and they
produce antimicrobial agents from an organosulfur compound released by Symbiodiniaceae (Raina et al. 2016,
2017). A growing body of knowledge suggests a central role of tripartite interactions between bacteria,
Symbiodiniaceae, and the coral host in nutrition, health and fitness (reviewed in Matthews et al. 2020).
However, given the intricacy of players' diversity and their metabolic capacities, linking partner identity to
function and holobiont phenotype proves particularly challenging.
Model organisms can help unravel holobiont complexity through manipulation. Comparison of different
host-Symbiodiniaceae combinations in the sea anemone Aiptasia revealed that heat-tolerance of the symbiont
is not linearly transferred to the host (Chakravarti et al. 2017; Gabay et al. 2019; Herrera et al. 2020). Thus
complex mechanisms where holobiont properties cannot be predicted as the "sum of its parts" require a more
holistic approach (Goulet et al. 2020). Such empirical testing of multi-partner interactions relies on the ability
to study the holobiont upon experimental manipulation, namely by removing and/or adding members (Jaspers
et al. 2019). Most of such studies in the field were conducted with non-calcifying species such as Aiptasia or
the hydroid polyp *Hydra*, which are well established, tractable model organisms (Weis et al. 2008; Galliot 2012).
Yet, while these models have proven instrumental for breakthrough discoveries in the cnidarian symbiosis field
(e.g., Weis 2008; Murillo-Rincon et al. 2017; Pietschke et al. 2017; Gabay et al. 2018), they lack key features of
reef-building corals such as calcification and the obligate endosymbiosis necessary to understand the ecology
of coral functioning. To include these critical features, research with corals is irreplaceable (Puntin et al. 2022b).
The reef-building coral *Galaxea fascicularis* (Linnaeus, 1767) has been proposed as a model species for coral
symbiosis research (Puntin et al. 2022a). This species is well represented in the literature, featured in studies
that characterize the coral gastric cavity (Agostini et al. 2012; Zhou et al. 2020) and the calcification processes
(Al-Horani et al. 2003, 2005, 2007), among others (Ferrier-Pagès et al. 1998; Niu et al. 2016; Miura et al. 2019).
Introducing *G. fascicularis* as a model system, we previously demonstrated its ease of rearing in simplified
systems (closed, small volume), compatibility with *ex-situ* reproduction, effective removal of the algal symbiont
through menthol bleaching, and subsequent reestablishment of the symbiosis with both cultured and
environmental Symbiodiniaceae in adult individuals (Puntin et al. 2022a). This demonstrated the potential to

experimentally produce a variety of coral-Symbiodiniaceae combinations to study symbiosis functioning and partner compatibility in a true reef-building coral. While recent coral probiotic approaches rapidly expand our knowledge on the functions of the bacterial fraction (Rosado et al. 2019; Peixoto et al. 2021), untangling the complexity in the holobiont requires detailed knowledge of the interrelationships that consider all three partners.

One of the main knowledge gap to advance mechanistic symbiosis research in the Galaxea model system is the effect of menthol bleaching of Symbiodiniaceae on the remaining coral microbiome. Menthol is becoming increasingly common in manipulatve experiments due to its efficacy while causing virtually no mortality (Wang et al. 2012; Matthews et al. 2015; Puntin et al. 2022a). To date, menthol bleaching has been used with a range of symbiotic cnidarians, including jellyfish (Röthig et al. 2021), anemones (Matthews et al. 2015; Dani et al. 2016), corallimorpharia (Lin et al. 2019), and nine species of reef-building corals (Wang et al. 2012, 2019; Puntin et al. 2022a; Scharfenstein et al. 2022; Chan et al. 2023). Yet, its impact on the bacterial fraction remains unknown. To address this knowledge gap, we characterized the bacterial microbiome of symbiotic and menthol-bleached *G. fascicularis* polyps from the central Red Sea and the South China Sea that were maintained in two separate facilities for several months.

# Materials and Methods

**Coral collection and long-term aquarium rearing**

Colonies of *Galaxea fascicularis* were collected from two locations: the Red Sea (hereafter referred as "Red Sea") and Hong Kong in the South China Sea (hereafter referred as "Hong Kong"). Red Sea colonies (n = 3) were collected from the central Saudi Arabian Red Sea at "Al Fahal" reef (N 22˚18.324' E 38˚57.930'), at 9-13 m depth in March 2019 (CITES permit 19-SA-000096-PD) and transported to the Ocean2100 aquarium facility at Justus Liebig University Giessen (Germany) (Schubert and Wilke 2018). In the aquarium system, light was provided by white and blue fluorescent lamps with a light:dark cycle of 12:12 h at 130-160 µmol photons $m^{-2}$ $s^{-1}$ to approximate light conditions at the collection site (Ziegler et al. 2015). Salinity was maintained around 35 and temperature at 26 °C. Colonies were fed daily with a combination of frozen copepods, Artemia, krill, and Mysis. Hong Kong colonies (n = 2) were collected from ≤ 5 m depth from Crescent Island (N 22° 31' 51.035", E 114° 18' 53.888) in June 2019 and transported to the University of Hong Kong (HKU), where they were maintained in a 500-L aquarium equipped with a filtration system and protein skimmer, and fed daily with Reef-Roids (Polyplab) and frozen artemia. Light intensity, salinity, and temperature conditions were consistent with those maintained in the Ocean2100 facility.

**Menthol bleaching**

At both locations, individual (clonal) polyps were mechanically isolated from their colony, mounted on coral glue (Red Sea colonies, JLU-Ocean2100; Grotech, Cora-Fix SuperFast) or attached to small ceramic tiles (Hong Kong colonies, HKU; Aron Alpha, GEL-10) (Fig. 1). After 10-14 days of healing, polyps were randomly divided between a 'symbiotic' and a 'bleached' group at each location. Both groups were maintained under the same conditions until healed, then the 'bleached' group was treated with menthol to chemically induce bleaching. Menthol treatment started after approximately seven (Red Sea) and three (Hong Kong) months of captivity, and it was replicated at the two facilities following a protocol modified from Wang et al. (2012). Specifically, three days treatment in 0.38 mM menthol solution in filtered (1.2 µm) artificial seawater (FASW) was followed by one day of rest and another day of menthol treatment. Menthol incubations lasted 8 h during the light period.

Bleaching was assessed in the Red Sea polyps by visual inspection under a fluorescence stereomicroscope (Leica MZ16 F) 10 days after the menthol treatment, when algal cells were not detectable in any polyp. At the same time point, Hong Kong polyps also appeared fully bleached under microscopic inspection (Olympus Optical, mod. CHK at 400×).

### Post-bleaching rearing conditions

To prevent coral from Symbiodiniaceae exposure and symbiosis re-establishment, all polyps were kept in simplified (see below for details) systems with FASW (1.2 µm) after the menthol bleaching treatment. Here, polyps were fed daily with one small frozen adult Artemia each, followed by partial (~10 %) water change after 2-3 h. At both facilities, temperature, light, and salinity were maintained consistent with the long-term rearing conditions, while the setups differed. Specifically, at the Ocean2100 facility, the polyps were distributed among eight 5-L glass tanks (20 cm × 30 cm) (four per treatment), each equipped with a small pump (Resun SP-500) in a temperature-controlled water bath. At HKU, symbiotic and bleached polyps were maintained in separate 600 ml glass jars, each holding ~6 polyps and equipped with magnetic stir bars for water flow inside a Plant Growth Chamber (Panasonic MLR-352H-PA).

**Figure 1** – Visual summary of the experimental design and data processing. Two samples (outlined polyps) were excluded from processing due to damage during shipping, while one sample (grey shaded) was omitted after rarefaction for alpha and beta diversity analysis owing to low sequencing depth. Symbiotic polyps with successful ITS2 sequencing are marked with their sample ID in bold.

### Sampling for microbial analysis

16S rRNA amplicon sequencing was employed to characterize the bacterial communities of symbiotic and menthol-bleached polyps, and ITS2 amplicon sequencing for the Symbiodiniaceae communities of symbiotic polyps of *G. fascicularis* colonies from the two geographic locations (Fig. 1).

At both locations, three polyps per colony (5 colonies: RS1, RS2, RS3, HK1, HK2) per state (2 states: symbiotic, bleached) were sampled on the 13th day after the menthol treatment (n = 15 bleached and 15 symbiotic polyps). The polyps were rinsed with seawater, separated from the substrate, placed in sterile tubes, and stored at -80 °C. For transport, polyps were stored in RNAlater (R0901-100ML, Sigma-Aldrich, Hong Kong S.A.R.). Two samples from Hong Kong (1 symbiotic and 1 bleached) were damaged during shipping and therefore excluded from processing. All samples were processed together for DNA extraction (University of

Derby's Aquatic Research Facility, UK) and subsequently sequenced in the same sequencing run (Bart's and the
London Genome Centre, Queen Mary, University of London).

**Bacterial and Symbiodiniaceae community analysis**
DNA was extracted using the Qiagen DNeasy 96 Blood & Tissue kit. About 10 mg of coral tissue per sample
was used as starting material. Extractions followed the user manual with centrifugations at working steps 10,
12, and 16 performed at 1500 $g$ at doubled centrifugation times. Extracted DNA was sent to the sequencing
facility for quality control, PCR, library preparation, and pair-end sequencing with Illumina MiSeq platform v3
(2 × 300 bp). The 16S rRNA gene region V5/V6 was amplified using the primers 784F and 1061R (Andersson et
al. 2008), while Symbiodiniaceae ITS2 region was amplified using the primers SYM_VAR_5.8S2 and SYM_VAR
(Hume et al. 2018). A contamination control consisting of pure RNAlater buffer was included in all steps.
Bacterial sequencing data were processed in Qiime2 (v.2021.11, Bolyen et al. 2019) and analyzed in R
(v.4.1.0, R Core Team 2021). After primer removal, forward and reverse reads were truncated to 232 and 234
nt respectively, paired, dereplicated, quality checked, cleaned, and clustered to amplicon sequence variants
(ASVs) using the denoise-paired method in DADA2 (Callahan et al. 2016). This resulted in a total of 138,620
sequences and 547 ASVs. ASVs were taxonomically assigned using a weighted classifier trained against the
SILVA 138 database (99 % clustering, full length) (Yilmaz et al. 2013) with the classify-sklearn method from 'q2-
feature-classifier' plug-in (Bokulich et al. 2018; Kaehler et al. 2019; Robeson et al. 2020). Then, sequences
assigned to "mitochondria", "chloroplast", "Archaea", "Eukaryota", or "unknown" at the phylum level were
removed. Sequences found in the control sample were considered potential lab contaminants and evaluated
based on presence/absence across the coral samples and habitat description (e.g., known contaminants) of
the closest BLASTn matches (GenBank) (for full details see https://zenodo.org/record/10551928,
"01_find_contaminants.R"). This led to the removal of five ASVs and their 18,990 sequences, and resulted in a
final data set of 112,789 sequences and 515 ASVs across 28 samples (after excluding the contamination
control).
Rarefaction and alpha and beta diversity calculations were performed with the R package 'phyloseq'
(v.1.38.0, McMurdie and Holmes 2013), 'metagMisc' (v.0.0.4, Mikryukov 2023) and 'btools' (v.0.0.1, Battaglia
2022). Samples were rarefied to 2,690 sequences (based on 1,000 iterations of random subsampling without
replacement), which caused the exclusion of one sample (symbiotic colony "RS3") due to low sequencing
depth. Rarefaction curves showed that for most samples the number of ASVs plateaued before the rarefaction
depth indicating that most of the diversity was captured and retained after rarefaction (Fig. S1). Alpha diversity
was estimated through multiple indices chosen for their complementarity and comparability with previous
studies (observed richness, Chao1, Shannon diversity, Simpson diversity, Pielou's evenness, and Faith's
phylogenetic diversity). Differences in alpha diversity between symbiotic and bleached individuals were tested
with t-tests or Mann-Whitney $U$ tests depending on data distribution and variance. Beta diversity based on
Bray-Curtis distances was visualized with non-metric multidimensional scaling (nMDS).
The contribution of host origin (Red Sea, Hong Kong) and symbiotic state (symbiotic, bleached) to
microbiome community structure was assessed using multivariate homogeneity of groups dispersions analysis
with the betadisper function (PERMDISP2), and one-way and two-way permutational multivariate analysis of
variance (PERMANOVA) on Bray-Curtis distances with the adonis2 function in the 'vegan' R package (v.2.5.7,
Dixon 2003; Oksanen et al. 2020). Relative abundance bubble plots of bacterial community composition at
bacterial family level and of core ASVs (occur in all groups by state and geographic origin), and the UpSet plot
(Lex et al. 2014) were generated from non-rarefied data. The UpSet plot was created with the package 'UpSetR'
(v.1.4.0, Conway et al. 2017). All other plots were created in 'ggplot2' (v.3.3.5, Wickham 2016).
For Symbiodiniaceae community analysis, raw ITS2 sequencing data were analyzed using the SymPortal
workflow remote instance (Hume et al. 2019). ITS2 sequencing produced poor results and it was only possible
to characterize eight of the symbiotic samples (6 Red Sea, 2 Hong Kong; see Fig. 1). Of these, two did not pass
quality check (i.e., they had < 200 sequences/genus) and SymPortal could not predict ITS2 type profiles.
Therefore, we report (post-MED) ITS2 sequences.

# Results

**Microbial diversity and richness were unaffected by menthol bleaching**

When considering all colonies together, alpha diversity remained similar between symbiotic and menthol-bleached samples across all diversity and richness indices tested (incl. observed richness, Chao1, Shannon diversity, Simpson diversity, Pielou's evenness, and Faith's phylogenetic diversity, see Table S1, S2), and regardless of their origin ($P_{\text{t-test}}$ or $P_{\text{Mann-Whitney U-test}}$ > 0.05; Fig. 2A). When considering each individual colony, alpha diversity was remarkably and consistently low in symbiotic RS1 polyps, while in their bleached counterparts it was in range with the other colonies (Fig. 2A, Fig. S2, Tab. S2). Within-colony difference between symbiotic and bleached polyps could only be tested for RS1 and RS2, and it was significant in RS1 across all alpha diversity indices tested ($P_{\text{Welch}}$ < 0.05, Tab. S3).

**Bacterial communities were generally uneven**

A small number of ASVs dominated the bacterial communities, where the 3 and 9 most abundant ASVs accounted for > 25 % and > 50 % of the total number of sequences, respectively (Fig. 2B). Evenness was on average lower among the symbiotic polyps. Specifically, in the symbiotic samples, the 5 (for Red Sea) and 4 (for Hong Kong) most abundant ASVs account for > 50 % of total reads, while in the bleached samples it took 7 (for Red Sea) and 12 (for Hong Kong) ASVs to pass the 50 % relative abundance threshold.

**Figure 2** - Diversity and evenness of the bacterial communities of symbiotic and menthol-bleached polyps of *Galaxea fascicularis*. (A) Comparison of observed number of ASVs, and Shannon and Simpson diversity index between symbiotic and bleached samples; (B) ASV accumulation curve for the whole data set, and separated by symbiotic state and geographic origin.

**Microbial community dissimilarity patterns differed by geographic origin of colonies**

Microbial communities of Red Sea samples showed significantly larger dissimilarities than those from Hong Kong, both within and between colonies (ANOVA on PERMDISP2, F = 118.7, P < 0.0001, Fig. 3A; Mann-Whitney U-test on pairwise Bray-Curtis distances, W = 102, P = 0.0001, Fig. 3B, Fig. S3).

Microbial communities of symbiotic Red Sea polyps clustered by colony, and Red Sea colonies appeared as different from each other as they were from those of Hong Kong colonies. Interestingly, one symbiotic colony

193  originating from the Red Sea did however share a similar microbial community with those from Hong Kong (Fig.
194  3A).

**Menthol-bleaching elicited stochastic changes in the microbial communities**

Changes in community composition between symbiotic states differed for coral colonies from the two
regions. For the Red Sea, there was no clear distinction in bacterial community composition between symbiotic
and bleached polyps in terms of location and dispersion in the ordination space, when considering all colonies
together (PERMANOVA, F = 0.76, PERMDISP2, F = 0.77, P > 0.05; Fig. 3C). For Hong Kong, the microbial
communities of symbiotic and bleached polyps were significantly different (PERMANOVA, F = 4.0, P < 0.01; Fig.
3D), while the differnce in dispersion was (just marginally) not statistically significant (PERMDISP2, F = 5.05 , P
= 0.057). However, while symbiotic polyps clustered by colony for both Red Sea and Hong Kong (indicating
similar microbial communities), bleached polyps showed no such clear grouping (Red Sea) or larger scattering
compared to symbiotic polyps (Hong Kong), and collectively had a significantly higher within-colony
dissimilarity ($P_{Mann-Whitney}$ = 0.0001, Fig. 3A,B, Fig. S3). This indicates random changes in the communities of the
menthol-bleached polyps.

**Figure 3** – Microbial community structure of *Galaxea fascicularis* from the Red Sea and Hong Kong. Non-
metric multi-dimensional scaling (nMDS) plot of bacterial community composition based on Bray-Curtis
dissimilarity for all polyps (A), Red Sea polyps (C), and Hong Kong polyps (D), and comparison of pairwise
dissimilarity between symbiotic and bleached polyps of the same colony (only groups with $n$ = 3
considered, but comparable results were found considering groups with $n$ < 3, see Fig. S3) (B). Ellipses =
95 % confidence intervals (A, C, D); colors (A, D) or shapes (C, D) denote colony identity, filled symbols =
symbiotic polyps, hollow symbols = bleached polyps.

### Three bacterial families dominated most microbial communities

A total of 138 bacterial families were found across the 28 sampled polyps. Of these, 104 bacterial families
occurred in symbiotic (74 in Red Sea, 66 in Hong Kong) and 109 in bleached (89 in Red Sea, 69 in Hong Kong)
polyps. The three most abundant families *Rhodobacteraceae* (25.8 %), *Alteromonadaceae* (14.6 %), and
*Moraxellaceae* (10.1 %) together represented > 50 % of all sequences, and the 11 most abundant families
represented > 75 % of all sequences. Symbiotic colonies were largely dominated by members of the two
bacterial families *Rhodobacteraceae* and *Alteromonadaceae*. Except for the two Red Sea colonies RS1 and RS3

that were dominated by *Moraxellaceae* and *Bacillaceae*, or that showed no consistently dominant family across individual polyps, respectively.

Bleached polyps were dominated by varying bacterial families, with inconsistent patterns between and within colonies and regions. In addition to the dominant families in symbiotic colonies, *Endozoicomonadaceae*, unclassified *Cellvibrionales*, and *Amoebophilaceae* became dominant in some bleached polyps. While members of the family *Endozoicomonadaceae* were the most abundant fraction in one bleached polyp (39 %; RS3), they were only present in 15 of the 28 polyps (7 symbiotic, 8 bleached), and with low relative abundance (mean 2.2 % in symbiotic, and 8.9 % in bleached). No single taxon (neither at ASV nor at bacterial familiy level) was significantly different in relative abundance between symbiotic and bleached polyps, regardless of whether samples from the Red Sea and Hong Kong were considered together or separately (Wilcoxon test with Benjamini-Hochman correction, all P > 0.05).

**Figure 4** – Relative abundance of bacterial families in symbiotic and bleached *Galaxea fascicularis* polyps from the Red Sea and Hong Kong. Bubble size is proportional to the relative abundance per polyp of the 17 most abundant families, which account for 83.7 % of the total number of reads; all other 121 families that had a relative abundance < 1 % are grouped as 'Others'.

**28 core ASVs were shared by all coral colonies and symbiotic states**

Altogether, there were more exclusive than shared ASVs between experimental groups (Fig. 5A). Specifically, 33.4 % and 52.9 % of ASVs were exclusively found in symbiotic or bleached samples, respectively, while 5.4 % of ASVs (28) were found in all groups (symbiotic state and geographic origin) and were considered core taxa. Although no ASV was found in all samples (nor in all samples of the same experimental group, Tab. S4), three ASVs occurred in ≥ 70 % of samples, and five additional ASVs occurred in ≥ 60 % of samples (Fig. 5B). The sequences of these eight ASVs, which also corresponded to the most abundant of the 28 core ASVs, were BLAST-searched against the NCBI Nucleotide collection (Tab. 1), which revealed that most of the matching sequences were from samples associated with marine environments and/or organisms. Among these, the most abundant bacterial sequences belonged to the genera *Alteromonas*, *Ruegeria*, and *Nautella* (Fig. 5B, Tab. 1). Interestingly, two ASVs (ASV_001 and ASV_006), both assigned to the genus *Ruegeria*, matched with strains isolated from aquarium-reared *Galaxea fascicularis* from the South China Sea (collected in Hainan Island, China,

250 Zhou et al. 2020) and Japan (Miura et al. 2019). Further, three ASVs (ASV_006, ASV_018, ASV_020) matched
251 with bacteria identified in *Acropora* spp. and *Pocillopora* spp. from the central Red Sea, amplified with the same
252 primer set used for this study (Tab. 1).

### All colonies of *Galaxea fascicularis* were dominated by *Cladocopium* spp. symbionts

Symbiodiniaceae composition was consistent in polyps from each colony and differed by region.
*Cladocopium* spp. ITS2 sequences accounted for > 92 % of the sequences in all samples but one (RS2, 76 %)
(Fig. S4). Of these, C1 was by large the dominant ITS2 sequence (~ 33 – 80 % relative abundance). Sequences
C1b, C41, and C41f were exclusively and consistently found in polyps of one Red Sea colony (RS3), where they
collectively accounted for ~ 26 % of reads. *Durusdinium* spp. sequences were found in only one colony from
the Red Sea (RS2), where they accounted for 4 % and 24 % of the sequences, and of which the most abundant
sequences were D1 and D4 (Fig. S4). One Red Sea sample (RS3) also hosted sequence A1 (genus *Symbiodinium*)
at ~ 5 % abundance. In both Hong Kong colonies C1 was the dominant sequence (68 – 80 %), with C1c present
in lower abundance (15 – 20 %) (Fig. S4).

**Figure 5** - Overview of total and core ASVs in *Galaxea fascicularis* symbiotic and menthol-bleached polyps
from the Red Sea and Hong Kong. (A) UpSet diagram showing the number of observed ASVs per group

and per intersection. (B) Relative abundance of core bacterial ASVs by host origin and symbiotic state,
considering only ASVs present in ≥ 70 % (top-darker grey block) and between 60 % and 70 % of samples
(bottom-lighter grey block); bubble size is proportional to the relative abundance of ASVs in each sample.

**Table 1** - Summary of NCBI BLASTn matches for the eight most common core ASVs (occur in both symbiotic
and bleached samples from both Red Sea and Hong Kong, and in > 60 % of all samples) in order of
abundance. For brevity, only the top three matches are reported (100 % query cover) with information
on isolation source and location, where coral species are highlighted in bold. Taxonomy is reported as the
lowest taxonomic level assigned by the Qiime2 classifier.

| Taxonomy | ASV no. | GenBank Accession no. | Identity (%) | Isolation source (location) |
|---|---|---|---|---|
| *Ruegeria* | ASV_001 | MK967091 | 100.00 | Jellyfish, *Aurelia aurita* polyp (Kiel Bight, Baltic Sea) |
| | | MT187950 | 100.00 | ***Galaxea fascicularis*** (Hainan, South China Sea) |
| | | MT187656 | 100.00 | ***Galaxea fascicularis*** (Hainan, South China Sea) |
| *Alteromonas* | ASV_003 | MT525287 | 100.00 | Marine sediment (South Korea) |
| | | MK967157 | 100.00 | Artificial Seawater (Kiel Bight, Baltic Sea) |
| | | MT515801 | 100.00 | Marine sediment (South Korea) |
| *Nautella* | ASV_005 | LC543501 | 100.00 | Coastal seawater (Japan) |
| | | MH556771 | 100.00 | Ciliate, *Euplotes vannus* in Yantai Institute of Coastal Zone Research (Shandong, China) |
| | | MK801649 | 100.00 | Unspecified coral from South China Sea (Guangdong, China) |
| *Ruegeria* | ASV_006 | MK736137 | 100.00 | ***Acropora hemprichii*** (Central Red Sea) |
| | | MK736054 | 100.00 | ***Pocillopora verrucosa*** (Central Red Sea) |
| | | MH807604 | 100.00 | ***Galaxea fascicularis*** (Aquarium, Japan) |
| *Alteromonas* | ASV_008 | MT472680 | 100.00 | Shrimp gut, *Penaeus vannamei* (India) |
| | | MN704564 | 100.00 | Undescribed source at marine molecular ecology lab (Zhejiang, China) |
| | | MH556744 | 100.00 | Ciliate, *Euplotes vannus* in Yantai Institute of Coastal Zone Research (Shandong, China) |
| unclassified *Gammaproteobacteria* | ASV_017 | AB294979 | 99.61 | Microbial mat at shallow submarine hot spring (Okinawa, Japan) |
| | | KU629853 | 99.23 | Seaweed, *Asparagopsis* sp. (Portugal) |
| | | GU061986 | 99.23 | Oceanic water (South China Sea) |
| unclassified *Cryomorphaceae* | ASV_018 | KY374050 | 100.00 | ***Acropora hyacinthus*** (Central Red Sea) |
| | | KF185498 | 99.60 | Marine snow (Adriatic Sea) |
| | | KU648372 | 95.65 | Seawater (Gullmarsfjorden, Sweden) |
| unclassified *Hyphomonadaceae* | ASV_020 | KY455482 | 100.00 | ***Acropora hemprichii*** (Central Red Sea) |
| | | KY373570 | 100.00 | ***Acropora hyacinthus*** (Central Red Sea) |
| | | CP017718 | 100.00 | Algal culture, coral reef substrate (Palmyra Atoll, Pacific Ocean) |

## Discussion

276    We employed 16S rRNA gene amplicon sequencing to characterize the bacterial communities of symbiotic
277   and menthol-bleached *G. fascicularis* polyps from the Red Sea and Hong Kong that were long-term aquarium
278   reared in separate facilities.

**Menthol bleaching led to stochastic changes in the microbiome of Galaxea**

Menthol bleaching was associated with changes in the bacterial communities that differed between individual polyps and produced stochastic configurations. Unlike previous studies on Aiptasia that compared bleached and symbiotic individuals and found significantly different microbiomes (Röthig et al. 2016; Curtis et al. 2023), we could not identify a "signature of bleaching" as no bacterial taxa showed differential abundance between symbiotic states in our Galaxea colonies. This surprisingly included Symbiodiniaceae-associated bacteria that we were expecting to be reduced after the physical removal of Symbiodiniaceae (Fig. S5; Supplementary Materials and Methods). Such results could be an artifact of whole-tissue sampling in this study. As different coral compartments host distinct bacterial communities (Sweet et al. 2011; Ainsworth et al. 2015; Apprill et al. 2016), changes at the level of the gastroderm, where Symbiodiniaceae and their bacterial communities are located, might have been masked. Alternatively, it might indicate that Symbiodiniaceae-associated bacteria only repesent a small proportion of the community in Galaxea, or that they were able to persist in the absence of Symbiodiniaceae.

The stochastic response to bleaching aligns well with the concept of an obligatory nature of the coral-algal symbiosis. In facultatively symbiotic cnidarians (such as the anemone Aiptasia), symbiotic and bleached can constitute alternative stable states, and it is thus possible to identify distinct and characteristic symbiotic and bleached microbiome configurations (Röthig et al. 2016; Curtis et al. 2023). Corals however depend on their algal symbiont for energy and other essential metabolic processes (Muscatine and Porter 1977; Muscatine 1990), and therefore the bleached state is not a stable alternative to the symbiotic state. In the bleached state the weakened coral host becomes progressively unable to regulate its microbial community, leaving room for the establishment of opportunistic bacteria, producing novel and stochastic combinations (Zaneveld et al. 2017).

The response of the bacterial microbiome to menthol bleaching differed between facilities. For Red Sea corals, in the nMDS-ordination space on microbial community composition (Fig. 3), bleached polyps moved towards the center of the plot to the other symbiotic colonies rather than spreading in any random direction. This, taken together with the observation that alpha diversity did not increase after bleaching (as otherwise expected with dysbiosis (Zaneveld et al. 2017)), points towards a "captivity" effect. Since the polyps were maintained together in filtered seawater in closed systems, their exposure to novel bacteria was limited, and bacteria shed by the other polyps likely constituted the predominant source of "novel" associates. In contrast, the response of the Hong Kong corals to menthol bleaching was directional and more uniform. We hypothesize that this latter might reflect a new stable state of the tank water, rather than of the holobiont. While rearing conditions were largely replicated between facilities, feed type, tank volume and filtration systems differed. Feed can introduce bacteria into the system (Hartman et al. 2020), and uneaten portions could promote microbial growth. Such effects would have been amplified by the smaller volume of the containers used in Hong Kong compared to Red Sea. However, it should be noted that our experimental design did not allow us to directly test these hypotheses. We therefore suggest that future studies incorporate an adequately replicated "facility" factor in their design, as well as food and seawater samples to better characterize the influence of rearing conditions on the host microbiome.

**The microbiome of long-term aquarium-reared *Galaxea fascicularis***

The *G. fascicularis* polyps hosted simple bacterial microbiomes which were composed of a relatively small number of bacterial taxa (10-78 ASVs). Species richness in our aquarium-reared polyps was thus almost two orders of magnitude lower than those of wild *G. fascicularis* colonies sampled in the South China Sea (646-1,459 OTUs, Li et al. 2013) and those reported from a wide range of species and locations, which is typically in the order of 100s to 1000s of bacterial taxa (e.g., Ziegler et al. 2016; Hernandez-Agreda et al. 2018; Pollock et al. 2018; Galand et al. 2023). Due to the absence of direct comparison with wild colonies we are unable to draw conclusions on whether captivity caused a reduction in bacterial diversity. However, we suspect that captivity favoured a simplification of the microbiome, as stable and homogenous environmental conditions decrease both chances and need for the association with functionally and taxonomically diverse microbial partners. In fact, decreases in metabolic diversity and species richness have consistently been reported for tropical reef-

building corals reared in closed systems (Kooperman et al. 2007; Vega Thurber et al. 2009; Pratte et al. 2015;
Damjanovic et al. 2020), and for the anemone Aiptasia already after a few days of captivity (Hartman et al.
2020). Such effects may also have been exacerbated by the use of filtered seawater during the bleaching phase,
which largely reduced the pool of available microbes (Dungan et al. 2021b). Additionally, as colony morphology
is a major factor affecting coral microbial communities (Morrow et al. 2022), a loss of bacterial species might
also be ascribed to reduced structural complexity, where single polyps have a simpler geometry with fewer
micro-environments and ecological niches compared to larger colonies (Putnam et al. 2017).

Although some may see this simplification of the microbiome as a problematic artefact associated with
captive corals, simplified microbiomes present the opportunity to identify essential associates and facilitate
the development of microbial manipulation protocols to unravel holobiont functioning (Jaspers et al. 2019;
Puntin et al. 2022b). While the majority of studies report corals as hosting complex and rich microbial
communities, the key functional players still remain elusive (Jaspers et al. 2019; Barreto et al. 2021). Culturing
corals in sterile seawater may help to limit the horizontal acquisition of transient microbes and thus favor
proliferation of core or stable members for detailed characterization (Dungan et al. 2021b). Also, a simplified
microbiome facilitates further elimination of bacterial populations to produce gnotobiotic or axenic hosts,
which could subsequently be re-inoculated to produce a range of host-bacteria combinations to test microbial
functions and inter-partner dynamics (Fraune et al. 2015; Murillo-Rincon et al. 2017; Jaspers et al. 2019;
Taubenheim et al. 2020). Reduced microbial complexity in captivity might therefore provide advantages for
these specific experimental approaches with the Galaxea model.

Interestingly, the coral colonies tested here maintained distinct bacterial microbiomes even after long-term
co-culturing, which supports a degree of host genotype effects controlling the microbiome composition as
previously reported from Hydrozoan corals in the field (Dubé et al. 2021). Surprisingly, the microbiome of one
Red Sea colony was highly similar to that of Hong Kong colonies, despite large differences in geographic and
environmental origin (Wepfer et al. 2020). These colonies were also maintained in separate facilities, but
rearing conditions were similar at both locations. In addition to the host phylogenetic basis of microbiome
composition (Pollock et al. 2018), the similarity in environmental conditions may have induced convergence of
microbial community composition (Dubé et al. 2021).

Besides host genotype, Symbiodiniaceae community composition observed herein could also explain
differences in bacterial community composition between Red Sea colonies (Littman et al. 2010; Bernasconi et
al. 2019). While only a small proportion of samples were successfully sequenced, we could identify patterns of
Symbiodiniaceae-bacteria co-occurrence that warrant further investigation. We therefore recommend that
future studies characterize a larger number of Galaxea holobionts at multiple locations across the species
distribution range to explore links between host-Symbiodiniaceae-bacteria associations. This could elucidate
the influence of each member on coral holobiont compositions and functioning.

**Core bacterial associates of Galaxea**

We defined core or stable microbial associates based on prevalence of ASVs across treatment groups and
polyps. Notably, no taxa was present in 100 % of the samples, suggesting a certain degree of microbiome
variability within this coral host species. Among the 28 core ASVs (occuring in all groups), the five most frequent
and abundant were assigned to the genera *Alteromonas* and *Ruegeria*. Both *Alteromonas* and *Ruegeria* are
common coral-associates reported from at least 20 other coral species, and sequences assigned to these two
genera ranked 6[th] and 33[rd] most abundant in the Coral Microbial Database (Huggett and Apprill 2019).

*Ruegeria* spp. are commonly and consistently found in association with *G. fascicularis* in wild and aquarium-
reared colonies, from Hong Kong to Japan and across seasons (Cai et al. 2018a, 2018b; Miura et al. 2019; Tang
et al. 2020; Kitamura et al. 2021). Indeed, the two *Ruegeria* core ASVs had identical sequences with *Ruegeria*
from *G. fascicularis* from Hainan and Japan that were maintained under aquarium conditions comparable to
ours (Zhou et al. 2020, Miura et al. 2019). This shows that the Galaxea-*Ruegeria* association is highly conserved
and therefore putatively biologically relevant. *Ruegeria* strains isolated from *G. fascicularis* were indeed
previously identified as potential probiotics, through inhibitory activity towards the coral pathogen *Vibrio*

*coralliilyticus* (Kitamura et al. 2021). The ubiquitous and persistent Galaxea-*Ruegeria* association thus warrants attention in future investigations.

The potential role of *Alteromonas* spp. in the Galaxea holobiont functioning also deserves attention. Despite their high abundance and prevalence in this study (and in corals in general), the role of *Alteromonas* spp. remains controversial. They have been considered pathogenic, owing to their co-occurrence with coral diseases (e.g., Sunagawa et al. 2009), but also listed as candidate probiotic for their free radical scavenging abilities (Raina et al. 2009; Dungan et al. 2021a), which could be linked to their consistent association with Symbiodiniaceae (Lawson et al. 2018; Nitschke et al. 2020).

Bacteria in the family *Endozoicomonadaceae* are the most prominent members of coral microbiomes in a range of coral species (Morrow et al. 2012; Bayer et al. 2013; Neave et al. 2017; Pogoreutz et al. 2022), which have been investigated for their involvement in holobiont metabolism, for example in the C and S cycles (Neave et al. 2016; Ide et al. 2022). Yet, *Endozoicomonadaceae* were only present in approximately half of Galaxea polyps at between 0.06 and 39.12 % (mean 5.78 %) relative abundance. This is slightly lower than in wild *G. fascicularis* from Hong Kong (< 10 % relative abundance) in which, however, *Endozoicomonas* spp. were present in all samples (Cai et al. 2018a, 2018b). Such scarcity and inconsistent presence therefore suggests that *Endozoicomonas* spp. might not play an essential role in the *G. fascicularis* under captive conditions. If further proven, a lack of reliance on *Endozoicomonas* spp. in captivity could offer insights into which functions benefit from microbial help in the wild, hence highlight the role of this bacterial associate in the coral holobiont in the wider context.

## Conclusions

Model organisms provide powerful tools for unraveling holobiont complexity. These models can be used to test hypotheses of functional relationships and inter-partner interactions through holobiont manipulation. To complement current cnidarian model systems such as Aiptasia and Hydra, we recently proposed the adoption of *Galaxea fascicularis* as a true coral model owing to its suitability to aquarium rearing and reproduction, and manipulation of its association with Symbiodiniaceae following menthol bleaching (Puntin et al. 2022a). However, how this bleaching treatment affected the bacterial microbiome remained to be explored.

In this study, we provided the first baseline assessment of the response of the Galaxea bacterial microbiome to menthol bleaching, and gain initial insights into the potential effects of long-term captivity in this coral species. Overall, menthol bleaching induced stochastic changes in the microbiome, indicating dysbiosis. The response of the bacterial microbiome to bleaching differed between the two facilities, likely reflecting differences in rearing conditions, which remain to be addressed. Bacterial communities of our captive Galaxea colonies were composed of fewer taxa than reported for wild corals, which is in line with decreasing microbial diversity of many captive organisms. Nevertheless, symbiotic polyps originating from different colonies maintained distinct community assemblies, and showed links to host and/or Symbiodiniaceae identity which warrant further investigation.

A simplified microbiome could facilitate both characterization and experimental manipulation, and guide the identification of essential ("core") members among the retained associates. In this regard, we identified *Ruegeria* spp. and *Alteromonas* spp. as candidate associates for further functional interrogation. In conclusion, our study contributes valuable information towards a better characterization of the Galaxea holobiont, as well as its continued development and establishment as a coral model system.

## Acknowledgements

We thank André Dietzmann, Kara E. Engelhardt, Patrick Schubert, and Christina Anding (all JLU) for assistance during experiments; and Talia Morris and Hugh Spencer (Austrop) for offering their office and kindness, allowing GP to work on this manuscript from a very special place.

## Data, scripts, code, and supplementary information availability

## Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article.

## Funding

This study is part of the 'Ocean2100' global change simulation project of the Colombian-German Center of Excellence in Marine Sciences (CEMarin) funded by the German Academic Exchange Service (DAAD). This study is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Project number 469364832 (M.Z.) – SPP 2299/Project number 441832482.

## References

Agostini S, Suzuki Y, Higuchi T, Casareto BE, Yoshinaga K, Nakano Y, Fujimura H (2012) Biological and chemical characteristics of the coral gastric cavity. Coral Reefs 31:147–156 http://link.springer.com/10.1007/s00338-011-0831-6

Ainsworth TD, Krause L, Bridge T, Torda G, Raina J-B, Zakrzewski M, Gates RD, Padilla-Gamiño JL, Spalding HL, Smith C, Woolsey ES, Bourne DG, Bongaerts P, Hoegh-Guldberg O, Leggat W (2015) The coral core microbiome identifies rare bacterial taxa as ubiquitous endosymbionts. ISME J 9:2261–2274 https://www.nature.com/articles/ismej201539

Al-Horani FA, Al-Moghrabi SM, de Beer D (2003) Microsensor study of photosynthesis and calcification in the scleractinian coral, *Galaxea fascicularis*: active internal carbon cycle. J Exp Mar Bio Ecol 288:1–15 https://linkinghub.elsevier.com/retrieve/pii/S0022098102005786

Al-Horani FA, Ferdelman T, Al-Moghrabi SM, de Beer D (2005) Spatial distribution of calcification and photosynthesis in the scleractinian coral *Galaxea fascicularis*. Coral Reefs 24:173–180 http://link.springer.com/10.1007/s00338-004-0461-3

Al-Horani FA, Tambutté É, Allemand D (2007) Dark calcification and the daily rhythm of calcification in the scleractinian coral, Galaxea fascicularis. Coral Reefs 26:531–538 http://link.springer.com/10.1007/s00338-007-0250-x

Andersson AF, Lindberg M, Jakobsson H, Bäckhed F, Nyrén P, Engstrand L (2008) Comparative Analysis of Human Gut Microbiota by Barcoded Pyrosequencing. PLoS One 3:e2836 https://dx.plos.org/10.1371/journal.pone.0002836

Apprill A, Weber LG, Santoro AE (2016) Distinguishing between Microbial Habitats Unravels Ecological Complexity in Coral Microbiomes. mSystems 1:e00143-16 http://msystems.asm.org/lookup/doi/10.1128/mSystems.00143-16

Barreto MM, Ziegler M, Venn A, Tambutté E, Zoccola D, Tambutté S, Allemand D, Antony CP, Voolstra CR, Aranda M
(2021) Effects of Ocean Acidification on Resident and Active Microbial Communities of Stylophora pistillata.
Front Microbiol 12:1–13 https://www.frontiersin.org/articles/10.3389/fmicb.2021.707674/full

Battaglia T (2022) btools: A suite of R function for all types of microbial diversity analyses.
https://rdrr.io/github/twbattaglia/btools/

Bayer T, Neave MJ, Alsheikh-Hussain A, Aranda M, Yum LK, Mincer T, Hughen K, Apprill A, Voolstra CR (2013) The
Microbiome of the Red Sea Coral Stylophora pistillata Is Dominated by Tissue-Associated Endozoicomonas
Bacteria. Appl Environ Microbiol 79:4759–4762 https://journals.asm.org/doi/10.1128/AEM.00695-13

Bernasconi R, Stat M, Koenders A, Huggett MJ (2019) Global Networks of Symbiodinium-Bacteria Within the Coral
Holobiont. Microb Ecol 77:794–807 http://link.springer.com/10.1007/s00248-018-1255-4

Bokulich NA, Kaehler BD, Rideout JR, Dillon M, Bolyen E, Knight R, Huttley GA, Caporaso JG (2018) Optimizing
taxonomic classification of marker-gene amplicon sequences with QIIME 2's q2-feature-classifier plugin.
Microbiome 6:90 https://doi.org/10.1186/s40168-018-0470-z

Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, Alexander H, Alm EJ, Arumugam M, Asnicar
F, Bai Y, Bisanz JE, Bittinger K, Brejnrod A, Brislawn CJ, Brown CT, Callahan BJ, Caraballo-Rodríguez AM, Chase
J, Cope EK, Da Silva R, Diener C, Dorrestein PC, Douglas GM, Durall DM, Duvallet C, Edwardson CF, Ernst M,
Estaki M, Fouquier J, Gauglitz JM, Gibbons SM, Gibson DL, Gonzalez A, Gorlick K, Guo J, Hillmann B, Holmes S,
Holste H, Huttenhower C, Huttley GA, Janssen S, Jarmusch AK, Jiang L, Kaehler BD, Kang K Bin, Keefe CR, Keim
P, Kelley ST, Knights D, Koester I, Kosciolek T, Kreps J, Langille MGI, Lee J, Ley R, Liu Y-X, Loftfield E, Lozupone
C, Maher M, Marotz C, Martin BD, McDonald D, McIver LJ, Melnik A V, Metcalf JL, Morgan SC, Morton JT,
Naimey AT, Navas-Molina JA, Nothias LF, Orchanian SB, Pearson T, Peoples SL, Petras D, Preuss ML, Pruesse E,
Rasmussen LB, Rivers A, Robeson MS, Rosenthal P, Segata N, Shaffer M, Shiffer A, Sinha R, Song SJ, Spear JR,
Swafford AD, Thompson LR, Torres PJ, Trinh P, Tripathi A, Turnbaugh PJ, Ul-Hasan S, van der Hooft JJJ, Vargas
F, Vázquez-Baeza Y, Vogtmann E, von Hippel M, Walters W, Wan Y, Wang M, Warren J, Weber KC, Williamson
CHD, Willis AD, Xu ZZ, Zaneveld JR, Zhang Y, Zhu Q, Knight R, Caporaso JG (2019) Reproducible, interactive,
scalable and extensible microbiome data science using QIIME 2. Nat Biotechnol 37:852–857
https://doi.org/10.1038/s41587-019-0209-9

Bourne DG, Morrow KM, Webster NS (2016) Insights into the Coral Microbiome: Underpinning the Health and
Resilience of Reef Ecosystems. Annu Rev Microbiol 70:317–340
http://www.annualreviews.org/doi/10.1146/annurev-micro-102215-095440

Cai L, Tian RM, Zhou G, Tong H, Wong YH, Zhang W, Chui APY, Xie JY, Qiu JW, Ang PO, Liu S, Huang H, Qian PY (2018a)
Exploring coral microbiome assemblages in the South China Sea. Sci Rep 8:1–13
http://dx.doi.org/10.1038/s41598-018-20515-w

Cai L, Zhou G, Tong H, Tian R-M, Zhang W, Ding W, Liu S, Huang H, Qian P-Y (2018b) Season structures prokaryotic
partners but not algal symbionts in subtropical hard corals. Appl Microbiol Biotechnol 102:4963–4973
http://link.springer.com/10.1007/s00253-018-8909-5

Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP (2016) DADA2: High-resolution sample
inference from Illumina amplicon data. Nat Methods 13:581–583
https://www.nature.com/articles/nmeth.3869

Certner RH, Vollmer S V. (2018) Inhibiting bacterial quorum sensing arrests coral disease development and disease-
associated microbes. Environ Microbiol 20:645–657 https://onlinelibrary.wiley.com/doi/full/10.1111/1462-
2920.13991

Chakravarti LJ, Beltran VH, van Oppen MJH (2017) Rapid thermal adaptation in photosymbionts of reef-building

501 corals. Glob Chang Biol 23:4675–4688 https://onlinelibrary.wiley.com/doi/10.1111/gcb.13702

502 Chan WY, Rudd D, Meyers L, Topa SH, van Oppen MJH (2023) Heat-evolved algal symbionts enhance bleaching
503 tolerance of adult corals without trade-off against growth. bioRxiv
504 https://www.biorxiv.org/content/10.1101/2023.06.17.545444v1

505 Connelly MT, McRae CJ, Liu P-J, Martin CE, Traylor-Knowles N (2022) Antibiotics Alter Pocillopora Coral-
506 Symbiodiniaceae-Bacteria Interactions and Cause Microbial Dysbiosis During Heat Stress. Front Mar Sci 8:1–
507 18 https://www.frontiersin.org/articles/10.3389/fmars.2021.814124/full

508 Conway JR, Lex A, Gehlenborg N (2017) UpSetR: an R package for the visualization of intersecting sets and their
509 properties. Bioinformatics 33:2938–2940 https://doi.org/10.1093/bioinformatics/btx364

510 Costanza R, de Groot R, Sutton P, van der Ploeg S, Anderson SJ, Kubiszewski I, Farber S, Turner RK (2014) Changes in
511 the global value of ecosystem services. Glob Environ Chang 26:152–158
512 http://dx.doi.org/10.1016/j.gloenvcha.2014.04.002

513 Curtis E, Moseley J, Racicot R, Wright RM (2023) Bacterial microbiome variation across symbiotic states and clonal
514 lines in a cnidarian model. Front Mar Sci 10:1–9
515 https://www.frontiersin.org/articles/10.3389/fmars.2023.1113043/full

516 Damjanovic K, Blackall LL, Peplow LM, van Oppen MJH (2020) Assessment of bacterial community composition within
517 and among Acropora loripes colonies in the wild and in captivity. Coral Reefs 39:1245–1255
518 https://link.springer.com/10.1007/s00338-020-01958-y

519 Dani V, Priouzeau F, Pagnotta S, Carette D, Laugier JP, Sabourault C (2016) Thermal and menthol stress induce
520 different cellular events during sea anemone bleaching. Symbiosis 69:175–192
521 http://dx.doi.org/10.1007/s13199-016-0406-y

522 Dixon P (2003) VEGAN, a package of R functions for community ecology. J Veg Sci 14:927–930
523 https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1654-1103.2003.tb02228.x

524 Dubé CE, Ziegler M, Mercière A, Boissin E, Planes S, Bourmaud CA-F, Voolstra CR (2021) Naturally occurring fire coral
525 clones demonstrate a genetic and environmental basis of microbiome composition. Nat Commun 12:6402
526 https://www.nature.com/articles/s41467-021-26543-x

527 Dungan AM, Bulach D, Lin H, van Oppen MJH, Blackall LL (2021a) Development of a free radical scavenging bacterial
528 consortium to mitigate oxidative stress in cnidarians. Microb Biotechnol 14:2025–2040
529 https://onlinelibrary.wiley.com/doi/10.1111/1751-7915.13877

530 Dungan AM, van Oppen MJH, Blackall LL (2021b) Short-Term Exposure to Sterile Seawater Reduces Bacterial
531 Community Diversity in the Sea Anemone, Exaiptasia diaphana. Front Mar Sci 7:1–10
532 https://www.frontiersin.org/articles/10.3389/fmars.2020.599314/full

533 Eakin CM, Devotta D, Heron S, Geiger E, De La Cour J, Gomez A, Skirving W, Baird A, Cantin N, Donner S, Gilmour J
534 (2022) The 2014-17 Global Coral Bleaching Event: The Most Severe and Widespread Coral Reef Destruction.
535 Res Sq 1–24 https://www.researchsquare.com/article/rs-1555992/v1

536 Eddy TD, Lam VWY, Reygondeau G, Cisneros-Montemayor AM, Greer K, Palomares MLD, Bruno JF, Ota Y, Cheung
537 WWL (2021) Global decline in capacity of coral reefs to provide ecosystem services. One Earth 4:1278–1285
538 https://linkinghub.elsevier.com/retrieve/pii/S2590332221004747

539 Ferrier-Pagès C, Gattuso J, Cauwet G, Jaubert J, Allemand D (1998) Release of dissolved organic carbon and nitrogen
540 by the zooxanthellate coral Galaxea fascicularis. Mar Ecol Prog Ser 172:265–274 http://www.int-

res.com/abstracts/meps/v172/p265-274/

Fraune S, Anton-Erxleben F, Augustin R, Franzenburg S, Knop M, Schröder K, Willoweit-Ohl D, Bosch TCG (2015) Bacteria–bacteria interactions within the microbiota of the ancestral metazoan *Hydra* contribute to fungal resistance. ISME J 9:1543–1556 http://www.nature.com/articles/ismej2014239

Gabay Y, Parkinson JE, Wilkinson SP, Weis VM, Davy SK (2019) Inter-partner specificity limits the acquisition of thermotolerant symbionts in a model cnidarian-dinoflagellate symbiosis. ISME J 13:2489–2499 https://doi.org/10.1038/s41396-019-0429-5

Gabay Y, Weis VM, Davy SK (2018) Symbiont Identity Influences Patterns of Symbiosis Establishment, Host Growth, and Asexual Reproduction in a Model Cnidarian-Dinoflagellate Symbiosis. Biol Bull 234:1–10 http://www.journals.uchicago.edu/doi/10.1086/696365

Galand PE, Ruscheweyh H-J, Salazar G, Hochart C, Henry N, Hume BCC, Oliveira PH, Perdereau A, Labadie K, Belser C, Boissin E, Romac S, Poulain J, Bourdin G, Iwankow G, Moulin C, Armstrong EJ, Paz-García DA, Ziegler M, Agostini S, Banaigs B, Boss E, Bowler C, de Vargas C, Douville E, Flores M, Forcioli D, Furla P, Gilson E, Lombard F, Pesant S, Reynaud S, Thomas OP, Troublé R, Zoccola D, Voolstra CR, Thurber RV, Sunagawa S, Wincker P, Allemand D, Planes S (2023) Diversity of the Pacific Ocean coral reef microbiome. Nat Commun 14:3039 https://www.nature.com/articles/s41467-023-38500-x

Galliot B (2012) *Hydra*, a fruitful model system for 270 years. Int J Dev Biol 56:411–423 http://www.intjdevbiol.com/paper.php?doi=120086bg

Goulet TL, Erill I, Ascunce MS, Finley SJ, Javan GT (2020) Conceptualization of the Holobiont Paradigm as It Pertains to Corals. Front Physiol 11:1–8 https://www.frontiersin.org/article/10.3389/fphys.2020.566968/full

Hartman LM, van Oppen MJH, Blackall LL (2020) Microbiota characterization of Exaiptasia diaphana from the Great Barrier Reef. Anim Microbiome 2:10 https://animalmicrobiome.biomedcentral.com/articles/10.1186/s42523-020-00029-5

Hernandez-Agreda A, Leggat W, Bongaerts P, Herrera C, Ainsworth TD (2018) Rethinking the Coral Microbiome: Simplicity Exists within a Diverse Microbial Biosphere. MBio 9:1–14 https://journals.asm.org/doi/10.1128/mBio.00812-18

Herrera M, Klein SG, Schmidt-Roach S, Campana S, Cziesielski MJ, Chen JE, Duarte CM, Aranda M (2020) Unfamiliar partnerships limit cnidarian holobiont acclimation to warming. Glob Chang Biol 26:5539–5553 https://onlinelibrary.wiley.com/doi/10.1111/gcb.15263

van Hooidonk R, Maynard J, Tamelander J, Gove J, Ahmadia G, Raymundo L, Williams G, Heron SF, Planes S (2016) Local-scale projections of coral reef futures and implications of the Paris Agreement. Sci Rep 6:39666 http://www.nature.com/articles/srep39666

Huggett MJ, Apprill A (2019) Coral microbiome database: Integration of sequences reveals high diversity and relatedness of coral-associated microbes. Environ Microbiol Rep 11:372–385 https://onlinelibrary.wiley.com/doi/10.1111/1758-2229.12686

Hughes TP, Barnes ML, Bellwood DR, Cinner JE, Cumming GS, Jackson JBC, Kleypas J, van de Leemput IA, Lough JM, Morrison TH, Palumbi SR, van Nes EH, Scheffer M (2017) Coral reefs in the Anthropocene. Nature 546:82–90 https://www.nature.com/articles/nature22901

Hume BCC, Smith EG, Ziegler M, Warrington HJM, Burt JA, LaJeunesse TC, Wiedenmann J, Voolstra CR (2019) SymPortal: A novel analytical framework and platform for coral algal symbiont next-generation sequencing ITS2 profiling. Mol Ecol Resour 19:1063–1080 https://onlinelibrary.wiley.com/doi/10.1111/1755-0998.13004

Hume BCC, Ziegler M, Poulain J, Pochon X, Romac S, Boissin E, de Vargas C, Planes S, Wincker P, Voolstra CR (2018)
An improved primer set and amplification protocol with increased specificity and sensitivity targeting the
*Symbiodinium* ITS2 region. PeerJ 6:e4816 https://peerj.com/articles/4816

Ide K, Nishikawa Y, Maruyama T, Tsukada Y, Kogawa M, Takeda H, Ito H, Wagatsuma R, Miyaoka R, Nakano Y, Kinjo
K, Ito M, Hosokawa M, Yura K, Suda S, Takeyama H (2022) Targeted single-cell genomics reveals novel host
adaptation strategies of the symbiotic bacteria Endozoicomonas in Acropora tenuis coral. Microbiome 10:1–
16 https://microbiomejournal.biomedcentral.com/articles/10.1186/s40168-022-01395-9

Jaspers C, Fraune S, Arnold AE, Miller DJ, Bosch TCG, Voolstra CR (2019) Resolving structure and function of
metaorganisms through a holistic framework combining reductionist and integrative approaches. Zoology
133:81–87 https://doi.org/10.1016/j.zool.2019.02.007

Kaehler BD, Bokulich N, McDonald D, Knight R, Caporaso JG, Huttley GA (2019) Species abundance information
improves sequence taxonomy classification accuracy. Nat Commun 10:

Kitamura R, Miura N, Ito M, Takagi T, Yamashiro H, Nishikawa Y, Nishimura Y, Kobayashi K, Kataoka M (2021) Specific
Detection of Coral-Associated Ruegeria, a Potential Probiotic Bacterium, in Corals and Subtropical Seawater.
Mar Biotechnol 23:576–589 https://doi.org/10.1007/s10126-021-10047-2

Kooperman N, Ben-Dov E, Kramarsky-Winter E, Barak Z, Kushmaro A (2007) Coral mucus-associated bacterial
communities from natural and aquarium environments. FEMS Microbiol Lett 276:106–113
https://academic.oup.com/femsle/article-lookup/doi/10.1111/j.1574-6968.2007.00921.x

Lawson CA, Raina J, Kahlke T, Seymour JR, Suggett DJ (2018) Defining the core microbiome of the symbiotic
dinoflagellate, Symbiodinium. Environ Microbiol Rep 10:7–11
https://onlinelibrary.wiley.com/doi/10.1111/1758-2229.12599

Lex A, Gehlenborg N, Strobelt H, Vuillemot R, Pfister H (2014) UpSet: Visualization of Intersecting Sets. IEEE Trans Vis
Comput Graph 20:1983–1992 http://ieeexplore.ieee.org/document/6876017/

Li J, Chen Q, Zhang S, Huang H, Yang J, Tian X-P, Long L-J (2013) Highly Heterogeneous Bacterial Communities
Associated with the South China Sea Reef Corals Porites lutea, Galaxea fascicularis and Acropora millepora.
PLoS One 8:e71301 https://dx.plos.org/10.1371/journal.pone.0071301

Lin M-F, Takahashi S, Forêt S, Davy SK, Miller DJ (2019) Transcriptomic analyses highlight the likely metabolic
consequences of colonization of a cnidarian host by native or non-native Symbiodinium species. Biol Open
8:1–11 https://journals.biologists.com/bio/article/doi/10.1242/bio.038281/259291/Transcriptomic-analyses-
highlight-the-likely

Littman RA, Bourne DG, Willis BL (2010) Responses of coral-associated bacterial communities to heat stress differ
with Symbiodinium type on the same coral host. Mol Ecol 19:1978–1990
https://onlinelibrary.wiley.com/doi/10.1111/j.1365-294X.2010.04620.x

Matthews JL, Raina J, Kahlke T, Seymour JR, Oppen MJH, Suggett DJ (2020) Symbiodiniaceae-bacteria interactions:
rethinking metabolite exchange in reef-building corals as multi-partner metabolic networks. Environ Microbiol
22:1675–1687 https://onlinelibrary.wiley.com/doi/10.1111/1462-2920.14918

Matthews JL, Sproles AE, Oakley CA, Grossman AR, Weis VM, Davy SK (2015) Menthol-induced bleaching rapidly and
effectively provides experimental aposymbiotic sea anemones (*Aiptasia* sp.) for symbiosis investigations. J Exp
Biol 219:306–310 http://jeb.biologists.org/cgi/doi/10.1242/jeb.128934

McMurdie PJ, Holmes S (2013) phyloseq: An R package for reproducible interactive analysis and graphics of
microbiome census data. PLoS One 8:e61217 http://dx.plos.org/10.1371/journal.pone.0061217

Mikryukov V (2023) metagMisc: Miscellaneous functions for metagenomic analysis. https://rdrr.io/github/vmikk/metagMisc/

Miura N, Motone K, Takagi T, Aburaya S, Watanabe S, Aoki W, Ueda M (2019) Ruegeria sp. Strains Isolated from the Reef-Building Coral Galaxea fascicularis Inhibit Growth of the Temperature-Dependent Pathogen Vibrio coralliilyticus. Mar Biotechnol 21:1–8 http://link.springer.com/10.1007/s10126-018-9853-1

Morrow KM, Moss AG, Chadwick NE, Liles MR (2012) Bacterial Associates of Two Caribbean Coral Species Reveal Species-Specific Distribution and Geographic Variability. Appl Environ Microbiol 78:6438–6449 https://journals.asm.org/doi/10.1128/AEM.01162-12

Morrow KM, Pankey MS, Lesser MP (2022) Community structure of coral microbiomes is dependent on host morphology. Microbiome 10:113 https://doi.org/10.1186/s40168-022-01308-w

Motone K, Takagi T, Aburaya S, Miura N, Aoki W, Ueda M (2020) A Zeaxanthin-Producing Bacterium Isolated from the Algal Phycosphere Protects Coral Endosymbionts from Environmental Stress. MBio 11:1–13 https://journals.asm.org/doi/10.1128/mBio.01019-19

Murillo-Rincon AP, Klimovich A, Pemöller E, Taubenheim J, Mortzfeld B, Augustin R, Bosch TCG (2017) Spontaneous body contractions are modulated by the microbiome of Hydra. Sci Rep 7:15937 https://www.nature.com/articles/s41598-017-16191-x

Muscatine L (1990) The role of symbiotic algae in carbon and energy flux in reef corals. Ecosyst world 25:75–87

Muscatine L, Porter JW (1977) Reef Corals: Mutualistic Symbioses Adapted to Nutrient-Poor Environments. Bioscience 27:454–460 https://academic.oup.com/bioscience/article-lookup/doi/10.2307/1297526

Neave MJ, Apprill A, Ferrier-Pagès C, Voolstra CR (2016) Diversity and function of prevalent symbiotic marine bacteria in the genus Endozoicomonas. Appl Microbiol Biotechnol 100:8315–8324 http://dx.doi.org/10.1007/s00253-016-7777-0

Neave MJ, Rachmawati R, Xun L, Michell CT, Bourne DG, Apprill A, Voolstra CR (2017) Differential specificity between closely related corals and abundant Endozoicomonas endosymbionts across global scales. ISME J 11:186–200 http://dx.doi.org/10.1038/ismej.2016.95

Nitschke MR, Fidalgo C, Simões J, Brandão C, Alves A, Serôdio J, Frommlet JC (2020) Symbiolite formation: a powerful in vitro model to untangle the role of bacterial communities in the photosynthesis-induced formation of microbialites. ISME J 14:1533–1546 https://www.nature.com/articles/s41396-020-0629-z

Niu W, Huang H, Lin R, Chen C-H, Shen K-N, Hsiao C-D (2016) The complete mitogenome of the Galaxy Coral, *Galaxea fascicularis* (Cnidaria: Oculinidae). Mitochondrial DNA Part B 1:10–11 https://www.tandfonline.com/doi/full/10.1080/23802359.2015.1137796

Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin PR, O'Hara RB, Simpson GL, Solymos P, Stevens MHH, Szoecs E, Wagner H (2020) vegan: Community Ecology Package. https://cran.r-project.org/package=vegan

Peixoto RS, Sweet M, Villela HDM, Cardoso P, Thomas T, Voolstra CR, Høj L, Bourne DG (2021) Coral Probiotics: Premise, Promise, Prospects. Annu Rev Anim Biosci 9:265–288 https://www.annualreviews.org/doi/10.1146/annurev-animal-090120-115444

Pietschke C, Treitz C, Forêt S, Schultze A, Künzel S, Tholey A, Bosch TCG, Fraune S (2017) Host modification of a bacterial quorum-sensing signal induces a phenotypic switch in bacterial symbionts. Proc Natl Acad Sci 114:E8488–E8497 https://pnas.org/doi/full/10.1073/pnas.1706879114

Pogoreutz C, Oakley CA, Rädecker N, Cárdenas A, Perna G, Xiang N, Peng L, Davy SK, Ngugi DK, Voolstra CR (2022)
Coral holobiont cues prime Endozoicomonas for a symbiotic lifestyle. ISME J 16:1883–1895
https://www.nature.com/articles/s41396-022-01226-7

Pogoreutz C, Voolstra CR, Rädecker N, Weis V (2020) The coral holobiont highlights the dependence of cnidarian
animal hosts on their associated microbes. Cellular dialogues in the holobiont. pp 91–118

Pollock FJ, McMinds R, Smith S, Bourne DG, Willis BL, Medina M, Thurber RV, Zaneveld JR (2018) Coral-associated
bacteria demonstrate phylosymbiosis and cophylogeny. Nat Commun 9:4921
http://www.nature.com/articles/s41467-018-07275-x

Pratte ZA, Richardson LL, Mills DK (2015) Microbiota shifts in the surface mucopolysaccharide layer of corals
transferred from natural to aquaria settings. J Invertebr Pathol 125:
http://dx.doi.org/10.1016/j.jip.2014.12.009

Puntin G (2024) sPuntinG/Galaxea_Microbiome_PCI: Galaxea microbiome analysis 1st peer-review.
https://doi.org/10.5281/zenodo.10551928 https://doi.org/10.5281/zenodo.10551928

Puntin G, Craggs J, Hayden R, Engelhardt KE, McIlroy S, Sweet M, Baker DM, Ziegler M (2022a) The reef-building coral
Galaxea fascicularis: a new model system for coral symbiosis research. Coral Reefs 1–33
https://doi.org/10.1101/2022.06.02.494472

Puntin G, Sweet M, Fraune S, Medina M, Sharp K, Weis VM, Ziegler M (2022b) Harnessing the Power of Model
Organisms To Unravel Microbial Functions in the Coral Holobiont. Microbiol Mol Biol Rev
https://journals.asm.org/doi/10.1128/mmbr.00053-22

Putnam HM, Barott KL, Ainsworth TD, Gates RD (2017) The Vulnerability and Resilience of Reef-Building Corals. Curr
Biol 27:R528–R540 http://dx.doi.org/10.1016/j.cub.2017.04.047

R Core Team (2021) R: A Language and Environment for Statistical Computing. https://www.r-project.org/
https://www.r-project.org/

Raina J-B, Clode PL, Cheong S, Bougoure J, Kilburn MR, Reeder A, Forêt S, Stat M, Beltran V, Thomas-Hall P, Tapiolas
D, Motti CM, Gong B, Pernice M, Marjo CE, Seymour JR, Willis BL, Bourne DG (2017) Subcellular tracking reveals
the location of dimethylsulfoniopropionate in microalgae and visualises its uptake by marine bacteria. Elife
6:1–17 https://elifesciences.org/articles/23008

Raina J-B, Tapiolas D, Motti CA, Foret S, Seemann T, Tebben J, Willis BL, Bourne DG (2016) Isolation of an
antimicrobial compound produced by bacteria associated with reef-building corals. PeerJ 4:e2275
https://peerj.com/articles/2275

Raina J-B, Tapiolas D, Willis BL, Bourne DG (2009) Coral-Associated Bacteria and Their Role in the Biogeochemical
Cycling of Sulfur. Appl Environ Microbiol 75:3492–3501 https://journals.asm.org/doi/10.1128/AEM.02567-08

Robbins SJ, Singleton CM, Chan CX, Messer LF, Geers AU, Ying H, Baker A, Bell SC, Morrow KM, Ragan MA, Miller DJ,
Forêt S, Ball E, Beeden R, Berumen M, Aranda M, Ravasi T, Bongaerts P, Hoegh-Guldberg O, Cooke I, Leggat B,
Sprungala S, Fitzgerald A, Shang C, Lundgren P, Fyffe T, Rubino F, van Oppen M, Weynberg K, Robbins SJ,
Singleton CM, Xin Chan C, Messer LF, Geers AU, Ying H, Baker A, Bell SC, Morrow KM, Ragan MA, Miller DJ,
Foret S, Voolstra CR, Tyson GW, Bourne DG, Voolstra CR, Tyson GW, Bourne DG (2019) A genomic view of the
reef-building coral *Porites lutea* and its microbial symbionts. Nat Microbiol 4:2090–2100
https://www.nature.com/articles/s41564-019-0532-4

Robeson MS, O\textquoterightRourke DR, Kaehler BD, Ziemski M, Dillon MR, Foster JT, Bokulich NA (2020) RESCRIPt:
Reproducible sequence taxonomy reference database management for the masses. bioRxiv

https://www.biorxiv.org/content/early/2020/10/05/2020.10.05.326504

Rohwer F, Seguritan V, Azam F, Knowlton N (2002) Diversity and distribution of coral-associated bacteria. Mar Ecol
Prog Ser 243:1–10 http://www.int-res.com/abstracts/meps/v243/p1-10/

Rosado PM, Leite DCA, Duarte GAS, Chaloub RM, Jospin G, Nunes da Rocha U, P. Saraiva J, Dini-Andreote F, Eisen JA,
Bourne DG, Peixoto RS (2019) Marine probiotics: increasing coral resistance to bleaching through microbiome
manipulation. ISME J 13:921–936 http://www.nature.com/articles/s41396-018-0323-6

Rosenberg E, Koren O, Reshef L, Efrony R, Zilber-Rosenberg I (2007) The role of microorganisms in coral health,
disease and evolution. Nat Rev Microbiol 5:355–362 https://www.nature.com/articles/nrmicro1635

Röthig T, Costa RM, Simona F, Baumgarten S, Torres AF, Radhakrishnan A, Aranda M, Voolstra CR (2016) Distinct
bacterial communities associated with the coral model *Aiptasia* in aposymbiotic and symbiotic states with
*Symbiodinium*. Front Mar Sci 3: http://journal.frontiersin.org/article/10.3389/fmars.2016.00234/full

Röthig T, Puntin G, Wong JCY, Burian A, McLeod W, Baker DM (2021) Holobiont nitrogen control and its potential for
eutrophication resistance in an obligate photosymbiotic jellyfish. Microbiome 9:127
https://microbiomejournal.biomedcentral.com/articles/10.1186/s40168-021-01075-0

Scharfenstein HJ, Chan WY, Buerger P, Humphrey C, van Oppen MJH (2022) Evidence for de novo acquisition of
microalgal symbionts by bleached adult corals. ISME J 16:1676–1679
https://www.nature.com/articles/s41396-022-01203-0

Schubert P, Wilke T (2018) Coral Microcosms: Challenges and Opportunities for Global Change Biology. Corals in a
Changing World. InTech, pp 143–175 http://www.intechopen.com/books/corals-in-a-changing-world/coral-
microcosms-challenges-and-opportunities-for-global-change-biology

Sunagawa S, DeSantis TZ, Piceno YM, Brodie EL, DeSalvo MK, Voolstra CR, Weil E, Andersen GL, Medina M (2009)
Bacterial diversity and White Plague Disease-associated community changes in the Caribbean coral
Montastraea faveolata. ISME J 3:512–521 http://www.nature.com/articles/ismej2008131

Sweet MJ, Croquer A, Bythell JC (2011) Bacterial assemblages differ between compartments within the coral
holobiont. Coral Reefs 30:39–52 http://link.springer.com/10.1007/s00338-010-0695-1

Tandon K, Lu C-Y, Chiang P-W, Wada N, Yang S-H, Chan Y-F, Chen P-Y, Chang H-Y, Chiou Y-J, Chou M-S, Chen W-M,
Tang S-L (2020) Comparative genomics: Dominant coral-bacterium Endozoicomonas acroporae metabolizes
dimethylsulfoniopropionate (DMSP). ISME J 14:1290–1303 http://dx.doi.org/10.1038/s41396-020-0610-x

Tang K, Zhan W, Zhou Y, Xu T, Chen X, Wang W, Zeng Z, Wang Y, Wang X (2020) Antagonism between coral pathogen
Vibrio coralliilyticus and other bacteria in the gastric cavity of scleractinian coral Galaxea fascicularis. Sci China
Earth Sci 63:157–166

Taubenheim J, Willoweit-Ohl D, Knop M, Franzenburg S, He J, Bosch TCG, Fraune S (2020) Bacteria- and temperature-
regulated peptides modulate β-catenin signaling in *Hydra*. Proc Natl Acad Sci U S A 117:21459–21468
www.pnas.org/cgi/doi/10.1073/pnas.2010945117

Tebben J, Motti CA, Siboni N, Tapiolas DM, Negri AP, Schupp PJ, Kitamura M, Hatta M, Steinberg PD, Harder T (2015)
Chemical mediation of coral larval settlement by crustose coralline algae. Sci Rep 5:10803
http://www.nature.com/articles/srep10803

Thompson JR, Rivera HE, Closek CJ, Medina M (2015) Microbes in the coral holobiont: partners through evolution,
development, and ecological interactions. Front Cell Infect Microbiol 4:
http://journal.frontiersin.org/article/10.3389/fcimb.2014.00176/abstract

Vega Thurber R, Willner-Hall D, Rodriguez-Mueller B, Desnues C, Edwards RA, Angly F, Dinsdale E, Kelly L, Rohwer F
(2009) Metagenomic analysis of stressed coral holobionts. Environ Microbiol 11:2148–2163

Wang J-T, Chen Y-Y, Tew KS, Meng P-J, Chen CA (2012) Physiological and biochemical performances of menthol-
induced aposymbiotic corals. PLoS One 7:e46406 https://dx.plos.org/10.1371/journal.pone.0046406

Wang JT, Wang YT, Keshavmurthy S, Meng PJ, Chen CA (2019) The coral *Platygyra verweyi* exhibits local adaptation
to long-term thermal stress through host-specific physiological and enzymatic response. Sci Rep 9:1–11
http://dx.doi.org/10.1038/s41598-019-49594-z

Webster NS, Smith LD, Heyward AJ, Watts JEM, Webb RI, Blackall LL, Negri AP (2004) Metamorphosis of a
Scleractinian Coral in Response to Microbial Biofilms. Appl Environ Microbiol 70:1213–1221

Weis VM (2008) Cellular mechanisms of Cnidarian bleaching: stress causes the collapse of symbiosis. J Exp Biol
211:3059–66 http://jeb.biologists.org/content/211/19/3059.abstract

Weis VM, Davy SK, Hoegh-Guldberg O, Rodriguez-Lanetty M, Pringle JR (2008) Cell biology in model systems as the
key        to        understanding        corals.        Trends        Ecol        Evol        23:369–376
https://linkinghub.elsevier.com/retrieve/pii/S0169534708001614

Wickham H (2016) Ggplot2: Elegant graphics for data analysis. Springer International Publishing, Cham, Switzerland
https://books.google.de/books?id=RTMFswEACAAJ

Yilmaz P, Parfrey LW, Yarza P, Gerken J, Pruesse E, Quast C, Schweer T, Peplies J, Ludwig W, Glöckner FO (2013) The
SILVA and "All-species Living Tree Project (LTP)" taxonomic frameworks. Nucleic Acids Res 42:D643–D648
https://doi.org/10.1093/nar/gkt1209

Zaneveld JR, McMinds R, Vega Thurber R (2017) Stress and stability: applying the Anna Karenina principle to animal
microbiomes. Nat Microbiol 2:17121 http://www.nature.com/articles/nmicrobiol2017121

Zhou Y, Tang K, Wang P, Wang W, Wang Y, Wang X (2020) Identification of bacteria-derived urease in the coral gastric
cavity. Sci China Earth Sci 63:1553–1563 https://link.springer.com/10.1007/s11430-020-9647-x

Ziegler M, Roder C, Büchel C, Voolstra C (2015) Niche acclimatization in Red Sea corals is dependent on flexibility of
host-symbiont        association.        Mar        Ecol        Prog        Ser        533:149–161        http://www.int-
res.com/abstracts/meps/v533/p149-161/

Ziegler M, Roik A, Porter A, Zubier K, Mudarris MS, Ormond R, Voolstra CR (2016) Coral microbial community
dynamics in response to anthropogenic impacts near a major city in the central Red Sea. Mar Pollut Bull
105:629–640 https://linkinghub.elsevier.com/retrieve/pii/S0025326X15302496

773