1 SUPREM: an engineered non-site-specific m^{6}A RNA 2 methyltransferase with highly improved efficiency

4 Yoshiki Ochiai[1], Ben E. Clifton[1], Madeleine Le Coz[2], Marco Terenzio[2], Paola Laurino[1]*

6 [1] Protein Engineering and Evolution Unit, Okinawa Institute of Science and Technology 7 Graduate University (OIST), Okinawa 904-0495, Japan.

8 [2] Molecular Neuroscience Unit, Okinawa Institute of Science and Technology Graduate 9 University (OIST), Okinawa 904-0495, Japan.

11 *Corresponding author: Paola Laurino

12 Email: paola.laurino@oist.jp

Abstract

15 m^{6}A RNA methylation plays a key role in RNA processing and translational regulation, 16 influencing both normal physiological and pathological processes. Yet, current techniques for 17 studying RNA methylation struggle to isolate the effects of individual m^{6}A modifications. 18 Engineering of RNA methyltransferases (RNMTs) could enable development of improved 19 synthetic biology tools to manipulate RNA methylation, but is challenging due to limited 20 understanding of structure-function relationships in RNMTs. Herein, using ancestral sequence 21 reconstruction we explore the sequence space of the bacterial DNA methyltransferase EcoGII 22 (M.EcoGII), a promising target for protein engineering due to its lack of sequence specificity 23 and its residual activity on RNA. We thereby created an efficient non-specific RNMT termed 24 SUPREM, which exhibits 8-fold higher expression levels, 7 °C higher thermostability, and 12- 25 fold greater m^{6}A RNA methylation activity compared with M.EcoGII. Immunofluorescent 26 staining confirmed SUPREM's higher RNA methylation activity compared with M.EcoGII in 27 mammalian cells. Additionally, Nanopore direct RNA sequencing highlighted that SUPREM 28 is capable of methylating a larger number of RNA methylation sites than M.EcoGII. Through 29 phylogenetic and mutational analysis, we identified a critical residue for the enhanced RNA 30 methylation activity of SUPREM. Collectively, our findings indicate that SUPREM holds 31 promise as a versatile tool for *in vivo* RNA methylation and labeling.

Introduction

RNA methyltransferases (RNMTs) are enzymes that transfer a methyl group from S-adenosyl methionine (SAM) to their RNA substrates, thus contributing to the complex biological regulation of RNAs. In eukaryotes, these enzymes have an important role in methylation of N^6-methyladenine (m^{6}A), which is the most prominent internal modification in eukaryotic mRNA (1). Several eukaryotic m^{6}A RNMTs have been well characterized, and methylate mRNA and non-coding RNA molecules at a specific recognition sequence motif (2, 3). The m^{6}A modification is recognized by m^{6}A binding proteins, which thereby regulate RNA metabolism (4–6), pre-mRNA splicing (7–9), mRNA localization (10, 11), and translation (12–14). The m^{6}A modification is involved in development (15, 16) and stress response (17, 18), and is also linked to diseases such as cancer (12, 19, 20), infectious diseases (21), and type II diabetes (22). Thus, m^{6}A modifications are critical for both physiological and pathological processes. However, previous research on the biological functions of m^{6}A has largely relied on the knockout or knockdown of m^{6}A RNMTs and other m^{6}A-associated proteins, which results in a global disruption of RNA m^{6}A modification states. Consequently, dissecting the biological consequences of specific m^{6}A modifications and distinguish the direct and indirect effects of m^{6}A dysregulation has remained a challenge (23, 24). Thus, there is a need for tools that allow targeted manipulation of individual m^{6}A modifications in living cells to investigate their biological functions.

Few tools for site-specific m^{6}A methylation and demethylation have been reported previously (25–27). These tools are based on fusion constructs of the eukaryotic m^{6}A RNMT METTL3 with RNA-binding proteins, such as dead Cas proteins (dCas9 or dCas13) (25, 26) or PUF proteins (27). Dead Cas proteins can be targeted to specific sites in mRNA using guide RNA, while the RNA sequence specificity of PUF proteins is programmable by substitutions at specific residues. However, these METTL3-based tools have three important limitations. First, the range of sequences that can be methylated is restricted by the sequence specificity of METTL3, preventing the methylation of non-coding RNA and non-canonical methylation sites in mRNA. Second, since METTL3 works as part of a large methyltransferase complex (28), the isolated protein has impaired methylation efficiency (26). Third, the use of METTL3-based m^{6}A editing potentially introduces unwanted and confounding interactions with cellular proteins because METTL3 is an endogenous protein in eukaryotic cells. Altogether, an optimal RNMT for m^{6}A manipulation tools would be an exogenous enzyme with relaxed RNA sequence specificity that does not rely on other proteins.

The bacterial methyltransferase M.EcoGII is a promising candidate for an RNMT that could fulfill these requirements. M.EcoGII was identified as a DNA methyltransferase from a prophage region of a pathogenic *Escherichia coli* strain (29), and is capable of performing m^6A methylation of both DNA (~90% of adenine bases *in vitro; ~85% in vivo*) and RNA (~30% of adenine bases *in vitro*) (30). Unusually, M.EcoGII exhibits non-site-specific methylation activity and can potentially methylate any adenine base in DNA and RNA (29). Additionally, in contrast to mammalian methyltransferase complexes, M.EcoGII works independently of any other proteins, making it easier to engineer and orthogonal to cellular systems. Indeed, the non-specific DNA methyltransferase activity of M.EcoGII has been utilized to investigate chromatin structures in mammalian cells (31, 32).

Due to M.EcoGII's low RNA methyltransferase activity, expression level, and solubility, further protein engineering would be required to develop this enzyme as a research tool to manipulate RNA methylation states. However, protein engineering of RNMTs, including M.EcoGII, presents several challenges. In the case of M.EcoGII, no experimental structure has been reported and the protein exhibits low homology to other protein structures, especially in the RNA recognition domain. Significant conformational changes that are expected to occur upon nucleic acid binding decrease the reliability of structural prediction tools such as AlphaFold2 for improving the function of the protein. On the other hand, screening or directed evolution of RNMTs is also impractical due to the absence of efficient high-throughput screening techniques for measuring RNA methylation activity. Based on these limitations, we hypothesized that sequence-based engineering tools such as ancestral sequence reconstruction (ASR) would be most effective for improving the properties of M.EcoGII. ASR uses the sequences of extant proteins to statistically infer the sequences of extinct ancestral proteins, which often show higher thermostability and solubility and expanded substrate specificity compared to existing proteins (33–36). Although the underlying reasons for the improved properties of reconstructed ancestral proteins are not entirely clear (37, 38), ASR has been successfully applied to improve the properties of various proteins, including those of relatively recent evolutionary origin (39, 40). As a protein engineering technique, ASR allows for the sampling of sequence space enriched in proteins with favorable properties based on phylogenetic information (37).

Here, we used ASR to engineer ancestral M.EcoGII variants with higher expression, thermostability, and RNA methylation activity. We reconstructed five ancestral M.EcoGII variants, and one of these variants, which we named SUPer RNA EcoGII Methyltransferase (SUPREM), showed a 12-fold increase in RNA methylation activity and an 8-fold increase in

expression compared with M.EcoGII. SUPREM activity was confirmed in an *in vivo* context

via transfection in mammalian cells, suggesting a potential use for this enzyme in biological

investigations. Nanopore direct RNA sequencing analysis revealed that SUPREM had less

biased methylation on RNA than M.EcoGII. We also investigated the amino acid residues

responsible for the improved RNA methylation activity of SUPREM by site-directed

mutagenesis. Together, our engineered M.EcoGII variant has great potential for developing

novel RNA biology tools to manipulate RNA methylation.

Results

Ancestral sequence reconstruction of M.EcoGII leads to variants with higher stability

To reconstruct the sequences of ancestral M.EcoGII variants, we first collected a non-

redundant set of homologous protein sequences of β-class methyltransferases, including

M.EcoGII, from the Restriction Enzyme Database (REBASE) (41). A sequence similarity

network of the resulting protein sequences showed that M.EcoGII and highly similar sequences

clustered most closely with the site-specific DNA methyltransferase, M.PflMI (27.43%

identity), followed by two sequence clusters exemplified by M.McrMPORF631P (21.63%

identity) and M.Blo16BORF1919P (20.70% identity) (Supplementary Figure S1). For

subsequent phylogenetic analysis, we focused on the 202 sequences that clustered together with

these four sequences in the sequence similarity network. Analysis using the PHAST database

showed that all genes of these proteins are encoded in phage-derivative regions of their

respective host genomes (42).

A maximum-likelihood tree was constructed from these 202 sequences (Figure 1A:

schematic tree, Supplementary Figure S2: full tree). Because M.PflMI is the most closely

related enzyme to M.EcoGII that has confirmed site-specific methylation activity (recognition

sequence: CCANNNNNTGG), we postulated that non-site-specific methylation activity

emerged between M.EcoGII and the last common ancestor of M.EcoGII and M.PflMI, and that

ancestral proteins between these two nodes might retain the desired non-site-specific

methylation activity. We therefore selected six ancestral nodes, evenly spaced between

M.EcoGII and the last common ancestor of M.EcoGII and M.PflMI, for experimental

characterization (Figure 1A). The selected nodes were robustly supported by ultrahigh

bootstrap values (>90%), and the Shimodaira-Hasegawa approximate likelihood ratio test (SH-

aLRT) (>80%), and the corresponding ancestral sequences were also reconstructed with high

statistical confidence, with mean posterior probability values (>90%) (Supplementary Table
S1).

These ancestral M.EcoGII sequences were cloned into the pET-28a(+) vector and
expressed in the *E. coli* SHuffle T7 strain. Five of the six ancestral proteins could be expressed
and purified. The ancestral proteins were expressed at higher levels than M.EcoGII (6-fold to
13-fold; Supplementary Figure S3A, Supplementary Table S1). In addition, the ancestral
proteins showed higher thermostability than M.EcoGII, as measured by differential scanning
fluorimetry. Whereas M.EcoGII exhibited a denaturation temperature (T_M) of 43 °C and would
therefore have marginal stability at physiological temperatures, the ancestral proteins showed
increases in T_M of 7–16 °C compared with M.EcoGII (Supplementary Figure S3B-C). Thus,
ASR succeeded in improving the expression level and thermostability of M.EcoGII.

***In vitro* activity assays of ancestral M.EcoGII variants show improved catalytic efficiency**
for RNA methylation

To investigate the DNA and RNA methylation activity of M.EcoGII and the ancestral
proteins, we used the MTase-Glo assay (43), a luminescence-based assay that can be used to
quantify *S*-adenosyl homocysteine (SAH), a byproduct of the SAM-dependent methylation
reaction. PCR-amplified DNA (3.5 kbp, 1841 adenine bases) or *in vitro* transcribed RNA (1.8
knt, 544 adenine bases) was used as a substrate for *in vitro* methylation reactions. All the
ancestral proteins showed DNA and RNA methylation activity. There was limited variation in
the DNA methyltransferase activities of the ancestral proteins, with two of them, Anc291 and
Anc289, showing a small but significant increase in DNA methylation activity compared with
M.EcoGII (Anc291, 2.4-fold, $P = 0.0099$; Anc289, 2.4-fold, $P = 0.0139$) (Figure 1B).
Surprisingly, a much larger variation in RNA methyltransferase activity was observed, with
one ancestral protein, Anc284, showing a 12-fold increase in RNA methylation activity
compared with M.EcoGII ($P < 0.0001$; Figure 1C). Time-course DNA and RNA methylation
assays confirmed that the product formation was linear over the course of these experiments
and that Anc291 and Anc284 showed higher initial velocity for RNA methylation than
M.EcoGII (Figure 1D, Supplementary Figure S4, Supplementary Table S3).

To detect and quantify the m6A modification directly, we next performed LC-MS
analysis. A short RNA substrate (294 bases) was methylated with M.EcoGII or Anc284,
digested to nucleosides, and analyzed by LC-MS. Treatment of RNA with M.EcoGII and
Anc284 resulted in the formation of m6A, as shown by comparison of peak elution volume
with an m6A standard and detection of the m6A $[M+H]^+$ ion by MS (Supplementary Figure S5),

indicating the presence of RNA m^6A methylation activity in Anc284. Furthermore, the fraction

of m^6A-methylated adenosine determined by peak area integration was significantly higher for

Anc284, compared with M.EcoGII ($35.2 \pm 3.3\%$ vs. 72.4 ± 0.1 %; $P = 0.0001$ by unpaired t-

test). Altogether, these results show that Anc284 has substantially higher RNA methylation

activity than M.EcoGII. Thus, we coined Anc284 as <u>SUP</u>er <u>R</u>na <u>E</u>cogii <u>M</u>ethyltransferase

(SUPREM).

Methylation activity of SUPREM in human cells

To check that M.EcoGII and SUPREM are able to methylate RNA in human cells,

M.EcoGII-eGFP, SUPREM-eGFP, and eGFP control constructs were transfected in HEK 293T

cells. To quantify the exogenous methyltransferase activity, m^6A methylation efficiency was

analyzed by immunofluorescence. M.EcoGII expression in cells only showed a trend towards

a 2.8-fold increased m^6A fluorescence in transfected cells by compared to eGFP control, which

was not significant (Figure 2A&B, Supplementary Figure S6), while SUPREM expression

resulted in a 5.4-fold increase, which was significant compared to both eGFP control and

M.EcoGII (Figure 2A&B, Supplementary Figure S6). Compared to the data obtained *in vitro*,

while both M.EcoGII and SUPREM were able to methylate RNA, M.EcoGII activity in cells

was relatively low compared to SUPREM, which exhibited a more robust methylation activity.

Thus, we have successfully engineered an RNMT that could be exploited for biological studies.

The non-specificity in the methylation motifs is retained in SUPREM

A decrease in the site-specificity of RNMTs is advantageous for engineering RNA

methylation states in that more RNA sequences can be targeted. M.EcoGII showed non-site-

specific methylation activity for DNA (29), while the specificity for RNA has not been

explored so far. To investigate the presence of non-site-specific RNA methylation activity by

M.EcoGII and ancestral proteins, we performed Oxford Nanopore direct RNA sequencing

(DRS) to characterize the site specificity of RNA methylation in M.EcoGII, Anc291, and

SUPREM. The RNA substrate (the same substrate used for MTase-Glo methylation assays;

1872 nt, 1468 adenine-containing 5-mer motifs) was methylated using each methyltransferase

and sequenced using an Oxford Nanopore MinION device. The Nanopolish and Nanocompore

pipeline was then used to identify modification sites (specifically, 5-mer nucleotide sequences

containing at least one m^6A modification) through comparison with an unmethylated control

(Figure 3A) (44).

SUPREM methylated a greater number of diverse motifs (523 motifs of adenine-
containing motifs / 623 motifs of total detected motifs; false positive rate (FPR): 16%) than
Anc291 (389 motifs / 472 motifs; FPR: 18%) and M.EcoGII (225 motifs / 247 motifs; FPR:
9%) (Figure 3A, B). Thirty-six percent of adenine-containing 5-mer motifs contained the m^6A
modification for SUPREM, compared to Anc291's 26% and M.EcoGII's 15%. Analysis of the
5-mer sequences modified by each methyltransferase showed that the methyltransferase
activity of M.EcoGII was largely non-site-specific, but displayed a slight bias against GC-rich
sequences (Figure 3B). On the other hand, this bias was reduced in SUPREM, with the
frequency of each base more similar to the expected value if all adenine bases were methylated
equally (Figure 3B). Thus, our Oxford Nanopore DRS analysis revealed the presence of a
certain degree of bias in M.EcoGII's methylation activity for RNA, while SUPREM showed
less bias against GC-rich motifs in its methylation activity on RNA.

Investigation of key residues for improved RNA methyltransferase activity

To investigate which amino acid residues are important for RNA methylation activity,
we compared multiple sequence alignments of sequences in the M.EcoGII clade (clade I) and
SUPREM clade (clade II) in our maximum-likelihood phylogenetic tree (Figure 4A). We chose
14 amino acids as candidate residues that might be responsible for the observed differences in
RNA methylation activity based on two conditions, being that the residues were highly
conserved among each clade, and that the residues differed between clades I and II. Fourteen
single-point mutants of SUPREM were constructed, of which 12 variants could be expressed,
and 11 variants could be successfully purified (the amino acid sequences of these variants are
given in Supplementary Table S4). Using the MTase-Glo assay to measure RNA methylation
activity, one substitution, E179Q, was found to significantly decrease the RNA methylation
activity of SUPREM (3.4-fold decrease, $P = 0.0079$) (Figure 4B), suggesting that this residue
is important for RNA binding and methylation.

Structural analysis of M.EcoGII and its ancestral variants

To gain a greater understanding of the structural basis for the non-site-specific
methylation activity of M.EcoGII and the differences in DNA and RNA methylation activity
between M.EcoGII and its ancestral variants, we aimed to generate structural models of
M.EcoGII, Anc291, and SUPREM. To date, experimental structures of M.EcoGII have not
been determined. We performed extensive crystallization screening of M.EcoGII, SUPREM,
and Anc291, but no crystals were obtained, most likely due to the conformational flexibility of

the long interdomain linker. Homology modeling was unfeasible because of the low sequence
identity (<20%) of M.EcoGII to proteins of known structure. Indeed, although the structure of
M.EcoGII was previously modeled using the structure of M.EcoP15I (PDB 4ZCF) as a
template, M.EcoP15I bears no recognizable sequence homology to M.EcoGII in the target
recognition domain (45). Based on these considerations, we decided to use AlphaFold2 for
structural modeling.

To this end, we first aimed to determine the oligomeric state of M.EcoGII and the
ancestral proteins. Although previous research suggested that M.EcoGII is a monomeric
protein based on native-mass spectrometry experiments (30), previously characterized β-class
methyltransferases have shown a dimeric structure (45–47), and monomeric structures appear
to be incompatible with catalytic activity (48). We used size-exclusion chromatography (SEC)
to estimate the molecular weight of M.EcoGII and the ancestral proteins based on a standard
curve and found that each protein eluted as a dimer (Figure 4C, Supplementary Figure S7).

Finally, structural models of M.EcoGII and ancestral protein homodimers were
constructed using AlphaFold2 multimer modeling (Figure 4D, Supplementary Figure S8) (49).
According to the structural prediction, each M.EcoGII monomer is composed of a Rossmann
fold catalytic domain (residues 1–134 and 260–349, mean pLDDT score 95.8) and a four-helix
bundle target recognition domain (residues 160–237, mean pLDDT score 94.4) containing a
helix-turn-helix motif. The two domains are connected by a long, flexible, and highly charged
linker, partly composed of a two-helix structure (mean pLDDT score 89.1 for the predicted
helical portion). A search of the Protein Data Bank for similar structures using the DALI server
showed that the predicted structure of the Rossmann fold domain of M.EcoGII shows the
highest similarity to other DNA methyltransferases such as M.MboIIA and M.CcrMI. In
contrast, the target recognition domain bears similarity to various helix-turn-helix domain
DNA-binding proteins (e.g., transcriptional regulators), but generally not to known
methyltransferase structures (Supplementary Table S5). The overall predicted structure of the
site-specific DNA methyltransferase M.PflMI, including its target recognition domain, is
similar to M.EcoGII, with the exception of the highly charged linker region, which is absent in
M.PflMI (Supplementary Figure S8E). The large positively charged surface of M.EcoGII
spanning the target recognition domain and interdomain linker is likely important for the non-
specific binding of DNA and RNA (Figure 4E), providing a large interaction surface that
effectively compensates for the lack of protein-DNA and protein-RNA interactions with
specific bases.

The ancestral proteins showed a similar overall predicted structure to M.EcoGII
(Supplementary Figure S8A). The E179 residue identified as being important for RNA binding
and/or methylation is located on the edge of the putative DNA/RNA binding pocket, based on
the structural models predicted by AlphaFold2 (Figure 4D). In addition, the amino acids (97-
110, 116-140) present on the interaction surface between the subunits are completely conserved
among M.EcoGII and ancestral proteins, suggesting that the dimer interface might be crucial
for these enzymes. Co-crystal structures of SUPREM and RNA would be required for a
detailed understanding of the mechanisms underlying the improved RNA methylation activity
and promiscuity compared with M.EcoGII.

Discussion

Growing recognition of the importance of RNA methylation, particularly m^{6}A
methylation, in a broad range of physiological and pathological processes has driven demand
for new tools for the manipulation and analysis of RNA methylation states (23, 24). M.EcoGII
was identified as a potentially useful tool to manipulate m^{6}A methylation in mammalian cells
based on its non-site-specific DNA/RNA methyltransferase activity and its non-mammalian
origin, but the enzyme has not yet been utilized for this purpose due to its low expression and
RNA methylation activity. In this work, we used ASR to engineer an M.EcoGII variant,
SUPREM, with a number of advantages over the wild-type enzyme, including a 12-fold
increase in RNA methylation activity, 8 °C increase in thermostability, an 8-fold increase in
expression, and highly relaxed sequence specificity, resulting in methylation of up to 72% of
adenine bases in single strand RNA. M.EcoGII and other RNMTs are generally challenging
targets for other protein engineering techniques such as rational design or directed evolution;
for example, the lack of experimental structural information on M.EcoGII and the complexity
of structure-function relationships in nucleic acid enzymes would have limited the potential of
rational protein engineering in this case. A computational approach for the engineering of
RNMT-RNA interaction would be challenging due to difficulty in predicting interactions
between proteins and RNA. In contrast, our work shows the power of sequence-based
engineering approaches such as ASR to enable rapid improvement of a broad range of protein
properties, including challenging cases where structural information is limited.

All of the ancestral variants of M.EcoGII characterized in this work showed higher
thermostability and expression than the wild-type protein, with ΔT_m between 7 and 16 ºC. It is
noteworthy that major improvements in thermostability were achieved despite the limited

phylogenetic diversity of the sequences in the M.EcoGII clade, which are mainly derived from prophage regions in Enterobacteria genomes. Previous studies that have used ASR for protein engineering have typically focused on more phylogenetically diverse and evolutionarily ancient protein families (50), based on the hypothesis that the ancient progenitors of modern proteins existed in thermophilic organisms and were therefore thermostable. In the case of M.EcoGII, however, there is little reason to assume *a priori* that there would be a trend in thermostability from the ancestor of the M.EcoGII clade towards the modern enzyme. Other recent studies have similarly shown that it is unnecessary to reconstruct evolutionarily ancient proteins to achieve substantial increases in thermostability by ASR (39, 40). For example, Livada *et al*. systematically characterized all 56 ancestral nodes and 57 extant nodes in a phylogeny of gene reductases and found that even the shallowest nodes (i.e., the most recent ancestral nodes, adjacent to the extant nodes at the tips of the tree) had a T_M up to 16 °C higher than their closest extant protein (40). The reason for this general trend of reconstructed ancestral proteins showing higher thermostability compared with extant proteins is still unclear, but the bias in maximum-likelihood ASR towards common amino acids (38) and consensus amino acids (35) at ambiguously reconstructed positions may be partially responsible. The ability of ASR to preserve coevolutionary relationships between residues also provides an advantage over related methods such as consensus design (38). Overall, our work contributes to a growing body of evidence that ASR can be successfully applied as a protein engineering tool to a diverse range of protein families, even those with limited phylogenetic diversity.

Substantial differences in RNA methyltransferase activity were also observed among the ancestral M.EcoGII variants, which likely reflects the neutral variation of the promiscuous RNA methyltransferase activity of M.EcoGII throughout its evolutionary history. Although the precise biological function of M.EcoGII is unknown, its DNA methyltransferase activity is more likely to be biologically relevant, given that (1) phage DNA methyltransferases have a general role in the protection of phage DNA from host restriction enzymes during infection (50); (2) M.EcoGII shows higher DNA methyltransferase activity than RNA methyltransferase activity (30); and (3) m^6A modification of mRNA has no known function in bacteria, although it is known to occur in Gram-negative bacteria (18). Thus, we hypothesize that the DNA methyltransferase activity of M.EcoGII represents its native activity and that its RNA methyltransferase activity represents a promiscuous activity that is not itself under selection pressure. In other words, we hypothesize that the RNA methyltransferase activity is a byproduct of non-specific DNA methyltransferase activity that is driven mainly by non-specific ionic interactions between the positively charged surface of M.EcoGII and the

negatively charged phosphate groups of DNA, which also enables binding of RNA. Consistent with the fact that the promiscuous activities of enzymes generally show greater variability among homologous enzymes than their native activities (51), the observed variation in RNA methyltransferase activity among the M.EcoGII variants (~12-fold) is greater than the variation in DNA methyltransferase activity (~3-fold). The fact that there is no trend in RNA methyltransferase activity from the earlier ancestral proteins towards M.EcoGII is also suggestive of neutral variation. Altogether, these considerations suggest that the ~12-fold increase in RNA methyltransferase activity of M.EcoGII achieved by ASR can be attributed to sampling of the neutral variation of RNA methyltransferase activity that occurred throughout its evolutionary history. More generally, these results suggest that ASR can be used to obtain enzyme variants with improved properties and increased activity, even if the desired activity is not the native enzyme activity.

The combination of non-site-specific and highly efficient m^{6}A methylation activity observed in SUPREM is unique among known RNMTs and should be broadly useful in applications that require unbiased and efficient labeling of RNA. For example, SUPREM could potentially be applied to proximity labeling for transcriptome-wide detection of RNA binding sites for RNA-binding proteins. Fusion of a specific RNA-binding protein to SUPREM would lead to increased m^{6}A methylation adjacent to the binding site of the RNA-binding protein, allowing detection of the binding site using techniques for m^{6}A detection, including Oxford Nanopore direct RNA sequencing. A similar approach, TRIBE, relies on fusion of an RNA-binding protein to the RNA-editing enzyme adenosine deaminase (ADAR) rather than an RNMT for proximity labeling of the RNA binding site (52). This method provides numerous advantages over the widely-used cross-linking immunoprecipitation (CLIP) method for detection of RNA-protein interactions, such as the relatively low amount of input material needed; however, this method may also be susceptible to bias due to the inherent substrate specificity of ADAR (52), a problem that could potentially be addressed using SUPREM. SUPREM could also be applied to site-specific modification of RNA through conjugation of SUPREM to sequence-programmable RNA-binding proteins, such as dCas13 and PUF proteins (26, 27). Unlike previous iterations that rely on fusion proteins of the human m^{6}A writer METTL3, SUPREM would also allow site-specific methylation of sites targeted by other m^{6}A writers such as METTL16, as well as non-canonical methylation sites and sites that are not naturally methylated (for example, to modulate mRNA stability, localization or translation efficiency). Finally, the improved thermostability, expression, and RNA methyltransferase activity of SUPREM make it a useful starting point for further protein engineering. For

example, engineering of the cofactor binding pocket of SUPREM to promote the binding of

synthetic, functionalized *S*-adenosyl-L-methionine analogs, an approach that has been applied

to sequence-specific DNA methyltransferases (53), would allow bioorthogonal sequence-

independent functionalization and labeling of mRNA. Altogether, we expect that SUPREM

will be a powerful enabling tool for the development of new methods for RNA manipulation

and analysis.

Materials and methods

Phylogenetic analysis

Protein BLAST was used to search the collection of amino acid sequences of type II class β

methyltransferases, to which M.EcoGII belongs, in the REBASE database (41) (retrieved on

2020-09-17) using M.EcoGII (GenBank accession number EGT69837.1) as a query sequence.

An e-value cutoff of 10^{-7} was used. Redundant sequences were omitted by CD-HIT using a

90% sequence identity threshold, giving a total of 2368 sequences (54). The sequence

similarity network was created by all-versus-all BLAST of the resulting sequences using an e-

value threshold of 10^{-30} and visualized in Cytoscape software. For ancestral sequence

reconstruction, multiple sequence alignments of the M.EcoGII-containing cluster (158

sequences) and the outgroup clusters (46 sequences) were constructed separately using

MUSCLE software (55). The aligned protein sequences were manually edited to remove two

outlier sequences, large insertions in the outgroup, and N-/C-terminal extensions. Each edited

multiple sequence alignment was combined by profile-profile alignment in MUSCLE. An

unrooted maximum-likelihood tree was constructed using IQ-TREE software (56). To assess

the robustness of the tree, inference of the phylogenetic tree was repeated five times using the

sample substitution model (LG+I+G4) or two other substitution models (WAG and VT). The

LG+I+G4 model was chosen as the best model based on IQ-tree calculation of Bayesian

information criterion (BIC) score. The robustness of each node was assessed using the

Shimodaira-Hasegawa approximate likelihood ratio test (SH-aLRT) and ultrafast bootstrap

analysis (UFboot) with 1000 replicates. Ancestral protein sequences were reconstructed using

the empirical Bayes method implemented in PAML using the LG substitution model (57). The

posterior probability distribution at each site for each ancestral node was also calculated using

PAML.

Cloning and mutagenesis

M.EcoGII and six ancestral protein sequences were reverse-translated into nucleotide

sequences that were codon optimized for expression in *E. coli*. A sequence encoding a C-

terminal His$_6$ tag with a linker sequence (N-GGAAALGHHHHHH-C) was added to each gene,

and the resulting DNA sequences were synthesized (Twist Bioscience, California, USA) and

cloned into the pET-28a(+) plasmid using In-Fusion cloning kit (Takara Bio Inc., Shiga, Japan)

or the iVEC3 method (58). In-frame cloning into the pET-28a(+) vector was confirmed by

DNA sequencing. Mutagenesis of SUPREM was performed using the PrimeStar Mutagenesis

method and PrimeStarMax polymerase (Takara Bio Inc., Shiga, Japan).

The DNA sequences corresponding to M.EcoGII and SUPREM were inserted into the

pCDNA3.1-eGFP-GSx3 plasmid using the In-Fusion cloning kit (Takara Bio Inc., Shiga,

Japan), resulting in a sequence encoding eGFP with a flexible linker sequence (GGGGS)$_3$

linked to the N-terminus of the methyltransferase.

Protein expression

Plasmids containing M.EcoGII and all ancestral protein genes were transformed into *E.*

coli strain SHuffle T7 (New England Biolabs Inc., Massachusetts, USA) and plated on LB agar

containing 30 μg/mL kanamycin and 2% glucose, incubating at 37 °C overnight. Colonies were

inoculated into 5-20 mL LB media with 30 μg/ml kanamycin and 2% glucose and incubated at

37 °C with shaking at 220 rpm overnight. The preculture was transferred into 0.5-2 L LB media

with 30 μg/mL kanamycin and 1% glucose, and the *E. coli* cells were grown at 37 °C with

shaking at 220 rpm until the OD$_{600}$ reached 0.5-0.8. Protein expression of M.EcoGII and the

ancestral proteins was induced by 1 mM isopropyl β-D-1-thiogalactopyranoside (IPTG) for 4

h at 25 °C. SUPREM variants were expressed with 1 mM IPTG for 18 h at 18 °C. Cells were

harvested and stored at -20 °C before protein purification.

Protein purification

Cells were resuspended with lysis buffer (20 mM HEPES pH 7.0, 250 mM NaCl, 50

mM imidazole with 1× Protease Inhibitor Cocktail (EDTA-free) (Nacalai Tesque Inc., Kyoto,

Japan) and 1 mg/ml lysozyme (Sigma-Aldrich, Missouri, USA) and lysed by sonication. The

lysate was centrifuged at 10000 × *g*, 4 °C for 30 min, and the supernatant was filtered through

a 0.45 μm filter. The initial protein purification was performed using an ÄKTA Pure 150

system (Cytiva Japan, Tokyo, Japan) and a 1 ml HisTrap FF column (Cytiva Japan, Tokyo,

Japan) with Ni wash buffer (20 mM HEPES pH 7.0, 250 mM NaCl, 20 mM imidazole),

followed with elution buffer (20 mM HEPES pH 7.0, 250 mM NaCl, 200 mM imidazole) as a

stepwise purification. Peak fractions were analyzed by SDS-PAGE. Fractions enriched in the
target protein were pooled on ice, filtered through a 0.22 μm filter, and purified by size
exclusion column chromatography (HiLoad 26/600 Superdex 75 pg, Cytiva Japan,
Tokyo, Japan) with working buffer (20 mM HEPES pH 7.0, 250 mM NaCl). Peak fractions
were analyzed by SDS-PAGE. Fractions enriched in the target protein were pooled on ice,
filtered through a 0.22 μm filter, and concentrated in an Amicon Ultra 10 kDa molecular weight
cutoff (MWCO) centrifugal concentration device (Merck Millipore) with working buffer. The
concentrated protein samples (> 1 mg/ml) were mixed with glycerol to reach 10% (v/v)
concentration, quickly frozen by liquid nitrogen, and stored at -80 ºC.

For SUPREM variants, cells were resuspended with lysis buffer (20 mM HEPES pH
7.0, 250 mM NaCl, 20 mM imidazole with 1× Protease Inhibitor Cocktail (EDTA-free)
(Nacalai Tesque Inc., Kyoto, Japan) and 1 mg/ml lysozyme (Sigma-Aldrich, Missouri, USA)
and lysed by sonication. The lysis supernatant was obtained by centrifugation at $10000 \times g$,
4 °C for 30 min. SUPREM variants were purified by affinity chromatography in an open
column using cOmplete His-Tag Purification Resin (Roche, Basel, Switzerland), eluting with
20 mM HEPES pH 7.0, 250 mM NaCl, and 100 mM imidazole at room temperature. Each
fraction was analyzed by SDS-PAGE. Fractions enriched in the target protein were pooled,
filtered through a 0.22 μm filter, and concentrated in an Amicon Ultra 10 kDa MWCO
centrifugal concentration device (Merck Millipore) with working buffer. The concentrated
protein samples (>1 mg/ml protein and < 1 mM imidazole) were mixed with glycerol to reach
10% (v/v) concentration, quickly frozen by liquid nitrogen, and stored at -80 ºC.

MTase-Glo assay

The DNA substrate encoding M.EcoGII (3569 bp, containing 1819 adenine bases) was
amplified by PCR and purified using a NucleoSpin Gel and PCR Clean-up Kit (MACHEREY-
NAGEL GmbH & Co. KG, Düren, Germany) with Milli-Q water. The RNA substrate encoding
the firefly luciferase gene (1872 bases, 544 adenine bases) was prepared by *in vitro*
transcription using HiScribe T7 High Yield RNA Synthesis Kit (New England Biolabs Inc.,
Massachusetts, USA) with 1 unit Rnase Inhibitor (New England Biolabs Inc., Massachusetts,
USA) and purified in RNase-free water using a NucleoSpin RNA Clean-up spin column kit
with a DNA removal column (MACHEREY-NAGEL GmbH & Co. KG, Düren, Germany). *In*
vitro methylation experiments were performed for 10 min (DNA) or 1 h (RNA) at 37 °C in 8
μL reaction mixtures containing 1× CutSmart buffer (New England Biolabs Inc.,
Massachusetts, USA) with 20 nM DNA or 50 nM RNA with 1 unit RNase Inhibitor (New

England Biolabs Inc., Massachusetts, USA), 100 nM methyltransferase, and 20 μM SAM. The

methylation reaction in the 8 μL reaction was quenched by heating at 80 °C for 1 min at each

time point. The methyltransferase reaction product *S*-adenosylhomocysteine was quantified

using MTase-Glo kit (Promega, Wisconsin, USA) following the manufacturer's protocol (43).

To normalize the SAH production rate in DNA/RNA methylation, a calibration curve for SAH

was constructed in two independent experiments using from 0 to 5 μM SAH according to the

manufacturer's protocol of the MTase-Glo kit (Promega, Wisconsin, USA). The signal was

detected using a Tecan Spark microplate reader (Tecan Japan Co., Ltd, Japan).

UPLC assay

The RNA substrate, a transcribed portion of the pET-28a(+) vector (294 bases, 80

adenine bases), was prepared by *in vitro* transcription using HiScribe T7 High Yield RNA

Synthesis Kit (New England Biolabs Inc., Massachusetts, USA) with 1 unit RNase Inhibitor

(New England Biolabs Inc., Massachusetts, USA) and purified in RNase-free water using a

NucleoSpin RNA Clean-up spin column kit with a DNA removal column (MACHEREY-

NAGEL GmbH & Co. KG, Düren, Germany). *In vitro* RNA methylation experiments were

performed for 2 h at 37 °C in 20 μL reaction mixtures containing 1× CutSmart buffer (New

England Biolabs Inc., Massachusetts, USA) with 1 μg RNA and 1 unit RNase Inhibitor (New

England Biolabs Inc., Massachusetts, USA), 5 μg methyltransferase or no enzyme, and 320

μM SAM (New England Biolabs Inc., Massachusetts, USA). The methylation reaction was

quenched by heating at 80 °C for 1 min. RNA was purified using an RNA Clean &

Concentrator-5 kit (Zymo Research, California, USA) into 20 μL RNase-free water, and then

digested from RNA to nucleosides using a Nucleoside Digestion Mix kit (New England

Biolabs Inc., Massachusetts, USA) at 37 ºC for 1 h. The digestion reaction was quenched with

33% ethanol as the final concentration. The solvent was removed using a centrifugal evaporator,

and the sample was dissolved in 100 μL of 40 mM ammonium acetate pH 6.0. LC-MS analysis

was performed in triplicate on a Waters UPLC H-class system (Waters Corporation,

Massachusetts, USA) using an ACQUITY UPLC BEH C18 column (2.1 mm × 50 mm, 1.7

μm) with a buffer system consisting of 40 mM ammonium acetate pH 6.0 (A) and 40%

acetonitrile (B). The nucleosides were eluted at a flow rate of 0.25 ml/min with a linear gradient

of mobile phase of 0% B (0-1.5 min), 0-30% B (1.5-15 min), and 0% B (15-30 min) for re-

equilibration. The nucleoside peaks were compared between enzyme-treated samples and

standard compounds to identify each peak.

Differential scanning fluorimetry assay

The DSF experiments were conducted using the StepOnePlus real-time PCR instrument (Applied Biosystems, Massachusetts, USA), as described previously (59). Reaction mixtures were prepared in 96-well Real-Time PCR Plates, comprising 5× SYPRO Orange (Sigma-Aldrich, Missouri, USA), 5 μM enzyme, and a total volume of 20 μL working buffer (20 mM HEPES pH 7.0, 250 mM NaCl). The plate was sealed with optically clear sealing film and centrifuged at 2000 × g for 1 min prior to loading into the real-time PCR instrument. The temperature was ramped at a rate of 1% (approximately 1.33 °C/min) from 4 ºC to 95 ºC. Fluorescence signals were monitored using the ROX channel. The melting temperature values were determined by calculating the derivative of fluorescence intensity with respect to temperature and fitting the resulting data to a quadratic equation within a 6 °C window around the T_M, utilizing R software.

Nanopore direct RNA sequencing

For direct RNA sequencing by MinION sequencer (Oxford Nanopore Technologies, Oxford, UK), *in vitro* RNA methylation experiments were performed for 2 h at 37 °C in 20 μL reaction mixtures containing 1× CutSmart buffer with 5 μg RNA substrate (1.8 knt. Same substrate of MTase-Glo experiment) and 1 unit RNase Inhibitor (New England Biolabs Inc., Massachusetts, USA), 1 μM methyltransferase or no enzyme as a negative control, and 320 μM SAM for two independent reaction mixture. The reaction mixture without enzyme was used as a negative control. The polyadenylated tail was added by incubation with 5 U of poly-A polymerase (New England Biolabs Inc., Massachusetts, USA) for 30 min at 37 °C. The methylated and poly-A-tailed RNA samples were purified using an RNA purification kit (Zymo Research, California, USA) with RNase-free water. The concentration and purity of the RNA samples were measured using a Qubit 4 fluorometer with the Qubit RNA IQ kit (Thermo Fisher Scientific, Massachusetts, USA) (RIN score > 8.5). The preparation of libraries for direct RNA sequencing was conducted using the SQK-RNA002 kit from 500 ng of *in vitro* transcribed RNA. The direct RNA sequencing was carried out on a flow cell R.9.4.1 (FLO-MIN106, Oxford Nanopore Technologies, Oxford, UK) by the Sequencing Section of the Okinawa Institute of Science and Technology, following the protocol provided by Oxford Nanopore Technologies.

Sequencing data analysis

The data analysis was performed using the Nanocompore pipeline (44). Base-called
fastq files were created using Guppy software (version 6.4.2) and were concatenated into one
file. The transcript data were aligned to a reference transcript sequence by minimap2 (version
2.20), following sorting and indexing in samtools (version 1.15.1). The read file was indexed
and resquiggled using the index and eventalign commands in Nanocompore (44). The m^6A
peaks were detected using the eventalign_collapse and sampcomp commands in Nanocompore
(44). The detected peaks were visualized using ggplot2 (60), and methylated 5-mer motifs
containing adenine bases were extracted from the total detected motifs and visualized using an
R script and Prism 9 software (GraphPad Software).

Size exclusion chromatography analysis
Fifty to one hundred μg SUPREM or M.EcoGII samples were injected into a Superdex
200 Increase 10/300 GL column (Cytiva Japan, Tokyo, Japan), and eluted at 0.75 ml/min flow
rate with 1 column volume of working buffer (20 mM HEPES pH 7.0, 250 mM NaCl). A
calibration curve was constructed using the elution volumes of the LMW Marker Kit (Cytiva
Japan, Tokyo, Japan).

Structural modeling and analysis
Dimer structures of methyltransferases were modeled by Alphafold (ver.2.2.1) using
the multimer prediction option (49). Protein structural models were visualized using PyMOL
software. Structures similar to M.EcoGII and its target recognition domain were searched using
the DALI server (61).

Immunofluorescence assay
HEK 293T cells were grown in DMEM high glucose (Nacalai Tesque, 08458-45)
supplemented with 10% fetal bovine serum (FBS, Gibco, 102-70-106) and 100 μg/mL
primocin (InvivoGen, ant-pm-1). Cells were seeded at an appropriate confluency on polylysine
(Sigma, P4832) coated coverslips one day before JetPei® (Polyplus, 101000053) transfection
with the different constructs, following the manufacturer's recommendations. 48 hours after
transfection, cells were fixed for 30 minutes at room temperature (RT) in 4% paraformaldehyde
in PBS under a fume hood, then permeabilized in 0.3% Triton in PBS for 30 minutes at RT.
Cells were blocked with 1% (v/v) BSA in PBS for 1 hour at RT and incubated overnight at
4 °C with primary anti-m^6A antibody (Synaptic systems, 202-003) diluted in the blocking
solution. The next day, cells were incubated with secondary conjugated Alexa Fluor 647

antibody (Invitrogen, A21245) for 1 hour in the blocking solution, and then with NucBlue™ (Invitrogen, R37606) for 5 minutes in PBS. Slides were mounted in Fluoromount G™ (Invitrogen, 00-4958-02), and fluorescence was observed with Zeiss LSM 900 confocal microscope. m^{6}A immunofluorescence staining was quantified using the software ImageJ. A mask was made using the GFP fluorescence driven by the transfected construct and m^{6}A intensity was quantified inside the mask and averaged by the cell area. Statistical analysis was performed using the software GraphPad Prism. Statistical significance was evaluated using one-way ANOVA followed by *Tukey* posttest.

DATA AVAILABILITY

Raw fastq files of the Nanopore direct RNA sequencing were deposited in the National Center of Biotechnology Information (BioProject PRJNA990656). The pET28a-SUPREM (ID 205023), pET28a-AncMEcoGII291 (Anc291; ID 205024), and pET28a-AncMEcoGII250 (Anc250; ID 205025) plasmids will be made available via Addgene upon publication.

Author Contribution

Conceptualization: PL; Supervision: BEC, PL; Investigation: YO; Analysis: BEC, YO, PL; Cell Experiments and analysis: MLC and MT; Writing – original draft: YO, BEC, PL; Writing – review and editing: BEC, YO, PL, MLC, MT.

Acknowledgements

We thank the Okinawa Institute of Science and Technology Sequencing Section for preparing the library and sequencing Oxford Nanopore MinION Sequencer. We are grateful for the help and support provided by the Scientific Computing and Data Analysis section of the Research Support Division at OIST.

Funding

Funding for the research was supported by Okinawa Institute of Science and Technology (OIST). PL acknowledges Kakenhi Grant (No. 90812256). BEC was supported by a JSPS Postdoctoral Fellowship for Overseas Researchers from the Japan Society for the Promotion of Science. This project was supported by an OIST Kick start-up grant.

600 **CONFLICT OF INTEREST**

601 None declared.

Figure 1. Ancestral sequence reconstruction of M.EcoGII. (A) Schematic maximum likelihood tree. The node values showed ultrafast bootstrap values calculated by IQ-TREE. **(B-C)** MTase-Glo assay of M.EcoGII and its ancestral proteins for (B) DNA and (C) RNA at a single time point. **(B)** The DNA methylation reaction was performed using 100 nM of methyltransferases, 50 nM DNA substrate (3.5 kbp containing 1844 adenine bases, Supplementary Table S2) and 10 μM SAM in 1× CutSmart buffer (NEB) at 37 °C for 10 min. **(C)** The RNA methylation reaction was performed using 100 nM of the corresponding methyltransferases, 25 nM RNA substrate (1.7 kb containing 544 adenine bases, sequence in Supplementary Table S2), and 10 μM SAM in 1× CutSmart buffer (NEB) at 37 °C for 60 min. Three independent experiments were performed, and each point showed an average of technical duplicates. *P* values were determined by the one-way ANOVA test with the Dunnett test. **(D)** Time course experiments for RNA substrates. The condition of RNA methylation reaction was the same as (C) except for reaction time. Three independent experiments were performed, and each point showed an average of technical duplicates. SAH production amount was calculated by SAH standard curve. The lines represent simple linear regression analysis

on the data from 0 to 120 min (SUPREM) or to 240 min (M.EcoGII, Anc291). **(E)** The fraction

of m^6A methylation in total adenine by UPLC analysis. The RNA methylation reaction was

performed using 5 μg of the corresponding methyltransferases, 1 μg RNA substrate (294 nt

containing 80 adenine bases, sequence in Supplementary Table S2) and 320 μM SAM (NEB)

in 1× CutSmart buffer (NEB) at 37 °C for 120 min. Each point showed three independent

methylation reactions. Error bars represent standard error of the mean (SEM).

Figure 2. Methylation activity in HEK293 cell. (A) Representative images (63x objective with a 2x digital zoom) of HEK293T cells transfected with either eGFP, M.EcoGII-eGFP or SUPREM-eGFP constructs (reporter visible in green). m^{6}A modification were detected via immunofluorescence (in red) and the nucleus was stained with NucBlue™ (in blue). The scale bar is 10 μm. (B) Quantification of m^{6}A fluorescence intensity in HEK293 transfected as described in (A). The bar chart indicates the normalized intensity by cell area. Statistical analysis was performed by one-way ANOVA followed by *Tukey* posttest *(*: <0.0332; **: <0.0021).*

637   **Figure 3. RNA methylation motif analysis. (A)** Mapping detected RNA methylation sites on

638   the 18 knt Fluc RNA sequence by MinION DRS and Nanocompore pipeline. **(B)** Mean RNA

639   base frequency in modified 5-mer motifs detected by DRS. The "Expectation" represents the

640   mean frequency of each base in 5-mers that contain at least one adenine within the RNA

641   substrate.

Figure 4. Mutational and structural analysis of SUPREM. (A) Candidate amino acid residues for mutational analysis. (Left) M.EcoGII clade (clade I, 46 sequences) and SUPREM clade (clade II, 19 sequences). (Right) Comparison of amino acid sequence logo of mutational candidate sites based on multiple sequence alignment of clade I and clade II. The height of the logo represents the frequency of amino acid residues at each position. **(B)** RNA methylation activity of SUPREM variants. The RNA methylation reaction was performed using 100 nM methyltransferase and 25 nM RNA substrate (1.7 kb. 544 adenine bases) with 10 μM SAM in 1× CutSmart buffer (NEB) at 37 ºC for 60 min. The luminescence was detected by MTase-Glo assay. Three or four independent experiments were performed, and each point shows an average of technical duplicates. *P* values were determined by the one-way ANOVA test with the Dunnett test. **(C)** Size exclusion chromatogram of SUPREM. SUPREM with blue dextran (for estimation of void volume) was loaded into a Superdex 200 Increase 10/300 GL column, and the samples were eluted with 20 mM HEPES pH 7.4 and 250 mM NaCl. The peak positions of low molecular weight markers were shown above. The two independent measurements were performed. **(D)** Dimer structural model of SUPREM generated by AlphaFold2 multimer

659   program (version 2.2.1). Green spheres show residues selected for mutational analysis (Fig. 3).

660   **(E)** APBS calculation of the electrostatic surface potential of SUPREM (red: negative charge;

661   blue: positive charge).

References

1. Dominissini,D., Moshitch-Moshkovitz,S., Schwartz,S., Salmon-Divon,M., Ungar,L., Osenberg,S., Cesarkas,K., Jacob-Hirsch,J., Amariglio,N., Kupiec,M., *et al.* (2012) Topology of the human and mouse m6A RNA methylomes revealed by m6A-seq. *Nature*, **485**, 201–206.

2. Liu,J., Yue,Y., Han,D., Wang,X., Fu,Y., Zhang,L., Jia,G., Yu,M., Lu,Z., Deng,X., *et al.* (2014) A METTL3-METTL14 complex mediates mammalian nuclear RNA N6-adenosine methylation. *Nat Chem Biol*, **10**, 93–95.

3. Warda,A.S., Kretschmer,J., Hackert,P., Lenz,C., Urlaub,H., Höbartner,C., Sloan,K.E. and Bohnsack,M.T. (2017) Human METTL16 is a N6-methyladenosine (m6A) methyltransferase that targets pre-mRNAs and various non-coding RNAs. *EMBO reports*, **18**, 2004–2014.

4. Wang,X., Lu,Z., Gomez,A., Hon,G.C., Yue,Y., Han,D., Fu,Y., Parisien,M., Dai,Q., Jia,G., *et al.* (2014) N6-methyladenosine-dependent regulation of messenger RNA stability. *Nature*, **505**, 117–120.

5. Geula,S., Moshitch-Moshkovitz,S., Dominissini,D., Mansour,A.A., Kol,N., Salmon-Divon,M., Hershkovitz,V., Peer,E., Mor,N., Manor,Y.S., *et al.* (2015) m6A mRNA methylation facilitates resolution of naïve pluripotency toward differentiation. *Science*, **347**, 1002–1006.

6. Edupuganti,R.R., Geiger,S., Lindeboom,R.G.H., Shi,H., Hsu,P.J., Lu,Z., Wang,S.-Y., Baltissen,M.P.A., Jansen,P.W.T.C., Rossa,M., *et al.* (2017) N6-methyladenosine (m6A) recruits and repels proteins to regulate mRNA homeostasis. *Nat Struct Mol Biol*, **24**, 870–878.

7. Pendleton,K.E., Chen,B., Liu,K., Hunter,O.V., Xie,Y., Tu,B.P. and Conrad,N.K. (2017) The U6 snRNA m6A Methyltransferase METTL16 Regulates SAM Synthetase Intron Retention. *Cell*, **169**, 824-835.e14.

8. Mendel,M., Delaney,K., Pandey,R.R., Chen,K.-M., Wenda,J.M., Vågbø,C.B., Steiner,F.A., Homolka,D. and Pillai,R.S. (2021) Splice site m6A methylation prevents binding of U2AF35 to inhibit RNA splicing. *Cell*, **184**, 3125-3142.e25.

9. Price,A.M., Hayer,K.E., McIntyre,A.B.R., Gokhale,N.S., Abebe,J.S., Della Fera,A.N., Mason,C.E., Horner,S.M., Wilson,A.C., Depledge,D.P., *et al.* (2020) Direct RNA sequencing reveals m6A modifications on adenovirus RNA are necessary for efficient splicing. *Nat Commun*, **11**, 6016.

10. Yu,J., Chen,M., Huang,H., Zhu,J., Song,H., Zhu,J., Park,J. and Ji,S.-J. (2018) Dynamic m6A modification regulates local translation of mRNA in axons. *Nucleic Acids Research*, **46**, 1412–1423.

11. Di Timoteo,G., Dattilo,D., Centrón-Broco,A., Colantoni,A., Guarnacci,M., Rossi,F., Incarnato,D., Oliviero,S., Fatica,A., Morlando,M., *et al.* (2020) Modulation of circRNA Metabolism by m6A Modification. *Cell Reports*, **31**, 107641.

12. Lin,S., Choe,J., Du,P., Triboulet,R. and Gregory,R.I. (2016) The m6A Methyltransferase
METTL3 Promotes Translation in Human Cancer Cells. *Molecular Cell*, **62**, 335–345.

13. Meyer,K.D., Patil,D.P., Zhou,J., Zinoviev,A., Skabkin,M.A., Elemento,O., Pestova,T.V.,
Qian,S.-B. and Jaffrey,S.R. (2015) 5' UTR m6A Promotes Cap-Independent
Translation. *Cell*, **163**, 999–1010.

14. Wang,X., Zhao,B.S., Roundtree,I.A., Lu,Z., Han,D., Ma,H., Weng,X., Chen,K., Shi,H. and
He,C. (2015) N6-methyladenosine Modulates Messenger RNA Translation Efficiency.
Cell, **161**, 1388–1399.

15. Zheng,G., Dahl,J.A., Niu,Y., Fedorcsak,P., Huang,C.-M., Li,C.J., Vågbø,C.B., Shi,Y.,
Wang,W.-L., Song,S.-H., *et al.* (2013) ALKBH5 Is a Mammalian RNA Demethylase that
Impacts RNA Metabolism and Mouse Fertility. *Molecular Cell*, **49**, 18–29.

16. Yoshinaga,M., Han,K., Morgens,D.W., Horii,T., Kobayashi,R., Tsuruyama,T., Hia,F.,
Yasukura,S., Kajiya,A., Cai,T., *et al.* (2022) The N6-methyladenosine
methyltransferase METTL16 enables erythropoiesis through safeguarding genome
integrity. *Nat Commun*, **13**, 6435.

17. Zhou,J., Wan,J., Gao,X., Zhang,X., Jaffrey,S.R. and Qian,S.-B. (2015) Dynamic m6A mRNA
methylation directs translational control of heat shock response. *Nature*, **526**, 591–
594.

18. Deng,X., Chen,K., Luo,G.-Z., Weng,X., Ji,Q., Zhou,T. and He,C. (2015) Widespread
occurrence of N6-methyladenosine in bacterial mRNA. *Nucleic Acids Research*, **43**,
6557–6567.

19. Liu,J., Eckert,M.A., Harada,B.T., Liu,S.-M., Lu,Z., Yu,K., Tienda,S.M., Chryplewicz,A.,
Zhu,A.C., Yang,Y., *et al.* (2018) m6A mRNA methylation regulates AKT activity to
promote the proliferation and tumorigenicity of endometrial cancer. *Nat Cell Biol*,
20, 1074–1083.

20. Zhang,C., Tunes,L., Hsieh,M.-H., Wang,P., Kumar,A., Khadgi,B.B., Yang,Y.-Y.,
Doxtader,K.A., Herrell,E., Koczy,O., *et al.* (2023) Cancer mutations rewire the RNA
methylation specificity of METTL3-METTL14. 10.1101/2023.03.16.532618.

21. Winkler,R., Gillis,E., Lasman,L., Safra,M., Geula,S., Soyris,C., Nachshon,A., Tai-
Schmiedel,J., Friedman,N., Le-Trilling,V.T.K., *et al.* (2019) m6A modification controls
the innate immune response to infection by targeting type I interferons. *Nat
Immunol*, **20**, 173–182.

22. De Jesus,D.F., Zhang,Z., Kahraman,S., Brown,N.K., Chen,M., Hu,J., Gupta,M.K., He,C. and
Kulkarni,R.N. (2019) m6A mRNA methylation regulates human β-cell biology in
physiological states and in type 2 diabetes. *Nat Metab*, **1**, 765–774.

23. Boulias,K. and Greer,E.L. (2023) Biological roles of adenine methylation in RNA. *Nat Rev
Genet*, **24**, 143–160.

24. Sendinc,E. and Shi,Y. (2023) RNA m6A methylation across the transcriptome. *Molecular*
Cell, **83**, 428–441.

25. Liu,X.-M., Zhou,J., Mao,Y., Ji,Q. and Qian,S.-B. (2019) Programmable RNA N6-
methyladenosine editing by CRISPR-Cas9 conjugates. *Nat Chem Biol*, **15**, 865–871.

26. Wilson,C., Chen,P.J., Miao,Z. and Liu,D.R. (2020) Programmable m6A modification of
cellular RNAs with a Cas13-directed methyltransferase. *Nat Biotechnol*, **38**, 1431–
1440.

27. Shinoda,K., Suda,A., Otonari,K., Futaki,S. and Imanishi,M. (2020) Programmable RNA
methylation and demethylation using PUF RNA binding proteins. *Chem. Commun.*,
56, 1365–1368.

28. Su,S., Li,S., Deng,T., Gao,M., Yin,Y., Wu,B., Peng,C., Liu,J., Ma,J. and Zhang,K. (2022) Cryo-
EM structures of human m6A writer complexes. *Cell Res*, **32**, 982–994.

29. Fang,G., Munera,D., Friedman,D.I., Mandlik,A., Chao,M.C., Banerjee,O., Feng,Z., Losic,B.,
Mahajan,M.C., Jabado,O.J., *et al.* (2012) Genome-wide mapping of methylated
adenine residues in pathogenic Escherichia coli using single-molecule real-time
sequencing. *Nat Biotechnol*, **30**, 1232–1239.

30. Murray,I.A., Morgan,R.D., Luyten,Y., Fomenkov,A., Corrêa,I.R., Dai,N., Allaw,M.B.,
Zhang,X., Cheng,X. and Roberts,R.J. (2018) The non-specific adenine DNA
methyltransferase M.EcoGII. *Nucleic Acids Research*, **46**, 840–848.

31. Sobecki,M., Souaid,C., Boulay,J., Guerineau,V., Noordermeer,D. and Crabbe,L. (2018)
MadID, a Versatile Approach to Map Protein-DNA Interactions, Highlights Telomere-
Nuclear Envelope Contact Sites in Human Cells. *Cell Reports*, **25**, 2891-2903.e5.

32. Shipony,Z., Marinov,G.K., Swaffer,M.P., Sinnott-Armstrong,N.A., Skotheim,J.M.,
Kundaje,A. and Greenleaf,W.J. (2020) Long-range single-molecule mapping of
chromatin accessibility in eukaryotes. *Nat Methods*, **17**, 319–327.

33. Zakas,P.M., Brown,H.C., Knight,K., Meeks,S.L., Spencer,H.T., Gaucher,E.A. and
Doering,C.B. (2017) Enhancing the pharmaceutical properties of protein drugs by
ancestral sequence reconstruction. *Nat Biotechnol*, **35**, 35–37.

34. Nakano,S., Niwa,M., Asano,Y. and Ito,S. (2019) Following the Evolutionary Track of a
Highly Specific l-Arginine Oxidase by Reconstruction and Biochemical Analysis of
Ancestral and Native Enzymes. *Applied and Environmental Microbiology*, **85**, e00459-
19.

35. Trudeau,D.L., Kaltenbach,M. and Tawfik,D.S. (2016) On the Potential Origins of the High
Stability of Reconstructed Ancestral Proteins. *Molecular Biology and Evolution*, **33**,
2633–2641.

36. Babkova,P., Sebestova,E., Brezovsky,J., Chaloupkova,R. and Damborsky,J. (2017)
Ancestral Haloalkane Dehalogenases Show Robustness and Unique Substrate
Specificity. *ChemBioChem*, **18**, 1448–1456.

37. Clifton,B.E., Kozome,D. and Laurino,P. (2022) Efficient Exploration of Sequence Space by
Sequence-Guided Protein Engineering and Design. *Biochemistry*,
10.1021/acs.biochem.1c00757.

38. Spence,M.A. (2021) Ancestral sequence reconstruction for protein engineers. *Current
Opinion in Structural Biology*.

39. Joho,Y., Vongsouthi,V., Spence,M.A., Ton,J., Gomez,C., Tan,L.L., Kaczmarski,J.A.,
Caputo,A.T., Royan,S., Jackson,C.J., *et al.* (2023) Ancestral Sequence Reconstruction
Identifies Structural Changes Underlying the Evolution of Ideonella sakaiensis PETase
and Variants with Improved Stability and Activity. *Biochemistry*, **62**, 437–450.

40. Livada,J., Vargas,A.M., Martinez,C.A. and Lewis,R.D. (2023) Ancestral Sequence
Reconstruction Enhances Gene Mining Efforts for Industrial Ene Reductases by
Expanding Enzyme Panels with Thermostable Catalysts. *ACS Catal.*, **13**, 2576–2585.

41. Roberts,R.J., Vincze,T., Posfai,J. and Macelis,D. (2022) REBASE: a database for DNA
restriction and modification: enzymes, genes and genomes. *Nucleic Acids Research*,
10.1093/nar/gkac975.

42. Arndt,D., Marcu,A., Liang,Y. and Wishart,D.S. (2019) PHAST, PHASTER and PHASTEST:
Tools for finding prophage in bacterial genomes. *Briefings in Bioinformatics*, **20**,
1560–1567.

43. Hsiao,K., Zegzouti,H. and Goueli,S.A. (2016) Methyltransferase-Glo: a universal,
bioluminescent and homogenous assay for monitoring all classes of
methyltransferases. *Epigenomics*, **8**, 321–339.

44. Leger,A., Amaral,P.P., Pandolfini,L., Capitanchik,C., Capraro,F., Miano,V., Migliori,V.,
Toolan-Kerr,P., Sideri,T., Enright,A.J., *et al.* (2021) RNA modifications detection by
comparative Nanopore direct RNA sequencing. *Nat Commun*, **12**, 7198.

45. Gupta,Y.K., Chan,S.-H., Xu,S. and Aggarwal,A.K. (2015) Structural basis of asymmetric
DNA methylation and ATP-triggered long-range diffusion by EcoP15I. *Nat Commun*,
6, 7363.

46. Horton,J.R., Woodcock,C.B., Opot,S.B., Reich,N.O., Zhang,X. and Cheng,X. (2019) The cell
cycle-regulated DNA adenine methyltransferase CcrM opens a bubble at its DNA
recognition site. *Nat Commun*, **10**, 4600.

47. Osipiuk,J., Walsh,M.A. and Joachimiak,A. (2003) Crystal structure of MboIIA
methyltransferase. *Nucleic Acids Research*, **31**, 5440–5448.

48. Woodcock,C.B., Horton,J.R., Zhang,X., Blumenthal,R.M. and Cheng,X. (2020) Beta class
amino methyltransferases from bacteria to humans: evolution and structural
consequences. *Nucleic Acids Research*, **48**, 10034–10044.

49. Evans,R., O'Neill,M., Pritzel,A., Antropova,N., Senior,A., Green,T., Žídek,A., Bates,R.,
Blackwell,S., Yim,J., *et al.* (2022) Protein complex prediction with AlphaFold-
Multimer. 10.1101/2021.10.04.463034.

50. Thomson,R.E.S., Carrera-Pacheco,S.E. and Gillam,E.M.J. (2022) Engineering functional
thermostable proteins using ancestral sequence reconstruction. *J Biol Chem*,
10.1016/j.jbc.2022.102435.

51. Khersonsky,O. and Tawfik,D.S. (2010) Enzyme Promiscuity: A Mechanistic and
Evolutionary Perspective. *Annual Review of Biochemistry*, **79**, 471–505.

52. McMahon,A.C., Rahman,R., Jin,H., Shen,J.L., Fieldsend,A., Luo,W. and Rosbash,M. (2016)
TRIBE: Hijacking an RNA-Editing Enzyme to Identify Cell-Specific Targets of RNA-
Binding Proteins. *Cell*, **165**, 742–753.

53. Stankevičius,V., Gibas,P., Masiulionytė,B., Gasiulė,L., Masevičius,V., Klimašauskas,S. and
Vilkaitis,G. (2022) Selective chemical tracking of Dnmt1 catalytic activity in live cells.
Molecular Cell, **82**, 1053-1065.e8.

54. Li,W. and Godzik,A. (2006) Cd-hit: a fast program for clustering and comparing large sets
of protein or nucleotide sequences. *Bioinformatics*, **22**, 1658–1659.

55. Madeira,F., Park,Y.M., Lee,J., Buso,N., Gur,T., Madhusoodanan,N., Basutkar,P.,
Tivey,A.R.N., Potter,S.C., Finn,R.D., *et al.* (2019) The EMBL-EBI search and sequence
analysis tools APIs in 2019. *Nucleic Acids Res*, **47**, W636–W641.

56. Nguyen,L.-T., Schmidt,H.A., von Haeseler,A. and Minh,B.Q. (2015) IQ-TREE: A Fast and
Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies.
Molecular Biology and Evolution, **32**, 268–274.

57. Yang,Z. (2007) PAML 4: Phylogenetic Analysis by Maximum Likelihood. *Molecular Biology
and Evolution*, **24**, 1586–1591.

58. Nozaki,S. and Niki,H. (2019) Exonuclease III (XthA) Enforces In Vivo DNA Cloning of
Escherichia coli To Create Cohesive Ends. *J Bacteriol*, **201**, e00660-18.

59. Clifton,B.E., Alcolombri,U., Jackson,C.J. and Laurino,P. (2023) Ultrahigh-affinity transport
proteins from ubiquitous marine bacteria reveal mechanisms and global patterns of
nutrient uptake. 10.1101/2023.02.16.528805.

60. Wickham,H. (2016) ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New
York.

61. Holm,L. (2022) Dali server: structural unification of protein families. *Nucleic Acids
Research*, **50**, W210–W215.