Effects of biological and environmental factors on orthoebolavirus serology

in bats in Guinea

Maëliss Champagne, [1,2] Julien Cappelle, [1] Alexandre Caron, [1,3] Thibault Pouliquen, [1,4]

Aboubacar Samoura, [5] Mohamed Idriss Doumbouya, [6] Guillaume Thaurignac, Abdoul Karim

Soumah[4], [2] Ahidjo Ayouba, [2] Alpha Kabinet Keita, [2,4] Martine Peeters, [2] Mathieu Bourgarel,

[1,7] Hélène M. De Nys [1,7]

1. ASTRE, CIRAD, INRAE, Université de Montpellier, Montpellier, France

2. TransVIHMI, Institut de Recherche pour le Développement (IRD), Université de

Montpellier, INSERM, Montpellier, France

3. Faculdade de Veterinária, Universidade Eduardo Mondlane, Maputo, Mozambique

4. Centre de Recherche et de Formation en Infectiologie de Guinée (CERFIG), Université

Gamal Abdel Nasser de Conakry, Conakry, Guinea

5. Office Guinéen des Parcs Nationaux et Réserves de Faune (OGPRF), Ministère de

l'Environnement, et du Développement Durable (MEDD), Conakry, Guinea

6. Direction Nationale des Services Vétérinaires, Ministère de l'Elevage et des Productions

Animales, Conakry, Guinea

7. CIRAD, UMR ASTRE, Harare, Zimbabwe

Correspondence should be addressed to Hélène M. De Nys; helene.de_nys@cirad.f

Abstract

Background

   Outbreaks of Ebola disease emerge regularly in human populations on the African continent. This highly virulent haemorrhagic fever can be caused by different orthoebolaviruses from the *Filoviridae* family. Several bat species are believed to carry and transmit some of these viruses to people or other animal species. However, clear evidence is still lacking. Natural circulation and zoonotic transmissions of viruses are generally complex as they often involve several host species and are influenced by biological, environmental and anthropogenic factors. A better understanding of the role of bats and other wildlife in the emergence of Ebola disease outbreaks is crucial for their prevention and early detection.

Methods

   In this longitudinal study we searched for antibodies against orthoebolaviruses in fruit bats in order to explore some of the factors which might influence their natural circulation. We performed serological tests on populations of 4 bat species (*Eidolon helvum*, *Hypsignathus monstrosus*, *Myonycteris angolensis* and *Rousettus aegyptiacus*) sampled longitudinally for 18 months (2018-2020) in Guinea. The analysis of 1,427 bat samples for antibodies directed against different orthoebolavirus species allowed us to test the influence of biological and environmental variables on seropositivity by using GLMM.

Results

   Results showed that the presence of antibodies against one of the orthoebolaviruses antigen is more frequent in the *Eidolon helvum* and *Rousettus aegyptiacus* species, in males, in sexually immature adults, and during the dry season.

Conclusion

   Our results suggest that host species, sex, age, reproductive life-cycle and season may play an important role in the circulation of filoviruses. This information can be used to guide

further sampling to specifically characterize the orthoebolavirus species underlying the production of these antibodies and to adjust surveillance protocols at the interface between bats, humans and other animals.

Keywords: Orthoebolavirus, longitudinal studies, serologic tests, Chiroptera, Guinea

Background

Since the first two outbreaks of Ebola disease (EBOD) in 1976 in Yambuku (Democratic Republic of Congo) and Nzara (South Sudan), more than 30 outbreaks have been officially identified in humans across Central and West Africa (1). The frequency of these outbreaks seems to have recently increased (1) .

Despite recent epidemics which originated from viral resurgence in human populations (2–4), most outbreaks are believed to have resulted from independent zoonotic transmissions, through direct or indirect contact with a putative animal reservoir or other infected animals (5). The reservoir species of viruses from the *Orthoebolavirus* genus remain unknown, but bats are suspected to play an important role in the ecology of these viruses and transmission mechanisms (6–8). It is hypothesized that bats might infect humans directly when hunted and handled (9), or indirectly through contact with bat excrement or fruit contaminated with saliva or excrement (10). Bats might also be an indirect source by transmitting viruses to bridge hosts which can act as amplifiers and be at the origin of human outbreaks (6,11). This has been suspected in Central Africa and Ivory Coast where apes have been confirmed as the source of human infection in several outbreaks (12,13).

Certain bat species are regarded as potential reservoir of orthoebolaviruses since the discovery of viral RNA and anti-Ebola virus (anti-EBOV) antibodies in 3 fruit bat species

(*Epomops franqueti, Hypsignathus monstrosus, Myonycteris torquata*) (6). However, viral

RNA was only detected in 13 of 279 tested specimens, and confirmed by sequence analysis in

7. Other studies have since shown that other frugivorous species have anti-orthoebolavirus

antibodies (*Rousettus aegyptiacus, Eidolon helvum, Epomophorus gambianus, Micropteropus*

pusillus and Myonycteris angolensis) and that insectivorous species (*Mops condylurus and*

Chaerephon pumilus) host a recently discovered *Orthoebolavirus* named Bombali virus

(7,8,14). This supports the fact that these wild bats are exposed to filoviruses and are able to

survive infection, as demonstrated by experimental inoculations for some of these species

(15,16). In addition, *Rousettus aegyptiacus* is the reservoir species of Marburg virus, another

filovirus (17), and other bat species have been shown to be exposed or infected with other

filoviruses such as Lloviu virus detected in Europe (18) and newly described viruses in China

(19). Infectious orthoebolaviruses have never been isolated from free ranging bats so far. The

viremic period might be short or viruses might end up circulating at a low level and might only

be detected for a limited period of time in tissues or oral and rectal swabs, as shown

experimentally in *R. aegyptiacus* inoculated with Ebola and Marburg virus (15,20).

The aerial lifestyle, gregariousness, longevity as well as immunological and metabolic

specificities of bats are traits that may impact the ecology of orthoebolaviruses. Biological and

phenological factors, influenced by seasonal or climate changes, are also important, like the

migratory nature of certain bat species or reproduction patterns, both influenced by the

availability of food resources (21,22). It has been shown that spillover events of Marburg virus

to human populations coincide with the birthing season and presence of juveniles in *Rousettus*

aegyptiacus populations (23). A recent study on an *Eidolon helvum* colony also showed an

increase in the presence of antibodies against orthoebolaviruses when juveniles turn into

immature adults (24). Other factors can influence the dynamics of orthoebolaviruses in their

natural hosts, for instance population size and geographic distribution seem to impact viral

richness within bat populations (25). All these factors influence to some extent viral circulation,

risk of spillover events and emergence and are important to understand to adjust orthoebolavirus

research and EBOD surveillance strategies.

The objective of this study was to investigate the effects of biological and

environmental factors on the presence of anti-orthoebolavirus antibodies and improve

understanding of dynamics of orthoebolavirus circulation in bats. We performed a longitudinal

serological survey of orthoebolavirus in four frugivorous bat species in Guinea, a country at

risk of EBOD outbreak. Based on the outcomes of the serological investigation, a subset of bats

was also screened for filovirus RNA.

Methods

Sample and data collection

We collected blood samples as well as oral and rectal swabs from four frugivorous

bat species, i.e. *Eidolon helvum (E. helvum), Hypsignathus monstrosus (H. monstrosus),*

Myonycteris angolensis (M. angolensis) and Rousettus aegyptiacus (R. aegyptiacus), in which

anti-orthoebolavirus antibodies have already been detected in the past (8). Capture and

sampling were carried out longitudinally over 13 sampling periods (i.e. 16 to 28 consecutive

days of sampling per period) between November 2018 and July 2020, following the protocols

already described in De Nys *et al.* (8). Bats were released directly after sampling. They were

captured in feeding and roosting areas located mainly in four study sites of the prefecture of

Macenta (Figure 1), in forested Guinea, the region where the 2013-2016 Ebola epidemic

emerged. Sites were visited every 2 to 4 weeks during the course of the study. The environment

was mainly constituted of highly transformed habitat, with secondary and degraded forests

being gradually converted into a mosaic of forests and crops over the last 20 years (L. Lee-

Cruz, pers. Comm.). It varied from secondary forest, fields, villages to more urbanised areas.

Climate is characterized by a wet (May to October) and a dry (November to April) season.

Capture site (geographic coordinates, ecological environment), morphology, sex, age

class, status of lactation and late gestation, and species (determined according to morphological

criteria (26,27) and combined to genetic confirmation for some specimens, Additional file 1

Methods S1) were also recorded. Age was categorised as follows: juvenile (articular epiphyses

not yet fully ossified), sexually immature adult (articular epiphyses ossified, but

underdeveloped sexual organs), sexually mature adult (articular epiphyses ossified and clearly

developed sexual organs).

This study was approved by the Comité National d'Ethique pour la Recherche en

Santé of Guinea (permits 011/CNERS/19; 025/CNERS/20; 019/CNERS/20) and the Office

Guinéen des Parcs et Réserves (permit 017/OGUIPAR/MEEF/2017).

137
138

139
140 **Figure 1.** Sampling sites, A: Map of Guinea with the prefecture of Macenta in grey; B: prefecture of Macenta
141 with sampling sites (dots) and number of sampled bats. Source of the basemap:
142 https://data.humdata.org/dataset/cod-ab-gin; license: https://data.humdata.org/faqs/licenses.

143
144
145

 Screening for Ebolaviruses Antibodies

We tested dried blood spots (DBS) for anti-orthoebolavirus IgG with a Luminex-

based serologic assay adapted for bats as described in De Nys *et al*. (8). The assay included ten

recombinant orthoebolavirus proteins, i.e. the glycoprotein (GP), nucleoprotein (NP), and/or

viral protein 40 (VP40), for different orthoebolaviruses: Ebola virus Kissidougou and Mayinga

strains (NP, GP K, GP M, VP40), Sudan virus (NP, GP, VP40), Bundibugyo virus (GP, VP40)

and Reston virus (GP). Recombinant proteins were produced in sf9 insect cell vectors and

purchased at Sinobiologicals (Beijing, China) except for REST GP (IBT (Gaithersburg, MD)).

The characteristics of the antigens of the different orthoebolavirus species used are as follows:

Ebola virus (EBOV) NP (amino acids [aa] 488 to 739 strain Mayinga 1976), VP40 (aa 31 to

326, strain Kissidougou-Makona 2014), GP K (aa 1 to 650, Kissidougou-Makona 2014), GP

M (aa 1 to 650, Mayinga 1976); Sudan virus (SUDV) NP (aa 361-738, strain Gulu), GP (aa 1

to 637, strain Uganda 2000), and VP40 (aa 31 to 326, strain Gulu); Bundibugyo virus (BDBV)

GP (aa 1 to 501, strain Uganda 2007), and VP40 (aa 31 to 326, strain Uganda 2007); and Reston

virus (RESTV) GP (aa 1 to 650). DBS were eluted and tested as previously described in 200μL

of incubation buffer (8,24,28,29). One hundred μL of sample, corresponding to the equivalent

of a final plasma dilution of 1/2000, was incubated with 50μL of magnetic beads coated with

recombinant protein (2 μg protein/1.25 × 106 beads) in 96-well flat-bottom chimney plates

(Greiner bio one, Frickenhausen, Germany) on a plate shaker at 300 rpm for 16 h at 4 °C in the

dark. After washing, 0.1 μg/mL of goat anti-bat biotin-labeled IgG (Euromedex,

Souffelweyersheim, France) was added to each well and incubated for 30 min at 300 rpm at

room temperature. After washing, we added 50 μL of 4 μg/mL streptavidin-R-phycoerythrin

(Fisher Scientific/Life Technologies, Illkirch, France) per well and incubated for 10 min at 300

rpm at room temperature. Reactions were read with BioPlex-200 (BioRad, Marnes-la-Coquette,

France) or MagPix (Luminex, Austin, TX, US). At least 100 events were read for each bead set

on a Bioplex-200 instrument, and results were expressed as median fluorescence intensity

(MFI) per 100 beads. Samples that showed positive signals were repeated in order to validate

the results.

To interpret the serological results in the absence of true positive controls, MFI cutoff

values for each antigen were determined using four methods in De Nys et al. (8) (See Table S1

in Additional file 1). In this study we opted for the cutoff values resulting from the least

stringent method, i.e. the mean plus four times the standard deviation of MFIs on negative

control samples (M + 4SD) as calculated in De Nys et al. 2018, to search for serological patterns

through a multivariate statistical analysis. Negative control samples (n = 145) were from

captive-born insectivorous bat species (*Carollia perspicillata*, n=103) hosted at the Parc

Zoologique de Montpellier (Montpellier, France) and two frugivorous bat species (*Pteropus*

giganteus, n=19, *Rousettus aegyptiacus*, n=23) hosted at Wilhelma Zoo and Botanical Garden

(Stuttgart, Germany). Samples were considered seropositive for an antigen if the MFI values

were above the cutoff value for this antigen. Samples were considered seropositive for an

orthoebolavirus if MFI values were above the cutoff value for at least two antigens of this

specific orthoebolavirus.

Multivariate statistical analyses

Proportions of samples positive to at least one orthoebolavirus antigen or to minimum

two antigens of the same orthoebolavirus are presented in a descriptive way to provide a

synthetic overview of the data and serological reactivity. Multivariate analyses were performed

for orthoebolavirus seropositivity as well as separately for the 3 antigens for which most

reactive samples were observed. We developed Generalized Linear Mixed Models (GLMM)

to test the impact of several drivers on the probability of a bat being positive by serology for

orthoebolavirus antigens (lme4 package in R software). We used results of the serological tests

as the response variable with a binomial distribution, and species, sex, age and season as explanatory variables with fixed effects (See Table S2 from Additional file 1). To account for clustered samples collected during the same sampling session at the same site, we included sampling period as a random effect to control for repeated measures from the same trapping session.

In order to select the best model, we used the glmulti function (Rsoftware, glmulti package (30)) to rank all possible models combining the explanatory variables by the Akaike Information Criterion (AIC). We selected the models with the lowest AIC (See Table S3 in Additional file 1). The odds ratios were estimated for each explanatory variable. We also used the emmeans and ggeffects functions to obtain estimated marginal means for the response variable and visualize the effect of each explanatory variable. Analysis were carried out on R software 4.1.2.

PCR screening for filoviruses

Oral and rectal swabs from a subset of bats were selected according to the analysis of serological results, extracted and tested for filovirus RNA by a semi-nested PCR following the protocols described in Djomsi et al. 2022 (24). The PCR targeted a 630 bp fragment of the L gene using degenerated primers that detect a wide diversity of filoviruses, i.e., Filo-MOD-FWD: TITTYTCHVTICAAAAICAYTGGG and FiloL.conR: ACCATCATRTTRCTIGGRAAKGCTTT in round 1 and Filo-MOD-FWD: TITTYTCHVTI-CAAAAICAYTGGG, and Filo-MOD-RVS: GCYTCISMIAIIGTTTGIACATT in round 2 (7)

Results

Bat sampling and serological results

We analyzed blood samples collected longitudinally from 1,427 wild frugivorous bats: 196 *E. helvum*, 186 *H. monstrosus*, 380 *M. angolensis* and 665 *R. aegyptiacus* (Table 1 and Table S4 from Additional file 1). Descriptive analyses show a total of 336/1427 (24%) samples that are reactive with at least one antigen. Proportions of positive samples per bat species, sex, age class and season are presented in Table 2. A total of 24/1427 (1.7%) bats are reactive with minimum 2 antigens of a specific orthoebolavirus and are thus seropositive for at least one of the orthoebolavirus species tested for (Table 3). Orthoebolavirus seroprevalence is highest in the *R. aegyptiacus* population (3%).

Table 1. Bats sampled for ebolavirus serology by species, sex and age class

Species	Total no. sampled	Sex (%)			Age (%)			
		Male	Female	UN*	Juvenile	Immature adult	Mature adult	UN
Eidolon helvum	196	91 (46.4)	104 (53.1)	1 (0.5)	17 (8.7)	36 (18.4)	117 (59.7)	26 (13.3)
Hypsignathus monstrosus	186	97 (52.1)	86 (46.2)	3 (1.6)	42 (22.6)	8 (4.3)	126 (67.7)	10 (5.4)
Myonycteris angolensis	380	176 (46.3)	203 (53.4)	1 (0.3)	86 (22.6)	41 (10.8)	244 (64.2)	9 (2.4)
Rousettus aegyptiacus	665	287 (43.2)	366 (55)	12 (1.8)	223 (33.5)	54 (8.2)	362(54.4)	26 (3.9)
Total	1427	651 (45.6)	759 (53.2)	17 (1.2)	368 (25.8)	139 (9.7)	849 (59.5)	71 (5)

* UN = undetermined

Table 2. Number and percentage of samples positive for at least 1 of the ebolavirus antigens according to the M+4SD cutoff method

Species	Total		
	Tot.	Pos.	% (95% CI)
*E. helvum**	196	69	**35** (29-42)
H. monstrosus	186	17	**9** (6-14)
M. angolensis	380	37	**10** (7-13)
R. aegyptiacus	665	213	**32** (29-36)
Total	1427	336	**24** (21-26)

Season	Dry			Wet		
	Tot.	Pos.	% (95% CI)	Tot.	Pos.	% (95% CI)
E. helvum	195	69	**35** (29-42)	1	0	**0** (0-95)
H. monstrosus	173	14	**8** (5-13)	13	3	**23** (8-50)
M. angolensis	183	8	**4** (2-8)	197	29	**15** (10-20)
R. aegyptiacus	388	148	**38** (33-43)	277	65	**23** (19-29)
Total	939	239	**25** (23-28)	488	97	**20** (17-24)

Sex	Male			Female			UN		
	Tot.	Pos.	% (95% CI)	Tot.	Pos.	% (95% CI)	Tot.	Pos.	% (95% CI)
E. helvum	91	43	**47** (37-57)	104	25	**24** (17-33)	1	1	**100** (5-100)
H. monstrosus	97	9	**9** (5-17)	86	8	**9** (5-17)	3	0	**0** (0-56)
M. angolensis	176	16	**9** (6-14)	203	21	**10** (7-15)	1	0	**0** (0-95)
R. aegyptiacus	287	100	**35** (30-41)	366	107	**29** (25-34)	12	6	**50** (25-75)
Total	651	168	**26** (23-29)	759	161	**21** (18-24)	17	7	**40** (22-64)

Age	Juvenile			Immature adult			Mature adult			UN		
	Tot.	Pos.	% (95% CI)	Tot.	Pos.	% (95% CI)	Tot.	Pos.	% (95% CI)	Tot.	Pos.	% (95% CI)
E. helvum	17	8	**47** (26-69)	36	15	**42** (27-58)	117	34	**29** (22-38)	26	12	**46** (29-65)
H. monstrosus	42	2	**5** (1-16)	8	1	**13** (1-47)	126	14	**11** (7-18)	10	0	**0** (0-28)
M. angolensis	86	7	**8** (4-16)	41	8	**20** (10-34)	244	19	**8** (5-12)	9	3	**33** (12-65)
R. aegyptiacus	223	58	**26** (21-32)	54	21	**39** (27-52)	362	121	**33** (29-38)	26	13	**50** (32-68)
Total	368	75	**20** (17-25)	139	45	**32** (25-41)	849	188	**22** (19-25)	71	28	**39** (29-51)

* *E.* =*Eidolon, H.= Hypsignathus, M. = Myonycteris, R. = Rousettus*, Pos. = positive, Tot. = total, UN =undetermined, CI = Confidence Interval

Table 3. Number and percentage of bats seropositive for orthoebolaviruses (positive for minimum two antigens of the same orthoebolavirus according to the M+4SD cutoff method)

		EBOV		SUDV		BDBV		Seropositive for min. 1 orthoebolavirus	
	Tot.	Pos.	% (95% CI)	Pos.	% (95% CI)	Pos.	% (95% CI)	Pos.	% (95% CI)
*E. helvum**	196	0	**0** (0-1.9)	2	**1** (0.3-3.6)	0	**0** (0-1.9)	2	**1** (0.3-3.6)
H. monstrosus	186	0	**0** (0-2)	0	**0** (0-2)	0	**0** (0-2)	0	**0** (0-2)
M. angolensis	380	0	**0** (0-1)	2	**0.5** (0.1-1.9)	0	**0** (0-1)	2	**0.5** (0.1-1.9)
R. aegyptiacus	665	9	**1.4** (0.7-2.6)	13	**2** (1.1-3.3)	1	**0.2** (0-0.8)	20	**3** (2-4.6)
Total	1427	9	**0.6** (0.3-1.2)	17	**1.2** (0.7-1.9)	1	**0.1** (0-0.4)	24	**1.7** (1.1-2.5)

* *E.* =*Eidolon, H.= Hypsignathus, M. = Myonycteris, R. = Rousettus*, Pos. = positive, Tot. = total, min. = minimum, CI = Confidence Interval

Regarding seroprevalences for each antigen of the different viruses, GPs were the recombinant proteins for which most reactions were observed: 9.8% of the samples showed reactivity against the EBOV Kissidougou strain, 18.8% against SUDV and 9.6% against BDBV versus 0.1 to 3.3 % against other antigens (Table 4). Lower reactivity for VP40 and NP antigens was observed for each bat species. Simultaneous reactivity to GP and NP of the same virus was rare and observed for 0.1% of the samples for EBOV and 0.6% for SUDV. The simultaneous reactivity to the same antigen of different viral lineages was frequent: 53% of the samples were reactive to GP of at least two orthoebolavirus species, 23% to VP40 and no cross-reactivity was observed for NP (See Table S5 from Additional file 1).

Table 4. Number (percentage) of bat blood samples reactive for each orthoebolavirus antigen according to the M+4SD cutoff method

Orthoebolavirus	Antigen*	E. helvum (n=196)		H. monstrosus (n=186)		M. angolensis (n=380)		R. aegyptiacus (n=665)		Total (n=1427)	
		no. (%)	95% CI	no. (%)	95% CI	no. (%)	95% CI	no. (%)	95% CI	no. (%)	95% CI
Ebola											
	NP	1 (0.5)	0.03-3	1 (0.5)	0.03-3	1 (0.3)	0.01-1	11 (2)	0.9-3	14 (1)	0.6-2
	GP K	35 (18)	13-24	5 (3)	1-6	7 (2)	0.9-4	93 (14)	12-17	140 (10)	8-11
	GP M	10 (5)	3-9	0	0-2	0	0-1	37 (6)	4-8	47 (3)	2-4
	VP40	1 (0.5)	0.3-3	0	0-2	1 (0.2)	0.01-1	17 (3)	2-4	19 (1)	0.9-2
	NP+GP	0	0-2	0	0-2	0	0-1	2 (0.3)	0.08-1	2 (0.1)	0.04-0.5
Sudan											
	NP	1 (0.5)	0.03-3	1 (0.5)	0.03-3	7 (2)	0.9-4	19 (3)	2-4	28 (2)	1-3
	GP	66 (34)	27-41	14 (8)	5-12	18 (5)	3-7	171(26)	23-29	269 (19)	17-21
	VP40	3 (2)	0.5-4	0	0-2	3 (0.8)	0.3-2	18 (3)	2-4	24 (2)	1-2
	NP+GP	1 (0.5)	0.03-3	0	0-2	1 (0.3)	0.01-1	7 (1)	0,5-2	9 (0.6)	0.3-1
Bundibugyo											
	GP	32 (16)	12-22	2 (1)	0.3-4	15 (4)	2-6	88 (13)	11-16	137 (10)	8-10
	VP40	0	0-2	0	0-2	1 (0.3)	0.01-2	7 (1)	0.5-2	8 (0.6)	0.3-1
Reston											
	GP	0	0-2	1 (0.5)	0.03-3	0	0-2	1 (0.2)	0.008-0.8	2 (0.1)	0.04-0.5

*NP = nucleoprotein; GP = glycoproteine (K = Kissidougou strain; M = Mayinga strain); VP = Viral protein

Factors influencing the presence of anti-orthoebolavirus antibodies in bats

We performed analysis on orthoebolavirus seropositivity, as well as on the presence of antibodies against the following 3 antigens: GP Ebola virus Kissidougou (GP EBOV-K), GP

Sudan virus (GP SUDV) and GP Bundibugyo virus (GP BDBV). The GLMM was not

converging for orthoebolavirus seropositivity, due to the low seroprevalence rates. However,

for each separate GP antigen we obtained fairly similar best fitted models (Table 5), i.e. the

Species, Sex and Age variables were consistently selected in the best models for each antigen.

The variable Season was also selected in the best models for GP EBOV-K and GP BDBV and

models with or without Season were near to equivalent for GP SUDV (Table 5).

276
277 Table 5. Best models obtained for the presence of antibodies against GP EBOV Kissidougou, GP BDBV and the
278 2 best models for GP SUDV
279

Ranking	Model								AIC
1	GP_EBOV-K ~ 1	+	Species	+	Age	+	Sex	+ Season	736.2752
1	GP_BDBV ~ 1	+	Species	+	Age	+	Sex	+ Season	740.2046
1	GP_SUDV ~ 1	+	Species	+	Age	+	Sex		1136.707
2	GP_SUDV ~ 1	+	Species	+	Age	+	Sex	+ Season	1136.824

281
Odds ratios (OR) for each explanatory variable selected in the best models are

displayed with confidence intervals in Table 6. Interestingly, *H. monstrosus* and *M. angolensis*

had consistently and significantly lower GP seroprevalences than *E. helvum* (OR ranging from

0.08 to 0.19 and 0.11 to 0.30 respectively, $p < 0.002$). Immature adults had consistently higher

GP seroprevalences than juveniles in all models (OR ranging from 1.84 to 2.33), although it

was only significant for GP EBOV-K and GP SUDV models ($p < 0.021$) but not for GP BDBV

($p = 0.069$). Males had consistently and significantly higher GP seroprevalences than females

for all the models (OR ranging from 1.56 to 1.72, $p < 0.034$).

294 Table 6. Regression table representing the odds ratios of the best statistical models for GP EBOV-K, GP BDBV
295 and GP SUDV
296

	GP EBOV-K* AIC=736.3			GP BDBV AIC=740.2			GP SUDV AIC=1136.7		
Variables	OR	95% CI	p-value*	OR	95% IC	p-value*	OR	95% IC	p-value*
Species									
E. helvum	—	—		—	—		—	—	
H. monstrosus	0.18	0.07, 0.49	<0.001	0.08	0.02, 0.36	<0.001	0.19	0.10, 0.37	<0.001
M. angolensis	0.13	0.05, 0.34	<0.001	0.30	0.14, 0.64	0.002	0.11	0.06, 0.20	<0.001
R. aegyptiacus	1.21	0.67, 2.17	0.5	1.31	0.73, 2.32	0.4	0.76	0.48, 1.21	0.3
Sex									
Female	—	—		—	—		—	—	
Male	1.72	1.15, 2.57	0.008	1.56	1.03, 2.36	0.034	1.70	1.10, 2.30	<0.001
Age									
Juvenile	—	—		—	—		—	—	
Immature	2.33	1.17, 4.62	0.016	1.91	0.95, 3.83	0.069	1.84	1.10, 3.10	0.021
Mature	1.55	0.94, 2.54	0.085	1.40	0.82, 2.39	0.2	1.03	0.72, 1.48	0.9
Season									
Wet	—	—		—	—				
Dry	2.10	1.01, 4.36	0.046	1.72	1.02, 2.92	0.043			

*significant <0.05; GP = glycoprotein; EBOV-K = Ebola virus Kissidougou strain; BDBV = Bundibugyo virus; SUDV = Sudan virus;

OR = Odds Ratio; CI = Confidence Interval

Similarly, the estimated marginal means of GP seroprevalences were significantly

higher for *R. aegyptiacus* and *E. helvum* compared to *H. monstrosus* and *M. angolensis* (Figure

2 and Additional file 1 Table S6) and for males compared to females (Figure 3), whatever the

antigen. GP seroprevalences were significantly higher for immature adults compared to

juveniles for GP EBOV-K and compared to juveniles and adults for GP SUDV (Figure 4). The

GP seroprevalences were also significantly higher in the dry season compared to the rainy

season for GP EBOV-K and GP BDBV (Figure 5).

Figure 2. Estimated marginal means (emm) of bat seroprevalence depending on the species and the antigens (estimated marginal means annotated with different letters are significantly different ($p < 0.05$))

Figure 3. Estimated marginal means of bat seroprevalence depending on the sex and the antigens (estimated marginal means annotated with different letters are significantly different ($p < 0.05$))

319 **Figure 4.** Estimated marginal means of bat seroprevalence depending on the age class and the antigens (estimated
320 marginal means annotated with different letters are significantly different (p < 0.05))
321

322
323 **Figure 5.** Estimated marginal means of bat seroprevalence depending on the season and the antigens (estimated
324 marginal means annotated with different letters are significantly different (p < 0.05))
325

 Filovirus RNA detection

Given the results of the serological and statistical analysis, oral (195) and rectal (141) swabs from 142 *R. aegyptiacus* and 53 *E. helvum* were screened by PCR. Juveniles and immature adults, as well as a subset of adults which were serologically positive to at least one antigen were selected (see Table S7 from Additional file 1). In addition, 64 *H. monstrosus* (64 oral and 63 rectal swabs) as well as 75 *M. angolensis* (75 oral and 75 rectal swabs) were also tested given that *H. monstrosus* is one of the 3 African fruit bat species in which viral EBOV RNA has been detected in previous studies (6) and that antibodies against orthoebolaviruses were detected in both species. All samples were negative.

Discussion

Although a number of cross-sectional studies and significant sampling efforts show the presence of antibodies against orthoebolaviruses in several bat species (8,14,28), only one reported detection of Ebola virus RNA in three fruit bats species (6). This low detection rate of viral genetic material makes it difficult to understand the dynamics of orthoebolavirus circulation in bats, the role of bats in orthoebolavirus maintenance and inter-species transmissions and related zoonotic risks. Antibody detection remains therefore a valuable indirect indicator of orthoebolavirus circulation in bats even if its interpretation remains challenging (8,31). Lack of positive controls, standardized cut-off determination methods, serological cross-reactivity, knowledge on antibody response dynamics in bats and viral molecular evidence generally complicate interpretation of results.

Here we opted for a longitudinal sampling strategy which allowed us to gain further insight into dynamics of orthoebolaviruses or closely related filoviruses in fruit bats, which is valuable for guiding further research efforts and surveillance strategies. Serological results are

in line with cross-sectional studies carried out in the same region (8,32), confirming

seropositivity for several orthoebolavirus antigens in the four bat species of interest, the

proportion of samples reacting with at least one antigen (23,5% versus 18,2% and 18,9% for

the least stringent cut-off) and simultaneous reactivity against same antigens of different

orthoebolavirus species. Antibody kinetics in human EBOD survivors have been shown to vary

for GP and NP antigens, with NP-antibodies declining faster (33). This might contribute to the

higher seroprevalence observed here for antibodies against GP antigens, and the rarity of

simultaneous reactivity to GP and NP antigens, also found in other studies (8,32). The lack of

specificity due to antibody cross-reactivity between filoviruses and the use of single GP

antigens per virus for statistical analysis might constitute limitations of this study. However,

clear and biologically meaningful patterns are observed with the methodology used. This

suggests that the serological results for the GP antigens as interpreted here do reflect past or

present infection with orthoebolavirus species tested for in this study or with one or several

other, perhaps not yet identified, filoviruses which induce antibodies that cross-react with

known orthoebolavirus antigens.

Our results show that the presence of antibodies against orthoebolavirus GP antigens

in fruit bats is best explained by bat species, sex, age as well as season. Best fitted models were

almost identical regardless of the orthoebolavirus tested for (GP EBOV-K, GP SUDV or GP

BDBV). We opted for interpretation of results in terms of orthoebolavirus GP in general and

not specific to orthoebolavirus species as the homogeneity of the model outputs for the different

orthoebolavirus GP antigens might be linked to cross-reactivity between GP antigens, as has

been described in previous studies (8,29,31).

E. helvum and *R. aegyptiacus* consistently showed significantly higher

orthoebolavirus GP seroprevalences than *H. monstrosus* and *M. angolensis* and *R. aegyptiacus*

had the highest seroprevalence when taking into account minimum 2 antigens per virus. *R.*

aegyptiacus are reservoir hosts of Marburg virus, a filovirus also known to circulate in West

Africa (17,34), and have been experimentally shown to be less likely to maintain

orthoebolaviruses (35). One explanation for our serological findings in *R. aegyptiacus* might

be that they reflect Marburg virus infection through IgG cross-reactivity with orthoebolavirus

antigens rather than infection with an orthoebolavirus. Serological IgG cross-reactivity between

orthoebolaviruses and Marburg virus has experimentally been shown to be very limited in bats

using indirect ELISA with infectious-based filovirus antigens (31). However, we used

recombinant viral protein antigens for our multiplex serological assays, hence cross-reaction of

anti-Marburg virus IgG might be higher compared to assays based on infectious-based antigens

(31). An alternative explanation for our serological results in *R. aegytpiacus* could be regular

exposure of *R. aegyptiacus* to known or unknown orthoebolaviruses through other infected

animals, leading to antibody response, and potential low-level replication followed by rapid

viral clearance. *R. aegyptiacus* is actually accountable for the highest prevalence of

orthoebolavirus seropositive bats (minimum 2 antigens) of our study. Despite the fact that *R.*

aegyptiacus might not be able to maintain known orthoebolaviruses infection, they have

actually been shown to produce virus-specific IgG when challenged experimentally with

orthoebolaviruses (35). Moreover, inoculation experiments on *R. aegyptiacus* use viral strains

originating from human epidemics, which might have been subject to mutations during

passages on cells. The extent to which other potentially circulating orthoebolaviruses are able

to cause more or less persistent infection in this species is unknown. Our results for *E. helvum*

are in line with other serological surveys which have detected antibodies against

orthoebolavirus GP in colonies from various geographical sites (8,24,36). An experimental

study has shown reduced susceptibility of Eidolon cells to EBOV (37). However, susceptibility

to other filoviruses might vary. In the same way as for *R. aegyptiacus*, the detected antibody

response might reflect exposure to orthoebolavirus(es) followed by limited replication and

rapid viral clearance or a certain degree of infection by one or several other closely related

filoviruses.

Beyond species-specific physiology and immunity, bat behaviour and ecology certainly also

contribute to inter-species variability in antibody responses. Large social groups have been

shown to allow higher rates of intra-species contacts which, for some pathogens, might facilitate

transmission and increase prevalence (38). *E. helvum* and *R. aegyptiacus* are social species

living in large colonies (tens or hundreds of thousands of individuals), therefore offering a large

potential for viral transmission, while *M. angolensis* and *H. monstrosus* live in smaller groups

(less than fifty for *M. angolensis* and twenty for *H. monstrosus*) (26). *H. monstrosus* is known

to form leks which gather thousands of individuals during the mating season in central Africa

(39), but leks seem to be smaller and sparse in forested Guinea (field observation). *E. helvum*

migrate long distances and form a panmictic population across most of the African continent

(40), which might also favor repeated viral reintroductions in colonies. Specific feeding and

roosting habits might lead to different degrees of inter-species contacts with other bat or non-

bat hosts of orthoebolaviruses and thus viral exposure. As a matter of fact, *R. aegyptiacus* and

H. monstrosus were observed sharing the same feeding sites in Macenta (T. Pouliquen, pers.

comm.), forested Guinea, hence exposure of one of these species to filoviruses hosted by the

other followed by immune reaction cannot be excluded. Actually, specific Marburg antibodies

have been documented in *H. monstrosus* in Gabon (41). Understanding the role of bats in

orthoebolavirus maintenance and transmission requires a perspective at the community level

rather than species level. Further studies on bat demographics, ecology, behaviour as well as

identification of the viruses responsible for the detected antibody responses are still needed to

elucidate these species differences in seroprevalence and their role in viral maintenance and

transmission, particularly in Guinea.

This study suggests for the first time that male fruit bats are more likely to be

seropositive against orthoebolavirus GP than females despite species-specific sampling

limitations. This gender difference was however not observed in a longitudinal study on *E.*

helvum in Cameroon (24). Further longitudinal species-specific studies focusing on this

question are needed to investigate this in different species, and to explore the explanatory

hypothesis based on observations made on humans. Studies on human populations have shown

that EBOV frequently persists in male survivors' semen, sometimes for several years as

suggested by the recent epidemic recrudescence in Guinea (2,42). This might result in repeated

antigenic stimulation and slower decrease of antibodies (33,43). We cannot exclude that this

scenario also occurs in some bat species, perhaps at specific times, and underlies higher GP

seroprevalences in males (under the assumption that antibodies decline over time when there is

no persistence). Behaviour such as mating and aggressive contacts might also increase the

frequency of close interactions of males with conspecifics compared to females and favor

microbial transmissions, as has been observed for other host species (44). In the same line, a

recent study on astro- and paramyxovirus circulation in bats has shown higher viral detection

rates in males (45).

The effect of age on the probability of detecting anti-orthoebolavirus GP antibodies,

with peak seroprevalences in the age class of immature adults, is in line with Marburg virus

infection dynamics where viral infection peaks in older juvenile (6 months old) *R. aegyptiacus*

(23) as well as with orthoebolavirus serological profiles described in *E. helvum* in Cameroun

(24,32). Passive immunity, made up of maternal antibodies to Marburg, wanes in juvenile *R.*

aegyptiacus aged 4 to 5 months, making them vulnerable to infection with the virus (46). This

could also be the case for other filoviruses in fruit bats and explain increases in orthoebolavirus

GP seroprevalence in immature adults following peaks of viral infections once older juveniles

(4 to 6 months) are losing their passive immunity. Juvenile bats sampled in the present study

were at least several months old as they were flying independently, which makes this hypothesis

plausible in explaining the observed pattern of increased GP seroprevalence in immature adults

compared to these older juveniles. Regardless of the underlying mechanisms, and similarly to

what has been observed in *E. helvum* (24), results suggest that a seroconversion against

orthoebolavirus or closely related filovirus takes place during the transition between juvenile

and immature adults. The duration of immunity against orthoebolaviruses in bats being a

knowledge gap, the lower antibody prevalence observed here in mature adults compared to

immature adults, especially for GP SUDV, could suggest a waning in time of the antibody

response, with potential re-stimulation in certain individuals, as suggested in humans (43).

Finally, orthoebolavirus GP seroprevalences seem to be generally higher during the dry

season (November to April) compared to the rainy season. *E. helvum,* which have relatively

high seroprevalence rates, are almost absent of the study site during the rainy season due to

their migratory behaviour. Reported seroprevalences may thus also reflect bat population

dynamics. Moreover, this seasonal pattern is most probably also related to other factors which

we couldn't measure in this study hence caution is needed with interpretation of this result.

Food abundance such as fruits can be limited during the dry season. Consequently, this could

lead to an exacerbated intra and interspecific competition for food (47) and increased viral

transmission followed by increased seroprevalences. Moreover, bat species studied here present

one or several birth pulses during this period, and for instance new *E. helvum* recruits reach

about 6-7 months of age at the beginning of the dry season. Rainfall and bat reproduction are

probably strongly interlinked (22,48), hence seasonal effects detected here are likely to reflect

the effect of reproductive life-cycle (and thus also age) on seroprevalence. The data per bat

species was insufficient to identify temporal patterns at a finer scale or assess the effect of

parameters such as period of copulation, parturition, lactation and weaning for each bat species.

However, other studies show that these periods can correspond to increased risks of viral

transmissions or impact viral maintenance (41,49,50).

Despite important sampling efforts over time, sample size remains small in relation to

the size of bat populations, and species-specific sample size was too limited to go into further

analysis at the bat species level and at a finer temporal scale. Nevertheless, this study provides

valuable information to guide further research on filovirus circulation in wildlife as well as

surveillance in the study region. Due to potential serological cross-reactivity between

orthoebolavirus species (29,31,33), serological observations may reflect past infections with

various orthoebolaviruses or closely related filoviruses, including undiscovered viruses. The

regular discovery of novel filovirus species across continents such as new filoviruses in China

(51,52), Bombali virus in four African countries (7,53–55), Lloviu virus in three European

countries (18,56,57), suggests that the full filovirus diversity has only been partly explored so

far. Molecular data which could help determine the filovirus species responsible for the

antibodies detected in our study and confirm serological interpretation remains limited. Future

identification of the virus(es) underlying these serological signals requires sequencing of viral

genetic material and virus isolation, and is essential to understand filovirus maintenance,

transmission dynamics and risks of inter-species spillover or emergence. Our results suggest

that intensive sampling efforts to search for evidence of and characterize active viral infection

should include a strong focus on older juveniles as well as male bats. A special attention should

be given to *E. helvum* and *R. aegyptiacus* which present significantly higher seroprevalences

compared to the other bat species, without excluding research on other species. In the same

way, surveillance of spillover events related to filovirus circulation in bats should be reinforced

where and when circulation and risks might be enhanced: at the interface with bat colonies

when older juveniles are highly prevalent, especially *E. helvum* (during the dry season for

forested Guinea) and *R. aegyptiacus* (already demonstrated for Marburg virus in *R.*

aegyptiacus). Based on these hypotheses, we screened a selection of individuals for filovirus RNA, but all samples were negative. Given the observed GP seroprevalence rates, we assume that the failure to detect filovirus RNA might be explained by either very low viral shedding at time of sampling, the circulation of unknown filoviruses which might not be detected by the PCR protocol used, or inappropriate sample type. Viral load might indeed be generally higher in internal organs than in rectal and oral swabs and it might thus be necessary to use the above-mentioned criteria to target a subset of individuals for more invasive sampling. Screening of *H. monstrosus* and *M. angolensis* which showed lower GP seroprevalence rates also revealed negative results. Nevertheless, EBOV RNA has been detected in organ samples from *H. monstrosus* in the past (6). Isolating filovirus nucleic acid from bats remains a challenge and when detected, prevalence rates based on rectal and oral swabs are often low (e.g. 1, 6 % for BOBV in Goldstein et al. 2018 (7) and 0,7% for Marburg virus in Paweska et al. 2020 (58)). Better targeted and larger sample sizes, combined with metagenomic approaches might be necessary to increase probability of filovirus RNA detection. Further experimental studies can also provide a clear insight into host competence for maintenance and transmission of known viruses and be complementary to field studies (59).

Conclusions

This study improved the understanding of filovirus dynamics in frugivorous bats by using a longitudinal approach that helps to test the role of potentially influential factors. It emphasizes the importance of host reproductive seasonality as a driver of pathogen circulation and thus spillover risks, which should be considered when designing sampling protocols for surveillance and research. Data can for instance be used to update orthoebolavirus spillover risk maps already developed for forested Guinea (60). The study was implemented in an ecosystem exposed to intensive human transformation influencing viral ecology. Substantial sampling

efforts need to be pursued and combined with data from ecological and behavioural studies on

bats and inter-species relations to allow for a broader comprehension of the ecology of

filoviruses at the community level. These efforts should also be replicated in different social-

ecological ecosystems with different exposure to human footprint for a comparative analysis.

List of abbreviations

AIC: Akaike Information Criterion

BDBV: Bundibugyo Ebola virus

bp: base pairs

EBOD: Ebola disease

EBOV: Ebolavirus

ELISA: Enzyme-Linked Immuno-Sorbent Assay

emm: Estimated Marginal Means

GLMM: Generalized Linear Mixed Models

GP: Glycoprotein

GP-K or GP EBOV-K: Glycoprotein Ebola virus Kissidougou strain

GP-M or GP EBOV-M: Glycoprotein Ebola virus Mayinga strain

IgG: Immunoglobulin G

MFI: Median Fluorescence Intensity

M + 4SD: mean plus four times the Standard Deviation

NP: Nucleoprotein

OR: Odds Ratios

p: p value (significant < 0.05)

PCR: Polymerase Chain Reaction

RESTV: Reston Ebola virus

RNA: Ribonucleic acid

SUDV: Sudan Ebola virus

VP40: Viral protein 40

Declarations

Ethics approval and consent to participate

This study was approved by the Comité National d'Ethique pour la Recherche en Santé (CNERS) in Guinea (permit number 025/CNERS/20), and the Office Guinéen des Parcs Nationaux et Réserves de Faune (OGPRF) of the Ministère de l'Environnement, et du Développement de la République de Guinée (permit 017OGUIPAR/MEEF/2017).

Availability of data and materials

The datasets generated and analysed during the current study are available in the following repository: https://doi.org/10.18167/DVN1/KMKUQ8.

Competing interests

The authors declare that they have no competing interests.

Funding

This study was supported by the projects Ebo-Sursy (https://rr-africa.oie.int/en/projects/ebo-sursy-en/) (FOOD/2016/379–660) and BCOMING (Horizon Europe project 101059483), both funded by the European Union, and the EbOHealth project, publicly funded by I-SITE MUSE (Montpellier Université d'Excellence https://muse.edu.umontpellier.fr/en/muse-isite - MUSE2018-EbOHEALTH) through the French National Research Agency (ANR) under the "Investissements d'avenir" program (grant number ANR-16-IDEX-0006).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Authors' contributions

MC has contributed to the design, acquisition, analysis, interpretation of the data, and drafting of the manuscript. AC, MB have contributed to the conception, design, analysis, interpretation and revision of the manuscript. AA, MP, AKK have contributed to the design, acquisition, analysis, interpretation of the data and revision of the manuscript. AS and MID have contributed to the data acquisition, interpretation and revision of the manuscript, TP and GT have contributed to the acquisition of the data and revision of the manuscript. JC and HDN have contributed to the conception, design, data acquisition, analysis, interpretation and drafting of the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We thank the Comité National d'Ethique pour la Recherche en Santé (CNERS) and the Office Guinéen des Parcs Nationaux et Réserves de Faune (OGPRF) of the Ministère de l'Environnement, et du Développement de la République de Guinée for authorising this study. We thank the Direction Nationale des Services Vétérinaires of the Ministère de l'Agriculture et de l'Elevage of Guinea, the OGPFR, and the CERFIG for their constant support and facilitation of the project, as well as all the field staff (Aboubacar Diallo, Pierre Kabinet Kamano, Mamadou Aliou Bah, Jean Paul Guilavogui, Mamadou Aliou Sow, Aboubacar Camara Bountouraby, Aboubacar Camara, Fassou Kourouma , KoïKoï Sakouvogui).

References

1. Centers for Disease Control and Prevention. 2022. [cited 2022 Feb 1]. History of Ebola Virus Disease (EVD) Outbreaks. Available from: https://www.cdc.gov/vhf/ebola/history/chronology.html#anchor_1526565058132

2. Keita AK, Koundouno FR, Faye M, Düx A, Hinzmann J, Diallo H, et al. Resurgence of Ebola virus in 2021 in Guinea suggests a new paradigm for outbreaks. Nature [Internet]. 2021 Sep 23 [cited 2021 Dec 28];597(7877):539–43. Available from: https://www.nature.com/articles/s41586-021-03901-9

3.   Mbala-Kingebeni P, Pratt C, Mutafali-Ruffin M, Pauthner MG, Bile F, Nkuba-Ndaye A, et al.
Ebola Virus Transmission Initiated by Relapse of Systemic Ebola Virus Disease. New England
Journal of Medicine [Internet]. 2021 Apr 1 [cited 2022 Feb 3];384(13):1240–7. Available
from: http://www.nejm.org/doi/10.1056/NEJMoa2024670

4.   Fairhead J, Leach M, Millimouno D. Spillover or endemic? Reconsidering the origins of Ebola
virus disease outbreaks by revisiting local accounts in light of new evidence from Guinea. BMJ
Glob Health [Internet]. 2021 Apr 1 [cited 2021 May 6];6(4):e005783. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/33893144

5.   Pigott DM, Golding N, Mylne A, Huang Z, Henry AJ, Weiss DJ, et al. Mapping the zoonotic
niche of Ebola virus disease in Africa. Elife. 2014;3:e04395.

6.   Leroy EM, Kumulungui B, Pourrut X, Rouquet P, Hassanin A, Yaba P, et al. Fruit bats as
reservoirs of Ebola virus. Nature [Internet]. 2005 Dec 1 [cited 2014 Jul 18];438(7068):575–6.
Available from: http://dx.doi.org/10.1038/438575a

7.   Goldstein T, Anthony SJ, Gbakima A, Bird BH, Bangura J, Tremeau-Bravard A, et al. The
discovery of Bombali virus adds further support for bats as hosts of ebolaviruses. Nat
Microbiol [Internet]. 2018 Oct 27 [cited 2018 Nov 12];3(10):1084–9. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/30150734

8.   De Nys HM, Kingebeni PM, Keita AK, Butel C, Thaurignac G, Villabona-Arenas CJ, et al.
Survey of Ebola Viruses in Frugivorous and Insectivorous Bats in Guinea, Cameroon, and the
Democratic Republic of the Congo, 2015–2017. Emerg Infect Dis [Internet]. 2018 Dec 17
[cited 2018 Nov 12];24(12). Available from: http://www.ncbi.nlm.nih.gov/pubmed/30307845

9.   Kamins AO, Rowcliffe JM, Ntiamoa-Baidu Y, Cunningham AA, Wood JLN, Restif O.
Characteristics and Risk Perceptions of Ghanaians Potentially Exposed to Bat-Borne Zoonoses
through Bushmeat. Ecohealth [Internet]. 2015 Mar 30 [cited 2017 Apr 6];12(1):104–20.
Available from: http://link.springer.com/10.1007/s10393-014-0977-0

10.   Leroy E, Gonzalez JP, Pourrut X. Ebolavirus and Other Filoviruses. In Springer, Berlin,
Heidelberg; 2007 [cited 2021 Dec 28]. p. 363–87. Available from:
http://link.springer.com/10.1007/978-3-540-70962-6_15

11.   Leendertz SAJ, Gogarten JF, Düx A, Calvignac-Spencer S, Leendertz FH. Assessing the
Evidence Supporting Fruit Bats as the Primary Reservoirs for Ebola Viruses. Ecohealth
[Internet]. 2016 Mar 13 [cited 2021 Dec 28];13(1):18–25. Available from:
http://link.springer.com/10.1007/s10393-015-1053-0

12.   Leroy EM, Rouquet P, Formenty P, Souquière S, Kilbourne A, Froment JM, et al. Multiple
Ebola virus transmission events and rapid decline of central African wildlife. Science
[Internet]. 2004 Jan 16 [cited 2016 Mar 23];303(5656):387–90. Available from:
http://science.sciencemag.org/content/303/5656/387.abstract

13.   Formenty P, Boesch C, Wyers M, Steiner C, Donati F, Dind F, et al. Ebola virus outbreak
among wild chimpanzees living in a rain forest of Cote d'Ivoire. J Infect Dis [Internet].
1999/02/13. 1999;179 Suppl:S120-6. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/9988175

14. Pourrut X, Délicat A, Rollin PE, Ksiazek TG, Gonzalez JP, Leroy EM. Spatial and temporal
patterns of Zaire ebolavirus antibody prevalence in the possible reservoir bat species. J Infect
Dis [Internet]. 2007 Nov 15 [cited 2017 Jun 30];196 Suppl 2(s2):S176-83. Available from:
https://academic.oup.com/jid/article-lookup/doi/10.1086/520541

15. Paweska JT, Storm N, Grobbelaar AA, Markotter W, Kemp A, Jansen van Vuren P.
Experimental Inoculation of Egyptian Fruit Bats (Rousettus aegyptiacus) with Ebola Virus.
Viruses [Internet]. 2016 Jan 22 [cited 2016 Feb 1];8(2):29. Available from:
http://www.mdpi.com/1999-4915/8/2/29/htm

16. Swanepoel R, Leman PA, Burt FJ, Zachariades NA, Braack LE, Ksiazek TG, et al.
Experimental inoculation of plants and animals with Ebola virus. Emerg Infect Dis [Internet].
1996 [cited 2016 Oct 21];2(4):321–5. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/8969248

17. Towner JS, Amman BR, Sealy TK, Carroll SAR, Comer JA, Kemp A, et al. Isolation of
Genetically Diverse Marburg Viruses from Egyptian Fruit Bats. Fouchier RAM, editor. PLoS
Pathog [Internet]. 2009 Jul 31 [cited 2016 Aug 30];5(7):e1000536. Available from:
http://dx.plos.org/10.1371/journal.ppat.1000536

18. Negredo A, Palacios G, Vázquez-Morón S, González F, Dopazo H, Molero F, et al. Discovery
of an Ebolavirus-Like Filovirus in Europe. Basler CF, editor. PLoS Pathog [Internet]. 2011 Oct
20 [cited 2016 Nov 8];7(10):e1002304. Available from:
http://dx.plos.org/10.1371/journal.ppat.1002304

19. Yang XL, Zhang YZ, Jiang RD, Guo H, Zhang W, Li B, et al. Genetically Diverse Filoviruses
in *Rousettus* and *Eonycteris* spp. Bats, China, 2009 and 2015. Emerg Infect Dis [Internet].
2017 Mar [cited 2017 Jun 1];23(3):482–6. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/28221123

20. Amman BR, Jones MEB, Sealy TK, Uebelhoer LS, Schuh AJ, Bird BH, et al. ORAL
SHEDDING OF MARBURG VIRUS IN EXPERIMENTALLY INFECTED EGYPTIAN
FRUIT BATS (*ROUSETTUS AEGYPTIACUS*). J Wildl Dis [Internet]. 2015 Jan [cited 2021
Jun 8];51(1):113–24. Available from: http://www.bioone.org/doi/10.7589/2014-08-198

21. Richter H V., Cumming GS. Food availability and annual migration of the straw-colored fruit
bat (Eidolon helvum). J Zool [Internet]. 2005 Dec 19 [cited 2018 Apr 13];268(1):35–44.
Available from: http://doi.wiley.com/10.1111/j.1469-7998.2005.00020.x

22. Cumming GS, Bernard RTF. Rainfall, food abundance and timing of parturition in African
bats. Oecologia [Internet]. 1997 Jul 18 [cited 2021 Dec 28];111(3):309–17. Available from:
http://link.springer.com/10.1007/s004420050240

23. Amman BR, Carroll SA, Reed ZD, Sealy TK, Balinandi S, Swanepoel R, et al. Seasonal Pulses
of Marburg Virus Circulation in Juvenile Rousettus aegyptiacus Bats Coincide with Periods of
Increased Risk of Human Infection. Kawaoka Y, editor. PLoS Pathog [Internet]. 2012 Oct 4
[cited 2016 Nov 3];8(10):e1002877. Available from:
http://dx.plos.org/10.1371/journal.ppat.1002877

24. Djomsi D, Mba Djonzo F, Ndong Bass I, Champagne M, Lacroix A, Thaurignac G, et al.
Dynamics of Antibodies to Ebolaviruses in an Eidolon helvum Bat Colony in Cameroon.

Viruses [Internet]. 2022 Mar 9 [cited 2022 Apr 9];14(3):560. Available from:
https://www.mdpi.com/1999-4915/14/3/560

25.   Maganga GD, Bourgarel M, Vallo P, Dallo TD, Ngoagouni C, Drexler JF, et al. Bat
Distribution Size or Shape as Determinant of Viral Richness in African Bats. Baker ML, editor.
PLoS One [Internet]. 2014 Jun 24 [cited 2021 Dec 28];9(6):e100172. Available from:
https://dx.plos.org/10.1371/journal.pone.0100172

26.   Rosevear DR. The bats of West Africa. London: British Museum (Natural History); 1965.

27.   Monadjem A, Taylor PJ, Cotterill FPD (Woody), Schoeman MC. Bats of Southern and Central
Africa [Internet]. Wits University Press; 2020 [cited 2023 Nov 30]. Available from:
http://www.jstor.org/stable/10.18772/22020085829

28.   Lacroix A, Mbala Kingebeni P, Ndimbo Kumugo SP, Lempu G, Butel C, Serrano L, et al.
Investigating the Circulation of Ebola Viruses in Bats during the Ebola Virus Disease
Outbreaks in the Equateur and North Kivu Provinces of the Democratic Republic of Congo
from 2018. Pathogens [Internet]. 2021 May 4 [cited 2021 May 6];10(5):557. Available from:
https://www.mdpi.com/2076-0817/10/5/557

29.   Ayouba A, Touré A, Butel C, Keita AK, Binetruy F, Sow MS, et al. Development of a
Sensitive and Specific Serological Assay Based on Luminex Technology for Detection of
Antibodies to Zaire Ebola Virus. Tang YW, editor. J Clin Microbiol [Internet]. 2017 Jan 26
[cited 2021 Dec 28];55(1):165–76. Available from:
https://journals.asm.org/doi/10.1128/JCM.01979-16

30.   Calcagno V, Mazancourt C de. glmulti : An R Package for Easy Automated Model Selection
with (Generalized) Linear Models. J Stat Softw [Internet]. 2010 May 31 [cited 2021 Dec
28];34(12):1–29. Available from: http://www.jstatsoft.org/v34/i12/

31.   Schuh AJ, Amman BR, Sealy TS, Flietstra TD, Guito JC, Nichol ST, et al. Comparative
analysis of serologic cross-reactivity using convalescent sera from filovirus-experimentally
infected fruit bats. Sci Rep [Internet]. 2019 Dec 30 [cited 2021 Dec 28];9(1):6707. Available
from: http://www.nature.com/articles/s41598-019-43156-z

32.   Peeters M, Champagne M, Ndong Bass I, Goumou S, Ndimbo Kumugo SP, Lacroix A, et al.
Extensive Survey and Analysis of Factors Associated with Presence of Antibodies to
Orthoebolaviruses in Bats from West and Central Africa. Viruses [Internet]. 2023 Sep 15 [cited
2023 Dec 14];15(9):1927. Available from: https://www.mdpi.com/1999-4915/15/9/1927

33.   Diallo MSK, Ayouba A, Keita AK, Thaurignac G, Sow MS, Kpamou C, et al. Temporal
evolution of the humoral antibody response after Ebola virus disease in Guinea: a 60-month
observational prospective cohort study. Lancet Microbe [Internet]. 2021 Dec 1 [cited 2022 Aug
29];2(12):e676–84. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S2666524721001701

34.   Amman BR, Bird BH, Bakarr IA, Bangura J, Schuh AJ, Johnny J, et al. Isolation of Angola-
like Marburg virus from Egyptian rousette bats from West Africa. Nat Commun [Internet].
2020 Jan 24 [cited 2023 Feb 28];11(1):510. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/31980636

35.   Jones M, Schuh A, Amman B, Sealy T, Zaki S, Nichol S, et al. Experimental Inoculation of
729   Egyptian Rousette Bats (Rousettus aegyptiacus) with Viruses of the Ebolavirus and
730   Marburgvirus Genera. Viruses [Internet]. 2015;7(7):3420–42. Available from:
731   http://www.mdpi.com/1999-4915/7/7/2779/

732   36.   Ogawa H, Miyamoto H, Nakayama E, Yoshida R, Nakamura I, Sawa H, et al.
733   Seroepidemiological Prevalence of Multiple Species of Filoviruses in Fruit Bats (Eidolon
734   helvum) Migrating in Africa. Journal of Infectious Diseases. 2015;212(Suppl 2):S101–8.

735   37.   Ng M, Ndungo E, Kaczmarek ME, Herbert AS, Binger T, Kuehne AI, et al. Filovirus receptor
736   NPC1 contributes to species-specific patterns of ebolavirus susceptibility in bats. Elife
737   [Internet]. 2015 Dec 23;4. Available from: https://elifesciences.org/articles/11785

738   38.   Nunn CL, Jordán F, McCabe CM, Verdolin JL, Fewell JH. Infectious disease and group size:
739   more than just a numbers game. Philosophical Transactions of the Royal Society B: Biological
740   Sciences [Internet]. 2015 May 26 [cited 2021 Dec 28];370(1669):20140111. Available from:
741   https://royalsocietypublishing.org/doi/10.1098/rstb.2014.0111

742   39.   Bradbury JW. Lek Mating Behavior in the Hammer-headed Bat. Z Tierpsychol [Internet]. 2010
743   Apr 26 [cited 2021 Dec 28];45(3):225–55. Available from:
744   https://onlinelibrary.wiley.com/doi/10.1111/j.1439-0310.1977.tb02120.x

745   40.   Peel AJ, Wood JLN, Baker KS, Breed AC, Carvalho A De, Fernández-Loras A, et al. How
746   Does Africa's Most Hunted Bat Vary Across the Continent? Population Traits of the Straw-
747   Coloured Fruit Bat (Eidolon helvum) and Its Interactions with Humans. Acta Chiropt
748   [Internet]. 2017 Jun 1 [cited 2023 Dec 8];19(1):77. Available from:
749   https://bioone.org/journals/acta-chiropterologica/volume-19/issue-
750   1/15081109ACC2017.19.1.006/How-Does-Africas-Most-Hunted-Bat-Vary-Across-the-
751   Continent/10.3161/15081109ACC2017.19.1.006.full

752   41.   Pourrut X, Souris M, Towner JS, Rollin PE, Nichol ST, Gonzalez JP, et al. Large serological
753   survey showing cocirculation of Ebola and Marburg viruses in Gabonese bat populations, and a
754   high seroprevalence of both viruses in Rousettus aegyptiacus. BMC Infect Dis [Internet]. 2009
755   Dec 28 [cited 2021 Dec 28];9(1):159. Available from:
756   https://bmcinfectdis.biomedcentral.com/articles/10.1186/1471-2334-9-159

757   42.   Thorson AE, Deen GF, Bernstein KT, Liu WJ, Yamba F, Habib N, et al. Persistence of Ebola
758   virus in semen among Ebola virus disease survivors in Sierra Leone: A cohort study of
759   frequency, duration, and risk factors. Kindrachuk J, editor. PLoS Med [Internet]. 2021 Feb 10
760   [cited 2021 Dec 28];18(2):e1003273. Available from:
761   https://dx.plos.org/10.1371/journal.pmed.1003273

762   43.   Adaken C, Scott JT, Sharma R, Gopal R, Dicks S, Niazi S, et al. Ebola virus antibody decay–
763   stimulation in a high proportion of survivors. Nature [Internet]. 2021 Feb 18 [cited 2021 Dec
764   28];590(7846):468–72. Available from: http://www.nature.com/articles/s41586-020-03146-y

765   44.   Schroeder P, Hopkins B, Jones J, Galloway T, Pike R, Rolfe S, et al. Temporal and spatial
766   Mycobacterium bovis prevalence patterns as evidenced in the All Wales Badgers Found Dead
767   (AWBFD) survey of infection 2014–2016. Sci Rep [Internet]. 2020 Sep 16 [cited 2023 Dec
768   8];10(1):15214. Available from: https://www.nature.com/articles/s41598-020-72297-9

769   45.   Hoarau AOG, Goodman SM, Al Halabi D, Ramasindrazana B, Lagadec E, Le Minter G, et al.
770   Investigation of astrovirus, coronavirus and paramyxovirus co-infections in bats in the western
771   Indian Ocean. Virol J [Internet]. 2021 Dec 12 [cited 2021 Dec 28];18(1):205. Available from:
772   https://virologyj.biomedcentral.com/articles/10.1186/s12985-021-01673-2

773   46.   Storm N, Jansen Van Vuren P, Markotter W, Paweska JT. Antibody Responses to Marburg
774   Virus in Egyptian Rousette Bats and Their Role in Protection against Infection. Viruses
775   [Internet]. 2018 [cited 2021 Dec 28];10(2). Available from:
776   http://www.ncbi.nlm.nih.gov/pubmed/29439384

777   47.   Buceta J, Johnson K, Jin Z, Wang Y, Wang Z, Drexler J. Modeling the Ebola zoonotic
778   dynamics: Interplay between enviroclimatic factors and bat ecology. Sun GQ, editor. PLoS
779   One [Internet]. 2017 Jun 12 [cited 2017 Jun 13];12(6):e0179559. Available from:
780   http://dx.plos.org/10.1371/journal.pone.0179559

781   48.   Happold DCD, Happold M. Reproductive strategies of bats in Africa. J Zool [Internet]. 1990
782   Dec 24 [cited 2021 Dec 28];222(4):557–83. Available from:
783   https://onlinelibrary.wiley.com/doi/10.1111/j.1469-7998.1990.tb06014.x

784   49.   Changula K, Kajihara M, Mori-Kajihara A, Eto Y, Miyamoto H, Yoshida R, et al.
785   Seroprevalence of Filovirus Infection of Rousettus aegyptiacus Bats in Zambia. J Infect Dis
786   [Internet]. 2018 Jun 8 [cited 2021 Oct 15];218(suppl_5):S312–7. Available from:
787   https://academic.oup.com/jid/advance-article/doi/10.1093/infdis/jiy266/5034832

788   50.   Baker KS, Suu-Ire R, Barr J, Hayman DTS, Broder CC, Horton DL, et al. Viral antibody
789   dynamics in a chiropteran host. Boots M, editor. Journal of Animal Ecology [Internet]. 2014
790   Mar 13 [cited 2021 Oct 15];83(2):415–28. Available from:
791   https://onlinelibrary.wiley.com/doi/10.1111/1365-2656.12153

792   51.   Yang XL, Tan CW, Anderson DE, Jiang RD, Li B, Zhang W, et al. Characterization of a
793   filovirus (Měnglà virus) from Rousettus bats in China. Nat Microbiol [Internet]. 2019 Jan 7
794   [cited 2023 Nov 17];4(3):390–5. Available from: https://www.nature.com/articles/s41564-
795   018-0328-y

796   52.   He B, Feng Y, Zhang H, Xu L, Yang W, Zhang Y, et al. Filovirus RNA in Fruit Bats, China. Emerg
797   Infect Dis [Internet]. 2015 Sep [cited 2023 Oct 3];21(9):1675–7. Available from:
798   http://wwwnc.cdc.gov/eid/article/21/9/15-0260_article.htm

799   53.   Kareinen L, Ogola J, Kivistö I, Smura T, Aaltonen K, Jääskeläinen AJ, et al. Range Expansion of
800   Bombali Virus in Mops condylurus Bats, Kenya, 2019. Emerg Infect Dis [Internet]. 2020 Dec
801   [cited 2021 Jan 28];26(12):3007–10. Available from:
802   http://www.ncbi.nlm.nih.gov/pubmed/33219788

803   54.   Karan LS, Makenov MT, Korneev MG, Sacko N, Boumbaly S, Yakovlev SA, et al. Bombali Virus
804   in *Mops condylurus* Bats, Guinea. Emerg Infect Dis [Internet]. 2019 Sep 17 [cited 2019 Aug
805   6];25(9). Available from: http://www.ncbi.nlm.nih.gov/pubmed/31310231

806   55.   Lebarbenchon C, Goodman SM, Hoarau AOG, Le Minter G, Dos Santos A, Schoeman MC, et al.
807   Bombali Ebolavirus in Mops condylurus Bats (Molossidae), Mozambique. Emerg Infect Dis

[Internet]. 2022 Dec [cited 2023 Nov 17];28(12):2583–5. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/36418002

56. Kemenesi G, Tóth GE, Mayora-Neto M, Scott S, Temperton N, Wright E, et al. Isolation of
infectious Lloviu virus from Schreiber's bats in Hungary. Nat Commun [Internet]. 2022 Dec 31
[cited 2022 Apr 4];13(1):1706. Available from: https://www.nature.com/articles/s41467-022-
29298-1

57. Tóth GE, Hume AJ, Suder EL, Zeghbib S, Ábrahám Á, Lanszki Z, et al. Isolation and genome
characterization of Lloviu virus from Italian Schreibers's bats. Sci Rep [Internet]. 2023 Jul 13
[cited 2023 Nov 17];13(1):11310. Available from: https://www.nature.com/articles/s41598-
023-38364-7

58. Pawęska JT, Storm N, Markotter W, Di Paola N, Wiley MR, Palacios G, et al. Shedding of
Marburg Virus in Naturally Infected Egyptian Rousette Bats, South Africa, 2017. Emerg Infect
Dis [Internet]. 2020 Dec [cited 2023 Feb 6];26(12):3051–5. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/33219802

59. Weber N, Nagy M, Markotter W, Schaer J, Puechmaille SJ, Sutton J, et al. Robust evidence for
bats as reservoir hosts is lacking in most African virus studies: a review and call to optimize
sampling and conserve bats. Biol Lett [Internet]. 2023 Nov [cited 2023 Dec
8];19(11):20230358. Available from: http://www.ncbi.nlm.nih.gov/pubmed/37964576

60. Lee-Cruz L, Lenormand M, Cappelle J, Caron A, De Nys H, Peeters M, et al. Mapping of Ebola
virus spillover: Suitability and seasonal variability at the landscape scale. Bird B, editor. PLoS
Negl Trop Dis [Internet]. 2021 Aug 23 [cited 2021 Dec 28];15(8):e0009683. Available from:
https://dx.plos.org/10.1371/journal.pntd.0009683

Additional file 1

PDF format .pdf

Methods S1. Molecular confirmation of bat species

Table S1. Mean fluorescence intensity cutoff values for different viral antigens, by
orthoebolavirus lineage and statistical method used to determine cutoff according to De Nys
et al, 2018.

Table S2. Characteristics of the variables used in the GLMM statistical models.

Table S3. Ranking of the top 5 best statistical models for the 3 glycoproteins by increasing
AIC value carried out on Rstudio software (version 1.4.1106)

Table S4. Longitudinal representation of the number of bats sampled per month for
orthoebolavirus serology (October 2018 - July 2020)

Table S5. Simultaneous reactivity of bat antibodies against antigens of different
orthoebolavirus lineages (M+4SD cutoff)

Table S6. p-values calculated with the estimated marginal means methods to compare

differences in antibody positivity according to age, sex, season and species (with the M+4SD

method) (significant p-value <0.05)

Table S7. Samples tested by PCR for filovirus RNA