Effects of biological and environmental factors on filovirus serology in bats

in Guinea

Maëliss Champagne, [1,2] Julien Cappelle, [1] Alexandre Caron, [1,3] Thibault Pouliquen, [1,4]

Aboubacar Samoura, [5] Mohamed Idriss Doumbouya, [6] Guillaume Thaurignac, [2] Ahidjo

Ayouba, [2] Alpha Kabinet Keita, [2,4] Martine Peeters, [2] Mathieu Bourgarel, [1,7] Hélène M. De

Nys [1,7]

1. ASTRE, CIRAD, INRAE, Université de Montpellier, Montpellier, France

2. TransVIHMI, Institut de Recherche pour le Développement (IRD), Université de

Montpellier, INSERM, Montpellier, France

3. Faculdade de Veterinária, Universidade Eduardo Mondlane, Maputo, Mozambique

4. Centre de Recherche et de Formation en Infectiologie de Guinée (CERFIG), Université

Gamal Abdel Nasser de Conakry, Conakry, Guinea

5. Office Guinéen des Parcs Nationaux et Réserves de Faune (OGPRF), Ministère de

l'Environnement, et du Développement Durable (MEDD), Conakry, Guinea

6. Direction Nationale des Services Vétérinaires, Ministère de l'Elevage et des Productions

Animales, Conakry, Guinea

7. CIRAD, UMR ASTRE, Harare, Zimbabwe

Correspondence should be addressed to Hélène M. De Nys; helene.de_nys@cirad.fr

Abstract

We investigated factors which might influence the circulation of ebolaviruses in fruit bats in Guinea in order to better understand their role in the ecology of ebolaviruses and evaluate risks of spill-over to other animals or humans. We performed serological tests on populations of 4 bat species (*Eidolon helvum*, *Hypsignathus monstrosus*, *Lissonycteris angolensis* and *Rousettus aegyptiacus*) sampled longitudinally for 18 months (2018-2020). The analysis of 1,427 bat samples for antibodies directed against different ebolavirus species allowed to test the influence of biological and environmental variables on seropositivity. Results showed that species (*Eidolon helvum and Rousettus aegyptiacus*), sex (males), age (sexually immature adults, and thus reproductive life-cycle), as well as season (dry) have a significant positive effect on the presence of antibodies against ebolaviruses in bats. These factors may play an important role in the circulation of filoviruses. Search for viral evidence and surveillance strategies should take these into consideration.

Introduction

Since the first two outbreaks of Ebola virus disease (EVD) in 1976 in Yambuku (DRC) and Nzara (South Sudan), 32 outbreaks have been officially identified in humans across Central and West Africa (1). The frequency of these outbreaks seems to have recently increased (1).

Despite recent epidemics which originated from viral resurgence in human populations (2–4), most outbreaks are believed to have resulted from independent zoonotic transmissions, through direct or indirect contact with the animal reservoir or other infected animals (5). The reservoir species of ebolaviruses remain unknown, but bats are suspected to play an important role in the ecology of the virus and transmission mechanisms (6–8). It is hypothesized that bats

might infect humans directly when hunted and handled (9), or indirectly through contact with

bat excrements or fruits contaminated with saliva or excrements (10). Bats might also be an

indirect source by transmitting viruses to bridge hosts which can act as amplifiers and be at the

origin of human outbreaks (6,11). This has been suspected in Central Africa and Ivory Coast

where apes have been confirmed as the source of infection in several outbreaks (12,13).

Certain bat species are regarded as potential reservoir of ebolaviruses since the

discovery of viral RNA and anti-Ebolavirus (anti-EBV) antibodies specific to the Zaire

ebolavirus in 3 fruit bat species (*Epomops franqueti*, *Hypsignathus monstrosus*, *Myonycteris*

torquata) (6). However, viral RNA was only detected in 13 of 279 tested specimens, and

confirmed by sequence analysis in 7. Other studies have since shown that other frugivorous

species have anti-EBV antibodies (*Rousettus aegyptiacus*, *Eidolon helvum*, *Epomophorus*

gambianus, *Micropteropus pusillus* and *Lissonycteris angolensis*) and that an insectivorous

species (*Mops sp.*) hosts a recently discovered ebolavirus named Bombali virus (7,8,14). This

supports the fact that these wild bats are exposed to filoviruses and are able to survive infection,

as demonstrated by experimental inoculations for some of these species (15,16). In addition,

Rousettus aegyptiacus are the reservoir species of Marburg virus, another filovirus (17), and

other bat species have been shown to be exposed or infected with other filoviruses such as

Lloviucuevavirus detected in Europe (18) and newly described viruses in China (19). Until

now, infectious ebolaviruses have never been isolated from bats. The viremic period might be

short or viruses might end up circulating at very low level and could therefore only be detected

for a limited period of time, as shown experimentally in *R. aegyptiacus* inoculated with

Marburg virus (15,20).

The aerial lifestyle, gregariousness, longevity as well as immunological and metabolic

specificities of bats are traits that may impact the ecology of ebolaviruses. Biological and

phenological factors, influenced by seasonal or climate changes, are also important, like the

migratory nature of certain bat species or reproduction patterns, both influenced by the

availability of food resources (21,22). It has been shown that spillover events of Marburg virus

to human populations coincide with the birthing season and presence of juveniles in *Rousettus*

aegyptiacus populations (23). A recent study on an *Eidolon helvum* colony also showed an

increase in the presence of antibodies against ebolaviruses when juveniles turn into immature

adults (24). Other factors can influence the dynamics of ebolaviruses in their natural hosts, for

instance population size and geographic distribution seem to impact viral richness within bat

populations (25). All these factors influence to some extent viral circulation, risk of spillover

events and emergence and are important to understand to adjust ebolaviruses research and EVD

surveillance strategies.

The objective of this study was to investigate the effects of biological and

environmental factors on the presence of anti-Ebola antibodies and improve understanding of

dynamics of ebolaviruses circulation in bats. We performed a longitudinal serological survey

of ebolaviruses in four frugivorous bat species in Guinea, a country at risk of EVD outbreak.

Based on the outcomes of the serological investigation, a subset of bats was also screened for

filoviruses RNA.

Methods

Sample and data collection

We collected blood samples as well as oral and rectal swabs from four frugivorous

bat species, i.e. *Eidolon helvum, Hypsignathus monstrosus, Lissonycteris angolensis and*

Rousettus aegyptiacus, already detected with anti-EBV antibodies (7). Capture and sampling

were carried out longitudinally over 13 sampling periods (i.e. 16 to 28 consecutive days of

sampling per period) between November 2018 and July 2020, following the protocols already

described in De Nys *et al.* (7). Bats were released directly after sampling. They were captured

in feeding and roosting areas located mainly in four study sites of the prefecture of Macenta

(Figure 1), in forested Guinea, the region where the 2013-2016 Ebola epidemic emerged. Sites

were visited every 2 to 4 weeks during the course of the study. The environment was mainly

constituted of highly transformed habitat, with secondary and degraded forests being gradually

converted into a mosaic of forests and crops over the last 20 years (L. Lee-Cruz, pers. Comm.).

It varied from secondary forest, fields, villages to more urbanised areas. Climate is

characterized by a wet (May to October) and a dry (November to April) season.

Capture site (geographic coordinates, ecological environment), morphology, sex, age

class, status of lactation and late gestation, and species (determined according to morphological

criteria combined to genetic confirmation for some specimens, Supporting information

Methods S1) were also recorded. Age was categorised as follows: juvenile (articular epiphyses

not yet fully ossified), sexually immature adult (articular epiphyses ossified, but

underdeveloped sexual organs), sexually mature adult (articular epiphyses ossified and clearly

developed sexual organs).

This study was approved by the Comité National d'Ethique pour la Recherche en

Santé of Guinea (permits 011/CNERS/19; 025/CNERS/20; 019/CNERS/20) and the Office

Guinéen des Parcs et Réserves (permit 017/OGUIPAR/MEEF/2017).

119
120
121

Figure 1. Sampling sites, A: Map of Guinea with the prefecture of Macenta in grey; B: prefecture of Macenta with sampling sites (dots) and number of sampled bats

Screening for Ebolaviruses Antibodies

We tested dried blood spots (DBS) for anti-EBVs IgG with a Luminex-based serologic assay adapted for bats as described in De Nys *et al*. (7). The assay included ten recombinant Ebola virus proteins, i.e. the glycoprotein (GP), nucleoprotein (NP), and/or viral protein 40 (VP40), for different ebolaviruses: Ebola Zaire Kissidougou and Mayinga strains (NP, GP K, GP M, VP40), Sudan (NP, GP, VP40), Bundibugyo (GP, VP40) and Reston (GP). Recombinant proteins were produced in sf9 insect cell vectors and purchased at Sinobiologicals (Beijing, China) except for REST GP (IBT (Gaithersburg, MD)). The characteristics of the antigens of the different ebolavirus species used are as follows: EBOV, previously called Zaire (NP, amino acids [aa] 488 to 739 strain Mayinga 1976; VP40, aa 31 to 326 strain Kissidougou-Makona 2014; GP-k (aa 1 to 650, Kissidougou-Makona 2014) GP-m (aa 1 to 650, Mayinga 1976); SUDV (NP (aa 361-738, strain Gulu), GP, aa 1 to 637, strain Uganda 2000; and VP40, aa 31 to 326, strain Gulu,); BDBV, Bundibugyo (GP, aa 1 to 501, strain Uganda 2007, and VP40, aa 31 to 326, strain Uganda 2007) and RESTV, Reston (GP, aa 1 to

650). DBS were eluted and tested as previously described in 200µL of incubation buffer (7,24,26,27). One hundred µL of sample, corresponding to the equivalent of a final plasma dilution of 1/2000, was incubated with 50µL of magnetic beads coated with recombinant protein (2 µg protein/1.25×10^6 beads) in 96-well flat-bottom chimney plates (Greiner bio one, Frickenhausen, Germany) on a plate shaker at 300 rpm for 16 h at 4 °C in the dark. After washing, 0.1 µg/mL of goat anti-bat biotin-labeled IgG (Euromedex, Souffelweyersheim, France) was added to each well and incubated for 30 min at 300 rpm at room temperature. After washing, we added 50 µL of 4 µg/mL streptavidin-R-phycoerythrin (Fisher Scientific/Life Technologies, Illkirch, France) per well and incubated for 10 min at 300 rpm at room temperature. Reactions were read with BioPlex-200 (BioRad, Marnes-la-Coquette, France) or MagPix (Luminex, Austin, TX, US). At least 100 events were read for each bead set on a Bioplex-200 instrument, and results were expressed as median fluorescence intensity (MFI) per 100 beads. Samples that showed positive signals were repeated in order to validate the results.

To interpret the serological results in the absence of true positive controls, MFI cutoff values for each antigen were determined using four methods in De Nys et al. (7) (See Table S1in the Supporting information). In this study we opted for the cutoff values resulting from the least stringent method, i.e. the mean plus four times the standard deviation of MFIs on negative control samples (M + 4SD) as calculated in De Nys et al. 2018, to search for serological patterns through a multivariate statistical analysis. Negative control samples (n = 145) were from captive-born insectivorous bat species (*Carollia perspicillata*, n=103) hosted at the Parc Zoologique de Montpellier (Montpellier, France) and two frugivorous bat species (*Pteropus giganteus*, n=19, *Rousettus aegyptiacus*, n=23) hosted at Wilhelma Zoo and Botanical Garden (Stuttgart, Germany). Samples were considered positive for the presence of antibodies against an antigen if the MFI values were above the cutoff value.

Multivariate statistical analyses

Proportions of samples positive to at least one antigen are presented in a descriptive way to provide a synthetic overview of the data and serological reactivity. Multivariate analyses were performed separately for the 3 antigens for which most reactive samples were observed. We developed Generalized Linear Mixed Models (GLMM) to test the impact of several drivers on the probability of a bat being positive by serology for EBV antigens (lme4 package in R software). We used results of the serological tests as the response variable with a binomial distribution, and species, sex, age and season as explanatory variables with fixed effects (See Table S2 from Supporting information). To account for clustered samples collected during the same sampling session at the same site, we included sampling period as a random effect to control for repeated measures from the same trapping session.

In order to select the best model, we used the glmulti function (Rsoftware, glmulti package (28)) to rank all possible models combining the explanatory variables by the Akaike Information Criterion (AIC). We selected the models with the lowest AIC (See Table S3 in Supporting information). The odds ratios were estimated for each explanatory variable. We also used the emmeans and ggeffects functions to obtain estimated marginal means for the response variable and visualize the effect of each explanatory variable. Analysis were carried out on R software 4.1.2.

PCR screening for filoviruses

Oral and rectal swabs from a subset of bats were selected according to the analysis of serological results, extracted and tested for filoviruses RNA by a semi-nested PCR following the protocols described in Djomsi et al. 2022 (24). The PCR targeted a 630 bp fragment of the L gene using degenerated primers that detect a wide diversity of filoviruses, i.e., Filo-MOD-FWD: TITTYTCHVTICAAAAICAYTGGG and FiloL.conR:

191 ACCATCATRTTRCTIGGRAAKGCTTT in round 1 and Filo-MOD-FWD: TITTYTCHVTI-

192 CAAAAICAYTGGG, and Filo-MOD-RVS: GCYTCISMIAIIGTTTGIACATT in round 2 (29)

Results

Bat sampling and serological results

We analyzed blood samples collected longitudinally from 1,427 wild frugivorous

bats: 196 *E. helvum*, 186 *H. monstrosus*, 380 *L. angolensis* and 665 *R. aegyptiacus* (Table 1

and Table S4 from Supporting information). Descriptive analyses show a total of 336/1427

(24%) samples that are reactive with at least one antigen. Proportions of positive samples per

bat species, sex, age class and season are presented in Table 2.

Table 1. Bats sampled for ebolavirus serology by species, sex and age class

Species	Total no. sampled	Sex (%)			Age (%)			
		Male	Female	UN*	Juvenile	Immature adult	Mature adult	UN
Eidolon helvum	196	91 (46.4)	104 (53.1)	1 (0.5)	17 (8.7)	36 (18.4)	117 (59.7)	26 (13.3)
Hypsignathus monstrosus	186	97 (52.1)	86 (46.2)	3 (1.6)	42 (22.6)	8 (4.3)	126 (67.7)	10 (5.4)
Lissonycteris angolensis	380	176 (46.3)	203 (53.4)	1 (0.3)	86 (22.6)	41 (10.8)	244 (64.2)	9 (2.4)
Rousettus aegyptiacus	665	287 (43.2)	366 (55)	12 (1.8)	223 (33.5)	54 (8.2)	362 (54.4)	26 (3.9)
Total	1427	651 (45.6)	759 (53.2)	17 (1.2)	368 (25.8)	139 (9.7)	849 (59.5)	71 (5)

204 * UN = undetermined

Table 2. Number and percentage of samples positive for at least 1 of the ebolavirus antigens according to the M+4SD cutoff method

Species	Total		
	Tot.	Pos.	% (95% CI)
*E. helvum**	196	69	**35** (29-42)
H. monstrosus	186	17	**9** (6-14)
L. angolensis	380	37	**10** (7-13)
R. aegyptiacus	665	213	**32** (29-36)
Total	1427	336	**24** (21-26)

Season	Dry			Wet		
	Tot.	Pos.	% (95% CI)	Tot.	Pos.	% (95% CI)
E. helvum	195	69	**35** (29-42)	1	0	**0** (0-95)
H. monstrosus	173	14	**8** (5-13)	13	3	**23** (8-50)
L. angolensis	183	8	**4** (2-8)	197	29	**15** (10-20)
R. aegyptiacus	388	148	**38** (33-43)	277	65	**23** (19-29)
Total	939	239	**25** (23-28)	488	97	**20** (17-24)

Sex	Male			Female			UN		
	Tot.	Pos.	% (95% CI)	Tot.	Pos.	% (95% CI)	Tot.	Pos.	% (95% CI)
E. helvum	91	43	**47** (37-57)	104	25	**24** (17-33)	1	1	**100** (5-100)
H. monstrosus	97	9	**9** (5-17)	86	8	**9** (5-17)	3	0	**0** (0-56)
L. angolensis	176	16	**9** (6-14)	203	21	**10** (7-15)	1	0	**0** (0-95)
R. aegyptiacus	287	100	**35** (30-41)	366	107	**29** (25-34)	12	6	**50** (25-75)
Total	651	168	**26** (23-29)	759	161	**21** (18-24)	17	7	**40** (22-64)

Age	Juvenile			Immature adult			Mature adult			UN		
	Tot.	Pos.	% (95% CI)	Tot.	Pos.	% (95% CI)	Tot.	Pos.	% (95% CI)	Tot.	Pos.	% (95% CI)
E. helvum	17	8	**47** (26-69)	36	15	**42** (27-58)	117	34	**29** (22-38)	26	12	**46** (29-65)
H. monstrosus	42	2	**5** (1-16)	8	1	**13** (1-47)	126	14	**11** (7-18)	10	0	**0** (0-28)
L. angolensis	86	7	**8** (4-16)	41	8	**20** (10-34)	244	19	**8** (5-12)	9	3	**33** (12-65)
R. aegyptiacus	223	58	**26** (21-32)	54	21	**39** (27-52)	362	121	**33** (29-38)	26	13	**50** (32-68)
Total	368	75	**20** (17-25)	139	45	**32** (25-41)	849	188	**22** (19-25)	71	28	**39** (29-51)

* E. =Eidolon, H.= Hypsignathus, L. = Lissonycteris, R. = Rousettus, Pos. = positive, Tot. = total, UN =undetermined

Regarding seroprevalences for each antigen of the different viruses, GPs were the recombinant proteins for which most reactions were observed: 9.8% of the samples showed reactivity against the Zaire Kissidougou strain, 18.8% against Sudan and 9.6% against Bundibugyo versus 0.1 to 3.3 % against other antigens (Table 3). Lower reactivity for VP40 and NP antigens was observed for each bat species. Simultaneous reactivity to GP and NP of the same virus was rare and observed for 0.1% of the samples for EBOV and 0.6% for SUDV.

The simultaneous reactivity to the same antigen of different viral lineages was frequent: 53% of the samples were reactive to GP of at least two ebolavirus species, 23% to VP40 and no cross-reactivity was observed for NP (See Table S5 from Supporting information).

Table 3. Number (percentage) of bat blood samples reactive for each ebolavirus antigen according to the M+4SD cutoff method

Ebolavirus	Antigen*	E. helvum (n=196) no. (%)	95% CI	H. monstrosus (n=186) no. (%)	95% CI	L. angolensis (n=380) no. (%)	95% CI	R. aegyptiacus (n=665) no. (%)	95% CI	Total (n=1427) no. (%)	95% CI
Zaïre											
	NP	1 (0.5)	0.03-3	1 (0.5)	0.03-3	1 (0.3)	0.01-1	11 (2)	0.9-3	14 (1)	0.6-2
	GP K	35 (18)	13-24	5 (3)	1-6	7 (2)	0.9-4	93 (14)	12-17	140 (10)	8-11
	GP M	10 (5)	3-9	0	0-2	0	0-1	37 (6)	4-8	47 (3)	2-4
	VP40	1 (0.5)	0.3-3	0	0-2	1 (0.2)	0.01-1	17 (3)	2-4	19 (1)	0.9-2
	NP+GP	0	0-2	0	0-2	0	0-1	2 (0.3)	0.08-1	2 (0.1)	0.04-0.5
Sudan											
	NP	1 (0.5)	0.03-3	1 (0.5)	0.03-3	7 (2)	0.9-4	19 (3)	2-4	28 (2)	1-3
	GP	66 (34)	27-41	14 (8)	5-12	18 (5)	3-7	171 (26)	23-29	269 (19)	17-21
	VP40	3 (2)	0.5-4	0	0-2	3 (0.8)	0.3-2	18 (3)	2-4	24 (2)	1-2
	NP+GP	1 (0.5)	0.03-3	0	0-2	1 (0.3)	0.01-1	7 (1)	0,5-2	9 (0.6)	0.3-1
Bundibugyo											
	GP	32 (16)	12-22	2 (1)	0.3-4	15 (4)	2-6	88 (13)	11-16	137 (10)	8-10
	VP40	0	0-2	0	0-2	1 (0.3)	0.01-2	7 (1)	0.5-2	8 (0.6)	0.3-1
Reston											
	GP	0	0-2	1 (0.5)	0.03-3	0	0-2	1 (0.2)	0.008-0.8	2 (0.1)	0.04-0.5

*NP = nucleoprotein; GP = glycoproteine (K = Kissidougou strain; M = Mayinga strain); VP = Viral protein

Factors influencing the presence of anti-EBV antibodies in bats

We performed analysis on the presence of antibodies against the following 3 antigens: GP Zaïre Kissidougou (GP EBOV-K), GP Sudan (GP SUDV) and GP Bundibugyo (GP BDBV). For each antigen we obtained fairly similar best fitted models (Table 4), i.e. the Species, Sex and Age variables were consistently selected in the best models for each antigen. The variable Season was also selected in the best models for GP EBOV-K and GP BDBV and models with or without Season were near to equivalent for GP SUDV (Table 4).

Table 4. Best models obtained for the presence of antibodies against GP Zaire Kissidougou, GP Bundibugyo and the 2 best models for GP Sudan

Ranking	Model										AIC
1	GP_EBOV-K	~ 1	+	Species	+	Age	+	Sex	+	Season	736.2752
1	GP_BDBV	~ 1	+	Species	+	Age	+	Sex	+	Season	740.2046
1	GP_SUDV	~ 1	+	Species	+	Age	+	Sex			1136.707
2	GP_SUDV	~ 1	+	Species	+	Age	+	Sex	+	Season	1136.824

Odds ratios (OR) for each explanatory variable selected in the best models are displayed with Confidence Intervals in Table 5. Interestingly, *H. monstrosus* and *L. angolensis* had consistently and significantly lower seroprevalences than *E. helvum* (OR ranging from 0.08 to 0.19 and 0.11 to 0.30 respectively, $p < 0.002$). Immature adults had consistently higher seroprevalences than juveniles in all models (OR ranging from 1.84 to 2.33), although it was only significant for GP EBOV-K and GP-SUDV models ($p < 0.021$) but not for GP-BDBV ($p = 0.069$). Males had consistently and significantly higher seroprevalences than females for all the models (OR ranging from 1.56 to 1.72, $p < 0.034$).

Table 5. Regression table representing the odds ratios of the best statistical models for GP EBOV-K, GP BDBV and GP SUDV

Variables	GP EBOV-K* AIC=736.3			GP BDBV AIC=740.2			GP SUDV AIC=1136.7		
	OR	95% CI	p-value*	OR	95% IC	p-value*	OR	95% IC	p-value*
Species									
E. helvum	—	—		—	—		—	—	
H. monstrosus	0.18	0.07, 0.49	<0.001	0.08	0.02, 0.36	<0.001	0.19	0.10, 0.37	<0.001
L. angolensis	0.13	0.05, 0.34	<0.001	0.30	0.14, 0.64	0.002	0.11	0.06, 0.20	<0.001
R. aegyptiacus	1.21	0.67, 2.17	0.5	1.31	0.73, 2.32	0.4	0.76	0.48, 1.21	0.3
Sex									
Female	—	—		—	—		—	—	
Male	1.72	1.15, 2.57	0.008	1.56	1.03, 2.36	0.034	1.70	1.10, 2.30	<0.001
Age									
Juvenile	—	—		—	—		—	—	
Immature	2.33	1.17, 4.62	0.016	1.91	0.95, 3.83	0.069	1.84	1.10, 3.10	0.021
Mature	1.55	0.94, 2.54	0.085	1.40	0.82, 2.39	0.2	1.03	0.72, 1.48	0.9
Season									
Wet	—	—		—	—				
Dry	2.10	1.01, 4.36	0.046	1.72	1.02, 2.92	0.043			

*significant <0.05; GP = glycoprotein; EBOV-K = Zaire Ebolavirus Kissidougou strain; BDBV = Bundibugyo Ebolavirus; SUDV = Sudan Ebolavirus; OR = Odds Ratio; CI = Confidence Interval

Similarly, the estimated marginal means of seroprevalences were significantly higher for *R. aegyptiacus* and *E. helvum* compared to *H. monstrosus* and *L. angolensis* (Figure 2 and Supporting information Table S6) and for males compared to females (Figure 3), whatever the antigen. Seroprevalences were significantly higher for immature adults compared to juveniles for GP EBOV-K and compared to juveniles and adults for GP SUDV (Figure 4). The seroprevalences were also significantly higher in the dry season compared to the rainy season for GP EBOV-K and GP BDBV (Figure 5).

299
300 **Figure 2.** Estimated marginal means (emm) of bat seroprevalence depending on the species and the antigens
301 (estimated marginal means annotated with different letters are significantly different (p < 0.05))
302
303

304
305 **Figure 3.** Estimated marginal means of bat seroprevalence depending on the sex and the antigens (estimated
306 marginal means annotated with different letters are significantly different (p < 0.05))
307
308

309
310 **Figure 4.** Estimated marginal means of bat seroprevalence depending on the age class and the antigens (estimated
311 marginal means annotated with different letters are significantly different ($p < 0.05$))

312
313
314
315

316
317 **Figure 5.** Estimated marginal means of bat seroprevalence depending on the season and the antigens (estimated
318 marginal means annotated with different letters are significantly different ($p < 0.05$))

319
320
321
322

 Filoviruses RNA detection

324   Given the results of the serological and statistical analysis, oral (195) and rectal (141) swabs

325   from 142 *R. aegyptiacus* and 53 *E. helvum* were screened by PCR. Juveniles and immature

326   adults, as well as a subset of adults which were serologically positive to at least one antigen

327   were selected (see Table S7 from Supporting information). All samples were negative.

329   **Discussion**

331   Although a number of cross-sectional studies and significant sampling efforts show

the presence of antibodies against ebolavirus in several bat species (7,14,26), only one reported

detection of Zaire Ebola virus RNA in three fruit bats species (6). This makes it difficult to

understand the dynamics of ebolavirus circulation in bats, the role of bats in ebolavirus

maintenance and inter-species transmissions and related zoonotic risks. Here we opted for a

longitudinal sampling strategy which allowed to gain further insight into dynamics of

ebolaviruses or closely related filoviruses in fruit bats, which is valuable for guiding further

research efforts and surveillance strategies. Serological results are in line with a previous cross-

sectional study carried out in the same region (7), confirming seropositivity for ebolavirus

antigens in the four bat species of interest, the proportion of samples reacting with at least one

antigen (23,5% versus 18,2% for the least stringent cut-off in (7)) and simultaneous reactivity

against same antigens of different ebolavirus lineages. Antibody kinetics in human ebolavirus

disease survivors have been shown to vary for GP and NP antigens, with NP-antibodies

declining faster (30). This might contribute to the higher prevalence observed here for

antibodies against GP antigens, and the rarity of simultaneous reactivity to GP and NP antigens.

Lack of positive controls, standardized cut-off determination methods and knowledge on

antibody response dynamics in bats complicate interpretation of results. We are aware that the

risk of false positives might constitute a limitation of this study. However, clear and biologically

meaningful patterns are observed with the methodology used in this study, which suggests that

the serological results for the GP antigens as interpreted here do reflect past or present infection

with one of the ebolavirus species tested in this study or cross-reactivity with another filovirus

not yet identified.

353   Our results show that the presence of antibodies against ebolaviruses in fruit bats is

best explained by bat species, sex, age as well as season. Best fitted models were almost

identical regardless of the ebolavirus tested for (GP EBOV-K, GP SUDV or GP BDBV). We

opted for interpretation of results in terms of ebolaviruses in general and not specific to

ebolavirus species as the homogeneity of the model outputs for the different ebolaviruses might

be linked to cross-reactivity between GP antigens, as has been described in previous studies

(7,27,31). *E. helvum* and *R. aegyptiacus* consistently showed significantly higher

seroprevalences than *H. monstrosus* and *L. angolensis*. Large social groups have been shown

to allow higher rates of inter individual contacts which, for some pathogens, might facilitate

transmission and increase prevalence (32). *E. helvum* and *R. aegyptiacus* are social species

living in large colonies (tens or hundreds of thousands of individuals), therefore offering a large

potential for viral transmission, while *L. angolensis* and *H. monstrosus* live in smaller groups

(less than fifty for *L. angolensis* and twenty for *H. monstrosus*) (33). *H. monstrosus* is known

to form leks which gather thousands of individuals during the mating season in central Africa

(34), but leks seem to be smaller and sparse in forested Guinea (field observation). *E. helvum*

migrate long distances and form a panmictic population across most of the African continent

(35), which might also favor repeated viral reintroductions in colonies. Beyond these factors,

species-specific physiology and immunity certainly also contribute to these differences.

Important to note is that *R. aegyptiacus* are reservoir hosts of Marburg virus, a filovirus also

known to circulate in West Africa (17,36), and have been experimentally shown to be less likely

to maintain ebolavirus (37). One explanation for our serological findings in *R. aegyptiacus* might be that they reflect Marburg virus circulation through IgG cross-reactivity with ebolaviruses rather than infection with an ebolavirus. However, this remains questionable as serological IgG cross-reactivity between ebolaviruses and Marburg virus has experimentally been shown to be very limited in bats using indirect ELISA with infectious-based filovirus antigens (31). To bear in mind is that specific feeding and roosting habits might lead to different degrees of inter-species contacts with other bat or non-bat hosts of ebolaviruses and thus viral exposure. As a matter of fact, *R. aegyptiacus* and *H. monstrosus* were observed sharing the same feeding sites in Macenta (T. Pouliquen, pers. comm.), forested Guinea, hence exposure of one of these species to filoviruses hosted by the other followed by immune reaction cannot be excluded. Actually, specific Marburg antibodies have been documented in *H. monstrosus* in Gabon (38). Understanding the role of bats in ebolavirus maintenance and transmission requires a perspective at the community level rather than species level. Further studies on bat demographics, ecology and behavior are still needed to elucidate these differences in seroprevalence, particularly in Guinea.

This study suggests for the first time that male fruit bats are more likely to be seropositive against ebolaviruses than females. Given species specific sampling limitations, species-specific studies focusing on this question should be done to confirm this for different species, and to explore the explanatory hypothesis based on observations made on humans. Studies on human populations have shown that ebolavirus frequently persists in male survivors' semen, sometimes for several years as suggested the recent epidemic recrudescence in Guinea (2,39). This might result in repeated antigenic stimulation and slower decrease of antibodies (30,40). We cannot exclude that this scenario also occurs in some bat species, perhaps at specific times, and underlies higher seroprevalences in males (under the assumption that antibodies decline over time when there is no persistence). Behaviour such as mating and

aggressive contacts might also increase the frequency of close interactions of males with

conspecifics compared to females and favor viral transmissions. In the same line, a recent study

on astro- and paramyxovirus circulation in bats has shown higher viral detection rates in males

(41).

The effect of age on the probability of detecting anti-EBV antibodies, with peak

seroprevalences in the age class of immature adults, is in line with Marburg virus infection

dynamics where viral infection peaks in older juvenile (6 months old) *R. aegyptiacus* (23) as

well as with ebolavirus serological profiles described in *E. helvum* in Cameroun (24). Passive

immunity, made up of maternal antibodies to Marburg, wanes in juvenile *R. aegyptiacus* aged

4 to 5 months, making them vulnerable to infection with the virus (42). This could also be the

case for other filoviruses in fruit bats and explain increases in ebolavirus antibody

seroprevalence in immature adults following peaks of viral infections once older juveniles (4

to 6 months) are losing their passive immunity. Juvenile bats sampled in the present study were

at least several months old as they were flying independently, which makes this hypothesis

plausible in explaining the observed pattern of increased seroprevalence in immature adults

compared to these older juveniles. Regardless of the underlying mechanisms, and similarly to

what has been observed in *E. helvum* (24), results suggest that a seroconversion against

ebolavirus or closely related filoviruses takes place during the transition between juvenile and

immature adults. The duration of immunity against ebolaviruses in bats being a knowledge gap,

the lower antibody prevalence observed here in mature adults compared to immature adults,

especially for GP SUDV, could suggest a waning in time of the antibody response, with

potential re-stimulation in certain individuals, as suggested in humans (40).

Finally, seroprevalences seem to be generally higher during the dry season (November

to April) compared to the rainy season. *E. helvum,* which have relatively high sero-prevalence

rates, are almost absent of the study site during the rainy season due to their migratory

behaviour. Reported seroprevalences may thus also reflect bat population dynamics. Moreover, this seasonal pattern is most probably also related to other factors which we couldn't measure in this study hence caution is needed with interpretation of this result. Food abundance such as fruits can be limited during the dry season. Consequently, this could lead to an exacerbated intra and interspecific competition for food (43) and increased viral transmission followed by increased seroprevalences. Moreover, bat species studied here present one or several birth pulses during this period, and for instance new *E. helvum* recruits reach about 6-7 months of age at the beginning of the dry season. Rainfall and bat reproduction are probably strongly interlinked (22,44), hence seasonal effects detected here are likely to reflect the effect of reproductive life-cycle (and thus also age) on seroprevalence. The data per bat species was insufficient to identify temporal patterns at a finer scale or assess the effect of parameters such as period of copulation, parturition, lactation and weaning for each bat species. However, other studies show that these periods can correspond to increased risks of viral transmissions or impact viral maintenance (38,45,46).

Despite important sampling efforts over time, sample size remains small given large bat population sizes, and species-specific sample size was too limited to go into further analysis at the bat species level and at a finer temporal scale. Nevertheless, this study provides valuable information to guide further research on ebolavirus circulation in wildlife as well as surveillance in the study region. Due to potential serological cross-reactivity between ebolavirus species (27,30,31), serological observations may reflect past infections with various ebolaviruses or closely related filoviruses, including undiscovered viruses, and unfortunately virologic data which could help with serological interpretation remains limited. Future identification of the virus(es) underlying these serological signals is essential to understand filovirus maintenance, transmission dynamics and risks of inter-species spillover or emergence. Our results suggest that intensive sampling efforts to search for evidence of and characterize active viral infection

should include a strong focus on older juveniles as well as male bats. A special attention should

be given to *E. helvum* and *R. aegyptiacus* which present significantly higher seroprevalences

compared to the other bat species, without excluding research on other species. In the same

way, surveillance of ebolavirus spillover events related to ebolavirus circulation in bats should

be reinforced where and when circulation and risks might be enhanced: at the interface with bat

colonies when older juveniles are highly prevalent, especially *E. helvum* (during the dry season

for forested Guinea) and *R. aegyptiacus* (already demonstrated for Marburg virus in *R.*

aegyptiacus). Based on these hypotheses, we screened a selection of individuals for filoviruses

RNA, but all samples were negative. Given the observed sero-prevalence rates, we assume that

the failure to detect filovirus RNA might be explained by either very low viral shedding at time

of sampling, the circulation of unknown filoviruses which might not be detected by the PCR

protocol used, or inappropriate sample type. Viral load might indeed be generally higher in

internal organs than in rectal and oral swabs and it might thus be necessary to use the above-

mentioned criteria to target a subset of individuals for more invasive sampling.

462   In conclusion, this study improved the understanding of filovirus dynamics in

frugivorous bats by using a longitudinal approach that helps to test the role of potentially

influential factors. It emphasizes the importance of host reproductive seasonality as a driver of

pathogen circulation and thus spillover risks, which should be considered when designing

sampling protocols for surveillance and research. Data can for instance be used to update

ebolavirus spillover risk maps already developed for forested Guinea (47). The study was

implemented in an ecosystem exposed to intensive human transformation influencing viral

ecology. Substantial sampling efforts need to be pursued and combined with data from

ecological and behavioural studies on bats and inter-species relations to allow for a broader

comprehension of the ecology of filoviruses at the community level. These efforts should also

be replicated in different social-ecological ecosystems with different exposure to human
footprint for a comparative analysis.

475 Acknowledgments

We thank the Comité National d'Ethique pour la Recherche en Santé (CNERS) and the Office
Guinéen des Parcs Nationaux et Réserves de Faune (OGPRF) of the Ministère de
l'Environnement, et du Développement de la République de Guinée for authorising this study.
We thank the Direction Nationale des Services Vétérinaires of the Ministère de l'Agriculture et
de l'Elevage of Guinea, the OGPFR, and the CERFIG for their constant support and facilitation
of the project, as well as all the field staff (Aboubacar Diallo, Pierre Kabinet Kamano,
Mamadou Aliou Bah, Jean Paul Guilavogui, Mamadou Aliou Sow, Aboubacar Camara
Bountouraby, Aboubacar Camara, Fassou Kourouma , KoïKoï Sakouvogui).

485 Data availability

486 The datasets generated and analysed during the current study are available in the Supporting
487 information or in the following repository: https://doi.org/10.18167/DVN1/KMKUQ8

489 Conflicts of Interest

490 The author(s) declare(s) that there is no conflict of interest regarding the publication of this
491 paper

493 Funding Statement

This research was carried out as part of the EBO-SURSY project (https://rr-
africa.oie.int/en/projects/ebo-sursy-en/), funded by the European Union Delegation under the

Agreement [FOOD/2016/379-660], and the EbOHealth project, publicly funded by I-SITE

MUSE (Montpellier Université d'Excellence https://muse.edu.umontpellier.fr/en/muse-isite -

MUSE2018-EbOHEALTH) through the French National Research Agency (ANR) under the

"Investissements d'avenir" program (grant number ANR-16-IDEX-0006).

The funders had no role in study design, data collection and analysis, decision to publish, or

preparation of the manuscript.

Supporting information

Methods S1. Molecular confirmation of bat species

Table S1. Mean fluorescence intensity cutoff values for different viral antigens, by Ebola
virus lineage and statistical method used to determine cutoff according to De Nys et al, 2018.

Table S2. Characteristics of the variables used in the GLMM statistical models.

Table S3. Ranking of the top 5 best statistical models for the 3 glycoproteins by increasing
AIC value carried out on Rstudio software (version 1.4.1106)

Table S4. Longitudinal representation of the number of bats sampled per month for
ebolavirus serology (October 2018 - July 2020)

Table S5. Simultaneous reactivity of bat antibodies against antigens of different ebolavirus
lineages (M+4SD cutoff)

Table S6. p-values calculated with the estimated marginal means methods to compare

differences in antibody positivity according to age, sex, season and species (with the M+4SD

method) (significant p-value <0.05)

Table S7. Samples tested by PCR for filovirus RNA

518
519
References

521 1.   Centers for Disease Control and Prevention. History of Ebola Virus Disease (EVD)

522   Outbreaks [Internet]. 2022. [cited 2022 Feb 1]. Available from:

523 https://www.cdc.gov/vhf/ebola/history/chronology.html#anchor_1526565058132

524 2. Keita AK, Koundouno FR, Faye M, Düx A, Hinzmann J, Diallo H, et al. Resurgence of

525   Ebola virus in 2021 in Guinea suggests a new paradigm for outbreaks. Nature

526   [Internet]. 2021 Sep 23 [cited 2021 Dec 28];597(7877):539–43. Available from:

527   https://www.nature.com/articles/s41586-021-03901-9

528 3. Fairhead J, Leach M, Millimouno D. Spillover or endemic? Reconsidering the origins

529   of Ebola virus disease outbreaks by revisiting local accounts in light of new evidence

530   from Guinea. BMJ Glob Heal [Internet]. 2021 Apr 1 [cited 2021 May 6];6(4):e005783.

531   Available from: http://www.ncbi.nlm.nih.gov/pubmed/33893144

532 4. Mbala-Kingebeni P, Pratt C, Mutafali-Ruffin M, Pauthner MG, Bile F, Nkuba-Ndaye

533   A, et al. Ebola Virus Transmission Initiated by Relapse of Systemic Ebola Virus

534   Disease. N Engl J Med [Internet]. 2021 Apr 1 [cited 2022 Feb 3];384(13):1240–7.

535   Available from: http://www.nejm.org/doi/10.1056/NEJMoa2024670

536 5. Pigott DM, Golding N, Mylne A, Huang Z, Henry AJ, Weiss DJ, et al. Mapping the

537   zoonotic niche of Ebola virus disease in Africa. Elife. 2014;3:e04395.

538 6. Leroy EM, Kumulungui B, Pourrut X, Rouquet P, Hassanin A, Yaba P, et al. Fruit bats

539   as reservoirs of Ebola virus. Nature [Internet]. 2005 Dec 1 [cited 2014 Jul

540   18];438(7068):575–6. Available from: http://dx.doi.org/10.1038/438575a

541 7. De Nys HM, Kingebeni PM, Keita AK, Butel C, Thaurignac G, Villabona-Arenas C-J,

542   et al. Survey of Ebola Viruses in Frugivorous and Insectivorous Bats in Guinea,

543   Cameroon, and the Democratic Republic of the Congo, 2015–2017. Emerg Infect Dis

544   [Internet]. 2018 Dec 17 [cited 2018 Nov 12];24(12). Available from:

545   http://www.ncbi.nlm.nih.gov/pubmed/30307845

546 8. Goldstein T, Anthony SJ, Gbakima A, Bird BH, Bangura J, Tremeau-Bravard A, et al.

547   The discovery of Bombali virus adds further support for bats as hosts of ebolaviruses.

Nat Microbiol [Internet]. 2018 Oct 27 [cited 2019 Mar 14];3(10):1084–9. Available

from: http://www.ncbi.nlm.nih.gov/pubmed/30150734

9.   Kamins AO, Rowcliffe JM, Ntiamoa-Baidu Y, Cunningham AA, Wood JLN, Restif O.

Characteristics and Risk Perceptions of Ghanaians Potentially Exposed to Bat-Borne

Zoonoses through Bushmeat. Ecohealth [Internet]. 2015 Mar 30 [cited 2017 Apr

6];12(1):104–20. Available from: http://link.springer.com/10.1007/s10393-014-0977-0

10.   Leroy E, Gonzalez JP, Pourrut X. Ebolavirus and Other Filoviruses. In Springer,

Berlin, Heidelberg; 2007 [cited 2021 Dec 28]. p. 363–87. Available from:

http://link.springer.com/10.1007/978-3-540-70962-6_15

11.   Leendertz SAJ, Gogarten JF, Düx A, Calvignac-Spencer S, Leendertz FH. Assessing

the Evidence Supporting Fruit Bats as the Primary Reservoirs for Ebola Viruses.

Ecohealth [Internet]. 2016 Mar 13 [cited 2021 Dec 28];13(1):18–25. Available from:

http://link.springer.com/10.1007/s10393-015-1053-0

12.   Leroy EM, Rouquet P, Formenty P, Souquière S, Kilbourne A, Froment J-M, et al.

Multiple Ebola virus transmission events and rapid decline of central African wildlife.

Science [Internet]. 2004 Jan 16 [cited 2016 Mar 23];303(5656):387–90. Available

from: http://science.sciencemag.org/content/303/5656/387.abstract

13.   Formenty P, Boesch C, Wyers M, Steiner C, Donati F, Dind F, et al. Ebola virus

outbreak among wild chimpanzees living in a rain forest of Cote d'Ivoire. J Infect Dis

[Internet]. 1999/02/13. 1999;179 Suppl:S120-6. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/9988175

14.   Pourrut X, Délicat A, Rollin PE, Ksiazek TG, Gonzalez J-P, Leroy EM. Spatial and

temporal patterns of Zaire ebolavirus antibody prevalence in the possible reservoir bat

species. J Infect Dis [Internet]. 2007 Nov 15 [cited 2017 Jun 30];196 Suppl 2(s2):S176-

83. Available from: https://academic.oup.com/jid/article-lookup/doi/10.1086/520541

15.   Paweska JT, Storm N, Grobbelaar AA, Markotter W, Kemp A, Jansen van Vuren P.

Experimental Inoculation of Egyptian Fruit Bats (Rousettus aegyptiacus) with Ebola

Virus. Viruses [Internet]. 2016 Jan 22 [cited 2016 Feb 1];8(2):29. Available from:

http://www.mdpi.com/1999-4915/8/2/29/htm

16.   Swanepoel R, Leman PA, Burt FJ, Zachariades NA, Braack LE, Ksiazek TG, et al.

Experimental inoculation of plants and animals with Ebola virus. Emerg Infect Dis

[Internet]. 1996 [cited 2016 Oct 21];2(4):321–5. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/8969248

17.   Towner JS, Amman BR, Sealy TK, Carroll SAR, Comer JA, Kemp A, et al. Isolation

of Genetically Diverse Marburg Viruses from Egyptian Fruit Bats. Fouchier RAM,

editor. PLoS Pathog [Internet]. 2009 Jul 31 [cited 2016 Aug 30];5(7):e1000536.

Available from: http://dx.plos.org/10.1371/journal.ppat.1000536

18.   Negredo A, Palacios G, Vázquez-Morón S, González F, Dopazo H, Molero F, et al.

Discovery of an Ebolavirus-Like Filovirus in Europe. Basler CF, editor. PLoS Pathog

[Internet]. 2011 Oct 20 [cited 2021 Jun 8];7(10):e1002304. Available from:

https://dx.plos.org/10.1371/journal.ppat.1002304

19.   Yang X-L, Zhang Y-Z, Jiang R-D, Guo H, Zhang W, Li B, et al. Genetically Diverse

Filoviruses in *Rousettus* and *Eonycteris* spp. Bats, China, 2009 and 2015. Emerg Infect

Dis [Internet]. 2017 Mar [cited 2017 Apr 18];23(3):482–6. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/28221123

20.   Amman BR, Jones MEB, Sealy TK, Uebelhoer LS, Schuh AJ, Bird BH, et al. ORAL

SHEDDING OF MARBURG VIRUS IN EXPERIMENTALLY INFECTED

EGYPTIAN FRUIT BATS (*ROUSETTUS AEGYPTIACUS*). J Wildl Dis [Internet].

2015 Jan [cited 2021 Jun 8];51(1):113–24. Available from:

http://www.bioone.org/doi/10.7589/2014-08-198

21.   Richter H V., Cumming GS. Food availability and annual migration of the straw-

599   colored fruit bat (Eidolon helvum). J Zool [Internet]. 2005 Dec 19 [cited 2018 Apr

600   13];268(1):35–44. Available from: http://doi.wiley.com/10.1111/j.1469-

601   7998.2005.00020.x

602   22.   Cumming GS, Bernard RTF. Rainfall, food abundance and timing of parturition in

603   African bats. Oecologia [Internet]. 1997 Jul 18 [cited 2021 Dec 28];111(3):309–17.

604   Available from: http://link.springer.com/10.1007/s004420050240

605   23.   Amman BR, Carroll SA, Reed ZD, Sealy TK, Balinandi S, Swanepoel R, et al.

606   Seasonal Pulses of Marburg Virus Circulation in Juvenile Rousettus aegyptiacus Bats

607   Coincide with Periods of Increased Risk of Human Infection. Kawaoka Y, editor. PLoS

608   Pathog [Internet]. 2012 Oct 4 [cited 2016 Nov 3];8(10):e1002877. Available from:

609   http://dx.plos.org/10.1371/journal.ppat.1002877

610   24.   Djomsi D, Mba Djonzo F, Ndong Bass I, Champagne M, Lacroix A, Thaurignac G, et

611   al. Dynamics of Antibodies to Ebolaviruses in an Eidolon helvum Bat Colony in

612   Cameroon. Viruses [Internet]. 2022 Mar 9 [cited 2022 Apr 9];14(3):560. Available

613   from: https://www.mdpi.com/1999-4915/14/3/560

614   25.   Maganga GD, Bourgarel M, Vallo P, Dallo TD, Ngoagouni C, Drexler JF, et al. Bat

615   Distribution Size or Shape as Determinant of Viral Richness in African Bats. Baker

616   ML, editor. PLoS One [Internet]. 2014 Jun 24 [cited 2021 Dec 28];9(6):e100172.

617   Available from: https://dx.plos.org/10.1371/journal.pone.0100172

618   26.   Lacroix A, Mbala Kingebeni P, Ndimbo Kumugo SP, Lempu G, Butel C, Serrano L, et

619   al. Investigating the Circulation of Ebola Viruses in Bats during the Ebola Virus

620   Disease Outbreaks in the Equateur and North Kivu Provinces of the Democratic

621   Republic of Congo from 2018. Pathogens [Internet]. 2021 May 4 [cited 2021 May

622   6];10(5):557. Available from: https://www.mdpi.com/2076-0817/10/5/557

623   27.   Ayouba A, Touré A, Butel C, Keita AK, Binetruy F, Sow MS, et al. Development of a

624   Sensitive and Specific Serological Assay Based on Luminex Technology for Detection

625   of Antibodies to Zaire Ebola Virus. Tang Y-W, editor. J Clin Microbiol [Internet].

626   2017 Jan [cited 2017 Apr 6];55(1):165–76. Available from:

627   http://jcm.asm.org/lookup/doi/10.1128/JCM.01979-16

628   28.   Calcagno V, Mazancourt C de. glmulti : An R Package for Easy Automated Model

629   Selection with (Generalized) Linear Models. J Stat Softw [Internet]. 2010 May 31

630   [cited 2021 Dec 28];34(12):1–29. Available from: http://www.jstatsoft.org/v34/i12/

631   29.   Goldstein T, Anthony SJ, Gbakima A, Bird BH, Bangura J, Tremeau-Bravard A, et al.

632   The discovery of Bombali virus adds further support for bats as hosts of ebolaviruses.

633   Nat Microbiol [Internet]. 2018 Oct 27 [cited 2018 Nov 12];3(10):1084–9. Available

634   from: http://www.ncbi.nlm.nih.gov/pubmed/30150734

635   30.   Diallo MSK, Ayouba A, Keita AK, Thaurignac G, Sow MS, Kpamou C, et al.

636   Temporal evolution of the humoral antibody response after Ebola virus disease in

637   Guinea: a 60-month observational prospective cohort study. The Lancet Microbe

638   [Internet]. 2021 Dec 1 [cited 2022 Aug 29];2(12):e676–84. Available from:

639   https://linkinghub.elsevier.com/retrieve/pii/S2666524721001701

640   31.   Schuh AJ, Amman BR, Sealy TS, Flietstra TD, Guito JC, Nichol ST, et al.

641   Comparative analysis of serologic cross-reactivity using convalescent sera from

642   filovirus-experimentally infected fruit bats. Sci Rep [Internet]. 2019 Dec 30 [cited 2019

643   May 3];9(1):6707. Available from: http://www.nature.com/articles/s41598-019-43156-

644   z

645   32.   Nunn CL, Jordán F, McCabe CM, Verdolin JL, Fewell JH. Infectious disease and

646   group size: more than just a numbers game. Philos Trans R Soc B Biol Sci [Internet].

647   2015 May 26 [cited 2021 Jun 8];370(1669):20140111. Available from:

https://royalsocietypublishing.org/doi/10.1098/rstb.2014.0111

33. Rosevear DR. The bats of West Africa. London: British Museum (Natural History);

1965.

34. Bradbury JW. Lek Mating Behavior in the Hammer-headed Bat. Z Tierpsychol

[Internet]. 2010 Apr 26 [cited 2021 Dec 28];45(3):225–55. Available from:

https://onlinelibrary.wiley.com/doi/10.1111/j.1439-0310.1977.tb02120.x

35. Peel AJ, Wood JLN, Baker KS, Breed AC, Carvalho A De, Fernández-Loras A, et al.

How Does Africa's Most Hunted Bat Vary Across the Continent? Population Traits of

the Straw-Coloured Fruit Bat (*Eidolon helvu* m) and Its Interactions with Humans.

Acta Chiropterologica. 2017;19(1).

36. Amman BR, Bird BH, Bakarr IA, Bangura J, Schuh AJ, Johnny J, et al. Isolation of

Angola-like Marburg virus from Egyptian rousette bats from West Africa. Nat

Commun [Internet]. 2020 Dec 24 [cited 2021 Dec 28];11(1):510. Available from:

http://www.nature.com/articles/s41467-020-14327-8

37. Jones M, Schuh A, Amman B, Sealy T, Zaki S, Nichol S, et al. Experimental

Inoculation of Egyptian Rousette Bats (Rousettus aegyptiacus) with Viruses of the

Ebolavirus and Marburgvirus Genera. Viruses [Internet]. 2015 Jun 25 [cited 2021 Jun

8];7(7):3420–42. Available from: http://www.mdpi.com/1999-4915/7/7/2779

38. Pourrut X, Souris M, Towner JS, Rollin PE, Nichol ST, Gonzalez J-P, et al. Large

serological survey showing cocirculation of Ebola and Marburg viruses in Gabonese

bat populations, and a high seroprevalence of both viruses in Rousettus aegyptiacus.

BMC Infect Dis [Internet]. 2009 [cited 2016 Jun 21];9:159. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/19785757

39. Thorson AE, Deen GF, Bernstein KT, Liu WJ, Yamba F, Habib N, et al. Persistence of

Ebola virus in semen among Ebola virus disease survivors in Sierra Leone: A cohort

study of frequency, duration, and risk factors. Kindrachuk J, editor. PLOS Med

[Internet]. 2021 Feb 10 [cited 2021 Dec 28];18(2):e1003273. Available from:

https://dx.plos.org/10.1371/journal.pmed.1003273

40.   Adaken C, Scott JT, Sharma R, Gopal R, Dicks S, Niazi S, et al. Ebola virus antibody

decay–stimulation in a high proportion of survivors. Nature [Internet]. 2021 Feb 18

[cited 2021 Dec 28];590(7846):468–72. Available from:

http://www.nature.com/articles/s41586-020-03146-y

41.   Hoarau AOG, Goodman SM, Al Halabi D, Ramasindrazana B, Lagadec E, Le Minter

G, et al. Investigation of astrovirus, coronavirus and paramyxovirus co-infections in

bats in the western Indian Ocean. Virol J [Internet]. 2021 Dec 12 [cited 2021 Oct

14];18(1):205. Available from:

https://virologyj.biomedcentral.com/articles/10.1186/s12985-021-01673-2

42.   Storm N, Jansen Van Vuren P, Markotter W, Paweska JT. Antibody Responses to

Marburg Virus in Egyptian Rousette Bats and Their Role in Protection against

Infection. Viruses [Internet]. 2018 [cited 2021 Dec 28];10(2). Available from:

http://www.ncbi.nlm.nih.gov/pubmed/29439384

43.   Buceta J, Johnson K, Jin Z, Wang Y, Wang Z, Drexler J. Modeling the Ebola zoonotic

dynamics: Interplay between enviroclimatic factors and bat ecology. Sun G-Q, editor.

PLoS One [Internet]. 2017 Jun 12 [cited 2017 Jun 13];12(6):e0179559. Available from:

http://dx.plos.org/10.1371/journal.pone.0179559

44.   Happold DCD, Happold M. Reproductive strategies of bats in Africa. J Zool [Internet].

1990 Dec 24 [cited 2021 Dec 28];222(4):557–83. Available from:

https://onlinelibrary.wiley.com/doi/10.1111/j.1469-7998.1990.tb06014.x

45.   Baker KS, Suu-Ire R, Barr J, Hayman DTS, Broder CC, Horton DL, et al. Viral

antibody dynamics in a chiropteran host. Boots M, editor. J Anim Ecol [Internet]. 2014

Mar 13 [cited 2021 Oct 15];83(2):415–28. Available from:

https://onlinelibrary.wiley.com/doi/10.1111/1365-2656.12153

46.   Changula K, Kajihara M, Mori-Kajihara A, Eto Y, Miyamoto H, Yoshida R, et al.

Seroprevalence of Filovirus Infection of Rousettus aegyptiacus Bats in Zambia. J Infect

Dis [Internet]. 2018 Jun 8 [cited 2021 Oct 15];218(suppl_5):S312–7. Available from:

https://academic.oup.com/jid/advance-article/doi/10.1093/infdis/jiy266/5034832

47.   Lee-Cruz L, Lenormand M, Cappelle J, Caron A, De Nys H, Peeters M, et al. Mapping

of Ebola virus spillover: Suitability and seasonal variability at the landscape scale. Bird

B, editor. PLoS Negl Trop Dis [Internet]. 2021 Aug 23 [cited 2021 Dec

28];15(8):e0009683. Available from: https://dx.plos.org/10.1371/journal.pntd.0009683

711
712