Title: Adaptation to oral infection in *D. melanogaster* through evolution of both resistance and disease tolerance mechanisms

Tânia F. Paulo[1*], Priscilla A. Akyaw[1,3,4], Tiago Paixão[1], Élio Sucena[1,2,3,4*]

1 – Instituto Gulbenkian de Ciência, Rua da quinta grande 6, 2780-156 Oeiras, Portugal

2 – Departamento de Biologia Animal, Faculdade de Ciências, Universidade de Lisboa, Edifício C2, Campo Grande, 1749-016 Lisbon, Portugal

3- cE3c: Centre for Ecology, Evolution and Environmental Changes, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1749-016 Lisbon, Portugal

4- CHANGE – Global Change and Sustainability Institute

* TFP: tfpaulo@igc.gulbenkian.pt; ES: jesucena@fc.ul.pt

Keywords: Experimental Evolution, Innate Immunity, *Drosophila*, *Pseudomonas entomophila*

Abstract

Pathogens exert strong selection on hosts, that evolve and deploy different defensive strategies, namely minimizing pathogen exposure (avoidance), directly promoting pathogen elimination (resistance), and/or managing the deleterious effects of illness (disease tolerance). However, how the response to pathogens partitions across these processes has never been directly assessed in a single system, let alone in the context of known adaptive trajectories under controlled selection regimes. Here, an experimental evolution system composed of *D. melanogaster* and its natural pathogen *P. entomophila* is used to determine the role of behavioural traits, and of resistance and disease tolerance mechanisms on host adaption. We compare a population adapted to oral infection with *P. entomophila* (BactOral) with its control population to find no evidence for behavioural change but measurable differences in both resistance and disease tolerance. In BactOral, we identify a relative decrease in bacterial loads correlated with an increase in gut production of specific AMPs, but no differences in bacterial intake, in differential gut cell renewal rate, or in the rate of defecation, pointing to a strengthening in resistance. Additionally, we posit that disease tolerance also contributes to the adaptive response of BactOral through a tighter control of its immune response and of the deleterious effects of exposure. This study reveals a genetically complex and mechanistically multi-layered response, possibly reflecting the structure of adaptation to infection in natural populations.

Introduction

Host-parasite interactions are main drivers of evolution[1,2]. In natural environments, parasites are constantly attempting to penetrate host defences, which consist on multiple mechanisms to fight infection[3]. Such mechanisms include behavioural strategies (e.g. avoidance)[4], the active reduction or elimination of pathogens (i.e. resistance) and/or the maintenance of homeostasis and fitness without interfering with pathogen burden (i.e. disease tolerance)[5–8]. While many studies in the past decades have explored the mechanistic and genetic bases of these different host defence strategies in diverse organisms [9–11], few have focused on how these processes can be shaped by evolution[12–15]. Experimental evolution is a powerful method by which it is possible to infer the evolutionary response of a host species to a specific pathogenic pressure under controlled conditions[16]. Previous studies of experimental evolution have addressed the impact of different hosts on parasite fitness[17], of different pathogens on the host's evolutionary trajectory[13,18,19], of immune priming[20,21], and of different routes of infection on host adaptation[19,22].

It is believed that oral exposure to pathogens constitutes a common process of infection in *D. melanogaster*[23,24], as it lives on decaying substrates. When feeding on pathogenic microorganisms, *Drosophila* deploys different responses that include avoidance behaviours[25,26], the barrier action of the peritrophic matrix[27], localized high acidity of the midgut[28], intensification of gut transit[29], local production of Anti-Microbial Peptides (AMPs)[30] and Reactive Oxygen Species (ROS) by gut epithelial cells[31,32], increase of enterocyte delamination and renewal rates[33–36]. In addition, it can activate a systemic humoral response, when the pathogen manages to cross the gut epithelium into the haemolymph[37,38] or otherwise provokes a systemic reaction[23,39–41]. In the specific case of *D. melanogaster* oral infection with the natural entomopathogen *Pseudomonas entomophila*, its high pathogenicity is linked to the production of toxins and virulence factors[42,43] causing translational arrest in host tissues and the inability to renew damaged gut epithelium[44,45]. These responses provide a clear picture of the multi-layered and mechanistically complex nature of the immune response, understood in its broad sense, as presented above[3].

In this work, we used a previously experimentally evolved outbred population of *D. melanogaster* (hereafter called "BactOral") showing a fast evolutionary response to a strong selection with oral *P. entomophila* infection[19,46], to investigate the mechanisms targeted by adaptation. We systematically tested for behaviour, resistance, and disease tolerance contributions, to provide a comprehensive account of the partitioning of these defense strategies in adaptation.

Results

Evolved population shows higher survival upon infection

In previous work[19,46], experimental evolution was conducted on an outbred population of *D. melanogaster* against oral infection either with the bacterium *P. entomophila* (hereafter, BactOral) or with a control solution (hereafter, Control). Since then, these populations[46] have been kept under relaxed selection for over 80 generations. To understand to which extent this relaxation affected the response to the original selection regime, we measured survival of Control and BactOral populations upon *P. entomophila* oral infection. We found significantly higher survival of BactOral for both sexes (Fig. 1, Infected Control females *versus* Infected BactOral females: *z.ratio*= 9.123, *p*< 0.0001; Infected Control males *versus* Infected BactOral males: *z.ratio*= 9.909, *p*< 0.0001)(Suppl. Table 1). Moreover, these differences are not due to an inherent frailty of the Control as compared to BactOral, since survival under uninfected conditions (for the equivalent time period) was similar in both populations (Fig. 1, Uninfected Control females *versus* Uninfected BactOral females: *z.ratio*= 0.004, *p*= 1; Uninfected Control males *versus* Uninfected BactOral males: *z.ratio*= -1.080, *p*= 0.9611)(Suppl. Table 1).

Taking these results into account, we undertook a systematic description of multiple traits previously associated to immune response to oral infections aiming to identify different processes targeted by selection and estimate their relative contribution.

Evolved population has not changed bacterial uptake or defecation rates

Our systematic approach started by screening behavioural traits that impact gut bacterial number: feeding and defecation. To test for differences in feeding behaviour, we performed a Capillary Feeding assay (CAFE)[47] on females of both Control and BactOral populations at different timepoints over 24 hours of exposure to the bacteria solution. We found no significant differences between the amount ingested by individuals of the two populations. Pairwise comparisons of their cumulative bacterial intake after 2 hours, 8 hours and 24 hours exhibited a comparable amount of total bacteria ingested (Suppl. Table 2 and Fig. 2A, Control *versus* BactOral at 2 hours, *t.ratio*= 0.097, *p*= 1; at 8 hours, *t.ratio*= 0.643, *p*= 0.9873; at 24 hours, *t.ratio*= 0.574, *p*= 0.9925).

Additionally, we estimated the average slopes from linear regression where microliters of bacteria eaten were modelled between different time points, concluding that the feeding rates of both populations decreased in similar ways as infection progresses. Thus, exposure affects feeding ability of both populations to a comparable degree and, consequently, adaptation to oral

infection with *P. entomophila* in the BactOral regime did not involve an alteration of feeding
behaviour.

Considering the absence of significant differences in bacterial intake, an alternative would
be that the BactOral population had evolved a more effective defecation rate, which would reduce
the time of bacterial permanence and proliferation in the gut. We established a protocol to measure
live bacteria loads in the faeces of individual flies upon oral infection through the quantification
of CFUs defecated over a six-hour period and the subsequent twelve hours post-exposure (Fig.
2B).

We observed a few instances where flies did not expel alive bacteria during that period
or did so in a quantity that fell below our detection threshold. Therefore, a zero-inflated model
with a negative binomial distribution was fitted on the counts data (accounting for overdispersion)
jointly with a binomial distribution on the zeros. For the counts portion of the model, none of the
three factors significantly predicted the measured bacterial loads (Suppl. Table 3, model
parameters - Population: *z.value*= 1.324, *p*= 0.185; Sex: *z.value*= 0.073, *p*= 0.942; Timepoint:
z.value= -0.359, *p*= 0.720). However, looking at the zero portion of the model, we find a
significant effect of Population (Population: *z.value*= -2.407, *p*= 0.016; Timepoint: *z.value*= -
0.006, *p*= 0.996).

We then performed post-hoc pairwise comparisons on these regressions and identified a
significant decrease in bacterial loads defecated by males of BactOral, between the two timepoints
(males: *z.ratio*= -2.719, *p*= 0.033), but no other significant effects among bacterial counts between
populations and/or timepoints, in a sex-dependent manner. However, when considering the zeros,
we found a significant difference in the proportion of BactOral flies with undetectable amounts
of bacteria in the faeces between the first and second timepoints (B6h– B18h: *z.ratio*= 4.092, *p*=
0.0002), as well as compared to Control at both timepoints (B6h– C6h: *z.ratio*= 3.612, *p*= 0.002;
B6h– C18h: *z.ratio*= 3.530, *p*= 0.002)(Suppl. Table 3).

In sum, we were not able to correlate the higher survival of the BactOral population upon
oral infection with a more efficient defecation. However, we observed that males of BactOral
decrease the amount of live bacteria they defecate as time progresses, hinting at a possible
resistance mechanism more evident in males of the adapted lines.

Evolved population shows a faster decrease and clearance of bacteria

Next, we studied the physiological responses to estimate the roles of resistance and
disease tolerance. First, we tested for structural changes in the gut itself that could influence the
"carrying capacity" of the structure and create difficulties in interpretating the quantification

experiments. We measured the capacity of BactOral to regenerate its gut epithelium upon

bacterial infection relative to Control, using initial length and extent of the shortening induced by

infection[36], as a proxy. For this, we measured the total length of the guts of females, before and

after 24 hours of oral infection with *P. entomophila* and performed multiple pairwise comparisons

on a linear model with mixed effects to pinpoint how population and infection status affected this

trait. We found a shortening of approximately 2 mm in the total length of infected guts by

comparing the post-infection timepoint of 24 hours to the initial timepoint, prior to exposure

(Suppl. Table 4 and Suppl. Fig. 1, Control before infection *versus* Control 24 hours post-infection:

t.ratio= 6.274, *p*<0.0001; BactOral before infection *versus* BactOral 24 hours post-infection:

t.ratio= 5.453, *p* <0.0001). Nonetheless, no significant differences between populations in the

total length of the guts could be detected both before and after infection (Without infection:

BactOral *versus* Control – *p*= 0.6719; after 24 hours of infection – *p*= 0.9925) (Suppl. Fig. 1).

To explore the dynamics of bacterial numbers inside the host, we performed a time-course

characterization of bacterial loads during and after oral infection by counting the number of

colony-forming units (CFUs) obtained from individual flies. In Figure 3, we show the progression

of bacterial loads from 1 hour until 52 hours post-infection (Fig. 3A) as well as the cumulative

number of individuals without detectable live bacteria (Fig. 3B). The infection dynamics of both

populations start with very high numbers of *P. entomophila* at early stages, confirming the

absence of differences in initial inoculum reported above (Table 1, parameter *a*: 3.072 $\pm$ 0.016).

Bacterial loads continuously decrease and, at approximately 20 hours of infection, Control and

BactOral population begin to differ significantly in a sex-dependent manner. From this point

onwards we observe a steady decrease of bacterial quantities in the two populations, always more

severe in males than in females, and in BactOral than in Control (Fig. 3A). In addition, the number

of flies that clear the infection also increases with time in both populations, but at a faster rate and

reaching higher levels in BactOral flies (Fig. 3B). By 52 hours post-infection, in both males and

females, the number of "cleared" flies in the BactOral populations was twice as high as that of its

Control counterpart (Fig. 3B). Together, these analyses indicate that the evolved population

eliminates bacteria faster and more efficiently than Control.

In order to pinpoint more accurately the parameters contributing to this overall difference,

we modelled the bacterial load and zero dynamics, and assessed differences in parameter

estimates considering the initial load or inoculum (*a*), the load in the long-term or set-point

bacterial load (*b*), the timepoint at which the load is reduced in half (*k*) and the highest rate of

decrease or slope at timepoint *k* (*h*) (see Materials and Methods for details). In Table 1, we show

the mean estimates, standard deviation, and high-density intervals (3% and 97%) for all four

parameters of the fitted sigmoidal model, for Control and BactOral populations. Overall, these

results show that sex has a significant effect on all curve parameters (*k*= -0.053 $\pm$ 0.068; *h*= -0.330

± 0.227; a = -0.130 ± 0.038; b = -0.147 ± 0.066), reinforcing the notion of a strong sexual

dimorphism in response to infections[48–50]. Additionally, populations differ on "time to half height"

(k= -0.108 ± 0.046) and final load (b= -0.285 ± 0.053), but not on initial load (a= -0.002 ± 0.024)

nor on highest rate of decrease (h= 0.042 ± 0.182).

Because this time-course experiment yielded a significant amount of zero counts,

corresponding to individual flies for which no CFUs could be counted, we treated this as an

independent dataset for which we developed a separate model focused on infection clearance

(Table 2). With that, we found that a_1 (final fraction of zeros) was significantly affected by both

sex and population (with no observable overlap between the mean estimates for each condition)

(a_1 Females= -0.781 ± 0.07; a_1 Males= -0.061 ± 0.069; a_1 Control= -0.700 ± 0.070; a_1 BactOral=

-0.144 ± 0.069), with more BactOral individuals having cleared the infection by the end of the

experiment and males more so than females. There were significant overlaps between the

estimates of the remaining parameters, evidencing a difference in clearance rate only in late stages

of bacterial infection. In sum, these data align with those of bacterial load dynamics, whereby

flies from BactOral population clear *P. entomophila* faster and better from 20h of exposure

onwards.

These results align with the hypothesis that resistance mechanisms participate in the

adapted response of BactOral. Indeed, bacterial numbers are affected differently between

BactOral and Control populations despite the equivalent initial bacterial inocula (Fig. 2A, Table

1) and defecation rates (Fig. 2B). The differences in number of detected live bacteria inside flies

suggests a direct bacterial control mechanism is operating more efficiently in BactOral

individuals.

Differences in AMP expression are consistent with distinct bacterial loads

dynamics

In an attempt to pinpoint which resistance mechanisms contribute to the differences in

bacterial clearance rates observed between populations, we infected female flies from Control and

BactOral and dissected their guts prior to exposure, and after 8, 24, and 32 hours of infection with

P. entomophila. We quantified expression levels for a panel of 10 AMPs covering different

molecular families, dedicated pathways and pathogen specificities, including *Attacin-A*,

Bomanins Short 2 and *3*, *Cecropin-A1*, *Defensin*, *Diptericin-A*, *Drosomycin* and *Drosomycin*-like

2, *3* and *5* (Fig. 4, Suppl. Table 5 and Supp. Fig. 2).

We first verified if the baseline levels of AMPs (i.e., expressed under homeostatic

conditions) differed between populations, by relativizing expression levels to the Control

population at the unchallenged timepoint. With this approach, we established that BactOral and Control express comparable amounts of 8 AMPs when unchallenged, the two exceptions being *Cecropin-A1* (*CecA*) with lower expression levels (*t.ratio*= 2.892, *p*= 0.0445) and *Drosomycin-like 5* (*Drsl5*) with higher levels (*t.ratio*= -4.686 , *p*= 0.0094) (Supp. Fig. 2).

In addition, and since our main goal was to characterize the temporal dynamics of AMP expression in both populations, we performed the analysis for the entirety of the time-course using the unchallenged timepoint as reference (Fig. 4 and Suppl. Table 5). Overall, we detect consistent dynamics with the reduction of expression happening sooner (by 32 hours) in BactOral than Control across 7 of the 10 AMPs tested, the exceptions being Attacin A, Defensin and Drosomycin-like 2 (Fig. 4). This earlier down-regulation in BactOral relative to Control (at 32-hours) of *BomaninS2* (*BomS2*) (*t.ratio*= 4.347, *p*= 0.0047), *BomaninS3* (*BomS3*) (*t.ratio*= 4.035, *p*= 0.0089), *CecropinA* (*t.ratio*= 4.529, *p* = 0.0010), *Diptericin* (*t.ratio*= 2.613, *p*= 0.0439), *Drosomycin* (*t.ratio*= 4.795, *p*= 0.0056), *Drosomycin-like 3* (*Drsl3*) (*t.ratio*= 3.194, *p*= 0.0105) and *Drosomycin-like 5* (*Drsl5*) (*t.ratio*= 4.266, *p*= 0.0021). Furthermore, we detected a significantly higher up-regulation of *AttacinA* and *CecA* in BactOral at 8 hours of infection (*AttacinA*: Control 8h *versus* BactOral 8h – *t.ratio*= -2.563, *p*= 0.0389; *CecA*: Control 8h *versus* BactOral 8h – *t.ratio*= - 3.034, *p*= 0.0123), consistent with the role of these peptides as main players against infections with Gram-negative bacteria[51,52]. A similar trend was observed for other tested AMPs such as *Defensin* and *Drsl2*, although these changes were not statistically significant (*Defensin*: Control 8h *versus* BactOral 8h – *t.ratio*= -2.101, *p*= 0.231; *Drsl2*: Control 8h *versus* BactOral 8 h.p.i. – *t.ratio*= -0.899, *p*= 0.465).

Principle Component Analysis (PCA) supports these trends (Suppl. Fig. 3). The two main Principle Components (PC1 and PC2) evidence the time-dependent differences observed between populations by showing an initial indistinguishable grouping of both, followed by a dissimilar change in expression patterns at 8 hours of infection which culminates with a maximum divergence between Control and BactOral at the later timepoint (32 hours).

Increase of AMP levels in the Control population phenocopies BactOral response to infection

Having correlated the improved bacterial clearance of BactOral (Fig. 3) with significant differences in expression levels of AMPs between populations (Fig. 4), we sought to functionally associate these two observations. For that, we conducted a priming experiment to induce higher basal levels of AMPs by exposing flies to a solution of heat-killed bacteria, which has been shown

to "artificially" increase AMP levels[53–55]. These primed flies were, subsequently, infected orally

with live *P. entomophila*, and followed daily for mortality.

Previous exposure to heat-killed bacteria increased survival to infection with live *P.*

entomophila in BactOral and Control populations, confirming the stronger immune response

expected upon priming (Suppl. Table 6 and Fig. 5, BactOral females fed with live bacteria

(control) *versus* BactOral females primed with heat-killed bacteria (primed): *z.ratio*= 7.193, *p*<

0.0001; BactOral males treated with control *versus* BactOral males primed with Heat-killed

bacteria: *z.ratio*= 4.368, *p*= 0.0001; Control females fed with control *versus* Control females :

z.ratio= 10.011 , *p*< 0.0001; Control males treated with control *versus* Control males primed :

z.ratio= 7.299 , *p*< 0.0001). Females from the Control population primed with heat-killed bacteria

increased their survival to oral infection, virtually phenocopying BactOral infected with live

bacteria, whereas in males, the same tendency is observed but to a lesser degree (Suppl. Table 6

and Fig. 5, BactOral females fed with control *versus* Control females primed: *z.ratio*= 1.629, *p*=

0.3620 and BactOral males fed with control *versus* Control males primed: *z.ratio*= - 4.547,

p<0001). This effect is supported by a time-course of relative expression levels of IMD pathway

key players, *Relish* (its devoted transcription factor) and its associated AMP, *Diptericin*, as well

as the Toll-activated AMP, *Drosomycin* (Suppl. Fig. 4). For both populations, we observed an

increase in levels of *Diptericin* to a comparable degree upon exposure to heat-killed *P.*

entomophila, but not of *Drosomycin*.

With this experiment we strengthened the case for a role of improved resistance in the

adapted phenotype of BactOral.

BactOral maintains higher survival upon immune response activation

To ascertain whether disease tolerance may also contribute to BactOral increased survival

upon oral infection, we continuously fed flies with a mixture of heat-killed *P. entomophila* and

followed survival.

As the experimental evolution protocol with which these populations were generated

consisted of selection until flies were approximately 10-day old, we analyzed survival after

exposure to heat-killed *P. entomophila* exclusively until day 14. With this we verified that, up to

this timepoint, both females and males of BactOral survive significantly more than Control after

this specific damage damage (Fig 6, Control - HK *versus* BactOral - HK females: *z.ratio*= 6.579,

p< 0.0001; males: *z.ratio*= 9.612, *p*< 0.0001). No significant differences exist under control

conditions (Control - PBS *versus* BactOral - PBS females: *z.ratio*= 1.392, *p*= 0.504; males:

z.ratio= 0.587, *p*= 0.936).

This result strongly suggests a role for disease tolerance, as exposure to heat-killed
bacteria is expected to trigger host responses. Both the deleterious effects of virulence factors (at
least partially) as well as of immunopathology should be at play and appear to participate
differentially between control and adapted populations in their survival. Together with the gene
expression data presented above (Fig. 4) showing that BactOral represses the activation of
immune effectors earlier upon infection, it is tempting to hypothesize that a similar process
happened during the course of this experiment, and a role may be attributed to tighter control and
negative regulation of immune over-activation.

Discussion

In this work, we enquired systematically for the bases of adaptation against an oral
bacterial infection in *D. melanogaster*, considering the three general layers that constitute the
immune response: behaviour, resistance and disease tolerance[3,8,56]. Upon experimental evolution
of an outbred population of *D. melanogaster* against oral infection with *P. entomophila*, and after
approximately 6 generations, the selected population (BactOral) showed increased survival upon
infection[19]. This phenotype was maintained throughout the selection experiment but, since then,
the population has been kept for more than 80 generations under relaxed selection. Despite the
fact that no physiological costs associated to this response could be detected[46], we tested the
immunocompetence of the evolved populations. The average survival rate after oral infection of
flies from BactOral was approximately 63% (50% for females and 75% for males) and
approximately 10% in the Control population (5% for females and 15% for males). Considering
that at the end of selection BactOral survival reached 90 %[19], this partial decrease in overall
survival after infection with bacteria can potentially underestimate the contributions of each of
the different mechanisms behind adaptation. Notwithstanding, we were able to ascertain the
relative importance of behavioural, resistance, and disease tolerance mechanisms for BactOral
adaptation to oral infection with *P. entomophila*.

No behavioural or structural differences between BactOral and Control populations

It is well described that the route of infection is of paramount importance to the outcome of host-
parasite interactions In *D. melanogaster*, the mode of entry of a bacterial pathogen into the host's
body elicits different physiological responses[19,57–61], which can start by the avoidance of the

pathogenic substrate, in the case of infection through oral route[26,62,63], both by inhibiting feeding

or choosing oviposition sites according to bacterial composition[26,53,62–64]. P. entomophila is a

natural pathogen of Drosophila that causes high mortality when ingested across diverse genetic

backgrounds and conditions[23,43,44,65,66]. Thus, it is plausible to consider that fasting could constitute

a general response against oral infection. The simplest behaviour that could explain the observed

differences in survival between populations would be that BactOral evolved a reduced intake

during exposure to bacteria, thereby reducing the "inoculum" size. Indeed, fastening is a prevalent

response to infection and behavioural avoidance is a common strategy to limit exposure to

infected substrates in different systems[4,62,63]. However, we showed similar amounts of bacteria

were ingested by females of both populations. With time, feeding rate decreases, which is

consistent with reports that *P. entomophila* ingestion can cause feeding inhibition (Figure 2A)[23,67].

Moreover, we consubstantiate the findings from the CAFE assay with the bacterial loads time-

course, that show equivalent amounts of *P. entomophila* colonies inside infected flies at early

timepoints. Our results indicate that behavioural inhibition of bacterial intake was not one of the

mechanisms that evolved in BactOral flies, eliminating fastening as a component of the adaptation

to *P. entomophila* exposure.

Previous reports have shown that pathogen intake can affect gut function and structure[34–

36,45] as well as influence the dynamics of exposure by altering defecation rates[29]. However, we

could not find evidence for any differences in these features between control and evolved

populations. Firstly, there are no changes in gut length in steady-state conditions that would

enable somehow BactOral to a priori face bacterial infection more favourably nor is there a

difference in their response to damage and capacity to restore lost cells more efficiently upon

infection. Increased enteric cell renewal, which often leads to a temporary shortening of gut

length[36], has been described as one of the main mechanisms of *D. melanogaster* defense against

oral bacterial infection[35,36,45]. In our experiments, we detected a significant decrease in total gut

length after 24 hours of oral infection but equivalent in both populations, not supporting a role for

renewal rate in BactOral adaptation.

These experiments, however, do not completely discard the hypothesis of differential gut

structure underlying the adapted phenotype of BactOral. For example, proportions of gut cell

types can change upon infection[68] and such qualitative shift could contribute to higher fitness

upon exposure to *P. entomophila* in the adapted population. Also, it could be that relevant changes

in gut length occur at later timepoints. Notwithstanding, it has been previously shown that

infection-induced gut cell loss starts as early as 2 hours after exposure, with a return to 75% of

the total original gut length after 24 hours[36]. This time window is well within the boundaries

defined in our setup. Another important aspect to note is that renewal capacity is sexually

dimorphic under diverse physiological contexts[69,70]. For instance, females have higher gut cell

renewal rates than males upon oral infection with *P. carotovorum* 15 [71], a bacterium that provokes
higher mortality in males than females, contrary to oral infection with *P. entomophila* (see
Results). The disparity between those and our observations may be explained by the different the
different mechanisms of virulence employed by these two pathogens to infect *D. melanogaster*[45].
Be as it may, we envision further experiments testing for potential differences in gut delamination
rates between evolved populations at distinct timepoints.

BactOral flies also did not expel pathogenic bacteria from their guts differently from
Control. However, BactOral shows a decrease in the number of alive bacteria defecated between
6- and 18-hours post-exposure, evidencing, to some extent, improved bacterial clearance and
suggesting a role for resistance. Since our protocol only detects live bacteria, we could potentially
be missing an important component of the response and future work should quantify defecation
of alive and dead bacteria, for example through fluorescent microscopy, to compare populations
with regards to the contribution of overall peristalsis to the elimination of total gut content[29]. In
addition, considering the differences in the proportion of individuals that defecated alive bacteria
below our detection limit (zeros), we hypothesize that BactOral either retain *P. entomophila* for
longer inside their guts before they begin to expel it (which could point to higher disease
tolerance), or reduce bacterial loads more effectively at earlier stages, even though with time
bacterial proliferation recovers (again suggesting stronger resistance).

A role for resistance

To further test the suggested role for resistance, we characterized the bacterial load
dynamics in evolved populations over 52 hours and determined that BactOral flies eliminate
pathogens in a more efficient way than Control. This significant change in bacterial loads
indicates that BactOral has evolved higher resistance mechanisms, that control earlier and clear
better gut bacterial loads. As this model was applied to the logarithm of mean bacterial counts at
each timepoint, it is possible that information regarding intra-populational variation is not
considered in the analysis. Nevertheless, the conclusions from the comparison of the bacterial
counts are further supported by the separate analysis on the number of "zeros" as a function of
time, which is a proxy for bacterial clearance.

Even though there is no discernible difference in the amount of CFUs infecting both
females and males until approximately 20h of infection, by this time individuals from BactOral
begin to exhibit gradually decreasing levels of bacteria. This trend aligns with the higher clearance
rate observed in the adapted population (and even higher in males than females) as measured by
the increase in the number of flies from which we cannot grow colonies of *P. entomophila*
("zeros"). Flies from the Control population control their bacterial loads (at around 20h of
infection) but do it at a slower pace and in a less effective way (i.e. many individuals exhibited

high bacterial counts by the end of the experiment). It is possible that our proxy for clearance (i.e.,
flies classified as "zeros"), could be overestimated due to a low detection threshold. Indeed, it has
been shown that *D. melanogaster* can endure chronic bacterial infections[72–74], albeit at rather low
levels, which can be the case in our experimental setup. Nonetheless, this holds true for both
populations, therefore not invalidating our interpretation of a different bacterial clearance rate
between populations.

Having shown that BactOral clears *P. entomophila* infection more rapidly, we
hypothesized that this response could rely on an increased production of immune effectors, such
as AMPs. Analysis of AMP gut gene expression showed that in unchallenged condition, only two
AMPs (*CecA1* and *Drsl5*) differed in expression between populations (down- and up-regulation
respectively), indicating that the base levels of AMP expression are unlikely to have been the
major target of selection. The detected expression difference of *Drsl5*, a gut-specific AMP
peptide, could represent a pre-emptive resistance mechanism acquired by BactOral, by
establishing a more antagonistic environment for *P. entomophila* proliferation in the guts of these
flies after ingestion. At 8 hours after exposure, two AMPs show a faster up-regulation while seven
of the panel members display a pattern of faster downregulation, in BactOral relative to Control.
These results suggest that a two-step process might be taking place, whereby the response is
initially (8h) more aggressive, permitting a more effective bacterial control, and is turned off
sooner (32h), limiting the secondary effects of prolonged immune responses (immunopathology).
Interestingly, Drosomycin-like genes (*Drsl2*, *Drsl3*, and *Drsl5*), whose expression is specific to
the gut epithelium and is regulated by JAK-STAT[35], show up-regulation after 8-hours of infection
in both populations. For all three genes, there is a slight tendency for a faster response that could
represent a more effective bacterial control in the BactOral population and a faster down-
regulation at the 32-hour timepoint that could limit self-inflicted damage. Likewise, we observed
a similar process of down-regulation of other AMPs specifically deployed to fight off infections
with Gram-negative pathogens (e.g. *CecA1* and *Diptericin*) at the 32-hour timepoint. In contrast,
some AMPs typically associated to response to Gram-positive bacteria, (e.g. *BomS2*, *BomS3* and
Drosomycin[51,75,76]), do not exhibit significant up-regulation after exposure to *P. entomophila* until
32-hours post-infection and only in the Control population.

Strikingly, and even though Diptericin has been shown to be specifically deployed by *D.*
melanogaster to resist *P. entomophila* infection[67], we did not detect a difference in the relative
expression of Diptericin between Control and BactOral until after 24 hours of infection. At this
timepoint, adapted flies are already returning to basal levels of AMP production whereas Control
flies keep actively fighting off the infection.

The same pattern is observed also in the priming experiment (Fig. 5 and Suppl. Fig. 4).
After priming with heat-killed bacteria and subsequent exposure to live bacteria, both populations
maintain an equivalent expression increase until 8 hours but from 24 hours onwards BactOral
shows, again, an earlier downregulation of *Diptericin* and a quicker return to homeostatic levels
of expression (Suppl. Fig. 4). In brief, the dynamics of *Diptericin* expression mimic the expected
effect of priming by establishing higher basal levels of this AMP with which to counteract an
incoming infection. This is not seen for the AMP *Drosomycin*, which again is consistent with our
previous result that this AMP is not deployed against infection with *P. entomophila* (Fig. 4 and
Suppl. Fig. 2).

With this we re-enforce the notion that this process could be an active response rather
than a consequence of lower bacterial loads, thus pointing to a potential role for disease tolerance
mechanisms in this population.

A role for Disease Tolerance

Disease tolerance encompasses many distinct mechanisms of action, and describes
multiple different physiological processes, which have in common the wide umbrella of
maintaining host fitness without directly impacting pathogen load[6,77]. Maintenance of fitness in
the face of stress-inducing or homeostatic conditions is assured by tissue damage control
mechanisms[6,77]. Furthermore, these damage control responses can be activated in the event of an
infection, either due to direct action of pathogens (i.e., injury-inducing toxins) or indirectly from
immune-derived damages (i.e., immunopathology).

With this in mind, a putative role for disease tolerance was tested by measuring survival
of the evolved populations to prolonged exposure to heat-killed bacteria. This protocol provided
a means by which to test the response to damage, whether provoked by pathogenicity of the
bacteria or by the host's own response, independently of resistance. A similar longevity was
observed for females and males of Control and BactOral after feeding on heat-killed bacteria, but
an interesting trend appeared when comparing survival profiles (Fig. 6). We found that BactOral
survived better to the stress of prolonged exposure to heat-killed *P. entomophila*, specifically
under an evolutionarily relevant timeframe, that is, the period when flies reproduced during the
selection experiment. Although both Control and BactOral individuals fully succumbed to the
chronic exposure to heat-killed bacteria, significant differences in survival distinguish the two
populations and their overall fitness. Physiological responses measured after flies are
approximately 10 days old, can potentially represent evolutionary costs[78–81],which are not the
focus of this study.

The improved response of adapted flies to exposure to damage-inducing substrate can be
explained by: 1) an increased capacity to withstand damage provoked by their own immune

response, 2) virulence factors present in inactivated bacterial supernatant, or 3) a correlation with

the previously identified tighter immune repression. Disentangling the relative contribution of

these mechanisms to the overall adapted phenotype would need further empirical characterization.

449   In sum, we were able to identify resistance and disease tolerance mechanisms as targets

of experimental evolution for increased survival to *P. entomophila* oral infection. Our

observations align with previous work that identifies a positive genetic correlation between these

two distinct immune defense strategies[82]. The increased survival of our experimentally adapted

population to oral infection directly relates to its capacity to decrease bacterial loads after

infection, independently of bacterial feeding rate, gut renewal capacity and bacterial defecation.

Previous work found resistance to be targeted by the selection of immune response in *D.*

melanogaster[79,83]. Our work reports one additional case of resistance evolution embedded in a

much wider ecological immunity context that exhaustively approaches the different processes by

which immunity may evolve in populations fighting off persistent infections. Indeed, in addition

to resistance, we could find a quantifiable role for disease tolerance in the evolved response of

BactOral, revealing the complex nature of this adaptative process[82], which is likely to rely also

on improved tissue repair coupled with a more controlled deployment of damage-inducing

immune effectors, such as AMPs[84,85]. Future work will focus on identifying the specific

mechanisms of disease tolerance and their physiological mode of action and interactions with

resistance.

Materials and Methods

Maintenance of Drosophila populations under relaxed selection

Experimentally evolved Control and BactOral populations were initially derived from a highly outbred population of *D. melanogaster*, as described in previous studies [19,46]. After selection was terminated, populations were kept at high census (1500-2000 individuals per cage), allowing for maintenance of high genetic variability and optimal larval development. For all generations, flies were kept under constant temperature (25°C), humidity (55–65%) and light-darkness cycle (12:12), and fed with a standard cornmeal-agar medium, consisting of 4.5% molasses, 7.5% sugar, 7% corn-flower, 2% granulated yeast extract, 1% agar and 0.25% nipagin, mixed in distilled water.

Each generation cycle lasted approximately 3 weeks, during which flies ecloded and remained undisturbed until they were 7-8 days old. At this time, fresh food was placed in the cages for 2 days, which allowed for oviposition of retained eggs. After this period (when flies were 9-11 days old), similarly to the protocol applied for selection, new fresh food was placed in the cages for controlled egg-lays (1-3 hours long) and establishment of the next generation.

Pathogen culture and infections

Oral infection protocol with *P. entomophila* was adapted from the one used in Martins and Faria et al[19]. Briefly, single bacterial colonies were grown in kick-start cultures (5mL of LB) for approximately 8 hours after which they were transferred into larger volumes (1 L of LB) for overnight growth, both periods at 29ºC. After centrifugation and resuspension of bacterial pellets, concentration was adjusted to OD_{600}= 100 in PBS and finally diluted 1:1 with a 5% sucrose solution. For infection, 3-5 days-old flies were separated by sex into groups of twenty with the use of CO_2 (at least 24 hours before the start of the experiments) and allowed to feed on a filter disk embedded with the bacterial solution described above for 24 hours (for survival assays) or for the amount of time specific to each experiment (see Results). After this period, infected flies were flipped onto clean food and survival was scored daily. Controls were fed a 1:1 PBS:5% sucrose solution.

For the artificial immune activation experiment, a solution of *P. entomophila* at OD_{600}= 100 was previously heat-killed for 1 hour at 55ºC, mixed 1:1 with a 5% sucrose solution and fed to flies for 2 hours, in a similar way as described above for regular oral infection. After this time, flies were transferred into tubes containing a live solution of *P. entomophila* for 24 hours, after which they were changed back to clean food and their survival scored.

For the experiment with prolonged exposure to heat-killed *P. entomophila* a similar
bacterial preparation and heat-killing protocol were performed as described above, but the final
dead bacteria suspension was mixed 1:1 with standard fly food and dispensed into fly vials. To
the final solution, extra agar powder was added to account for viscosity and to optimize texture
and left to cool down and solidify for at least 2 hours, before being fed to flies. The same protocol
was performed for the control treatment, in which fly food was mixed with sterile 1X PBS.
Exposure to these treatments was constant throughout the experiment, during which replicates of
15 females and 15 males were co-housed in vials, exclusively feeding on the dead-bacteria or
PBS mixtures. Survival scoring began at the start of exposure and lasted 14 days. During the
treatment, flies were frequently flipped onto fresh food mixture and survival scoring continued
daily.

Survival analysis was done for the multiple experiments by fitting data with Cox
Proportional Hazards models (with *coxme* or *coxph* functions from the *survival* package[86]) using
population, treatment, and sex as fixed factors and replicate (or replicate nested within block) as
a random factor. To assess the effect sizes of each factor, type 3 ANOVAs (*Anova* from the *car*
package[87]) were ran on the survival models. Additionally, multiple comparisons with *tukey*
adjustments were done on all the models to disentangle which specific pairwise comparisons were
significantly different (*emmeans* function[88]). Finally, the *cld* function of the *multcomp* package[89]
(with *sidak* confidence-level adjustments) was used to summarize the results of the multiple
comparisons across conditions. Survival models used were:

immunocompetence (Fig. 1)- *coxme*(~Population*Treatment*Sex+(1|Replicate));

priming (Fig. 5)- *coxme*(~Population*Treatment*Sex+(1|block/Replicate)) and

immune over-activation (Fig. 6)– *coxme*(~Population*Treatment*Sex+(1|Replicate), data =
survival_HKexposure).

Bacterial defecation assay

To have a measure of the rate of gut purge/defecation a change was introduced to the
bacterial infection protocol, whereby the feeding period was reduced to a 3-hour period. After
this time, flies were surface sterilized and separated individually into plastic spectrophotometry
cuvettes, previously filled with fly food. After 6 hours (6 hours.), flies were changed into fresh
cuvettes where they remained for an additional 12 hours (18 hours.), after which they were
discarded. Bacterial quantification was performed on the interior surface of the cuvettes, for both
timepoints, as described below.

Statistical analysis on number of defecated bacteria was done by fitting a zero-inflated on

the CFU counts and zeros (using the *zeroinfl* function[90] from the *pscl* package[91]). This model

allowed for the fitting of a negative-binomial distribution on the counts data and of a binomial

distribution on the zeros (*zeroinfl*(formula = Counts ~ Population * Sex * Timepoint | Population

* Timepoint, dist = "negbin")). To assess differences between specific pairwise comparisons we

used *emmeans* with *tukey* adjustments, on both distributions of the zero-inflated model. In the

instances where flies died during either timepoint, their count at that time was removed from

further analysis and the linear regression was not included in Figure 2B.

Quantification of bacterial loads

To quantify bacterial loads we adapted an established protocol[19,92]. In short, individual

flies were surface sterilized (washed in EtOH 70%, bleach 60% and two washes with MiliQ water)

and placed in 96 deep-well plates, where they were homogenized with glass beads in 50 µL of

LB using a TissueLyser II (Qiagen). Individual samples were serially diluted 5 times with 1:10

dilution steps, plated as 4 µL droplets in LB+Rifampicin, and left to grow overnight at 29ºC to

optimize colony size for counting. For the bacterial defecation assay, cuvettes that contained flies

for each of the timepoints were filled with 1mL of PBS and incubated at room temperature for 15

minutes, after which each sample was diluted up to 4 times (1:5 dilutions) and platted as droplets

in LB+Rifampicin.

Analysis of the time-course of infection progression was performed by dividing the data

obtained from the bacterial quantification into "counts" and "zeros" and using different models

to examine their dynamics in our evolved populations. For the counts, the modelling of the mean

of the load dynamics during the time-course was done using a sigmoidal function that considers

four parameters: a the log (initial load), k the timepoint at which the load is half of the maximum

value (in this case, a), h the slope at the k timepoint (steepest slope of the curve), and b the log

(final load). A linear model was applied to each of the parameters which was fitted using a Monte

Carlo Markov Chain approach. To estimate the impact of Sex and Population on the likely

distribution of the parameters, a normal error structure was assumed, using Control females as

reference. For the "zeros", a modified logistic model with four parameters was ran: a_0 initial

fraction of zeros, k_0 rate of zero disappearance, k_1 rate of zero re-appearance, t_0 timepoint at which

zeros re-appear and a_1 final fraction of zeros. Similarly, to the "counts" analysis, linear models

were used to assess the impact of Sex and Population on all four parameters.

Quantification of food intake by CApillary FEeder Assay - CAFE

To measure the amount of bacterial solution ingested by single flies, an adapted CAFE[47]

assay was performed. We substituted the food of small fly vials with agarose, to ensure humidity,

and placed a glass capillary in each lid, filled with 5% sucrose solution. We allowed individual

flies to habituate to this experimental setup for 24 hours before the infection. After this period,

capillaries filled with *P. entomophila* solution (prepared as described before) were given to

individual females to feed on and consumption estimated, at different timepoints. To facilitate

scoring in capillaries, the bacterial solution was mixed with blue food colouring. Controls were

fed dyed solution with equivalent dilution but without bacteria.

Modelling of cumulative amounts of bacteria eaten over time was done with a

Generalized Linear Model with a Gamma distribution, (*glm*(Quantity ~ Population*Timepoint))

using the *glm* function from the base *stat* R package[93]. For better visualization of feeding rates,

linear regressions were drawn between mean values of microliters eaten cumulatively by each

population at each timepoint, using the *stat_poly_eq* function of the *ggpmisc* package[94]. Finally,

multiple comparisons were done with *emmeans*[88], using *tukey* adjustments, to disentangle

differences between populations across timepoints.

RNA extractions and quantitative PCR

For RNA extractions, flies from Control and BactOral population were infected as

described above and, at different timepoints, pools of females were collected and homogeneized

in 500 µL of Trizol. RNA extractions were performed using a Direct-zol™ RNA MiniPrep w/

Zymo-Spin™ IIC Columns. After precipitation, a DNase I (RQ1 RNASE-FREE DNASE 1* from

Promega®) treatment was done to all samples, followed by reverse transcription using the

Thermo Scientific® RevertAid H Minus cDNA kit. cDNA was finally diluted 1:5 for qPCR.

For quantification of gene expression, qPCRs were performed using SYBR™ Green

Master Mix (Thermo Scientific®) and reactions ran on 384-well plates (Applied Biosystems®).

The PCR conditions used in all experiments were: initial denaturation/ enzyme activation, 95ºC

for 10'; followed by 45 cycles of denaturation, 95ºC for 10"; annealing, 60ºC for 10"; extension,

72ºC for 30". Sequences of primers used for qPCRs are shown in Supp. Table. 1. Either EIF2

(eIF-2alpha) or Rpl32 were used as reference.

Analysis of gene expression differences was done with the relative quantification

technique ($\Delta\Delta$Ct)[95]. Briefly, to the average of technical replicates for each candidate gene we

subtracted the average of the Ct values of the respective sample's house-keeping gene (EIF2 or

Rpl32) (ΔCt) and normalized this value to the ΔCt of the respective reference condition. For the

time-course analysis, the reference was the "Unchallenged" timepoint while for comparisons

between "Unchallenged" conditions among populations, Control was used as reference. Finally,

we used reverse logarithm to transform the final gene expression values into fold change levels

($2^{-\Delta\Delta Ct}$) and analysed the Log$_2$(foldchange) differences across Populations and/or Timepoints.

Statistical analysis on gene expression was done by running Linear Models (from base R

stats[93]) on the Log$_2$(foldchange) values of each individual gene (*lm*(logfold_AMP ~ Population *

Timepoint), followed by an ANOVA (type three) from the car package[87]. To individualize

significant pairwise comparisons, *emmeans* was ran on the linear models (with a Benjamini-

Hochberg adjustment). Finally, the *cld* function of the multcomp package[89] (with sidak

confidence-level adjustments) was used to summarize the results of the multiple comparisons

across different conditions.

Measurement of gut length and crop size

Females were individually dissected in PBS 1X, and their guts removed for imaging by

adapting a protocol described in[96]. Dissections started by fixing flies in the head with a Minutien

pin and gradually separating the gut from the remaining tissues, in a posterior-to-anterior

direction. After the gut was exposed, we added a few drops of a solution of 5% Acetic Acid and

50%EtOH, which slightly whitens the tissues, allowing for better visualization and a slight tissue

fixation. After having isolated the guts, they were individually transferred to a coverslip and

photographed using a uEye camera installed on an Olympus SZX7 stereoscope.

Images were processed in FIJI v2.0.0 and measurements were collected by drawing

consecutive transects along the total length of guts, as illustrated in Supplementary figure 1A.

Each measurement was repeated 3 times, to account for errors in the manual drawing of the

transects and averaged to obtain a final gut length in millimetres (mm).

Differences in lengths of guts between populations were analysed with a Linear Mixed-

Effects Model (*lmer*(Length ~ Population * Treatment + (1|Block)) with *lmer* from the *lmerTest*

package[97]. The model was followed by multiple comparisons with *emmeans*[88], with a *tukey*

adjustment, for assessment of individual pairwise differences.

Statistical analysis and graphical representations

All datasets (excluding survival data) were tested for Normality before further analysis,

using the Shapiro-Wilk Test (*shapiro.test* function of the *stat* R package), for determination of

the most suitable type of statistical test to perform in each case. All model formulas used for

statistical analysis are discriminated throughout the Results sections. All graphical representations

were performed using the *ggplot2* package[98].

Statistical analysis and graphics for all experiments, excluding analysis of the bacterial

loads time-course modelling analysis (Fig. 5), were done on R version 4.2.1, through RStudio[99]

version 1.3.959. For the bacterial dynamics time-course, models were implemented and fitted in

Python[100], using the probabilistic programming language package *pymc3*[101] version 3.11.

Acknowledgements

We thank Dr Bruno Lemaitre for stocks, Liliana Vieira and Sandra Crisóstomo for technical assistance, André Barros and David Duneau for help with analyses, and Elvira Lafuente, Diogo Roque and Patrícia Duarte for critical comments and discussions that improved the manuscript. This work was supported by Instituto Gulbenkian de Ciência/Fundação Calouste Gulbenkian and FCT-Fundação para a Ciência e a Tecnologia (Portugal) UID/Multi/04555/2013 and UIDB/00329/2020. TFP (BD/128432/2017 and COVID/BD/151645/2021) and to PAA (BD/06404/2020) were supported FCT—Fundação para a Ciência e a Tecnologia (Portugal); and by C.S. CONGENTO, project LISBOA-01-0145-FEDER-022170, co-financed by Lisboa Regional Operational Programme (Lisboa 2020), under the Portugal 2020 Partnership Agreement, through the European Regional Development Fund (ERDF), and Foundation for Science and Technology (Portugal).

Author Contributions

Conceptualization: Tânia F. Paulo and Élio Sucena.

Data curation: Tânia F. Paulo

Formal analysis: Tânia F. Paulo, Tiago Paixão

Funding acquisition: Élio Sucena.

Investigation: Tânia F. Paulo, Priscilla A. Akyaw and Élio Sucena.

Methodology: Tânia F. Paulo, Priscilla A. Akyaw, Tiago Paixão and Élio Sucena.

Resources: Élio Sucena and Tiago Paixão.

Supervision: Élio Sucena.

Visualization: Tânia F. Paulo.

Writing – original draft: Tânia F. Paulo and Élio Sucena.

Writing – review & editing: Tânia F. Paulo, Priscilla A. Akyaw, Tiago Paixão and Élio Sucena.

References

(1)   Fischer, O.; Schmid-Hempel, P. Selection by parasites may increase host recombination frequency. *Biol. Lett.* **2005**. https://doi.org/10.1098/rsbl.2005.0296.

(2)   Schmid-Hempel, P.; Ebert, D. On the evolutionary ecology of specific immune defence. *Trends Ecol. Evol.* **2003**, *18* (1), 27–32. https://doi.org/10.1016/S0169-5347(02)00013-7.

(3)   Schulenburg, H.; Kurtz, J.; Moret, Y.; Siva-Jothy, M. T. Introduction. Ecological immunology. *Philos. Trans. R. Soc. B Biol. Sci.* **2008**, *364* (1513), 3–14. https://doi.org/10.1098/rstb.2008.0249.

(4)   Gibson, A. K.; Amoroso, C. R. Evolution and ecology of parasite avoidance. *Annu. Rev. Ecol. Evol. Syst.* **2022**, *53* (1), 47–67. https://doi.org/10.1146/annurev-ecolsys-102220-020636.

(5)   Kutzer, M. A. M.; Armitage, S. A. O. Maximising fitness in the face of parasites: a review of host tolerance. *Zoology* **2016**, *119* (4), 281–289. https://doi.org/10.1016/j.zool.2016.05.011.

(6)   Medzhitov, R.; Schneider, D. S.; Soares, M. P. Disease tolerance as a defense strategy. *Science* **2012**, *335* (6071), 936–941. https://doi.org/10.1126/science.1214935.

(7)   Ayres, J. S.; Schneider, D. S. Two ways to survive an infection: what resistance and tolerance can teach us about treatments for infectious diseases. *Nat. Rev. Immunol.* **2008**, *8* (11), 889–895. https://doi.org/10.1038/nri2432.

(8)   Råberg, L.; Graham, A. L.; Read, A. F. Decomposing health: tolerance and resistance to parasites in animals. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **2009**, *364* (1513), 37–49. https://doi.org/10.1098/rstb.2008.0184.

(9)   Menge, D. M.; Zhong, D.; Guda, T.; Gouagna, L.; Githure, J.; Beier, J.; Yan, G. Quantitative Trait Loci controlling refractoriness to *Plasmodium falciparum* in natural *Anopheles Gambiae* mosquitoes from a malaria-endemic region in western Kenya. *Genetics* **2006**, *173* (1), 235–241. https://doi.org/10.1534/genetics.105.055129.

(10)   Kim, D. H.; Feinbaum, R.; Alloing, G.; Emerson, F. E.; Garsin, D. A.; Inoue, H.; Tanaka-Hino, M.; Hisamoto, N.; Matsumoto, K.; Tan, M.-W.; Ausubel, F. M. A conserved P38 MAP kinase pathway in *Caenorhabditis elegans* innate immunity. *Science* **2002**, *297* (5581), 623–626. https://doi.org/10.1126/science.1073759.

(11)   Scharsack, J. P.; Kalbe, M.; Harrod, C.; Rauch, G. Habitat-Specific Adaptation of immune responses of stickleback (*Gasterosteus Aculeatus*) lake and river ecotypes. *Proc. R. Soc. B Biol. Sci.* **2007**, *274* (1617), 1523–1532. https://doi.org/10.1098/rspb.2007.0210.

(12)   Faria, V. G.; Martins, N. E.; Magalhães, S.; Paulo, T. F.; Nolte, V.; Schlötterer, C.; Sucena, É.;

Teixeira, L. Drosophila adaptation to viral infection through defensive symbiont evolution. *PLOS Genet.* **2016**, *12* (9), e1006297. https://doi.org/10.1371/journal.pgen.1006297.

(13)   Martins, N. E.; Faria, V. G.; Nolte, V.; Schlotterer, C.; Teixeira, L.; Sucena, E.; Magalhaes, S. Host adaptation to viruses relies on few genes with different cross-resistance properties. *Proc. Natl. Acad. Sci.* **2014**, *111* (16), 5938–5943. https://doi.org/10.1073/pnas.1400378111.

(14)   Gupta, V.; Venkatesan, S.; Chatterjee, M.; Syed, Z. A.; Nivsarkar, V.; Prasad, N. G. No apparent cost of evolved immune response in *Drosophila melanogaster. Evolution* **2016**, *70* (4), 934–943. https://doi.org/10.1111/evo.12896.

(15)   Zbinden, M.; Haag, C. R.; Ebert, D. Experimental evolution of field populations of *Daphnia magna* in response to parasite treatment. *J. Evol. Biol.* **2008**, *21* (4), 1068–1078. https://doi.org/10.1111/j.1420-9101.2008.01541.x.

(16)   Kawecki, T. J.; Lenski, R. E.; Ebert, D.; Hollis, B.; Olivieri, I.; Whitlock, M. C. Experimental evolution. *Trends Ecol. Evol.* **2012**, *27* (10), 547–560. https://doi.org/10.1016/j.tree.2012.06.001.

(17)   Ebert, D. Experimental evolution of parasites. *Science* **1998**, *282* (5393), 1432–1436. https://doi.org/10.1126/science.282.5393.1432.

(18)   Marxer, M.; Barribeau, S.; Schmid-Hempel, P. Experimental evolution of a trypanosome parasite of bumblebees and its implications for infection success and host immune response. *Evol. Biol.* **2016**, *43* (2), 160–170. https://doi.org/10.1007/s11692-015-9366-z.

(19)   Martins, N. E.; Faria, V. G.; Teixeira, L.; Magalhães, S.; Sucena, É. Host adaptation is contingent upon the infection route taken by pathogens. *PLoS Pathog.* **2013**, *9* (9), e1003601. https://doi.org/10.1371/journal.ppat.1003601.

(20)   Khan, I.; Prakash, A.; Agashe, D. Experimental evolution of insect immune memory versus pathogen resistance. *Proc. R. Soc. B Biol. Sci.* **2017**, *284* (1869), 20171583. https://doi.org/10.1098/rspb.2017.1583.

(21)   Cabrera, K.; Hoard, D. S.; Gibson, O.; Martinez, D. I.; Wunderlich, Z. *Drosophila* immune priming to *Enterococcus faecalis* relies on immune tolerance rather than resistance. *PLOS Pathog.* **2023**, *19* (8), e1011567. https://doi.org/10.1371/journal.ppat.1011567.

(22)   Mondotte, J. A.; Saleh, M.-C. Chapter Eleven - Antiviral immune response and the route of infection in *Drosophila melanogaster*. In *Advances in Virus Research*; Kielian, M., Mettenleiter, T. C., Roossinck, M. J., Eds.; Academic Press, 2018; Vol. 100, pp 247–278. https://doi.org/10.1016/bs.aivir.2017.10.006.

(23)   Vodovar, N.; Vinals, M.; Liehl, P.; Basset, A.; Degrouard, J.; Spellman, P.; Boccard, F.;

Lemaitre, B. *Drosophila* host defense after oral infection by an entomopathogenic *Pseudomonas* species. *Proc. Natl. Acad. Sci. U. S. A.* **2005**, *102* (32), 11414–11419. https://doi.org/10.1073/pnas.0502240102.

(24)   Chandler, J. A.; Lang, J. M.; Bhatnagar, S.; Eisen, J. A.; Kopp, A. Bacterial communities of diverse *Drosophila* species: Ecological Context of a Host–Microbe Model System. *PLOS Genet.* **2011**, *7* (9), e1002272. https://doi.org/10.1371/journal.pgen.1002272.

(25)   Charroux, B.; Daian, F.; Royet, J. *Drosophila* aversive behavior toward *Erwinia carotovora carotovora* is mediated by bitter neurons and leukokinin. *iScience* **2020**, *23* (6), 101152. https://doi.org/10.1016/j.isci.2020.101152.

(26)   Soldano, A.; Alpizar, Y. A.; Boonen, B.; Franco, L.; López-Requena, A.; Liu, G.; Mora, N.; Yaksi, E.; Voets, T.; Vennekens, R.; Hassan, B. A.; Talavera, K. Gustatory-Mediated Avoidance of bacterial lipopolysaccharides via TRPA1 activation in *Drosophila*. *eLife* **2016**, *5* (JUN2016), 1–15. https://doi.org/10.7554/eLife.13133.

(27)   Kuraishi, T.; Binggeli, O.; Opota, O.; Buchon, N.; Lemaitre, B. Genetic Evidence for a protective role of the peritrophic matrix against intestinal bacterial infection in *Drosophila melanogaster*. *Proc. Natl. Acad. Sci. U. S. A.* **2011**, *108* (38), 15966–15971. https://doi.org/10.1073/pnas.1105994108.

(28)   Overend, G.; Luo, Y.; Henderson, L.; Douglas, A. E.; Davies, S. A.; Dow, J. A. T. Molecular mechanism and functional significance of acid generation in the *Drosophila* midgut. *Nat. Publ. Group* **2016**, No. May, 1–11. https://doi.org/10.1038/srep27242.

(29)   Du, E. J.; Ahn, T. J.; Kwon, I.; Lee, J. H.; Park, J. H.; Park, S. H.; Kang, T. M.; Cho, H.; Kim, T. J.; Kim, H. W.; Jun, Y.; Lee, H. J.; Lee, Y. S.; Kwon, J. Y.; Kang, K. J. TrpA1 regulates defecation of food-borne pathogens under the control of the Duox pathway. *PLoS Genet.* **2016**, *12* (1), 1–21. https://doi.org/10.1371/journal.pgen.1005773.

(30)   Fink, C.; Hoffmann, J.; Knop, M.; Li, Y.; Isermann, K.; Roeder, T. Intestinal FoxO signaling is required to survive oral infection in *Drosophila*. *Mucosal Immunol.* **2016**, *9* (4), 927–936. https://doi.org/10.1038/mi.2015.112.

(31)   Ha, E.-M.; Oh, C.-T.; Bae, Y. S.; Lee, W.-J. A direct role for dual Oxidase in *Drosophila* gut immunity. *Science* **2005**, *310* (5749), 847–850. https://doi.org/10.1126/science.1117311.

(32)   Ha, E.-M.; Lee, K.-A.; Seo, Y. Y.; Kim, S.-H.; Lim, J.-H.; Oh, B.-H.; Kim, J.; Lee, W.-J. Coordination of multiple dual Oxidase–regulatory pathways in responses to commensal and infectious microbes in *Drosophila* gut. *Nat. Immunol.* **2009**, *10* (9), 949–957. https://doi.org/10.1038/ni.1765.

(33)   Tian, A.; Morejon, V.; Kohoutek, S.; Huang, Y.-C.; Deng, W.-M.; Jiang, J. Damage-induced regeneration of the intestinal stem cell pool through enteroblast mitosis in the *Drosophila* midgut. *EMBO J.* **2022**, e110834. https://doi.org/10.15252/embj.2022110834.

(34)   Bonfini, A.; Liu, X.; Buchon, N. From Pathogens to Microbiota: How Drosophila intestinal stem cells react to gut microbes. *Dev. Comp. Immunol.* **2015**. https://doi.org/10.1016/j.dci.2016.02.008.

(35)   Buchon, N.; Broderick, N. A.; Poidevin, M.; Pradervand, S.; Lemaitre, B. *Drosophila* intestinal response to bacterial infection: Activation of host defense and stem cell proliferation. *Cell Host Microbe* **2009**, *5* (2), 200–211. https://doi.org/10.1016/j.chom.2009.01.003.

(36)   Buchon, N.; Broderick, N. A.; Kuraishi, T.; Lemaitre, B. *Drosophila* EGFR pathway coordinates stem cell proliferation and gut remodeling following infection. *BMC Biol.* **2010**, *8* (1), 152. https://doi.org/10.1186/1741-7007-8-152.

(37)   Nehme, N. T.; Liégeois, S.; Kele, B.; Giammarinaro, P.; Pradel, E.; Hoffmann, J. A.; Ewbank, J. J.; Ferrandon, D. A model of bacterial intestinal infections in *Drosophila melanogaster. PLoS Pathog.* **2007**, *3* (11), 1694–1709. https://doi.org/10.1371/journal.ppat.0030173.

(38)   Limmer, S.; Haller, S.; Drenkard, E.; Lee, J.; Yu, S.; Kocks, C.; Ausubel, F. M.; Ferrandon, D. *Pseudomonas aeruginosa* RhlR is required to neutralize the cellular immune response in a *Drosophila melanogaster* oral infection model. *Proc. Natl. Acad. Sci. U. S. A.* **2011**, *108* (42), 17378–17383. https://doi.org/10.1073/pnas.1114907108.

(39)   Yang, S.; Zhao, Y.; Yu, J.; Fan, Z.; Gong, S.; Tang, H.; Pan, L. Sugar alcohols of Polyol pathway serve as Alarmins to mediate local-systemic innate immune communication in *Drosophila. Cell Host Microbe* **2019**, *26* (2), 240-251.e8. https://doi.org/10.1016/j.chom.2019.07.001.

(40)   Zaidman-Rémy, A.; Hervé, M.; Poidevin, M.; Pili-Floury, S.; Kim, M.-S.; Blanot, D.; Oh, B.-H.; Ueda, R.; Mengin-Lecreulx, D.; Lemaitre, B. The *Drosophila* Amidase PGRP-LB modulates the immune response to bacterial infection. *Immunity* **2006**, *24* (4), 463–473. https://doi.org/10.1016/j.immuni.2006.02.012.

(41)   Basset, A.; Khush, R. S.; Braun, A.; Gardan, L.; Boccard, F.; Hoffmann, J. A.; Lemaitre, B. The phytopathogenic bacteria *Erwinia carotovora* infects *Drosophila* and activates an immune response. *Proc. Natl. Acad. Sci.* **2000**, *97* (7), 3376–3381. https://doi.org/10.1073/pnas.97.7.3376.

(42)   Vodovar, N.; Vallenet, D.; Cruveiller, S.; Rouy, Z.; Barbe, V.; Acosta, C.; Cattolico, L.; Jubin, C.; Lajus, A.; Segurens, B.; Vacherie, B.; Wincker, P.; Weissenbach, J.; Lemaitre, B.; Médigue, C.; Boccard, F. Complete genome sequence of the entomopathogenic and metabolically versatile soil bacterium *Pseudomonas entomophila. Nat. Biotechnol.* **2006**, *24* (6), 673–679. https://doi.org/10.1038/nbt1212.

(43)   Dieppois, G.; Opota, O.; Lalucat, J.; Lemaitre, B. *Pseudomonas entomophila*: a versatile bacterium with entomopathogenic properties. In *Pseudomonas*; Springer Netherlands: Dordrecht, 2015; pp 25–49. https://doi.org/10.1007/978-94-017-9555-5_2.

(44)   Opota, O.; Vallet-Gély, I.; Vincentelli, R.; Kellenberger, C.; Iacovache, I.; Gonzalez, M. R.; Roussel, A.; Goot, F.-G. van der; Lemaitre, B. Monalysin, a novel ß-pore-forming toxin from the *Drosophila* pathogen *Pseudomonas entomophila*, contributes to host intestinal damage and lethality. *PLOS Pathog.* **2011**, *7* (9), e1002259. https://doi.org/10.1371/journal.ppat.1002259.

(45)   Chakrabarti, S.; Liehl, P.; Buchon, N.; Lemaitre, B. Infection-induced host translational blockage inhibits immune responses and epithelial renewal in the *Drosophila* gut. *Cell Host Microbe* **2012**, *12* (1), 60–70. https://doi.org/10.1016/j.chom.2012.06.001.

(46)   Faria, V. G.; Martins, N. E.; Paulo, T.; Teixeira, L.; Sucena, É.; Magalhães, S. Evolution of *Drosophila* resistance against different pathogens and infection routes entails no detectable maintenance costs. *Evolution* **2015**, *69* (11), 2799–2809. https://doi.org/10.1111/evo.12782.

(47)   Ja, W. W.; Carvalho, G. B.; Mak, E. M.; De La Rosa, N. N.; Fang, A. Y.; Liong, J. C.; Brummel, T.; Benzer, S. Prandiology of *Drosophila* and the CAFE assay. *Proc. Natl. Acad. Sci. U. S. A.* **2007**, *104* (20), 8253–8256. https://doi.org/10.1073/pnas.0702726104.

(48)   Vincent, C. M.; Dionne, M. S. Disparate regulation of IMD signaling drives sex differences in infection pathology in *Drosophila melanogaster. Proc. Natl. Acad. Sci.* **2021**, *118* (32), e2026554118. https://doi.org/10.1073/pnas.2026554118.

(49)   Belmonte, R. L.; Corbally, M.-K.; Duneau, D. F.; Regan, J. C. Sexual dimorphisms in innate immunity and responses to infection in *Drosophila melanogaster. Front. Immunol.* **2020**, *10*.

(50)   Duneau, D. F.; Kondolf, H. C.; Im, J. H.; Ortiz, G. A.; Chow, C.; Fox, M. A.; Eugénio, A. T.; Revah, J.; Buchon, N.; Lazzaro, B. P. The Toll pathway underlies host sexual dimorphism in resistance to both Gram-negative and Gram-positive bacteria in mated *Drosophila. BMC Biol.* **2017**, *15* (1), 124. https://doi.org/10.1186/s12915-017-0466-3.

(51)   Hanson, M. A.; Dostálová, A.; Ceroni, C.; Poidevin, M.; Kondo, S.; Lemaitre, B. Synergy and remarkable specificity of antimicrobial peptides *in vivo* using a systematic knockout approach. *eLife* **2019**, *8*, e44341. https://doi.org/10.7554/eLife.44341.

(52)   Carboni, A. L.; Hanson, M. A.; Lindsay, S. A.; Wasserman, S. A.; Lemaitre, B. Cecropins contribute to *Drosophila* host defense against a subset of fungal and Gram-negative bacterial infection. *Genetics* **2022**, *220* (1), iyab188. https://doi.org/10.1093/genetics/iyab188.

(53)   Keesey, I. W.; Koerte, S.; Khallaf, M. A.; Retzke, T.; Guillou, A.; Grosse-Wilde, E.; Buchon,

N.; Knaden, M.; Hansson, B. S. Pathogenic bacteria enhance dispersal through alteration of *Drosophila* social communication. *Nat. Commun.* **2017**, *8* (1), 265. https://doi.org/10.1038/s41467-017-00334-9.

(54) Patrnogic, J.; Castillo, J. C.; Shokal, U.; Yadav, S.; Kenney, E.; Heryanto, C.; Ozakman, Y.; Eleftherianos, I. Pre-exposure to non-pathogenic bacteria does not protect *Drosophila* against the entomopathogenic bacterium *Photorhabdus*. *PLOS ONE* **2018**, *13* (10), e0205256. https://doi.org/10.1371/journal.pone.0205256.

(55) Wen, Y.; He, Z.; Xu, T.; Jiao, Y.; Liu, X.; Wang, Y.-F.; Yu, X.-Q. Ingestion of killed bacteria activates antimicrobial Peptide Genes in *Drosophila melanogaster* and protects flies from septic infection. *Dev. Comp. Immunol.* **2019**, *95*, 10–18. https://doi.org/10.1016/j.dci.2019.02.001.

(56) Paul Schmid-Hempel. Evolutionary ecology of insect immune defenses. *Annu Rev Entomol* **2005**, *50*, 529–551. https://doi.org/10.1146/annurev.ento.50.071803.130420.

(57) Eleftherianos, I.; Tafesh-Edwards, G.; Mohamed, A. Pathogen infection routes and host innate immunity: lessons from insects. *Immunol. Lett.* **2022**, *247*, 46–51. https://doi.org/10.1016/j.imlet.2022.05.006.

(58) Gupta, V.; Vasanthakrishnan, R. B.; Siva-Jothy, J.; Monteith, K. M.; Brown, S. P.; Vale, P. F. The route of infection determines *Wolbachia* antibacterial protection in *Drosophila*. *Proc. R. Soc. B Biol. Sci.* **2017**, *284* (1856), 20170809. https://doi.org/10.1098/rspb.2017.0809.

(59) Vizoso, D. B.; Ebert, D. Phenotypic plasticity of host–parasite interactions in response to the route of infection. *J. Evol. Biol.* **2005**, *18* (4), 911–921. https://doi.org/10.1111/j.1420-9101.2005.00920.x.

(60) Dekmak, A. S.; Yang, X.; Dohna, H. Z.; Buchon, N.; Osta, M. A. The Route of infection influences the contribution of key immunity genes to antibacterial defense in *Anopheles gambiae. J. Innate Immun.* **2021**, *13* (2), 107–126. https://doi.org/10.1159/000511401.

(61) Hall, M. D.; Bento, G.; Ebert, D. The evolutionary consequences of stepwise infection processes. *Trends Ecol. Evol.* **2017**, *xx*, 1–12. https://doi.org/10.1016/j.tree.2017.05.009.

(62) Vale, P. F.; Jardine, M. D. Infection avoidance behavior: viral exposure reduces the motivation to forage in female *Drosophila melanogaster*. *Fly (Austin)* **2017**, *11* (1), 3–9. https://doi.org/10.1080/19336934.2016.1207029.

(63) Kobler, J. M.; Rodriguez Jimenez, F. J.; Petcu, I.; Grunwald Kadow, I. C. Immune receptor signaling and the mushroom body mediate post-ingestion pathogen avoidance. *Curr. Biol.* **2020**, *30* (23), 4693-4709.e3. https://doi.org/10.1016/j.cub.2020.09.022.

(64) Stensmyr, M. C.; Dweck, H. K. M.; Farhan, A.; Ibba, I.; Strutz, A.; Mukunda, L.; Linz, J.;

Grabe, V.; Steck, K.; Lavista-Llanos, S.; Wicher, D.; Sachse, S.; Knaden, M.; Becher, P. G.; Seki, Y.; Hansson, B. S. A conserved dedicated olfactory circuit for detecting harmful microbes in D*rosophila*. *Cell* **2012**, *151* (6), 1345–1357. https://doi.org/10.1016/j.cell.2012.09.046.

(65)   Ahlawat, N.; Geeta Arun, M.; Maggu, K.; Jigisha; Singh, A.; Prasad, N. G. *Drosophila melanogaster* hosts coevolving with *Pseudomonas entomophila* pathogen show sex-specific patterns of local adaptation. *BMC Ecol. Evol.* **2022**, *22* (1), 77. https://doi.org/10.1186/s12862-022-02031-8.

(66)   Faria, V. G.; Martins, N. E.; Schlötterer, C.; Sucena, É. Readapting to DCV infection without *Wolbachia*: frequency changes of *Drosophila* antiviral alleles can replace endosymbiont protection. *Genome Biol. Evol.* **2018**, *10* (7), 1783–1791. https://doi.org/10.1093/gbe/evy137.

(67)   Liehl, P.; Blight, M.; Vodovar, N.; Boccard, F.; Lemaitre, B. Prevalence of local immune Response against oral infection in a *Drosophila/Pseudomonas* infection model. *PLoS Pathog.* **2006**, *2* (6), e56. https://doi.org/10.1371/journal.ppat.0020056.

(68)   Liu, X.; Nagy, P.; Bonfini, A.; Houtz, P.; Bing, X.-L.; Yang, X.; Buchon, N. Microbes affect gut epithelial cell composition through immune-dependent regulation of intestinal stem cell differentiation. *Cell Rep.* **2022**, *38* (13). https://doi.org/10.1016/j.celrep.2022.110572.

(69)   Hudry, B.; Khadayate, S.; Miguel-Aliaga, I. The sexual identity of adult intestinal stem cells controls organ size and plasticity. *Nature* **2016**, *530* (7590), 344–348. https://doi.org/10.1038/nature16953.

(70)   Hudry, B.; de Goeij, E.; Mineo, A.; Gaspar, P.; Hadjieconomou, D.; Studd, C.; Mokochinski, J. B.; Kramer, H. B.; Plaçais, P.-Y.; Preat, T.; Miguel-Aliaga, I. Sex differences in intestinal carbohydrate metabolism promote food intake and sperm maturation. *Cell* **2019**, *178* (4), 901-918.e16. https://doi.org/10.1016/j.cell.2019.07.029.

(71)   Regan, J. C.; Khericha, M.; Dobson, A. J.; Bolukbasi, E.; Rattanavirotkul, N.; Partridge, L. Sex difference in pathology of the ageing gut mediates the greater response of female lifespan to dietary restriction. *eLife* **2016**, *5*, e10956. https://doi.org/10.7554/eLife.10956.

(72)   Wukitch, A. M.; Lawrence, M. M.; Satriale, F. P.; Patel, A.; Ginder, G. M.; Van Beek, E. J.; Gilani, O.; Chambers, M. C. Impact of chronic infection on resistance and tolerance to secondary infection in *Drosophila melanogaster*. *Infect. Immun.* **2023**, e0036022. https://doi.org/10.1128/iai.00360-22.

(73)   Chambers, M. C.; Jacobson, E.; Khalil, S.; Lazzaro, B. P. Consequences of chronic bacterial infection in *Drosophila melanogaster*. *PLoS ONE* **2019**, *14* (10), e0224440. https://doi.org/10.1371/journal.pone.0224440.

(74)   Sibley, C. D.; Duan, K.; Fischer, C.; Parkins, M. D.; Storey, D. G.; Rabin, H. R.; Surette, M. G. Discerning the complexity of community interactions using a *Drosophila* model of polymicrobial infections. *PLOS Pathog.* **2008**, *4* (10), e1000184. https://doi.org/10.1371/journal.ppat.1000184.

(75)   Lin, S. J. H.; Fulzele, A.; Cohen, L. B.; Bennett, E. J.; Wasserman, S. A. Bombardier enables delivery of short-form Bomanins in the *Drosophila* Toll response. *Front. Immunol.* **2019**, *10*, 3040. https://doi.org/10.3389/fimmu.2019.03040.

(76)   Lindsay, S. A.; Lin, S. J. H.; Wasserman, S. A. Short-form Bomanins mediate humoral immunity in *Drosophila*. *J. Innate Immun.* **2018**, *10* (4), 306–314. https://doi.org/10.1159/000489831.

(77)   Martins, R.; Carlos, A. R.; Braza, F.; Thompson, J. A.; Bastos-Amador, P.; Ramos, S.; Soares, M. P. Disease tolerance as an inherent component of immunity. *Annu. Rev. Immunol.* **2019**, *37* (1), 405–437. https://doi.org/10.1146/annurev-immunol-042718-041739.

(78)   Vijendravarma, R. K.; Kraaijeveld, A. R.; Godfray, H. C. J. Experimental evolution shows *Drosophila melanogaster* resistance to a microsporidian pathogen has fitness costs. *Evol. Int. J. Org. Evol.* **2009**, *63* (1), 104–114. https://doi.org/10.1111/j.1558-5646.2008.00516.x.

(79)   Kraaijeveld, A. R.; Godfray, H. C. J. Trade-off between parasitoid resistance and larval competitive ability in *Drosophila melanogaster*. *Nature* **1997**, *389* (6648), 278–280. https://doi.org/10.1038/38483.

(80)   Hanson, M. A.; Lemaitre, B.; Unckless, R. L. Dynamic Evolution of Antimicrobial Peptides Underscores Trade-Offs Between Immunity and Ecological Fitness. *Front. Immunol.* **2019**, *10*, 2620. https://doi.org/10.3389/fimmu.2019.02620.

(81)   Lochmiller, R. L.; Deerenberg, C. Trade-offs in evolutionary immunology: just what is the cost of immunity? *Oikos* **2000**, *88* (1), 87–98. https://doi.org/doi:10.1034/j.1600-0706.2000.880110.x.

(82)   Howick, V. M.; Lazzaro, B. P. The genetic architecture of defence as resistance to and tolerance of bacterial infection in *Drosophila melanogaster*. *Mol. Ecol.* **2017**, *26* (6), 1533–1546. https://doi.org/10.1111/mec.14017.

(83)   Leitão, A. B.; Arunkumar, R.; Day, J. P.; Geldman, E. M.; Morin-Poulard, I.; Crozatier, M.; Jiggins, F. M. Constitutive activation of cellular immunity underlies the evolution of resistance to infection in *Drosophila*. *eLife* **2020**, *9*, e59095. https://doi.org/10.7554/eLife.59095.

(84)   Lhocine, N.; Ribeiro, P. S.; Buchon, N.; Wepf, A.; Wilson, R.; Tenev, T.; Lemaitre, B.; Gstaiger, M.; Meier, P.; Leulier, F. PIMS modulates immune tolerance by negatively regulating *Drosophila* innate immune signaling. *Cell Host Microbe* **2008**, *4* (2), 147–158. https://doi.org/10.1016/j.chom.2008.07.004.

(85)   Bonnay, F.; Cohen-Berros, E.; Hoffmann, M.; Kim, S. Y.; Boulianne, G. L.; Hoffmann, J. A.; Matt, N.; Reichhart, J.-M. Big Bang gene modulates gut immune tolerance in *Drosophila*. *Proc. Natl. Acad. Sci. U. S. A.* **2013**, *110* (8), 2957–2962. https://doi.org/10.1073/pnas.1221910110.

(86)   T.M. Therneau. _A package for survival analysis in R_., 2022. https://CRAN.R-project.org/package=survival.

(87)   John Fox; Sanford Weisberg. An {R} Companion to applied regression, 2019. https://socialsciences.mcmaster.ca/jfox/Books/Companion/.

(88)   R. Lenth. _emmeans: Estimated marginal means, aka Least-squares means_., 2022. https://CRAN.R-project.org/package=emmeans.

(89)   Torsten Hothorn; Frank Bretz; Peter Westfall. Simultaneous inference in general parametric models. *Biom. J.* **2008**, *50* (3), 346–363.

(90)   Achim Zeileis; Christian Kleiber; Simon Jackman. Regression models for count data in R. *J. Stat. Softw.* **2008**, *27* (8).

(91)   Simon Jackman. Pscl: classes and methods for R developed in the political science computational laboratory, 2020. https://github.com/atahk/pscl/.

(92)   Neyen, C.; Bretscher, A. J.; Binggeli, O.; Lemaitre, B. Methods to study *Drosophila* immunity. *Methods* **2014**, *68* (1), 116–128. https://doi.org/10.1016/j.ymeth.2014.02.023.

(93)   R Core Team. R: A language and environment for statistical computing., 2022. https://www.R-project.org/.

(94)   Aphalo P. _ggpmisc: Miscellaneous extensions to 'ggplot2'_, 2022. https://CRAN.R-project.org/package=ggpmisc.

(95)   Livak, K. J.; Schmittgen, T. D. Analysis of relative gene expression data using Real-time Quantitative PCR and the $2-\Delta\Delta CT$ method. *Methods* **2001**, *25* (4), 402–408. https://doi.org/10.1006/meth.2001.1262.

(96)   Micchelli, C. A. Whole-mount immunostaining of the adult *Drosophila* gastrointestinal tract. *Methods* **2014**, *68* (1), 273–279. https://doi.org/10.1016/j.ymeth.2014.03.022.

(97)   Kuznetsova A; Brockhoff PB; Christensen RHB. LmerTest Package: Tests in linear mixed effects models. *J. Stat. Softw.* **2017**, *82* (12), 1–26. https://doi.org/10.18637/jss.v082.i13.

(98)   H. Wickham. Ggplot2: Elegant graphics for data analysis, 2016. https://ggplot2.tidyverse.org.

(99)   RStudio Team. RStudio: Integrated development environment for R, 2020. http://www.rstudio.com/.

(100)   Van Rossum; Guido and Drake; Fred L. Python 3 reference manual, 2009.

(101)   Salvatier J.; Wiecki T.V.; Fonnesbeck C. Probabilistic programming in Python using PyMC3. *PeerJ Comput. Sci. 2 E55* **2016**. https://doi.org/10.7717/peerj-cs.55.

973 **FIGURES AND TABLES**

975 **Figure 1.**

980 **Figure 2.**

984   **Figure 3.**

989   **Figure 4.**

992 **Figure 5.**

996 **Figure 6.**

997   **Table 1.**

parameter	mean	sd	hdi 3%	hdi 97%
Ref. *k*	3.143	0.033	3.080	3.204
Ref. *h*	-2.309	0.144	-2.577	-2.043
Ref. *a*	3.072	0.016	3.043	3.104
Ref. *b*	2.361	0.034	2.298	2.425
Sex *k*	-0.053	0.068	-0.172	0.068
Pop. *k*	-0.108	0.046	-0.193	-0.021
Sex *h*	-0.330	0.227	-0.769	0.089
Pop. *h*	0.042	0.182	-0.301	0.376
Sex *a*	-0.130	0.038	-0.195	-0.061
Pop. *a*	-0.002	0.024	-0.047	0.043
Sex *b*	-0.147	0.066	-0.276	-0.035
Pop. *b*	-0.285	0.053	-0.384	-0.187
ss	3.671	0.062	3.552	3.785

1000

1001

1002

1003

1004

1005

1006

1007

Table 1. Parameter estimates for the model sigmoidal function fitted to the time-course of bacterial counts. A sigmoidal function of 4 parameters (*a*, *k*, *h*, *b*) was fitted to the bacterial dynamics time-course shown in Figure 3A, for quantification of differences between Control and BactOral CFU counts. Control females were used as reference on which to estimate the contribution of tested variables (variable "Sex": males and variable "Population": BactOral) on the parameters. Effects (Sex and Population) on parameters were considered significant when the 94% highest posterior density interval (hdi) did not include zero. The "ss" parameter represents the estimate of the standard deviation of the normal error structure.

1017   **Table 2.**

1018

parameter	mean	sd	hdi 3%	hdi 97%
a_0 **Control**	-1.214	0.570	-2.262	-0.127
a_0 **BactOral**	-0.731	0.570	-1.816	0.324
a_0 **Female**	-0.984	0.569	-2.073	0.075
a_0 **Male**	-0.959	0.569	-2.064	0.079
a_1 **Female**	-0.781	0.071	-0.914	-0.645
a_1 **Male**	-0.061	0.069	-0.188	0.074
a_1 **Control**	-0.700	0.070	-0.833	-0.570
a_1 **BactOral**	-0.144	0.069	-0.268	-0.010
k_0 **Female**	0.086	0.097	-0.096	0.269
k_0 **Male**	0.085	0.095	-0.088	0.268
k_0 **Control**	0.066	0.097	-0.114	0.250
k_0 **BactOral**	0.104	0.095	-0.076	0.281
k_1 **Female**	-0.161	0.103	-0.354	0.030
k_1 **Male**	-0.232	0.103	-0.426	-0.037
k_1 **Control**	-0.161	0.101	-0.355	0.027
k_1 **BactOral**	-0.231	0.105	-0.426	-0.033
t_0 **Female**	0.152	0.119	-0.071	0.372
t_0 **Male**	0.253	0.116	0.031	0.462
t_0 **Control**	0.168	0.119	-0.047	0.397
t_0 **BactOral**	0.238	0.116	0.020	0.453

1019

Table 2. Parameter estimates for the model logistic function fitted to the zero counts of the time-course of bacterial counts. An adapted logistic function of 4 parameters (a_0, a_1, k_0, k_1) was fitted to the progression of zero counts shown in Fig. 3B), for quantification of bacterial clearance differences between Control and BactOral. Single estimates were generated for the contribution of each variable (Sex: females and males; Population: Control and BactOral) to the model parameters. Effects were considered significant when was considered when the mean estimate ± standard deviation did not overlap for the same parameter between each variable, which is the case for a_1 when considering Population and Sex. High-density intervals of 3% and 97% show that all our estimated parameters have a high-density probability (i.e. credibility) as they are all included in them.

1020

1021

1022

1023

FIGURE LEGENDS

Figure 1. Survival upon infection with *P. entomophila*.
Females (top) and males (bottom) of Control (grey) and BactOral (yellow) populations were orally infected with a culture of *P. entomophila* OD_{600}=50 (dashed lines and circles) or a control solution (solid lines and squares) for 24 hours, after which they were flipped onto clean food and their survival measured daily. We identify a significantly higher survival after infection in BactOral (in both sexes) as compared with Control, in a treatment-dependent manner. Results from multiple comparisons are shown as letters in the plots.

Figure 2. Evolved and control populations eat and defecate equivalent amounts of bacteria.
A) CAFE assay. Females of Control and BactOral populations were fed separately *P. entomophila* solution through a graduated glass capillary and the amount of bacterial solution ingested (in μL) per fly was registered at 2, 8 and 24 hours. Cumulative number of μL ingested per fly showed no statistically significant difference in amounts ingested by either population at the different time-points. Linear regressions between averaged values eaten by each population between time-points show a decrease in feeding as infection progresses (equations shown in plot for the periods, 2–8 hours (above lines) and 8–24 hours (below lines). **B)** Defecation assay. Females (top) and males (bottom) from Control (grey) and BactOral (yellow) populations were fed *P. entomophila* for 3 hours, after which the number *P. entomophila* defecated was estimated after 6 hours and 12 hours. Lines connecting the two points provide a sense of individual defecation rate within this time window. Linear regressions between the two counts of the same fly show a significant decrease in the number of bacteria defecated during the initial 6 hours and the following 12 hours by males of BactOral (shown with *). Additionally, there is a significant difference in the number of BactOral flies that defecated below our detection threshold (zero CFU counts) at 6 hours compared to 18 hours and compared to Control for both timepoints.

Figure 3. Bacterial load dynamics. Individual flies from Control (grey) and BactOral (yellow) were collected upon exposure to *P. entomophila* at different timepoints for assessment of the number of Colony Forming Units (CFUs) inside their bodies. After the end of the infection period (24 hours), flies were changed onto clean food and collection continued until 52 hours post-infection. **A)** Log_2 of the number of CFUs found for females (top) and males (bottom) of either Control (grey) or BactOral (yellow) populations, with the average value at each time-point evidenced in red (triangles for Control and circles for BactOral). There are no detectable

1059 differences in bacterial loads between both populations until 20 hours of infection, when BactOral

1060 starts to evidence a bigger decrease in bacterial counts. **B)** Cumulative number of individual flies

1061 from which no CFUs were obtained increases as time progresses, with the BactOral reaching

1062 significantly higher final values.

1063

Figure 4. AMP expression differs between populations.

Guts from Control (grey) and BactOral (yellow) female flies were dissected before exposure

("Unc.") and at different timepoints after oral infection (8, 24 and 32 hours) with *P. entomophila.*

Relative gene expression, normalized to housekeeping gene *EIF2,* was performed for a

representative panel of 10 AMPs. We detected a significantly higher up-regulation in BactOral

compared to Control for *AttacinA* and *CecropinA* at 8-hours of infection. Results from these

comparisons are shown in the letters above the plots.

1071

1072

Figure 5. Control flies primed with heat-killed bacteria phenocopy BactOral survival to oral

infection with *P. entomophila*. Control (grey) and BactOral (yellow) females (left) and males

(right) were fed a solution of heat-killed bacteria for 2 hours before oral infection (dashed lines)

or directly infected with live *P. entomophila* (full lines) for 24 hours, after which they were

changed onto clean food and their survival measured daily. Previous exposure to heat-killed

bacteria before infection increased survival in both populations, leading additionally Control

females to phenocopy BactOral survival levels upon infection. Results from multiple comparisons

are shown as letters in the plots.

1081

1082

Figure 6. BactOral displays higher survival than Control upon immune system activation

without bacterial proliferation. Daily survival was measured for Control (grey) and BactOral

(yellow) populations fed a mixture of food with heat-killed bacteria or PBS continuously BactOral

females and males exhibit a slower mortality compared to Control flies until around day 10. This

difference fades away for both sexes after 14 days. Survival comparisons between populations

and treatments are shown as lower-case letters.

1089

1092

Supplementary Figure 1.

No detectable differences in gut structure between Control and BactOral after oral infection

with *P. entomophila*. Females of Control (grey) and BactOral (yellow) were fed normal food

("Without infection") or orally infected with *P. entomophila* for 24 hours, after which their **A)**

guts or **B)** crops were dissected and photographed. Gut length (in mm) was measured by drawing

consecutive longitudinal transects along the guts thrice and averaging those measurements to

reach final length per sample. Crop perimeter was measured in a similar way, by drawing

segments around the border of each crop. **A)** Exposure to bacteria shortens the guts significantly

in both populations (Control before infection *versus* Control 24 hours post-infection: *t.ratio* =

6.274, *p* <0.0001; BactOral before infection *versus* BactOral 24 hours post-infection: *t.ratio* =

5.453, *p* <0.0001). No differences were detected in gut lengths between populations within each

timepoint (Without infection: BactOral *versus* Control – *p* = 0.6719; after 24 hours of infection –

p = 0.9925), although a significant effect of treatment was observed in both Control (Without

infection *versus* after 24 hours of infection – *p* < 0.0001) and BactOral (*p* < 0.0001), showing that

gut shortening due to infection affected both regimes in a similar manner. **B)** Dissected guts of

females from Control and BactOral populations were photographed and scaled for posterior

length quantification. Manual transects were drawn longitudinally along the guts (as shown

above) and total size of the cumulative transects was calculated in FIJI. Three independent

measurements were done on each gut and averaged to obtain a final value per sample (Fig. 2 A)).

1112

1113

Supplementary Figure 2.

AMP expression differences between populations under homeostatic conditions. Guts from

Control (grey) and BactOral (yellow) female flies were dissected at different timepoints after oral

infection with *P. entomophila* and relative quantification of gene expression was performed.

Expression levels of each gene in BactOral samples were normalized to expression levels of

Control samples. Log$_2$(FoldChange) of these differences were fitted on a linear model with

Population as factor and followed by multiple comparisons (*emmeans*) to uncover pairwise

differences. Results from these comparisons are shown in the letters above the plots. There are

significantly lower levels of CecropinA (CecA) (*t.ratio*= 2.892, *p* = 0.0445) and significantly

higher levels of *Drosomycin-like 5* (Drsl5) (*t.ratio*= -4.686 , *p* = 0.0094) being expressed in guts

of BactOral flies under unchallenged conditions.

1125

1126

1127

1128

Supplementary Figure 3.

Principle Component Analysis on relative gene expression levels in evolved populations. (A)

Relative AMP expression levels of Control and BactOral flies upon oral infection with *P.*

entomophila showed significant differences at various timepoints (see Fig. 4 and Supp. Fig. 2).

(B-C) Relative influence of each AMP to the establishment of PC1 and PC2 (see Supp. fig. 3 A).

Importantly, Cecropin A1 represents the largest contributor to PC1 (**B**) whereas Drosomycin

represents the largest contributor, although negatively correlated, to the variation in PC2 (**C**).

1136

1137

1138

1139

Supplementary Figure 4.

Quantification of immunity-related gene expression upon artificial immune stimulation and

infection. To test for individual pairwise differences, relative log2 (FoldChange) expression

levels of each gene were fitted onto a linear model with Population and Timepoints as factors and

followed by multiple comparisons (*emmeans*). No differences are observed between non-

challenged individuals or between individuals after 2 hours of exposure. However, for Diptericin

Unchallenged versus 2 HK show differences in both populations (Control: *t.ratio*= -4.778 , *p*=

0.0001; BactOral: *t.ratio*= -3.122, *p*= 0.0052).

1148