TITLE:

Evolution of resistance and disease tolerance mechanisms to oral

bacterial infection in *D. melanogaster*

AUTHORS:

Tânia F. Paulo[1*#], Priscilla A. Akyaw[1,3,4], Tiago Paixão[1], Élio Sucena[1,2,3,4*]

1 – Instituto Gulbenkian de Ciência, Rua da quinta grande 6, 2780-156 Oeiras, Portugal

2 – Departamento de Biologia Animal, Faculdade de Ciências, Universidade de Lisboa, Edifício C2, Campo Grande, 1749-016 Lisbon, Portugal

3- cE3c: Centre for Ecology, Evolution and Environmental Changes, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1749-016 Lisbon, Portugal

4- CHANGE – Global Change and Sustainability Institute

* TFP: tania.paulo@childrens.harvard.edu; ES: jesucena@fc.ul.pt

present adress: Boston Children´s Hospital, Harvard Medical School, Harvard, USA.

KEYWORDS:

Experimental Evolution, Innate Immunity, *Drosophila*, *Pseudomonas entomophila*

ABSTRACT

Pathogens exert strong selection on hosts, that evolve and deploy different defensive strategies, namely minimizing pathogen exposure (avoidance), directly promoting pathogen elimination (resistance), and/or managing the deleterious effects of illness (disease tolerance). However, how the response to pathogens partitions across these processes has never been directly assessed in a single system, let alone in the context of known adaptive trajectories under controlled selection regimes. Here, an experimental evolution system composed of *D. melanogaster* and its natural pathogen *P. entomophila* is used to independently assess the role of behavioural traits, and of resistance and disease tolerance mechanisms on host evolution. We compare one replicate of a population adapted to oral infection with *P. entomophila* (BactOral) to a replicate of its control population to find no evidence for behavioural change but measurable differences in both resistance and disease tolerance. In BactOral, we identify a relative decrease in bacterial loads correlated with an increase in gut production of specific AMPs, but no differences in bacterial intake, in gut cell renewal rate, or in the rate of bacterial defecation, pointing to a strengthening in resistance. Additionally, we posit that disease tolerance also contributes to the adaptive response of the BactOral population through a tighter control of its immune response and of the deleterious effects of exposure. This study reveals a genetically complex and mechanistically multi-layered response, possibly reflecting the structure of adaptation to infection in natural populations.

BACKGROUND

Host-parasite interactions are main drivers of evolution[1,2]. In natural environments, parasites are constantly attempting to penetrate host defences, which consist of multiple mechanisms to fight infections[3]. Such mechanisms include behavioural strategies (e.g. avoidance)[4], the active reduction or elimination of pathogens (i.e. resistance) and/or the maintenance of homeostasis and fitness without interfering with pathogen burden (i.e. disease tolerance)[5–8]. While many studies in the past decades have explored the mechanistic and genetic bases of these different host defence strategies in diverse organisms [9–11], few have focused on how these processes can be shaped by evolution[12–15]. Experimental evolution is a powerful method by which it is possible to infer the evolutionary response of a host species to a specific pathogenic pressure under controlled conditions[16]. Previous studies of experimental evolution have addressed the impact of different hosts on parasite fitness[17], of different pathogens on the host's evolutionary trajectory[13,18,19], of immune priming[20,21], and of different routes of infection on host adaptation[19,22].

It is postulated that oral exposure to pathogens constitutes a common process of infection in *D. melanogaster*[23,24], as it lives on decaying substrates. When feeding on pathogenic microorganisms, *Drosophila* deploys different responses that include avoidance behaviours[25,26], the barrier action of the peritrophic matrix[27], localized high acidity of the midgut[28], intensification of gut transit[29], local production of Anti-Microbial Peptides (AMPs)[30] and Reactive Oxygen Species (ROS) by gut epithelial cells[31,32] and increase of enterocyte delamination and renewal rates[33–36]. In addition, *Drosophila* can activate a systemic humoral response, when the pathogen manages to cross the gut epithelium into the haemolymph[37,38] or otherwise provokes a systemic reaction[23,39–41]. In the specific case of oral infection with the natural entomopathogen *Pseudomonas entomophila*, its high pathogenicity to *D. melanogaster* is linked to the production of toxins and virulence factors[42,43] causing translational arrest in host tissues and the inability to renew damaged gut epithelium[44,45]. The

above-presented plethora of possible host responses provides a clear picture of the multi-

layered and mechanistically complex nature of the immune response, understood in its broad

sense[3].

In this work, we used a previously experimentally evolved outbred population of *D.*

melanogaster (hereafter called "BactOral") showing a fast evolutionary response to a strong

selection against oral *P. entomophila* infection[19,46], to investigate the mechanisms targeted by

adaptation. We systematically tested for behaviour, resistance, and disease tolerance

contributions, to provide a comprehensive account of the partitioning of these defense

strategies in adaptation.

82   RESULTS

83   Evolved population shows higher survival upon infection

In previous work[19,46], experimental evolution was conducted on an outbred population

of *D. melanogaster* against oral infection either with the bacterium *P. entomophila* (hereafter,

BactOral) or with a control solution (hereafter, Control). Since then, these populations[46] have

been kept under relaxed selection for over 80 generations. To understand to which extent this

relaxation affected the response to the original selection regime, we measured survival of

Control and BactOral populations upon *P. entomophila* oral infection. We found significantly

higher survival of BactOral for both sexes (Fig. 1, Infected Control females *versus* Infected

BactOral females: *z.ratio*= 9.123, *p*< 0.0001; Infected Control males *versus* Infected BactOral

males: *z.ratio*= 9.909, *p*< 0.0001)(Suppl. Table 1). Moreover, these differences are not due to

an inherent frailty of the Control as compared to BactOral. Indeed, survival under uninfected

conditions for both long term longevity experiments (Suppl. Fig. 1 and Suppl. Table 2) and

over the equivalent time period shows similar trends in both populations (Fig. 1, Uninfected

Control females *versus* Uninfected BactOral females: *z.ratio*= 0.004, *p*= 1; Uninfected Control

males *versus* Uninfected BactOral males: *z.ratio*= -1.080, *p*= 0.9611)(Suppl. Table 1).

Taking these results into account, we undertook a systematic description of multiple

traits previously associated to immune response to oral infections aiming to identify different

processes targeted by selection and estimate their relative contribution.

Evolved population has not changed bacterial uptake or defecation rates

Our systematic approach started by screening behavioural traits that impact bacterial

numbers in the gut: feeding and defecation. To test for differences in feeding behaviour, we

performed a Capillary Feeding assay (CAFE)[47] on females of both Control and BactOral

populations at different timepoints over 24 hours of exposure to bacterial suspension. We

allowed individual flies to feed on a *P. entomophila* suspension mixed with food coloring in

graduated glass capillaries and measured the amounts eaten by each female at 2, 8 and 24

hours of exposure. We found no significant differences between the amount ingested by

individuals of the two populations (power analysis for a t-test: *Cohen's d*= 1,06). Pairwise

comparisons of their cumulative bacterial intake after 2 hours, 8 hours and 24 hours exhibited

a comparable amount of total bacteria ingested (Suppl. Table 3 and Fig. 2A, Control *versus*

BactOral at 2 hours, *t*= 0.097, *p*= 1; at 8 hours, *t*= 0.643, *p*= 0.9873; at 24 hours, *t*= 0.574, *p*=

0.9925).

Additionally, we estimated the average slopes from linear regression where microliters

of bacteria eaten were modelled between different time points, concluding that the bacterial

quantities ingested by both populations remain undistinguishable as the infection progresses.

Thus, exposure affects feeding ability of both populations to a comparable degree and,

consequently, adaptation to oral infection with *P. entomophila* in the BactOral regime did not

involve an alteration of feeding behaviour.

Considering the absence of significant differences in bacterial intake, an alternative

would be that the BactOral population had evolved a more effective bacterial defecation, which

would reduce the time of pathogen permanence and proliferation in the gut. We established a

protocol to measure live bacteria loads in the faeces, hence considering the number of

bacteria defecated as a proxy for defecation rate itself, assuming that no interaction between

the two parameters exists. To this purpose, we took individual flies upon oral infection and

quantified the CFUs defecated over a six-hour and the subsequent twelve hour-periods, post-

exposure (Fig. 2B).

We observed a few instances where flies did not expel alive bacteria during that period

or did so in a quantity that fell below our detection threshold, resulting in a few counts of zero.

Therefore, a zero-inflated model with a negative binomial distribution was fitted on the counts

data (accounting for overdispersion) jointly with a binomial distribution on the zeros. For the

counts portion of the model, none of the three factors significantly predicted the measured

bacterial loads (Suppl. Table 4, model parameters - Population: *z.value*= 1.324, *p*= 0.185;

Sex: *z.value*= 0.073, *p*= 0.942; Timepoint: *z.value*= -0.359, *p*= 0.720). However, looking at the

zero portion of the model, we find a significant effect of Population (Population: *z.value*= -

2.407, *p*= 0.016; Timepoint: *z.value*= -0.006, *p*= 0.996).

We then performed post-hoc pairwise comparisons on these regressions and identified

a significant decrease in bacterial loads defecated by males of BactOral, between the two

timepoints (males: *z.ratio*= -2.719, *p*= 0.033), but no other significant effects among bacterial

counts between populations and/or timepoints, in a sex-dependent manner. However, when

considering the zeros, we found a significant difference in the proportion of BactOral flies with

undetectable amounts of bacteria in the faeces between the first and second timepoints (B6h–

B18h: *z.ratio*= 4.092, *p*= 0.0002), as well as compared to Control at both timepoints (B6h–

C6h: *z.ratio*= 3.612, *p*= 0.002; B6h– C18h: *z.ratio*= 3.530, *p*= 0.002)(Suppl. Table 4).

These results demonstrate that both Control and BactOral populations defecate alive

bacteria in comparable amounts upon infection. Nonetheless, we detected that males of

BactOral decrease the amount of live bacteria they defecate as time progresses, hinting at a

possible resistance mechanism more evident in males of the adapted lines, that was not

significant in females. This observation is corroborated by the difference in the number of

"zeros" found between both populations, with BactOral exhibiting more instances of expelling

undetectable bacterial loads.

Evolved population shows a faster decrease and clearance of bacteria

Next, we studied the physiological responses to estimate the roles of resistance and

disease tolerance. First, we tested for structural changes in the gut itself that could influence

the "carrying capacity" of the structure and create difficulties in interpreting the quantification

experiments. We measured the capacity of BactOral to regenerate its gut epithelium upon

bacterial infection relative to Control, using initial length and extent of the shortening induced

by infection[36], as a proxy. For this, we measured the total length of the guts of females, before

and after 24 hours of oral infection with *P. entomophila* and performed multiple pairwise

comparisons on a linear model with mixed effects to pinpoint how population and infection

status affected this trait. We found a shortening of approximately 2 mm in the total length of

infected guts by comparing the post-infection timepoint of 24 hours to the initial timepoint, prior

to exposure, in both populations (Suppl. Table 5 and Suppl. Fig. 2, Control before infection

versus Control 24 hours post-infection: $t= 6.274$, $p<0.0001$; BactOral before infection *versus*

BactOral 24 hours post-infection: $t = 5.453$, $p <0.0001$). Nonetheless, no significant differences

between populations in the total length of the guts could be detected both before and after

infection (Without infection: BactOral *versus* Control – $p= 0.6719$; after 24 hours of infection –

$p= 0.9925$), (power analysis for a multiple linear regression: *Cohen's d* for Population= 0,71;

d for Treatment= 0,71) (Suppl. Fig. 2).

To explore the dynamics of bacterial proliferation inside the host, we performed a time-

course characterization of bacterial loads during and after oral infection by counting the

number of colony-forming units (CFUs) obtained from individual flies. In Figure 3, we show the

progression of bacterial loads from 1 hour until 52 hours post-infection (Fig. 3A) as well as the

cumulative number of individuals without detectable live bacteria (Fig. 3B). The infection

dynamics of both populations start with very high numbers of *P. entomophila* at early stages,

confirming the absence of differences in initial inoculum reported above (Table 1, parameter

a: 3.072± 0.016). Bacterial loads continuously decrease and, at approximately 20 hours of

infection, Control and BactOral population begin to differ significantly in a sex-dependent

manner. From this point onwards we observe a steady decrease of bacterial quantities in the

two populations, always more severe in males than in females, and in BactOral than in Control

flies (Fig. 3A). In addition, the number of flies that clear the infection also increases with time

in both populations, but at a faster rate and reaching higher levels in BactOral (Fig. 3B). By 52

hours post-infection, in both males and females, the number of "cleared" flies in the BactOral

population was twice as high as that of its Control counterpart (Fig. 3B). Together, these

analyses indicate that the evolved population eliminates bacteria faster and more efficiently

than Control.

In order to pinpoint more accurately the parameters contributing to this overall

difference, we modelled the bacterial load and zero dynamics, and assessed differences in

parameter estimates considering the initial load or inoculum (*a*), the load in the long-term or

set-point bacterial load (*b*), the timepoint at which the load is reduced in half (*k*) and the highest

rate of decrease or slope at timepoint *k* (*h*) (see Materials and Methods for details). In Table

1, we show the mean estimates, standard deviation, and high-density intervals (3% and 97%)

for all four parameters of the fitted sigmoidal model, for Control and BactOral populations.

Overall, these results show that sex has a significant effect on all curve parameters (*k*= -0.053

± 0.068; *h*= -0.330 ± 0.227; *a*= -0.130 ± 0.038; *b* = -0.147 ± 0.066), reinforcing the notion of a

strong sexual dimorphism in response to infections[48–50]. Additionally, populations differ on

"time to half height" (*k*= -0.108 ± 0.046) and final load (*b*= -0.285 ± 0.053), but not on initial

load (*a*= -0.002 ± 0.024) nor on highest rate of decrease (*h*= 0.042 ± 0.182).

Because this time-course experiment yielded a significant amount of zero counts,

corresponding to individual flies for which no CFUs could be counted, we treated this as an

independent dataset for which we developed a separate model focused on infection clearance

(Table 2). With that, we found that a_1 (final fraction of zeros) was significantly affected by both

sex and population (with no observable overlap between the mean estimates for each

condition) (a_1 Females= -0.781 ± 0.07; a_1 Males= -0.061 ± 0.069; a_1 Control= -0.700 ± 0.070;

a_1 BactOral= -0.144 ± 0.069), with more BactOral individuals having cleared the infection by

the end of the experiment, and males more so than females. There were significant overlaps

between the estimates of the remaining parameters, evidencing a difference in clearance rate

only in late stages of bacterial infection. In sum, these data align with those of bacterial load

dynamics, whereby flies from BactOral population clear *P. entomophila* faster and better from

20h of exposure onwards.

These results support the hypothesis that resistance mechanisms participate in the

adapted response of BactOral. Indeed, bacterial numbers are affected differently between

BactOral and Control populations despite the equivalent initial bacterial inocula (Fig. 2A, Table

1) and rates of live bacteria defecation (Fig. 2B). The differences in number of detected live

bacteria inside flies suggests a direct bacterial control mechanism is operating more efficiently

in BactOral individuals.

Differences in AMP expression are consistent with distinct bacterial loads dynamics

In an attempt to pinpoint which resistance mechanisms contribute to the differences in

bacterial clearance rates observed between populations, we infected female flies from Control

and BactOral and dissected their guts prior to exposure, and after 8, 24, and 32 hours of

infection with *P. entomophila*. We quantified expression levels for a panel of 10 AMPs covering

different molecular families, dedicated pathways and pathogen specificities, including *Attacin-*

A, *Bomanins Short 2* and *3*, *Cecropin-A1*, *Defensin*, *Diptericin-A*, *Drosomycin* and

Drosomycin-like *2*, *3* and *5* (Fig. 4, Suppl. Tables 6-7, and Supp. Fig. 3).

We first verified if the baseline levels of AMPs (i.e., expressed under homeostatic

conditions) differed between populations, by relativizing expression levels to the Control

population at the unchallenged timepoint. With this approach, we established that BactOral

and Control populations express comparable amounts of 8 AMPs when unchallenged, the two

exceptions being *Cecropin-A1* (*CecA*) with lower expression levels ($t = 2.892$, $p = 0.0445$) and

Drosomycin-like 5 (*Drsl5*) with higher levels ($t = -4.686$, $p = 0.0094$) (Supp. Fig. 2) in the

Control compared to the BactOral population.

In addition, and since our main goal was to characterize the temporal dynamics of AMP

expression in both populations, we performed the analysis for the entirety of the time-course

using the unchallenged timepoint as reference (power analysis for a t-test: *Cohen's d*= 0,62)

(Fig. 4 and Suppl. Tables 6-7). Overall, we detect consistent dynamics with the reduction of

expression happening sooner (by 32 hours) in BactOral than Control across 7 of the 10 AMPs

tested, the exceptions being *Attacin A*, *Defensin* and *Drosomycin-like 2* (*Drsl2*) (Fig. 4). This

earlier down-regulation in BactOral relative to Control flies (at 32-hours) of *BomaninS2*

(*BomS2*) ($t = 4.347$, $p = 0.0047$), *BomaninS3* (*BomS3*) ($t = 4.035$, $p = 0.0089$), *CecA* ($t = 4.529$,

$p = 0.0010$), *Diptericin* ($t = 2.613$, $p = 0.0439$), *Drosomycin* ($t = 4.795$, $p = 0.0056$), *Drosomycin-*

like 3 (*Drsl3*) ($t = 3.194$, $p = 0.0105$) and *Drsl5* ($t = 4.266$, $p = 0.0021$). Furthermore, we detected

a significantly higher up-regulation of *AttacinA* and *CecA* in BactOral at 8 hours of infection

(*AttacinA*: Control 8h *versus* BactOral 8h – $t = -2.563$, $p = 0.0389$; *CecA*: Control 8h *versus*

BactOral 8h – $t = - 3.034$, $p = 0.0123$), consistent with the role of these peptides as main players

against infections with Gram-negative bacteria[51,52]. A similar trend was observed for other

tested AMPs such as *Defensin* and *Drsl2*, although these changes were not statistically

significant (*Defensin*: Control 8h *versus* BactOral 8h – $t = -2.101$, $p = 0.231$; *Drsl2*: Control 8h

versus BactOral 8 h.p.i. – $t = -0.899$, $p = 0.465$).

Principal Component Analysis (PCA) supports these trends (Suppl. Fig. 4). The two

main Principal Components (PC1 and PC2) evidence the time-dependent differences

observed between populations by showing an initial indistinguishable grouping of both,

followed by a dissimilar change in expression patterns at 8 hours of infection which culminates

with a maximum divergence between Control and BactOral at the later timepoint (32 hours).

Increase of AMP levels in the Control population phenocopies

BactOral response to infection

Having correlated the improved bacterial clearance of BactOral flies (Fig. 3) with

significant differences in expression levels of AMPs between populations (Fig. 4), we sought

to functionally associate these two observations. For that, we conducted a priming experiment

to induce higher basal levels of AMPs by feeding flies a solution of heat-killed bacteria. These

primed flies were, subsequently, infected orally with live *P. entomophila*, and followed daily for

mortality. This exposure to dead bacteria has been shown to "artificially" increase AMP

levels[53–55]. We expect the priming with heat-killed bacteria to increase the basal expression of

AMPs and thus provoke, particularly in Control flies, an up-regulation of the immune system,

allowing test of the influence of this heightened condition in survival to oral infection.

Flies previously fed with heat-killed bacteria increased survival to infection with live *P.*

entomophila in BactOral and Control populations, confirming the stronger immune response

expected upon priming (Suppl. Table 8 and Fig. 5, BactOral females fed with live bacteria

(control) *versus* BactOral females primed with heat-killed bacteria (primed): *z.ratio*= 7.193, *p*<

0.0001; BactOral males treated with control *versus* BactOral males primed with Heat-killed

bacteria: *z.ratio*= 4.368, *p*= 0.0001; Control females fed with control *versus* Control females :

z.ratio= 10.011 , *p*< 0.0001; Control males treated with control *versus* Control males primed :

z.ratio= 7.299 , *p*< 0.0001). Females from the Control population primed with heat-killed

bacteria increased their survival to oral infection, virtually phenocopying BactOral infected with

live bacteria, whereas in males, the same tendency is observed but to a lesser degree (Suppl.

Table 8 and Fig. 5, BactOral females fed with control *versus* Control females primed: *z.ratio*=

1.629, *p*= 0.3620 and BactOral males fed with control *versus* Control males primed: *z.ratio*= -

4.547, *p*<0001). This effect is supported by a time-course of relative expression levels of IMD

pathway key players, *Relish* (its devoted transcription factor) and its associated AMP,

Diptericin, as well as the Toll-activated AMP, *Drosomycin* (Suppl. Fig. 5). For both populations,

we observed an increase in levels of *Diptericin* to a comparable degree upon exposure to

heat-killed *P. entomophila*, but not of *Drosomycin*.

With this experiment we strengthened the case for a role of improved resistance in the

adapted phenotype of the BactOral population.

BactOral maintains higher survival upon immune response

activation

To ascertain whether disease tolerance may also contribute to BactOral increased

survival upon oral infection, we continuously fed flies with a mixture of heat-killed *P.*

entomophila and followed survival (Fig. 6).

With this setup we verified that both females and males of BactOral survive significantly

more than Control after the specific damages induced by constant exposure to heat-killed

bacteria. Importantly, the survival differences detected between populations occur up until the

time at which flies were selected to reproduce during the experimental evolution protocol (10

days upon infection), evidencing the adaptive nature of this response. (Suppl. Table 9 and Fig

6, Control - HK *versus* BactOral - HK females: *z.ratio*= 6.579, *p*< 0.0001; males: *z.ratio*=

9.612, *p*< 0.0001). No significant differences exist under control conditions (Control - PBS

versus BactOral - PBS females: *z.ratio*= 1.392, *p*= 0.504; males: *z.ratio*= 0.587, *p*= 0.936).

DISCUSSION

In this work, we enquired systematically for the bases of adaptation of one evolved line

of *D. melanogaster*, against an oral bacterial infection considering the three general layers

that constitute the immune response: behaviour, resistance and disease tolerance[3,8,56]. Upon

experimental evolution of an outbred population of *D. melanogaster* against oral infection with

P. entomophila, and after approximately 6 generations, the selected population (BactOral)

increased survival upon infection[19]. This phenotype was maintained throughout the selection

experiment but, since then, the population has been kept for more than 80 generations under

relaxed selection. Despite the fact that no physiological costs associated to this response

could be detected in any of the original four replicates[46], we tested the immunocompetence of

one selected representative replicate per evolved population. The average survival rate after

oral infection of flies from BactOral was approximately 63% (50% for females and 75% for

males) and approximately 10% in the Control population (5% for females and 15% for males).

Considering that at the end of selection, BactOral survival reached 90 %[19], this partial

decrease in overall survival after infection with bacteria can potentially underestimate the

contributions of each of the different mechanisms behind adaptation. Importantly, these losses

could be attributed to genetic drift acting on the populations during the long relaxed-selection

period they sustained. Notwithstanding, we established that it is methodologically possible to

ascertain the relative importance of behavioural, resistance, and disease tolerance

mechanisms.

No role for behavioural or structural differences

It is well described that the route of infection is of paramount importance to the outcome

of host-parasite interactions. In *D. melanogaster*, the mode of entry of a bacterial pathogen

into the host's body elicits different physiological responses[19,57–61], which can start by the

avoidance of the pathogenic substrate, in the case of infection through oral route[26,62,63], both

by inhibiting feeding or choosing oviposition sites according to bacterial composition[26,53,62–64].

P. entomophila is a natural pathogen of Drosophila that causes high mortality when ingested

across diverse genetic backgrounds and conditions[23,43,44,65,66]. Thus, it is plausible to consider

that fasting could constitute a general response against oral infection. The simplest behaviour

that could explain the observed differences in survival between populations would be that

BactOral flies evolved a reduced intake during exposure to bacteria, thereby reducing their

"inoculum" size. Indeed, fasting is a prevalent response to infection and behavioural avoidance

is a common strategy to limit exposure to infected substrates in different systems[4,62,63].

However, we showed similar amounts of bacteria were ingested by females of both

populations (Figure 2A)[23,67]. Moreover, we consubstantiate the findings from the CAFE assay

with the bacterial loads time-course, that show equivalent amounts of *P. entomophila* colonies

inside infected flies at early timepoints. Our results indicate that behavioural inhibition of

bacterial intake was not one of the mechanisms that evolved in BactOral flies, eliminating

fasting as a component of the adaptation to *P. entomophila* exposure.

Previous reports have shown that pathogen intake can affect gut function and

structure[34–36,45] as well as influence the dynamics of exposure by altering defecation rates[29].

However, we could not find evidence for any differences in these features between control

and evolved populations. Firstly, there are no changes in gut length in steady-state conditions

that would somehow enable the BactOral population to *a priori* face bacterial infection more

favourably. Increased enteric cell renewal, which often leads to a temporary shortening of gut

length[36], has been described as one of the main mechanisms of *D. melanogaster* defense

against oral bacterial infection[35,36,45]. In our experiments, we detected a significant decrease

in total gut length after 24 hours of oral infection, but this change was equivalent in both

populations, not supporting a role for altered renewal rate in BactOral adaptation.

These experiments, however, do not completely discard the hypothesis of differential

gut structure underlying the adapted phenotype of BactOral. For example, proportions of gut

cell types can change upon infection[68] and such qualitative shift could contribute to higher

fitness upon exposure to *P. entomophila* in the adapted population. Also, it could be that

relevant changes in gut length occur at later timepoints. Notwithstanding, it has been

previously shown that infection-induced gut cell loss starts as early as 2 hours after exposure,

with a return to 75% of the total original gut length after 24 hours[36]. This time window is well

within the boundaries defined in our setup. Another important aspect to note is that renewal

capacity is sexually dimorphic under diverse physiological contexts[69,70]. For instance, females

have higher gut cell renewal rates than males upon oral infection with *P. carotovorum* 15 [71], a

bacterium that provokes higher mortality in males than females, contrary to oral infection with

P. entomophila (see Results). The disparity between those and our observations may be

explained by the different mechanisms of virulence employed by these two pathogens to infect

D. melanogaster[45]. Be as it may, we envision further experiments testing for potential

differences in gut delamination rates between evolved populations at distinct timepoints.

BactOral flies also did not expel pathogenic bacteria from their guts differently from

Control. However, males of BactOral show a decrease in the number of alive bacteria

defecated between 6- and 18-hours post-exposure, evidencing, to some extent, improved

bacterial clearance and suggesting a role for resistance. Since our protocol only detects live

bacteria, we could potentially be missing an important component of the response and future

work should quantify defecation of alive and dead bacteria, for example through fluorescent

microscopy, to compare populations with regards to the contribution of overall peristalsis to

the elimination of total gut content[29]. In addition, considering the slightly higher number of of

BactOral flies that defecated alive bacteria below our detection limit (zeros), we hypothesize

that this population either retains *P. entomophila* for longer inside guts before they begin to

expel it (which could point to higher disease tolerance), or reduce bacterial loads more

effectively at earlier stages, even though, with time, bacterial proliferation recovers (again

suggesting stronger resistance).

A role for resistance

To further test the suggested role for resistance, we characterized the bacterial load

dynamics in evolved populations over 52 hours and determined that BactOral flies eliminate

pathogens in a more efficient way than Control. This significant change in bacterial loads

indicates that BactOral has evolved higher resistance, that control earlier and clear better gut

bacterial loads. As this model was applied to the logarithm of mean bacterial counts at each

timepoint, it is possible that information regarding intra-populational variation is not considered

in the analysis. Nevertheless, the conclusions from the comparison of the bacterial counts are

further supported by the separate analysis on the number of "zeros" as a function of time,

which is a proxy for bacterial clearance.

Even though there is no discernible difference in the amount of CFUs infecting both

males and females between both populations until approximately 20h of infection, by this time

individuals from BactOral begin to exhibit gradually decreasing levels of bacteria. This trend

aligns with the higher clearance rate observed in the adapted population (and even higher in

males than females) as measured by the increase in the number of flies from which we cannot

grow colonies of *P. entomophila* ("zeros"). Flies from the Control population control their

bacterial loads (at around 20h of infection) but do it at a slower pace and in a less effective

way (i.e. many individuals exhibited high bacterial counts by the end of the experiment). It is

possible that our proxy for clearance (i.e., flies classified as "zeros"), could be overestimated

due to a low detection threshold. Indeed, it has been shown that *D. melanogaster* can endure

chronic bacterial infections[72–74], albeit at rather low levels, which can be the case in our

experimental setup. Nonetheless, this holds true for both populations, therefore not

invalidating our interpretation of a different bacterial clearance rate between genotypes.

Having shown that BactOral clears *P. entomophila* infection more rapidly, we

hypothesized that this response could rely on an increased production of immune effectors,

such as AMPs. Analysis of AMP gut gene expression showed that in unchallenged condition,

only two AMPs (*CecA1* and *Drsl5*) differed in expression between populations (down- and up-

regulation respectively), indicating that the base levels of AMP expression are unlikely to have

been a major target of selection. The detected expression difference of *Drsl5*, a gut-specific

AMP peptide, could represent a pre-emptive resistance mechanism acquired by BactOral, by

establishing a more antagonistic environment for *P. entomophila* proliferation in the guts of

these flies after ingestion.

At 8 hours of exposure, comparing BactOral to Control, two AMPs show a significantly

faster up-regulation while seven of the panel members display a pattern of faster

downregulation, later on (32h). These results suggest that a two-step process might be taking

place, whereby the response is initially (8h) more aggressive, permitting a more effective

bacterial control, and is turned off sooner (32h), limiting the secondary effects of prolonged

immune responses (immunopathology). Interestingly, *Drosomycin-like* genes (*Drsl2*, *Drsl3*,

and *Drsl5*), whose expression is specific to the gut epithelium and are regulated by JAK-

STAT[35], show up-regulation after 8-hours of infection in both populations. For all three genes,

there is a slight tendency for a faster response that could represent a more effective bacterial

control in the BactOral population and a faster down-regulation at the 32-hour timepoint that

could limit self-inflicted damage. Likewise, we observed a similar process of down-regulation

of other AMPs specifically deployed to fight off infections with Gram-negative pathogens (e.g.

CecA1 and *Diptericin*) at the 32-hour timepoint. In contrast, some AMPs typically associated

to response to Gram-positive bacteria, (e.g. *BomS2, BomS3* and *Drosomycin*[51,75,76]), do not

exhibit significant up-regulation after exposure to *P. entomophila* until 32-hours post-infection

and only in the Control population.

Strikingly, and even though *Diptericin* has been shown to be deployed by *D.*

melanogaster to resist *P. entomophila* infection[67], we did not detect a difference in the relative

expression of this gene between Control and BactOral until after 24 hours of infection. At this

timepoint, adapted flies are already returning to basal levels of AMP production whereas

Control flies keep actively fighting off the infection.

The same pattern is observed also in the priming experiment (Fig. 5 and Suppl. Fig.

5). After priming with heat-killed bacteria and subsequent exposure to live bacteria, both

populations maintain an equivalent expression increase until 8 hours but from 24 hours

onwards BactOral shows, again, an earlier downregulation of *Diptericin* and a quicker return

to homeostatic levels of expression (Suppl. Fig. 5). In brief, the dynamics of *Diptericin*

expression mimic the expected effect of priming by establishing higher basal levels of this

AMP with which to counteract an incoming infection. This is not seen for the AMP *Drosomycin*,

which again is consistent with our previous result that this AMP is not deployed against

infection with *P. entomophila* (Fig. 4 and Suppl. Fig. 3).

With this we re-enforce the notion that the process of earlier AMP down-regulation

could be an active response rather than a consequence of lower bacterial loads. This points

to a potential role for disease tolerance mechanisms in this population through the prevention

of AMP-induced immunopathology.

A role for Disease Tolerance

Disease tolerance encompasses many distinct mechanisms of action, and describes

multiple different physiological processes, which have in common the wide umbrella of

maintaining host fitness without directly impacting pathogen load[6,77]. Maintenance of fitness in

the face of stress-inducing or homeostatic conditions is assured by tissue damage control

mechanisms[6,77]. Furthermore, these damage control responses can be activated in the event

of an infection, either due to direct action of pathogens (i.e., injury-inducing toxins) or indirectly

from immune-derived damages (i.e., immunopathology).

With this in mind, a putative role for disease tolerance was tested by measuring survival

of the evolved populations to prolonged exposure to heat-killed bacteria. This protocol

provided a means by which to test the response to damage, whether provoked by

pathogenicity of the bacteria or by the host's own response, independently of active pathogen

elimination. Although we hypothesize that the heat-killing treatment was insufficient to

denature all the virulence factors present in the suspension, we are yet to disentangle the

relative contribution of virulence factors versus immunopathology to the longevity

measurements. A similar longevity was observed for females and males of Control and

BactOral after feeding on heat-killed bacteria, but an interesting trend appeared when

comparing survival profiles (Fig. 6). We found that BactOral survived better to the stress of

prolonged exposure to heat-killed *P. entomophila*, specifically under an evolutionarily relevant

timeframe, that is, the period when flies reproduced during the selection experiment. Although

both Control and BactOral individuals fully succumbed to the chronic exposure to heat-killed

bacteria, significant differences in survival distinguish the two populations. Physiological

responses measured approximately at 10 days post-infection (the age at which populations

were selected to reproduce), can potentially represent evolutionary costs[78–81], which are not

the focus of this study.

Additionally, together with the gene expression data presented in Fig. 4, showing that

BactOral represses the activation of immune effectors earlier upon infection, it is tempting to

hypothesize that a similar process happened during the course of this experiment, and a role

may be attributed to tighter control and negative regulation of immune over-activation.

The improved response of adapted flies to exposure to heat-killed bacterial substrate

can be explained by a higher capacity to withstand damage, whether 1) provoked by their own

immune response, or 2) caused by virulence factors present in inactivated bacterial

supernatant. Additionally, process 1) could be correlated with the previously identified tighter

immune repression. Disentangling the relative contribution of these mechanisms to the overall

adapted phenotype would need further empirical characterization.

CONCLUSION

In sum, we were able to identify resistance and disease tolerance mechanisms as

targets of experimental evolution for increased survival to *P. entomophila* oral infection. Our

observations align with previous work that identifies a positive genetic correlation between

these two distinct immune defense strategies[82]. The increased survival of an experimentally

adapted population to oral infection directly relates to its capacity to decrease bacterial loads

after infection, independently of bacterial feeding rate, gut renewal capacity and bacterial

defecation. Previous work found resistance to be targeted by selection upon immune

challenge in *D. melanogaster*[79,83]. Our work adds to this approach by approaching the other

mechanisms at play in immunity, namely disease tolerance and behaviour, in the framework

of a single evolutionarily coherent system. We report one additional case of resistance

evolution that exhaustively approaches the different processes by which immunity may evolve

in populations fighting off persistent infections. In addition to resistance, we find a quantifiable

role for disease tolerance in the evolved response of BactOral, revealing the complex nature

of this adaptative process[82], which is likely to rely also on improved tissue repair coupled with

a more controlled deployment of damage-inducing immune effectors, such as AMPs[84,85]. This

interplay between disease tolerance and resistance, underscores the complex mechanistic

nature of immune response evolution with, undoubtedly, parallels in other host-pathogen

systems. Future work will focus on identifying the specific mechanisms of disease tolerance

and their physiological mode of action and interactions with resistance and their potential

evolutionary repeatability.

METHODS

Maintenance of Drosophila populations under relaxed selection

Experimentally evolved Control and BactOral populations were initially derived from a highly outbred population of *D. melanogaster*, as described in previous studies [19,46]. During experimental evolution, four replicates of each selection regime were maintained by orally infecting 310 females and 310 males per replicate, with a suspension of *P. entomophila* previously determined to cause approximately 66% of mortality in the starting population. After selection was terminated, populations were kept at high census (1500-2000 individuals per cage), allowing for maintenance of high genetic variability and optimal larval development. For all generations, flies were kept under constant temperature (25°C), humidity (55–65%) and light-darkness cycle (12:12), and fed with a standard cornmeal-agar medium, consisting of 4.5% molasses, 7.5% sugar, 7% corn-flower, 2% granulated yeast extract, 1% agar and 0.25% nipagin, mixed in distilled water. Control and BactOral were under selection for 24 generations and under relaxed selection for approximately 80 generations, after which one replicate of each regime was singled out to perform mechanistic experiments.

Each generation cycle lasted approximately 3 weeks, during which flies eclosed and remained undisturbed until they were 7-8 days old. At this time, fresh food was placed in the cages for 2 days, which allowed for oviposition of retained eggs. After this period (when flies were 9-11 days old), similarly to the protocol applied for selection, new fresh food was placed in the cages for controlled egg-lays (1-3 hours long) and establishment of the next generation.

Pathogen culture and infections

Oral infection protocol with *P. entomophila* (Rifampicin resistant strain kindly provided by Bruno Lemaitre) was adapted from the one used in Martins and Faria et al[19]. Briefly, single bacterial colonies were grown in kick-start cultures (5mL of LB) for approximately 8 hours after which they were transferred into larger volumes (1 L of LB) for overnight growth, both periods at 29°C. After centrifugation and resuspension of bacterial pellets, concentration was adjusted

to OD_{600}= 100 and finally diluted 1:1 with a 5% sucrose solution. For infection, 3-5 days-old

flies were separated by sex into groups of twenty with the use of CO_2 (at least 24 hours before

the start of the experiments) and allowed to feed on a filter disk embedded with the bacterial

solution described above for 24 hours (for survival assays) or for the amount of time specific

to each experiment (see Results). After this period, infected flies were flipped onto clean food

and survival was scored daily. Controls were fed a 1:1 LB/PBS:5% sucrose solution.

For the priming experiment, a solution of *P. entomophila* at OD_{600}= 100 was previously

heat-killed for 1 hour at 55ºC, mixed 1:1 with a 5% sucrose solution and fed to flies for 2 hours,

in a similar way as described above for regular oral infection. Efficiency was confirmed by

plating the heat-killed culture and confirm absence of bacterial growth. After this time, flies

were transferred into tubes containing a live solution of *P. entomophila* for 24 hours, after

which they were changed back to clean food and their survival scored.

For the experiment with prolonged exposure to heat-killed *P. entomophila* a similar

bacterial preparation and heat-killing protocol were performed as described above, but the

final dead bacteria suspension was mixed 1:1 with standard fly food and dispensed into fly

vials. To the final solution, extra agar powder was added to account for viscosity and to

optimize texture and left to cool down and solidify for at least 2 hours, before being fed to flies.

The same protocol was performed for the control treatment, in which fly food was mixed with

sterile 1X PBS. Exposure to these treatments was constant throughout the experiment, during

which replicates of 15 females and 15 males were co-housed in vials, exclusively feeding on

the dead-bacteria or PBS mixtures. Survival scoring began at the start of exposure and lasted

14 days. During the treatment, flies were frequently flipped onto fresh food mixture and survival

scoring continued daily. Overall efficiency of the bacterial heat-killing protocol was tested by

culturing the heat-killed suspension and confirm the absence of bacterial growth, prior to

storage at -20 ºC, which anteceded the mixing with the food or sucrose solution.

Survival analysis was done for the multiple experiments by fitting data with Cox

Proportional Hazards models (with *coxme* or *coxph* functions from the *survival* package[86])

using population, treatment, and sex as fixed factors and replicate (or replicate nested within

block) as a random factor. To assess the effect sizes of each factor, type 3 ANOVAs (*Anova*

from the *car* package[87]) were ran on the survival models. Additionally, multiple comparisons

with *tukey* adjustments were done on all the models to disentangle which specific pairwise

comparisons were significantly different (*emmeans* function[88]). Finally, the *cld* function of the

multcomp package[89] (with *sidak* confidence-level adjustments) was used to summarize the

results of the multiple comparisons across conditions. Survival models used were:

-    immunocompetence (Fig. 1)- *coxme*(~Population*Treatment*Sex+(1|Replicate));

-    priming (Fig. 5)- *coxme*(~Population*Treatment*Sex+(1|block/Replicate)) and

-    immune over-activation (Fig. 6)- *coxme*(~Population*Treatment*Sex+(1|Replicate),

data = survival_HKexposure).

Bacterial defecation assay

To have a measure of the rate of gut purge/defecation a change was introduced to the

bacterial infection protocol, whereby the feeding period was reduced to a 3-hour period. After

this time, flies were surface sterilized and separated individually into plastic spectrophotometry

cuvettes, previously filled with fly food. After 6 hours (6 hours.), the same flies were changed

into fresh cuvettes where they remained for an additional 12 hours (18 hours.), after which

they were discarded. Bacterial quantification was performed for both timepoints by washing

the interior surface of the cuvettes with PBS and plating, as described below.

Statistical analysis on number of defecated bacteria was done by fitting a zero-inflated

on the CFU counts and zeros (using the *zeroinfl* function[90] from the *pscl* package[91]). This

model allowed for the fitting of a negative-binomial distribution on the counts data and of a

binomial distribution on the zeros (*zeroinfl* (formula= Counts ~ Population * Sex * Timepoint |

Population * Timepoint, dist= "negbin")). To assess differences between specific pairwise

comparisons we used *emmeans* with *tukey* adjustments, on both distributions of the zero-

inflated model. In the instances where flies died during either timepoint, their count at that time

was removed from further analysis and the linear regression was not included in Figure 2B.

Quantification of food intake by CApillary FEeder Assay - CAFE

To measure the amount of bacterial solution ingested by single flies, an adapted

CAFE[47] assay was performed. We substituted the food of small fly vials with agarose, to

ensure humidity, and placed a glass capillary in each lid, filled with 5% sucrose solution. We

allowed individual flies to habituate to this experimental setup for 24 hours before the infection.

After this period, capillaries filled with *P. entomophila* suspension (prepared as described

before) were given to individual females to feed on and consumption estimated, at different

timepoints. To facilitate scoring in capillaries, the bacterial solution was mixed with blue food

colouring.

Modelling of cumulative amounts of bacteria eaten over time was done with a

Generalized Linear Model with a Gamma distribution, (*glm*(Quantity ~ Population*Timepoint))

using the *glm* function from the base *stat* R package[93]. For better visualization of feeding over

time, linear regressions were drawn between mean values of microliters eaten cumulatively

by each population at each timepoint, using the *stat_poly_eq* function of the *ggpmisc*

package[94]. Finally, multiple comparisons were done with *emmeans*[88], using *tukey*

adjustments, to disentangle differences between populations across timepoints.

Measurement of gut length

Females were individually dissected in PBS 1X, and their guts removed for imaging by

adapting a protocol described in[96]. Dissections started by fixing flies in the head with a Minutien

pin and gradually separating the gut from the remaining tissues, in a posterior-to-anterior

direction. After the gut was exposed, we added a few drops of a solution of 5% Acetic Acid

and 50%EtOH, which slightly whitens the tissues, allowing for better visualization and a slight

tissue fixation. After having isolated the guts, they were individually transferred to a coverslip

and photographed using a uEye camera installed on an Olympus SZX7 stereoscope.

Images were processed in FIJI v2.0.0 and measurements were collected by drawing

consecutive transects along the total length of guts, as illustrated in Supplementary figure 2A.

Each individual gut measurement was repeated 3 times, to account for errors in the manual

drawing of the transects and averaged to obtain a final gut length in millimeters (mm).

Differences in lengths of guts between populations were analysed with a Linear Mixed-

Effects Model (*lmer*(Length ~ Population * Treatment + (1|Block)) with *lmer* from the *lmerTest*

package[97]. The model was followed by multiple comparisons with *emmeans*[88], with a *tukey*

adjustment, for assessment of individual pairwise differences.

Quantification of bacterial loads

To quantify bacterial loads, we adapted an established protocol[19,92]. In short, individual

flies were surface sterilized (washed in EtOH 70%, bleach 60% and two washes with MiliQ

water) and placed in 96 deep-well plates, where they were homogenized with glass beads in

50 µL of LB using a TissueLyser II (Qiagen). Individual samples were serially diluted 5 times

with 1:10 dilution steps, plated as 4 µL droplets in LB+Rifampicin, and left to grow overnight

at 29ºC to optimize colony size for counting. For the bacterial defecation assay, cuvettes that

contained flies for each of the timepoints were filled with 1mL of PBS and incubated at room

temperature for 15 minutes, after which each sample was diluted up to 4 times (1:5 dilutions)

and platted as droplets in LB+Rifampicin.

Analysis of the time-course of infection progression was performed by dividing the data

obtained from the bacterial quantification into "counts" and "zeros" and using different models

to examine their dynamics in our evolved populations. For the counts, the modelling of the

mean of the load dynamics during the time-course was done using a sigmoidal function that

considers four parameters: a the log (initial load), k the timepoint at which the load is half of

the maximum value (in this case, a), h the slope at the k timepoint (steepest slope of the

curve), and *b* the log (final load). A linear model was applied to each of the parameters which

was fitted using a Monte Carlo Markov Chain approach. To estimate the impact of Sex and

Population on the likely distribution of the parameters, a normal error structure was assumed,

using Control females as reference. For the "zeros", a modified logistic model with four

parameters was ran: a_0 initial fraction of zeros, k_0 rate of zero disappearance, k_1 rate of zero

re-appearance, t_0 timepoint at which zeros re-appear and a_1 final fraction of zeros. Similarly,

to the "counts" analysis, linear models were used to assess the impact of Sex and Population

on all four parameters.

RNA extractions and quantitative PCR

For RNA extractions, flies from Control and BactOral population were infected as

described above and, at different timepoints, pools of 10 females were collected and

homogenized in 500 μL of Trizol. RNA extractions were performed using a Direct-zol™ RNA

MiniPrep w/ Zymo-Spin™ IIC Columns. After precipitation, a DNase I (RQ1 RNASE-FREE

DNASE 1* from Promega®) treatment was done to all samples, followed by reverse

transcription using the Thermo Scientific® RevertAid H Minus cDNA kit. cDNA was finally

diluted 1:5 for qPCR.

For quantification of gene expression, qPCRs were performed using SYBRTM Green

Master Mix (Thermo Scientific®) and reactions ran on 384-well plates (Applied Biosystems®).

The PCR conditions used in all experiments were: initial denaturation/ enzyme activation, 95ºC

for 10'; followed by 45 cycles of denaturation, 95ºC for 10"; annealing, 60ºC for 10"; extension,

72ºC for 30". Sequences of primers used for qPCRs are shown in Supp. Table. 1. Either EIF2

(eIF-2alpha) or Rpl32 were used as reference.

Analysis of gene expression differences was done with the relative quantification

technique (DDCt)[95]. Briefly, to the average of technical replicates for each candidate gene we

subtracted the average of the Ct values of the respective sample's house-keeping gene (EIF2

or Rpl32) (DCt) and normalized this value to the DCt of the respective reference condition. For

the time-course analysis, the reference was the "Unchallenged" timepoint while for

comparisons between "Unchallenged" conditions among populations, Control was used as

reference. Finally, we used reverse logarithm to transform the final gene expression values

into fold change levels (2^{-DDCt}) and analysed the Log_2(foldchange) differences across

Populations and/or Timepoints.

Statistical analysis on gene expression was done by running Linear Models (from base

R stats[93]) on the Log_2(foldchange) values of each individual gene (*lm*(logfold_AMP ~

Population * Timepoint), followed by an ANOVA (type three) from the car package[87]. To

individualize significant pairwise comparisons, *emmeans* was ran on the linear models (with a

Benjamini-Hochberg adjustment). Finally, the *cld* function of the multcomp package[89] (with

sidak confidence-level adjustments) was used to summarize the results of the multiple

comparisons across different conditions.

672 Statistical analysis and graphical representations

All datasets (excluding survival data) were tested for Normality before further analysis,

using the Shapiro-Wilk Test (*shapiro.test* function of the *stat* R package), for determination of

the most suitable type of statistical test to perform in each case. All model formulas used for

statistical analysis are discriminated throughout the Results sections. All graphical

representations were performed using the *ggplot2* package[98].

Statistical analysis and graphics for all experiments, excluding analysis of the bacterial

loads time-course modelling analysis (Fig. 5), were done on R version 4.2.1, through RStudio[99]

version 1.3.959. For the bacterial dynamics time-course, models were implemented and fitted

in Python[100], using the probabilistic programming language package *pymc3*[101] version 3.11.

Sensitivity (power) calculations for the CAFE and qPCR datasets was calculated using

G*Power 3.1 (https://www.psychologie.hhu.de/arbeitsgruppen/allgemeine-psychologie-und-

arbeitspsychologie/gpower).

Sensitivity analysis for gut length dataset was performed by simulation, using Python

packages *scipy*, *numpy*, and *statsmodels*. Briefly: a synthetic dataset of the same sample

sizes and structure as the real data but with a specified effect in both factors (Treatment and

Genotype) was generated. Multiple linear regression was then performed and the significance

(*p-value*) of both factors was recorded. This was performed 10^5 times, and the fraction of

significant tests was recorded (power, independently for both factors). A numerical search for

the minimum effect sizes that achieve 80% power on both factors was then performed.

ACKNOWLEDGEMENTS

We thank Dr Bruno Lemaitre for stocks, Liliana Vieira and Sandra Crisóstomo for

technical assistance, André Barros and David Duneau for help with analyses, and Elvira

Lafuente, Diogo Roque and Patrícia Duarte for critical comments and discussions that improved

the manuscript. This work was supported by Instituto Gulbenkian de Ciência/Fundação Calouste

Gulbenkian and FCT-Fundação para a Ciência e a Tecnologia (Portugal) UID/Multi/04555/2013

and UIDB/00329/2020. TFP (BD/128432/2017 and COVID/BD/151645/2021) and to PAA

(BD/06404/2020) were supported FCT—Fundação para a Ciência e a Tecnologia (Portugal); and

by C.S. CONGENTO, project LISBOA-01-0145-FEDER-022170, co-financed by Lisboa Regional

Operational Programme (Lisboa 2020), under the Portugal 2020 Partnership Agreement, through

the European Regional Development Fund (ERDF), and Foundation for Science and Technology

(Portugal).

DECLARATION

The authors declare that this work is not under review elsewhere and that none of its authors has any conflict of interest to declare.

DATA AVAILABILITY

All relevant data are within the manuscript and its Supporting Information files.

AUTHOR CONTRIBUTIONS

Conceptualization: Tânia F. Paulo and Élio Sucena.

Data curation: Tânia F. Paulo

Formal analysis: Tânia F. Paulo, Tiago Paixão

Funding acquisition: Élio Sucena.

Investigation: Tânia F. Paulo, Priscilla A. Akyaw and Élio Sucena.

Methodology: Tânia F. Paulo, Priscilla A. Akyaw, Tiago Paixão and Élio Sucena.

Resources: Élio Sucena and Tiago Paixão.

Supervision: Élio Sucena.

Visualization: Tânia F. Paulo.

Writing – original draft: Tânia F. Paulo and Élio Sucena.

Writing – review & editing: Tânia F. Paulo, Priscilla A. Akyaw, Tiago Paixão and Élio Sucena.

REFERENCES

(1)   Fischer, O.; Schmid-Hempel, P. Selection by Parasites May Increase Host Recombination Frequency. *Biol. Lett.* **2005**. https://doi.org/10.1098/rsbl.2005.0296.

(2)   Schmid-Hempel, P.; Ebert, D. On the Evolutionary Ecology of Specific Immune Defence. *Trends Ecol. Evol.* **2003**, *18* (1), 27–32. https://doi.org/10.1016/S0169-5347(02)00013-7.

(3)   Schulenburg, H.; Kurtz, J.; Moret, Y.; Siva-Jothy, M. T. Introduction. Ecological Immunology. *Philos. Trans. R. Soc. B Biol. Sci.* **2008**, *364* (1513), 3–14. https://doi.org/10.1098/rstb.2008.0249.

(4)   Gibson, A. K.; Amoroso, C. R. Evolution and Ecology of Parasite Avoidance. *Annu. Rev. Ecol. Evol. Syst.* **2022**, *53* (1), 47–67. https://doi.org/10.1146/annurev-ecolsys-102220-020636.

(5)   Kutzer, M. A. M.; Armitage, S. A. O. Maximising Fitness in the Face of Parasites: A Review of Host Tolerance. *Zoology* **2016**, *119* (4), 281–289. https://doi.org/10.1016/j.zool.2016.05.011.

(6)   Medzhitov, R.; Schneider, D. S.; Soares, M. P. Disease Tolerance as a Defense Strategy. *Science* **2012**, *335* (6071), 936–941. https://doi.org/10.1126/science.1214935.

(7)   Ayres, J. S.; Schneider, D. S. Two Ways to Survive an Infection: What Resistance and Tolerance Can Teach Us about Treatments for Infectious Diseases. *Nat. Rev. Immunol.* **2008**, *8* (11), 889–895. https://doi.org/10.1038/nri2432.

(8)   Råberg, L.; Graham, A. L.; Read, A. F. Decomposing Health: Tolerance and Resistance to Parasites in Animals. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **2009**, *364* (1513), 37–49. https://doi.org/10.1098/rstb.2008.0184.

(9)   Menge, D. M.; Zhong, D.; Guda, T.; Gouagna, L.; Githure, J.; Beier, J.; Yan, G. Quantitative Trait Loci Controlling Refractoriness to Plasmodium Falciparum in Natural Anopheles Gambiae Mosquitoes from a Malaria-Endemic Region in Western Kenya. *Genetics* **2006**, *173* (1), 235–241. https://doi.org/10.1534/genetics.105.055129.

(10)   Kim, D. H.; Feinbaum, R.; Alloing, G.; Emerson, F. E.; Garsin, D. A.; Inoue, H.; Tanaka-Hino, M.; Hisamoto, N.; Matsumoto, K.; Tan, M.-W.; Ausubel, F. M. A Conserved P38 MAP Kinase Pathway in Caenorhabditis Elegans Innate Immunity. *Science* **2002**, *297* (5581), 623–626. https://doi.org/10.1126/science.1073759.

(11)   Scharsack, J. P.; Kalbe, M.; Harrod, C.; Rauch, G. Habitat-Specific Adaptation of Immune Responses of Stickleback (Gasterosteus Aculeatus) Lake and River Ecotypes.

Proc. R. Soc. B Biol. Sci. **2007**, *274* (1617), 1523–1532.

https://doi.org/10.1098/rspb.2007.0210.

(12)   Faria, V. G.; Martins, N. E.; Magalhães, S.; Paulo, T. F.; Nolte, V.; Schlötterer, C.;

Sucena, É.; Teixeira, L. Drosophila Adaptation to Viral Infection through Defensive Symbiont

Evolution. *PLOS Genet.* **2016**, *12* (9), e1006297.

https://doi.org/10.1371/journal.pgen.1006297.

(13)   Martins, N. E.; Faria, V. G.; Nolte, V.; Schlotterer, C.; Teixeira, L.; Sucena, E.;

Magalhaes, S. Host Adaptation to Viruses Relies on Few Genes with Different Cross-

Resistance Properties. *Proc. Natl. Acad. Sci.* **2014**, *111* (16), 5938–5943.

https://doi.org/10.1073/pnas.1400378111.

(14)   Gupta, V.; Venkatesan, S.; Chatterjee, M.; Syed, Z. A.; Nivsarkar, V.; Prasad, N. G.

No Apparent Cost of Evolved Immune Response in Drosophila Melanogaster. *Evolution*

2016, *70* (4), 934–943. https://doi.org/10.1111/evo.12896.

(15)   Zbinden, M.; Haag, C. R.; Ebert, D. Experimental Evolution of Field Populations of

Daphnia Magna in Response to Parasite Treatment. *J. Evol. Biol.* **2008**, *21* (4), 1068–1078.

https://doi.org/10.1111/j.1420-9101.2008.01541.x.

(16)   Kawecki, T. J.; Lenski, R. E.; Ebert, D.; Hollis, B.; Olivieri, I.; Whitlock, M. C.

Experimental Evolution. *Trends Ecol. Evol.* **2012**, *27* (10), 547–560.

https://doi.org/10.1016/j.tree.2012.06.001.

(17)   Ebert, D. Experimental Evolution of Parasites. *Science* **1998**, *282* (5393), 1432–

1436. https://doi.org/10.1126/science.282.5393.1432.

(18)   Marxer, M.; Barribeau, S.; Schmid-Hempel, P. Experimental Evolution of a

Trypanosome Parasite of Bumblebees and Its Implications for Infection Success and Host

Immune Response. *Evol. Biol.* **2016**, *43* (2), 160–170. https://doi.org/10.1007/s11692-015-

9366-z.

(19)   Martins, N. E.; Faria, V. G.; Teixeira, L.; Magalhães, S.; Sucena, É. Host Adaptation

Is Contingent upon the Infection Route Taken by Pathogens. *PLoS Pathog.* **2013**, *9* (9),

e1003601. https://doi.org/10.1371/journal.ppat.1003601.

(20)   Khan, I.; Prakash, A.; Agashe, D. Experimental Evolution of Insect Immune Memory

versus Pathogen Resistance. *Proc. R. Soc. B Biol. Sci.* **2017**, *284* (1869), 20171583.

https://doi.org/10.1098/rspb.2017.1583.

(21)   Cabrera, K.; Hoard, D. S.; Gibson, O.; Martinez, D. I.; Wunderlich, Z. Drosophila

Immune Priming to Enterococcus Faecalis Relies on Immune Tolerance Rather than

Resistance. *PLOS Pathog.* **2023**, *19* (8), e1011567.

https://doi.org/10.1371/journal.ppat.1011567.

(22)   Mondotte, J. A.; Saleh, M.-C. Chapter Eleven - Antiviral Immune Response and the

Route of Infection in Drosophila Melanogaster. In *Advances in Virus Research*; Kielian, M.,

Mettenleiter, T. C., Roossinck, M. J., Eds.; Academic Press, 2018; Vol. 100, pp 247–278.

https://doi.org/10.1016/bs.aivir.2017.10.006.

(23)   Vodovar, N.; Vinals, M.; Liehl, P.; Basset, A.; Degrouard, J.; Spellman, P.; Boccard,

F.; Lemaitre, B. Drosophila Host Defense after Oral Infection by an Entomopathogenic

Pseudomonas Species. *Proc. Natl. Acad. Sci. U. S. A.* **2005**, *102* (32), 11414–11419.

https://doi.org/10.1073/pnas.0502240102.

(24)   Chandler, J. A.; Lang, J. M.; Bhatnagar, S.; Eisen, J. A.; Kopp, A. Bacterial

Communities of Diverse Drosophila Species: Ecological Context of a Host–Microbe Model

System. *PLOS Genet.* **2011**, *7* (9), e1002272. https://doi.org/10.1371/journal.pgen.1002272.

(25)   Charroux, B.; Daian, F.; Royet, J. Drosophila Aversive Behavior toward Erwinia

Carotovora Carotovora Is Mediated by Bitter Neurons and Leukokinin. *iScience* **2020**, *23* (6),

101152. https://doi.org/10.1016/j.isci.2020.101152.

(26)   Soldano, A.; Alpizar, Y. A.; Boonen, B.; Franco, L.; López-Requena, A.; Liu, G.;

Mora, N.; Yaksi, E.; Voets, T.; Vennekens, R.; Hassan, B. A.; Talavera, K. Gustatory-

Mediated Avoidance of Bacterial Lipopolysaccharides via TRPA1 Activation in Drosophila.

eLife **2016**, *5* (JUN2016), 1–15. https://doi.org/10.7554/eLife.13133.

(27)   Kuraishi, T.; Binggeli, O.; Opota, O.; Buchon, N.; Lemaitre, B. Genetic Evidence for a

Protective Role of the Peritrophic Matrix against Intestinal Bacterial Infection in Drosophila

Melanogaster. *Proc. Natl. Acad. Sci. U. S. A.* **2011**, *108* (38), 15966–15971.

https://doi.org/10.1073/pnas.1105994108.

(28)   Overend, G.; Luo, Y.; Henderson, L.; Douglas, A. E.; Davies, S. A.; Dow, J. A. T.

Molecular Mechanism and Functional Significance of Acid Generation in the Drosophila

Midgut. *Nat. Publ. Group* **2016**, No. May, 1–11. https://doi.org/10.1038/srep27242.

(29)   Du, E. J.; Ahn, T. J.; Kwon, I.; Lee, J. H.; Park, J. H.; Park, S. H.; Kang, T. M.; Cho,

H.; Kim, T. J.; Kim, H. W.; Jun, Y.; Lee, H. J.; Lee, Y. S.; Kwon, J. Y.; Kang, K. J. TrpA1

Regulates Defecation of Food-Borne Pathogens under the Control of the Duox Pathway.

PLoS Genet. **2016**, *12* (1), 1–21. https://doi.org/10.1371/journal.pgen.1005773.

(30)   Fink, C.; Hoffmann, J.; Knop, M.; Li, Y.; Isermann, K.; Roeder, T. Intestinal FoxO

Signaling Is Required to Survive Oral Infection in Drosophila. *Mucosal Immunol.* **2016**, *9* (4),

927–936. https://doi.org/10.1038/mi.2015.112.

(31)   Ha, E.-M.; Oh, C.-T.; Bae, Y. S.; Lee, W.-J. A Direct Role for Dual Oxidase in

Drosophila Gut Immunity. *Science* **2005**, *310* (5749), 847–850.

https://doi.org/10.1126/science.1117311.

(32)   Ha, E.-M.; Lee, K.-A.; Seo, Y. Y.; Kim, S.-H.; Lim, J.-H.; Oh, B.-H.; Kim, J.; Lee, W.-J.

Coordination of Multiple Dual Oxidase–Regulatory Pathways in Responses to Commensal

and Infectious Microbes in Drosophila Gut. *Nat. Immunol.* **2009**, *10* (9), 949–957.

https://doi.org/10.1038/ni.1765.

(33)   Tian, A.; Morejon, V.; Kohoutek, S.; Huang, Y.-C.; Deng, W.-M.; Jiang, J. Damage-

Induced Regeneration of the Intestinal Stem Cell Pool through Enteroblast Mitosis in the

Drosophila Midgut. *EMBO J.* **2022**, e110834. https://doi.org/10.15252/embj.2022110834.

(34)   Bonfini, A.; Liu, X.; Buchon, N. From Pathogens to Microbiota: How Drosophila

Intestinal Stem Cells React to Gut Microbes. *Dev. Comp. Immunol.* **2015**.

https://doi.org/10.1016/j.dci.2016.02.008.

(35)   Buchon, N.; Broderick, N. A.; Poidevin, M.; Pradervand, S.; Lemaitre, B. Drosophila

Intestinal Response to Bacterial Infection: Activation of Host Defense and Stem Cell

Proliferation. *Cell Host Microbe* **2009**, *5* (2), 200–211.

https://doi.org/10.1016/j.chom.2009.01.003.

(36)   Buchon, N.; Broderick, N. A.; Kuraishi, T.; Lemaitre, B. Drosophila EGFR Pathway

Coordinates Stem Cell Proliferation and Gut Remodeling Following Infection. *BMC Biol.*

2010, *8* (1), 152. https://doi.org/10.1186/1741-7007-8-152.

(37)   Nehme, N. T.; Liégeois, S.; Kele, B.; Giammarinaro, P.; Pradel, E.; Hoffmann, J. A.;

Ewbank, J. J.; Ferrandon, D. A Model of Bacterial Intestinal Infections in Drosophila

Melanogaster. *PLoS Pathog.* **2007**, *3* (11), 1694–1709.

https://doi.org/10.1371/journal.ppat.0030173.

(38)   Limmer, S.; Haller, S.; Drenkard, E.; Lee, J.; Yu, S.; Kocks, C.; Ausubel, F. M.;

Ferrandon, D. Pseudomonas Aeruginosa RhlR Is Required to Neutralize the Cellular

Immune Response in a Drosophila Melanogaster Oral Infection Model. *Proc. Natl. Acad. Sci.*

U. S. A. **2011**, *108* (42), 17378–17383. https://doi.org/10.1073/pnas.1114907108.

(39)   Yang, S.; Zhao, Y.; Yu, J.; Fan, Z.; Gong, S.; Tang, H.; Pan, L. Sugar Alcohols of

Polyol Pathway Serve as Alarmins to Mediate Local-Systemic Innate Immune

Communication in Drosophila. *Cell Host Microbe* **2019**, *26* (2), 240-251.e8.

https://doi.org/10.1016/j.chom.2019.07.001.

(40)   Zaidman-Rémy, A.; Hervé, M.; Poidevin, M.; Pili-Floury, S.; Kim, M.-S.; Blanot, D.;

Oh, B.-H.; Ueda, R.; Mengin-Lecreulx, D.; Lemaitre, B. The Drosophila Amidase PGRP-LB

Modulates the Immune Response to Bacterial Infection. *Immunity* **2006**, *24* (4), 463–473.

https://doi.org/10.1016/j.immuni.2006.02.012.

(41)   Basset, A.; Khush, R. S.; Braun, A.; Gardan, L.; Boccard, F.; Hoffmann, J. A.;

Lemaitre, B. The Phytopathogenic Bacteria Erwinia Carotovora Infects Drosophila and

Activates an Immune Response. *Proc. Natl. Acad. Sci.* **2000**, *97* (7), 3376–3381.

https://doi.org/10.1073/pnas.97.7.3376.

(42)   Vodovar, N.; Vallenet, D.; Cruveiller, S.; Rouy, Z.; Barbe, V.; Acosta, C.; Cattolico, L.;

Jubin, C.; Lajus, A.; Segurens, B.; Vacherie, B.; Wincker, P.; Weissenbach, J.; Lemaitre, B.;

Médigue, C.; Boccard, F. Complete Genome Sequence of the Entomopathogenic and

Metabolically Versatile Soil Bacterium Pseudomonas Entomophila. *Nat. Biotechnol.* **2006**, *24*

(6), 673–679. https://doi.org/10.1038/nbt1212.

(43)   Dieppois, G.; Opota, O.; Lalucat, J.; Lemaitre, B. Pseudomonas Entomophila: A

Versatile Bacterium with Entomopathogenic Properties. In *Pseudomonas*; Springer

Netherlands: Dordrecht, 2015; pp 25–49. https://doi.org/10.1007/978-94-017-9555-5_2.

(44)   Opota, O.; Vallet-Gély, I.; Vincentelli, R.; Kellenberger, C.; Iacovache, I.; Gonzalez,

M. R.; Roussel, A.; Goot, F.-G. van der; Lemaitre, B. Monalysin, a Novel ß-Pore-Forming

Toxin from the Drosophila Pathogen Pseudomonas Entomophila, Contributes to Host

Intestinal Damage and Lethality. *PLOS Pathog.* **2011**, *7* (9), e1002259.

https://doi.org/10.1371/journal.ppat.1002259.

(45)   Chakrabarti, S.; Liehl, P.; Buchon, N.; Lemaitre, B. Infection-Induced Host

Translational Blockage Inhibits Immune Responses and Epithelial Renewal in the Drosophila

Gut. *Cell Host Microbe* **2012**, *12* (1), 60–70. https://doi.org/10.1016/j.chom.2012.06.001.

(46)   Faria, V. G.; Martins, N. E.; Paulo, T.; Teixeira, L.; Sucena, É.; Magalhães, S.

Evolution of Drosophila Resistance against Different Pathogens and Infection Routes Entails

No Detectable Maintenance Costs. *Evolution* **2015**, *69* (11), 2799–2809.

https://doi.org/10.1111/evo.12782.

(47)   Ja, W. W.; Carvalho, G. B.; Mak, E. M.; De La Rosa, N. N.; Fang, A. Y.; Liong, J. C.;

Brummel, T.; Benzer, S. Prandiology of Drosophila and the CAFE Assay. *Proc. Natl. Acad.*

Sci. U. S. A. **2007**, *104* (20), 8253–8256. https://doi.org/10.1073/pnas.0702726104.

(48)   Vincent, C. M.; Dionne, M. S. Disparate Regulation of IMD Signaling Drives Sex

Differences in Infection Pathology in Drosophila Melanogaster. *Proc. Natl. Acad. Sci.* **2021**,

118 (32), e2026554118. https://doi.org/10.1073/pnas.2026554118.

(49)   Belmonte, R. L.; Corbally, M.-K.; Duneau, D. F.; Regan, J. C. Sexual Dimorphisms in

Innate Immunity and Responses to Infection in Drosophila Melanogaster. *Front. Immunol.*

2020, *10*.

(50)   Duneau, D. F.; Kondolf, H. C.; Im, J. H.; Ortiz, G. A.; Chow, C.; Fox, M. A.; Eugénio,

A. T.; Revah, J.; Buchon, N.; Lazzaro, B. P. The Toll Pathway Underlies Host Sexual

Dimorphism in Resistance to Both Gram-Negative and Gram-Positive Bacteria in Mated

Drosophila. *BMC Biol.* **2017**, *15* (1), 124. https://doi.org/10.1186/s12915-017-0466-3.

(51)   Hanson, M. A.; Dostálová, A.; Ceroni, C.; Poidevin, M.; Kondo, S.; Lemaitre, B.

Synergy and Remarkable Specificity of Antimicrobial Peptides in Vivo Using a Systematic

Knockout Approach. *eLife* **2019**, *8*, e44341. https://doi.org/10.7554/eLife.44341.

(52)   Carboni, A. L.; Hanson, M. A.; Lindsay, S. A.; Wasserman, S. A.; Lemaitre, B.

Cecropins Contribute to Drosophila Host Defense against a Subset of Fungal and Gram-

Negative Bacterial Infection. *Genetics* **2022**, *220* (1), iyab188.

https://doi.org/10.1093/genetics/iyab188.

(53)   Keesey, I. W.; Koerte, S.; Khallaf, M. A.; Retzke, T.; Guillou, A.; Grosse-Wilde, E.;

Buchon, N.; Knaden, M.; Hansson, B. S. Pathogenic Bacteria Enhance Dispersal through

Alteration of Drosophila Social Communication. *Nat. Commun.* **2017**, *8* (1), 265.

https://doi.org/10.1038/s41467-017-00334-9.

(54)   Patrnogic, J.; Castillo, J. C.; Shokal, U.; Yadav, S.; Kenney, E.; Heryanto, C.;

Ozakman, Y.; Eleftherianos, I. Pre-Exposure to Non-Pathogenic Bacteria Does Not Protect

Drosophila against the Entomopathogenic Bacterium Photorhabdus. *PLOS ONE* **2018**, *13*

(10), e0205256. https://doi.org/10.1371/journal.pone.0205256.

(55)   Wen, Y.; He, Z.; Xu, T.; Jiao, Y.; Liu, X.; Wang, Y.-F.; Yu, X.-Q. Ingestion of Killed

Bacteria Activates Antimicrobial Peptide Genes in Drosophila Melanogaster and Protects

Flies from Septic Infection. *Dev. Comp. Immunol.* **2019**, *95*, 10–18.

https://doi.org/10.1016/j.dci.2019.02.001.

(56)   Paul Schmid-Hempel. Evolutionary Ecology of Insect Immune Defenses. *Annu Rev*

Entomol **2005**, *50*, 529–551. https://doi.org/10.1146/annurev.ento.50.071803.130420.

(57)   Eleftherianos, I.; Tafesh-Edwards, G.; Mohamed, A. Pathogen Infection Routes and

Host Innate Immunity: Lessons from Insects. *Immunol. Lett.* **2022**, *247*, 46–51.

https://doi.org/10.1016/j.imlet.2022.05.006.

(58)   Gupta, V.; Vasanthakrishnan, R. B.; Siva-Jothy, J.; Monteith, K. M.; Brown, S. P.;

Vale, P. F. The Route of Infection Determines Wolbachia Antibacterial Protection in

Drosophila. *Proc. R. Soc. B Biol. Sci.* **2017**, *284* (1856), 20170809.

https://doi.org/10.1098/rspb.2017.0809.

(59)   Vizoso, D. B.; Ebert, D. Phenotypic Plasticity of Host–Parasite Interactions in

Response to the Route of Infection. *J. Evol. Biol.* **2005**, *18* (4), 911–921.

https://doi.org/10.1111/j.1420-9101.2005.00920.x.

(60) Dekmak, A. S.; Yang, X.; Dohna, H. Z.; Buchon, N.; Osta, M. A. The Route of

Infection Influences the Contribution of Key Immunity Genes to Antibacterial Defense in

Anopheles Gambiae. *J. Innate Immun.* **2021**, *13* (2), 107–126.

https://doi.org/10.1159/000511401.

(61) Hall, M. D.; Bento, G.; Ebert, D. The Evolutionary Consequences of Stepwise

Infection Processes. *Trends Ecol. Evol.* **2017**, *xx*, 1–12.

https://doi.org/10.1016/j.tree.2017.05.009.

(62) Vale, P. F.; Jardine, M. D. Infection Avoidance Behavior: Viral Exposure Reduces the

Motivation to Forage in Female Drosophila Melanogaster. *Fly (Austin)* **2017**, *11* (1), 3–9.

https://doi.org/10.1080/19336934.2016.1207029.

(63) Kobler, J. M.; Rodriguez Jimenez, F. J.; Petcu, I.; Grunwald Kadow, I. C. Immune

Receptor Signaling and the Mushroom Body Mediate Post-Ingestion Pathogen Avoidance.

Curr. Biol. **2020**, *30* (23), 4693-4709.e3. https://doi.org/10.1016/j.cub.2020.09.022.

(64) Stensmyr, M. C.; Dweck, H. K. M.; Farhan, A.; Ibba, I.; Strutz, A.; Mukunda, L.; Linz,

J.; Grabe, V.; Steck, K.; Lavista-Llanos, S.; Wicher, D.; Sachse, S.; Knaden, M.; Becher, P.

G.; Seki, Y.; Hansson, B. S. A Conserved Dedicated Olfactory Circuit for Detecting Harmful

Microbes in Drosophila. *Cell* **2012**, *151* (6), 1345–1357.

https://doi.org/10.1016/j.cell.2012.09.046.

(65) Ahlawat, N.; Geeta Arun, M.; Maggu, K.; Jigisha; Singh, A.; Prasad, N. G. Drosophila

Melanogaster Hosts Coevolving with Pseudomonas Entomophila Pathogen Show Sex-

Specific Patterns of Local Adaptation. *BMC Ecol. Evol.* **2022**, *22* (1), 77.

https://doi.org/10.1186/s12862-022-02031-8.

(66) Faria, V. G.; Martins, N. E.; Schlötterer, C.; Sucena, É. Readapting to DCV Infection

without Wolbachia: Frequency Changes of Drosophila Antiviral Alleles Can Replace

Endosymbiont Protection. *Genome Biol. Evol.* **2018**, *10* (7), 1783–1791.

https://doi.org/10.1093/gbe/evy137.

(67) Liehl, P.; Blight, M.; Vodovar, N.; Boccard, F.; Lemaitre, B. Prevalence of Local

Immune Response against Oral Infection in a Drosophila/Pseudomonas Infection Model.

PLoS Pathog. **2006**, *2* (6), e56. https://doi.org/10.1371/journal.ppat.0020056.

(68) Liu, X.; Nagy, P.; Bonfini, A.; Houtz, P.; Bing, X.-L.; Yang, X.; Buchon, N. Microbes

Affect Gut Epithelial Cell Composition through Immune-Dependent Regulation of Intestinal

Stem Cell Differentiation. *Cell Rep.* **2022**, *38* (13).

https://doi.org/10.1016/j.celrep.2022.110572.

(69)   Hudry, B.; Khadayate, S.; Miguel-Aliaga, I. The Sexual Identity of Adult Intestinal
Stem Cells Controls Organ Size and Plasticity. *Nature* **2016**, *530* (7590), 344–348.
https://doi.org/10.1038/nature16953.

(70)   Hudry, B.; de Goeij, E.; Mineo, A.; Gaspar, P.; Hadjieconomou, D.; Studd, C.;
Mokochinski, J. B.; Kramer, H. B.; Plaçais, P.-Y.; Preat, T.; Miguel-Aliaga, I. Sex Differences
in Intestinal Carbohydrate Metabolism Promote Food Intake and Sperm Maturation. *Cell*
2019, *178* (4), 901-918.e16. https://doi.org/10.1016/j.cell.2019.07.029.

(71)   Regan, J. C.; Khericha, M.; Dobson, A. J.; Bolukbasi, E.; Rattanavirotkul, N.;
Partridge, L. Sex Difference in Pathology of the Ageing Gut Mediates the Greater Response
of Female Lifespan to Dietary Restriction. *eLife* **2016**, *5*, e10956.
https://doi.org/10.7554/eLife.10956.

(72)   Wukitch, A. M.; Lawrence, M. M.; Satriale, F. P.; Patel, A.; Ginder, G. M.; Van Beek,
E. J.; Gilani, O.; Chambers, M. C. Impact of Chronic Infection on Resistance and Tolerance
to Secondary Infection in Drosophila Melanogaster. *Infect. Immun.* **2023**, e0036022.
https://doi.org/10.1128/iai.00360-22.

(73)   Chambers, M. C.; Jacobson, E.; Khalil, S.; Lazzaro, B. P. Consequences of Chronic
Bacterial Infection in Drosophila Melanogaster. *PLoS ONE* **2019**, *14* (10), e0224440.
https://doi.org/10.1371/journal.pone.0224440.

(74)   Sibley, C. D.; Duan, K.; Fischer, C.; Parkins, M. D.; Storey, D. G.; Rabin, H. R.;
Surette, M. G. Discerning the Complexity of Community Interactions Using a Drosophila
Model of Polymicrobial Infections. *PLOS Pathog.* **2008**, *4* (10), e1000184.
https://doi.org/10.1371/journal.ppat.1000184.

(75)   Lin, S. J. H.; Fulzele, A.; Cohen, L. B.; Bennett, E. J.; Wasserman, S. A. Bombardier
Enables Delivery of Short-Form Bomanins in the Drosophila Toll Response. *Front. Immunol.*
2019, *10*, 3040. https://doi.org/10.3389/fimmu.2019.03040.

(76)   Lindsay, S. A.; Lin, S. J. H.; Wasserman, S. A. Short-Form Bomanins Mediate
Humoral Immunity in Drosophila. *J. Innate Immun.* **2018**, *10* (4), 306–314.
https://doi.org/10.1159/000489831.

(77)   Martins, R.; Carlos, A. R.; Braza, F.; Thompson, J. A.; Bastos-Amador, P.; Ramos,
S.; Soares, M. P. Disease Tolerance as an Inherent Component of Immunity. *Annu. Rev.*
Immunol. **2019**, *37* (1), 405–437. https://doi.org/10.1146/annurev-immunol-042718-041739.

(78)   Vijendravarma, R. K.; Kraaijeveld, A. R.; Godfray, H. C. J. Experimental Evolution
Shows Drosophila Melanogaster Resistance to a Microsporidian Pathogen Has Fitness
Costs. *Evol. Int. J. Org. Evol.* **2009**, *63* (1), 104–114. https://doi.org/10.1111/j.1558-
5646.2008.00516.x.

(79) Kraaijeveld, A. R.; Godfray, H. C. J. Trade-off between Parasitoid Resistance and

Larval Competitive Ability in Drosophila Melanogaster. *Nature* **1997**, *389* (6648), 278–280.

https://doi.org/10.1038/38483.

(80) Hanson, M. A.; Lemaitre, B.; Unckless, R. L. Dynamic Evolution of Antimicrobial

Peptides Underscores Trade-Offs Between Immunity and Ecological Fitness. *Front.*

Immunol. **2019**, *10*, 2620. https://doi.org/10.3389/fimmu.2019.02620.

(81) Lochmiller, R. L.; Deerenberg, C. Trade-Offs in Evolutionary Immunology: Just What

Is the Cost of Immunity? *Oikos* **2000**, *88* (1), 87–98. https://doi.org/doi:10.1034/j.1600-

0706.2000.880110.x.

(82) Howick, V. M.; Lazzaro, B. P. The Genetic Architecture of Defence as Resistance to

and Tolerance of Bacterial Infection in Drosophila Melanogaster. *Mol. Ecol.* **2017**, *26* (6),

1533–1546. https://doi.org/10.1111/mec.14017.

(83) Leitão, A. B.; Arunkumar, R.; Day, J. P.; Geldman, E. M.; Morin-Poulard, I.; Crozatier,

M.; Jiggins, F. M. Constitutive Activation of Cellular Immunity Underlies the Evolution of

Resistance to Infection in Drosophila. *eLife* **2020**, *9*, e59095.

https://doi.org/10.7554/eLife.59095.

(84) Lhocine, N.; Ribeiro, P. S.; Buchon, N.; Wepf, A.; Wilson, R.; Tenev, T.; Lemaitre, B.;

Gstaiger, M.; Meier, P.; Leulier, F. PIMS Modulates Immune Tolerance by Negatively

Regulating Drosophila Innate Immune Signaling. *Cell Host Microbe* **2008**, *4* (2), 147–158.

https://doi.org/10.1016/j.chom.2008.07.004.

(85) Bonnay, F.; Cohen-Berros, E.; Hoffmann, M.; Kim, S. Y.; Boulianne, G. L.; Hoffmann,

J. A.; Matt, N.; Reichhart, J.-M. Big Bang Gene Modulates Gut Immune Tolerance in

Drosophila. *Proc. Natl. Acad. Sci. U. S. A.* **2013**, *110* (8), 2957–2962.

https://doi.org/10.1073/pnas.1221910110.

(86) T.M. Therneau. _A Package for Survival Analysis in R_., 2022. https://CRAN.R-

project.org/package=survival.

(87) John Fox; Sanford Weisberg. An {R} Companion to Applied Regression, 2019.

https://socialsciences.mcmaster.ca/jfox/Books/Companion/.

(88) R. Lenth. _emmeans: Estimated Marginal Means, Aka Least-Squares Means_.,

2022. https://CRAN.R-project.org/package=emmeans.

(89) Torsten Hothorn; Frank Bretz; Peter Westfall. Simultaneous Inference in General

Parametric Models. *Biom. J.* **2008**, *50* (3), 346–363.

(90) Achim Zeileis; Christian Kleiber; Simon Jackman. Regression Models for Count Data

in R. *J. Stat. Softw.* **2008**, *27* (8).

(91) Simon Jackman. Pscl: Classes and Methods for R Developed in the Political Science

Computational Laboratory, 2020. https://github.com/atahk/pscl/.

(92) Neyen, C.; Bretscher, A. J.; Binggeli, O.; Lemaitre, B. Methods to Study Drosophila

Immunity. *Methods* **2014**, *68* (1), 116–128. https://doi.org/10.1016/j.ymeth.2014.02.023.

(93) R Core Team. R: A Language and Environment for Statistical Computing., 2022.

https://www.R-project.org/.

(94) Aphalo P. _ggpmisc: Miscellaneous Extensions to 'ggplot2'_, 2022. https://CRAN.R-

project.org/package=ggpmisc.

(95) Livak, K. J.; Schmittgen, T. D. Analysis of Relative Gene Expression Data Using

Real-Time Quantitative PCR and the 2−ΔΔCT Method. *Methods* **2001**, *25* (4), 402–408.

https://doi.org/10.1006/meth.2001.1262.

(96) Micchelli, C. A. Whole-Mount Immunostaining of the Adult Drosophila

Gastrointestinal Tract. *Methods* **2014**, *68* (1), 273–279.

https://doi.org/10.1016/j.ymeth.2014.03.022.

(97) Kuznetsova A; Brockhoff PB; Christensen RHB. LmerTest Package: Tests in Linear

Mixed Effects Models. *J. Stat. Softw.* **2017**, *82* (12), 1–26.

https://doi.org/10.18637/jss.v082.i13.

(98) H. Wickham. Ggplot2: Elegant Graphics for Data Analysis, 2016.

https://ggplot2.tidyverse.org.

(99) RStudio Team. RStudio: Integrated Development Environment for R, 2020.

http://www.rstudio.com/.

(100) Van Rossum; Guido and Drake; Fred L. Python 3 Reference Manual, 2009.

(101) Salvatier J.; Wiecki T.V.; Fonnesbeck C. Probabilistic Programming in Python Using

PyMC3. *PeerJ Comput. Sci. 2 E55* **2016**. https://doi.org/10.7717/peerj-cs.55.

1059

FIGURE AND TABLE LEGENDS

Figure 1.

Figure 1. Survival upon infection with *P. entomophila.*

Females (left) and males (right) of Control (grey) and BactOral (yellow) populations were orally infected with a culture of *P. entomophila* OD_{600}=50 (dashed lines) or a control solution (solid lines) for 24 hours, after which they were flipped onto clean food and their survival measured daily. We identify a significantly higher survival after infection in BactOral (in both sexes) as compared with Control, in a treatment-dependent manner. Results from multiple comparisons are shown as letters in the plots. Sample sizes were 40 uninfected and 180 infected individuals, for each sex per population.

1072　**Figure 2.**

1073

1074
1075
1076

Figure 2. Evolved and control populations eat and defecate equivalent amounts of

bacteria.

A) CAFE assay. Females of Control and BactOral populations were fed individually *P. entomophila*

solution through a graduated glass capillary and the amount of bacterial solution ingested (in µL) per

fly was registered at 2, 8 and 24 hours. Cumulative number of µL ingested per fly showed no statistically

significant difference in amounts ingested by either population at the different time-points. Linear

regressions between averaged values eaten by each population between time-points show a

comparable decrease in feeding as infection progresses (equations shown in plot for the periods, 2–8

hours (above lines) and 8–24 hours (below lines). 15 females were used from each population. **B)**

Quantification of defecated bacteria. Females (top) and males (bottom) from Control (grey) and

BactOral (yellow) populations were fed *P. entomophila* for 3 hours, after which the number *P.*

entomophila defecated was estimated after 6 hours and 12 hours. Lines connecting the two points

provide a sense of individual rates of live bacteria defecation within this time window. Linear regressions

between the two counts of the same fly show a significant decrease in the number of bacteria defecated

during the initial 6 hours and the following 12 hours by males of BactOral (shown with *). Additionally,

there is a significant difference in the number of BactOral flies that defecated below our detection

threshold (zero CFU counts) at 6 hours compared to 18 hours and compared to Control for both

timepoints. Samples sizes are shown at the top of each panel.

1095

1096 **Figure 3.**

1097
1098

Figure 3. Bacterial load dynamics.

Individual flies from Control (grey) and BactOral (yellow) were collected upon exposure to *P.*

entomophila at different timepoints for assessment of the number of Colony Forming Units (CFUs)

inside their bodies. After the end of the infection period (24 hours), flies were changed onto clean food

and collection continued until 52 hours post-infection. **A)** Log$_2$ of the number of CFUs found for females

(top) and males (bottom) of either Control (grey) or BactOral (yellow) populations, with the mean value

at each time-point evidenced in red (triangles for Control and circles for BactOral). There are no

detectable differences in bacterial loads between both populations until 20 hours of infection, when

BactOral starts to evidence a bigger decrease in bacterial counts. **B)** Cumulative number of individual

flies from which no CFUs were obtained increases as time progresses, with the BactOral reaching

significantly higher final values.

Numbers at the top of each panel correspond to the sample size at each timepoint, with BactOral

population in yellow and Control population in grey.

Figure 4.

1113

Figure 4. AMP expression differs between populations.

Guts from Control (grey) and BactOral (yellow) female flies (3 samples containing 10 guts, each) were

dissected before exposure ("Unc.") and at different timepoints after oral infection (8, 24 and 32 hours)

with *P. entomophila.* Relative gene expression, normalized to housekeeping gene *EIF2,* was performed

for a representative panel of 10 AMPs. We detected a significantly higher up-regulation in BactOral

compared to Control for *AttacinA* and *CecropinA* at 8-hours of infection. Results from these

comparisons are shown in the letters above the plots.

1121

 Figure 5.

 Figure 5. Control flies primed with heat-killed bacteria phenocopy BactOral survival to

 oral infection with *P. entomophila*.

 Control (grey) and BactOral (yellow) females (left) and males (right) were fed a solution of heat-killed

 bacteria for 2 hours before oral infection (dashed lines) or directly infected with live *P. entomophila* (full

 lines) for 24 hours, after which they were changed onto clean food and their survival measured daily.

 Previous exposure to heat-killed bacteria before infection increased survival in both populations, leading

 additionally Control females to phenocopy BactOral survival levels upon infection. Results from multiple

 comparisons are shown as letters in the plots. Numbers at the end of each trajectory correspond to the

 sample size, with BactOral population in yellow and Control population in grey.

Figure 6.

Figure 6. BactOral displays higher survival than Control upon immune system

activation without bacterial proliferation.

Daily survival was measured for Control (grey) and BactOral (yellow) populations fed a mixture of food

with heat-killed bacteria or PBS continuously. BactOral females and males exhibit a slower mortality

compared to Control flies until around day 8. This difference fades away for both sexes after that period.

Survival comparisons between populations and treatments are shown as lower-case letters. Sample

sizes were 150 individuals, for each sex, per population, per treatment.

Table 1.

parameter	mean	sd	hdi 3%	hdi 97%
Ref. k	3.143	0.033	3.080	3.204
Ref. h	-2.309	0.144	-2.577	-2.043
Ref. a	3.072	0.016	3.043	3.104
Ref. b	2.361	0.034	2.298	2.425
Sex k	-0.053	0.068	-0.172	0.068
Pop. k	-0.108	0.046	-0.193	-0.021
Sex h	-0.330	0.227	-0.769	0.089
Pop. h	0.042	0.182	-0.301	0.376
Sex a	-0.130	0.038	-0.195	-0.061
Pop. a	-0.002	0.024	-0.047	0.043
Sex b	-0.147	0.066	-0.276	-0.035
Pop. b	-0.285	0.053	-0.384	-0.187
ss	3.671	0.062	3.552	3.785

Table 1. Parameter estimates for the model sigmoidal function fitted to the time-course of bacterial counts.

A sigmoidal function of 4 parameters (a, k, h, b) was fitted to the bacterial dynamics time-course shown in Figure 3A, for quantification of differences between Control and BactOral CFU counts. Control females were used as reference on which to estimate the contribution of tested variables (variable "Sex": males and variable "Population": BactOral) on the parameters. Effects (Sex and Population) on parameters were considered significant when the 94% highest posterior density interval (hdi) did not include zero. The "ss" parameter represents the estimate of the standard deviation of the normal error structure.

1174 **Table 2.**

1175

1176

parameter	mean	sd	hdi 3%	hdi 97%
a_0 **Control**	-1.214	0.570	-2.262	-0.127
a_0 **BactOral**	-0.731	0.570	-1.816	0.324
a_0 **Female**	-0.984	0.569	-2.073	0.075
a_0 **Male**	-0.959	0.569	-2.064	0.079
a_1 **Female**	-0.781	0.071	-0.914	-0.645
a_1 **Male**	-0.061	0.069	-0.188	0.074
a_1 **Control**	-0.700	0.070	-0.833	-0.570
a_1 **BactOral**	-0.144	0.069	-0.268	-0.010
k_0 **Female**	0.086	0.097	-0.096	0.269
k_0 **Male**	0.085	0.095	-0.088	0.268
k_0 **Control**	0.066	0.097	-0.114	0.250
k_0 **BactOral**	0.104	0.095	-0.076	0.281
k_1 **Female**	-0.161	0.103	-0.354	0.030
k_1 **Male**	-0.232	0.103	-0.426	-0.037
k_1 **Control**	-0.161	0.101	-0.355	0.027
k_1 **BactOral**	-0.231	0.105	-0.426	-0.033
t_0 **Female**	0.152	0.119	-0.071	0.372
t_0 **Male**	0.253	0.116	0.031	0.462
t_0 **Control**	0.168	0.119	-0.047	0.397
t_0 **BactOral**	0.238	0.116	0.020	0.453

1177

1178

Table 2. Parameter estimates for the model logistic function fitted to the zero counts

of the time-course of bacterial counts.

An adapted logistic function of 4 parameters (a_0, a_1, k_0, k_1) was fitted to the progression of zero counts

shown in Fig. 3B), for quantification of bacterial clearance differences between Control and BactOral.

Single estimates were generated for the contribution of each variable (Sex: females and males;

Population: Control and BactOral) to the model parameters. Effects were considered significant when

was considered when the mean estimate ± standard deviation did not overlap for the same parameter

between each variable, which is the case for a_1 when considering Population and Sex. High-density

intervals of 3% and 97% show that all our estimated parameters have a high-density probability (i.e.

credibility) as they are all included in them.

1189

SUPPLEMENTARY FIGURE AND TABLE LEGENDS

1190

1191

1192 **Supplementary Figure 1.**

1193

Supplementary Figure 1. Longevity of BactOral and Control populations.

Females (full lines) and males (dashed lines) of Control (grey) and BactOral (yellow) populations, are shown. No

differences can be found between populations within sex (Uninfected Control females versus Uninfected BactOral

females: z.ratio= 1.214, p= 0.6180; Uninfected Control males versus Uninfected BactOral males: z.ratio= 0.615 ,

p= 0.9272).

Results from multiple comparisons are shown as lower-case letters in the plots.

Supplementary Figure 2.

1201

Supplementary Figure 2. No detectable differences in gut structure between Control

and BactOral after oral infection with *P. entomophila*.

Females of Control (grey) and BactOral (yellow) were fed normal food ("Unchallenged") or orally infected with *P.*

entomophila for 24 hours, after which their **A)** guts were dissected and photographed **B)**. Gut length (in mm) was

measured by drawing consecutive longitudinal transects along the guts thrice and averaging those

measurements to reach final length per sample. **A)** Exposure to bacteria shortens the guts significantly in both

populations (Control before infection *versus* Control 24 hours post-infection: t= 6.274, p<0.0001; BactOral before

infection *versus* BactOral 24 hours post-infection: t= 5.453, p<0.0001). No differences were detected in gut

lengths between populations within each timepoint (Without infection: BactOral *versus* Control – p= 0.6719; after

24 hours of infection – p= 0.9925), although a significant effect of treatment was observed in both Control

(Without infection *versus* after 24 hours of infection – p< 0.0001) and BactOral (p< 0.0001), showing that gut

shortening due to infection affected both regimes in a similar manner. **B)** Dissected guts of females from Control

and BactOral populations were photographed and scaled for posterior length quantification. Manual transects

were drawn longitudinally along the guts (as shown above) and total size of the cumulative transects was

calculated in FIJI. Three independent measurements were done on each gut and averaged to obtain a final

value per sample (Supp. Fig. 2 A)).

Samples sizes are shown at the top of the panel.

Supplementary Figure 3.

Supplementary Figure 3. AMP expression differences between populations under

homeostatic conditions.

Guts from Control (grey) and BactOral (yellow) female flies (3 samples containing 10 guts, each) were dissected

at different timepoints after oral infection with *P. entomophila* and relative quantification of gene expression was

performed. Expression levels of each gene in BactOral samples were normalized to expression levels of Control

samples. Log_2(FoldChange) of these differences were fitted on a linear model with Population as factor and

followed by multiple comparisons (*emmeans*) to uncover pairwise differences. Results from these comparisons

are shown in the letters above the plots. There are significantly lower levels of CecropinA (CecA) (*t*= 2.892, *p*=

0.0445) and significantly higher levels of *Drosomycin-like 5* (Drsl5) (*t*= -4.686 , *p*= 0.0094) being expressed in

guts of BactOral flies under unchallenged conditions.

1233   **Supplementary Figure 4.**

1234

Supplementary Figure 4. Principal Component Analysis on relative gene expression

levels in evolved populations.

(A) Relative AMP expression levels of Control and BactOral flies upon oral infection with *P. entomophila* showed

significant differences at various timepoints (see Fig. 4 and Supp. Fig. 2). **(B-C)** Relative influence of each AMP to

the establishment of PC1 and PC2 (see Supp. Fig. 3 A). Importantly, Cecropin A1 represents the largest contributor

to PC1 (**B**) whereas Drosomycin represents the largest contributor, although negatively correlated, to the variation

in PC2 (**C**).

1242

1243

1244   Supplementary Figure 5.

1245

1246

1247

1248   Supplementary Figure 5. Quantification of immunity-related gene expression upon

1249   artificial immune stimulation and infection.

RNA from whole-females from Control (grey) and BactOral (yellow) (4 samples containing 10 flies, each) was

extracted at different timepoints. Relative gene expression, normalized to housekeeping gene *rpl32,* was performed

for *Diptericin*, *Drosomycin* and *Relish*. Results from these comparisons are shown in the letters above the plots.

To test for individual pairwise differences, relative Log2 (FoldChange) expression levels of each gene were fitted

onto a linear model with Population and Timepoints as factors and followed by multiple comparisons (*emmeans*).

No differences are observed between non-challenged individuals or between individuals after 2 hours of exposure.

However, for Diptericin Unchallenged versus 2 HK show differences in both populations (Control: t= -4.778 , p=

0.0001; BactOral: t= -3.122, p= 0.0052).

SUPPLEMENTARY TABLES

Supplementary Table 1:

A)
Model: coxme(Surv(time, status) ~ Population * Treatment * Sex + (1|Replicate))

fixed coefficients	coef	exp(coef)	se(coef)	*z*	*p*
Population-B	-0.005	0.995	1.418	0.00	1.0e+00
Treatment-Pe	5.022	151.786	1.008	4.98	**6.2e-07**
Sex-m	0.712	2.038	1.229	0.58	5.6e-01
Population-B:Treatment-Pe	-1.250	0.287	1.425	-0.88	3.8e-01
Population-B: Sex-m	0.916	2.499	1.650	0.56	5.8e-01
Treatment-Pe:Sex-m	-1.542	0.214	1.236	-1.25	2.1e-01
Population-B:Treatment-Pe:Sex-m	-1.452	0.234	1.665	-0.87	3.8e-01
Random effects					
Group	**Variable**	**Std Dev**	**Variance**		
Replicate	Intercept	0.106	0.011		

B)
Analysis of Deviance Table (Type III tests)

	Df	Chisq	Pr(>Chisq)
Population	1	0.0000	0.997
Treatment	1	24.845	**6.214e-07 *****
Sex	1	0.308	0.563
Population:Treatment	1	0.770	0.380
Population:Sex	1	0.302	0.579
Treatment:Sex	1	1.558	0.212
Population:Treatment:Sex	1	0.760	0.383

Supplementary Table 1. Statistical analysis of survival after oral infection with *P. entomophila* (immunocompetence, shown in Fig. 1).

A) Model estimates and **B)** ANOVA estimates for survival analysis testing immunocompetence of Control and BactOral populations.

Supplementary Table 2:

A)
Model: coxph(Surv(time, status) ~ Population * Sex)

	coef	exp(coef)	se(coef)	z	Pr(>|z|)
Population-B	0.271	1.312	0.224	1.214	0.225
Sex-m	1.069	2.912	0.215	4.970	**6.7e-07 *****
Population-B: Sex-m	-0.389	0.677	0.295	-1.321	0.186

B)
Analysis of Deviance Table (Type III tests)

	Chisq	Df	Pr(>Chisq)
Population	1.489	1	0.222
Sex	26.067	1	**3.298e-07 *****
Population:Sex	1.759	1	0.185

Supplementary Table 2. Statistical analysis of survival under uninfected conditions (Longevity, shown in Supp. Fig. 1).

A) Model estimates and **B)** ANOVA estimates for survival analysis testing longevity of Control and BactOral populations.

Supplementary Table 3:

A)
Model: glmer(µL.eaten ~ Population*factor(Timepoint) + (1|Replicate fly), family = "Gamma")

Fixed effects:	Estimate	Std. Error	t-value	Pr(>|z)
(Intercept)	1.56	0.104	14.940	**< 2e-16 *****
Population-BactOral	-0.015	0.140	-0.104	0.918
Timepoint-8	-0.836	0.109	-7.657	**1.91e-14*****
Timepoint-24	-1.156	0.102	-11.290	**< 2e-16 *****
Population-B:Timepoint-8	-0.028	0.153	-0.181	0.856
Population-B:Timepoint-24	-0.005	0.144	-0.035	0.972
Random effects:				
Groups	**Name**	**Variance**	**Std. Dev.**	
Replicate fly	(Intercept)	0..004	0.065	
Residual	--	0.059	0.243	

B)
Analysis of Deviance Table (Type III tests)

	Chisq	Df	Pr(>Chisq)
(Intercept)	223.2	1	**<2e-16 *****
Population	0.0107	1	0.918
Timepoint	153.151	2	**<2e-16 *****
Population:Timepoint	0.110	2	0.947

Supplementary Table 3. Analysis of the cumulative amounts of bacteria ingested by evolved populations (CAFE assay, shown in Fig. 2 A).

A) Model estimates and **B)** ANOVA estimates for the generalized linear model testing the effects of population and timepoint on ingested bacterial mixture.

Supplementary Table 4:

Model: zeroinfl(formula = avg_geom_cfus ~ Population * Sex * factor(Timepoint) | Population * factor(Timepoint), dist = "negbin")

Count model coefficients (negbin with log link)				
	Estimate	**Std. Error**	**z value**	**Pr(>|z|)**
(Intercept)	13.721	0.482	28.447	**<2e-16 *****
Population-C	0.816	0.616	1.324	0.185
Sex-m	0.047	0.647	0.073	0.942
Timepoint-18	-0.251	0.699	-0.359	0.720
Population-C:Sex-m	-0.910	0.840	-1.083	0.279
Population-C:Timepoint-18	-0.579	0.905	-0.640	0.522
Sex-m:Timepoint-18	-0.950	0.911	-1.043	0.297
Population-C:Sex-m:Timepoint-18	2.171	1.204	1.803	0.071
Log(theta)	-1.031	0.106	-9.700	**<2e-16 *****
Zero-inflation model coefficients (binomial with logit link):				
	Estimate	**Std. Error**	**z value**	**Pr(>|z|)**
(Intercept)	-0.828	0.351	-2.357	**0.018 ***
Population-C	-3.004	1.248	-2.407	**0.016***
Timepoint-18	-18.771	3347.432	-0.006	0.996
Population-C:Timepoint-18	18.876	3347.433	0.006	0.996

Supplementary Table 4. Analysis of quantities of defecated bacteria by evolved populations (CFUs, shown in Fig. 2 B)).

A) Model estimates and for the zero-inflated model testing the effects of population, sex and timepoint on amount of defecated live bacteria and instances of non- measurable bacteria (zeros).

Supplementary Table 5:

A)
Model: lmer(Lenght ~ Population*Treatment + (1|Round), REML=TRUE)

Fixed effects	Estimate	Std. Error	df	t-value	Pr(>|t|)
(Intercept)	6.794	0.202	61	33.565	**< 2e-16 *****
Population-B	-0.314	0.276	61	-1.136	0.260
Treatment-Pe	-2.084	0.320	61	-6.513	**1.58e-08 *****
Population-B:Treatment-Pe	0.407	0.436	61	0.933	0.354
Random effects					
Groups	Name	Variance	Std Dev		
Round	(Intercept)	0	0		
Residual	---	0.7376	0.8588		

B)
Analysis of Deviance Table (Type III Wald chisquare tests)

	Chisq	Df	Pr(>Chisq)
(Intercept)	1126.6148	1	**< 2.2e-16 *****
Population	1.2914	1	0.256
Treatment	42.4139	1	**7.386e-11 *****
Population:Treatment	0.8710	1	0.351

Supplementary Table 5. Analysis of total gut lengths of females of evolved populations before and after 24 hours of infection (shown in Supp. fig. 1 A)).

A) Model estimates and **B)** ANOVA estimates for the linear model testing the effects of population and treatment on total length of guts.

Supplementary Table 6:

A)
Model: lm(LogFold_gene ~ Population * factor(Timepoint))
AttacinA

Coefficients	Estimate	Std. Error	t-value	Pr(>|t|)
(Intercept)	-3.333e-10	8.287e-01	0.00	1.00
Population-B	3.333e-10	1.172e+00	0.00	1.00
Timepoint-8	5.914e+00	1.172e+00	5.047	0.000119 ***
Timepoint-24	6.171e+00	1.172e+00	5.266	7.69e-05 ***
Timepoint-32	7.375e+00	1.172e+00	6.293	1.07e-05 ***
Population-B: Timepoint-8	3.004e+00	1.657e+00	1.813	0.088712 .
Population-B: Timepoint-24	-1.412e+00	1.657e+00	-0.852	0.406965
Population-B: Timepoint-32	-1.950e+00	1.657e+00	-1.177	0.256541

B)
Anova Table (Type III tests)

	Sum Sq	Df	F-value	Pr(>F)
(Intercept)	0.0	1	0.00	1.0
Population	0.0	1	0.00	1.0
Timepoint	98.317	3	15.908	**4.647e-05 ***
Population:Timepoint	22.181	3	3.589	**0.03718 ***
Residuals	32.962	16		

C)
BomaninS2

Coefficients	Estimate	Std. Error	t-value	Pr(>|t|)
(Intercept)	-5.881e-16	8.017e-01	0.0	1.0
Population-B	1.358e-15	1.134e+00	0.0	1.0
Timepoint-8	1.563e+00	1.134e+00	1.378	0.187
Timepoint-24	2.062e+00	1.134e+00	1.818	0.088.
Timepoint-32	**6.049e+00**	1.134e+00	5.335	**6.7e-05 ***
Population-B: Timepoint-8	5.023e-01	1.603e+00	0.313	0.758
Population-B: Timepoint-24	2.832e-01	1.603e+00	0.177	0.862
Population-B: Timepoint-32	-4.928e+00	1.603e+00	-3.074	**0.007 **

D)
Anova Table (Type III tests)

	Sum Sq	Df	F-value	Pr(>F)
(Intercept)	0.0	1	0.0	1.0
Population	0.0	1	0.0	1.0
Timepoint	59.677	3	10.3161	**0.0005071 ***
Population:Timepoint	30.495	3	5.2715	**0.0101493 ***
Residuals	30.852	16		

E)
BomaninS3

Coefficients	Estimate	Std. Error	t-value	Pr(>|t|)
(Intercept)	-8.158e-16	7.573e-01	0.0	1.0
Population-B	1.282e+00	1.071e+00	1.197	0.249
Timepoint-8	-1.355e-17	1.071e+00	0.0	1.0
Timepoint-24	1.124e-15	1.071e+00	0.0	1.0
Timepoint-32	1.014e-15	1.071e+00	0.0	1.0
Population-B: Timepoint-8	7.635e-02	1.515e+00	0.050	0.960
Population-B: Timepoint-24	2.302e-01	1.515e+00	0.152	0.881
Population-B: Timepoint-32	-4.322e+00	1.515e+00	-2.853	**0.012***

F)
Anova Table (Type III tests)

	Sum Sq	Df	F-value	Pr(>F)
(Intercept)	0.0	1	0.0	1.0
Population	2.467	1	1.434	0.249
Timepoint	0.0	3	0.0	1.0
Population:Timepoint	22.057	3	4.273	**0.021 ***
Residuals	27.529	16		

G)
CecropinA1

Coefficients	Estimate	Std. Error	t-value	Pr(>|t|)
(Intercept)	-1.813e-16	8.111e-01	0.0	1.0
Population-B	-2.107e+00	1.147e+00	-1.837	0.085.
Timepoint-8	5.263e-16	1.147e+00	0.0	1.0
Timepoint-24	-3.400e-16	1.147e+00	0.0	1.0
Timepoint-32	-5.934e-16	1.147e+00	0.0	1.0
Population-B: Timepoint-8	3.480e+00	1.622e+00	2.146	**0.048 ***
Population-B: Timepoint-24	-7.721e-01	1.622e+00	-0.476	0.641
Population-B: Timepoint-32	-5.195e+00	1.622e+00	3.202	**0.006****

H)
Anova Table (Type III tests)

	Sum Sq	Df	F-value	Pr(>F)
(Intercept)	0.0	1	0.0	1.0
Population	6.660	1	3.375	0.085.
Timepoint	0.0	3	0.0	1.0
Population:Timepoint	57.223	3	9.665	**0.0007*****
Residuals	31.576	16		

I)
<u>Defensin</u>

Coefficients	Estimate	Std. Error	t-value	Pr(>|t|)
(Intercept)	-4.532e-16	1.167e+00	0.0	1.0
Population-B	-1.155e+00	1.651e+00	-0.700	0.4942
Timepoint-8	-3.333e-10	1.651e+00	0.0	1.0
Timepoint-24	-3.333e-10	1.651e+00	0.0	1.0
Timepoint-32	6.636e-16	1.651e+00	0.0	1.0
Population-B: Timepoint-8	3.468e+00	2.335e+00	1.485	0.1569
Population-B: Timepoint-24	-1.377e-01	2.335e+00	-0.059	0.9537
Population-B: Timepoint-32	-4.353e+00	2.335e+00	-1.865	0.0807 .

J)
Anova Table (Type III tests)

	Sum Sq	Df	F-value	Pr(>F)
(Intercept)	0.0	1	0.0	1.0
Population	2.001	1	0.490	0.494
Timepoint	0.0	3	0.0	1.0
Population:Timepoint	46.098	3	3.759	**0.032 ***
Residuals	65.404	16		

K)
<u>Diptericin</u>

Coefficients	Estimate	Std. Error	t-value	Pr(>|t|)
(Intercept)	2.617e-15	1.001e+00	0.0	1.0
Population-B	3.333e-10	1.416e+00	0.0	1.0
Timepoint-8	5.259e+00	1.416e+00	3.713	**0.002 ** **
Timepoint-24	4.907e+00	1.416e+00	3.465	**0.003 ** **
Timepoint-32	7.187e+00	1.416e+00	5.075	**0.000113 *** **
Population-B: Timepoint-8	2.307e+00	2.003e+00	1.152	0.266
Population-B: Timepoint-24	1.076e-01	2.003e+00	0.054	0.958
Population-B: Timepoint-32	-3.701e+00	2.003e+00	-1.848	0.083 .

L)
Anova Table (Type III tests)

	Sum Sq	Df	F-value	Pr(>F)
(Intercept)	0.0	1	0.0	1.0
Population	0.0	1	0.0	1.0
Timepoint	84.314	3	9.3416	**0.0008357 *** **
Population:Timepoint	27.922	3	3.0937	0.057 .
Residuals	48.137	16		

M)
Drosomycin

Coefficients	Estimate	Std. Error	t-value	Pr(>|t|)
(Intercept)	-3.333e-10	6.719e-01	0.0	1.0
Population-B	3.333e-10	9.503e-01	0.0	1.0
Timepoint-8	3.984e-01	9.503e-01	0.419	0.681
Timepoint-24	2.392e-01	9.503e-01	0.252	0.804
Timepoint-32	3.783e+00	9.503e-01	3.981	**0.001 ****
Population-B: Timepoint-8	-1.971e-01	1.344e+00	-0.147	0.885
Population-B: Timepoint-24	4.762e-01	1.344e+00	0.354	0.728
Population-B: Timepoint-32	-4.556e+00	1.344e+00	-3.390	**0.004****

N)
Anova Table (Type III tests)

	Sum Sq	Df	F-value	Pr(>F)
(Intercept)	0.0	1	0.0	1.0
Population	0.0	1	0.0	1.0
Timepoint	28.931	3	7.1198	**0.003 ****
Population:Timepoint	24.678	3	6.0731	**0.006 ****
Residuals	21.672	16		

O)
Drosomycin-like2

Coefficients	Estimate	Std. Error	t-value	Pr(>|t|)
(Intercept)	1.807e-15	9.335e-01	0.0	1.0
Population-B	3.333e-10	1.320e+00	0.0	1.0
Timepoint-8	5.925e+00	1.320e+00	4.488	**0.0004 *****
Timepoint-24	6.968e+00	1.320e+00	5.278	**7.50e-05 *****
Timepoint-32	8.724e+00	1.320e+00	6.609	**6.01e-06 *****
Population-B: Timepoint-8	1.187e+00	1.867e+00	0.636	0.534
Population-B: Timepoint-24	1.857e+00	1.867e+00	0.995	0.335
Population-B: Timepoint-32	-3.125e+00	1.867e+00	-1.674	0.114

P)
Anova Table (Type III tests)

	Sum Sq	Df	F-value	Pr(>F)
(Intercept)	0.0	1	0.0	1.0
Population	0.0	1	0.0	1.0
Timepoint	128.832	3	16.428	**3.843e-05 *****
Population:Timepoint	21.934	3	2.797	0.074 .
Residuals	41.826	16		

Q)
Drosomycin-like3

Coefficients	Estimate	Std. Error	t-value	Pr(>|t|)
(Intercept)	5.143e-16	6.851e-01	0.0	1.0
Population-B	1.147e-15	9.688e-01	0.0	1.0
Timepoint-8	4.849e+00	9.688e-01	5.005	**0.0001***
Timepoint-24	6.738e+00	9.688e-01	6.955	**3.24e-06 ***
Timepoint-32	8.622e+00	9.688e-01	8.900	**1.36e-07 ***
Population-B: Timepoint-8	1.678e+00	1.370e+00	1.225	0.238
Population-B: Timepoint-24	1.297e+00	1.370e+00	0.947	0.358
Population-B: Timepoint-32	-3.095e+00	1.370e+00	-2.259	**0.038 ***

R)
Anova Table (Type III tests)

	Sum Sq	Df	F-value	Pr(>F)
(Intercept)	0.0	1	0.0	1.0
Population	0.0	1	0.0	1.0
Timepoint	123.465	3	29.230	**9.967e-07 ***
Population:Timepoint	21.108	3	4.9972	**0.01238 ***
Residuals	22.528	16		

S)
Drosomycin-like5

Coefficients	Estimate	Std. Error	t-value	Pr(>|t|)
(Intercept)	5.815e-16	6.871e-01	0.0	1.0
Population-B	3.333e-10	9.717e-01	0.0	1.0
Timepoint-8	2.630e+00	9.717e-01	2.706	**0.016 ***
Timepoint-24	5.041e+00	9.717e-01	5.188	**8.98e-05 ***
Timepoint-32	7.878e+00	9.717e-01	8.108	**4.66e-07 ***
Population-B: Timepoint-8	1.179e+00	1.374e+00	0.858	0.404
Population-B: Timepoint-24	7.405e-01	1.374e+00	0.539	0.597
Population-B: Timepoint-32	-4.145e+00	1.374e+00	-3.016	**0.008 ****

T)
Anova Table (Type III tests)

	Sum Sq	Df	F-value	Pr(>F)
(Intercept)	0.0	1	0.0	1.0
Population	0.0	1	0.0	1.0
Timepoint	101.853	3	23.972	**3.678e-06 ***
Population:Timepoint	26.821	3	6.313	**0.00497 ****
Residuals	22.660	16		

Supplementary Table 6. Analysis of relative gene expression levels in guts of females of evolved populations at several timepoints throughout infection (shown in Fig. 4).

Model estimates and ANOVA estimates for the linear models testing the effects of population and timepoint on expression levels of multiple AMPs during the course of oral bacterial infection.

Supplementary Table 7

Primer name	5'-Sequence-3'
rpl32 F01	CCAGTCGGATCGATATGCTAA
rpl32 R01	GTTCGATCCGTSACCGATGT
eIF(-2alpha) F01	TCGCATCAACCTGATAGCAC
eIF(-2alpha) R01	ATCGTACTCGCTGGTCTTGG
relish F01	ACAAAGGACGAGGATTCCGTGG
relish R01	TGATTCAGCAGCGAACAGAGCC
drosomycin F01	CTCTTCGCTGTCCTGATGCTGG
drosomycin R01	GACAGGTCTCGTTGTCCCAGAC
diptericin F01	TACTTTGGCTTATCCGATGCCC
diptericin R01	GTGCTGTCCATATCCTCCATTC
cecropin a1 F01	GTTTTCGTCGCTCTCATTCTGG
cecropin a1 R01	CTTGTTGAGCGATTCCCAGTC
defensin F01	TCGCTTTTGCTCTGCTTGCTTG
defensin R01	GTTGCAGTAGCCGCCTTTGAAC
drsl2 F01	CCTGGCTGTGATGACAATTGTTG
drsl2 R01	ACTGATATGTCCCTCCTCCTTGC
drsl3 F01	TCCTAATGGAGGCCAACACTG
drsl3 R01	TCCCTCCTCAATGCAGAGACG
drsl5 F01	TCCTGGCTGTGATGACCATCTTC
drsl5 R01	ACTGGATCGTCCTTCCTCCTTG
boms2 F01	AGTCGTCACCGTCTTTGTGTT
boms2 R01	GCAGTATTTGCAGTCCCCGTTG
boms3 F01	CTATCACTCGCCTTCGTTTTGG
boms3 R01	TCACATTGCAGACGCGGCAATC
attacin a1 F01	CATTACCGAGGCACTTCCCAC
attacin a1 R01	GAAAACCTGACCCACCACATTG

Supplementary Table 7. Primer sequences used in Real-Time quantitative PCRs (shown in Fig. 4).

The first four genes listed (shown in bold and italics) were used as reference (house-keeping) for relative quantifications.

Supplementary Table 8:

A)
Model: coxme(Surv(time,status)~Population*Treatment*Sex+(1|Block/Replicate))

Fixed coefficients	coef	exp(coef)	se(coef)	z	p
Population-B	-1.065	0.345	0.156	-6.82	**8.8e-12**
Treatment-HK	-1.324	0.266	0.132	-10.01	**0.0e+00**
Sex-m	-0.195	0.823	0.130	-1.50	1.3e-01
Population-B:Treatment-HK	-0.115	0.891	0.239	-0.48	6.3e-01
Population-B: Sex-m	-0.792	0.453	0.256	-3.09	**2.0e-03**
Treatment-HK: Sex-m	0.382	1.465	0.184	2.08	**3.8e-02**
Population-B:Treatment-HK:Sex-m	-0.172	0.842	0.391	-0.44	6.6e-01
Random effects					
Group	Variable	Std Dev	Variance		
Block/Replicate	(Intercept)	0.152	0.023		
Block	(Intercept)	0.173	0.030		

B)
Analysis of Deviance Table (Type III tests)

	Df	Chisq	Pr(>Chisq)
Population	1	46.577	**8.808e-12 *** **
Treatment	1	100.214	**< 2.2e-16 *** **
Sex	1	2.238	0.135
Population:Treatment	1	0.233	0.639
Population:Sex	1	9.553	**0.002 ** **
Treatment:Sex	1	4.308	**0.038 * **
Population:Treatment:Sex	1	0.194	0.659

Supplementary Table 8. Statistical analysis of survival of evolved populations after priming with heat-killed bacteria (shown in Fig. 5).

A) Model estimates and **B)** ANOVA estimates for survival analysis testing the effect of priming on infection in Control and BactOral.

Supplementary Table 9:
A)
Model: coxme(Surv(time,status)~Population*Treatment*Sex+(1|Replicate))

Fixed coefficients	coef	exp(coef)	se(coef)	z	p
Population-B	-1.137	0.321	0.817	-1.39	0.160
Treatment-Pe WT	-5.354	211.507	0.430	12.45	**0.000**
Sex-m	1.032	2.807	0.479	2.16	**0.031**
Population-B: Treatment-Pe WT	0.359	1.432	0.824	0.44	0.660
Population-B: Sex-m	0.917	2.502	0.898	1.02	0.310
Treatment-Pe WT: Sex-m	0.141	1.151	0.494	0.28	0.780
Population-B: Treatment-Pe WT: Sex-m	-1.348	0.260	0.913	-1.48	0.140
Random effects					
Group	**Variable**	**Std Dev**	**Variance**		
Replicate	(Intercept)	4.149e-03	1.721e-05		

B)
Analysis of Deviance Table (Type III tests)

	Df	Chisq	Pr(>Chisq)
Population	1	1.937	0.164
Treatment	1	154.932	**< 2e-16 *****
Sex	1	4.648	**0.0312 ***
Population:Treatment	1	0.1897	0.663
Population:Sex	1	1.044	0.307
Treatment:Sex	1	0.081	0.776
Population:Treatment:Sex	1	2.178	0.140

Supplementary Table 9. Statistical analysis of survival of evolved populations after prolonged exposure to heat-killed bacteria (shown in Fig. 6).

A) Model estimates and **B)** ANOVA estimates for survival analysis testing the effects of population, treatment and sex on survival to prolonged exposure to heat-killed *P. entomophila*, for 14 days.