1
2
3 **Targeted mutagenesis of *Medicago truncatula Nodule-specific Cysteine-rich* (*NCR*) genes**
4 **using the *Agrobacterium rhizogenes*-mediated CRISPR/Cas9 system**
5
6
7

8 Berivan Güngör[1,2*], János Barnabás Biró[1*], Ágota Domonkos[2], Beatrix Horváth[2], Péter Kaló[1,2]
9
10
11
12 1 Institute of Plant Biology, Biological Research Centre, Eötvös Lóránd Research Network,
13 Szeged, Hungary
14 2 Institute of Genetics and Biotechnology, Hungarian University of Agriculture and Life
15 Sciences, Gödöllő, Hungary
30 * B.G. and J.B.B. contributed equally to this work.
31
32 Author for correspondence: Péter Kaló, Email: kalo.peter@brc.hu
33
34 ORCID IDs:
35
36 Peter Kalo: 0000-0002-0404-8904
37 Berivan Güngör: 0000-0002-5612-1130
38 János Barnabás Biró: 0000-0001-8851-0387
39 Agota Domonkos: 0000-0003-4017-0605
40 Beatrix Horvath: 0000-0001-8499-568X
41
42
43

Abstract

The host-produced nodule specific cysteine-rich (NCR) peptides control the terminal

differentiation of endosymbiotic rhizobia in the nodules of IRLC legumes. Although the

Medicago truncatula genome encodes about 700 NCR peptides, only few of them have been

proved to be crucial for nitrogen-fixing symbiosis. In this study, we applied the CRISPR/Cas9

gene editing technology to generate knock-out mutants of *NCR* genes for which no genetic or

functional data were previously available. We have developed a workflow to analyse the

mutation and the symbiotic phenotype of individual nodules formed on *Agrobacterium*

rhizogenes-mediated transgenic hairy roots. The selected *NCR* genes were successfully edited

by the CRISPR/Cas9 system and nodules formed on knockout hairy roots showed wild type

phenotype indicating that peptides NCR068, NCR089, NCR128 and NCR161 are not

essential for symbiosis between *M. truncatula* Jemalong and *Sinorhizobium medicae*

WSM419. We regenerated stable mutants edited for the *NCR068* from hairy roots obtained

from *A. rhizogenes*-mediated transformation. The analysis of the symbiotic phenotype of

stable *ncr068* mutants showed that peptide NCR068 is not required for symbiosis with *S.*

meliloti strains 2011 and FSM-MA either. Our study reports that gene editing can help to

elicit the role of particular NCRs in symbiotic nitrogen fixation.

Introduction

Most legumes establish nitrogen-fixing symbiotic interaction with soil bacteria termed rhizobia. Compatible interaction between the host legumes and their rhizobia partner induces the formation of a specialised organ on the root, referred to as nodules. Rhizobia colonize the symbiotic cells of the nodules and differentiate into their symbiotic form, the bacteroids which convert the atmospheric nitrogen into ammonia[1]. The indeterminate nodules formed on the roots of temperate legumes, such as the model legume *Medicago truncatula*, possess a persistent apical meristem[2]. The continuous activity of the meristem (zone I) produces cells creating a developmental gradient in the nodules, whereupon these nodules are cylindrical in shape. The new cells mature, many of them become infected in zone II (infection zone) wherein bacteria are engulfed by a plant-derived membrane forming new organelles, the symbiosomes. Infected host cells and rhizobia differentiate in interzone (IZ) and the reduction of nitrogen takes place in zone III (nitrogen fixation zone). The completely developed and effectively nitrogen-fixing nodules produce leghemoglobin which gives wild type nodules their pink colour.

Rhizobia in the nodules of legumes belonging to the inverted repeat–lacking clade (IRLC), including *M. truncatula*, and some Dalbergoid species undergo irreversible differentiation mediated by nodule-specific cysteine-rich (NCR) peptides[3–5]. These host produced peptides contain four or six cysteine residues in conserved positions and possess signal peptides which target them to the symbiosomes through a nodule-specific protein secretory pathway[6,7] where they govern the transition of rhizobia into bacteroids. The *M. truncatula* genome encodes a large family of NCR peptides with more than 700 members[8,9]. Although these high number of peptides show remarkable amino acid sequence variation, their function was initially presumed redundant. Nevertheless, recent studies have identified some peptides, NCR169, NCR211, NCR247, NCR343 and NCR-new35, crucial for the bacteroid differentiation and persistence in *M. truncatula* nodules but also found peptides (NCR178, NCR341, NCR344 and NCR345) which are not essential for nitrogen-fixing symbiosis[10–15]. The identification of most of these essential peptides was random during the forward genetic analyses of mutagenized *M. truncatula* populations. Therefore, the reverse genetic approach provides a comprehensive and targeted way to analyse the function of the *NCR* gene family members. The recently developed CRISPR/Cas9 (Clustered Regularly Interspaced Short Palindromic Repeats/CRISPR-associated protein 9) system is a simple and precise tool for targeted gene editing and analyse gene function in various plants[16]. Several reports described the application

of CRISPR/Cas9 gene editing system in legume species (soybean[17]; peanut[18]; pea[19]; *Lotus*
japonicus[20] including *M. truncatula*[21–24]. Unfortunately, most legume species are largely
recalcitrant to somatic embryogenesis and *in vitro* plant regeneration, therefore generating
stable transformant legume plants using the *Agrobacterium tumefaciens*-mediated
transformation system is a cumbersome and time-consuming process. In contrast, the *A.*
rhizogenes-mediated hairy root transformation does not need complete plant regeneration,
therefore it provides an efficient tool for a fast and large-scale functional analysis of root-
related traits, such as symbiotic nodule development. Despite its robustness, the *A.*
rhizogenes-mediated hairy root transformation has some disadvantages. For instance, the
transformation efficiency varies, and gene transfer is not achieved in all root cells[25], or the
emerged hairy roots develop from multiple cells with different gene editing events which can
result in mosaicism of transformed and untransformed cells or even multiple genotypes. This
could cause trouble during gene editing of a wild-type genome to generate knockouts because
a small amount of residual wild-type cells could affect the phenotype to be analysed.
Therefore, the removal of untransformed or partially transformed roots and analysing the
genotype of the whole hairy root is particularly important.
The CRISPR/Cas9 gene editing system, have been applied successfully to knockout and
analyse the requirement of peptides NCR247, NFS1 and NFS2[12,26,27] using *A. rhizogenes*-
mediated hairy root transformation. In these reports, the targeted *NCR* genes were selected
based on preliminary genetic or functional studies. In this study, we optimized the
CRISPR/Cas9 system using hairy root transformation to generate mutation in selected *NCR*
genes and developed a workflow to conduct their functional analysis for symbiosis.
119
120

Results
Testing the efficiency of a modified vector for CRISPR/Cas9 gene editing system to
target *NCR169*
To analyse the function of *NCR* genes, the CRISPR/Cas9 gene editing technology was used to
generate composite *Medicago truncatula* plants mutated in hairy roots. In order to monitor the
presence of CRISPR constructs, delivered into the genome of *M. truncatula* root cells using
the *Agrobacterium rhizogenes*-mediated hairy-root system[28], the gene coding for the DsRed
fluorescent transformation marker was introduced into the CRISPR/Cas9 plant transformation
vector pKSE401 resulting in the vector pKSE401-RR. The constitutively active red
fluorescent marker allowed us to remove non-transgenic roots at the early stages to promote

the growth of transgenic hairy roots. To test the efficiency of targeted mutagenesis of
pKSE401-RR, we designed three different single guide RNA (sgRNA) constructs targeting
the *NCR169* gene, previously proved to be essential for effective nitrogen-fixing symbiosis[10].
Each sgRNA (NCR169a, b and c) was designed directing to the first half of the coding
sequence of *NCR169* but the initial 5' nucleotides for the protospacer sequences of NCR169b
and NCR169c were C instead of the optimal G(/A) which is preferred by the AtU6 promoter
(Fig. S1a;[29]). Nevertheless, we used these two suboptimal sgRNA constructs as well because
the short, 189 bp long coding sequence of *NCR169* constrained the available optimal target
sequences. The three constructs carrying each sgRNAs were introduced into roots of wild-
type *M. truncatula* 2HA genotype using *A. rhizogenes*-mediated hairy root transformation. At
least ten transformed roots for each construct, detected by the DsRed fluorescent marker, were
sampled and used for genomic DNA isolation. The region surrounding the targeted sequence
in *NCR169* was amplified by PCR. The sequence analysis of the amplicons detected only
wild-type genotypes on all the NCR169b-transformed roots and roots transformed with the
construct NCR169c were heterogeneous of wild-type and various mutant genotypes.
However, the construct NCR169a generated mutations at higher efficiency and beside
chimeric roots having both wild-type and distinct mutations, we could identify roots with
homozygous or biallelic mutations for *NCR169* (Fig. S1d). The results indicated that the
presence of G at the 5' end of the protospacer sequence (G(/A)N19NGG) is crucial and using
C in this position greatly reduced the efficiency of genome editing which restricts the number
of possible target sites in the case of small genes such as *NCR*s. In addition, the sequence
analysis of amplicons containing the targeted region revealed that the sequences obtained by
Sanger method composed of multiplex sequence traces around the break site (Fig. S1b). An
online tool, TIDE, enabling the decomposition of mixed sequence signals was applied[30] that
predicted the types of the mutation and the size of the deletion or insertion. For instance, the
decomposition of one of the amplicons of NCR169c disclosed the mixture of wild-type and
CRISPR/Cas9 edited sequences containing a 7 bp deletion and/or a one bp insertion (Fig.
S1c).
In agreement with a previous report[22], we also found that different genome editing events
occurred in distinct hairy roots of a single plant which necessitates the analysis of individual
nodules separately. These findings led us to the conclusions that the effectiveness of genome
editing and the reliability of the identification of the nature of genome editing events needed
to be improved in our system. Therefore (i) we replaced the Arabidopsis U6-26 promoter with
the *M. truncatula* U6 (MtU6.6) promoter driving sgRNA synthesis, (ii) we developed a

preparation procedure allowing phenotypic and genotypic analysis of individual nodules, and

(iii) applied an NGS amplicon sequencing approach to assess allele distribution in individual

nodules.

Inducing mutations in *NCR169* and *NCR-new35* using the optimized CRISPR/Cas9-

mediated genome editing system

Previous studies found that the *M. truncatula* U6 promoter enhanced the expression efficiency

of guide or hairpin RNAs in legume species[21,31]. To increase the efficacy of targeted

mutagenesis by CRISPR/Cas9, the Arabidopsis U6-26 promoter was replaced with MtU6.6 in

the pKSE401-RR resulting in the vector pKSE466-RR. We inserted the sgRNAs NCR169a

and another one, targeting the *NCR-new35*, of which requirement for effective symbiosis has

been recently demonstrated[13], into pKSE466-RR, respectively. To validate the efficiency of

the pKSE466-RR vector construct, the empty vector and pKSE466-RR-NCR169a were

introduced into wild-type *M. truncatula* 2HA plants using *A. rhizogenes*-mediated hairy root

transformation and plants were inoculated with *S. medicae* WSM419. Nodules developed on

DsRed fluorescent transformed roots were harvested 3-4 weeks post inoculation (wpi) with

rhizobia, individually fixed and sections were stained with nucleic acid-binding dye SYTO13.

Following image capturing, genomic DNA was purified from nodule sections and the regions

surrounding the targeted sequences were amplified by PCR.

The elongated nodules developed on empty vector-transformed roots were pink and showed

the characteristic zonation of indeterminate nodules (Fig. 1a and d). The nodules on the roots

of *M. truncatula* 2HA transformed with the pKSE466-RR-NCR169a were either pink and

elongated, suggesting that these nodules were functional wild-type (Fig. 2g, i and k), or white

undeveloped, implying that these were ineffective symbiotic nodules (Fig. 1b and 2b, d and f).

To determine the nature of the mutations at the targeted region, the sequence analysis of the

amplicons, generated from randomly selected elongated and undeveloped nodules, was

carried out. The targeted amplicon sequences were determined using a next-generation

sequencing (NGS) approach and the quantitative allele composition and the verification of

genome editing mutations were analysed using the online tool CRISPResso2[32]. The analysis

revealed that all the white undeveloped nodules carried few base pair biallelic insertions or

deletions at a high ratio and the very few reads of wild-type alleles identified in these nodules

were probably derived from sporadic wild-type cells or the trace amount of wild-type

contamination (Fig. 1b, Fig. 2a-f). The morphology and bacterial occupancy of these

SYTO13-stained undeveloped nodules were very similar to the nodulation phenotype of the

deletion mutant *dnf7-2* showing no fluorescence in the nodule region proximal to the root

(Fig. 1e). In addition, these plants, compared with empty-vector-transformed plants, showed

the symptoms of nitrogen starvation (yellowish leaves and stunted growth habit) indicating

the defect of symbiotic nitrogen fixation of these nodules (Fig. 1b). Contrary to these, the

sequence analysis of the randomly selected elongated pink nodules showed that they either

carried almost exclusively wild-type alleles (Fig. 2g and h) or they were heterozygous for

example for a 4 bp deletion and the wild-type alleles (Fig. 2i and j). We also found a wild-

type nodule that predominantly carried a homozygous deletion of 3 base pairs resulting in a

deletion of the first single Glu residue of the mature peptide of NCR169 which mutation did

not abolish the symbiotic function of NCR169 (Fig. 2k and l).

We also applied the developed CRISPR/Cas9 toolkit to target *NCR-new35*, a recently

identified *NCR* gene which is crucial for the persistence of bacteroids in *M. truncatula* cv.

Jemalong nodules[13]. The empty vector and the pKSE466-RR-NCRnew35 constructs were

introduced into the roots of *M. truncatula* 2HA and *M. truncatula* ecotype R108 plants using

A. rhizogenes-mediated hairy root transformation. Nodules on empty vector-transformed roots

displayed the characteristic zonation of indeterminate nodules showing cells occupied by

rhizobia (Figure S2e and m). In contrast, undeveloped nodules were observed on roots of *M.*

truncatula 2HA and R108 plants transformed with the editing construct of *NCR-new35*

(Figures S2 and Fig. S3g-r and Fig. 4e-h). The sequence analysis showed that undeveloped

nodules, randomly collected from transformed roots of *M. truncatula* 2HA and R108 plants,

carried biallelic or monoallelic homozygous mutations in the amplicons containing the target

sequence in *MtNCR-new35* (Fig. 4q and r). SYTO13-staining revealed that these nodules did

not show the typical zonation and only their apical part fluoresced indicating that the region

corresponding to the prospective nitrogen fixation zone was devoid of rhizobia (Figure S2f-h

and n-p). The morphology and bacterial occupancy of genome edited nodules were in

agreement with the nodulation phenotype of *Mtsym19* and *Mtsym20* nodules defective in

NCR-new35[13]. These results also revealed that NCR-new35 is crucial for the symbiotic

interactions not only between *M. truncatula* 2HA and *S. medicae* WSM419 but between *M.*

truncatula R108 and *S. meliloti* FSM-MA as well.

Selecting of *NCR* genes to be targeted by CRISPR/Cas9 editing

The *M. truncatula* genome contains a large gene family coding for more than 700 NCR

peptides but only few of them, NCR169, NCR211, NCR247, NCR343 and NCR-new35, have

been proved to be crucial for symbiosis[10–15]. The sequences of the essential *NCR* genes are

highly variable, and their common feature is that they contain four cysteine residues in

conserved positions. The peptide NCR247 is unique among the crucial and most of the other

NCR peptides because it contains five residues between the third and fourth conserved

cysteines compared with the canonical four internal residues[13]. In addition, no common

feature could be identified among few NCR peptides found to be non-essential for

symbiosis[10,11,13]. Therefore, it was not straightforward to ascertain which additional NCR

peptides are essential and should be targeted. Despite all that, we have set up a selection

pipeline to identify *NCR* genes to be targeted. We have analysed several features of more than

700 predicted *NCR* genes identified in the *M. truncatula* genome[9]. The signal peptides of

NCRs are crucial for delivering the peptides to the bacteroids[4,6,7], therefore, (i) first we

selected 655 *NCR* genes coding for peptides possessing a predicted signal peptide (Fig. 3a).

The pool of *NCR* genes to be targeted (ii) were further enriched by selecting the encoded

peptides with four cysteine residues in conserved positions. (iii) We have analysed the

expression profile of the diminished list of 248 *NCR* genes in the database obtained by laser-

capture microdissection of *M. truncatula* nodules[33]. We identified 11 *NCRs* with no reads,

evidently seem to be pseudogenes, and 65 *NCRs* which are expressed at low level (< 10 000

reads) in nodules compared with other NCRs. The recently identified *NCR-new3*5 shows the

lowest expression level among the crucial NCRs[13] and thus, we have selected genes from the

172 *NCRs* showing similar or higher expression level as *NCR-new35* (Fig. 3c and d). (iv)

These *NCR* genes were analysed to predict whether they had contained sequences which were

amenable to CRISPR-based editing. The cysteine residues are essential for the function of

NCR peptides *in planta* [10,13], therefore we have prioritised those *NCRs* which could be

targeted prior to the fourth conserved cysteines to generate presumably non-functional

peptides. We have further searched the *M. truncatula* genome using an online prediction tool

CRISPOR[34] for off-target sites that could be recognized by sgRNAs designed for *NCR* genes.

The isoeletric points (pI) of crucial NCR peptides vary between 4.78 and 10.15 implying that

anionic, neutral and cationic NCR peptides could be essential for nitrogen-fixing symbiosis in

M. truncatula. For this reason, the gene of NCR peptides with wide range of charges were

selected to be edited (Fig. 3b). (v) Finally, we have selected four *NCR* genes, *NCR068*,

NCR089, *NCR128* and *NCR161*, to be targeted by CRISPR/Cas9 editing. The pI of the

encoded mature peptides varied between 5.75 and 8.34 (Fig. 3b). *NCR128* and *NCR161* are

induced in the late infection zone and showed expression in the interzone but *NCR068* and

NCR089 are preferentially expressed in the interzone and at lower level in the nitrogen

fixation zone based on the LCM transcription data (Fig. 3c,[33]). The transcription activity of

267 the four selected *NCRs* was lower than *NCR169* and *NCR211* but similar or higher than *NCR-*

268 *new35* (Fig. 3d).

270 **Inducing mutations in selected *NCR* genes using the CRISPR/Cas9 system**

271 To induce mutations in genes *NCR068*, *NCR089*, *NCR128* and *NCR161* by CRISPR/Cas9-

272 mediated cleavage, sgRNA constructs targeting a single site of each gene were designed. The

273 designed sgRNAs targeted either the upstream region of the gene close to the sequence coding

274 for the first residues of the mature peptide or at least at a site prior the sequence coding for the

275 last cysteine residue (Fig. 3e). These sgRNA constructs along with the control empty vector

276 and the constructs targeting *NCRnew-35* and *NCR169* were introduced into *M. truncatula*

277 2HA roots using *A. rhizogenes*-mediated hairy root transformation. Transformed plants were

278 inoculated with *S. medicae* WSM419 at 2-3 weeks after removal of non-transformed roots and

279 the symbiotic phenotype of transformed plants were scored at 4-7 wpi. The aerial part of

280 plants transformed with empty vector or the constructs targeting *NCR068* did not show the

281 symptoms of nitrogen deficiency (Fig. 1a and c) and plants transformed with constructs

282 targeting genes *NCR089*, *NCR128* and *NCR161* displayed the similar growth habit (data not

283 shown) indicating the effective symbiotic nitrogen fixation capacity of these plants. Nodules

284 formed on roots transformed with sgRNA constructs of the four selected *NCRs* were

285 elongated and pink suggesting that they were functional nodules (Fig. 4j, l, n and p). The

286 staining of nodule sections with SYTO13 revealed that the bacterial colonization of nodules

287 targeted for mutagenesis of genes *NCR068, NCR089*, *NCR128* and *NCR161* was identical to

288 the wild-type indeterminate nodules developed on empty-vector transformed roots (Fig. 4i-p).

289 Contrary, 2HA plants carrying mutations in *NCR169* or *NCR-new35* in each hairy root

290 displayed retarded growth, showed the symptoms of nitrogen starvation characteristically

291 (Fig. 1b) and slightly elongated white nodules were found on their roots (Fig. 4f, Figures S2

292 and S3). In addition, the occupancy of nodules mutagenized for *NCR169* and *NCRnew-35*

293 showed the phenotype characteristic for the previously described deletion mutants[10,13] (Fig.

294 1b, 4e-h, Figures S2 and S3).

295 To verify the result of CRISPR/Cas9 genome editing, the targeted regions were amplified

296 from nodule sections following microscopic analysis and sequenced using a MiSeq platform.

297 The sequence analysis of the amplicons was carried out with the online tool CRISPResso2.

298 The Cas9-induced mutations were predominantly indels, with frequencies between 47 and

299 100% of total reads (Table 1.). The genotyping also revealed that individual nodules were

300 often chimeric containing more than two alleles or the ratio of the two alleles deviated from

the expected 1:1 ratio indicating that the tissues of these nodules on hairy roots originated

from multiple founder cells with different genetic background. The identified indels generated

frameshift mutations in the targeted *NCR* genes presumably producing malfunctioning

peptides. Nevertheless, all the nodules carrying biallelic mutations in the targeted *NCRs* were

elongated and pink, indicating the presence of leghemoglobin, and were fully colonized by

rhizobia indicating that peptides NCR068, NCR089, NCR128 and NCR161 are not required

for the effective symbiotic nitrogen fixation between *M. truncatula* 2HA and *S. medicae*

WSM419 (Fig. 4i-p. and S4).

310   The nodulation phenotype of stable mutant of *NCR068* regenerated from gene edited

311   hairy roots is not strain dependent

To analyse the symbiotic phenotype of genome editing mutants in more detail, we regenerated

transgenic plants from hairy roots mutagenized for *NCR* genes. As a proof of concept, we cut

root segments that had been transformed with the editing construct of *NCR068* and stable

transformants were regenerated using a slightly modified formerly published protocol[35]. DNA

was purified from the leaves of hairy root-derived regenerated plants and the sequence

analysis revealed that all the plants were homozygous containing a single adenine insertion in

NCR068 (Fig. 5g). The one bp insertion generates a frame shift in the coding sequence of

NCR068 and the mutant sequence encodes a chimera peptide containing 27 NCR068-specific

and 23 non-NCR068-specific residues before termination. The fourth cysteine residue in the

conserved position is lacking in the chimeric NCR068 peptide and based on the necessity of

cysteines for the function of NCRs *in planta* [10,13], we reckoned that the mutant NCR068

peptide does not function correctly.

To confirm the symbiotic phenotype of regenerated *ncr068* mutant, T1 seeds were

synchronously germinated and planted into substrate with no nitrogen content. Three wpi with

S. medicae WSM419, *ncr068* stable transformants did not show symptoms of nitrogen

starvation and developed like wild type *M. truncatula* 2HA plants (Fig. 5a). Nodules, formed

on the roots of stable transformant *ncr068* mutant plants were elongated similarly what we

found on the hairy roots of composite plants induced by the transformation construct targeting

NCR068 (Fig. 5c and e). The staining of *ncr068* nodule sections with SYTO13 revealed

infected cells in interzone and nitrogen fixation zone fully occupied with elongated bacteria,

similarly as it was found in wild-type nodules and *NCR068*-targeted hairy roots (Fig. 1d and f,

5b and d). The nodulation phenotype of the regenerated mutant *ncr068* confirmed that the

peptide NCR068 is not essential for the nitrogen-fixing symbiosis between *M. truncatula* and

S. medicae WSM419.

Different symbiotic efficiency has been reported for *Sinorhizobium* sp.[36,37], therefore we

assayed the strain-dependent nodulation phenotype of regenerated *ncr068* mutant in response

to *S. meliloti* strains 2011 and FSM-MA compared with *S. medicae* WSM419. The growth

habit of wild-type *M. truncatula* 2HA and *ncr068* mutant plants was similar following

inoculation with any of the tested strains (Fig. 6). We observed a similar shoot dry weight of

wild-type and *ncr068* plants inoculated with *S. meliloti* FSM-MA or *S. medicae* WSM419

(Figure S5). In agreement with published results[38], the performance of wild-type and, in our

study *ncr068* plants as well, inoculated with *S. medicae* WSM419 superseded the plants

inoculated with *S. meliloti* 2011 (Figure S5). The *dnf7-2* mutant showed the previously

described ineffective symbiotic phenotype with all three tested rhizobia strains (Fig. 6). In line

with the growth habit, the nodules developed on wt and *ncr068* roots inoculated with the

highly compatible strains *S. medicae* WSM419 and *S. meliloti* FSM-MA were cylindrical,

pink and fully colonized by rhizobia but nodules elicited by the less effective strain *S. meliloti*

2011 were only slightly elongated (Fig. 6). In brief, no difference in plant development and

nodule structure was noticed between wt and *ncr068* plants with the tested rhizobia which

suggests that the symbiotic phenotype of plants lacking NCR068 is not strain-dependent.

Discussion

NCR peptides, present solely in IRLC clade and Dalbergoid legumes, modulate the

differentiation of nitrogen-fixing endosymbiont bacteria [4,5,8]. *NCR*s compose a large gene

family with more than 700 members[9] in *M. truncatula* and because this large number of

NCRs, their function was thought to be redundant. The identification of few individual NCRs

crucial for nitrogen-fixing symbiosis disproved this concept [10–15]. These essential NCRs

except for NCR247, which was analysed by reverse genetic approach, were identified

fortuitously using deletion mutant collections of *M. truncatula*. In addition, the large deletions

in mutants *dnf4-1* and NF-FN9363 unveiled NCR peptides, NCR178, NCR341, NCR344 and

NCR345, which are not crucial for the effective nitrogen-fixing symbiosis with the studied

rhizobia strains[11,13]. Moreover, no specific chemical properties, charge or amino acid

composition have been observed for the NCRs which would discriminate between crucial and

non-crucial NCR peptides. Therefore, the targeted gene editing is the most promising tool to

ascertain the need of a particular NCR peptide for nitrogen-fixing symbiosis. In this study, we

used the CRISPR/Cas9 technology in the model legume *M. truncatula* to knock-out four

selected *NCR* genes encoding for peptides with four conserved cysteine residues representing

the anionic, neutral and cationic members of the gene family.

The *A. rhizogenes*-mediated hairy root transformation produces chimeric root system

composed of different ratio of transformed and non-transformed roots depending on the

efficiency of transformation. To identify transformed roots in a convenient and non-

destructive way and improve gene editing efficiency, a modified vector system carrying the

DsRed transformation visible marker and single-site gRNAs under the control of the *M.*

truncatula U6 promoter were applied to generate mutations in the selected *NCR* genes in *M.*

truncatula 2HA and R108 plants. Despite of its robustness, the *A. rhizogenes*-mediated hairy

root transformation system has the disadvantage of occasionally emerged chimeric roots of a

mixture of transformed and -non-transformed cells. In addition, the CRISPR/Cas9 system is

not necessarily competent to edit the target sites in every cell or independent gene editing

events occur in several transformed cells. Our results provided evidence of both cases; we

found chimeric roots containing the mixture of wild-type and mutant alleles with different

ratio, these roots mostly retained wild-type nodulation phenotype, and we detected roots

having multi-allelic mutations at target sites. To eliminate these interfering effects and reveal

the association between CRISPR/Cas9-induced mutations and the symbiotic phenotype

without a doubt, we have sequenced the amplified fragment population of the targeted sites of

SYTO13-stained and examined sections of individual nodules. This enabled the precise

analysis of a large set of nodules and gene editing events could be coupled with the phenotype

of each nodule.

We used two essential NCRs, NCR169 and NCR-new35, to prove the efficiency of the

CRISPR/Cas9 gene editing tool in our *A. rhizogenes*-mediated hairy root transformation

system and selected four NCRs (NCR068, NCR089, NCR128 and NCR161) to assay their

requirement for symbiotic nitrogen fixation. We generated mutant versions of the four

previously non-investigated *NCR*s and the analysis of the colonization of the corresponding

nodules indicated that these four NCRs are not essential for the symbiotic interaction between

M. truncatula 2HA and *S. medicae* WSM419. The removal of variable number of non-

transgenic roots from each plant and the potential presence of root cells with non-edited

genome did not enable the reliable analysis of the nitrogen starvation phenotype (stunted

growth, yellow leaves) of the shoots of transformed plants. The nodules of all the previously

identified mutants defective in essential *NCRs* were white and slightly elongated and showed

defects in colonization of the region corresponding to the nitrogen fixation zone[10,11,13–15].

Therefore, we assessed the symbiotic phenotype of genome edited nodules mostly based on

their size, colour, and the colonization of the nitrogen fixation zone. The coordinated

sequence and phenotype analyses which we carried out on every single nodule, clearly

demonstrated that CRISPR/Cas9-induced mutations generating corrupted open reading frames

in the coding sequence of *NCR068*, *NCR089*, *NCR128* and *NCR161* did not affect the

nodulation phenotype of gene edited roots, thus these NCR peptides are not essential for the

nitrogen-fixing symbiosis between *M. truncatula* 2HA and *S. medicae* WSM419. Although

the four target NCR genes were selected based on the common features of the encoded

essential peptides (four cysteine residues in conserved positions, predicted signal peptide,

nodule-specific expression, unique sequence, various pI;[10,11,13]), these criteria were not

adequate to identify essentials NCR peptides. This conclusion was supported by the genetic

studies that identified non-essential peptides, NCR341, NCR344 and NCR345, possessing

these features. The four NCR peptides targeted in this study extend the group of NCRs proved

to be non-required for nitrogen fixation under the symbiotic conditions used in this study. Our

findings show that peptides NCR068, NCR089, NCR128 and NCR161 are either not crucial

or have minor or redundant roles in symbiosis. Alternatively, these peptides may be required

under specific symbiotic or environmental conditions or essential for strain-specific symbiotic

interaction.

We have demonstrated that the *A. rhizogenes*-mediated hairy root transformation was

successfully adapted to screen the requirement of NCR peptides for nitrogen fixation. We

have regenerated plants of gene edited knockout of *NCR068* gene directly from transformed

hairy root segments to carry out a more detailed analysis of CRISPR/Cas9-induced mutations

in stable transformants. A previous proteomic study identified peptide NCR068 in

bacteroids[39] and we anticipated that NCR068 may have a crucial role in symbiosis. The

mutant plants carrying a frame-shift mutation, induced by the insertion of a single base pair,

in *NCR068* in their germline showed similar nodulation phenotype to nodules found on *A.*

rhizogenes-mediated mutant hairy roots proving the efficiency and usability of CRISPR/Cas9-

mediated mutagenesis using hairy root transformation. The stable *ncr068* mutant enabled us

to assay the symbiotic phenotype with *S. medicae* strain WSM419 and two *S. meliloti* strains,

2011, a laboratory reference strain and FSM-MA, which is highly effective symbiotic partner

of *M. truncatula* A17 and R108[37]. All the three tested rhizobia strains induced the formation

of wild-type nodules on the root of *ncr068*, indicating that its symbiotic phenotype is not

strain-specific.

Our study demonstrates that an optimized gene editing CRISPR/Cas9 vector construct and the

developed workflow (Fig. S6) to analyse nodules developed on hairy roots mediated by *A.*

rhizogenes- transformation is an effective system to analyse the symbiotic function of *NCR*

genes. Although the selected NCR peptides were found to be non-essential for nitrogen-fixing

symbiosis, the application of the technology can identify additional indispensable and crucial

NCR peptides for nitrogen-fixing symbiosis.

Materials and Methods

Plant material, growth conditions, bacterial strains and nodulation assay. *Medicago*

truncatula cv. Jemalong 2HA line[40] (2HA) and the *M. truncatula* R108 accession (R108)

were used for transformation experiments. For sterilization and scarification, seeds were

soaked in concentrated sulfuric acid for 8 minutes, then rinsed with distilled water. Sterilized

seeds were vernalized for 3 days on inverted 0.8 % (w/v) water agar plates at 4 °C and

germinated overnight at room temperature before transformation. Seedlings were grown in

zeolite substrate (Geoproduct Kft., Mád, Hungary) for five days before inoculation with

rhizobia. Plants were grown in SANYO HM350 environmental test chambers at 23 °C at 16/8

hours light/dark periods.

M. truncatula 2HA plants were inoculated with *Sinorhizobium (Ensiferi) medicae* strain

WSM419 (SmWSM419) carrying pXLGD4 plasmid[41], constitutively expressing the *lacZ*

reporter gene under the control of the *hemA* promoter. To test strain specific symbiotic

phenotype of *ncr068* mutants, T1 plants were inoculated with *S. meliloti* 2011, *S. meliloti*

FSM-MA and SmWSM419 (pXLGD4) strains. *M. truncatula* R108 plants were inoculated

with *S. meliloti* FSM-MA. The rhizobium strains were grown overnight in liquid TA medium.

The bacteria were pelleted, resuspended and diluted to OD_{600} 0.1 in nitrogen free liquid

Fahraeus medium [42] and used for inoculation.

Construction of the CRISPR/Cas9 binary vector used for gene editing

For CRISPR/Cas9-based gene editing experiments, a modified version of the pKSE401

vector[43] was generated. DsRed fluorescent protein under the control of the 830 bp promoter of

UBQ10 gene from *Arabidopsis thaliana* was inserted into the *EcoR*I digested vector to allow

the identification of transformed roots based on their fluorescence. This generated construct

(pKSE401_RR) was further modified by replacing the AtU6 promoter with MtU6.6

promoter[31,44] resulting the construct pKSE466_RR. The MtU6.6 fragment was amplified from

the genomic DNA of *M. truncatula* A17 using the primers MtU6.6_prom_InF_F1 and

MtU6.6_prom_InF_R along with the guide cassette fragment amplified from pKSE401_RR

using the primers pKSE401_empty_InF_F2 and pKSE401_InF_R2 (Table S1) and the

amplified fragments were assembled into the *Hind*III digested pKSE401_RR vector backbone

using In-Fusion cloning (Takara Bio, Kusatsu, Japan).

Selection of targeted genes and generating CRISPR/Cas9 constructs

To select the *M. truncatula NCR* genes to be targeted, we have analysed the previously

collected and described NCR peptide sequences[9]. During the selection, first the signal peptide

prediction was repeated with the SignalP 5.0 tool[45]. If the prediction did not detect signal

peptide, the whole peptide sequence was considered as mature peptide. Further on, NCR

peptides possessing four cysteine residues in conserved position $(X_n\text{-}C\text{-}X_5\text{-}C\text{-}X_n\text{-}C\text{-}X_4\text{-}C\text{-}X_n)$[8]

were selected. The transcriptional activity of this set of *NCR* genes was analysed in the

RNAseq data of *M. truncatula* nodules[33] and genes with more than 10 000 reads were

retained. The final set of 157 *NCR* genes was classified based on their isoelectric point[9] and

few genes showing low similarity to other *NCR*s were selected from each category and were

selected to design sgRNA constructs. All sgRNAs were designed using the online tool

available at http://crispor.tefor.net/ [34] using the *M. truncatula* A17 (MtrunA17r5.0) genome as

reference[46]. In each case, the targeted region was in the first half of the coding sequence of the

NCR gene and upstream from the last conserved cysteine residue. To minimize the risk of off-

target mutations all selected sgRNA had at least two mismatches comparing with potential

off-target sites (Table S2). Targets with GN_{19} NGG motif were preferred for optimal

expression controlled by the MtU6.6 promoter. Plasmid constructs were assembled following

the protocol described by Xing at al.[43]

***Agrobacterium rhizogenes*-mediated hairy root transformation**

CRISPR constructs were introduced into the ARqua1 strain of *Agrobacterium rhizogenes* and

used for hairy root transformation of *M. truncatula* 2HA and R108 seedlings as described by

Boisson-Dernier et al.[28]. Each CRISPR construct was transformed into the roots of at least 35

plants in transformation and experiments were repeated at least twice. Following

transformation, the seedlings were grown for one week in square petri dishes containing

antibiotic-free Fahraeus medium and then seedlings were transferred to selective Fahraeus

medium supplemented with 25 mg/l kanamycin. To prevent root growth into the agar, the

roots were carefully positioned between two layers of Whatman filter paper. Following the

plant growth for additional two weeks, non-transgenic roots, identified based on the absence

of red fluorescence, were excised. After then, plants were transferred into zeolite
supplemented with Fahraeus medium containing 2 mM NH_4NO_3. Plants were further grown
in plastic boxes to preserve rhizobium-free conditions and inoculated with rhizobia after 2-3
weeks. Prior inoculation with rhizobia, newly emerged non-transgenic roots were excised
again.
509

Embryonic induction and plant regeneration from hairy-root cultures
Plantlets were regenerated from transformed hairy roots grown on plate under sterile
conditions using a slightly modified protocol described previously[35]. Three-week-old
transgenic hairy roots were identified by DsRed fluorescence using a Leica MZ10-F
fluorescence stereo microscope (Leica Microsystems GmbH, Wetzlar, Germany) and cut into
approximately 0,5-1 cm segments and placed on modified MTR-2 medium (Murashige and
Skoog medium including vitamins supplemented with 5.0 mg/l 2,4-Dichlorophenoxyacetic
acid, 0.5 mg/l 6-Benzylaminopurine (BAP), 3 % (w/v) sucrose, 50 mg/l kanamycin, 250 mg/l
cefotaxime solidified with 0.8 % (w/v) agar) to induce primary calli. After 2-3 weeks of
incubation in dark at room temperature, calli were transferred to the modified MTR-3 medium
supplemented with BAP (Murashige and Skoog medium including vitamins supplemented
with 0.5 mg/l BAP, 2 % (w/v) sucrose, 50 mg/l kanamycin, 250 mg/l cefotaxime solidified
with 0.8 % (w/v) agar) to induce embryonic calli and plates were kept in plant growing
chambers with a photopheriod of 16 h of light and 8 h darkness at 24 ^{o}C. The calli were
transferred to fresh medium in 1-2-week intervals. Shoots developed on MTR-3 + BAP
medium were put on MTR-3 medium (without selective agent) in glass tubes and transferred
to fresh medium on every 2-5 weeks to promote development and induce roots. Well-grown
shoots were propagated by cutting the apical 2-3 cm of the shoots and they were transferred
into separate tubes. After rooting, the T0 regenerated plants were transferred into jars
containing zeolite, watered with Fahraeus medium supplemented with 2 mM NH_4NO_3. The
jars were kept in growth chambers under the same condition as described above and humidity
in the jars was gradually decreased. Once new healthy leaves emerged, the plantlets were
inoculated and tested for nodulation phenotype or alternatively, they were transferred to
nutrient-rich soil in the greenhouse for seed collection.
534

Phenotypic characterization of transformed plants and microscopy analysis
To observe intact roots and nodules and detect fluorescence of transformed roots, a Leica
MZ10F fluorescent dissecting microscope equipped with ET DsRED, and GFP LP filters was

used (Leica Microsystems). Transgenic nodules were harvested, promptly immersed into ice-

cold 1xPBS (pH 7.4) and fixed with 4% paraformaldehyde. Fixed nodules were embedded in

5% (w/v) agarose gel prepared in 1xPBS and 100 µm tick longitudinal sections were prepared

with Leica VT 1200S vibratome. Nodule sections were stained in 1xPBS solution containing

5 µM SYTO13 dye (Thermo Fisher Scientific, Waltham, MA, USA) for 30 minutes at room

temperature. Images of stained nodule sections were captured using a Leica SP5 AOBS

confocal laser scanning microscope (Leica Microsystems).

DNA isolation from sections of transgenic nodules

To purify DNA from fixed nodule sections, 2-8 sections per nodule sample were collected

after SYTO13 staining and microscopic analysis and transferred into nodule lysis buffer [100

mM Tris-HCl (pH 8), 0,5M EDTA (pH 8), 0.2 % SDS, 5 mM NaCl, Proteinase K 0.02 µg/µl].

The samples were homogenized using glass pestles and incubated at 56 °C overnight.

Proteinase K was inactivated by incubating the samples at 95 °C for 15 minutes. Samples

were spin down by microcentrifuge and cooled them to room temperature. Subsequently 2 µL

RNase A (from 10 mg/ml stock) was added and incubated at room temperature for 2 min.

After that, 200 µL binding buffer [2.55M Guanidium-HCl, 10 % Triton X-100, 37 % ethanol]

was added and briefly vortexed, and 200 µL of 96 % ethanol was added to the mix and

vortexed immediately. The mixture was centrifuge at 8000 rpm for 30 second to pellet the

tissue debris. The supernatants were then transferred to QIAprep Spin Miniprep columns

(Qiagen, Hilden, Germany). Following centrifugation at 8000 rpm for 1 min, the flow-through

was discarded and columns were washed two times with 500 µL washing solution (10 mM

Tris-HCl pH 7.5, 80% ethanol). After the second washing step flow-through was discarded

and columns were centrifuged at 13.000 rpm for 3 min. DNA elution was performed by

incubating the columns with 35-50 µL of de-ionized water at 50 °C for 3 minutes and eluted

DNA was collected by centrifugation at 13,000 rpm for 1 min.

Sequence analysis of targeted genes in gene-edited nodules

The analysis of the result of target site mutagenesis of transgenic roots targeted by the

sgRNAs NCR169a, NCR169b and NCR169c was performed by Sanger sequencing of PCR

fragments amplified from the target region using ncr169_F1 and ncr169_R17 primer pair. To

analyse the edits of targeted regions of genes NCR169, NCR-new-35, NCR068, NCR128,

NCR089 and NCR161 in nodule samples, the PCR amplified fragments were analysed by

next-generation sequencing (NGS). The genomic regions of appr. 200 bp, including the target

sites, were amplified using GoTaq™ G2 polymerase (Promega corp. Madison WI, USA)

according to the manufacturer's instructions and gene specific primer pairs containing 5'

adaptor sequences (Table S1) The samples were sequenced on Illumina platform, generating

10 000 reads per sample. NGS sequence analysis was carried out by using CRISPresso2

program[32] using default parameters.

Acknowledgements

This work was supported by the Hungarian National Research Fund/National Research,

Development and Innovation Office grants OTKA 119652, 129547 and 132646 as well as by

the Collaborative Research Programme ICGEB Research Grant HUN17-03. We thank H. Cs.

Tolnainé for her skillful technical assistance and Z. Tóth for support in confocal microscopy

imaging.

Author contributions

J.B.B. and P.K conceived the project. B.G., J.B.B., D.Á. and B.H. performed the experiments

and analysed the data J.B.B. and P.K. wrote the manuscript that all authors reviewed.

Data availability

All data supporting the findings of this study are available within the paper and the

corresponding supplementary materials published online.

Competing interests

The authors declare no competing interest.

References

1.   Jones, K. M., Kobayashi, H., Davies, B. W., Taga, M. E. & Walker, G. C. How

600   rhizobial symbionts invade plants: The Sinorhizobium - Medicago model. *Nat. Rev.*

601   *Microbiol.* **5**, 619–633 (2007).

602   2.   Vasse, J., De Billy, F., Camut, S. & Truchet, G. Correlation between ultrastructural

603   differentiation of bacteriods and nitrogen fixation in alfalfa nodules. *J. Bacteriol.* **172**,

604   4295–4306 (1990).

3.   Mergaert, P. *et al.* Eukaryotic control on bacterial cell cycle and differentiation in the

Rhizobium –legume symbiosis. *Proc. Natl. Acad. Sci.* **103**, 5230–5235 (2006).

4.   Van de Velde, W. *et al.* Plant Peptides Govern Terminal Differentiation of Bacteria in

Symbiosis. *Science (80-.).* **327**, 1122–1126 (2010).

5.   Czernic, P. *et al.* Convergent Evolution of Endosymbiont Differentiation in

Dalbergioid and Inverted Repeat-Lacking Clade Legumes Mediated by Nodule-

Specific Cysteine-Rich Peptides. *Plant Physiol.* **169**, 1254–1265 (2015).

6.   Wang, D. *et al.* A Nodule-Specific Protein Secretory Pathway Required for Nitrogen-

Fixing Symbiosis. *Science (80-.).* **327**, 1126–1129 (2010).

7.   Yang, J. *et al.* A signal peptide peptidase is required for ER-symbiosome proximal

association and protein secretion. *Nat. Commun.* **14**, 4355 (2023).

8.   Mergaert, P. *et al.* A novel family in *Medicago truncatula* consisting of more than 300

nodule-specific genes coding for small, secreted polypeptides with conserved cysteine

motifs. *Plant Physiol.* **132**, 161–173 (2003).

9.   Montiel, J. *et al.* Morphotype of bacteroids in different legumes correlates with the

number and type of symbiotic NCR peptides. *Proc. Natl. Acad. Sci. U. S. A.* **114**,

5041–5046 (2017).

10. Horvath, B. *et al.* Loss of the nodule-specific cysteine rich peptide, NCR169, abolishes

symbiotic nitrogen fixation in the *Medicago truncatula dnf7* mutant. *Proc. Natl. Acad.*

Sci. U. S. A. **112**, 15232–15237 (2015).

11. Kim, M. *et al.* An antimicrobial peptide essential for bacterial survival in the nitrogen-

fixing symbiosis. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 15238–15243 (2015).

12. Sankari, S. *et al.* A haem-sequestering plant peptide promotes iron uptake in symbiotic

bacteria. *Nat. Microbiol.* **7**, 1453–1465 (2022).

13. Horváth, B. *et al.* The *Medicago truncatula* nodule☐specific cysteine☐rich peptides,

NCR343 and NCR☐new35 are required for the maintenance of rhizobia in

nitrogen☐fixing nodules. *New Phytol.* **239**, 1974–1988 (2023).

14. Zhang, R. *et al.* Nodule-Specific Cysteine-Rich Peptide 343 is required for symbiotic

nitrogen fixation in *Medicago truncatula. Plant Physiol.* (2023)

doi:10.1093/plphys/kiad454.

15. Gao, F. *et al.* NCR343 is required to maintain the viability of differentiated bacteroids

in nodule cells in *Medicago truncatula. New Phytol.* (2023) doi:10.1111/nph.19180.

16. Zhang, Y., Malzahn, A. A., Sretenovic, S. & Qi, Y. The emerging and uncultivated

potential of CRISPR technology in plant science. *Nat. Plants* **5**, 778–794 (2019).

17. Le, H. *et al.* CRISPR/Cas9-Mediated Knockout of Galactinol Synthase-Encoding

640   Genes Reduces Raffinose Family Oligosaccharide Levels in Soybean Seeds. *Front.*

641   *Plant Sci.* **11**, (2020).

642 18. Shu, H., Luo, Z., Peng, Z. & Wang, J. The application of CRISPR/Cas9 in hairy roots

643   to explore the functions of *AhNFR1* and *AhNFR5* genes during peanut nodulation.

644   *BMC Plant Biol.* **20**, 417 (2020).

645 19. Li, G. *et al.* Development of an Agrobacterium-mediated CRISPR/Cas9 system in pea

646   (*Pisum sativum* L.). *Crop J.* **11**, 132–139 (2023).

647 20. Wang, L. *et al.* CRISPR/Cas9 knockout of leghemoglobin genes in *Lotus japonicus*

648   uncovers their synergistic roles in symbiotic nitrogen fixation. *New Phytol.* **224**, 818–

649   832 (2019).

650 21. Meng, Y. *et al.* Targeted mutagenesis by CRISPR/Cas9 system in the model legume

651   *Medicago truncatula. Plant Cell Rep.* **36**, 371–374 (2017).

652 22. Zhang, H. *et al.* Efficient Generation of CRISPR/Cas9-Mediated Homozygous/Biallelic

653   *Medicago truncatula* Mutants Using a Hairy Root System. *Front. Plant Sci.* **11**, 1–11

654   (2020).

655 23. Confalonieri, M. *et al.* CRISPR/Cas9-Mediated Targeted Mutagenesis of *CYP93E2*

656   Modulates the Triterpene Saponin Biosynthesis in *Medicago truncatula. Front. Plant*

657   *Sci.* **12**, 1–16 (2021).

658 24. Jain, D., Jones, L. & Roy, S. Gene editing to improve legume-rhizobia symbiosis in a

659   changing climate. *Curr. Opin. Plant Biol.* **71**, 102324 (2023).

660 25. Limpens, E. *et al.* RNA interference in *Agrobacterium rhizogenes*-transformed roots of

661   Arabidopsis and *Medicago truncatula. J. Exp. Bot.* **55**, 983–992 (2004).

662 26. Wang, Q. *et al.* Host-secreted antimicrobial peptide enforces symbiotic selectivity in

663   *Medicago truncatula. Proc. Natl. Acad. Sci. U. S. A.* **114**, 6854–6859 (2017).

664 27. Yang, S. *et al.* Microsymbiont discrimination mediated by a host-secreted peptide in

665   Medicago truncatula. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 6848–6853 (2017).

666 28. Boisson-Dernier, A. *et al. Agrobacterium rhizogenes*-transformed roots of *Medicago*

667   *truncatula* for the study of nitrogen-fixing and endomycorrhizal symbiotic associations.

668   *Mol. Plant-Microbe Interact.* **14**, 695–700 (2001).

669 29. Wang, D. *et al.* Optimized CRISPR guide RNA design for two high-fidelity Cas9

670   variants by deep learning. *Nat. Commun.* **10**, 1–14 (2019).

671 30. Brinkman, E. K., Chen, T., Amendola, M. & Van Steensel, B. Easy quantitative

672   assessment of genome editing by sequence trace decomposition. *Nucleic Acids Res.* **42**,

673   1–8 (2014).

31.   Jacobs, T. B., LaFayette, P. R., Schmitz, R. J. & Parrott, W. A. Targeted genome
modifications in soybean with CRISPR/Cas9. *BMC Biotechnol.* **15**, 1–10 (2015).

32.   Clement, K. *et al.* CRISPResso2 provides accurate and rapid genome editing sequence
analysis. *Nat. Biotechnol.* **37**, 224–226 (2019).

33.   Roux, B. *et al.* An integrated analysis of plant and bacterial gene expression in
symbiotic root nodules using laser-capture microdissection coupled to RNA
sequencing. *Plant J.* **77**, 817–837 (2014).

34.   Concordet, J. P. & Haeussler, M. CRISPOR: Intuitive guide selection for
CRISPR/Cas9 genome editing experiments and screens. *Nucleic Acids Res.* **46**, W242–
W245 (2018).

35.   Crane, C., Wright, E., Dixon, R. A. & Wang, Z. Y. Transgenic *Medicago truncatula*
plants obtained from Agrobacterium tumefaciens-transformed roots and *Agrobacterium*
rhizogenes-transformed hairy roots. *Planta* **223**, 1344–1354 (2006).

36.   Terpolilli, J. J., O'Hara, G. W., Tiwari, R. P., Dilworth, M. J. & Howieson, J. G. The
model legume *Medicago truncatula* A17 is poorly matched for N 2 fixation with the
sequenced microsymbiont Sinorhizobium meliloti 1021. *New Phytol.* **179**, 62–66
(2008).

37.   Kazmierczak, T. *et al.* Specific Host-Responsive Associations Between *Medicago*
truncatula Accessions and Sinorhizobium Strains. *Mol. Plant-Microbe Interact.* **30**,
399–409 (2017).

38.   Larrainzar, E., Gil-Quintana, E., Seminario, A., Arrese-Igor, C. & González, E. M.
Nodule carbohydrate catabolism is enhanced in the *Medicago truncatula* A17-
Sinorhizobium medicae WSM419 symbiosis. *Front. Microbiol.* **5**, (2014).

39.   Durgo, H. *et al.* Identification of nodule-specific cysteine-rich plant peptides in
endosymbiotic bacteria. *Proteomics* **15**, 2291–2295 (2015).

40.   Rose, R. J., Nolan, K. E. & Bicego, L. The development of the highly regenerable seed
line Jemalong 2HA for transformation of *Medicago truncatula* - Implications for
regenerability via somatic embryogenesis. *J. Plant Physiol.* **155**, 788–791 (1999).

41.   Leong, S. A., Williams, P. H. & Ditta, G. S. Analysis of the 5' regulatory region of the
gene for δ-aminolevulink add synthetase of Rhizobium meliloti. *Nucleic Acids Res.* **13**,
5965–5976 (1985).

42.   Fahraeus, G. The Infection of Clover Root Hairs by Nodule Bacteria Studied by a
Simple Glass Slide Technique. *J. Gen. Microbiol.* **16**, 374–381 (1957).

43.   Xing, H. L. *et al.* A CRISPR/Cas9 toolkit for multiplex genome editing in plants. *BMC*

Plant Biol. **14**, (2014).

44.   Kim, G. B. & Nam, Y. W. Isolation and Characterization of *Medicago truncatula* U6

Promoters for the Construction of Small Hairpin RNA-Mediated Gene Silencing

Vectors. *Plant Mol. Biol. Report.* **31**, 581–593 (2013).

45.   Almagro Armenteros, J. J. *et al.* SignalP 5.0 improves signal peptide predictions using

deep neural networks. *Nat. Biotechnol.* **37**, 420–423 (2019).

46.   Pecrix, Y. *et al.* Whole-genome landscape of *Medicago truncatula* symbiotic genes.

Nat. Plants **4**, 1017–1025 (2018).

Figure legends

Figure 1. Plant gross and nodulation phenotype of *Agrobacterium rhizogenes*-mediated

transgenic *M. truncatula* Jemalong 2HA hairy-roots mutagenized with CRISPR/Cas9 gene

editing. No sign of nitrogen deficiency was observed on plants transformed with empty vector

(**a**) or the construct targeting *NCR068* (**c**) 4 weeks post inoculation with *S. medicae* WSM419.

Transgenic roots were identified by fluorescence of DsRed protein. Elongated pink nodules,

showing the characteristic features of indeterminate nodules, developed on transgenic roots.

These nodules were fully colonized by rhizobia visualized by the fluorescent dye SYT013 (**d**

and **f**). (**b**) Targeted mutagenesis of *NCR169* resulted in transgenic plants showing the

symptoms of nitrogen starvation. Small, spherical and white ineffective nodules were formed

on *NCR169*-targeted roots. (**e**) *ncr169* mutant nodules did not contain infected cells in the

region corresponding to the nitrogen fixation zone of wild-type plants. Inset images in (**d-f**)

show the higher magnification of cells in nodule regions indicated by white rectangles. (**g** and

h) The targeted regions were amplified and sequenced from the transgenic nodules. The

sequence analysis revealed biallelic mutations causing frameshifts in edited *NCR169* (g) and

NCR068 (h) genes. ZII: infection zone; IZ: interzone; ZIII: nitrogen fixation zone; EV, empty

vector

Figure 2. The nodulation phenotype of hairy roots transformed with the construct pKSE466-

RR-MtNCR169a was associated with the proportion of wild-type *NCR169* allele or with the

presence of mutant NCR169 lacking a single residue. White (Fix-) and pink (Fix+) nodules

were randomly harvested 3-5 wpi with *S. medicae* WSM419 and the result of gene editing

was identified by the sequencing of the targeted region using an Illumina platform. The ratios

of reads assigned to mutant (brown ochre) and wild-type (blue) alleles are presented in pie

chart. Fix- nodules (**b**, **d**, and **f**) carried almost exclusively insertions or deletions resulting in

frameshift in the coding sequence of *NCR169* (**a**, **c** and **e**). Contrary, pink nodules (**g**, **i** and **k**)

carried either unmodified alleles on a large scale (>40%) (**h** and **j**) or a deletion of 3 base pairs

that did not result in the loss of function of NCR169 (**l**).

Figure 3. Selection of *NCR* genes to be targeted by CRISPR/Cas9 mutagenesis. (**a**) 679 *M.*

truncatula NCR genes were grouped based on the presence of a signal peptide in their

encoded peptide sequence and containing four cysteine residues in conserved positions. (**b**)

the 248 NCR peptides complying with all the three features (**a**) were classified based on the

charge of the encoded mature peptides. NCR peptides crucial for symbiosis are highlighted in

red, peptides targeted in this study by gene editing are in black and peptides found to be non-

essential in the former study[13] are in purple. (**c**) Relative spatial expression of four *NCR* genes

essential for symbiosis and four *NCRs* mutagenized using CRISPR/Cas9 method in this study.

Expression was determined by RNAseq of nodule zones obtained with laser-capture

microdissection described in the study[33]. (**d**) The total read number of the *NCR* transcripts

identified by RNAseq and analysed in panel (**c**). All the four *NCR*s selected for gene editing

show higher transcriptional activity than *NCR-new35* showing the lowest expression activity

among the crucial *NCR* genes. (**e**) The cleavage sites of Cas9 on the sequences of *NCR* genes

targeted by CRISPR/Cas6 gene editing in this study. The target sites of each guideRNA were

designed upstream of the nucleotides coding for the fourth conserved cysteines of each NCR

peptides. ZIII: nitrogen fixation zone; IZ: interzone; IID: distal part of infection zone; IIP:

proximal part of infection zone; pI: isoelectric point

Figure 4. (a-p) Nodulation phenotype of *M. truncatula* 2HA (**e, f, i-p**) and R108 (**g** and **h**)

plants mutagenized using the CRISPR/Cas9 method with *A. rhizogenes*-mediated hairy root

transformation. Transformation with gene editing construct designed to target *NCR-new35* (**e-**

h) resulted in ineffective white and slightly elongated nodules while the homozygous or

biallelic mutant alleles of *NCR068* (**i** and **j**), *NC089* (**k** and **l**), *NCR128* (**m** and **n**) and

NCR161 (**o** and **p**) were similar to the elongated and pink nodules formed on empty vector-

transformed roots of *M. truncatula* 2HA (a and b) and R108 (**c** and **d**) plants. Images of

nodule sections following SYTO13 staining and whole nodules were taken 4-5 weeks post

inoculation with *S. medicae* WSM419 used for *M. truncatula* 2HA and *S. meliloti* FSM-MA

for R108 plants. (**q-v**) Allele-specific results of CRISPR/Cas9 editing of the targeted *NCR*

genes identified by next generation sequencing on Illumine platform. Black and brown

rectangles show deleted and inserted nucleotides, respectively.

Figure 5. (**a**) Stable *ncr068* mutants regenerated from segments of transgenic hairy roots with

induced mutations in *NCR068* using the CRISPR method retained wild type symbiotic

phenotype similar wild wild-type (WT) *M. truncatula* 2HA plants 3 wpi with *S. medicae*

WSM419. (**b** and **c**) The gene edited nodules developed on *A. rhizogenes*-transformed hairy-

roots were elongated and pink with colonized cells in the nitrogen fixation zone. (**d** and **e**)

Nodules developed on the roots of regenerated stable *ncr068* mutants were similar to the

nodules found on hairy-roots mediated by *A. rhizogenes*. (**f**) The selected hairy root

(NCR068-5_S138) was chimeric and composed of cells with homozygous insertions of A or

G in the targeted region of *NCR068*. (**g**) The regenerated stable *ncr068* mutants (Ac2211)

carried a homozygous A insertion causing a frameshift in the coding sequence of *NCR068*. (**h**)

The alignment of WT and mutant NCR068 peptide sequences encoded by the mutated gene in

the stable *ncr068* mutant. Identical residues are highlighted by black background, conserved

Cys residues are in red. SP, signal peptide; MP, mature peptide

Figure 6. Stable *ncr068* T1 mutant plants inoculated with three rhizobia strains show wild-

type symbiotic phenotype. Wild-type *M. truncatula* 2HA (WT), the deletion mutant deficient

in *NCR169* (*dnf7-2*) and *ncr068* mutant plants were inoculated with *S. medicae* WSM419, *S.*

meliloti FSM-MA or *S. meliloti* 2011 and scored for symbiotic phenotype at 5 wpi. WT and

ncr068 plants showed similar growth habit but presented less effective symbiotic interaction

with *S. meliloti* 2011 compared with the two other strains. Mutant *dnf7-2* showed nitrogen

deficiency with all three tested rhizobia strains. SYTO13-staining revealed a similar

colonization of WT and *ncr068* nodules while only bacteria in infection threads and sporadic

saprophytic bacteria fluoresced in the basal part of *dnf7-2* nodules.

Table 1. Editing efficiency of different targeted *NCR* genes in nodules harvested from hairy

roots mediated by *A. rhizogenes* transformation. Randomly selected nodules were harvested

3-5 wpi with *S. medicae* WSM419 and total DNA was isolated from fixed nodule sections.

The PCR amplified target regions were sequenced on Illumina platform and the reads were

analysed by using the CRISPresso2 tool. The gene *NCR169* was targeted for gene editing as a

control. Three pinkish elongated nodules were selected randomly from hairy roots and also

three small white nodules were picked randomly from plants which did not form any pink
nodules on their roots.
*One of the pinkish nodules carried mutant alleles on large proportion (94%). This mutation
was a three bp deletion that resulted in an in frame deletion of the first Glu residue of the
mature peptide NCR169. The lack a of this single residue probably did not abolish the
function of peptide NCR169.
814
815
816 **Additional information**
817
818 **Supplementary information**
819
Fig. S1. Analysis of three sgRNAs targeting *NCR169*.
Fig. S2. The symbiotic phenotype of nodules formed on *M. truncatula* Jemalong and R108 hairy roots
edited for *NCR-new35*.
Fig. S3. The symbiotic phenotype of nodules formed on *M. truncatula* Jemalong and R108 hairy roots
edited for *NCR169* and *NCR-new35*.
Fig. S4. The nodulation phenotype of hairy roots edited for *NCR089*, *NCR0128* and *NCR161*.
Fig. S5. The shoot biomass of stable *ncr068* mutant plants inoculated with three rhizobia strains.
Fig. S6. The workflow applied for simultaneous genetic and phenotypic analysis of gene edited
individual nodules.
Table S1. List of primers used in this study.
Table S2. Off-target prediction using the CRISPOR tool.
831
832
833

a *EV*   **b** *NCR169*   **c** *NCR068*

dsRed   dsRed   dsRed

d   **e**   **f**

ZII
IZ
ZIII

0.5 mm   0.1 mm   0.5 mm   0.1 mm   0.5 mm   0.1 mm

g

NCR169 **-7/+1 bp**

C T T G T T G T A A T C T T C G A A G A T A T A G T G C A T A T A A G A T A A G T T
sgRNA

h

NCR068 **-1/+1 bp**

C C T G A A T A T T T G A A G T G T C A A G T G G T A T T G T G C A T A T G A T T
sgRNA

Insertion
Deletion
Quantification window

a WT   *ncr068* T1

b SYTO13   **c**   **d** SYTO13   **e**

f *NCR068-5_S138* +1 A/G
TTGAAGTGTCAAGGTGGTAT

g *ncr068* T₁ (Ac2211) +1 A
TTGAAGTGTCAAGGTGGTAT

Insertion

h

NCR068 (WT)
ncr068 T₁ (Ac2211)

```
                SP                              MP
        1         10          20          30          40
NCR068 (WT)        MAQILMFVYFLIIFSLSFLVESIKIFTEHRCRTDADCPAR
ncr068 T₁ (Ac2211) MAQILMFVYFLIIFSLSFLVESIKIFTEHRCRTDADCPAR
                   ************************************* *

                   50          60          70
NCR068 (WT)        ELPEYLKCQGGMCRLLIKKD
ncr068 T₁ (Ac2211) ELPEYLKCCRWYVSAFNKEGLNQKIIIFPYIK
                   *******  :  .  *  : . :
```

S. medicae WSM419			S. meliloti FSM-MA			S. meliloti 2011		
WT	dnf7-2	ncr068 T1	WT	dnf7-2	ncr068 T1	WT	dnf7-2	ncr068 T1

	NCR169		NCR068	NCR089	NCR128	NCR161
Number of sequenced nodules	3 white	3 pink	15	13	10	20
Number nodules carrying max. 50% WT allele	3	2	15	13	9	19
Number of fully mutant nodules (mismatch/indel)	3	0*	14	13	9	12
Number of fully mutant nodules (only indel)	3	0*	7	13	9	12
Nodule samples carrying only indel alleles	**100%**	**0%**	**47%**	**100%**	**90%**	**60%**