Separating cognitive and motor processes in the behaving mouse

Munib A. Hasnain[1,3*], Jaclyn E. Birnbaum[2,3*], Juan Luis Ugarte Nunez[1], Emma K. Hartman[1], Chandramouli Chandrasekaran[4,5,6], Michael N. Economo[1,3,6#]

[*] These authors contributed equally. Order determined via coin flip.

[#] Corresponding author

Email: mne@bu.edu

[1] *Department of Biomedical Engineering, Boston University, Boston, MA*

[2] *Graduate Program for Neuroscience, Boston University, Boston, MA*

[3] *Center for Neurophotonics, Boston University, Boston, MA*

[4] *Department of Psychological and Brain Sciences, Boston University, Boston, MA*

[5] *Department of Neurobiology & Anatomy, Boston University, Boston, MA*

[6] *Center for Systems Neuroscience, Boston University, Boston, MA*

Abstract

The cognitive processes supporting complex animal behavior are closely associated with ubiquitous movements responsible for our posture, facial expressions, ability to actively sample our sensory environments, and other central processes. These movements are strongly related to neural activity across much of the brain, making it challenging to dissociate the neural dynamics that support cognitive processes from those supporting movements when they are highly correlated in time. Of critical importance is whether the dynamics supporting cognitive processes and related movements are separable, or if they are both driven by common neural mechanisms. Here, we demonstrate how the separability of cognitive and motor processes can be assessed, and, when separable, how each component can be isolated. We establish a novel two-context behavioral task in mice that involves multiple cognitive processes and show that commonly observed dynamics taken to support cognitive processes are strongly contaminated by movements. When cognitive and motor components are isolated using our analytical approach, we find that they exhibit distinct dynamical trajectories. Further, properly accounting for movement revealed that separate populations of cells encode cognitive and motor variables, in contrast to the 'mixed selectivity' reported by prior work. Accurately isolating the dynamics associated with particular cognitive and motor processes will be essential for developing conceptual and computational models of neural circuit function and evaluating the function of the cell types of which neural circuits are composed.

Introduction

Goal-directed behavior follows from an interplay between cognitive and motor processes in the mammalian brain. For example, while dribbling down the court, a basketball player must track the players around them, decide which play to run, plan their next move, execute fast and accurate movements, and flexibly adapt their movements according to the reactions of other players. The neural representations of these cognitive and motor processes are often distributed across many of the same brain areas[1–6] and engaged concurrently[7–9]. For these reasons, disentangling the neural signatures of specific cognitive and motor processes is challenging.

Many behavioral tasks are designed with this issue in mind, aiming to isolate constituent neural processes by separating them into discrete temporal epochs. For example, in sensory-guided decision-making tasks, subjects are often trained to withhold responses while stimuli are presented so that perceptual decision making is isolated in time from associated actions. Analogous paradigms have been developed to uncover the neural underpinnings of working memory[10–12], motor planning[13,14], contextual encoding[15,16], reward prediction[17,18], and other cognitive processes[19–22].

These paradigms often rely on the assumption that experimental subjects withhold motor output during task epochs in which instructed responses are absent. However, uninstructed movements not required for task completion – such as changes in posture, facial expressions, and gaze – are commonly observed in rodents, humans, and non-human primates during learned behavioral tasks[23–29]. Importantly, uninstructed movements explain much of the variance observed in brain-wide neural activity[27,30–32] and can be strongly correlated with task variables[12,28,33–38].

These observations imply a serious challenge for experimental studies addressing a wide-ranging set of cognitive processes. If uninstructed movements are correlated with a latent cognitive variable of interest – a stimulus perceived, a decision made, a memory stored, or a motor plan formed – then the neural dynamics leading to, or resulting from, uninstructed movements can be easily misconstrued as responsible for that cognitive process. One common approach to isolate putative cognitive dynamics is to measure and 'regress out' movements[24,30,39–44], but this approach assumes that cognitive and motor processes are driven by independent neural dynamics. Some cognitive processes may be inherently embodied, such that their associated neural dynamics are linked to overt movement[45,46]. Regressing out neural dynamics associated with embodied movements would lead to the inadvertent removal of the precise dynamics one wishes to study. Whether cognitive and motor dynamics are separable remains an open question and likely depends on the brain area, behavior, and cognitive process of interest. Methods both for evaluating separability within specific experimental paradigms and for effectively dissociating cognitive and motor dynamics, when separable, are lacking.

Here, we address the separability of dynamics associated with cognitive processes (defined broadly here as those not directly related to motor commands, efference copies, or sensory reafference) and correlated movements. We developed a behavioral paradigm in which mice perform sensory-guided movements involving multiple cognitive signals and idiosyncratic, task-correlated uninstructed movements. We build upon work showing that different neural processes can be multiplexed in a single brain region[39,40,47–50] to develop a novel method for assessing the separability of cognitive and motor dynamics – and for isolating each component, when they are separable. This approach is simple to adopt and model-free, precluding the need for parameter adjustment or assumptions of linearity.

We find that some cognitive signals are separable from dynamics associated with co-occurring movements while others are largely inseparable. When dynamics are separable, examining the 'cognitive' component of neural dynamics revealed trajectories that differed markedly from estimates of the same dynamics when corrupted by uninstructed movements. Strikingly, we found that cognitive and motor variables were largely encoded by separate populations of cells, in contrast to previous work that did not account for uninstructed movements. Together, these results highlight the importance of critical consideration of the relationship between cognition and movements to better understand the neural dynamics supporting complex processes and how they map onto the myriad cell types comprising neural circuits.

Results

Task-switching behavioral paradigm

We first designed a behavioral paradigm associated with multiple cognitive processes. In this paradigm, head-fixed mice performed two directional licking tasks that alternated block-wise within each behavioral session. These tasks varied in their cognitive demands but require the same instructed motor output.

This paradigm employed an established delayed-response task (DR) in which motor planning is temporally dissociated from movement execution (**Fig. 1a**). In this task, an auditory stimulus indicated the location of a reward. After a brief delay epoch, an auditory 'go cue' instructed the mouse to move. If the mouse performed a directional tongue movement to the correct target, a water reward was delivered. In separate blocks of trials, mice performed a water-cued (WC) task in which all auditory stimuli were omitted. Instead, a drop of water was presented at a random point in time at a randomly selected reward port. Animals consumed water immediately upon its presentation. Mice received no explicit cues signaling DR and WC blocks. The DR task required a perceptual decision that guides the planning of a subsequent motor action. In the WC task, uncertainty in the timing and location of reward prevented formation of a motor plan prior to reward presentation, and instead, mice detected reward availability using olfactory and/or vibrissal cues[51]. This structure required animals to maintain a representation of the current block identity (context) for successful task execution.

We examined uninstructed movements in this behavioral paradigm using high-speed video. We tracked movement of the tongue, jaw, nose, and paws[52] and calculated frame-by-frame motion energy to quantify movement in a feature-agnostic manner[31] (**Fig. 1d-f**). We found that mice performed uninstructed movements in both contexts that varied in both their identity and timing across trials and contexts (**Fig. 1e; Supplementary Movie 1**).

Prior work has established the necessity of the antero-lateral motor cortex (ALM) and tongue-jaw motor cortex (tjM1) for the planning and execution of tongue movements[14,53–56]. To assess cortical involvement in the initiation of instructed movements (directional licking) in both behavioral tasks, we used optogenetic photoinactivation to inhibit the ALM and tjM1 at the go cue (DR) or water presentation (WC) bilaterally in VGAT-ChR2-EYFP mice (2 mice, 6 sessions). Simultaneous photoinactivation of both regions at the go cue impaired the initiation of movement in both contexts (**Fig. 1b,c**; DR left trials: $p = 4\times10^{-5}$, DR right trials: $p = 0.02$, WC left trials: $p = 0.02$, WC right trials: $p = 0.04$, paired *t*-test; **EDFig. 1c,d**). To assess cortical involvement in the expression of uninstructed movements, we bilaterally silenced the ALM and tjM1 during the delay epoch of DR trials (2 mice, ALM: 15 sessions, tjM1: 9 sessions). Uninstructed movements were suppressed by photoinactivation of either area (**EDFig. 1b,** ALM: $p=3\times10^{-4}$; tjM1: $p=0.002$, paired *t*-test) and more strongly suppressed by concurrent photoinactivation of both areas (**Fig. 1g-h,** 2 mice, 14 sessions; $p=3\times10^{-7}$), similar to previous observations[34] (but see ref[57]). In the DR task, delay photoinactivation of both areas led to a significant decrease in overall performance (**Fig. 1i**, All trials: $p=0.002$, paired *t*-test; and **EDFig. 1b**) as expected from the disruption of choice/planning dynamics[14]. These results indicate that the ALM and tjM1 are involved in the generation of uninstructed movements and the planning and execution of instructed directional tongue movements in the two-context paradigm.

Neural dynamics related to task variables

While movement- and task-related information are encoded in both the ALM and tjM1 in the DR task[53,58] and other directional licking tasks[55], task variables are more strongly represented in the ALM[55,56,59] and so we focused our subsequent analyses there. We recorded activity extracellularly in the ALM with high-density silicon probes during the two-context paradigm to track dynamics associated with planning, context, and the execution of movements (**Fig. 2**; DR task only: 9 mice, 25 sessions, 1651 units including 483 well-isolated single units; task switching (DR and WC tasks): 6 mice, 12 sessions, 522 units including 214 well-isolated single units). As reported previously, we found that choice selectivity (right vs. left DR trials) was widespread across all task epochs (sample: 36%, delay: 42%, response: 58% of single units; 9 mice, 483 single units). Individual units showed a variety of activity patterns across trial types, including delay epoch preparatory activity preceding

134 instructed movements, a key signature of motor planning[54] (**Fig. 2b,** *left*). Individual units were also selective for
135 behavioral context during all task epochs

137
Figure 1 – Cortical dependence and uninstructed movements during a two-context task. a. *Top*, Schematic of the two-context task. In
delayed-response (DR) trials, an auditory stimulus was presented during the sample epoch (1.3 s) instructing mice to lick for reward to the
right (white noise) or left (pure tone). Mice were trained to withhold their response during a delay epoch (0.9 s) and initiated their response
following an auditory go cue. In WC trials, mice were presented with a water reward at a random port at random inter-reward intervals. The
WC task was introduced once mice became experts in the DR task. *Bottom*, 20-trial trailing average of correct, error, and ignore rates for an
example session. **b.** Performance for an example session separated by trial type (left vs. right), control vs. photoinactivation trials, and by
context. Photoinactivation (1 s) was initiated at the go cue. Blue dots indicate right licks and red dots indicate left licks. Grey bars, correct
trials; black bars, error trials. **c.** Proportion of trials with correct licks within 600 ms of the go cue on photoinactivation trials. Each point
indicates mean value for a session. *Top*, DR context. *Bottom*, WC context. Asterisks denote significant differences ($p < 0.05$) between
control and photoinactivation trials (DR left trials: $p = 4 \times 10^{-5}$, DR right trials: $p = 0.02$, WC left trials: $p = 0.02$, WC right trials: $p = 0.04$,
paired t-test). **d.** Behavior was tracked with high-speed video from side (*left*) and bottom (*right*) views. **e.** Trajectories of delay epoch
uninstructed movements from an example session overlayed. Uninstructed movements are highly variable across trials and across time.
Top, jaw, nose, and paw speed for an example trial. *Bottom*, feature overlay for a subset of trials in an example session. At each time bin, t,
an [r, g, b] color value was encoded as [jaw$_t$, nose$_t$, paw$_t$]. **f.** Schematic of motion energy calculation. Example frames depicting high and
low motion energy are shown. **g-i.** Bilateral motor cortex photoinactivation during the delay epoch (n=2 mice). **g.** Motion energy on
control (*top*) and photoinactivation (*bottom*) trials for an example session. On photoinactivation trials, light was delivered for 0.8 seconds
beginning at the start of the delay epoch. **h.** Average delay epoch motion energy on single sessions for control and photoinactivation trials.
Red points indicate left trials and blue points indicate right trials for a single session. **i.** Performance for control and delay epoch

photoinactivation trials. Mice revert to a default choice following delay epoch photoinactivation. Each point indicates average performance
over all sessions for each mouse. Error bars indicate standard deviation across all sessions.

and during the inter-trial interval (ITI) (26% of single units; 6 mice, 214 single units), suggesting persistent
coding of context (**Fig. 2b,** *right*).

We next sought to understand how the selectivity patterns observed in individual cells were encoded at the
population level. We examined neural activity in state space, in which the firing rate of each unit represents one
dimension. We defined one-dimensional coding directions (CD) within the high-dimensional state space that best
encoded task variables[54,60–62] (**Fig. 2c**). We first examined putative neural correlates of three cognitive processes –
choice, urgency, and contextual encoding.

We defined a coding direction for left-right trial type, CD_{choice}, as the direction in state space that best
differentiates activity on left versus right trials during the delay epoch of DR trials. Projections along CD_{choice}
revealed that choice selectivity in the ALM emerges early in the sample epoch and continues to grow throughout
the delay epoch before collapsing at the onset of the response epoch (**Fig. 2d**). Such preparatory activity is
predictive of animals' upcoming instructed responses[13,54,58,60,63,64] and has been taken to represent a neural
correlate of a motor plan[54,65] or choice[32,61,66]. We next defined CD_{ramp} as the direction that captures trial-type
independent increases or decreases in preparatory activity between the onset of the stimulus and the go cue in DR
trials[60,61,67,68]. This non-selective ramping has been thought to represent an urgency (or timing) signal[69–72]. Finally,
we defined $CD_{context}$, the direction that best differentiates activity on DR and WC trials during inter-trial intervals.
Projections along $CD_{context}$ revealed that context is encoded in the ALM throughout each trial (**Fig. 2d**) with strong
modulation at the onset of the response epoch.

178
Figure 2 – Single cell activity and population dynamics encode task-relevant cognitive and motor processes. Schematic of silicon
probe recordings in the anterolateral motor cortex (ALM). **b.** Spike rasters (*top*) and peri-stimulus time histograms (PSTH; *bottom*) for
four example units in the ALM. Units show selectivity for trial type (left vs. right) and context (DR vs. WC) during all task epochs. **c.**
Schematic illustrating identification of coding directions (CD). Coding directions are defined as directions in state space that maximally
separate activity between trajectories defined by trial types (CD_{choice}, CD_{action}), context ($CD_{context}$), or time points (CD_{ramp}). **d.** Projections
along each of four defined CDs. Gray shaded regions indicate time points used for CD estimation. Mean and 95% CI of bootstrap
distribution shown.

In addition to neural activity putatively related to these cognitive processes, we also examined neural signals
associated with the execution of movements. We defined an additional coding direction, CD_{action}, as the direction
that captures trial-type selective changes in neural activity that emerge after the go cue. Projections of neural
activity along CD_{action} robustly encoded movement direction in the ALM, as expected. We focus our analyses on
these four population level signals, which putatively encode ALM dynamics associated with cognitive and motor
processes in the two-context task-switching paradigm.

Uninstructed movements related to preparatory dynamics

We next examined uninstructed movements in the DR task and their correlation with both task variables and
preparatory dynamics (**Fig. 3**). We found that animals frequently performed uninstructed movements that were
correlated with trial type (**Fig. 3a**) and time within each trial. On single trials, upcoming choice could be decoded
from uninstructed movements beginning immediately after the presentation of the sample tone, with accuracy
increasing throughout the delay epoch (**Fig. 3b**).

Not only were uninstructed movements related to the animal's upcoming actions, but they were highly correlated
with preparatory dynamics. The onset and magnitude of uninstructed movements coincided with the onset and
magnitude of choice selectivity and ramping (**Fig. 3c,e**). We used cross-validated multiple linear regression to
predict projections along CD_{choice} and CD_{ramp} on a trial-by-trial basis from captured kinematic features. Model
predictions were often highly correlated with motor planning signals, indicating a close relationship between
animals' uninstructed movements and preparatory dynamics (**Fig. 3d,f**; CD_{choice}: $R^2 = 0.41 \pm 0.23$; CD_{ramp}: $R^2 =$
0.48 ± 0.23), although a high degree of variability was observed in the features that were most predictive across
animals and sessions (**EDFig. 2a-c**).

To further understand the relationship between uninstructed movements and preparatory dynamics, we trained
additional animals on a randomized delay task in which the timing of the go cue cannot be anticipated[64].
Incorporating uncertainty into the length of the delay epoch leads to a qualitative change in population-level
choice selectivity; selectivity at both the single-neuron level and in CD_{choice} projections emerges earlier than in the
standard DR task and ramps up to a stable fixed point, suggesting that motor plans are prepared prior to the
earliest possible go cue time[64] (**Fig. 3g,** *middle*). We found that the timing of uninstructed movements, too, shifted
in an analogous fashion (**Fig. 3g,** *bottom*). Projections along CD_{choice} remained predictable from uninstructed
movements on a single- trial level (**Fig. 3h**; average R^2 across sessions – CD_{choice}: 0.54 ± 0.25; 2 animals, 4
sessions), providing further evidence of the tight link between preparatory dynamics and uninstructed movements.
216

Uninstructed movements related to behavioral context

We next compared uninstructed movements across task blocks in the two-context paradigm (**Fig. 4**). We found
that animals perform qualitatively different uninstructed movements in each behavioral context, despite both
contexts requiring instructed movements to the same targets (**Fig. 4a**). Context could be decoded from kinematic
features of movement across trial epochs – including during the ITI, in which all external contextual cues are
absent (**Fig. 4b,** *top*). The time-course of context decoding from neural population data and from movement
kinematics was also similar (**Fig. 4b,** *bottom*).

We sought to understand whether trial-to-trial variability in the neural representation of context-encoding could be
predicted from uninstructed movements. We again trained a multiple linear regression model to predict single-
trial projections along $CD_{context}$ from kinematic features and found that, like putative choice and urgency
dynamics, context-selective signals could be predicted with high fidelity (average R^2 across sessions – **Fig. 4c,d**;
$CD_{context}$: $R^2 = 0.49 \pm 0.22$; 6 animals, 12 sessions).

Taken together, these findings suggest that dynamics along CD_{choice}, CD_{ramp}, and $CD_{context}$ could be associated with
uninstructed movements and/or the cognitive processes related to the anticipation and planning of upcoming
instructed movements and encoding of task context.

Figure 3 - Uninstructed movements are tightly linked to putative planning dynamics. a. Choice-dependent uninstructed movements. Overlayed trajectories from a random subset of DR trials in an example session. Uninstructed movements are prominent on left, but not right, trials during the delay epoch. **b.** Choice (left vs. right) decoding from kinematic features and neural population data. Gray lines denote shuffled choice labels. Mean and 95% CI of across sessions/shuffles shown. **c.** *Top*, averaged projection onto CD$_{choice}$ on left (*red*) and right (*blue*) trials for the example session depicted in (a). Shaded area, 95% CI across trials. *Bottom*, average motion energy for all correct left and right trials. **d.** Observed and video predictions of single-trial projections of neural activity onto CD$_{choice}$ (see **Methods**) for the same example session as in (a). *Top*, Heatmap of actual and predicted single-trial projections. *Middle*, trial-averaged CD$_{choice}$ projections and predictions. *Bottom*, scatter plot of the average delay epoch projection of neural data onto CD$_{choice}$ versus corresponding video predictions. Dots denote single trials and dashed line denotes linear fit. *Inset*, R² values for all sessions (n = 25). Open circle denotes the example session in (a). **e.** Same as (c) but for CD$_{ramp}$ (different example session). Left and right hits are grouped (*purple*). **f.** Same as (d) but for CD$_{ramp}$. Same session as (e). **g.** *Top*, schematic of the randomized delay task. The delay epoch duration was randomly selected from six values (see **Methods**) with frequencies following an exponential distribution. *Middle*, motion energy averaged across sessions for randomized delay (*gray*) and fixed delay (*purple*) tasks (n = 12 sessions, *randomized delay*; n = 25, *fixed*). *Bottom*, selectivity of all single units pooled across sessions (n = 41 units, *randomized delay*; n = 283, *fixed*). Vertical lines indicate time of delay onset (black), go cue for the fixed delay (*purple*) and or go cue for the randomized delay (*gray*). Only trials with a delay duration of 1.2 s are shown for the randomized delay task for clarity. Shaded areas represent 95% CI in all panels. **h.** Same as (d) for an example session of the randomized delay task. *Top*, trials with delay durations of 0.3, 0.6, 1.2, and 1.8 s are shown. *Middle* and *bottom*, trials with delay duration of 1.2 s only.

Figure 4 - **Uninstructed movements are closely related to the neural encoding of context. a.** Schematic of tasks (*top*) and context-dependent uninstructed movements. Jaw/nose/paw movements on an example trial (*middle*) and overlayed trajectories of uninstructed movements across an example session (*bottom*) for DR task (*left*) and WC task (*right*). **b.** Context decoding from high-speed video (*top*) and neural data (*bottom*) as a function of time across all sessions. Gray lines denote shuffled context labels; shaded area, 95% confidence intervals (CI) across sessions. **c.** *Left*, heatmap of single-trial projections of neural data onto $CD_{context}$. *Right*, $CD_{context}$ projections predicted from video in an example session. **d.** *Left*, trial-averaged projection of neural data onto $CD_{context}$ and predictions from video (*dotted lines*) for the example session illustrated in (c). Shaded area, 95% CI across trials. *Right*, scatter plot of average projection of neural data onto $CD_{context}$ during the ITI versus average video prediction. Dots denote single trials; dashed line, line of best fit. *Inset*, R^2 values for all sessions (n = 12). Filled circle denotes the example session in (c).

Subspace decomposition of neural activity

We next examined whether neural signals responsible for the encoding of task variables could be isolated from movement-related signals, despite their correlation in time. We hypothesized that if the neural dynamics associated with cognitive and motor processes are separable, their components in the ALM should be driven by distinct latent signals. Previous theoretical work, later confirmed in primates, suggests that movement planning and execution dynamics can be confined to orthogonal 'null' and 'potent' subspaces of neural activity, respectively[39,40,50]. According to this model, changes in activity along potent dimensions (within the potent subspace) lead to changes in muscle activation, and, thus, movement. Changes along null dimensions (within the null subspace) relate to cognitive processes. We applied this computational framework to determine whether neural dynamics associated with instructed and uninstructed movement could be separated from dynamics associated with the encoding of choice, urgency, and context. Null and potent subspaces were first identified using trial-averaged data from DR and WC trials within each session (see **Fig. 5a** and **Methods**) using procedures described previously[40]. In this formalism, null and potent subspaces are constrained to be orthogonal and are jointly identified with the objective of maximizing the variance explained in the delay epoch (during motor planning) and response epochs (during movement), respectively.

We found that dynamics within null and potent subspaces were confined to the delay and response epochs, respectively, closely resembling results observed in primates (**Fig. 5b-d**). Surprisingly, examining the relationship between activity within the null and potent subspaces and uninstructed movements, we observed that activity within both subspaces was highly correlated with uninstructed movements on single trials (**Fig. 5b,e,f**). One likely explanation for this observation is that decomposing trial-averaged data in this way makes two assumptions that are unlikely to hold. First, this approach assumes that planning- and movement-related processes are strictly

283   confined to distinct task epochs. Uninstructed movements, however, were expressed across all task epochs (**Figs.**
284   **1,3,4**). Second, this approach assumes that cognitive processes and movement do not co-occur in time – for
285   example, that animals did not express uninstructed movements while planning their response or engage cognitive
286   processes during response epoch movements. To address these shortcomings, we sought an analytical approach
287   for identifying null and potent subspaces that avoids these assumptions.

Subspace decomposition of single-trial neural activity

We reasoned that the potent subspace should be defined using single-trial data to account for the trial-to-trial
variability in instructed and uninstructed movements. We used motion energy to identify when animals were
moving and stationary on a moment-to-moment basis (**see Methods**). Then, null and potent subspaces were
identified as the subspaces that maximized variance of neural activity during periods of stationarity and
movement, respectively, during all task epochs (**Fig. 5h**). Following this approach for subspace identification, we
found that neural dynamics in the potent subspace showed high similarity to motion energy on single trials.
Activity in the null subspace was only weakly correlated with movement (**Fig. 5i,l,m**).

Uninstructed movements were commonly observed during the delay epoch of DR trials, during motor planning,
and were highly correlated with putative choice and urgency signals (**Fig. 3c-h**). We examined whether the potent
subspace inadvertently captured choice or urgency dynamics that are correlated in time with movements. To
examine this possibility, we estimated null and potent subspaces using trials from the WC context only. Here, the
potent subspace is determined in a context in which choice and urgency signals are absent[23] (**Fig. 2d, EDFig. 3c-**
f), precluding the possibility that choice and urgency dynamics are inadvertently captured by the potent subspace.
This approach yielded estimates of the null and potent subspaces that were highly similar (**EDFig. 3d-f**). Further,
examining null and potent subspaces estimated only using data recorded during the response epoch of both tasks,
when planning and urgency dynamics should be minimal, again yielded similar results (**EDFig. 3g-i**).
Importantly, these observations suggest that the potent subspace indeed captures movement-related signals, and
not separable cognitive signals correlated in time with movements. We also determined whether null and potent
subspaces could be better estimated from latent dynamics estimated using factor analysis (FA), Gaussian
processes factor analysis[73] (GPFA), and latent factor analysis via dynamical systems[74] (LFADS), but found that
these approaches introduced aberrant dynamics (**see Methods**).

We hypothesized that activity in the null and potent subspaces, estimated on single trials, should capture variance
in neural activity across all task epochs. Animals may think and move simultaneously; thus, it follows that
cognitive dynamics may be present during the response epoch. Additionally, uninstructed movements occur
during both the sample and delay epochs, concurrent with stimulus and choice encoding - thus there should be
potent dynamics during all task epochs. As expected, activity within null and potent subspaces captured variance
across epochs (**Fig. 5k**). Interestingly, the proportion of units with activity largely confined to the potent subspace
was much larger than expected by chance (**Fig. 5n** and **EDFig. 4b;** p=6x10^{-9}; two-sample Kolmogorov-Smirnov
test), and much larger than when subspaces were identified from trial-averaged data here (**Fig. 5g**) and in non-
human primates[39]. This observation indicates the identification of cellular populations engaged solely in motor
processes that could not be identified without properly accounting for uninstructed movements.

Cognitive and motor dynamics during motor planning

The potent subspace may contain three classes of movement-related neural dynamics: (1) motor commands (and
efferent copies of motor commands), (2) sensory feedback related to ongoing movements, and (3) cognitive
signals that occur *exclusively during periods of movement*. Therefore, we reasoned that if a particular cognitive
process was "embodied," in the sense that it is always associated with some externally visible movement, then we
would expect to find its associated neural dynamics wholly contained within the potent subspace (**Fig. 6c**). In
contrast, cognitive signals driven by processes which are independent of movements should be wholly confined to
the null subspace (**Fig. 6a**). Between these extremes, we expect to observe dynamics within both subspaces when
a cognitive process influences the expression of uninstructed movements that are driven by distinct latent
dynamics (**Fig. 6b**).

Figure 5 – Subspace decomposition of neural activity using trial-averaged and single-trial data. a. Schematic of the approach for
estimating null (*green*) and potent (*pink*) subspaces from trial-averaged neural data. Null and potent subspaces are determined as the
orthogonal subspaces that maximally contain the variance of neural activity in the delay and response epochs, respectively. Trials from
both DR and WC contexts were used to identify subspaces. **b.** Activity during DR trials within the null and potent subspaces determined
from trial-averaged data. *Left*, motion energy across trials. *Middle*, sum squared magnitude of activity in the potent subspace. *Right*, sum
squared magnitude of activity in the null subspace. Trials sorted by average delay epoch motion energy. **c.** Sum squared magnitude of
activity in null and potent subspaces during DR lick-left and lick-right trials for an example session. Mean and standard error across trials
shown. **d.** Normalized variance explained of the neural data during DR and WC trials by the activity in the null and potent subspaces
during the delay and response epochs. Null and potent subspaces selectively capture activity in the delay and response epochs, as expected.
e. Variance explained (R^2) of motion energy by the sum squared magnitude of activity in the null and potent subspaces on DR and WC
single trials. Each point represents the mean across trials for a session. **f.** Cross-correlation between motion energy and activity in the null
and potent subspaces on DR and WC single trials. Mean and 95% CI across sessions shown. **g.** Subspace alignment for all single units
across sessions (see **Methods**). Values closer to 1 indicate that more variance of a unit's activity is contained within the null subspace. **h.**
Schematic of the approach for estimating null and potent subspaces from single-trial data. Motion energy is used to annotate when an
animal is moving (*green*) or stationary (*pink*). These labels are then applied to single-trial neural data. Null and potent subspaces are
determined as the orthogonal subspaces that maximally capture the variance of neural activity during periods of quiescence and movement,
respectively. Trials from both DR and WC contexts were used to identify subspaces. **i-n.** Same as (b-f) but using the single-trial approach
for estimating null and potent subspaces.

To evaluate these possibilities, we examined whether putative cognitive signals (illustrated in **Fig. 2**) evolved within the potent and/or null subspaces. We found choice selectivity – dynamics that differentiate left and right DR trials – in both the null and potent subspaces, suggesting that the encoding of choice is separable from dynamics related to the execution of choice-related uninstructed movements. Interestingly, the time course of choice dynamics in

Figure 6 – Schematic of potential relationships between cognitive and movement-related dynamics. a. Some cognitive and movement-related dynamics may be separable and independent from one another. *Top*, toy schematic where a cognitive process and a motor process are each governed by a single latent variable (L_1 and L_2). In this case, the value of the latent cognitive variable (L_1) is independent of the latent motor variable (L_2). *Bottom*, cognitive dynamics, denoted by the green circle, are governed by (black arrow) the latent factor, L_1, while the movement-related dynamics, denoted by the pink circle are governed by a separate, independent latent factor, L_2. **b.** Same as (a) for the separable and dependent case. The value of the latent cognitive variable (L_1) biases (grey dotted arrow) the value of the latent motor variable (L_2). **c.** Same as (b) for the inseparable case. Cognitive and motor dynamics are governed by a single latent variable such that these processes have a nearly one-to-one relationship.

each subspace was distinct, as was their expression on error trials. Selectivity emerged quickly following stimulus presentation in the null subspace and changed little through the delay epoch (**Fig. 7a;** $p=0.07$; paired *t*-test comparing selectivity during last 100ms of sample and delay epochs) consistent with a cognitive signal driven by available sensory evidence. On error trials, selectivity in the null subspace initially followed the same trajectory as correct trials but collapsed following stimulus offset. In the potent subspace, in contrast, selectivity continued to increase monotonically during the delay epoch ($p=3\times10^{-13}$; paired *t*-test comparing selectivity during last 100ms of sample and delay epochs), and no potent subspace selectivity emerged on error trials. These observations suggest that sensory stimuli initially drive appropriate dynamics within the null subspace on error trials, but that choice is not encoded stably and does not engage potent dynamics related to uninstructed movements.

Although the trajectory of choice dynamics in each subspace shared some similarities in trial-averaged data (**Fig. 7a**), they differed markedly on single trials in each subspace. The onset and magnitude of activity along the component of CD_{choice} within the potent subspace tracked trial-type selective motion energy on a moment-by-moment basis (**Fig. 7b, c**). Activity along the component of CD_{choice} in the null subspace tracked choice independently of ongoing uninstructed movements and, importantly, independently of activity within the potent subspace, suggesting that choice-correlated movements are not a result of planning dynamics that 'leak' into the potent subspace. Importantly, we found that many single neurons contribute to either the null or potent subspace representations of CD_{choice}, but not both ($p=1\times10^{-4}$, dip=0.04; Hartigan's dip test of unimodality), providing additional evidence of dynamics that are not only separable, but encoded by distinct populations of neurons (**Fig. 7d**) within the ALM microcircuit.

385
Figure 7 – Subspace decomposition allows for the re-examination of population measures of motor planning. a. Selectivity (neural
projections on lick-right trials minus lick-left trials) of null (*left*) and potent (*right*) subspace activity when projected along CD_{choice}. Mean
and 5-95% CI of the bootstrap distribution for correct (*solid*) and error (*dashed*) trials shown. Horizontal bars (solid lines, correct trials;
solid lines with black dashes, incorrect trials) above plot indicate when the selectivity trace is significantly different from zero ($p<0.05$,
one-sided test, bootstrap). **b.** Projections along CD_{choice} in the potent subspace (*middle*), but not the null subspace (*right*), follow the time-
course and magnitude of motion energy (*left*) on single lick-left and lick-right trials (example session). **c.** Variance explained (R^2) of
motion energy by projections along CD_{choice} in the null or potent subspace on single trials. Each point is the average across all trials in a
session. **d.** Distribution of single unit alignment with CD_{choice} (see **Methods**). This distribution was significantly different than that
expected from a unimodal distribution, indicating the presence of heterogeneous populations contributing to either the null or potent
representations of choice ($p=1\times10^{-4}$, dip=0.04; Hartigan's dip test of unimodality). **e.** Projection of null (*left*) and potent (*right*) subspace
activity along CD_{ramp} on lick-left and lick-right trials. Mean and 95% CI of the bootstrap distribution for correct trials shown. **f-h**. Same as
(b-d) for CD_{ramp}. **h.** Distribution of single unit alignment with CD_{ramp}. This distribution was not significantly different than that expected
from a unimodal distribution ($p=0.3$, dip=0.02; Hartigan's dip test of unimodality).

Taking a similar approach, we found that ramping dynamics were mostly contained within the potent subspace
(**Fig. 7e**) – perhaps surprising given that ramping activity has been interpreted as an urgency or timing signal[69–72].
This observation remained consistent when estimating null and potent subspaces using only WC trials or using
only the response epoch of WC and DR trials (**EDFig. 6a, 6d**), when ramping dynamics were not explicitly
observed. As a further test of the existence of null-subspace ramping dynamics, we explicitly identified the
dimension in the null (and potent) subspace that maximized ramping dynamics – which could be different than the
dimension that maximized ramping within the full activity space – and again failed to identify a prominent
ramping signal (**EDFig. 6c**). The relative paucity of ramping dynamics in the null subspace suggests that they are
not readily separable from associated movements in the ALM (see **Discussion**).

A persistent null-space representation of context.

Next, we examined projections of activity along $CD_{context}$ in each subspace (**Fig. 8a**). We found robust contextual
selectivity in both subspaces consistent with the interpretation that both context-correlated movements and a
cognitive representation of context and are both encoded within the ALM. The transient, response-epoch
dynamics observed along $CD_{context}$ (**Fig 2d,** *far right*) were entirely contained within the potent subspace (**Fig. 8a**)
likely indicative of subtle context-specific differences in instructed movements. We then compared context
selectivity (the difference in projections onto $CD_{context}$ on DR vs. WC trials) on trials with high motion energy
during the ITI ('Move' trials) and trials with minimal ITI motion energy ('Quiet' trials). Context selectivity within
the null

419
Figure 8 – A persistent, cognitive representation of context in the null subspace. a. Projections along $CD_{context}$ found using all neural
activity (*top*), activity in the null subspace (*middle*), or the potent subspace (*bottom*). Solid lines denote the mean projection across sessions
(n = 12) for DR hits (*purple*) and WC hits (*orange*); shaded area, 95% CI across sessions. Transient response epoch dynamics are absent in
the null subspace. **b.** Selectivity in $CD_{context}$ projections (DR – WC trials) on 'move' (*solid lines*) or 'quiet' (*dashed*) trials. Lines denote the
mean selectivity across sessions (n = 12) in the full space of neural activity (*top*), activity within the null subspace (*middle*), and potent
subspace (*bottom*); shaded area, 95% CI across sessions. Yellow region denotes ITI used to define 'move' and 'quiet' trials. **c.** Average
$CD_{context}$ selectivity during the ITI (yellow shaded region in *(b)*) for 'move' (*filled bars*) vs. 'quiet' (*open bars*) trials. Dots denote single
sessions; bars, mean across sessions. Asterisks denote significant differences ($p < 0.01$) in $CD_{context}$ selectivity between 'move' and 'quiet'
trials (*p*-values from left to right, Full population: $p = 0.005$, Null: $p = 0.5$, Potent: $p = 2 \times 10^{-4}$; *paired t-test*).

subspace was indistinguishable in high- and low-movement trials. In contrast, context selectivity within the potent
subspace was considerably larger on 'Move' trials, demonstrating that a null – but not a potent – representation of
context persists in the absence of movement (**Figs. 8b,c**; Selectivity on 'Move' vs. 'Quiet' trials – Full population:
$p = 0.005$, Null: $p = 0.5$, Potent: $p = 2 \times 10^{-4}$; paired *t*-test). These observations suggest that the analysis of context-
dependent dynamics without subspace decomposition indeed spuriously conflated separate latent dynamics
responsible for a cognitive representation of context (contained within the null subspace) and related to context-
dependent movements (contained in the potent subspace). Decomposing neural activity into null and potent
subspaces enables the identification of separable motor and cognitive components of contextual encoding.

Encoding of tongue kinematics in the potent subspace

Finally, we asked whether trial-type dependent activity following the go cue relates to movement, according to
previous suppositions[58,63] (**EDFig. 7**). Left-right selectivity along CD_{action} existed within both the null and potent
subspaces (**EDFig. 7a**), although the magnitude of selectivity in the null subspace was less than that within potent
subspace. On error trials, the activity along CD_{action} flipped to resemble that of the other trial type in the null
subspace but flipped asymmetrically within the potent subspace. This asymmetry closely corresponded to an
asymmetry in tongue angle, with incorrect movements directed to less extreme angles when directed to the left, on
average (**EDFig. 7b,c**). The component of CD_{action} within the potent subspace also better tracked moment-to-
moment changes in tongue angle during the response epoch ($p = 2 \times 10^{-8}$; paired *t*-test) (**EDFig. 7d,e**). The
similarity between tongue angle and dynamics along CD_{action} again suggests that neural dynamics in the potent
subspace relate to movement kinematics. The interpretation of action dynamics within the null subspace remains
unclear but could relate to the motor plan for directional licking.

Discussion

We aimed to understand whether the neural correlates of three cognitive variables commonly examined in
sensorimotor decision-making tasks – choice, urgency, and context – were separable from the neural encoding of
related uninstructed movements in the mouse ALM. We addressed this question by adapting an analytical
formalism in which neural activity is decomposed into orthogonal subspaces – a movement-potent subspace
containing dynamics related to the execution of movements and a complementary movement-null subspace
containing dynamics related to cognitive processes[39,40,50]. Extending upon this framework allowed us to consider
single-trial neural data recorded in the presence of uninstructed movements that can vary dramatically across trials
in their timing and identity (**Fig. 5**).

Separability of choice, urgency, and context encoding from movement-dynamics

Using this approach, we demonstrated that the ALM contains neural dynamics related to encoding of choice and
context that could be readily separated from movements, despite both cognitive variables being strongly
correlated with movements in time (**Fig. 7** and **Fig. 8**). Choice-selective signals were present in both null and
potent subspaces (**Fig. 7a**), consistent with the interpretation that a cognitive representation of choice within the
null subspace biases the probability and identity of uninstructed movements in a choice-selective manner.
Supporting this interpretation, null and potent subspace dynamics followed similar trajectories on average but
differed markedly at the single-trial level (**Fig. 7b**). Further, null subspace encoding of choice exhibited features
absent from encoding within the potent subspace. Choice encoding strengthened only with the availability of
sensory evidence, while potent subspace encoding continued to increase monotonically after sensory stimuli were
removed, mirroring the temporal evolution of animal's uninstructed movements. On error trials, stimulus-driven
choice encoding initially evolved correctly – but only within the null subspace – before decaying prior to the go
cue (**Fig. 7a**). Together, these results suggest cognitive and motor processes that are governed by separable latent
processes (**Fig. 6b**).

Context was stably encoded within the null subspace during all trial epochs and the inter-trial interval (**Fig. 8a-c**).
Although context was strongly represented in the potent subspace as well, these dynamics were abolished in the
absence of movement and were modulated by animals' instructed movements. These results suggest a persistent
cognitive encoding of context in the null subspace, and distinct potent-subspace dynamics related to context-
dependent movements, again indicative of related, but separable, latent processes (**Fig. 6b**).

Surprisingly, trial-averaged ramping dynamics proposed to underlie urgency (or timing)[69–72] were principally
represented in the potent subspace (**Fig. 7e**) indicating the possible absence of a cognitive representation of
urgency that can be dissociated from movements in the ALM. Although it remains possible that one exists in
another brain region, our results suggest the embodiment of urgency in our behavioral paradigm[75,76].

In contrast to these cognitive variables, encoding of kinematic features of movements, which were not used to
determine subspaces, was confined to the potent subspace (**EDFig. 7**)

Subspace decomposition

The approach to subspace decomposition presented here represents a means for assessing the issue of separability,
and for isolating separable dynamics into motor- and non-motor components for further analyses across a wide
range of experimental preparations. This method only requires annotation of when animals are moving and
utilizes a robust analytical approach not sensitive to parameter tuning.

A number of alternative methods have been proposed to evaluate neural dynamics associated with cognitive
processes in the presence of movement. The most common approach, which assumes separability of cognitive and
motor dynamics, is to parameterize movements as fully as possible and 'regress them out.' However,
parameterization of movement, particularly the orofacial and postural movements that have been associated with
strong, brain-wide neural dynamics[5,30,31], can be challenging. Further, this approach typically assumes a linear
relationship between neural activity and kinematic (or electromyographic) variables – an assumption unlikely to
hold for common movements mediated by central pattern generators and other highly nonlinear mechanisms[77–79].
Subspace decomposition is straightforward to implement under a range of experimental conditions and does not
preclude complex, nonlinear relationships between neural activity and movement variables, as expected for many

orofacial movements[77]. Methods for properly interpreting cognitive signals in the presence of related movements
on single trials will be vital for increasingly common experimental paradigms examining freely moving animals
performing complex, naturalistic behaviors[80–90].

Determination of null and potent subspaces as the orthogonal subspaces containing the most variance in neural
activity during periods of stationarity and movement is conservative in the assignment of dynamics to the null
subspace. It is more likely that cognitive dynamics are misassigned to the potent subspace than movement-related
dynamics to the null subspace. For example, dynamics associated with "embodied" cognitive processes that
always occur during periods of movement will only be represented in the potent subspace. Thus, the existence of
dynamics within the null subspace are highly likely to indicate cognitive signals. Nevertheless, we found that the
potent subspace captures similar dynamics when estimated from data recorded during the response epoch
('instructed' movements) or from WC trials, where uninstructed movements occur in the absence of choice and
urgency signals (**EDFig. 3, 6a,d**). This observation argues against the possibility that additional cognitive
dynamics associated with choice and urgency were inadvertently assigned to the potent subspace due to their
correlation in time with uninstructed movements.

That many individual neurons contribute solely to one subspace (**Fig. 5n, 6d,h**) suggests that the complementary
subspaces we identify map to distinct cell types within the underlying cortical circuit[62,91]. This observation that
individual neurons only appear to code for either cognitive or motor variables when uninstructed movements are
accounted for (**Fig. 5n, 6d,h**) underscores the importance of properly considering movements in future work
focused on the roles of anatomically- or transcriptomically-defined cell-types in neural computation.

Our approach to subspace decomposition makes several simplifying assumptions. First, movements of the
posterior torso, hindlimbs, or tail of the animal were not captured via videography. It remains possible, albeit
unlikely, that animals commonly perform movements confined to these portions of the body without concomitant
movement of the forepaws, head, neck, face, or whiskers[26]. Periods of stationarity were no doubt algorithmically
classified imperfectly and could have contained subtle movements below our threshold for detection. Second, we
did not consider potential time lags between motion energy captured via videography and the dynamics associated
with motor and/or sensory signals[92]. However, these time lags, on the order of a few tens of milliseconds, should
be much shorter than the timescale of dynamics explored in this study. Third, the set of signals related to
movement and/or cognitive processes may also be better described as occupying nonlinear manifolds rather than
linear subspaces[93,94]. Subspaces (or manifolds) could also shift dynamically during behavior following changes in
the activation of upstream or downstream brain regions[95,96] or changes in context[16,48,97]. Considering this
additional complexity may improve estimates of the dynamics associated with specific neural processes.

Variable relationships between cognition and movement

The dynamics associated with some movements and cognitive processes may be largely independent (**Fig. 6a**) –
perhaps in the case of the locomotor patterns of an individual thinking about where they might spend an
upcoming holiday while walking down the street. On the other hand, some cognitive processes may be embodied,
in the sense that they are tightly linked to externally observable changes in the body (**Fig. 6c**), such as the
relationship between arousal and changes in pupil diameter[98–100]. Between these extremes may be movements
which are correlated with, but separable from, cognitive processes. The probability and identity of uninstructed
movements in this study appeared related to choice and context (**Fig. 6b**), but with a high degree of trial-to-trial
and moment-to-moment variability unlikely to reflect commensurate variability in the cognitive processes to
which they relate. Correlations between cognitive processes and movements may also differ considerably in
highly trained, head-fixed animals compared to naturalistic settings[83,88]. This high degree of variability in these
relationships underscores the importance of developing and utilizing tools for assessing whether the neural
activity supporting cognitive processes can be dissociated from those related to movements in particular
experimental paradigms of interest.

The tight link between cognition and movement might suggest that some cognition-associated movements may be
beneficial for behavior. For example, postural adjustments may be highly related to decisions or motor plans,
enabling faster reaction times and/or more accurate movements[5,33]. Movements are also essential for some

cognitive processes – those supporting active sensation of the environment surely facilitate the construction and
continual updating of internal models of the environment[30,46]. However, just as separability may not be consistent
across movements and cognitive processes, not all cognition-associated movements may have a functional
role[30,101]. A poker player's 'tell' may be highly related to their cognitive state, yet may not directly aid the player
in the game. Regardless of their functional relationship, understanding the separability of cognitive processes and
associated movements in a wide variety of experimental settings will be essential for the interpretation of neural
dynamics observed during behavior.

Methods

Animals
This study used data collected from 13 mice; both male and female animals older than 8 weeks were used. Six
animals were used for the two-context task and were either C57BL/6J (JAX 000664) or VGAT-ChR2-EYFP (-/-)
(bred in-house). An additional three animals, either C57BL/6J or VGAT-ChR2-EYFP (-/-) were trained only on
the delayed response task. A separate cohort of two animals were used in the randomized delay task (VGAT-
ChR2-EYFP (+/-). Finally, two VGAT-ChR2-EYFP (+/-) animals were used in optogenetics experiments. Mice
were housed in a 12-hour reverse dark/light cycle room with ad libitum access to food. Access to water was
restricted during behavioral and electrophysiology experiments, for details see section ***Mouse behavior.*** Sample
sizes for all experiments were determined based on what is standard in the field; sample sizes were not determined
using any statistical tests.
565
Surgical procedures
All performed surgical procedures were in accordance with protocols approved by Boston University Institutional
Animal Care and Use Committee. For post-operative analgesia, mice were given ketaprofen (0.1 mL of 1 mg/mL
solution) and buprenorphine (0.07 mL of 0.03 mg/mL solution) prior to the start of all surgical procedures. Mice
were anesthetized with 1-2% isoflurane and placed on a heating pad in a stereotaxic apparatus. Artificial tears
(Akorn Sodium Chloride 5% Opthalamic Ointment) were applied to their eyes and a local anesthetic was injected
under the skin (Bupivacaine; 0.1 mL of 5 mg/mL solution) above the skull. The skin overlying the skull was
removed to expose the ALM (AP: +2.5 mm, ML: +1.5 mm), bregma, and lambda. The periosteum was removed
and the skin was secured to the skull with cyanoacrylate (Krazy Glue) around the margins. For electrophysiology
experiments, a headbar was implanted just posterior to bregma and secured with superglue and dental cement
(Jet™). Wells to hold cortex buffer during electrophysiology recordings were sculpted using dental cement and a
thin layer of superglue was applied over any exposed bone.

For optogenetics experiments, after headbar implantation, bone overlying frontal cortex was thinned bilaterally
with a dental drill and removed, exposing the frontal cortex. A glass window was then implanted over each
hemisphere and secured to the skull with cyanoacrylate[102]. Before performing photoinactivation experiments, any
bone regrowth was removed.
582
Mouse behavior
Following surgery, mice were allowed to recover for ~1 week and then water restricted, receiving 1.0 mL of water
per day. Behavioral training commenced after animals had been water restricted for 3-4 days. If animals received
less than 1.0 mL during training, they were supplemented with additional water.

All mice used in experiments were first trained on the delayed-response task with a fixed delay epoch until they
reached at least 70% accuracy. At the beginning of each trial, one of two auditory tones lasting 1.3 s were played;
the tone indicating a 'right' trial was white noise while the tone indicating a 'left' trial was a pure tone (8 kHz).
The delay epoch (0.9 s for 12 mice, 0.7 s for 1 mouse and linearly time warped to 0.9 s) started after the
completion of the sample tone. Following the delay epoch, an auditory go cue was played (swept-frequency
cosine [chirp], 10 ms) after which animals were rewarded with ~3 µL of water for contacting the correct lickport.
Lickport contacts before the response epoch resulted in the current task epoch restarting, ensuring that they could
not progress to the next task epoch until they had stopped licking for the specified length of the current epoch. If

the animal did not respond within 3 s of the go cue, this was considered an 'ignore' trial, although responses were
typically registered within 300 ms of the go cue.

Mice used for the two-context task *(***Figs. 1,2,4,8** and **EDFig. 1-3***)* were introduced to the water-cued context after
they were fully trained on the delayed-response context and at least two days before the first electrophysiology
recording session. A behavioral session began with ~100 DR trials and was then followed with interleaved blocks
of WC and DR trials. Each interleaved block was 10-25 trials, with no fixed block duration. All sessions started
with DR trials because sessions that began with WC trials had to be terminated early due to a high 'no response'
rate, likely due to the mouse becoming sated early in the session *(data not shown)*. In a WC trial, all auditory cues
were omitted and a ~3 µL water reward was presented randomly from either lickport. Trials in which the animal
contacted the lickport prior to the reward ('early lick') were omitted. Inter-trial intervals were randomly drawn
from an exponential distribution with mean 1.5 s. There were no explicit cues during the ITI to indicate to the
animal which context block it was in.

For the randomized delay task *(***Fig. 3g-h***)*, mice were first fully trained on the fixed delay DR task (delay duration
of 0.9 s). The delay epoch length was then randomized – the duration was randomly selected from six possible
values (0.3, 0.6, 1.2, 1.8, 2.4, and 3.6 s). The probability of each delay duration being selected was assigned such
that it resembled a cumulative distribution function of an exponential distribution with $\tau = 0.9$ s, as in ref [64].
Animals were trained on this version of the task until they became experts ($> 70\%$ accuracy and $< 20\%$ early lick
rate).
613
Videography analysis
High-speed video was captured (400 Hz frame rate) from two cameras (FLIR, Blackfly® monochrome camera,
Model number: BFS-U3-16S2M-CS). One provided a side view of the mouse and the other provided a bottom
view. We tracked the movements of specific body features using DeepLabCut[52] (**Figs. 1,3,4**). The tongue, jaw,
and nose were tracked using both cameras while the paws were only tracked using the bottom view. Position and
velocity of each tracked feature was calculated from each camera. The x and y position of each kinematic feature
was extracted from the output of DeepLabCut. Missing values were filled in with the nearest available value for
all features except for the tongue. The velocity of each feature was then calculated as the first-order derivative of
the position vector. Tongue angle and length were found using the bottom camera. Tongue angle was defined as
the angle between the vector pointing from the jaw to the tip of the tongue and the vector defining the direction
the mouse was facing. Tongue length was calculated as the Euclidean distance from the jaw to the tip of the
tongue.

Plots of kinematic feature overlays (**Fig. 1e, 3a, 4a and EDFig. 2**) were generated by plotting an [r, g, b] value for
each timepoint, *t*, where the values were specified by [KinFeature1$_t$, KinFeature2$_t$, KinFeature3$_t$]. All kinematic
features (speed or motion energy) were first standardized by taking the 99th percentile across time and trials and
normalizing to this value.
630
Motion energy
The motion energy for a given frame and pixel was defined as the absolute value of the difference between the
median value of the pixel across the next 5 frames (12.5 ms) and the median value of the pixel across the previous
5 frames (12.5 ms). Motion energy for each frame was then converted to a single value by taking the 99th
percentile across pixels. A threshold above which an animal was classed as moving, below which stationary, was
defined on a per session basis and set using a manual approach. Motion energy distributions, per session, were
bimodal, showing a large peak with little variance at low motion energy values and a second, smaller peak, with
high variance at large values. The threshold was set as the motion energy value separating these two mixtures. We
found that this method of setting the threshold captured both short and long duration movements. Alternative
methods such as defining baseline period of movements with which to compare against were less successful due
to the variability in timing of the movements across trials.
642
Photoinactivation experiments

Optogenetic photoinactivation was deployed on ~30% of trials selected at random. A 'masking flash' (470 mm
LED's LUXEON REBEL LED) controlled by an Arduino Teensy microcontroller (100 ms pulses, at 10 Hz) was
delivered constantly for the duration of the session to prevent mice from differentiating control and
photoinactivation trials. A 488 nm laser (Coherent OBIS 488 nm LX 150 mW Laser System) was used for all
photoinactivation experiments. ChR2-assisted photoinactivation (**Fig. 1b,c,g-i; EDFig. 1b-d**) was performed
through a cranial window[102] (see ***Surgical Procedures***), which replaced bone over the frontal cortex. Light was
delivered either at the start of the delay epoch, or at the start of the response epoch per session. A power density of
1.5 mW/mm^2 was used for all photoinactivation experiments.

For delay epoch photoinactivation (**Fig. 1g-i; EDFig. 1b**), light was delivered at the onset of the delay epoch and
lasted for 0.6 s followed by a 0.2 s ramp down. We targeted one of three regions per session: bilateral motor
cortex, bilateral ALM, or bilateral tjM1. A scanning galvo system (THORLABS GPS011) was used to
simultaneously target both hemispheres by scanning at 40 Hz. The beam (2 mm diameter) was centered around
the following locations: ALM (AP 2.5 mm, ML 1.5 mm), tjM1 (AP 1.75 mm, ML 2.25 mm), motor cortex (i.e.
ALM and tjM1; AP 2.25 mm, ML 1.85 mm). Due to their proximity, when specifically targeting ALM or tjM1,
Kwik-Cast™ (World Precision Instruments) was applied to the surrounding regions to prevent photoinactivation
of   other   regions.
For photoinactivation at the go cue (**Fig. 1b,c; EDFig. 1c,d**), light was delivered to the motor cortex for 0.8 s
followed by a 0.2 s ramp down beginning at the go cue presentation.

Electrophysiology recordings
Extracellular recordings were performed in the ALM using one of two types of silicon probes: H2 probes, which
have two shanks, each with 32 channels with 25 µm spacing (Cambridge Neurotech) or Neuropixels 1.0[103] which
have one shank and allow recording from 384 channels arranged in a checkerboard pattern (IMEC). For
recordings with H2 probes, the 64-channel voltage signals were multiplexed using a Whisper recording system
(Neural Circuits, LLC), recorded on a PXI-8133 board (National Instruments) and digitized at 14 bits. The signals
were demultiplexed into 64 voltage traces sampled at 25 kHz and stored for offline analysis.

At least 6 hours before recording, a small craniotomy (1-1.5 mm diameter) was made over ALM (AP 2.5 mm, ML
1.5 mm). 2-5 recordings were performed on a single craniotomy on consecutive days. After inserting the probes
between 900 and 1100 µm (MPM System, New Scale Technologies), brain tissue was allowed to settle for at least
5   minutes   before   starting   recordings.   All   recordings   were   made   using   SpikeGLX
(https://github.com/billkarsh/SpikeGLX).

Behavioral analysis
All sessions used for behavioral analysis had at least 40 correct DR trials for each direction (left or right) and 20
correct WC trials for each direction, excluding early lick and ignore trials, which were omitted from all analyses.
To assess the impact of ALM photoinactivation at the go cue, we first calculated the percent of trials with a
correct lick within 600 ms of the go cue (**Fig. 1c**). To find the fraction of time with the tongue visible during the
photoinactivation period (**EDFig. 1d**), for each trial we found the number of time-points within the 1 s after the go
cue where the tongue was labelled as visible by DeepLabCut and divided that by the total number of time-points
within the photoinactivation period. This was then averaged for all control or photoinactivation trials for a given
session.

To assess the impact of bilateral MC/ALM/tjM1 photoinactivation during the delay epoch, the average motion
energy during the delay epoch was calculated separately for right and left control vs. photoinactivation trials (**Fig.
1h, EDFig. 1b**).

Electrophysiology recording analysis
JRCLUST[104]   (https://github.com/JaneliaSciComp/JRCLUST)   and/or   Kilosort3
(https://github.com/MouseLand/Kilosort) with manual curation using Phy (https://github.com/cortex-lab/phy)
were used for spike sorting. For the DR task we recorded 1651 units (483 single units) in ALM from 25 sessions
using 9 mice. 12 of these sessions in 6 of the mice were utilized for the analyses of the two-context task. 522 units
(214 well-isolated single units) were recorded in these sessions. Finally for the randomized delay task we

recorded 226 units (41 well-isolated single units) in ALM from 4 sessions using 2 mice. All units were included
in analyses with the exceptions of subspace alignment (**Fig. 5g,n** and **Fig. 7d,h**) and single-unit selectivity
analyses (**Fig. 3g** and **EDFig. 3c**) where only well-isolated units were included.

To find choice-selective neurons (**Fig. 2**), forty trials were subsampled for both right and left correct trials and a
ranksum test was used to compare spike counts for each unit during the sample, delay, or response epochs.
Selective units were those with significantly different spike counts ($p < 0.05$, ranksum test). Context selective
units (**Fig. 2**) were defined in the same manner but comparing spike counts during the ITI (the 300 ms preceding
the sample tone onset).

Selectivity in the neural population (**Fig. 3g, EDFig. 3c**) was calculated as the difference in spike rate on
preferred – non-preferred trials in choice-selective units, with *preferred* trials referring to the trial type with a
higher spike rate for each unit.

Coding direction analysis
Coding directions (CD) were defined as directions in neural activity state space, defined by trial-averaged data,
that maximally separated trajectories of two conditions. Trial-averaged neural activity for each condition *c* was
formatted in a matrix $\mathbf{X}_c \in \square^{T \times N}$, where T is the number of time bins and N the number of units. Each $\mathbf{X}_c$ was
baseline subtracted and scaled by baseline standard deviation before estimating CDs.

CD_{choice} and CD_{action} were calculated as:
709

$$v = \frac{\overline{x}_{lickright} - \overline{x}_{lickleft}}{\sqrt{\mathrm{var}(\overline{x}_{lickright}) + \mathrm{var}(\overline{x}_{lickleft})}}$$

$$CD = \frac{v}{\sum |v|}$$

For each unit, the mean spike rate difference between right lick trials, $\overline{x}_{lickright}$, and left lick trials, $\overline{x}_{lickleft}$, was
calculated in a 400 ms time interval. CD_{choice} was calculated in the last 400 ms of the delay epoch and CD_{action} was
calculated in the first 400 ms of the response epoch. The vector v was then obtained by normalizing by the square
root of the sum of the variances across conditions. Finally, v was normalized by its L_1 norm to ensure projections
do not scale by the length of v, the number of units simultaneously recorded.

CD_{ramp} was calculated as:
717

$$v = \frac{\overline{x}_{delay} - \overline{x}_{ITI}}{\sqrt{\mathrm{var}(\overline{x}_{delay}) + \mathrm{var}(\overline{x}_{ITI})}}$$

$$CD = \frac{v}{\sum |v|}$$

720   This calculation was similar to the calculations of CD_{choice} and CD_{action} but incorporated data from all correct DR
721   trials during the last 400 ms of the delay epoch, $\overline{x}_{delay}$, and during the ITI (300 ms preceding the sample tone
722   onset), $\overline{x}_{ITI}$.

723   $CD_{context}$ was calculated as:
724

$$v = \frac{\overline{x}_{DR} - \overline{x}_{WC}}{\sqrt{\mathrm{var}(\overline{x}_{DR}) + \mathrm{var}(\overline{x}_{WC})}}$$

$$CD = \frac{v}{\sum |v|}$$

726
727
This calculation was similar to the calculation of other CDs but incorporated data from correct DR, $\overline{x}_{DR}$, and WC,
$\overline{x}_{WC}$, trials during the ITI.

CD$_{action}$ was orthogonalized to CD$_{choice}$ to exclusively capture response epoch selectivity that emerges after the go
cue. CD$_{ramp}$ was orthogonalized to CD$_{action}$ and CD$_{choice}$ to remove selectivity that may emerge during the response
epoch that is also captured in CD$_{action}$. Orthogonalization was performed using the Gram-Schmidt process.

Projections of the population activity along the CD were calculated as:
734

$$proj = CD^T * \overline{x}$$

735
For the randomized delay task, trials with non-1.2 s delay lengths were used to calculate CD$_{choice}$ (fit trials),
always using the last 600 ms of the delay epoch as the time interval for calculation[64]. Population activity from
trials with 1.2 s delays (test trials) were then projected along the CD$_{choice}$ (**Figure 3h - *middle***). For visualization
purposes, trials with all delay lengths (including fit trials) were projected along CD$_{choice}$ in **Figure 3h – *top*.**

Choice/context decoding from neural population or kinematic features
We trained logistic regression models to predict either animal choice (**Fig. 3b**) or behavioral context (**Fig. 4b**)
from either kinematic features or single trial neural activity. The kinematic regressors were made up of the x and y
positions and velocities of DLC tracked features (tongue, jaw, and nose), as well as the tongue length, angle, and
motion energy. The neural regressors were the firing rates of the units simultaneously recorded within each
session. Animal choice was encoded as -1 on right lick trials and 1 on left lick trials. Context was encoded as 1 on
DR trials and -1 on WC trials. Both correct and error trials were used for this analysis. A separate model was
trained at each time bin for both neural and kinematic decoding. Models were trained with ridge regularization
and 4-fold cross-validation with 30% of trials held out for testing. Chance accuracy was obtained by shuffling
choice/context labels across trials. Accuracy of the prediction was defined as:
750

$$Accuracy = \frac{Number\ of\ correct\ predictions}{Total\ number\ of\ predictions}$$

751
Predicting projections along coding directions from kinematic features
We trained support vector machines (SVM) to predict projections along CD$_{choice}$, CD$_{ramp}$, and CD$_{context}$. Correct
DR trials were used for predicting CD$_{choice}$ and CD$_{ramp}$ projections; correct DR and WC trials were used for
predicting CD$_{context}$. For predicting CD$_{context}$, the regressors were made up of the x and y positions and velocities of
DLC tracked features (tongue, jaw, and nose), as well as the tongue length, angle, and motion energy (totaling 60
regressors). Only the sample and delay epoch were considered when predicting CD$_{choice}$ and CD$_{ramp}$ projections;
because of this, any tongue-related metrics were not included as regressors in these analyses (totaling 55
regressors). Projections and kinematic features were first mean-centered and scaled to unit variance before input
to the regression model. The previous B bins of kinematic data were used to predict the current bin's neural data
(B=6, each bin is 5 ms). Models were trained with ridge regularization and 4-fold cross-validation, however
similar results were obtained with unregularized regression. The models were tested on held-out testing data (40%
of trials for CD$_{choice}$ and CD$_{ramp}$; tested on 30% of trials for CD$_{context}$). To assess the prediction quality between
projections and model predictions, we calculated the variance explained as:
765

$$R^2 = 1 - \frac{\sum_t (y_t - \hat{y}_t)^2}{\sum_t (y_t - \overline{y})^2}$$

where y_i is the value of the projection at time t, $\hat{y}_t$ is the prediction, and $\square$ is the mean of the projection across all
time.

When predicting projections along CD$_{\text{choice}}$ during the randomized delay task, unregularized models were used
(**Fig. 3h**). This was due to the small numbers of trials for each delay length which precluded the partitioning of
data into 3 separate sets (training, validation, and testing trials). Instead, models were trained with linear
regression and 4-fold cross-validation and tested on held-out data from each cross-validation fold. For assessing
prediction quality between projections and model predictions, trials with 1.2 s delay epochs were used.

To understand which groups of kinematic features were most informative for predicting projections along
CD$_{\text{choice}}$, or CD$_{\text{context}}$ (**EDFig. 2**), the absolute value of the beta coefficients for each kinematic regressor in the
trained model was first obtained. For each feature group (jaw, nose, motion energy), the average beta coefficient
across all regressors belonging to this kinematic feature group (*8*nBins* for jaw, *6*nBins* for nose, and *1*nBins*
for motion energy) was calculated. The average beta coefficient for each feature group was then expressed as a
fraction of the total of all beta coefficients.

Subspace identification
Null and potent subspaces were identified following the general approach described in ref[40]. For identifying
subspaces using trial-averaged data (**Fig. 5a**), neural activity was first trial-averaged within trial type. Activity
across trial types was then concatenated, providing a matrix $\mathbf{X} \in \square^{(T*C) \times N}$, where T is the number of time bins, C
the number of trial types, and N the number of units. Trial-averaged neural activity was soft-normalized, then
mean-centered as described in ref[40]. The matrix $\mathbf{X}$ was further divided into two matrices, $\mathbf{X}_{\text{null}}$ and $\mathbf{X}_{\text{potent}}$. $\mathbf{X}_{\text{null}}$
contained activity from [-1,0] seconds relative to the go cue, and $\mathbf{X}_{\text{potent}}$ [0,1] seconds relative to the go cue. Null
and potent subspaces were then identified by solving the following optimization problem:
789

$$[\hat{\boldsymbol{Q}}_{null}, \hat{\boldsymbol{Q}}_{potent}] = argmax_{[\boldsymbol{Q}_{null}, \boldsymbol{Q}_{potent}]} \; \frac{1}{2} \frac{Tr(\boldsymbol{Q}_{null}^T \boldsymbol{C}_{null} \boldsymbol{Q}_{null})}{\sum_{i=1}^{d_{null}} \sigma_{null}(i)} + \frac{1}{2} \frac{Tr(\boldsymbol{Q}_{potent}^T \boldsymbol{C}_{potent} \boldsymbol{Q}_{potent})}{\sum_{i=1}^{d_{potent}} \sigma_{potent}(i)}$$

$$subject \; to \quad \boldsymbol{Q}_{null}^T \boldsymbol{Q}_{potent} = 0, \quad \boldsymbol{Q}_{null}^T \boldsymbol{Q}_{null} = I, \quad \boldsymbol{Q}_{potent}^T \boldsymbol{Q}_{potent} = I$$

791
792
where $\mathbf{C}_{\text{null}}$ and $\mathbf{C}_{\text{potent}}$ are the covariances of $\mathbf{X}_{\text{null}}$ and $\mathbf{X}_{\text{potent}}$, σ_{null} and σ_{potent} are the singular values of $\mathbf{C}_{\text{null}}$ and
$\mathbf{C}_{\text{potent}}$, and d_{null} and d_{potent} are the dimensionality of the subspaces. This optimization problem jointly identifies null
and potent subspaces that maximize the variance explained during the delay (null) and response (potent) epochs.
The dimensionality of each space was chosen to be half the number of simultaneously recorded neurons, or
twenty, whichever was smaller. Therefore, the dimensionality of the full population was equal to the
dimensionality of the null or potent subspaces for sessions containing forty or fewer neurons. We capped
dimensionality of subspaces at twenty for computational tractability. Optimization was performed using the
manopt[105] toolbox for MATLAB.

For identifying subspaces using single-trial data (**Fig 5h**), single-trial neural activity was first binned in 5 ms
intervals and smoothed with a causal Gaussian kernel with an s.d. of 35 ms. $\mathbf{X}_{\text{null}} \in \square^{(T\_n*K) \times N}$ and $\mathbf{X}_{\text{potent}} \in \square^{(T\_p*K)}$
$^{\times N}$ were defined using the single trial neural activity when the animal was stationary or moving (see ***Motion***
Energy), respectively. T_n is the number of time bins annotated as the animal being stationary, T_p the number of
time points annotated as the animal moving, and K the number of trials used for subspace identification. All
correct and error trials from DR and WC contexts were used unless specified otherwise. Once the data was
partitioned into moving and stationary time points, subspace identification was carried out as described for trial-
averaged data.

Our decision to use binned and smoothed single-trial firing rates rather than inferring smooth single-trial firing
rates using techniques such as FA, GPFA[73] and LFADS[74] came from the observation that, when using these
techniques, an additional mode of dynamics not present in the original data was observed. To prevent our results

from being biased by the technique used to pre-process single-trial firing rates, we opted to used binned and
smoothed data instead.

We quantified the normalized variance explained in a subspace as in refs[40,106] (**Fig. 5d,k**):
815

$$normVE = \frac{Tr(\mathbf{Q}^T \mathbf{C} \mathbf{Q})}{\sum_{i=1}^{d} \sigma(i)}$$

816
817   where **Q** is the subspace, **C** is the covariance of neural activity, and σ are the singular values of **C**. This
818   normalization provides the maximum variance that can be captured by d dimensions.

819   Single units were reconstructed from null and potent subspaces according to the following equation:
820

$$\mathbf{X}_{recon} = \mathbf{X}_{subspace} \mathbf{Q}$$

821
where $\mathbf{X}_{recon}$ is the reconstructed neural activity, $\mathbf{X}_{subspace}$ is the projected neural activity within the null or potent
subspace, and **Q** is either the null or potent subspace.

Coding directions within null and potent subspaces were obtained by projecting $\mathbf{X}_{recon}$ along coding directions
obtained from the full neural population (**Fig. 7**).

To ask whether null and potent subspaces independently contained dynamics similar to that captured by coding
directions, but along a different direction contained wholly within one subspace, we separately identified coding
directions from null and potent subspace activity ($\mathbf{X}_{subspace}$). Activity within each subspace was then projected
along the respective coding directions (**EDFig. 5b,c**).
830
831   Subspace alignment for an individual unit was calculated as:
832

$$A = \frac{VE_{null} - VE_{potent}}{VE_{null} + VE_{potent}}$$

833
834   where VE is the variance explained of each individual unit by the null or potent subspace, or by the activity along
835   a coding direction within the null or potent subspace. VE for a single unit was calculated as:
836

$$R^2 / VE = 1 - \frac{\sum_t (\mathbf{x}_t - \hat{\mathbf{x}}_t)^2}{\sum_t (\mathbf{x}_t - \overline{\mathbf{x}})^2}$$

where x_t is the trial-averaged firing rate, $\hat{\mathbf{x}}$ the reconstructed trial-averaged firing rate, and t the time bin.

To ask if the subspace alignment distributions were different than expected by chance, we compared the
distributions to those obtained from null and potent subspaces sampled randomly as described in ref [40]. Each
element comprising a subspace were randomly sampled from a normal distribution with zero mean and unit
variance but were biased by the covariance structure of the neural activity for a given session. Neural activity
from each session was projected along the randomly sampled subspaces and alignment indices were calculated.
This procedure was repeated 1000 times to generate null distributions (**EDFig. 4**).

Subspaces were identified using data from DR and WC correct and error trials (**Fig. 5,7,8**). Control analyses were
performed using separate sets of trials to assess if the potent subspace erroneously contained null-subspace
dynamics. First, we estimated subspaces using WC trials only (**EDFig. 3, 5**). Second, we estimated subspaces
using DR and WC trials, but restricted time points used to those in the response epoch only (**ED Fig. 3,5**). Both
controls allowed us to estimate null and potent subspaces in the presence of uninstructed movements, but in the
absence of planning dynamics. These controls thus allowed us to measure the degree to which the potent subspace
erroneously captured null dynamics.

CD_{Context} selectivity on 'Move' vs. 'Quiet' trials

For each session, DR and WC trials were first separated into 'Move' and 'Quiet' trials. 'Move' trials were those where the average motion energy in the ITI was greater than the movement threshold for that session, 'Quiet' trials were those where it was lower than the movement threshold. The last 40 trials in each session were excluded from analysis to account for the decrease in uninstructed movements that are observed towards the end of behavioral sessions as animals become sated and disengaged. 50% of 'Move' trials and 50% of 'Quiet' trials were randomly sampled from DR and WC trials separately to ensure that 'Move' and 'Quiet' trials did not contain an overrepresentation of one context. These trials were then pooled together and used to calculate $CD_{Context}$ (fit trials). Population activity from the remaining 'Move' and 'Quiet' trials (test trials) was then projected along $CD_{Context}$. Selectivity was defined as the trial-averaged projection along $CD_{Context}$ on DR – WC trials (**Fig. 8b**).

Hierarchical bootstrapping

Projections along coding directions were obtained via a hierarchical bootstrapping procedure[107,108] (**Figs. 2, 7** and **EDFigs. 3, 4, 5**). Pseudopopulations were constructed by randomly sampling with replacement M mice, 2 sessions per sampled mouse, 50 correct trials of each type, 20 error trials of each type, and 20 neurons. M is the number of mice in the original cohort. Bootstrapping was repeated for 1000 iterations. For all figures, mean and 95% confidence intervals (shaded area) of the bootstrap distribution are shown. For selectivity, 5-95% confidence intervals are shown to indicate where projections significantly differ from zero ($p<0.05$, one-sided test, bootstrap).

Statistics

No statistical methods were used to determine sample sizes. All two-sample *t*-tests were Students *t*-tests. All *t*-tests were two-sided unless stated otherwise.

DATA AVAILABILITY
Primary and derived data described in this study will be made available on Figshare upon publication.

CODE AVAILABILITY
MATLAB code for subspace identification is available at https://github.com/economolab/subspaceID. Custom MATLAB code used for analyses will be made available on Github upon publication.

COMPETING INTERESTS
The authors declare no competing interests.

AUTHOR CONTRIBUTIONS
MAH, JEB, and MNE conceived of the project. MNE and CC supervised research. MAH, JEB, and MNE designed experiments. MAH, JEB, JLUN, and EKH performed experiments. MAH, JEB, and MNE analyzed data. MAH, JEB, CC, and MNE wrote the manuscript.

ACKNOWLEDGMENTS
The authors thank, Karel Svoboda, Brian DePasquale, Shaul Druckmann, and Ben Scott for helpful discussions. We thank Tim Wang for helpful comments on the manuscript. This work was supported by the Whitehall Foundation, the Klingenstein Fund, the Simons Foundation, and NIH R01NS121409.

References

1. Alexander, G. E. & Crutcher, M. D. Neural representations of the target (goal) of visually guided arm

    movements in three motor areas of the monkey. *Journal of Neurophysiology* **64**, 164–178 (1990).

2. Cisek, P. & Kalaska, J. F. Neural Correlates of Reaching Decisions in Dorsal Premotor Cortex:

    Specification of Multiple Direction Choices and Final Selection of Action. *Neuron* **45**, 801–814 (2005).

3.   Lebedev, M. A. & Wise, S. P. Insights into Seeing and Grasping: Distinguishing the Neural Correlates of

Perception and Action. *Behavioral and Cognitive Neuroscience Reviews* **1**, 108–129 (2002).

4.   Chandrasekaran, C., Peixoto, D., Newsome, W. T. & Shenoy, K. V. Laminar differences in decision-related

neural activity in dorsal premotor cortex. *Nat Commun* **8**, 614 (2017).

5.   Chen, S. *et al.* Brain-wide neural activity underlying memory-guided movement. 2023.03.01.530520

Preprint at https://doi.org/10.1101/2023.03.01.530520 (2023).

6.   Basso, M. A. & May, P. J. Circuits for Action and Cognition: A View from the Superior Colliculus. *Annu*

Rev Vis Sci **3**, 197–226 (2017).

7.   Zimnik, A. J. & Churchland, M. M. Independent generation of sequence elements by motor cortex. *Nat*

Neurosci **24**, 412–424 (2021).

8.   Ames, K. C., Ryu, S. I. & Shenoy, K. V. Simultaneous motor preparation and execution in a last-moment

reach correction task. *Nat Commun* **10**, 2718 (2019).

9.   Flash, T. & Henis, E. Arm Trajectory Modifications During Reaching Towards Visual Targets. *Journal of*

Cognitive Neuroscience **3**, 220–230 (1991).

10. Fuster, J. M. & Alexander, G. E. Neuron activity related to short-term memory. *Science* **173**, 652–654

(1971).

11. Romo, R., Brody, C. D., Hernández, A. & Lemus, L. Neuronal correlates of parametric working memory in

the prefrontal cortex. *Nature* **399**, 470–473 (1999).

12. Erlich, J. C., Bialek, M. & Brody, C. D. A Cortical Substrate for Memory-Guided Orienting in the Rat.

Neuron **72**, 330–343 (2011).

13. Tanji, J. & Evarts, E. V. Anticipatory activity of motor cortex neurons in relation to direction of an intended

movement. *Journal of Neurophysiology* **39**, 1062–1068 (1976).

14. Guo, Z. V. *et al.* Flow of Cortical Activity Underlying a Tactile Decision in Mice. *Neuron* **81**, 179–194

(2014).

15. Wallis, J. D., Anderson, K. C. & Miller, E. K. Single neurons in prefrontal cortex encode abstract rules.

Nature **411**, 953–956 (2001).

16. Mante, V., Sussillo, D., Shenoy, K. V. & Newsome, W. T. Context-dependent computation by recurrent
dynamics in prefrontal cortex. *Nature* **503**, 78–84 (2013).

17. Schultz, W., Dayan, P. & Montague, P. R. A Neural Substrate of Prediction and Reward. *Science* **275**,
1593–1599 (1997).

18. Watabe-Uchida, M., Eshel, N. & Uchida, N. Neural Circuitry of Reward Prediction Error. *Annual Review of*
Neuroscience **40**, 373–394 (2017).

19. Freedman, D. J., Riesenhuber, M., Poggio, T. & Miller, E. K. Categorical representation of visual stimuli in
the primate prefrontal cortex. *Science* **291**, 312–316 (2001).

20. Carr, M. F., Jadhav, S. P. & Frank, L. M. Hippocampal replay in the awake state: a potential substrate for
memory consolidation and retrieval. *Nat Neurosci* **14**, 147–153 (2011).

21. Levy, S. *et al.* Cell-Type-Specific Outcome Representation in the Primary Motor Cortex. *Neuron* **107**, 954-
971.e9 (2020).

22. Padoa-Schioppa, C. & Assad, J. A. Neurons in Orbitofrontal Cortex Encode Economic Value. *Nature* **441**,
223–226 (2006).

23. Zagha, E. The Importance of Accounting for Movement When Relating Neuronal Activity to Sensory and
Cognitive Processes | Journal of Neuroscience. https://www.jneurosci.org/content/42/8/1375#ref-90 (2022).

24. Drew, P. J., Winder, A. T. & Zhang, Q. Twitches, Blinks, and Fidgets: Important Generators of Ongoing
Neural Activity. *Neuroscientist* **25**, 298–313 (2019).

25. Galton, F. The Measure of Fidget. *Nature* **32**, 174–175 (1885).

26. Hulsey, D., Zumwalt, K., Mazzucato, L., McCormick, D. A. & Jaramillo, S. Decision-making dynamics are
predicted by arousal and uninstructed movements. 2023.03.02.530651 Preprint at
https://doi.org/10.1101/2023.03.02.530651 (2023).

27. Tremblay, S., Testard, C., DiTullio, R. W., Inchauspé, J. & Petrides, M. Neural cognitive signals during
spontaneous movements in the macaque. *Nat Neurosci* **26**, 295–305 (2023).

28. Hafed, Z. M., Lovejoy, L. P. & Krauzlis, R. J. Modulation of Microsaccades in Monkey during a Covert
Visual Attention Task. *J Neurosci* **31**, 15219–15230 (2011).

29.   Corneil, B. D., Munoz, D. P., Chapman, B. B., Admans, T. & Cushing, S. L. Neuromuscular consequences
of reflexive covert orienting. *Nat Neurosci* **11**, 13–15 (2008).

30.   Musall, S., Kaufman, M. T., Juavinett, A. L., Gluf, S. & Churchland, A. K. Single-trial neural dynamics are
dominated by richly varied movements. *Nat Neurosci* **22**, 1677–1686 (2019).

31.   Stringer, C. Spontaneous behaviors drive multidimensional, brainwide activity | Science.
https://www.science.org/doi/10.1126/science.aav7893 (2019).

32.   Steinmetz, N. A., Zatka-Haas, P., Carandini, M. & Harris, K. D. Distributed coding of choice, action and
engagement across the mouse brain. *Nature* **576**, 266–273 (2019).

33.   Mangin, E. N., Chen, J., Lin, J. & Li, N. Behavioral measurements of motor readiness in mice.
2023.02.03.527054 Preprint at https://doi.org/10.1101/2023.02.03.527054 (2023).

34.   Terada, S.-I., Kobayashi, K. & Matsuzaki, M. Transition of distinct context-dependent ensembles from
secondary to primary motor cortex in skilled motor performance. *Cell Reports* **41**, 111494 (2022).

35.   Lowet, E. *et al.* Enhanced Neural Processing by Covert Attention only during Microsaccades Directed
toward the Attended Stimulus. *Neuron* **99**, 207-214.e3 (2018).

36.   Popescu, S. T. & Wexler, M. Spontaneous Body Movements in Spatial Cognition. *Front Psychol* **3**, 136
(2012).

37.   Hajnal, J. V. *et al.* Artifacts due to stimulus correlated motion in functional imaging of the brain. *Magn*
Reson Med **31**, 283–291 (1994).

38.   Dominiak, S. E. *et al.* Whisking Asymmetry Signals Motor Preparation and the Behavioral State of Mice. *J.*
Neurosci. **39**, 9818–9830 (2019).

39.   Kaufman, M. T., Churchland, M. M., Ryu, S. I. & Shenoy, K. V. Cortical activity in the null space:
permitting preparation without movement. *Nat Neurosci* **17**, 440–448 (2014).

40.   Elsayed, G. F., Lara, A. H., Kaufman, M. T., Churchland, M. M. & Cunningham, J. P. Reorganization
between preparatory and movement population responses in motor cortex. *Nat Commun* **7**, 13239 (2016).

41.   Liu, T. T., Nalci, A. & Falahpour, M. The global signal in fMRI: Nuisance or Information? *NeuroImage*
150, 213–229 (2017).

971 42. Friston, K. J., Williams, S., Howard, R., Frackowiak, R. S. J. & Turner, R. Movement-Related effects in

972   fMRI time-series. *Magnetic Resonance in Medicine* **35**, 346–355 (1996).

973 43. Bullmore, E. T. *et al.* Methods for diagnosis and treatment of stimulus☐correlated motion in generic brain

974   activation studies using fMRI. *Hum Brain Mapp* **7**, 38–48 (1998).

975 44. Power, J. D., Schlaggar, B. L. & Petersen, S. E. Recent progress and outstanding issues in motion correction

976   in resting state fMRI. *NeuroImage* **0**, 536 (2015).

977 45. Cisek, P. & Kalaska, J. F. Neural Mechanisms for Interacting with a World Full of Action Choices. *Annual*

978   *Review of Neuroscience* **33**, 269–298 (2010).

979 46. Cisek, P. & Pastor-Bernier, A. On the challenges and mechanisms of embodied decisions. *Philosophical*

980   *Transactions of the Royal Society B: Biological Sciences* **369**, 20130479 (2014).

981 47. Koay, S. A., Charles, A. S., Thiberge, S. Y., Brody, C. D. & Tank, D. W. Sequential and efficient neural-

982   population coding of complex task information. *Neuron* **110**, 328-349.e11 (2022).

983 48. Aoi, M. C., Mante, V. & Pillow, J. W. Prefrontal cortex exhibits multidimensional dynamic encoding during

984   decision-making. *Nat Neurosci* **23**, 1410–1420 (2020).

985 49. Hennig, J. A. *et al.* Learning is shaped by abrupt changes in neural engagement. *Nat Neurosci* **24**, 727–736

986   (2021).

987 50. Druckmann, S. & Chklovskii, D. B. Neuronal Circuits Underlying Persistent Representations Despite Time

988   Varying Activity. *Current Biology* **22**, 2095–2103 (2012).

989 51. Galiñanes, G. L., Bonardi, C. & Huber, D. Directional Reaching for Water as a Cortex-Dependent

990   Behavioral Framework for Mice. *Cell Reports* **22**, 2767–2783 (2018).

991 52. Mathis, A. *et al.* DeepLabCut: markerless pose estimation of user-defined body parts with deep learning.

992   *Nat Neurosci* **21**, 1281–1289 (2018).

993 53. Mayrhofer, J. M. *et al.* Distinct Contributions of Whisker Sensory Cortex and Tongue-Jaw Motor Cortex in

994   a Goal-Directed Sensorimotor Transformation. *Neuron* **103**, 1034-1043.e5 (2019).

995 54. Inagaki, H. K. *et al.* Neural Algorithms and Circuits for Motor Planning. *Annual Review of Neuroscience*

996   **45**, 249–271 (2022).

55. Xu, D. *et al.* Cortical processing of flexible and context-dependent sensorimotor sequences. *Nature* **603**,

464–469 (2022).

56. Chen, T.-W., Li, N., Daie, K. & Svoboda, K. A Map of Anticipatory Activity in Mouse Motor Cortex.

Neuron **94**, 866-879.e4 (2017).

57. Yin, X., Wang, Y., Li, J. & Guo, Z. V. Lateralization of short-term memory in the frontal cortex. *Cell*

Reports **40**, 111190 (2022).

58. Li, N., Chen, T.-W., Guo, Z. V., Gerfen, C. R. & Svoboda, K. A motor cortex circuit for motor planning and

movement. *Nature* **519**, 51–56 (2015).

59. Esmaeili, V. *et al.* Rapid suppression and sustained activation of distinct cortical regions for a delayed

sensory-triggered motor response. *Neuron* **109**, 2183-2201.e9 (2021).

60. Li, N., Daie, K., Svoboda, K. & Druckmann, S. Robust neuronal dynamics in premotor cortex during motor

planning. *Nature* **532**, 459–464 (2016).

61. Yang, W., Tipparaju, S. L., Chen, G. & Li, N. Thalamus-driven functional populations in frontal cortex

support decision-making. *Nat Neurosci* **25**, 1339–1352 (2022).

62. Economo, M. N. *et al.* Distinct descending motor cortex pathways and their roles in movement. *Nature* **563**,

79–84 (2018).

63. Svoboda, K. & Li, N. Neural mechanisms of movement planning: motor cortex and beyond. *Current*

Opinion in Neurobiology **49**, 33–41 (2018).

64. Inagaki, H. K., Fontolan, L., Romani, S. & Svoboda, K. Discrete attractor dynamics underlies persistent

activity in the frontal cortex. *Nature* **566**, 212–217 (2019).

65. Shenoy, K. V., Sahani, M. & Churchland, M. M. Cortical Control of Arm Movements: A Dynamical

Systems Perspective. *Annual Review of Neuroscience* **36**, 337–359 (2013).

66. Orlandi, J. G., Abdolrahmani, M., Aoki, R., Lyamzin, D. R. & Benucci, A. Distributed context-dependent

choice information in mouse posterior cortex. *Nat Commun* **14**, 192 (2023).

67. Inagaki, H. K., Inagaki, M., Romani, S. & Svoboda, K. Low-Dimensional and Monotonic Preparatory

Activity in Mouse Anterior Lateral Motor Cortex. *J. Neurosci.* **38**, 4163–4185 (2018).

68.   Finkelstein, A. *et al.* Attractor dynamics gate cortical information flow during decision-making. *Nat*

1024   *Neurosci* **24**, 843–850 (2021).

1025   69.   Narayanan, N. S. Ramping activity is a cortical mechanism of temporal control of action. *Current Opinion*

1026   *in Behavioral Sciences* **8**, 226–230 (2016).

1027   70.   Cisek, P., Puskas, G. A. & El-Murr, S. Decisions in Changing Conditions: The Urgency-Gating Model. *J.*

1028   *Neurosci.* **29**, 11560–11571 (2009).

1029   71.   Tanaka, M. Cognitive Signals in the Primate Motor Thalamus Predict Saccade Timing. *J. Neurosci.* **27**,

1030   12109–12118 (2007).

1031   72.   Maimon, G. & Assad, J. A. A cognitive signal for the proactive timing of action in macaque LIP. *Nat*

1032   *Neurosci* **9**, 948–955 (2006).

1033   73.   Yu, B. M. *et al.* Gaussian-Process Factor Analysis for Low-Dimensional Single-Trial Analysis of Neural

1034   Population Activity. *Journal of Neurophysiology* **102**, 614–635 (2009).

1035   74.   Pandarinath, C. *et al.* Inferring single-trial neural population dynamics using sequential auto-encoders. *Nat*

1036   *Methods* **15**, 805–815 (2018).

1037   75.   Kawai, R. *et al.* Motor cortex is required for learning but not for executing a motor skill. *Neuron* **86**, 800–

1038   812 (2015).

1039   76.   Safaie, M. *et al.* Turning the body into a clock: Accurate timing is facilitated by simple stereotyped

1040   interactions with the environment. *Proceedings of the National Academy of Sciences* **117**, 13084–13093

1041   (2020).

1042   77.   Moore, J. D., Kleinfeld, D. & Wang, F. How the brainstem controls orofacial behaviors comprised of

1043   rhythmic actions. *Trends in Neurosciences* **37**, 370–380 (2014).

1044   78.   Arber, S. & Costa, R. M. Networking brainstem and basal ganglia circuits for movement. *Nat Rev Neurosci*

1045   **23**, 342–360 (2022).

1046   79.   Merel, J., Botvinick, M. & Wayne, G. Hierarchical motor control in mammals and machines. *Nat Commun*

1047   **10**, 5489 (2019).

1048   80.   Hausmann, S. B., Vargas, A. M., Mathis, A. & Mathis, M. W. Measuring and modeling the motor system

1049   with machine learning. *Current Opinion in Neurobiology* **70**, 11–23 (2021).

1050   81.   Pereira, T. D., Shaevitz, J. W. & Murthy, M. Quantifying behavior to understand the brain. *Nat Neurosci* **23**,

1051   1537–1549 (2020).

1052   82.   Anderson, D. J. & Perona, P. Toward a Science of Computational Ethology. *Neuron* **84**, 18–31 (2014).

1053   83.   Miller, C. T. *et al.* Natural behavior is the language of the brain. *Current Biology* **32**, R482–R493 (2022).

1054   84.   Krakauer, J. W., Ghazanfar, A. A., Gomez-Marin, A., MacIver, M. A. & Poeppel, D. Neuroscience Needs

1055   Behavior: Correcting a Reductionist Bias. *Neuron* **93**, 480–490 (2017).

1056   85.   Datta, S. R., Anderson, D. J., Branson, K., Perona, P. & Leifer, A. Computational Neuroethology: A Call to

1057   Action. *Neuron* **104**, 11–24 (2019).

1058   86.   Parker, P. R. L., Brown, M. A., Smear, M. C. & Niell, C. M. Movement-Related Signals in Sensory Areas:

1059   Roles in Natural Behavior. *Trends in Neurosciences* **43**, 581–595 (2020).

1060   87.   Gomez-Marin, A. & Ghazanfar, A. A. The Life of Behavior. *Neuron* **104**, 25–36 (2019).

1061   88.   Dennis, E. J. *et al.* Systems Neuroscience of Natural Behaviors in Rodents. *J. Neurosci.* **41**, 911–919 (2021).

1062   89.   Testard, C. *et al.* Neural signatures of natural behavior in socializing macaques. 2023.07.05.547833 Preprint

1063   at https://doi.org/10.1101/2023.07.05.547833 (2023).

1064   90.   Marshall, J. D. *et al.* Continuous Whole-Body 3D Kinematic Recordings across the Rodent Behavioral

1065   Repertoire. *Neuron* **109**, 420-437.e8 (2021).

1066   91.   Christensen, A. J., Ott, T. & Kepecs, A. Cognition and the single neuron: How cell types construct the

1067   dynamic computations of frontal cortex. *Current Opinion in Neurobiology* **77**, 102630 (2022).

1068   92.   Wang, Z. A. *et al.* Not everything, not everywhere, not all at once: a study of brain-wide encoding of

1069   movement. 2023.06.08.544257 Preprint at https://doi.org/10.1101/2023.06.08.544257 (2023).

1070   93.   Gallego, J. A., Perich, M. G., Miller, L. E. & Solla, S. A. Neural Manifolds for the Control of Movement.

1071   *Neuron* **94**, 978–984 (2017).

1072   94.   Fortunato, C. *et al.* Nonlinear manifolds underlie neural population activity during behaviour.

1073   2023.07.18.549575 Preprint at https://doi.org/10.1101/2023.07.18.549575 (2023).

95. Dahmen, D. *et al.* Strong and localized recurrence controls dimensionality of neural activity across brain

areas. 2020.11.02.365072 Preprint at https://doi.org/10.1101/2020.11.02.365072 (2023).

96. Morales-Gregorio, A. *et al.* Neural manifolds in V1 change with top-down signals from V4 targeting the

foveal region. 2023.06.14.544966 Preprint at https://doi.org/10.1101/2023.06.14.544966 (2023).

97. Sohn, H., Narain, D., Meirhaeghe, N. & Jazayeri, M. Bayesian Computation through Cortical Latent

Dynamics. *Neuron* **103**, 934-947.e5 (2019).

98. Joshi, S. & Gold, J. I. Pupil Size as a Window on Neural Substrates of Cognition. *Trends in Cognitive*

Sciences **24**, 466–480 (2020).

99. Lee, C. R. & Margolis, D. J. Pupil Dynamics Reflect Behavioral Choice and Learning in a Go/NoGo Tactile

Decision-Making Task in Mice. *Frontiers in Behavioral Neuroscience* **10**, (2016).

100. Bradley, M. M., Miccoli, L., Escrig, M. A. & Lang, P. J. The pupil as a measure of emotional arousal and

autonomic activation. *Psychophysiology* **45**, 602–607 (2008).

101. Yin, C. *et al.* Engaged decision-makers align spontaneous movements to stereotyped task demands.

2023.06.26.546404 Preprint at https://doi.org/10.1101/2023.06.26.546404 (2023).

102. Kim, T. H. *et al.* Long-Term Optical Access to an Estimated One Million Neurons in the Live Mouse

Cortex. *Cell Reports* **17**, 3385–3394 (2016).

103. Jun, J. J. *et al.* Fully integrated silicon probes for high-density recording of neural activity. *Nature* **551**, 232–

236 (2017).

104. Jun, J. J. *et al.* Real-time spike sorting platform for high-density extracellular probes with ground-truth

validation and drift correction. *bioRxiv* 101030 (2017) doi:10.1101/101030.

105. Boumal, N., Mishra, B., Absil, P.-A. & Sepulchre, R. Manopt, a Matlab Toolbox for Optimization on

Manifolds. *Journal of Machine Learning Research* **15**, 1455–1459 (2014).

106. Jiang, X., Saggar, H., Ryu, S. I., Shenoy, K. V. & Kao, J. C. Structure in Neural Activity during Observed

and Executed Movements Is Shared at the Neural Population Level, Not in Single Neurons. *Cell Reports* **32**,

108006 (2020).

107. Aarts, E., Verhage, M., Veenvliet, J. V., Dolan, C. V. & van der Sluis, S. A solution to dependency: using

multilevel analysis to accommodate nested data. *Nat Neurosci* **17**, 491–496 (2014).

108. de Leeuw, J. & Meijer, E. *Handbook of Multilevel Analysis*. vols 401–433 (Springer, New York, 2008).

1102

1103

1104   **Extended Data**

1105

Extended Data Fig. 1 – Effects of photoinactivation of ALM and tjM1 on uninstructed movements and tongue kinematics. a. Percentage of time spent moving (see **Methods**) in a session determined by video recordings using the side view, bottom view, and both views. Points indicate each of the 12 randomly selected sessions used for this analysis. **b.** Delay epoch photoinactivation results are not exclusive to full MC photoinactivation (*top*, copied from **Fig. 1g-i,** n=2 mice, 14 sessions). Similar results were obtained when photoinactivating ALM (*middle*, n=2 mice, 15 sessions) and tjM1 (*bottom*, n=2 mice, 9 sessions) photoinactivation. **c.** Tongue length during control and go cue photoinactivation trials for delayed-response (*left*) and water-cued (*right*) contexts. Blue traces indicate right lickport contacts, red traces indicate left contacts, and black traces indicate no contact. Vertical dashed line indicates go cue or water drop onset. Blue shaded region indicates photoinactivation period. **d.** Fraction of time with tongue visible during photoinactivation period for DR trials (*left*) and WC trials (*right*). Asterisks denote significant differences ($p < 0.01$) between control and photoinactivation trials (*p*-values from left to right: 1.017e-08, 2.030e-06, 0.481, 0.002; paired t-test).

Extended Data Fig. 2 - Session-by-session variability of the relationship between kinematic features and putative cognitive dynamics. a. Regression weights for each group of kinematic predictors for predicting CD$_{choice}$ projections, as a fraction of all predictor coefficients (see **Methods**). Sessions are sorted in descending order by motion energy fraction. Outlined bars indicate example sessions shown in (b-c). **b.** Example session of uninstructed movements where motion energy made up the largest fraction of regression weights for predicting CD$_{choice}$ projections. *Top*, overlayed jaw/nose speed or motion energy for a subset trials. *Bottom*, two example trials of kinematic feature trajectories. **c.** Same as (b) but for an example session where jaw and nose features made up a larger fraction of regression weights. **d.** Same as (a) but for predicting CD$_{context}$ projections. **e, f.** Same as (b-c) but for two example sessions with different regression weight fractions for predicting CD$_{context}$ projections.

Extended Data Fig. 3 – Control analyses for subspace decomposition. a-b. Null and potent subspaces estimated as in **Fig. 5g-l** using DR and WC trials. **a.** Variance explained (R^2) of motion energy by the sum squared magnitude of activity in the null and potent subspaces on single trials. Each point is the mean across trials for a session. **b.** *Left*, motion energy on single trials for an example session. *Middle*, sum squared magnitude of activity in the potent subspace on single trials. *Right*, sum squared magnitude of activity in the null subspace. Trials sorted by average delay epoch motion energy. **c.** Selectivity (left vs. right) of the neural population during WC trials. Mean and 95% CI across sessions shown. **d-e.** Same as (a-b) but the null and potent subspaces have been estimated using WC trials only. This allows for the estimation of a "null" model of movements in the absence of goal-directed behavior. **f.** Normalized magnitude of activity in the null space (*left*) or potent space (*right*) when estimated using DR and WC trials versus when estimated using WC trials only. Circles are average activity per trial for an example session. **g-h.** Same as (a-b), but the null and potent subspaces have been estimated using data restricted to the response epoch of DR and WC trials. Only the delay epoch shown for (b,e,h) to highlight the methods ability to capture delay epoch uninstructed movements in the potent subspace under various subspace estimation conditions. **i.** Magnitude of activity in the null space (*left*) or potent space (*right*) when estimated using DR and WC trials versus when estimated using data from the response epoch of DR and WC trials. Circles are average activity per trial for an example session.

1145
Extended Data Fig. 4 – Random subspaces show single units contribute both to null and potent subspaces. In order to ask whether
the subspace alignment distributions seen in our data (**Fig. 5g,n**) were different from those expected by chance, we compared these
distributions against alignment distributions obtained when neural activity is decomposed into random null and potent subspaces. Each
element of the random subspaces was drawn from a normal distribution with zero mean and unit variance (see **Methods**). Null distributions
of alignment indices are shown for **a.** trial-averaged data and **b.** single-trial data. Both alignment distributions differed from those obtained
from random subspaces (Trial-averaged data, $p=6\times10^{-8}$; Single-trial data, $p=6\times10^{-9}$; two-sample Kolmogorov-Smirnov test).

1152

1153

1154

1155

EDFigure 5 – Alternative procedure for estimating motor planning CDs from subspaces. a. Projections of null and potent subspace activity along CD_{ramp} for each of the three control conditions. Null and potent subspaces were identified using both DR and WC trials (*left*), WC trials only (*middle*), and the response epoch of DR and WC trials (*right*). Mean and 95% CI of bootstrap distribution shown. **b.** Selectivity in null (*left*) and potent (*right*) components of CDchoice projections when CDchoice is determined from subspace activity, rather than from the full neural population. Mean and 95% CI of bootstrap distribution shown. **c.** Projections along null (*left*) and potent (*right*) components of CDramp when CDramp is determined from subspace activity, rather than from the full neural population. Mean and 95% CI of bootstrap distribution shown. **d.** Projections of potent subspace activity along CDchoice, CDaction, and CDramp when determining subspaces from WC trials only. CDs were determined using the full neural population. Mean and 95% CI of bootstrap distribution shown.

EDFigure 6 – **Relationship between tongue angle and neural activity in the null and potent subspaces. a.** Projections along potent (*top*) and null (*bottom*) components of CD_{action}. Correct trials shown in solid lines and error trials shown in dashed lines. **b.** Tongue angle for an example session for correct and error trials. **c.** Tongue angle on correct and error right and left trials. Tongue angle was linearly time warped to allow for averaging over trials and sessions. Mean and s.e.m. across sessions shown. **d.** Tongue angle (*left*) and predictions from the full population neural activity (*middle left*), null subspace activity (*middle right*), and potent subspace activity (*right*) for an example session. **e.** Variance explained (R^2) between tongue angle and predictions from null (*green*) and potent (pink) subspaces. Asterisks denote significant differences between predictions from null and potent subspaces ($p=2 \times 10^{-8}$, paired t-test).

Supplementary Movie 1 – Uninstructed movements vary in their identity and timing. Example trials in which uninstructed movements vary in their identity (across rows) and timing (across columns). Traces represent the y-position of the feature within the video frame. All example trials taken from the same mouse and session.