Functional Diversity of Dopamine Axons in Prefrontal Cortex During Classical Conditioning

Kenta Abe[1#], Yuki Kambe[2#], Kei Majima[3,4#], Zijing Hu[5,6#], Makoto Ohtake[1], Ali Momennezhad[2], Hideki Izumi[7], Takuma Tanaka[7], Ashley Matunis[1,8], Emma Stacy[1,8], Takahide Itokazu[9], Takashi R. Sato[1,11*], Tatsuo K. Sato[2,4,5,6,10*]

1. Department of Neuroscience, Medical University of South Carolina, Charleston, South Carolina 29425, United States
2. Department of Pharmacology, Kagoshima University, Kagoshima, 890-8544, Japan
3. Institute for Quantum Life Science, National Institutes for Quantum Science and Technology, Chiba 263-8555, Japan
4. Japan Science and Technology PRESTO, Saitama 332-0012, Japan
5. Department of Physiology, Monash University, Clayton, 3800, Australia
6. Neuroscience Program, Biomedicine Discovery Institute, Monash University, Clayton, 3800, Australia
7. Faculty of Data Science, Shiga University, Shiga, 522-8522, Japan
8. Department of Biology, College of Charleston, Charleston, South Carolina 29424, United States
9. Department of Neuro-Medical Science, Osaka University, Osaka, 565-0871, Japan
10. Japan Science and Technology FOREST, Saitama 332-0012, Japan
11. Lead contact

# These authors contributed equally to this work.
* For correspondence:
satot@musc.edu (TRS) and tatsuo.sato@m.kufm.kagoshima-u.ac.jp (TKS)

Number of pages: 57
Number of figures, tables, multimedia, and 3D models: 4, 0, 0, 0
Number of words for abstract, introduction, and discussion: 206, 742, 1458

Conflicts of Interest: The authors declare no conflicts of interest.

Acknowledgements: We thank J. Zhang for technical help. This work was supported by grants from JSPS KAKENHI (20K16465) and JST PRESTO to K.M. (JPMJPR2128), JSPS KAKENHI to T.I. (JP21K07459), BBRF Young Investigator Grant (29268), National Institute on Drug Abuse (COCA Pilot grant, P50DA046373), National Institute of Aging (R03 AG070517), National Institute of Neurological Disorders and Stroke (R21 NS125571, R01 NS131549), and NIH COBRE in Neurodevelopment and its Disorders (P20 GM148302) to T.R.S., and JST PRESTO (JPMJPR1883), JST FOREST (JPMJFR2245), NHMRC Ideas Grant (APP1184899), and KAKENHI (20K23378) to T.K.S.

Abstract

Midbrain dopamine neurons impact neural processing in the prefrontal cortex (PFC) through mesocortical projections. However, the signals conveyed by dopamine projections to the PFC remain unclear, particularly at the single-axon level. Here, we investigated dopaminergic axonal activity in the medial PFC (mPFC) during reward and aversive processing. By optimizing microprism-mediated two-photon calcium imaging of dopamine axon terminals, we found diverse activity in dopamine axons responsive to both reward and aversive stimuli. Some axons exhibited a preference for reward, while others favored aversive stimuli, and there was a strong bias for the latter at the population level. Long-term longitudinal imaging revealed that the preference was maintained in reward- and aversive-preferring axons throughout classical conditioning in which rewarding and aversive stimuli were paired with preceding auditory cues. However, as mice learned to discriminate reward or aversive cues, a cue activity preference gradually developed only in aversive-preferring axons. We inferred the trial-by-trial cue discrimination based on machine learning using anticipatory licking or facial expressions, and found that successful discrimination was accompanied by sharper selectivity for the aversive cue in aversive-preferring axons. Our findings indicate that a group of mesocortical dopamine axons encodes aversive-related signals, which are modulated by both classical conditioning across days and trial-by-trial discrimination within a day.

Impact statement

Two-photon calcium imaging revealed that many mesocortical dopamine axons show enhanced selectivity for aversive cue processing during classical conditioning.

Introduction

The prefrontal cortex (PFC) contributes to a variety of higher cognitive functions, achieving the flexible control of behaviors that enables animals to adapt to a changing environment (Miller and Cohen, 2001; Fuster, 2015). The PFC is involved, for instance, in stimulus selection, working memory, rule switching, and decision making (Miller and Wallis, 2009). PFC processing and circuits are highly sensitive to neuromodulators, including dopamine (Seamans and Yang, 2004; Arnsten et al., 2012). Indeed, studies using pharmacological or optogenetic manipulation of dopamine signaling have suggested roles of dopamine in gating sensory signals (Popescu et al., 2016; Vander Weele et al., 2018), maintaining working memory (Sawaguchi and Goldman-Rakic, 1994), and relaying decisions to motor structures (Ott et al., 2014). Consistently, dysregulation of dopamine signaling in the PFC has been suggested to underlie a wide array of neuropsychiatric disorders, including schizophrenia, depression, attention-deficit/hyperactivity disorder, and post-traumatic stress disorder (Okubo et al., 1997; Lindstrom et al., 1999; Granon et al., 2000; Arnsten and Dudley, 2005; Howes and Kapur, 2009; Hoexter et al., 2012; Grace, 2016).

The PFC receives dopaminergic inputs from a subset of dopamine neurons in the midbrain, but the information encoded by this subset *in vivo* remains unclear. Decades of investigations have revealed that midbrain dopamine neurons in the ventral tegmental area (VTA) generally encode reward prediction errors (Schultz et al., 1997): the neurons increase their firing to unexpected reward delivery and shift their response to cues that precede reward delivery after instrumental learning or classical conditioning (Rescorla and Wagner, 1972; Sutton and Barto, 1981). However, several studies have reported that a subpopulation of dopamine neurons show phasic responses to aversive stimuli as a part of salience signaling (Chiodo et al., 1980; Mantz et al., 1989; Guarraci and Kapp, 1999; Matsumoto and Hikosaka, 2009), implying that midbrain dopamine neurons may not be functionally homogeneous. Indeed, depending on the projection target, dopamine neurons can have distinct molecular, anatomical, and electrophysiological features (Lammel et al., 2008;

Poulin et al., 2018). Dopamine neurons that project to the PFC might locate primarily to the medial

posterior VTA (Lammel et al., 2008) and show different genetic profiles from other dopamine

neurons (Poulin et al., 2018). In addition, optogenetic stimulation of the PFC-projecting DA

neurons does not reinforce specific actions (Popescu et al., 2016; Ellwood et al., 2017; Vander

Weele et al., 2018). Moreover, these neurons might respond not only to rewarding stimuli but also

to aversive stimuli. Microdialysis, amperometry, and voltammetry measurements in the PFC have

demonstrated an increase of dopamine in response to appetitive stimuli (Hernandez and Hoebel,

1990; Ahn and Phillips, 1999; St Onge et al., 2012), aversive stimuli (Thierry et al., 1976;

Abercrombie et al., 1989; Finlay et al., 1995; Vander Weele et al., 2018), or both (Bassareo et al.,

2002). Similarly, measurements of the bulk calcium activity of mesocortical dopaminergic fibers

have shown responses to appetitive (Ellwood et al., 2017) and aversive (Kim et al., 2016) stimuli.

This apparent discrepancy is difficult to reconcile because none of these approaches could

investigate the activity of individual dopamine neurons. Moreover, most previous studies

evaluated the effects of either a rewarding or an aversive stimulus, rather than both. Consequently,

it remains unknown whether the same or different mesocortical dopamine neurons respond to

behaviorally opposing stimuli. It is also not known how these dopamine neurons change their

response during classical conditioning, where rewarding or aversive stimuli are paired with

conditioned cues.

To address these knowledge gaps, we developed an approach for imaging individual dopamine

axons based on *in vivo* two-photon imaging with a microprism (Low et al., 2014). We optimized

the microprism design and imaged dopamine axon terminals expressing genetically encoded

calcium sensors in the mouse medial PFC (mPFC). We then head-fixed the mice to give rewards

or aversive stimuli (water drops or electrical shocks) and trained the mice to associate the stimuli

with preceding auditory cues (classical conditioning). During classical conditioning, we tracked

the activity of dopamine axons over a period of days. We found that the dopamine axons showed

diverse preferences for unconditioned (rewarding or aversive) stimuli. Through the classical

conditioning, activity preferences for conditioned auditory cues were enhanced only for aversive-

preferring axons. Moreover, in aversive-preferring axons, a machine learning-based analysis

revealed that cue activity became more selective when the behavior of animals was judged as

correct. We conclude that mesocortical dopamine axon activity is involved in aversive-related

processing that is modulated by both classical conditioning across days and trial-by-trial

judgements of conditioned cues within a day.

Results

Two-photon imaging shows dopaminergic axons in the mPFC of
awake mice

To investigate the signal sent by dopamine neurons to the mPFC in mice, we developed an

approach based on two-photon imaging using a microprism (Andermann et al., 2013; Low et al.,

2014). We first expressed axon-jGCaMP8m, an axon-targeted (Broussard et al., 2018) genetically

encoded calcium sensor (Zhang et al., 2023), in dopamine neurons in the VTA. We injected Cre-

dependent AAV into the midbrain regions of transgenic mice (DAT-Cre), which express Cre-

recombinase in dopamine neurons (Kim et al., 2016) (Figure 1A, see Methods). After 2–3 weeks,

using sectioned slices, we confirmed that GCaMP expression in cell bodies in the VTA (and

substantia nigra pars compacta [SNc]) (Figure 1C) coincides with the expression of tyrosine

hydroxylase, an endogenous marker for dopamine neurons (Figure 1D). Dopamine neurons in

the VTA are known to project sparsely to the mPFC, including the superficial layers (Vander

Weele et al., 2018) (Figure 1–figure supplement 1), but the mPFC itself is located deep in the

medial bank (Figure 1B), rendering two-photon imaging of GCaMP (which is typically excited at

920–980 nm) infeasible. Therefore, we inserted a microprism into the longitudinal fissure between

the two medial banks (two hemispheres) to optically access the mPFC (Figure 1E–F). The right-

angle microprism bends the optical axis within the brain, providing optical access to the fissure

wall and the mPFC surface (Low et al., 2014). We optimized the microprism assembly (Figure 1–

figure supplement 2C) in order to reach up to 2 mm in depth from the dorsal surface (Figure 1F).

The assembly incorporated double-layer glass at the top (Komiyama et al., 2010), stabilizing the

brain from both the medial and dorsal sides, which significantly reduced the movement of the

brain (Figure 1–figure supplement 2E-F). Through the microprism, we could visualize GCaMP-

expressing axons in the superficial layers of the mPFC in live animals (at a depth of 30–100 μm,

Figure 1G, Figure 1–figure supplement 3). Sparse axons in the superficial layers are

advantageous for two-photon imaging, achieving low background noise. In contrast, axons in the

deep layers, which are known to be denser in sectioned slices (Figure 1–figure supplement 3),

could not be visualized under our experiments. The GCaMP signal can indicate the calcium influx

into axons and terminals, which is triggered by axonal action potentials (Petreanu et al., 2012;

Howe and Dombeck, 2016; Lutas et al., 2019), thereby providing a measure of the activity of

dopamine neurons that send projections to the mPFC. In contrast, when we inserted a gradient

refractive index (GRIN) lens into the mPFC (Kamigaki and Dan, 2017), we could not reliably

visualize GCaMP-expressing dopamine axons, unlike the case for dopamine axons in the basal

amygdala (Lutas et al., 2019). This difference might indicate that the dopamine axons in the mPFC

have weaker signals requiring a lens with a larger numerical aperture (GRIN lens: NA 0.5 vs.

Nikon objective lens: NA 0.8) or that these axons are less resilient to mechanical damage in close

vicinity.

Dopaminergic axons in the mPFC have diverse responses to rewarding and aversive stimuli

Using our imaging approach, we first investigated whether individual dopamine axons respond to

unexpected rewards (Schultz et al., 1997) and unexpected aversive stimuli (Vander Weele et al.,

2018). As a reward, we delivered drops of water through a spout with random timing to water-

deprived mice (Figure 2A). In response to the reward delivery, the mice licked the water spout,

and we filmed this behavior to quantify the tongue position (Figure 2B and Figure 2–figure

supplement 1). Upon the delivery of a water drop, the mice started licking (licking latency: 0.538

± 0.065 s, n = 8 animals) (Otis et al., 2017). Two-photon calcium imaging revealed that the water

reward evoked brief calcium transients in many dopamine axons (40.1% of axons in 8 animals,

example in Figure 2I). The brief calcium response to the reward is consistent with increased

phasic firing in dopamine neurons at the time of unexpected reward, as previously reported in

many studies in primates (Schultz et al., 1997) and rodents (Engelhard et al., 2019; Amo et al.,

2022).

In contrast to conventional midbrain dopamine neurons, mPFC dopamine axons are proposed to

play a key role in aversive processing (Vander Weele et al., 2019). To investigate the calcium

response to an unexpected aversive stimulus, we delivered mild electrical shocks to the tail of the

mice (Kim et al., 2016; Patriarchi et al., 2018; Lutas et al., 2019) (Figure 2A) that were randomly

interleaved with reward delivery (one shock for every seven rewards on average). The mild shock

evoked calcium transients in many dopamine projections (Figure 2F, G, H), together with

locomotion (Figure 2B). These transients could simply reflect locomotion initiation, similar to

dopamine axons in the dorsal striatum (Howe and Dombeck, 2016). To explore this possibility,

we investigated whether locomotion without aversive stimuli is accompanied by increased calcium

activity. We found no significant calcium increase at the time point of spontaneous locomotion

initiation (Figure 2– figure supplement 2). Therefore, unlike the axons projecting from the SNc to

the dorsal striatum (Howe and Dombeck, 2016; Ma et al., 2022), mPFC dopamine axons do not

encode the initiation of movement; rather, these axons respond to the aversive stimulus.

Some previous studies have demonstrated that the overall dopamine release at the mPFC or the

summed activity of mPFC dopamine axons exhibits a strong response to aversive stimuli (e.g.,

tail shock), but little to rewards (Kim et al., 2016). We evaluated the preference of individual axons

for rewarding and aversive signals at a single-axon resolution by computing the polar angle for

individual axons on a Cartesian representation of reward and shock activity (Figure 2J, K, Figure

2–figure supplement 3). In the polar representation, an angle of 0° indicates a strong preference

for reward information, whereas 90° indicates a preference for aversive information. The polar

angle distribution revealed that a significant number of axons preferred aversive stimuli, although

some preferred reward. As a result, probability density, estimated by kernel smoothing, showed

a bimodal distribution (Figure 2K, solid line) with a trough at around 45–50°. In addition, axons

showing significant responses were categorized into two clusters based on k-means clustering

(Figure 2J), the separation of which coincided roughly with 45–50° (Figure 2K). Thereafter, we

refer to these clusters as aversive- or reward-preferring axons (colored magenta or cyan,

respectively). These clusters do not respond exclusively to one hedonic valence (rewarding or

aversive stimuli), as evident from the broad angle distributions. We could not find any anatomical

patterns for aversive- or reward-preferring axons. These axons were present in either half of the

prism view (i.e., anterior or posterior; ventral or dorsal), implying no obvious functional projection

patterns within the mPFC. We note that the strength of preference could be quantitatively changed.

Indeed, we found that the reward response to 10 µL nearly reached saturation, but the aversive

response could be further increased at a stronger current (Figure 2–figure supplement 4).

Therefore, the exclusive preference for aversive stimuli observed in some studies might possibly

be explained by a smaller reward volume and/or stronger aversive stimulus. It may also be

possible that the relative frequency of the aversive stimulus could influence the aversive

preference. Moreover, measured signals in these studies may arise from deep layers and be

different from the superficial axons that we image. Altogether, our two-photon imaging revealed,

for the first time, that individual axons in the superficial layers show diverse preferences for

rewarding and aversive stimuli.

Aversive cue processing is enhanced in aversive-preferring axons during classical conditioning.

How do the reward and aversive activities of individual axons change while animals are learning that the reward and aversive events are preceded and predicted by sensory cues? This paradigm, known as classical conditioning, is a key framework for capturing learning-related changes in midbrain dopamine neurons (Sutton and Barto, 1981; Schultz et al., 1997) and mPFC neurons (Takehara-Nishiuchi and McNaughton, 2008; Otis et al., 2017). We presented mice with a 2-s pure tone as a conditioned stimulus (CS_{reward} and $CS_{aversive}$: 9 and 13 kHz, or 13 and 9 kHz), and then, after a 1-s delay, we presented either a rewarding or an aversive unconditioned stimulus (Figure 3A). Previous work has shown that mice are able to discriminate between two tones that differ by more than 7% (de Hoz and Nelken, 2014). Indeed the mice learned the contingency between the conditioned stimulus (tone) and the outcome (reward or electrical shock), which was reflected in changes in their behavior throughout this conditioning process (Figure 3B, C). To quantify such behavioral changes during learning, we separated the learning into three phases in addition to the first day (Figure 3D–G, for six phases, Figure 3–figure supplement 1). On the first day, the animals licked the water spout only after the reward was delivered (first day; Figure 3B). However, during the middle and late phases, animals gradually showed licking behavior even before the reward delivery, representing an anticipation of reward (Figure 3D). We observed this anticipatory licking more frequently after CS_{reward} than $CS_{aversive}$ (Figure 3D vs. 3E, $p = 0.031$ for the middle phase, $p = 0.031$ for the late phase, Wilcoxon signed-rank test, $n = 6$ animals), indicating that the animals behaviorally learned to discriminate the two conditioned stimulus tones. Similarly, running before the delivery of the unconditioned stimulus was more frequent after $CS_{aversive}$ than CS_{reward} at the late phase (Figure 3F vs. 3G, $p = 0.031$, Wilcoxon signed-rank test, $n = 6$ animals), again indicating that the two conditioned auditory cues were behaviorally discriminated. Therefore, as in previous studies, anticipatory licking (Otis et al., 2017) and

anticipatory running (Lutas et al., 2019) can capture whether animals behaviorally discriminate

conditioned cues in classical conditioning.

Through the classical conditioning paradigm, our long-term two-photon imaging revealed that

aversive-preferring dopamine axons maintained their preference for the unconditioned response

but enhanced their selectivity for the aversive cue activity (Figure 3H). We evaluated the activity

change at the time of the unconditioned stimuli (US) throughout the learning process for aversive-

and reward-preferring axons (Figure 3J, magenta and cyan, respectively). On the first day,

aversive-preferring dopamine axons showed stronger activity for the aversive stimuli (Figure 3J,

top, Figure 3–figure supplement 2A), similar to the response without classical conditioning (Figure

2J). Across learning, the activity for the rewarding and aversive unconditioned stimuli gradually

decreased (Figure 3K, magenta, $p = 0.002$ for rewarding stimuli, $p = 0.007$ for aversive stimuli, n

= 47; Wilcoxon signed-rank test, comparison between the first day and the last phase),

maintaining similar preferences for rewarding and aversive stimuli (Figure 3L, magenta, $p = 0.24$,

$n = 47$; circular statistics, comparison between the first day and the last phase). Similarly, reward-

preferring axons maintained their preferences over the course of classical conditioning (Figure 3L,

cyan, $p = 0.77$, $n = 12$; circular statistics, Figure 3–figure supplement 2B).

Next, we quantified the activity change at the time of the conditioned auditory cues (CS_{reward} and

$CS_{aversive}$, Figure 3I) throughout the learning process. On the first day, aversive-preferring axons

already showed a transient response to conditioned cues, implying that the conditioned stimulus

response was not acquired through learning (first day in Figure 3H, M, and Figure 3–figure

supplement 2A). In addition, the conditioned stimulus response showed no particular

reward/aversive preference (Figure 3N, first day, for aversive-preferring axons, $p = 0.66$, $n = 47$),

indicating that aversive-preferring axons did not distinguish the two conditioned cues. However,

at the later phases of learning, the conditioned stimulus response was enhanced for $CS_{aversive}$ in

aversive-preferring axons (Figure 3M, magenta, $p < 0.0001$, $n = 47$; Wilcoxon signed-rank test,

comparison between the first day and the last phase) and slightly attenuated for CS_{reward} ($p <$

0.001), resulting in a stronger preference for aversive processing (late phase in Figure 3N,

magenta, p < 0.001). In contrast, for reward-preferring axons, the conditioned stimulus response

increased both for $CS_{aversive}$ (Figure 3M, dotted cyan line, non-significantly, p = 0.09, n = 12,

Wilcoxon signed-rank test) and for CS_{reward} (solid cyan line, significantly, p < 0.007), resulting in

an unchanged preference (Figure 3N, cyan, p = 0.77). Consistently, these changes in CS and US

responses across the training phases were evident in the population-averaged calcium response

(Figure 3–figure supplement 2).

We also tested whether the dopamine axons showed suppressed activity when the predicted

reward was omitted, one of the major features of reward prediction error coding (Schultz et al.,

1997; Engelhard et al., 2019; Amo et al., 2022). Such activity suppression has been detected with

GCaMP6m at cell bodies of dopamine neurons (Engelhard et al., 2019) as reduced signal at 0–4

s after the delivery of reward. Therefore, we included one condition for an unexpected reward

omission on the last day of the late phase of the classical conditioning (Figure 3–figure

supplement 3). We found that upon the reward omission, the reward-preferring dopamine axons

did not show activity suppression, indicating that the mPFC dopamine axons do not respond to

reward omission.

Taken together, our two-photon imaging revealed that a minority of mPFC dopamine axons prefer

reward activity (reward-preferring axons), and that these axons are not involved in reward

prediction error in a classical learning paradigm. In contrast, the majority of dopamine axons are

strongly involved in aversive-related processing (aversive-preferring axons), and the preference

for the aversive cue is enhanced through classical conditioning.

Dopamine axons show enhanced selectivity of cue activity in trials with correct discrimination

In the classical conditioning paradigm, an enhanced preference of aversive-preferring dopamine

axons for aversive cues (Figure 3N) was accompanied by improved behavioral discrimination of

the two conditioned cues (Figure 3D-G). Based on this finding, can correct cue discrimination be

accompanied by an enhanced neural preference when animals make trial-by-trial judgements in

discriminating cues even after conditioning?

To investigate trial-by-trial judgements of conditioned cues, we classified the trials into four groups

(Figure 4A) based on correct or incorrect discriminating behavior. First, we focused on the

presence or absence of anticipatory licking, as the licking behavior can discriminate the two

conditioned stimulus tones, particularly at the late phase of learning (Figure 4–figure supplement

1, based on the random forest classifier). The first group exhibited licking after CS_{reward} (correct

reward discrimination), the second group exhibited no licking after CS_{reward} (incorrect reward

discrimination), the third group displayed no licking after $CS_{aversive}$ (correct aversive discrimination),

and the fourth group displayed licking after $CS_{aversive}$ (incorrect aversive discrimination). The

classification is invalid when animals make random guesses (discrimination of 50%), so we

focused on results from the late phase of learning (or the middle phase if there were no errors in

anticipatory licking in the late phase).

Can the axonal response to conditioned cues be impacted by whether animals discriminate the

cues correctly or incorrectly? An incorrect discrimination of the aversive cue is accompanied by

the presence of anticipatory licking, resulting in error trials in our machine learning-based analysis.

Such error trials (Figure 4A, fourth group) occurred in 1.6% of cases, showing a weaker aversive

cue response than correct trials (third group) ($p < 0.0001$, n = 44 magenta axons, Wilcoxon

signed-rank test, Figure 4B, right). In contrast, the absence of anticipatory licking despite the

reward-predictive cue comprises another type of error (second group, 49.0%). In such error trials,

the reward cue response was not significantly different from that in the correct trials (first group)

($p = 0.26$ n = 44 axons, Wilcoxon signed-rank test, Figure 4B, left). Overall, the reward/aversive

preference was stronger in correct discrimination trials than in incorrect trials (left vs. right in

Figure 4C, magenta, 82.2° ± 1.1° vs. 70.0° ± 7.6°, $p = 0.049$, circular statistics).

In addition to anticipatory licking, the discrimination of predictive cues can be inferred by the facial

expressions of mice. Facial expressions of mice can capture emotional states (Dolensek et al.,

2020), and have been used to make binary judgements of the presence or absence of pain with

the application of a deep neural network (Tuttle et al., 2018). In this study, we combined a

pretrained deep neural network (ResNet3D) (Tran et al., 2018) and a machine learning classifier

(random forest classifier) (Breiman, 2001) to make binary judgements of whether the animals

experienced reward or aversive conditions, based on facial expressions during the cue

presentation (Figure 4–figure supplement 2). The percentage of errors in discrimination during

the 2-s cue presentation was 16.0% ± 1.4% (n = 6 animals), comparable to the result for

anticipatory licking during the cue plus delay periods (25.3% ± 6.1%). However, discrimination

based on facial expressions resulted in a higher number of error trials in aversive conditions than

discrimination based on licking (14.1% vs. 1.6%), and a lower number in reward conditions (18.0%

vs. 49.0%). This discrepancy might be explained either by temporal discrepancy between the cue

period (facial expression) and the delay period (most cases of anticipatory licking) or by the fact

that anticipatory licking might represent reward uncertainty rather than reward expectation

(Ogawa et al., 2013).

Correct cue-discrimination based on facial expressions analysis also revealed sharper selectivity

for reward or aversive cues. A trial-by-trial error analysis revealed that the axonal activity to

$CS_{aversive}$ was enhanced in the correct trials in the aversive-preferring axons (Figure 4D, right, p =

0.023, n = 47 axons, Wilcoxon signed-rank test), consistent with the analysis based on

anticipatory licking (Figure 4B, right). In addition, the response to CS_{reward} was significantly weaker

in the correct trials (Figure 4D, left, p = 0.019, n = 47 axons, Wilcoxon signed-rank test). As a

result, the reward/aversive preference was stronger in correct discrimination trials than in incorrect

trials for the aversive-preferring axons (left vs. right in Figure 4E, magenta, 84.7° ± 3.0° vs. 81.7°

± 4.0°, p = 0.019, circular statistics).

In contrast to the aversive-preferring axons, correct discrimination had no effect on the CS activity

of the reward-preferring axons. We found that the response to CS_{reward} and $CS_{aversive}$ was not

significantly different between correct and incorrect judgement trials (cyan points in Figure 4B and

D, anticipatory licking: $CS_{aversive}$ p = 0.052, facial expressions: CS_{rewad} p = 0.08, $CS_{aversive}$ p = 0.20,

n = 12, Wilcoxon signed-rank test) except for the CS_{reward} response based on anticipatory licking

(p = 0.016, n= 12). As a result, selectivity for CS_{reward}/$CS_{aversive}$ was not improved in correct trials

(anticipatory licking: p = 0.15, facial expressions: p = 0.39, cyan points in Figure 4C, E). Therefore,

correct/incorrect discrimination impacts aversive- and reward-preferring axons differentially.

Altogether, when animals exhibited the correct behavioral response (either anticipatory licking or

facial expression), aversive-preferring but not reward-preferring axons showed a higher selectivity

for aversive cue processing (Figure 4C, E).

347 Discussion

Dopamine projections to the mPFC are considered one of the key neuromodulators that enable

flexibility in neural processing of the mPFC. However, due to technical difficulties in recording the

dopamine neurons of specific projections, little is known about the signals conveyed by mPFC

projections, including the basic question of whether individual projections signal reward or

aversive information. In this study, we optimized a two-photon imaging approach based on a

microprism to image the calcium activity of dopaminergic axons in the mPFC. We uncovered

differences in reward/aversive preferences in individual dopamine axons with an overall

preference for aversive stimuli. In addition, we demonstrated that aversive-preferring axons

responded equally to reward- and aversive-predictive conditioned cues in classical conditioning

on the first day; however, this response became strongly biased toward the aversive conditioned

cue through the conditioning. Finally, based on a trial-by-trial analysis of the animals' behavior

following reward- or aversive-predictive cues, we found that aversive-preferring axons exhibited

higher selectivity for cues when the cues were successfully discriminated behaviorally.

Our study revealed functional diversity in mPFC-projecting dopamine axons by addressing a long-

standing question of whether dopamine neurons send reward- or aversion-related signals to the

mPFC (Vander Weele et al., 2019; Verharen et al., 2020). The activity of mPFC-projecting

dopamine neurons can be investigated extracellularly by incorporating antidromic stimulation

(Mantz et al., 1989), but this approach is laborious. Thus, many studies have used more

technically feasible but less direct approaches, particularly for awake animals, such as measuring

dopamine release with microdialysis (Abercrombie et al., 1989; Bassareo et al., 2002), measuring

catecholamine release with fast-scan cyclic voltammetry (recently combined with optogenetic and

pharmacological identification (Vander Weele et al., 2018), and measuring bulk calcium activity

from dopamine axons with fiber photometry (Kim et al., 2016; Ellwood et al., 2017). These studies

have led to somewhat inconsistent conclusions: some studies have reported reward signals

whereas others have reported aversive signals. Reconciling these findings is challenging, as

different studies have used different approaches to assess the effects of either a rewarding or an

aversive stimulus, but not both. Our two-photon imaging approach provided a unique opportunity

to compare rewarding and aversive signals of individual projection neurons (i.e., individual axon

projections). Our comparison revealed diversity in the dopamine axons, and that many dopamine

axons responded to both rewarding and aversive stimuli, with a strong bias for aversive stimuli at

the population level. However, this population bias was not fixed; rather, the bias depended on

both the reward volume and the intensity of the aversive stimulus. In addition, the bias might also

rely on of the frequency of rewarding and aversive events, which our study could not address; we

presented aversive stimuli less frequently to keep mice engaged. All of these parameters may

collectively explain why some studies have reported a strong response to aversive stimuli but little

response to rewards. Our study revealed functional diversity in dopamine axons in the superficial

layers, but did not address whether this diversity could also be found in axons in the deep layers.

It is not clear whether single axons have branches in both the superficial and deep layers, even

in anatomical studies that used single-cell tracing of dopamine neurons (Matsuda et al., 2009;

Aransay et al., 2015). Further investigation into these layers may reveal a richer functional

diversity in dopamine axons in the mPFC.

The firing of mPFC-projecting dopamine neurons cannot be simply explained by value coding

performed by conventional midbrain dopamine neurons; their firing rates increase in response to

opposite hedonic valences, rewarding and aversive stimuli, exhibiting diverse preference. Their

firing might be captured by salience coding, which includes motivational salience signals and

alerting signals (Bromberg-Martin et al., 2010). Similarly, salience coding may explain the

response of aversive-preferring axons to aversive-predictive cues. On the first day of classical

conditioning, when the animals had not yet established a link between conditioned and

unconditioned stimuli, the aversive-preferring axons showed a transient activity increase to two

types of conditioned cues with no bias, implying that activity serves as an alerting signal (Vander

Weele et al., 2018). After days of training, the activity became strongly biased toward the aversive

cue, indicating that the activity might additionally encode a motivational salience signal

(Bromberg-Martin et al., 2010; Lee et al., 2021). In our study, motivational salience might appear

to play a significant role in the processing of the aversive predictive cue. Specifically, in aversive-

preferring axons, the activity associated to the aversive cue shows a distinct pattern in the initial

trial of each session, compared to the activities related to the reward cue and unconditioned cues

(as depicted by magenta points in Figure 3–figure supplement 4). The response of the aversive-

preferring axons may be useful for the recipient mPFC to allocate its resources to the most salient

outcomes and their predictors, a proposed role of the mPFC (Desimone and Duncan, 1995; Miller

and Cohen, 2001; Ridderinkhof et al., 2004; Isoda and Hikosaka, 2007; Nee et al., 2011;

Bissonette et al., 2013; Sharpe et al., 2019). The response of reward-preferring axons also does

not follow value-coding, considering the absence of reward omission suppression. It is an open

question whether saliency coding plays more important roles in aversive processing than in

reward processing. To clarify the detailed nature of the saliency coding, together with the

functional diversity of mPFC dopamine axons, further study is necessary. Such a study should

include different types of unconditioned stimuli, systematically vary the physical features of

conditioned and unconditioned stimuli, and separate the motivational salience and alerting signals

with different task designs.

Consistent with salience coding without hedonic valences, phasic optogenetic stimulation of

dopamine axons in the mPFC does not reinforce or suppress any behavioral actions (Popescu et

al., 2016; Ellwood et al., 2017; Vander Weele et al., 2018) (but see (Gunaydin et al., 2014)).

Instead, optogenetic stimulation can increase the signal-to-noise ratio of aversive processing in

mPFC neurons for competitive situations in which reward and appetitive cues are simultaneously

presented (Vander Weele et al., 2018). Our results are consistent with the recent view that

dopamine at the mPFC gates sensory inputs for aversive processing (Ott and Nieder, 2019;

Vander Weele et al., 2019).

Using aversive classical conditioning, we revealed that aversive learning can induce activity

changes in dopamine axons. The classical conditioning is a form of aversive learning distinct from

instrumental aversive learning [including punishment and active avoidance (Jean-Richard-Dit-

Bressel et al., 2018)]. Although all types of aversive learning are processed in the mPFC and

dopamine systems, each type may be expected to include distinct neural circuits. Further

research is necessary to reveal the detailed processing of aversive learning in the mPFC and

dopamine projections.

Our study provides new insights into the functional diversity of dopamine neurons that constitute

mesocortical pathways. Previous studies employing fiber photometry imaging have identified

functional diversity among dopamine neurons with distinct projection pathways; dopamine axons

in the ventral nucleus accumbens medial shell (de Jong et al., 2019; Yuan et al., 2019), in the tail

of the striatum (Menegas et al., 2017), and in the basal amygdala (Lutas et al., 2019) do not show

activity that matches reward prediction error coding, but instead show increased activity for

aversive stimuli (Verharen et al., 2020). Our two-photon imaging results demonstrate that even

the same projection-defined dopamine neurons can be inhomogeneous, with some preferring

aversive signals and others preferring reward signals. The aversive response found in some

dopamine pathways, including mesocortical dopamine projections, might be linked to glutamate

co-release, as vesicular glutamate transporter 2 (*Vglut2*) genes are expressed in dopamine

neurons projecting to the ventral nucleus accumbens medial shell, the tail of the striatum, and the

mPFC (Poulin et al., 2018), and as AMPA-receptor-mediated excitatory postsynaptic currents

have been confirmed upon the stimulation of dopamine axon terminals in the basal amygdala

(Lutas et al., 2019). These glutamate co-releasing dopamine axons might even be collaterals of

the same dopamine neurons; a single-cell tracing study showed examples of dopamine neurons

with their axon collaterals in the PFC and the basal amygdala or those in the PFC and nucleus

accumbens shell (Aransay et al., 2015). Meanwhile, dopamine axons projecting to the PFC are

not just from *Vglut2*-expressing neurons but also from *Vglut2*-negative neurons (Poulin et al.,

2018). One possible scenario is that *Vglut2* may be expressed in aversive-preferring axons in the

mPFC but not in reward-preferring axons. This scenario might be in line with a recent fibermetry

study on genetic features of dopamine neurons in the SNc, where $Vglut2^+$ dopamine neurons

showed a strong aversive and weak reward response as a population, whereas $Vglut2^-/Calb1^+$

dopamine neurons showed a weak aversive and strong reward response (Azcorra et al., 2023).

As of now, it is not clear how functional diversity within the same mesocortical pathway is linked

to molecular diversity. Clarifying such a link will further advance our understanding of distinct

dopamine subsystems and may shed light on how dopamine subsystems are dysregulated in
prefrontal psychiatric diseases.

Acknowledgements

We thank J. Zhang for technical help. This work was supported by grants from JSPS KAKENHI
(20K16465) and JST PRESTO to K.M. (JPMJPR2128), JSPS KAKENHI to T.I. (JP21K07459),
BBRF Young Investigator Grant (29268), National Institute on Drug Abuse (COCA Pilot grant,
P50DA046373), National Institute of Aging (R03 AG070517), National Institute of Neurological
Disorders and Stroke (R21 NS125571, R01 NS131549), and NIH COBRE in Neurodevelopment
and its Disorders (P20 GM148302) to T.R.S., and JST PRESTO (JPMJPR1883), JST FOREST
(JPMJFR2245), NHMRC Ideas Grant (APP1184899), and KAKENHI (20K23378) to T.K.S.

Declaration of interests

475 The authors declare no conflicts of interest.

Author contributions

478 T.R.S. and T.K.S. conceived the project; T.R.S. and T.K.S. designed the experiments. K. A.,
479 Y.K., K.M., and Z.H. contributed the project equally. K.A., Y.K., Z.H., O.M., T.I., T.R.S. and
480 T.K.S. carried out experiments; K.M., A.M., H.I., T.T., T.R.S. and T.K.S. prepared the analysis
481 codes. K.A., Z.H., A.M., A.M., E.S., T.R.S. and T.K.S. analysed the data. T.R.S. and T.K.S
482 interpreted the data and wrote the manuscript.

Materials and Methods

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rabbit Polyclonal anti-GFP	Thermo Fisher Scientific	Cat# A-11122, RRID:AB_221569
Sheep Polyclonal anti-Tyrosine Hydroxylase	Abcam	Cat# ab113, RRID:AB_297905
Donkey anti-Rabbit IgG (H+L) Antibody, Alexa Fluor™ 488 Conjugated	Thermo Fisher Scientific	Cat# A-21206, RRID:AB_2535792
Donkey Anti-Sheep IgG (H+L) Antibody, Alexa Fluor 568 Conjugated	Thermo Fisher Scientific	Cat# A-21099, RRID:AB_2535753
Bacterial and virus strains		
AAV2/1-hSynapsin1-FLEx-axon-jGCaMP8m-WPRE-SV40	University of South Carolina Viral Vectors Core	Addgene #216533
Bacterial and virus strains		
Normal Donkey Serum	Sigma-Aldrich	Cat# D9663, RRID:AB_2810235
Deposited data		
Mouse Brain Connectivity Atlas	Allen Brain Map	https://connectivity.brain-map.org/
Experimental models: Organisms/strains		
Mouse: Slc6a3$^{tm1.1(cre)Bkmn}$	The Jackson Laboratory	JAX: 006660
Software and algorithms		
MATLAB	Mathworks	RRID: SCR_001622 https://www.mathworks.com
Python	Mathworks	RRID:SCR_008394 https://www.anaconda.com/
Suite2p	Pachitariu et al., (2016)	https://github.com/MouseLand/suite2p
DeepLabCut	Mathis et al., 2018	https://github.com/DeepLabCut/DeepLabCut
B-spline Grid, Image and Point based Registration	Dirk-Jan Kroon	https://jp.mathworks.com/matlabcentral/fileexchange/20057-b-spline-grid-image-and-point-based-registration
Pytorch, TorchVision	Meta AI	https://pytorch.org/

EXPERIMENTAL MODEL DETAILS

Animals

All experimental procedures were approved by the Medical University of South Carolina, Monash University, Kagoshima University, and local institutions supervising animal experiments. Heterozygous dopamine transporter (DAT)-Cre mice (Slc6a3$^{tm1.1(cre)Bkmn}$, Jackson Laboratory, #006660, crossed with wild-type C57BL/6) was used in this study, including 12 mice for two-photon imaging and 10 for histology. Previous research utilized the same mouse line to express GCaMP6f in dopamine axon terminals in the mPFC that could be detected by one-photon fiber photometry (Kim et al., 2016). Mice of both sexes, aged >8 weeks were included. The mice were maintained in group housing (up to five mice per cage) and experiments were performed during the dark period of a 12-h light/12-h dark cycle.

METHOD DETAILS

Surgery

All surgical procedures were performed aseptically, with the mice under anesthesia with isoflurane. Lidocaine (subcutaneously at the incision), atropine (0.3 mg/kg, intraperitoneally), caprofen (5 mg/kg, intraperitoneally), and dexamethasone (2 mg/kg, intraperitoneally) were applied to prevent pain and brain edema. After surgery, the mice were allowed to recover for at least three days. No experimenter blinding was done.

Headplate implant and virus injection

A custom-made headpost was glued and cemented to the skull, and then, a small craniotomy (<0.5 mm) was performed over the VTA (~2.9–3.5 mm posterior and ~0.5 mm lateral from the

bregma). Inside the small craniotomy, axon-GCaMP virus (AAV2/1-hSynapsin1-FLEx-axon-

jGCaMP8m-WPRE-SV40) was volume-injected (Nanoject III, Drummond Scientific) to the VTA

through a pulled capillary glass (40–60 nL/site; depth: 4,200–4,400 µm; 15 min/injection). After

the injection, the craniotomy was sealed with a small piece of cover glass and silicon sealant

(Kwik-Cast) and animals were returned to their home cage.

For axon-GcaMP, we synthesized an axon-jGCaMP8m construct based on GAP43 (Broussard et

al., 2018), a linker (Broussard et al., 2018), and jGCaMP8m (Zhang et al., 2023), together with

restriction sites for SpeI and AscI. Then, the construct was inserted into a hSynapsin1-FLEx

vector, to make hSynapsin1-FLEx-axon-jGCaMP8m-WPRE-SV40. The plasmid has been

deposited into Addgene (#216533).

Microprism implant

After a 3-week waiting period of adeno-associated virus (AAV) expression, a microprism was

inserted for two-photon imaging as described previously (Low et al., 2014). A rectangular

craniotomy (4 x 2 mm) was made over the bilateral PFC (~1.5–3.5 mm anterior from the bregma),

and the dura was removed over the right hemisphere. Then, a microprism implant assembly was

inserted into the subdural space within the fissure (Figure 1B, E, F). The microprism was centered

~2.5 mm anterior to the bregma to avoid damaging bridging veins. Once implanted, the prism sat

flush against the opposing fissure wall, which contained the medial wall of the PFC (mainly the

prelimbic area) in the left hemisphere. The front face of the prism was oriented along the midline.

The assembly consisted of a right-angle microprism (2 x 2 x 1 mm, Prism RA N-BK7, Tower

Optical Corp.) and two coverslip layers (top layer: 4.5 x 3.0 mm, bottom layer: 3.6 x 1.8 mm), which

were glued by ultraviolet curing optical adhesive (Norland #81). The top layer of glass was

cemented to the skull with dental acrylic. Our assembly design (microprism of 2 x 2 x 1 mm, plus

double-layer glass) is different from the original report (microprism of 1.5 x 1.5 x 1.5 mm plus

single-layer glass) (Low et al., 2014) for the following reasons. First, the thinner microprism (1

mm in the anterior-posterior axis) was easier to insert into the bank, avoiding superficial veins

branching from the superior sagittal sinus. Second, the longer prism (2 mm in the dorsal-ventral

axis) could spare a wider imageable region below the superior sagittal sinus. Third, the double-

layer glass helped suppress brain movements.

Behavior

After the microprism implant surgery, the mice were allowed to recover in their home cages for

one week. After recovery, the mice underwent water scheduling (receiving 0.8–1 mL of water per

day). Then, the mice were pretrained for head fixation and for drinking water from a spout on a

linear passive treadmill (SpeedBelt, Phenosys) in a sound-proof blackout box for two days. After

the initial days of reward only, the animals received infrequent electrical shocks interspersed with

the reward. Once the animals experienced both reward and shock conditions, we started the two-

photon imaging sessions.

To monitor licking behavior, the face of each mouse was filmed with a camera at 65 Hz (CM3-U3-

13Y3M-CS, FLIR) using infrared illumination (850-nm light-emitting diode, IR30, CMVision or

M850F2, Thorlabs). To detect locomotion, the running speed on the treadmill was recorded at 30

kHz.

Rewarding and aversive stimuli

The mice received rewarding or aversive stimuli with unpredictable timing. The stimuli were

administered in a randomized order (rewarding stimuli: 7 out of 9 cases; aversive stimuli: 1 out of

9; control period: 1 out of 9), with a randomized inter-trial interval of 55–65 s. The mice exhibited

comfortable behavior on the treadmill for 1.5–2 h.

As a reward, 10 µL of sugar water was delivered through a water spout (Figure 2A), controlled by

a TTL pulse (200 ms) delivered to a syringe pump (PHM-107, Med Associates, Inc., USA). Based

on previous literature, a 10-µL reward is relatively large (Tsutsui-Kimura et al., 2020). Animals

typically underwent 100–200 reward trails. In some experiments, the reward volume was varied

between 0 and 15 μL (Figure 2–figure supplement 4). As an aversive stimulus (Kim et al., 2016;

de Jong et al., 2019; Lutas et al., 2019), a 1-s, 0.2-mA electrical current was delivered via a

stimulator (AM2100, A-M systems, USA) between two electrode pads attached to the mouse's

tail (Figure 2A). This current was considered to be mild, just strong enough to evoke locomotion.

When the current was doubled (Figure 2–figure supplement 4), the locomotion tended to become

stronger, but some animals stopped drinking water. Similarly, when the frequency of the aversive

stimuli was increased (e.g., 50% of trials), some mice were no longer motivated to drink the reward

water.

Classical conditioning

After three days of reward and aversive stimulus sessions, we trained the mice in reward and

aversive trace conditioning. The structure of the task is the same as that for the reward and

aversive stimulus sessions (reward condition: 7 out of 9 cases; aversive condition: 1 out of 9;

control condition: 1 out of 9; inter-trial interval: 55–65 s), except that auditory stimuli (9 or 13 kHz,

2 s) were presented 3 s before presenting the unconditioned stimuli (rewarding or aversive

stimulus). Anticipatory behavioral responses confirmed that the mice could discriminate the tone

frequency differences (Fig. 3D-G), consistent with a previous report showing that mice can

discriminate frequency differences down to 4–7% (de Hoz and Nelken, 2014). In three animals, 9

kHz tone was used for the reward-predictive cue, 13 kHz for the aversive-predictive cue. In the

remaining three animals, the tone association was reversed.

We separated the learning into three phases in addition to the first day. In the late phase,

anticipatory licking and running reach a saturation level, as evidenced by learning curves

spanning six periods (Figure 3–figure supplement 1).

When the anticipatory licking was stably manifested (Figure 3D, late phase), we included one

condition for an unexpected reward omission among the seven reward conditions (Figure 3–figure

supplement 3) and continued for two more days.

Timing control

Synchronization of the two-photon imaging, behavior camera image acquisition, reward delivery,

aversive stimulation, and sound presentation were achieved using digital and analogue output

from a National Instruments board (NI USB-6229), which was controlled by a custom-made

MATLAB program. We also recorded continuous signals (sampled at 30 kHz, PCIe-6363, National

Instruments) of the treadmill speed, the frame timing of two-photon imaging, the frame timing of

a behavior camera, copies of command waves to the syringe pump, the stimulator, and the

speaker.

Two-photon imaging

In vivo two photon imaging was performed using a table-mounted microscope   (Bergamo II,

Thorlabs or MOM, Sutter Instruments) and a data acquisition system. The light source was a

pulsed Ti:sapphire laser (MaiTai DeepSee eHP, SpectraPhysics, or Chameleon Ultra II,

Coherent) with dispersion compensation, with the laser wavelength set to 980 nm (Hasegawa et

al., 2017; Itokazu et al., 2018), which causes a higher fluorescent change in the GCaMP signal

and less scattering in the tissue than 920 nm. The laser power at the apochromatic objective lens

(16×, 0.80 NA, Nikon) was <70 mW, and we saw no bleaching. Green fluorescent photons were

filtered (ET525/70 m-2p) and collected by a hybrid photodetector (R11322U-40-01, Hamamatsu

Photonics) (Tischbirek et al., 2015) and a high-speed current amplifier (DHPCA-100, Femto).

Imaging frames were acquired at ~60 Hz and were downsampled offline. Images were collected

at a depth of 30–100 µm from the dural surface (up to ~200 x 200 µm). The small field of view at

a high sampling rate makes it possible to collect weak signals from small structures, as in spine

functional imaging (Jia et al., 2014).

Imaging fields were searched based on the presence of fiber morphology with at least occasional

calcium transients in the fibers, not based on the behavioral correlation of the transients. Fiber

morphology and spontaneous calcium transients were not reliably visible in axons deeper than

100 µm in a live-view mode, possibly because of the low signal-to-noise ratio. For each mouse,

imaging was performed for a single field per day in order to gain a sufficient number of repeats

with a 1-min inter-trial interval. In reward/aversive preference characterization (Figure 2), 1–2 sites

were imaged on different days. For the classical conditioning, only a single site was imaged during

the course of conditioning. Once the imaging site was determined on the first day, the reference

image of two-photon imaging was captured, in addition to the surface vessel image of one-photon

imaging. On subsequent days, these images were used to return to the same imaging site and

depth, comparing and overlaying the reference image and the ongoing imaging view.

Calcium imaging data analysis

Data processing

Imaging data was processed for motion correction and registration. Axons were detected for

region-of-interest (ROI) drawing using Suite2p (Pachitariu et al., 2016) and a custom-made

MATLAB program (Itokazu et al., 2018). A fluorescent trace for each ROI was generated, and

then the trace was normalized by the baseline fluorescence (F0, set as the 50^{th} percentile

fluorescence over a 30-s sliding window in order to remove any slow drifts in the baseline) to

produce a $\Delta F/F$ trace.

Dopamine axons were sparsely labeled in the mPFC, but the same axons needed to be excluded

based on correlation analysis among pairs (Petreanu et al., 2012; Sun et al., 2016; Itokazu et al.,

2018). The correlation coefficients of $\Delta F/F$ traces were calculated for axons in each plane, and

pairs showing a higher correlation (>0.65) (Itokazu et al., 2018) were considered to arise from the

same axon. The high correlation pairs were grouped into clusters, and the ROI with the largest

$\Delta F/F$ signal in each cluster was assigned to represent the cluster. The aforementioned procedure

was iterated repeatedly until the correlation between all remaining pairs fell below the threshold

of 0.65. Our results remained similar for different correlation threshold.

Reward, aversive, cue, and locomotion activity

For each axon, reward and aversive activity were evaluated. Reward activity was quantified as

an increase in ΔF/F by comparing the average ΔF/F between the control range (−2 to 0 s from the

onset of the reward TTL to a syringe pump) and the signal range (0 to 2 s). Similarly, aversive

activity was quantified as an increase in ΔF/F, based on the difference between the average ΔF/F

between the control range (−2 to 0 s from the onset of the electrical shock TTL) and the signal

range (0 to 2 s). Axons were considered to exhibit a significant response if the magnitude of either

activity was statistically larger than that of the baseline activity (Wilcoxon signed-rank test;

$p < 0.05$). Significant axons were classified as either reward-preferring (cyan) or aversive-

preferring clusters (magenta) based on k-means clustering, the separation of which coincided

approximately with the unity line of the reward/aversive scatter plot, as shown in Figure 2J.

The locomotion activity was quantified as an increase in ΔF/F by comparing the average ΔF/F

between the control range (−2 to 0 s from the locomotion initiation) and the signal range (0 to 2 s).

The locomotion initiation is defined in the "Running detection" section below.

During classical conditioning, activity was evaluated in a similar manner. For the conditioned cue

activity, the activity increase was computed by comparing the average ΔF/F between the control

range (−2 to 0 s from the onset of the predictive cue) and the signal range (0 to 2 s from the cue

onset). For the unconditioned response activity (reward or aversive), we compared the control

range (−2 to 0 s from the onset of the predictive cue) and the signal range (0 to 2 s from the onset

of the unconditioned stimulus). To investigate the preference for reward or aversive processing,

we used scatter plots (Figure 3I, J), similar to Figure 2J. The color-coded classification

(cyan/magenta) was based on k-means clustering, using the responses before classical

conditioning (Figure 2J).

Evaluation of brain movement

To compare the amount of brain movement between the two different microprism assemblies

(Figure 1–figure supplement 2), we obtained x- and y-axis shifts of acquired images caused by

the brain movement. The shifts were computed by the Suite2p program and used for image

registration (Pachitariu et al., 2016). We quantified the brain shift using two metrics: root mean

square and large transient movement. First, the root mean square was computed based on

sequential shifts in pixel in x- and y-dimensions that were combined trigonometrically (Figure 1–

figure supplement 2B, D). Second, to detect large transient movement events, combined brain

shift traces were filtered (Butterworth, at 1.5 Hz), and events larger than 5 μm (16 pixels) were

detected as movement events (black dots in B).

Behavioral analysis

Licking detection

To track the movement of the tongue, videos of orofacial movement (65 Hz, side view) were

processed using DeepLabCut (Mathis et al., 2018) (Figure 2–figure supplement 1). The tip of the

tongue, the location of the water spout and the position of the nose were labelled in randomly

selected ~200 frames from 6 animals. In frames when the tongue was inside the mouth and was

not visible, we estimated its location from the lips and jaw, instead of not labelling the tongue in

these frames. This estimation prevented DeepLabCut from making a completely wrong guess in

labelling the tongue for these frames.

The learning process was divided into three equal-duration periods. We confirmed that the

division into six periods resulted in a saturating discrimination curve for anticipatory licking in the

fifth and sixth periods (Figure 3–figure supplement 1). These last two periods in the six-period

division correspond to the 'late phase" of the three-period division that we used.

Running detection

The speed of treadmill was monitored as the output from a SpeedBelt apparatus (Phenosys). The

locomotion period was defined as the duration in which the treadmill speed was above the median

+ 0.5 x standard deviation for more than 200 ms. Then, the initiation of the locomotion period was

defined as a time point preceded by a non-locomotion period (when the running speed is below

the threshold) of at least 0.5 s.

Error analysis

To investigate how the cue discrimination of the animals impacts dopamine axonal activity, we

separated the trials into those with correct discriminating behavior and those with incorrect

behavior for reward and aversive conditions (Figure 4A). We used two types of discriminating

behaviors, anticipatory licking and facial expressions (see below). We analyzed the late phase of

the classical conditioning when animals showed robust anticipatory licking (Figure 4–figure

supplement 1) or facial expressions (Figure 4–figure supplement 2E). In one animal, anticipatory

licking was not seen in the aversive condition (the fourth from the top of Figure 4A) during the late

phase, so we analyzed the middle phase for that animal.

Machine learning analysis of facial expressions, licking, and running

The anticipation of animals regarding upcoming unconditioned stimuli (reward or electrical shock)

was quantified based on auditory predictive cues using a machine learning classifier (random

forest classifier) (Breiman, 2001).

Facial expressions were filmed by an infrared camera and analyzed with a random forest classifier

combined with a deep neural network (Figure 4–figure supplement 2A-D). First, features of facial

expressions were extracted from a given temporal series of frames (i.e., a video) using a deep

neural network model, the ResNet3D model. The ResNet3D model is a pretrained network

consisting of 18 layers, optimized for videos and provided by PyTorch (Tran et al., 2018). The

output from the final convolutional layer was fed into the random forest classifier. In our study,

training was performed not on the pretrained ResNet3D, but on the random forest classifier. The

random forest classifier was trained and tested with independent trials by five-fold cross-validation

within each day. To prevent the random forest classifier from being overfit, only the top 400

features of the input were used, which were ranked by the F-value. To train the random forest

classifier equally to the reward and aversive conditions despite their imbalanced frequency (7 or

1 out of 8 trials, Figure 3A), an ensemble training technique was used (Wallace et al., 2011). the

discrimination accuracy for reward and aversive conditions was computed separately and an

average was taken with equal weights as a final discrimination accuracy. The equal weights

prevented the accuracy computation from being dominated by the reward condition, which

occurred more frequently than the aversive condition. To investigate the time course of the

discrimination accuracy, accuracy computation was performed for a 500-ms time window instead

of a 2-s window, and the window was systematically shifted by 160 ms (Figure 4–figure

supplement 2E).

The discrimination accuracy based on anticipatory licking was also computed (Figure 4–figure

supplement 1). To enable a comparison among facial features and licking, the random forest

classifier was used. Instead of 400 features (facial expressions), the random forest classifier was

fed with one feature (either the number of licking instances).

Histology

Animals were perfused with 4% paraformaldehyde (PFA) in phosphate-buffered saline (PBS).

GCaMP or tyrosine hydroxylase immunostaining was performed using standard procedures

(Figure 1C, D). Coronal slices (thickness, 30 μm) were cut using a cryostat (Leica Microsystems)

and blocked in carrier solution (5% bovine serum albumin; 0.3% Triton X-100 in 0.1 M PBS) for

2 h at room temperature on a shaker. For GFP staining, slices were then incubated with anti-

green fluorescent protein (GFP) primary antibody (anti-GFP, 1:1000, A11122, Invitrogen) for 18 h

at 4°C on a shaker. After three rinses with 0.1 M PBS for 30 min, sections were incubated with

Alexa-Fluor-488-conjugated donkey anti-rabbit secondary antibody (Invitrogen, 1:500 in carrier

solution) for 1 h at room temperature on a shaker. For tyrosine hydroxylase staining, additional

incubation with anti- tyrosine hydroxylase (TH) primary antibody (anti-TH, 1:200, ab113) and

Alexa-Fluor-568-conjugated donkey anti-sheep secondary antibody (Invitrogen, 1:500) was

included. Cell nuclei were stained with DAPI (1:1000; D523, Dojindo). After a few additional rinses

for 30 min in 0.1 M PBS were performed, slices were mounted on slide glasses for imaging.

Images were acquired using a confocal laser-scanning microscopy (FV3000, Olympus) and a

fluorescence microscope (VS200, Olympus).

EXPERIMENTAL DESIGN AND STATISTICAL ANALYSIS

Data are described as the median ± s.e.m. unless otherwise noted. Statistical significance

was assessed using the non-parametric Wilcoxon signed-rank test, unless stated otherwise.

Significance levels of data were denoted as $* P < 0.05$, $** P < 0.01$ and $*** P < 0.001$. $P >$

0.05 was insignificant and was denoted as n.s.

RESOURCE AVAILABILITY

Lead contact

Further information and requests for reagents may be directed to the Lead Contact, Takashi R

Sato (satot@musc.edu).

Materials availability

The plasmid construct has been deposited to Addgene (#216533).

Data and code availability

All data and corresponding analysis codes reported in this study are available at GitHub.

https://github.com/pharmedku/2024-elife-da-axon

References

Abercrombie ED, Keefe KA, DiFrischia DS, Zigmond MJ (1989) Differential effect of stress on in vivo dopamine release in striatum, nucleus accumbens, and medial frontal cortex. J Neurochem 52:1655-1658.

Ahn S, Phillips AG (1999) Dopaminergic correlates of sensory-specific satiety in the medial prefrontal cortex and nucleus accumbens of the rat. J Neurosci 19:RC29.

Amo R, Matias S, Yamanaka A, Tanaka KF, Uchida N, Watabe-Uchida M (2022) A gradual temporal shift of dopamine responses mirrors the progression of temporal difference error in machine learning. Nat Neurosci 25:1082-1092.

Andermann ML, Gilfoy NB, Goldey GJ, Sachdev RN, Wolfel M, McCormick DA, Reid RC, Levene MJ (2013) Chronic cellular imaging of entire cortical columns in awake mice using microprisms. Neuron 80:900-913.

Aransay A, Rodriguez-Lopez C, Garcia-Amado M, Clasca F, Prensa L (2015) Long-range projection neurons of the mouse ventral tegmental area: a single-cell axon tracing analysis. Front Neuroanat 9:59.

Arnsten AF, Dudley AG (2005) Methylphenidate improves prefrontal cortical cognitive function through alpha2 adrenoceptor and dopamine D1 receptor actions: Relevance to therapeutic effects in Attention Deficit Hyperactivity Disorder. Behav Brain Funct 1:2.

Arnsten AF, Wang MJ, Paspalas CD (2012) Neuromodulation of thought: flexibilities and vulnerabilities in prefrontal cortical network synapses. Neuron 76:223-239.

Azcorra M, Gaertner Z, Davidson C, He Q, Kim H, Nagappan S, Hayes CK, Ramakrishnan C, Fenno L, Kim YS, Deisseroth K, Longnecker R, Awatramani R, Dombeck DA (2023) Unique functional responses differentially map onto genetic subtypes of dopamine neurons. Nat Neurosci 26:1762-1774.

Bassareo V, De Luca MA, Di Chiara G (2002) Differential Expression of Motivational Stimulus Properties by Dopamine in Nucleus Accumbens Shell versus Core and Prefrontal Cortex. J Neurosci 22:4709-4719.

Bissonette GB, Powell EM, Roesch MR (2013) Neural structures underlying set-shifting: roles of medial prefrontal cortex and anterior cingulate cortex. Behav Brain Res 250:91-101.

Breiman L (2001) Random Forests. Machine Learning 45:5-32.

Bromberg-Martin ES, Matsumoto M, Hikosaka O (2010) Dopamine in motivational control: rewarding, aversive, and alerting. Neuron 68:815-834.

Broussard GJ, Liang Y, Fridman M, Unger EK, Meng G, Xiao X, Ji N, Petreanu L, Tian L (2018) In vivo measurement of afferent activity with axon-specific calcium imaging. Nat Neurosci 21:1272-1280.

Chiodo LA, Antelman SM, Caggiula AR, Lineberry CG (1980) Sensory stimuli alter the discharge rate of dopamine (DA) neurons: evidence for two functional types of DA cells in the substantia nigra. Brain Res 189:544-549.

de Hoz L, Nelken I (2014) Frequency tuning in the behaving mouse: different bandwidths for discrimination and generalization. PLoS One 9:e91676.

de Jong JW, Afjei SA, Pollak Dorocic I, Peck JR, Liu C, Kim CK, Tian L, Deisseroth K, Lammel S (2019) A Neural Circuit Mechanism for Encoding Aversive Stimuli in the Mesolimbic Dopamine System. Neuron 101:133-151 e137.

Desimone R, Duncan J (1995) Neural mechanisms of selective visual attention. Annu Rev Neurosci 18:193-222.

Dolensek N, Gehrlach DA, Klein AS, Gogolla N (2020) Facial expressions of emotion states and their neuronal correlates in mice. Science 368:89-94.

Ellwood IT, Patel T, Wadia V, Lee AT, Liptak AT, Bender KJ, Sohal VS (2017) Tonic or Phasic
Stimulation of Dopaminergic Projections to Prefrontal Cortex Causes Mice to Maintain or
Deviate from Previously Learned Behavioral Strategies. J Neurosci 37:8315-8329.
Engelhard B, Finkelstein J, Cox J, Fleming W, Jang HJ, Ornelas S, Koay SA, Thiberge SY, Daw
ND, Tank DW, Witten IB (2019) Specialized coding of sensory, motor and cognitive
variables in VTA dopamine neurons. Nature 570:509-513.
Finlay JM, Zigmond MJ, Abercrombie ED (1995) Increased dopamine and norepinephrine release
in medial prefrontal cortex induced by acute and chronic stress: effects of diazepam.
Neuroscience 64:619-628.
Fuster JM (2015) The prefrontal cortex: Academic Press/ Elsevier.
Grace AA (2016) Dysregulation of the dopamine system in the pathophysiology of schizophrenia
and depression. Nat Rev Neurosci 17:524-532.
Granon S, Passetti F, Thomas KL, Dalley JW, Everitt BJ, Robbins TW (2000) Enhanced and
impaired attentional performance after infusion of D1 dopaminergic receptor agents into
rat prefrontal cortex. J Neurosci 20:1208-1215.
Guarraci FA, Kapp BS (1999) An electrophysiological characterization of ventral tegmental area
dopaminergic neurons during differential pavlovian fear conditioning in the awake rabbit.
Behav Brain Res 99:169-179.
Gunaydin LA, Grosenick L, Finkelstein JC, Kauvar IV, Fenno LE, Adhikari A, Lammel S,
Mirzabekov JJ, Airan RD, Zalocusky KA, Tye KM, Anikeeva P, Malenka RC, Deisseroth
K (2014) Natural neural projection dynamics underlying social behavior. Cell 157:1535-
1551.
Hasegawa M, Majima K, Itokazu T, Maki T, Albrecht UR, Castner N, Izumo M, Sohya K, Sato TK,
Kamitani Y, Sato TR (2017) Selective Suppression of Local Circuits during Movement
Preparation in the Mouse Motor Cortex. Cell Rep 18:2676-2686.
Hernandez L, Hoebel BG (1990) Feeding can enhance dopamine turnover in the prefrontal cortex.
Brain Res Bull 25:975-979.
Hoexter MQ, Fadel G, Felicio AC, Calzavara MB, Batista IR, Reis MA, Shih MC, Pitman RK,
Andreoli SB, Mello MF, Mari JJ, Bressan RA (2012) Higher striatal dopamine transporter
density in PTSD: an in vivo SPECT study with [(99m)Tc]TRODAT-1. Psychopharmacology
(Berl) 224:337-345.
Howe MW, Dombeck DA (2016) Rapid signalling in distinct dopaminergic axons during
locomotion and reward. Nature 535:505-510.
Howes OD, Kapur S (2009) The dopamine hypothesis of schizophrenia: version III--the final
common pathway. Schizophr Bull 35:549-562.
Isoda M, Hikosaka O (2007) Switching from automatic to controlled action by monkey medial
frontal cortex. Nat Neurosci 10:240-248.
Itokazu T, Hasegawa M, Kimura R, Osaki H, Albrecht UR, Sohya K, Chakrabarti S, Itoh H, Ito T,
Sato TK, Sato TR (2018) Streamlined sensory motor communication through cortical
reciprocal connectivity in a visually guided eye movement task. Nat Commun 9:338.
Jean-Richard-Dit-Bressel P, Killcross S, McNally GP (2018) Behavioral and neurobiological
mechanisms of punishment: implications for psychiatric disorders.
Neuropsychopharmacology 43:1639-1650.
Jia H, Varga Z, Sakmann B, Konnerth A (2014) Linear integration of spine Ca2+ signals in layer
4 cortical neurons in vivo. Proc Natl Acad Sci U S A 111:9277-9282.
Kamigaki T, Dan Y (2017) Delay activity of specific prefrontal interneuron subtypes modulates
memory-guided behavior. Nat Neurosci 20:854-863.
Kim CK, Yang SJ, Pichamoorthy N, Young NP, Kauvar I, Jennings JH, Lerner TN, Berndt A, Lee
SY, Ramakrishnan C, Davidson TJ, Inoue M, Bito H, Deisseroth K (2016) Simultaneous
fast measurement of circuit dynamics at multiple sites across the mammalian brain. Nat
Methods 13:325-328.

Komiyama T, Sato TR, O'Connor DH, Zhang YX, Huber D, Hooks BM, Gabitto M, Svoboda K
(2010) Learning-related fine-scale specificity imaged in motor cortex circuits of behaving
mice. Nature 464:1182-1186.
Lammel S, Hetzel A, Hackel O, Jones I, Liss B, Roeper J (2008) Unique properties of
mesoprefrontal neurons within a dual mesocorticolimbic dopamine system. Neuron
57:760-773.
Lee JY, Jun H, Soma S, Nakazono T, Shiraiwa K, Dasgupta A, Nakagawa T, Xie JL, Chavez J,
Romo R, Yungblut S, Hagihara M, Murata K, Igarashi KM (2021) Dopamine facilitates
associative memory encoding in the entorhinal cortex. Nature 598:321-326.
Lindstrom LH, Gefvert O, Hagberg G, Lundberg T, Bergstrom M, Hartvig P, Langstrom B (1999)
Increased dopamine synthesis rate in medial prefrontal cortex and striatum in
schizophrenia indicated by L-(beta-11C) DOPA and PET. Biol Psychiatry 46:681-688.
Low RJ, Gu Y, Tank DW (2014) Cellular resolution optical access to brain regions in fissures:
imaging medial prefrontal cortex and grid cells in entorhinal cortex. Proc Natl Acad Sci U
S A 111:18739-18744.
Lutas A, Kucukdereli H, Alturkistani O, Carty C, Sugden AU, Fernando K, Diaz V, Flores-
Maldonado V, Andermann ML (2019) State-specific gating of salient cues by midbrain
dopaminergic input to basal amygdala. Nat Neurosci 22:1820-1833.
Ma L, Day-Cooney J, Benavides OJ, Muniak MA, Qin M, Ding JB, Mao T, Zhong H (2022)
Locomotion activates PKA through dopamine and adenosine in striatal neurons. Nature
611:762-768.
Mantz J, Thierry AM, Glowinski J (1989) Effect of noxious tail pinch on the discharge rate of
mesocortical and mesolimbic dopamine neurons: selective activation of the mesocortical
system. Brain Res 476:377-381.
Mathis A, Mamidanna P, Cury KM, Abe T, Murthy VN, Mathis MW, Bethge M (2018) DeepLabCut:
markerless pose estimation of user-defined body parts with deep learning. Nat Neurosci
21:1281-1289.
Matsuda W, Furuta T, Nakamura KC, Hioki H, Fujiyama F, Arai R, Kaneko T (2009) Single
nigrostriatal dopaminergic neurons form widely spread and highly dense axonal
arborizations in the neostriatum. J Neurosci 29:444-453.
Matsumoto M, Hikosaka O (2009) Two types of dopamine neuron distinctly convey positive and
negative motivational signals. Nature 459:837-841.
Menegas W, Babayan BM, Uchida N, Watabe-Uchida M (2017) Opposite initialization to novel
cues in dopamine signaling in ventral and posterior striatum in mice. Elife 6.
Miller EK, Cohen JD (2001) An integrative theory of prefrontal cortex function. Annu Rev Neurosci
24:167-202.
Miller EK, Wallis JD (2009) Executive Function and Higher-Order Cognition: Definition and Neural
Substrates: Oxford: Academic Press.
Nee DE, Kastner S, Brown JW (2011) Functional heterogeneity of conflict, error, task-switching,
and unexpectedness effects within medial prefrontal cortex. Neuroimage 54:528-540.
Ogawa M, van der Meer MA, Esber GR, Cerri DH, Stalnaker TA, Schoenbaum G (2013) Risk-
responsive orbitofrontal neurons track acquired salience. Neuron 77:251-258.
Okubo Y, Suhara T, Sudo Y, Toru M (1997) Possible role of dopamine D1 receptors in
schizophrenia. Mol Psychiatry 2:291-292.
Otis JM, Namboodiri VM, Matan AM, Voets ES, Mohorn EP, Kosyk O, McHenry JA, Robinson JE,
Resendez SL, Rossi MA, Stuber GD (2017) Prefrontal cortex output circuits guide reward
seeking through divergent cue encoding. Nature 543:103-107.
Ott T, Nieder A (2019) Dopamine and Cognitive Control in Prefrontal Cortex. Trends Cogn Sci
23:213-234.
Ott T, Jacob SN, Nieder A (2014) Dopamine receptors differentially enhance rule coding in
primate prefrontal cortex neurons. Neuron 84:1317-1328.

Pachitariu M, Stringer C, Dipoppa M, Schröder S, Rossi L, Dalgleish H, Carandini M, Harris KD
(2016) Suite2p: beyond 10,000 neurons with standard two-photon microscopy. bioRxiv.
Patriarchi T, Cho JR, Merten K, Howe MW, Marley A, Xiong WH, Folk RW, Broussard GJ, Liang
R, Jang MJ, Zhong H, Dombeck D, von Zastrow M, Nimmerjahn A, Gradinaru V, Williams
JT, Tian L (2018) Ultrafast neuronal imaging of dopamine dynamics with designed
genetically encoded sensors. Science 360.
Petreanu L, Gutnisky DA, Huber D, Xu NL, O'Connor DH, Tian L, Looger L, Svoboda K (2012)
Activity in motor-sensory projections reveals distributed coding in somatosensation.
Nature 489:299-303.
Popescu AT, Zhou MR, Poo MM (2016) Phasic dopamine release in the medial prefrontal cortex
enhances stimulus discrimination. Proc Natl Acad Sci U S A 113:E3169-3176.
Poulin JF, Caronia G, Hofer C, Cui Q, Helm B, Ramakrishnan C, Chan CS, Dombeck DA,
Deisseroth K, Awatramani R (2018) Mapping projections of molecularly defined dopamine
neuron subtypes using intersectional genetic approaches. Nat Neurosci 21:1260-1271.
Rescorla RA, Wagner AR (1972) Classical Conditioning II. Current Research and Theory,:
Appleton-Century- Crofts, New York.
Ridderinkhof KR, Ullsperger M, Crone EA, Nieuwenhuis S (2004) The role of the medial frontal
cortex in cognitive control. Science 306:443-447.
Sawaguchi T, Goldman-Rakic PS (1994) The role of D1-dopamine receptor in working memory:
local injections of dopamine antagonists into the prefrontal cortex of rhesus monkeys
performing an oculomotor delayed-response task. J Neurophysiol 71:515-528.
Schultz W, Dayan P, Montague PR (1997) A neural substrate of prediction and reward. Science
275:1593-1599.
Seamans JK, Yang CR (2004) The principal features and mechanisms of dopamine modulation
in the prefrontal cortex. Prog Neurobiol 74:1-58.
Sharpe MJ, Stalnaker T, Schuck NW, Killcross S, Schoenbaum G, Niv Y (2019) An Integrated
Model of Action Selection: Distinct Modes of Cortical Control of Striatal Decision Making.
Annu Rev Psychol 70:53-76.
St Onge JR, Ahn S, Phillips AG, Floresco SB (2012) Dynamic fluctuations in dopamine efflux in
the prefrontal cortex and nucleus accumbens during risk-based decision making. J
Neurosci 32:16880-16891.
Sun W, Tan Z, Mensh BD, Ji N (2016) Thalamus provides layer 4 of primary visual cortex with
orientation- and direction-tuned inputs. Nat Neurosci 19:308-315.
Sutton RS, Barto AG (1981) Toward a modern theory of adaptive networks: expectation and
prediction. Psychol Rev 88:135-170.
Takehara-Nishiuchi K, McNaughton BL (2008) Spontaneous changes of neocortical code for
associative memory during consolidation. Science 322:960-963.
Thierry AM, Tassin JP, Blanc G, Glowinski J (1976) Selective activation of mesocortical DA
system by stress. Nature 263:242-244.
Tischbirek C, Birkner A, Jia H, Sakmann B, Konnerth A (2015) Deep two-photon brain imaging
with a red-shifted fluorometric Ca2+ indicator. Proc Natl Acad Sci U S A 112:11377-11382.
Tran D, Wang H, Torresani L, Ray J, LeCun Y, Paluri M (2018) A Closer Look at Spatiotemporal
Convolutions for Action Recognition. In: 2018 IEEE/CVF Conference on Computer Vision
and Pattern Recognition, pp 6450-6459.
Tsutsui-Kimura I, Matsumoto H, Akiti K, Yamada MM, Uchida N, Watabe-Uchida M (2020) Distinct
temporal difference error signals in dopamine axons in three regions of the striatum in a
decision-making task. Elife 9.
Tuttle AH, Molinaro MJ, Jethwa JF, Sotocinal SG, Prieto JC, Styner MA, Mogil JS, Zylka MJ
(2018) A deep neural network to assess spontaneous pain from mouse facial expressions.
Mol Pain 14:1744806918763658.

Vander Weele CM, Siciliano CA, Tye KM (2019) Dopamine tunes prefrontal outputs to orchestrate
aversive processing. Brain Res 1713:16-31.
Vander Weele CM, Siciliano CA, Matthews GA, Namburi P, Izadmehr EM, Espinel IC, Nieh EH,
Schut EHS, Padilla-Coreano N, Burgos-Robles A, Chang CJ, Kimchi EY, Beyeler A,
Wichmann R, Wildes CP, Tye KM (2018) Dopamine enhances signal-to-noise ratio in
cortical-brainstem encoding of aversive stimuli. Nature 563:397-401.
Verharen JPH, Zhu Y, Lammel S (2020) Aversion hot spots in the dopamine system. Curr Opin
Neurobiol 64:46-52.
Wallace BC, Small K, Brodley CE, Trikalinos TA (2011) Class Imbalance, Redux. In: 2011 IEEE
11th International Conference on Data Mining, pp 754-763.
Yuan L, Dou YN, Sun YG (2019) Topography of Reward and Aversion Encoding in the Mesolimbic
Dopaminergic System. J Neurosci 39:6472-6481.
Zhang Y et al. (2023) Fast and sensitive GCaMP calcium indicators for imaging neural populations.
Nature 615:884-891.
970

Figure 1: Two-photon imaging of dopaminergic axons projecting to the mPFC.

(A, B) Experimental design. The activity of midbrain dopamine neurons projecting to the mPFC

was measured by two-photon calcium imaging of their axons. The axons were accessed through

a microprism that bends the optical axis inside the brain (black arrows in B). (C) GCaMP was

expressed virally in dopamine neurons in DAT-Cre transgenic mice. A coronal section shows

GCaMP expression in the VTA, demonstrating that AAV-axon-DIO-jGCaMP8m was injected into

the VTA. (D) jGCaMP8m-expressing neurons were positive for tyrosine hydroxylase (TH), a

marker for dopamine neurons. (E, F) Dorsal view of a mouse head implanted with a microprism

assembly. The microprism was 1 x 2 mm. (G) An example *in vivo* image of jGCaMP8m-expressing

axons.

Figure 2: Dopaminergic axonal response to unexpected rewarding or aversive stimuli.

(A) Experimental design. Mice were placed under a two-photon microscope on a linear treadmill

and were given unexpected rewarding (water drops) or aversive (electrical shock to the tail) stimuli.

The mouse's face was filmed with an infrared camera to track the tongue position. (B) Example

behavioral response to rewarding (top, tongue position) or aversive (bottom, treadmill speed)

stimuli. (C) Average behavioral response on a single day for the same animal shown in B. (D)

Representative image for dopamine axons labeled with jGCaMP8m. Scale bar: 50 µm. (E, F)

Heatmaps of rewarding (E) or aversive (F) stimuli for the same imaging plane shown in D. (G–I)

Calcium response for rewarding (left) and aversive (right) stimuli of three example axons. (J)

Comparison between reward (x-axis) and aversive (y-axis) responses for dopamine axons (n =

162). Statistically significant axons were labeled in either cyan (reward-preferring axons, n = 25)

or magenta (aversive-preferring axons, n = 75). Vector averages representing reward-preferring

axons and aversive-preferring axons are depicted as cyan and magenta arrows. (K) Histogram of

the polar angle of the scatter plot in J (n = 162). The solid line indicates probability density,

estimated by kernel smoothing. A value of 0° represents axons that solely prefer rewards,

whereas 90° represents those that solely prefer aversive stimuli.

Figure 3: Classical conditioning induced behavioral and neural changes.

(A) Experimental design. Auditory cues were presented before unconditioned stimuli (rewarding

or aversive stimuli). (B, C) Behavioral changes in one example animal across 12 days (B: licking,

C: running). In the reward condition, the animal gradually developed anticipatory licking (B, left).

In the aversive condition, the animal usually ran after the shock delivery, but sometimes even

before the delivery (C, right). Licking and running traces were normalized to the instantaneous

maximum values for this animal and then averaged over a single day. (D-G) Anticipatory behavior

during classical learning (n = 6 animals). In the late phase of learning, anticipatory licking was

primarily observed in the reward condition (D) but not in the aversive condition (E). Anticipatory

running was seen more often in the aversive condition (G) than in the reward condition (F). (H)

Activity change in one example axon across 12 days. The axon was from the same animal shown

in B, C. (I, J) Learning induced changes in response to conditioned cues (I) and unconditioned

stimuli (J) for aversive-preferring axons (magenta) and reward-preferring axons (cyan) together

with non-significant axons (black). Aversive- and reward-preferring axons were defined before the

start of classical training. The x-axis represents the reward condition, and the y-axis represents

the aversive condition. n = 47 for aversive-preferring axons, n = 12 for reward-preferring axons.

Vector averages representing aversive-preferring axons and reward-preferring axons are

depicted as magenta and cyan arrows, overlaid in each panel. (K) Learning induced a change in

the amplitude of unconditioned response of aversive-preferring axons for the aversive condition

(magenta solid line) and reward condition (magenta dotted line) and that of reward-preferring

axons for reward condition (cyan solid line) and aversive condition (cyan dotted line). Number of

axons is the same as I, J. (L) The polar angle of the scatter plot in J. The magenta line represents

aversive-preferring axons and the cyan line represents reward-preferring axons. (M) Similar to K,

but for the conditioned response. (N) Similar to L, but for the conditioned response.

Figure 4: Axonal cue response in trials with correct or incorrect cue discrimination.

(A) Classification of trials based on the behavioral response that occurred between the conditioned stimulus onset and unconditioned stimulus onset. Such behaviors include anticipatory licking or facial expressions. (B) Comparison of cue response between correct (x-axis) and incorrect (y-axis) trials based on anticipatory licking (magenta, aversive-preferring axons, n = 44; cyan, reward-preferring axons, n = 12). The left panel shows the reward cue response, and the right panel shows the aversive cue response. (C) Preference for reward or aversive predictive

cues in correct (left panel) and incorrect (right) discrimination. Magenta represents aversive-

preferring axons (n = 44) and cyan represents reward-preferring axons (n = 12). Vector averages

representing aversive-preferring axons and reward-preferring axons are depicted as magenta and

cyan arrows. Note stronger preference for aversive cue as a population in correct discrimination.

(D) Similar to B, but based on facial expressions (magenta, aversive-preferring axons, n = 47;

cyan, reward-preferring axons, n = 12) (E) Similar to C, but based on facial expressions (magenta,

aversive-preferring axons, n = 47; cyan, reward-preferring axons, n = 12). As in C, preference for

the aversive cue is stronger in correct discrimination.

Figure 1–figure supplement 1: Sparse dopaminergic projections to the the mPFC.
(A) Example experiment of AAV-GFP injection into the VTA in the Allen Brain Atlas Data Portal (experiment No. 156314762). (B) Sparse GFP-expressing axons in the mPFC region, including the prelimbic area (the same experiment shown in A). (C) Sparse jGCaMP8m expression at the mPFC was observed in our experiment. The approximate position of the image is indicated as a square in B. (D) The signal of projected GFP axons at the prelimbic area was 1.3% ± 0.9% of that at the caudoputamen. All experiments at the Allen Portal used wild-type or tyrosine hydroxylase-Cre mice that received an AAV-GFP injection into the VTA (experiment No. 156314762, 120572378, 127796728).

Figure 1– figure supplement 2: Double-layer glass significantly reduces brain movement.
(A) An assembly design similar to Low et al., 2014. (B) Brain movement during imaging with the assembly design similar to Low's. The brain shift was computed by the Suite2p program, and shifts in pixels in the x and y dimensions were combined trigonometrically. The frame rate was 58 Hz (200,000 frames corresponds to ~57 minutes). Each pixel represents 0.31 μm. Movement events larger than 16 pixels (4.96 μm) were detected as red dots. (C) Our own assembly design with an additional layer of glass. (D) Similar to B, but with our assembly design. (E) The frequency of movement events larger than 5 μm for Low's design (standard: n = 3) and ours (extra layer: n=5) for the initial (left panel) and latter (right panel) halves of imaging. The root mean square is statistically smaller with our design (initial half: p = 0.002, latter half: p = 0.002, t-test). (F) Similar to E, but for root mean square of the brain shift. The events are less frequent with our design (initial half: p = 0.018, latter half: p = 0.005, t-test).

Figure 1- figure supplement 3: Long-term imaging of dopamine axons across days.
(A) The imaged acquired on the 1st day. (B) Similar to A, but on the 4th day. (C) Similar to A, but on the 9th day. The same imaging plane as Figure 1G.

Figure 2–figure supplement 1: Lick detection using DeepLabCut.
(A) Sample frames used by DeepLabCut. The tongue tip was labeled as a red dot and the water port as a blue dot (left). When the tongue was not visible (right), we intentionally trained DeepLabCut to consider the tongue position as the inside of the oral cavity (red dot). (B) Example of a tongue position trace. Gray shaded period is expanded in C. (C) Tongue movement that crossed the water port position (blue line) was counted as a lick (black ticks).

Figure 2–figure supplement 2: Insignificant locomotion activity of dopaminergic axons. No significant increase in locomotion initiation (for aversive-preferring axons, p = 0.583, n = 75; for reward-preferring axons, p = 0.92, n = 25; for all axons, p = 0.23, n = 162, Wilcoxon signed-rank text).

Figure 2–figure supplement 3: Comparison between reward (x-axis) and aversive (y-axis) responses for each mouse (n = 8 animals), similar to Figure 2J (for all animals).

Figure 2–figure supplement 4: Dopaminergic axonal activity depends on the reward volume and shock current.

(A) Axonal activity increased as the reward volume increased. Differences were significant between 0 and 5 μL (p < 0.001), between 2.5 μL and 5 μL (p < 0.001), and between 5 μL and 10 μL (p < 0.001, n = 39, 2 animals). However, the differences were not significant between 0 μL and 2.5 μL (p = 0.06) or between 10 μL and 15 μL (p = 0.97), indicating saturation at 10 μL. (B) Axonal activity increased as the electrical current increased. The difference was significant between 0 and 0.2 mA (p < 0.001, n = 61, 2 animals) and between 0.2 and 0.4 mA (p < 0.001). * indicates p < 0.05, ** P < 0.01, and *** P < 0.001.

Figure 3–figure supplement 1: Anticipatory licking and running during the classical conditioning ($n = 6$). (A) The conditioning period is divided into six phases plus the first day, instead of three phases (Fig. 3D-G). Licking rate was saturated at 5^{th} and 6^{th} phases. (B) Similar to A, but for anticipatory running.

Figure 3–figure supplement 2: Population activity of aversive- and reward-preferring axons throughout learning (aversive-preferring axons, n = 47; reward-preferring axons, n =12).
(A-D) Averaged calcium traces of aversive-preferring axons for the first day (A), early phase (B), middle phase (C), and late phase (D) of classical conditioning for the reward condition (left column) and aversive condition (right column). (E-H) Similar to A-D, but for reward-preferring axons.

Figure 3–figure supplement 3: Activity is not suppressed during reward omission.
(A) Population calcium trace for reward-preferring axons aligned to reward omission (n = 12 reward-preferring axons). (B) No significant change occurs during reward omission (for reward-preferring axons, p = 0.62, n = 12; Wilcoxon signed-rank test). Control and omission 2 s-periods are indicated in A.

Figure 3–figure supplement 4: Response in the first trials in each session.
(A) Comparison between CS_{reward} responses in the first trial and the remaining trials. (B) Similar to A, but for US_{reward}. (C) Similar to A, but for $CS_{aversive}$. (D) Similar to A, but for $US_{aversive}$. Note that for aversive-preferring axons, the response in the first trial was stronger for CS_{reward} responses (A, $p = 0.0002$, $n = 47$, Wilcoxon signed-rank test) and weaker for $CS_{aversive}$ responses (C, $p = 0.002$, $n = 47$). The response in the first trial was stronger for US_{reward} response (B, $p = 0.002$, $n = 47$), and was not significantly different for $US_{aversive}$ (D, $p = 0.36$, $n = 47$ aversive-preferring axons). For reward-preferring axons, there was no significant difference between the first trial and the rest (CS_{reward}, $p = 0.23$; $CS_{aversive}$, $p = 0.38$; US_{reward}, $p = 0.85$; $US_{aversive}$, $p = 1.00$).

Figure 4–figure supplement 1: Machine learning discriminates auditory cues based on anticipatory licking. On the first day, cue discrimination was nearly 50%, indicating that discrimination is random. However, as the training proceeded, discrimination was improved, reaching 79.2 ± 6.3% at the late phase (n = 6 animals).

A

Video frames during 2-s cue presentation

B All pixels

>400 features

Deep neural network (3D ResNet)

C

400 features

Machine learning classifier (random forest)

D

Prediction (cross-validated)

Is this cue period
for reward or aversive stimuli?

E

Figure 4–figure supplement 2: Discrimination based on facial expressions.
(A) A frame sequence during the cue presentation was analyzed. (B) All pixel values were inputted to a pretrained deep neural network, resulting in more than 400 features. (C) The top 400 features were fed into a machine learning classifier (random forest). (D) The recall rate of the classifier was computed as the discrimination rate. (E) The time course of discrimination was computed (one example animal). High discrimination was achieved only in the late phase of classical conditioning, even though the early phase was analyzed based on its own classifier trained with frames from the early phase.