Harnessing Agent-Based Modeling in CellAgentChat to Unravel Cell-Cell Interactions from Single-Cell Data

Vishvak Raghavan[1,2,3], Yue Li[1,3*], and Jun Ding[1,2,3*]

[1]School of Computer Science, McGill University, Montreal, QC, Canada
[2]Meakins-Christie Laboratories, Translational Research in Respiratory Diseases Program, Research Institute of the McGill University Health Centre, Montreal, QC, Canada
[3]Mila - Quebec Artificial Intelligence Institute, Montreal, QC, Canada

*Corresponding authors, YL (yueli@cs.mcgill.ca); JD (jun.ding@mcgill.ca)

ABSTRACT

Understanding cell-cell interactions (CCIs) is essential yet challenging due to the inherent intricacy and diversity of cellular dynamics. Existing approaches often analyze global patterns of CCIs using statistical frameworks, missing the nuances of individual cell behavior due to their focus on aggregate data. This makes them insensitive in complex environments where the detailed dynamics of cell interactions matter. We introduce CellAgentChat, an agent-based model (ABM) designed to decipher CCIs from single-cell RNA sequencing and spatial transcriptomics data. This approach models biological systems as collections of autonomous agents governed by biologically inspired principles and rules. Validated against seven diverse single-cell datasets, CellAgentChat demonstrates its effectiveness in detecting intricate signaling events across different cell populations. Moreover, CellAgentChat offers the ability to generate animated visualizations of single-cell interactions and provides flexibility in modifying agent behavior rules, facilitating thorough exploration of both close and distant cellular communications. Furthermore, CellAgentChat leverages ABM features to enable intuitive in silico perturbations via agent rule modifications, pioneering new avenues for innovative intervention strategies. This ABM method empowers an in-depth understanding of cellular signaling interactions across various biological contexts, thereby enhancing in-silico studies for cellular communication-based therapies.

Keywords:

Cell-cell interaction, Agent-based model, Single-cell, Single-cell RNA-sequencing, Spatial transcriptomics data, Cellular dynamics, In silico perturbation.

INTRODUCTION

Cells, as the fundamental units of most life forms, are dynamic and continuously adapt their states in response to changes in their microenvironment[1,2]. In multicellular organisms, cellular activities rely on the coordination of cell-cell interactions (CCIs) across different cell types and tissues[3-6]. CCIs are critical in numerous biological processes, including cell differentiation[7], disease pathogenesis[8], and immune responses[9]. Signaling molecules, such as ligands and receptors, facilitate cellular interactions and influence cell behavior, leading to dynamic changes in cellular functions[3]. By studying these interactions, we can gain insights into the mechanisms underlying biological processes, such as organ development and tumor growth, ultimately informing the study of various disorders and diseases and aiding in the identification of potential diagnostic and therapeutic strategies[9].

However, inferring CCIs in biological processes is challenging due to cellular heterogeneity in most biological systems[10]. The advent of single-cell sequencing, particularly single-cell RNA-sequencing (scRNA-seq) and spatial transcriptomics measurements like 10x genomics Visium[11], Slide-Seq[12], MERFISH[13], SeqFISH+[14], Stereo-seq[15] and Xenium[16] offers unparalleled opportunities to investigate CCIs in diverse biological processes. Numerous tools have been developed in recent years for CCI inference by integrating known ligand-receptor (LR) pairs with gene expression data from scRNA-seq data, including state-of-the-art methods such as CellPhoneDB[17] and CellChat[18]. Despite advancements, earlier methods and versions primarily rely on unimodal scRNA-seq data, which restricts their capacity to infer CCIs due to the absence of spatial distance information between cells[17-19]. As CCIs in tissue environments are heavily influenced by spatial structures and can occur at both long and short ranges[20], integrating spatial information is essential for studying endocrine and paracrine signaling, respectively. Thus, several recent methods, such as COMMOT[21], NICHES[22] and CellChat v2[23], have been developed for inferring cellular interactions from single-cell spatial measurements.

These existing methods still face significant drawbacks regardless of whether they integrate spatial information or not. Specifically, existing methods such as CellPhoneDB and CellChat depend on mean expression values computed from single-cell clusters and therefore focus on cell population levels, inferring CCIs between cell populations (clusters). Such approaches fail to fully exploit the single-cell resolution provided by the original measurements, resulting in a loss of the intricate signaling patterns that exists between cells. NICHES, COMMOT, and Scriabin[24] are newer methods for analyzing LR interactions at the single-cell level. COMMOT applies collective optimal transport with spatial data to infer CCIs yet cannot use scRNA-seq data alone. Conversely, Scriabin assesses CCIs using solely scRNA-seq data, lacking spatial insights. Additionally, no method yet models both long-range and short-range interactions effectively. These limitations hamper the accurate discovery of CCIs in various biological processes[25]. Finally, CCIs play crucial roles in the pathogenesis of many diseases[8], and the analysis of these interactions holds the potential for developing effective therapeutics. However, existing methods like COMMOT and scSeqComm[26], despite their ability to capture the effects of CCIs on downstream genes, do not support in-silico perturbation to utilize these impacts for exploring potential treatments, such as receptor antagonists.

Agent-based model (ABM) simulates the actions and interactions of autonomous agents, through which it helps understand the behaviour and outcomes of a complex system[27]. In ABMs, agents are typically individuals or entities endowed with specific characteristics, behaviors, and rules. These agents can interact with each other and their environment, often leading to emergent

behaviors and patterns at the macroscopic level. Therefore, ABM allows for a straightforward yet
effective depiction of cellular interactions, embracing the concept of emergence. In the context of
this study, this translates to intricate CCI patterns evolving from basic, local rules applied to
individual cells' interaction behaviors. This ABM methodology starkly diverges from existing top-
down methods described above that infer interactions through global statistical correlations, often
overlooking localized cellular dynamics and individual properties. Although ABMs hold great
potential, to our knowledge, they have not been fully utilized for inferring CCIs from single-cell
data. Due to their unique characteristics, ABMs are well-suited for exploring complex CCIs and
their impact on spatial and temporal cellular dynamics in biological systems. First, ABMs capture
emergent phenomena arising from collective agent interactions, revealing insights into cellular
community function[27,28]. Second, they represent heterogeneous cells as agents with different
statuses, enabling accurate modeling of diverse cell populations and capturing variations in cell
types, states, and behaviors critical for understanding biological processes[27,28]. Third, ABMs, like
many machine learning approaches, offer explicit integration of temporal and spatial information,
a crucial capability for studying complex biological processes such as tissue development, immune
responses, and tumor growth[27,28]. Lastly, ABMs facilitate the straightforward and effective
adjustment of agent rules, enabling perturbations in agent behavior. This allows for predictions of
alterations in cellular interactions and facilitates the search for potential therapeutics, like receptor
antagonists[29], based on modified cell agent behavior rules, such as receptor blocking.

In this study, we introduce CellAgentChat, a comprehensive ABM framework for inferring and
visualizing CCIs from scRNA-seq and spatial transcriptomics data. Leveraging the inherent
advantages of ABMs, CellAgentChat offers versatile functionalities for CCI inference and
visualization. By adopting an agent-based perspective, CellAgentChat seamlessly integrates
spatial and biological data, providing a more precise and comprehensive understanding of cellular
interaction dynamics at both individual cell and system levels. Moreover, CellAgentChat enables
in silico perturbations and in-depth analysis of the downstream effects of cellular interactions on
gene expression and facilitates the identification of potential therapeutic receptor targets. Notably,
the flexible framework of CellAgentChat allows for diverse functionalities by adjusting the
behavior rules associated with cell agents. This feature enables the identification and modeling of
long-range and short-range LR interactions. Finally, CellAgentChat has the capability to analyze
temporal dynamics, shedding light on how CCIs influence cellular behavior over short time
frames. Our evaluation of CellAgentChat, using several publicly available single-cell sequencing
and spatial transcriptomics datasets, reveal superior CCI inference accuracy compared to other
benchmarked methods. Together, we introduce a new paradigm of framework based on ABMs for
future advancements in the study of cellular interactions implicated in complex biological
processes.

RESULTS

Overview of CellAgentChat

CellAgentChat constitutes a comprehensive ABM framework integrating gene expression data and
existing knowledge of signaling LR interactions to compute the probabilities of CCI. Figure 1
schematically outlines the CellAgentChat methodology. Utilizing the principles of ABM, we
characterize each cell agent through various cellular attributes (states), including cell identities
(e.g., cell type or clusters), gene expression profiles, LR universe and spatial coordinates (Fig. 1a).

We quantify cellular interactions between sender and receiver cells based on the number of ligands
secreted by the sender cells and subsequently received by the receiver cells. This process hinges
upon three interrelated components: ligand diffusion rate (γ^l), receptor receiving rate (α^r), and
receptor conversion rate (β^r) (Fig. 1b, c, d).

133
Figure 1: Overview of CellAgentChat. a, CellAgentChat requires the following agent states for each cell: gene
expression derived from scRNA-seq data, its cell type/cluster information, and the ligand-receptor database (by default
uses CellTalkDB). Optionally, users can provide spatial coordinates of each agent (from spatial transcriptomics data).
b, The receptor receiving rate is the probability that an interaction is received by a receptor, calculated by non-

parametric min-max normalization of the receptor expression. **c,** The inputs of our deep learning model are the ligand
and receptor's gene expression, and the outputs are the gene expression of all genes. The model incorporates prior
knowledge of TF-receptor and TF-gene interactions from pre-existing databases to create a partially connected
feedforward network. For a receptor, we compute the perturbed expression by permuting the feature variables of each
receptor and scaling down the expression, thereby simulating receptor blocking. The receptor conversion $\boldsymbol{\beta^r}$ of the
receptor is estimated from the total change of the predicted expression after the perturbation. **d,** Calculating the ligand-
receptor score between two cells for a specific ligand-receptor pair requires the ligand diffusion rate γ^l, the receptor
receiving rate α^r, and the receptor conversion rate β^r. If spatial data is provided, the ligand diffusion rate γ^l
incorporates the Euclidean distance d between the two cells. The τ hyperparameter controls the degrees of spatial
freedom for the distance (default $\tau =2$). The δ hyperparameter (default $\delta=1$) controls the decay rate of ligands and can
be used to prioritize long-range ($0 < \delta < 1$) or short-range ($1 < \delta < 10$) interactions. x^l is the normalized ligand
expression. Using the LR score, we can calculate the interaction score between two clusters/cell types and the cell
receiving score (CRS) for a cell. After a permutation test, significant interactions are identified and used to calculate
the cell type pair score (CTPS) between two clusters. **e,** CellAgentChat enables agent-based animation, visualizing
cell type communication, finding significant LR pairs, identifying long-range and short-range interactions and in
silico-permutations to simulate blocking receptors to identify the most perturbed downstream genes to discover
therapeutic drugs.

The initial step in our approach involves the estimation of the receptor receiving rate (α^r), which
signifies the fraction of secreted ligands received by the receptor on the receiver cell (Fig. 1b). We
then leverage a neural network to evaluate the receptor conversion rate (β^r), which measures the
potency of a cellular interaction by studying its effect on downstream gene expression dynamics
(Fig. 1c). Considering the possibility that a similar interaction strength between LR pairs can yield
diverse impacts on downstream gene expression dynamics[3], it becomes crucial to assess the
conversion rate for each receptor. Next, our approach involves the computation of the ligand
diffusion rate ($\gamma^l = x^l/d^{\tau \times \delta}$), which quantifies the ligands secreted by the sender cell that reach
the target receiver cell (Fig. 1d; Eq 1 in Methods). This rate is a function of both ligand expression,
x^l and intercellular distance, d. The ligand diffusion rate also governed by two parameters: τ and
δ. τ represents the degrees of spatial freedom concerning the single-cell data, typically set to two
for spatial transcriptomics data that is derived from two-dimensional slices. δ signifies the decay
rate of ligand diffusion which prioritizes interactions over long or short distances. When δ is lower
than 1, it weakens the decay rate, giving precedence to long-range interactions. Conversely, when
δ exceeds 1, it amplifies the decay rate, favoring short-range interactions. A $\delta = 1$ (default) equally
prioritizes long and short-range interactions (no preference). Consequently, adjusting the δ
parameter can aid in identifying LR pairs more inclined to interact over long-range or short-range
distances. This could also identify proper δ rates for different ligands. We quantify the interaction
between a specific LR pair from sender and receiver cell agents using LR scores, which depend
on the above rates (Fig. 1d; Eq 4). The LR score for two cluster pairs is the Interaction Score (IS)
(Fig. 1d; Eq 7). We consider an IS for an LR pair significant if its IS score significantly surpasses
the background score derived from a permutation test. The cumulative interaction between two
cell clusters is computed as the total number of significant LR pairs (Eq 8). CellAgentChat also
computes the cell receiving score (CRS) (Eq 5) to measure the received interactions for each
individual cell. Together, CellAgentChat offers a suite of downstream functionalities (Fig. 1e),
several of which are distinctive and not available in existing methodologies.

CellAgentChat identifies signaling pathways involved in mouse hippocampus development
using high resolution spatial transcriptomics data
We first studied CCI in in mouse hippocampus development using time-series single-cell spatial
transcriptomics data obtained from the Stereo-seq platform to accurately define the cellular
transcriptomics and cellular composition in mouse neurogenesis[15]. Stereo-seq attains a nanoscale

resolution of 500 nm, facilitating precise spatial pinpointing and characterization of individual
cells[15]. We aimed to identify the key signaling pathways involved in the developmental lineage
between cells at embryonic day E12.5, E14.5 and E16.5, which respectively contained 3,671 cells,
3,648 cells and 19,419 cells across 3 slices, from a total of 11 cell type populations (Fig. 2a).

CellAgentChat calculates the interaction score based on the spatial coordinates of cells and the
distance between individual cells when spatial information is provided (Fig. 1). When considering
the spatial coordinates of cells, CellAgentChat identified 1,531 significant interactions involving
579 LR pairs at E12.5 (Fig. 2b, c). Microglial (Micro), Fibroblasts (Fibro), Erythrocytes (Ery),
Endothelial cells (Endo) and Glioblasts (GlioB) were involved in the greatest number of
interactions (Fig. 2b). The well-established description of Micro and other vascular cells' role in
promoting and regulating neurogenesis is widely acknowledged[30-32]. Micro also contributes to
regulating the size of the neural precursor pool, which is crucial in the process of neurogenesis[30].
Upon examining the LR pairs with the highest involvement as inferred by CellAgentChat, it
becomes apparent that the *Apoe* signaling pathway played a significant role (Fig. 2c). Interestingly,
the top four LR pairs all centered around the *Apoe* ligand, with *Apoe-Lrp1* ranking at the forefront.
The potential regulatory role of *Apoe* in mouse neurogenesis has been previously documented,
including its involvement in Micro expression to mitigate neurodegenerative diseases[33,34]. Further,
we see heavy collagen signaling (Fig. 2c). *Col1a1*, *Col1a2* and *Col8a1* are all commonly expressed
by Fibro to enhance hippocampal neurogenesis in mice[35,36]. We also see *Pf4* signaling (Fig. 2c),
which has also been shown to enhance hippocampal neurogenesis in mice[37].

At E14.5 and E16.5, we observe a continuation of the similar trends seen at E12.5. There is
consistently strong interaction among Micro and vascular cells such as Ery, Endo, and Fibro (Fig.
2d, f), highlighting their crucial role throughout neurogenesis[30-32]. Additionally, there remains
significant *Apoe* signaling within Micro and collagen signaling originating from Fibro (Fig. 2e, g).
However, notable changes emerge at E14.5. Firstly, there is increased signaling within glioblasts
(GlioB) and neuroblast cells (NeuB) (Fig. 2d, e). Beginning at E14, radial glial cells (RGCs)
produce intermediate progenitor cells like GlioB and NeuB before these cells terminally
differentiate into specific neuronal types[38,39]. This phenomenon persists until E18, explaining the
continued trend observed at E16.5 (Fig. 2f, g). Furthermore, there is an increase in interactions
involving RGCs at E16.5 (Fig. 2f, g), coinciding with their transition from producing neurons to
generating astrocytes and later oligodendrocytes from E16 until birth[38,39].

The data presented in Fig. 2f illustrates the communication patterns at stage E16.5, combining
three spatial sections into a cohesive analysis. We also analyzed the results inferred by
CellAgentChat for each of the three slices at E16.5 individually (Supplementary Fig. 1). Across
all three slices, we observe remarkably consistent communicative patterns, mirroring those seen
in the combined analysis presented in Figure 2f. This consistency underscores the robustness of
the tool, CellAgentChat. Within each of the three slices, robust signaling is evident among Micro,
Ery, Endo, and Fibro cells, consistent with previous discussions (Supplementary Fig. 1a, b, c).
Additionally, we observe strong correlations among these communication patterns across the slices
(Supplementary Fig. 1d).

Since spatial information is not always provided, to showcases its broad applicability, we also
applied CellAgentChat to the Stereo-seq mouse hippocampus development without using the
spatial information. This led to consistent findings to those obtained from the spatial

transcriptomics analysis including the crucial role of Micro and vascular cells and the increased
levels of collagen and *Apoe* signaling to regulate neurogenesis (Supplementary Fig. 2).

Figure 2: Analysis of cell communications in developing mouse hippocampus. a, UMAP visualization of scRNA-seq data across time points. (**b, d, f,)** Heatmaps display the communication network inferred by CellAgentChat at E12.5, E14.5 and E16.5, respectively. (**c, e, g,)** Dot plots illustrate the top 25 LR pairs between cell types inferred by CellAgentChat at E12.5, E14.5 and E16.5, respectively.

To systematically assess the reliability of the LR interaction pairs predicted by CellAgentChat, we further employed Reactome pathway enrichment analysis. We compared the enrichment P-values of these pathways' terms using the significant LR pairs with high scores (**Methods**). Many pathways associated with the LRs predicted by CellAgentChat are related to neurogenesis and are present in all time points and scenarios (spatial and non-spatial). These pathways included axon guidance, nervous system development and developmental biology. One of the most significant Reactome pathway in all time points was extracellular matrix (ECM) organization (Supplementary Fig. 3). Additional pathways involving ECM interactions had high adjusted P-values, indicating that these interactions play a role in neurogenesis (Supplementary Fig. 3). Studies have shown that the ECM influences and regulates many aspects of neurogenesis in mouse and other species[40,41]. Pathways involving collagen were also present (Supplementary Fig. 3), including collagen degradation which supports the collagen signaling discussed earlier. Collagen is the most abundant component in the ECM and is broken down with the help of Micro cells expressing *Mmp9* to enhance neurogenesis[42]. Finally, we see *EPH-Ephrin* signaling (Supplementary Fig. 3), which regulates cell migrating neurons in nervous system development[43].

Together, we showcased the utility of CellAgentChat for inferring CCIs with high-resolution spatial transcriptomics data by recapitulating the key time-dependent signaling events during the developmental process of neurogenesis in mice. Moreover, its consistent performance across various samples, including different time-points and technical variations, speaks to its reliability and robustness. Additionally, CellAgentChat can seamlessly integrate technical samples, such as different spatial slices, into a unified analysis framework, thus facilitating a comprehensive understanding of complex cellular interactions and signaling dynamics in spatially resolved datasets. CellAgentChat demonstrates versatility in dealing with scRNA-seq datasets with and without spatial information.

CellAgentChat finds signaling pathways in spatial transcriptomics data at multi-cell/spot resolution

To showcase the broad applicability of CellAgentChat, we further explored its utility in signaling analysis using Visium spatial transcriptomics data, where each spatial spot contains multiple cells. This approach aimed to demonstrate that CellAgentChat can efficiently operate at the spot-level in single-cell spatial datasets, which remain commonly adopted for single-cell spatial studies. In this application, we term spots as agents in the ABM. We applied CellAgentChat to a dataset of Squamous Cell Carcinomas (SCC), which comprises 3,295 spots from a total of thirteen distinct cell populations, corresponding to three normal keratinocyte (NKer) clusters, two tumor keratinocyte (TKer) clusters, and three myeloid dendritic cell (MDC) clusters (Fig. 3a). Additionally, the dataset includes several other cell types, such as endothelial (Endo), fibroblast (Fib), T-cells, B-cells, and melanocytes (Mel). When considering the spatial coordinates of spots, CellAgentChat identified 7,973 significant interactions involving 1521 LR pairs (Fig. 3b, c). Only a few cell types were involved in these interactions. NKer3, MDC3, T cells, and TKer2 participated in the most interactions (Fig. 3b). Furthermore, CellAgentChat identified that three out of the four LR pairs engaged in the most interactions prominently featured *CD44* as the receptor, with *PKM-*

CD44 emerging as the top pair (Fig. 3c). This discovery aligns with previous studies demonstrating
the pivotal role of this LR interaction in the progression of SCC[44]. Additionally, the *MMP1-CD44*
pair was involved in numerous interactions, with *MMP1* highly expressed in TKer subpopulations
and *CD44* overexpressed in cancer cells (Fig. 3c). These findings align with previous studies
linking *MMP1* expression to SCC, highlighting the significance of *CD44* in cancer cells[45,46]. The
LR pair *HLA-B-CANX* also stood out as a leading LR interaction, notably elevated in T cells and
TKer2 cells (Fig. 3c). This compelling association has been demonstrated to be linked with SCC[47-
49].

287
288
289 **Figure 3: Analysis of Cell-Cell Communications in Squamous Cell Carcinoma (SCC) using CellAgentChat. a,**
290 UMAP visualization of spot resolution single-cell transcriptomics data across time points. **b,** Heatmap displays the
291 communication network between cell types, inferred by CellAgentChat. **c,** Dot plot illustrates the top 25 LR pairs
292 involved in the communication between cell types inferred by CellAgentChat. **d,** Reactome pathways analysis using
293 the ligand-receptor interactions inferred by CellAgentChat. Pathways highlighted in red are linked with SCC.

294
Additionally, we employed Reactome pathway enrichment analysis to assess the reliability of the
LR interaction pairs predicted by CellAgentChat. Many pathways associated with the LRs
predicted by CellAgentChat are relevant to the disease of interest. The most significant Reactome
pathway was ECM organization (Fig. 3d). Additional pathways involving ECM interactions had

high -log[10] adjusted P-values, indicating that these interactions play a role in SCC. Studies have
shown that the ECM regulates cell behavior and tumor progression in SCC[50]. The immune system
pathway was also observed (Fig. 3d). Cancer patients often experience an elevated immune
response, with T-cells playing a critical role in recognizing and responding to tumor antigens in all
types of cancers, including SCC[51]. MET signaling pathways were also discovered (Fig. 3d). MET
signaling pathway regulates a range of downstream processes, including proliferation and
invasiveness of tumor cells, and researchers have associated it with poor prognosis in SCC[52,53].
Lastly, interaction pathways involving Laminin were discovered (Fig. 3d). Laminins constitute
integral components of ECM basement membranes, directly governing essential biological
processes linked to morphogenesis, including proliferation, adhesion, migration, angiogenesis, and
cell survival[54]. Deviations in laminin expression patterns and functionality are implicated in
pathological conditions, such as carcinogenesis, due to their ability to modulate critical cellular
processes, thereby influencing various cellular behaviors[54-56]. Furthermore, numerous studies have
indicated that alterations in laminin expression patterns and activity within tumor tissues, including
SCC, correlate with patient outcomes, such as tumor invasiveness and prognosis[54-56]. The inclusion
of this pathway is supported by the identification of *LAMC2-ITGB6* as a key LR interaction
involving TKer2 cells (Fig. 3c).

Effective in silico receptor blocking by CellAgentChat for analyzing perturbed downstream
genes and potential therapeutics

We harnessed CellAgentChat to build upon previous research on breast cancer exploring the
Xenium human breast cancer dataset[16], aiming to identify significant receptors related to the
disease and cultivate innovative therapeutic approaches. The first step of our process entailed using
CellAgentChat to pinpoint 13 key receptors involved in the top 25 LR interactions for the cellular
communications in the dataset (Supplementary Fig. 4d, 5a). We then simulated the blocking of
these receptors individually, akin to the actions of receptor inhibitor drugs. We deployed a
feedforward neural network that was designed to calculate each receptor's conversion rate and was
trained to predict the expression of all genes using the ligand and receptor expression as inputs.
The network connections were constrained to the known LR interactions, LR-TF, and TF-genes
along the forward pass, respectively. (Fig. 4a). Using the trained network, we predicted the
potential impact of receptor blocking on downstream gene expression and identified the most
perturbed downstream genes for each in silico blocked receptor. We subsequently assessed these
perturbed genes using DisGeNET[57] enrichment analysis to evaluate the potential therapeutic
implications of blocking specific receptors (Fig. 4a). Our analysis with CellAgentChat indicated
that blocking 9 of the 13 identified receptors—specifically EGFR, PDCD1, PECAM1, SDC4,
CXCR4, AVPR1A, CTLA4, CD80, and CD93 (hereafter referred to as candidate receptors)—
significantly disrupts genes associated with breast cancer as listed in DisGeNET (Fig. 4b). Further
investigation showed that the blockade of four specific receptors—PECAM1, CXCR4, CTLA4,
and CD80—also resulted in substantial expression changes of downstream genes that significantly
overlapped with breast carcinoma target genes in DisGeNET (binomial test; $P < 0.05$)

In an ablation study to highlight the benefits of integrating biological priors, we tested a fully
connected feedforward network without the LR-TF and TF-target prior constraints that were used
to define connectivity in our primary analysis. This approach aimed to demonstrate the impact of
our initial incorporation of biological insights. The removal of these biological constraints resulted
in a noticeable decrease in specificity: four receptors previously identified as nonsignificant were
now falsely marked as significant (Supplementary Fig. 5b).

The cell receiving score (CRS) of each cell, as identified from our animation platform, reveals that
tumor cells, tumor-associated stromal cells, endothelial cells, and macrophages demonstrate the
highest degrees of involvement in interactions with the candidate receptors (Supplementary Fig.
6). We have also observed significant interactions among these populations, bolstering this
conclusion (Supplementary Fig. 4c). The substantial engagement of tumorous and immune cells
in interactions with the candidate receptors further validates the precision of our cellular interaction
analysis and underscores the potential effectiveness of receptor inhibitor drugs that target these
critical cellular interactions. Following the blocking of receptors, interactions among these cell
types will be entirely inhibited, resulting in a significant alteration of the cells' interaction profile.

Although studies have associated all of the candidate receptors examined in this study with breast
cancer, the candidate receptors we identified—*EGFR*, *PDCD1* (*PD-1*), and *CTLA4*—appear to
play a particularly crucial role in breast cancer development[58-60] (Fig. 4c-e). Furthermore,
therapeutic treatments targeting the three receptors in breast cancer have been previously
documented. Both Lapatinib and Pembrolizumab are FDA approved drugs designed to specifically
target and disrupt pathways involving *EGFR* and *PD-1*. Ipilimumab and tremelimumab are
approved by the FDA to suppress the role of *CTLA4* in melanoma[60]. Although ipilimumab and
tremelimumab are not FDA-approved for breast cancer treatment, studies have shown their
positive impact in this context[60]. This corroborates the potential application of CellAgentChat to
identify receptor-mediated therapeutics. Moreover, blocking the *EGFR*, *PDCD1* (*PD-1*), and
CTLA4 receptors in silico perturbs the disease genes associated with breast cancer (from
DisGeNET) (Fig. 4c-e). In particular, when *EGFR* was blocked, *TPD52* emerged as the fifth most
affected gene, previously linked with breast cancer[61] (Fig. 4c). Similarly, *CDH1*, another gene
associated with breast cancer[62], was the fourth most affected gene when *PDCD1* (*PD-1*) was
blocked (Fig. 4d), and *EPCAM*[63] was the third most affected gene when *CTLA4* was blocked (Fig.
4e). Numerous perturbed genes notably influenced immune cell types like mast cells and
macrophages (Fig. 4c-e; left). This observation aligns with the context of breast cancer, where
immune cells play a crucial role in the tumor microenvironment and the immune response against
cancer cells. All other breast cancer disease target genes perturbed by blocking the candidate
receptors are highlighted in red in (Fig. 4c-e, Supplementary Fig. 5c-e, 7a-c).

Finally, to validate the accuracy of the candidate drug targets captured by CellAgentChat, we
conducted a survival analysis of breast cancer patients over time, comparing populations with high
and low expression levels of each candidate receptor. The analysis revealed statistically significant
differences in survival probabilities over 150 months for patients with varying levels of expression
of *EGFR*, *PDCD1* (*PD-1*), and *CTLA4* (Kaplan-Meier log-rank test; $P = 1.3 \times 10^{-10}$, $P = 3.4 \times 10^{-8}$,
$P = 1.6 \times 10^{-9}$, respectively) (Fig. 4c-e). Additionally, survival analysis of all other candidate
receptors exhibited statistically significant differences in survival probabilities between the two
patient groups (Supplementary Fig. 5c-e, 7a-c). These findings indicate that the identified
candidate receptors would serve as suitable therapeutic targets for treating breast cancer.

Figure 4: Identification of key downstream genes related to breast cancer after in silico blockage of top interacting receptors. a, Schema of the in-silico perturbation analysis using our neural network to block receptors and identify the most perturbed downstream genes linked with breast cancer. **b,** Enrichment for the breast cancer and breast carcinoma gene sets of the significantly perturbed genes based on the in-silico perturbation of the selected 13

receptors. The enrichment is represented as the -log$_{10}$ adjusted P-values (binomial test; FDR corrected-Bonferroni
Correction). *EGFR*, *PDCD1*, *PECAM1*, *SDC4*, *CXCR4*, *AVPR1A*, *CTLA4*, *CD80* and *CD93*, highlighted in red, have
significant -log$_{10}$ adjusted P-values for the breast cancer gene set enrichment. (**c, d, e,**) Matrix plots display the mean
change in expression of each target gene for each cell type for, *EGFR*, *PDCD1* (*PD-1*) and *CTLA4*, respectively (left).
Identification of the top 50 downstream genes with the highest percent change in absolute expression upon blocking
EGFR, *PDCD1* (*PD-1*) and *CTLA4*, respectively (middle). A red highlight indicates genes associated with breast
cancer. Survival analysis displaying the probability of survival of high vs. low receptor expression groups for *EGFR*,
PDCD1 (*PD-1*) and *CTLA4*, respectively (right).
395

396 In addition to using CellAgentChat to perform in-silico perturbation analysis on high-resolution
397 spatial transcriptomics data of a human breast cancer dataset, we also demonstrate the in-silico
398 perturbation capabilities of our model on a spot-level resolution pancreatic ductal adenocarcinoma
399 (PDAC) dataset, where we identified *NOTCH3*, *PLAUR*, *ITGA11* and *CD74* as potential
400 therapeutic targets (see Supplementary Text; Supplementary Figs. 8 and 9).

401 CellAgentChat facilitates the agent-based visualization of cellular interactions associated
402 with individual cells
403
We employed an ABM approach through CellAgentChat, allowing us to examine individual cell
interactions rather than solely focusing on cell populations. We applied the method to a spot-level
human pancreatic ductal adenocarcinoma (PDAC) dataset[64]. After preprocessing the PDAC
dataset, we discerned six distinct cell populations: ductal (Duct), three subpopulations of acinar
(Acin1, Acin2, and Acin3), stromal (Strom) cells, and two classes of cancer (Can1 and Can2)
(Supplementary Fig. 4b, e, f). To effectively exhibit the insights that CellAgentChat can provide
into individual cell interactions, we identified twenty most receiving spots (group A, red circles)
based on the cell receiving score (CRS) (**Methods**) and the least receiving spots (group B, black
circles) with the lowest CRS within the second cancer cell class (green cells) (Fig. 5a).

The spots with higher CRS (more interactive) generally reside in the outer layer of the tissue
(invasive front), consistent with the discovery from many existing studies[65]. These two cell groups
of distinct interacting patterns within the same cluster also demonstrated significant transcriptomic
differences. Group A showed higher mean expression levels of receptors involved in the top 25 LR
pairs than Group B (Fig. 5b). For example, *CD63* was highly expressed in both groups, with group
A having the highest expression levels (Fig. 5b). Additionally, receptors such as *LRP1*, *SDC1*,
ITGA3, *NOTCH3*, and *CD74* displayed notable differences in mean expression between the two
groups, with group A again exhibiting higher expression (Fig. 5b). Next, we systematically
explored the differentially expressed (DE) genes of these two groups. To identify the DE genes for
each group (A and B), we compare the spots within the group with the rest of the spots in the
dataset. The differential genes associated with group A show significantly higher expression in
group A compared with group B (Wilcoxon one-sided test; P =1.22x10^{-48}) (Fig. 5c). In contrast,
the expression of group B DE genes is relatively similar across groups (Fig. 5c).

We showed the top 50 DE genes for both groups by comparing the expression of spots in groups
A and B (Fig. 5d). While both groups showed high expression levels of *S100A6* and *ACTB*, group
B cells exhibited elevated expression of *KRT19* (Fig. 5d). Conversely, *COL1A1*, responsible for
type I collagen production, was particularly prominent in group A cells (Fig. 5d). Notably,
COL1A1, a gene found enriched in PDAC tumor cells, is associated with tumor growth and
metastasis in cells[66].
432

Figure 5: Application of CellAgentChat to detect variations in intercellular communication among individual

cancer cells in PDAC. a, CellAgentChat's animation platform enables the detection of communication strength
variations in individual cancer cells. The platform highlights the cell receiving score (CRS) (**Methods**) of cancer
subpopulation one cells (colored green). It identifies the top 20 cells with the highest CRS (group A) marked by red
circles and the bottom 20 cells with the lowest CRS (group B) marked by black circles. **b,** Expression analysis of
receptors involved in the top 25 ligand-receptor interactions reveals that Group A, consisting of the top 20 spots with
the highest CRS identified by CellAgentChat's animation platform, shows significantly higher expression of key
receptors compared to Group B, consisting of the bottom 20 receiving cancer spots (Wilcoxon test one-sided; $P <$
0.05). **c,** Mean expression of differentially expressed (DE) genes up-regulated for group A spots relative to rest of the
spots (top) and up-regulated for group B cells relative to rest of the spots (bottom). **d,** Heatmaps display the
differentially expressed (DE) genes for group A (left) and group B (right). The DE genes of group A cells are mainly
expressed by group A and not by group B (Wilcoxon test one-sided; $P < 0.001$). Both groups express the DE genes of
group B but are more significantly expressed by group A spots (Wilcoxon test one-sided; $P < 0.05$). **e,** The results of
the Reactome pathway analysis using group A DE genes (on the left) and group B DE genes (on the right). Pathways
affected by PDAC are more associated with group A DE genes (Wilcoxon test one-sided; $P = 0.0313$). Red highlights
indicate shared pathways.

To identify the biological functions associated with these groups, we conducted Reactome pathway
enrichment on the top 50 DE genes for both groups (Fig. 5e). Intriguingly, group A was associated
with several pathways related to the extracellular matrix (ECM) and collagen, known to be crucial
in PDAC tumor progression[67]. On the other hand, group B showed less involvement in ECM-
related pathways, aligning with their lower level of interaction compared to group A (Fig. 5e).
Examinations of CCIs revealed that cells of the same cluster/type can display diverse interaction
behaviors and play varying roles in pathogenesis and disease progression.

CellAgentChat can be tuned to identify long and short-range interactions
Our understanding of the specific distances at which LR pairs operate is evolving but remains
incomplete. While some interactions occur closely between cells, others span longer distances,
influenced by factors such as specific ligands and receptors, tissue environment, and physiological
conditions. The ABM employed by CellAgentChat provides nuanced representations of cell agent
behaviors, including the diffusion of signaling ligands. As a result, this facilitates modeling of both
short- and long-range interactions at the single-cell level by adjusting the delta (δ) factor to
manipulate the ligand diffusion rates (**Methods**).

Using CellAgentChat, we embarked on an investigation on the spatial distance tendencies of
cellular communication within the mouse somatosensory cortex leveraging data from SeqFISH+
technology (Supplementary Fig. 10). To this end, we utilized the CellChat database, which
provides known annotations of the distance ranges for each ligand-receptor (LR) pair. LR pairs are
categorized as either "Secreted Signaling" (indicating long-range interactions) or "cell-cell
contact" (indicating short-range interactions). To identify short-range interactions, we used a δ=10,
to constrain interactions to shorter distances and used δ=0.1 to identify long-range interactions. To
validate our results, we compared the top interacting LR pairs with CellChat's annotations. Under
our short-range mode, the majority of our top 10, 15, 20 and 25 LR pairs were annotated as "cell-
cell contact" (short-range) (Fig. 6a). Specifically, 80% of the pairs within the top 15 LR pairs
identified in the short-range mode fell into the "cell-cell contact" category (Fig. 6a). In the long-
range mode, most of the top LR pairs identified were classified as "secreted signaling" (long-
range), with 75% of pairs within the top 15 LR pairs falling into this category (Fig. 6a). Recently,
Liu et al.[68] have developed a method to identify long-range and short-range interactions using
optimal transport (OT). Compared to that approach, CellAgentChat more accurately captured
short-range interactions and identified long-range interactions to a similar degree of accuracy (Fig.
6b).

Figure 6: Application of CellAgentChat to infer long-range and short-range interactions in adult mouse cortex.
a, Bar plot illustrates the percentage of interactions found by CellAgentChat annotated by CellChat's database as either cell-cell contact or secreted signaling with a delta (δ) value set to 10 (top) and 0.1 (bottom). **b,** Bar plot illustrates the percentage of interactions found by CellAgentChat annotated by CellChat's database as either cell-cell contact or secreted signaling in comparison to an optimal transport-based approach[68]. **c,** CellAgentChat's animation platform depicts the cell receiving score (CRS) for the *Sema4a-Plxna4* ligand-receptor pair using the short-range mode (δ =10).

The colour of the cells (circles) represents the cell cluster. This animation depicts the release of *Sema4a* (ligand) from
cells in cluster 9 (represented by turquoise cells). In the animation, cell size corresponds to the cell receiving score
(CRS). Cells closer to cluster 9 cells exhibit higher CRS (within red square) than cells that are further away (within
blue square), which are smaller in size. **d,** Reactome pathway analysis of the receptors involved in the top 25 LR
interactions in the long-range mode (top) and short-range mode (bottom). **e,** Heatmap displaying the cell
communication between clusters after prescribing specific delta values based on identifying key short-range and long-
range interactions.

In our short-range mode, we detected LR pairs involving the *Sema4* pathway, such as *Sema4a-*
Plxna4. Previous research has linked these pairs to axon guidance in the mouse cortex, occurring
over short distances[69]. We further employed our animation platform to visualize the CRS of
individual cells for the *Sema4a-Plxna4* LR pair in the short-range mode originating from cluster 9
cells (turquoise cells) (Fig. 6c). The results highlighted higher CRS amongst cells close in
proximity to the cluster 9 cells represented by larger sizes (within the red box), compared to cells
situated further away (blue box). This observation underscores the capabilities of CellAgentChat
in modeling and identifying short-range interactions.

We conducted pathway enrichment comparisons of the LR interaction pairs predicted by
CellAgentChat for both long- and short-range interactions (Fig. 6d). The genes associated with
short-range interactions showed enrichment in pathways related to cell adhesion and cell-cell
junctions. In contrast, genes associated with long-range interactions were enriched in numerous
signaling pathways known for their wide regulatory range (Fig. 6d).

Ultimately, we leverage our understanding of the above discovered short-range and long-range LR
pairs to refine the ligand diffusion rate with greater precision. Having pinpointed distinct LR pairs
engaging in either long-range or short-range interactions, we assign specific delta values to these
ligands, to control the decay rate between cell i and j in computing the ligand diffusion rate $\gamma_{i,j}^l$
(Eqn. 1). This enables us to more accurately depict and simulate their actual decay rates during the
computation of CCIs. By integrating the recommended decay rates for the most impactful long
and short-acting ligands, we detect both long-range and short-range interactions concurrently (Fig.
6e). This reveals a communication pattern resembling interactions observed solely with scRNA-
seq data (Fig. 6e, Supplementary Fig. 10b), where spatial constraints are absent, thus more
accurately facilitating long-range interactions. Nevertheless, the communication pattern also
continues to exhibit numerous short-range interactions between proximate cells (e.g., cluster 9-
cluster 9 cells, cluster 3-cluster 3 cells) (Fig. 6e).

CellAgentChat empowers dynamic simulations using agent-based modeling
CellAgentChat provides dynamic simulation capabilities to analyze the impact of CCIs on cell
states over short durations. At each time point (time step), CCIs are computed as described earlier,
followed by the prediction of gene expression levels for each cell using our trained neural network
(**Methods**). Due to uncertainties and the influence of various factors that could affect the actual
magnitude of gene expression changes, directly substituting existing gene expression values with
predicted ones is impractical. Instead, we opt for assessing the directional change of each gene—
whether they increase or decrease—which proves to be more robust and easier to predict[70]
(Supplementary Fig. 11a). Genes with significant changes are identified for each cell at each time
step, and their expression is adjusted within the ABM for the next time step (**Methods**)
(Supplementary Fig. 11a). This iterative process is repeated across all time steps, reflecting the
dynamic modulation of gene expression by CCIs.

To validate the accuracy of our dynamic simulations, we determined the correlation between the
pseudotime trajectory of each gene, and its trajectory as established by the dynamic simulation of
CellAgentChat. Across all genes, we observed a correlation between the pseudotime trajectory
and the ABM dynamic trajectory (Supplementary Fig. 11b). Additionally, we examined the
trajectory of highly interacting ligands and receptors, finding strong agreement between the
pseudotime trajectory and the dynamic ABM trajectory for genes like *ITGA6* (receptor) and *MMP1*
(ligand), which showed overall increasing trends over time ($r^2 = 0.93$ and $r^2 = 0.97$, respectively)
(Supplementary Fig. 11c, d). Similarly, both trajectories indicated a decreasing expression trend
for *CALR* (ligand) over time ($r^2 = 0.90$) (Supplementary Fig. 11e). Moreover, CellAgentChat
accurately identified the initial decrease of expression of *CD44* (receptor) followed by a rapid
increase ($r^2 = 0.89$) (Supplementary Fig. 11f). These findings highlight the effectiveness of
CellAgentChat in capturing dynamic cellular changes resulting from CCIs within a time frame
comparable to the pseudotime represented by the cells.

Benchmarking of CellAgentChat with other cell–cell communication inference tools
We undertook a comparative analysis of CellAgentChat against five prominent CCI inference
methods based on both scRNA-seq and spatially resolved transcriptomics data: CellPhoneDB
(v5)[17], CellChat (v2)[18,23], NICHES[22], COMMOT[21] and Scriabin[24]. In our comprehensive
evaluation of methods, we benchmarked them rigorously from several key perspectives. First, due
to the absence of ground truth, we compared the agreement of the cell-cell communication
networks between each benchmarked method with the consensus of all methods. Secondly, we
examined the consistency of CellAgentChat's predictions with spatial information, focusing on
the spatial proximity of stronger interactions. Specifically, we measured the Pearson correlation
between the probability of interaction and the inverse distance between cells, grounding this
analysis on the established understanding that cells are more likely to interact when they are in
closer proximity[3]. Thirdly, we assessed the functional relevance of the predicted LR pairs by
conducting Reactome pathway enrichment analysis. Finally, we measured the overlap of inferred
LR pairs across different methods.

We began by benchmarking across methods using only scRNA-seq datasets to validate the
robustness of our CCI calculation framework before integrating spatial data. We assessed the cell-
cell communication networks generated by CellAgentChat and compared them with the outcomes
of other methods. Due to the absence of a definitive ground-truth, we adopted a
consensus/ensemble approach. This involved normalizing the interaction matrices computed by
each method and averaging them to create an ensemble interaction matrix, serving as our new
consensus reference. Subsequently, we computed the Pearson correlation between the ensemble
interaction matrix and the inferred communication patterns of each method for comparison. Across
all datasets, the results from CellAgentChat demonstrated superiority and consistency with the
communication networks inferred by CellPhoneDB, CellChat, and NICHES, showing a higher
median Pearson Correlation than the other three methods (Fig. 7a (top)). On examining individual
datasets, CellAgentChat consistently outperformed CellPhoneDB and Scriabin in all cases
(Wilcoxon one-sided test; $P < 0.05$) (Supplementary Fig. 12, 13, 14). Moreover, CellAgentChat,
CellChat, and NICHES showcased remarkably consistent and high-performance results across all
evaluated metrics (Supplementary Fig. 12, 13, 14). Upon incorporating spatially resolved
transcriptomics data, CellAgentChat continued to perform with a high degree of consistency
among the state-of-the-art methods. Specifically, CellAgentChat maintained superior performance
compared to CellPhoneDB and also outperformed CellChat (Wilcoxon one-sided test; $P < 0.05$)

(Fig. 7a (bottom), Supplementary Fig. 12, 15, 16). In comparison to NICHES and COMMOT,
CellAgentChat performed comparably, further demonstrating its status among the state-of-the-art
methods.

580
Figure 7: Benchmarking of CellAgentChat with existing state-of-the-art methods. a, Boxplot comparing the
Pearson correlation between inferred cell communication by individual methods and the ensemble of all methods, for
both non-spatial (top) and spatial data (bottom). **b,** Boxplot comparing the Pearson correlation between the inferred
cell communication for each method and the inverse cell-distance matrix (top), across datasets. Bottom plot in panel
shows the average rank across datasets. **c,** The Reactome pathway analysis, performed on receptors sourced from the

top 100 ligand-receptor pairs inferred by each method across various datasets using spatial transcriptomics data,
revealed that the pairs identified by CellAgentChat exhibit a greater level of statistical significance.

To validate that the CCIs inferred by CellAgentChat are spatially informed, we computed the
Pearson correlation between the predicted communication matrix and the cell-cell inverse distance
matrix. Indeed, cells predicted by CellAgentChat as highly interacting were significantly closer in
proximity compared to predictions from other methods (Fig. 7b). We demonstrated that our
spatially informed predictions outperformed those of CellPhoneDB and CellChat (Wilcoxon one-
sided test; $P < 0.05$) (Fig. 7b), showing superior performance across every dataset compared to
CellPhoneDB and in all but one dataset compared to CellChat (Supplementary Fig. 17).
Furthermore, we show comparable performance compared to NICHES and COMMOT (Fig. 7b,
Supplementary Fig. 17). Overall, CellAgentChat and COMMOT had the joint highest average rank
across all datasets (Fig. 7b (bottom)).

After validating the accuracy of CellAgentChat in predicting cell-cell interactions (CCIs) through
consensus and spatial proximity metrics, we proceeded to conduct pathway enrichment analysis to
further confirm the biological relevance of the inferred LR pairs. This step serves as an additional
layer of validation. We conducted computational validations, as previously described, comparing
the enrichment P-values for relevant pathways using ligands and receptors from the top 100
significant LR pairs from each method. The pathway enrichment analysis conducted for the results
using only scRNA-seq data underscored CellAgentChat's superiority over other methods.
CellAgentChat performed better than CellPhoneDB and Scriabin across all datasets
(Supplementary Fig. 18). Additionally, CellAgentChat outperformed CellChat in five out of the
seven datasets and showed comparable performance in the remaining two (Supplementary Fig.
18). When compared to NICHES, CellAgentChat performed better on the PDAC, breast cancer,
and E16.5 mouse hippocampus datasets, while demonstrating comparable performance on the rest
(Supplementary Fig. 18). Upon integrating spatial data, pathway enrichment analysis continued to
highlight CellAgentChat's superiority over CellPhoneDB and CellChat (Fig. 7c). CellAgentChat
continues to perform better than CellPhoneDB in all datasets (Fig. 7c, Supplementary Fig. 18b).
CellAgentChat also demonstrated superior performance to CellChat, NICHES COMMOT in the
majority of datasets (Fig. 7c, Supplementary Fig. 18b). Overall, pathway enrichment analysis
supported the CCI inference precision of CellAgentChat over other benchmarked methods.

Finally, we calculated the overlap in LR pairs among the top 100 identified LR pairs across the six
methods. We observed that the overlap between CellAgentChat and the other methods, particularly
CellChat, NICHES and COMMOT was high (Supplementary Fig. 19). This serves as another
confirmation of our method's performance, indicating that the LR signaling predictions made by
CellAgentChat align with those of other state-of-the-art methods.

Building on the findings from the preceding benchmarking sections, which underscore
CellAgentChat's superior, or at least comparable performance compared to benchmarked methods,
it is crucial to recognize that beyond these achievements, CellAgentChat also introduces distinct
functionalities not available in other tools. Unique to CellAgentChat is its integrated ABM
animation platform, which facilitates the visualization of individual-level cellular interactions and
supports dynamic simulations. Additionally, through easy and flexible adjustment of agent
behavior, CellAgentChat empowers the use of in silico receptor blocking to explore potential
interventions effectively and serves as a valuable tool for identifying both long and short-range
interactions.

DISCUSSION

DISCUSSION
We introduced CellAgentChat, an agent-based model (ABM) capable of elucidating cell-cell
interactions (CCI) from single-cell data, both with and without spatial information. As a novel
approach, CellAgentChat employs individual cell agents guided by simple behavior rules to
investigate the complexity of cellular interactions within the system. In contrast to the existing
methods for predicting CCI, CellAgentChat offers three unique features. Firstly, it offers a
dynamic visualization of cellular interactions related to individual cell agents. Through its cellular
interaction animation, one can explore cellular heterogeneity within a population, often uncovering
insights at the cluster level that are impossible to obtain (Fig. 5). Thus, CellAgentChat provides
both a cluster- and cell-specific view of cellular communication. Secondly, CellAgentChat enables
efficient in silico receptor blocking, facilitated by modulating the agent rules (for instance, receptor
receiving rate), allowing for a systematic assessment of a receptor blocking's impact on
downstream gene expression dynamics. CellAgentChat can propose systematic strategies for
potential therapeutic interventions by investigating disease-related genes.

Using CellAgentChat, we identified key LR pairs known to play crucial roles in neural
development and cancer. We identify the *Apoe* and collagen signaling pathways among microglial
and vascular cells in mouse neurogenesis (Fig. 2). Both pathways have been shown to play critical
roles in neurogenesis[33-36]. Furthermore, we identified key time-dependent signaling pathways such
as radial glial cells (RGCs) role in producing intermediate progenitor cells and neurons.
CellAgentChat also revealed a significant *LAMC2-ITGB6* interaction among tumor keratinocytes
(TKer cells) in squamous cell carcinoma (Fig. 3). This interaction has previously been linked to
critical roles in tumor progression and migration, suggesting its involvement in the early stages of
cancer development[55]. In addition, CellAgentChat confirmed the selection of three receptors,
EGFR, *PDCD1* (*PD-1*) and *CTLA4*, among others for receptor-targeting treatment options in
breast cancer (Fig. 4). Thirdly, researchers can tailor CellAgentChat to prioritize long- or short-
range interactions controlled by the diffusion rate employed in quantifying cellular interactions.
Using this approach, we identified *Sema4a-Plxna4* as a short-range interaction involved in axon-
guidance in the mouse cortex[69] (Fig. 6). Finally, we showed that CellAgentChat surpasses several
state-of-the-art methods across multiple metrics, including the generation of cell-cell
communication networks, alignment with spatial information, Reactome pathway enrichment
analysis, and overlap of ligand-receptor pairs across different methods (Fig. 7).

In future work, we consider the following extensions of CellAgentChat. By representing cells as
individual agents with distinct statuses, ABMs enable precise modeling of heterogeneous cell
populations, effectively capturing the diversity in cell types, states, and behaviors essential for
comprehending biological phenomena[27,28]. Expanding CellAgentChat's capabilities to include cell
movement and even cell death would be natural extensions, enhancing its capacity to simulate
realistic scenarios. Although technically challenging, further improvements by incorporating
ligands, receptors, and other cofactor molecules as agents in addition to cells would allow for a
more comprehensive and detailed simulation of biological systems' complex interactions and
dynamics. By considering these additional components as agents, the model could capture the
intricate interplay between cells and their surrounding molecular environment, providing a more
nuanced understanding of the underlying mechanisms at play. It would also improve the efficiency
of in silico perturbations such as receptor blocking. Given the prevalent issue of low gene coverage
in many single-cell resolution spatial datasets, like those generated with SeqFISH+ and Xenium,
spot-level technology remains a popular choice among researchers. In our framework, an agent

can represent individual cells or spots, adapting for analytical flexibility. While this representation
may overlook potential interactions between multiple cells within the same spot, especially since
many CCIs occur within a 250µm range[71], our approach simplifies the analysis of complex CCIs
in spot-level data and demonstrates superior interaction inference compared to other methods.
Users inclined towards analyzing spot-decomposed cell interactions, rather than staying at the spot
level, can employ tools such as SpatialScope[72] to deconvolve spot-level spatial transcriptomics
data, achieving single-cell resolution. With this enhanced resolution, CellAgentChat can be run to
explore and understand CCI dynamics in greater detail.

METHODS

Ligand-Receptor database

In this study, we used the ligand-receptor (LR) interactions from CellTalkDB[73], a curated database
of almost 3400 human and over 2000 mouse literature-supported interactions, as the default LR
interaction resource for our CellAgentChat method. The interactions from the curated database
specify a comprehensive collection of expert curated LR interactions. Besides the default
CellTalkDB LR interactions, CellAgentChat also allows for customized LR interactions that users
obtain from other sources or manually curate.

Single-cell sequencing data preprocessing

We preprocessed the single-cell RNA-sequencing (scRNA-seq) data using the Scanpy[74] pipeline.
We began by filtering out cells with more than 20% mitochondrial reads, which are indicative of
poor-quality cells[75,76]. We also removed genes not expressed by a minimum of 3 cells. Finally, cell
counts are normalized against the library size and transformed into log space (with the *log1p*
function). As an ablation study, we show that using normalized gene expression values provide
better results than using raw unprocessed counts (Supplementary Fig. 20). The spatial coordinates
of each cell are incorporated into the "anndata" object (in h5ad format) with the gene expression.

Scoring cell-cell interactions with the agent-based model

700 To build the agent-based model (ABM), we need to define the states and behavior of each agent.
701 Agents here represent the cells in the biosystem. The agent states describe the state and features
702 associated with each cell, while the agent rules define how each agent behaves.

Cell Agent State Description

As we are primarily interested in studying the interactions between cells (or clusters), we describe
each cell agent with characteristics and features related to cellular interactions. The remainder of
this section provides the detailed cell agent state description. (1) *Cell Agent Expression*: The
expression value of all genes, including ligands and receptors, is stored for each cell. The gene
expression profile presents the quantitative description of the cell agent state, which is essential to
infer the interaction behaviors. (2) *Cell Agent Spatial Position*: We use spatial transcriptomics data
to record the position of each cell agent. All the cells are placed on the grid at their spatial
coordinate (if provided). Otherwise, CellAgentChat randomly distributes all cells on the grid. (3)
Cell Agent Batch: To utilize multiple samples (biological replicates or technical samples)
concurrently, we designate the batch of each cell to confine interactions solely between cells of the
same batch. (4) *Cell Agent Ligand-Receptor Universe*: Every cell agent has the same LR universe
using interactions curated from CellTalkDB or user-customized LR interactions. Users can modify
the LR Universe by blocking a specific receptor or LR interaction (see Agent-Based Animation of
CCIs). By doing so, they can remove all interactions associated with ligands or receptors (or

specific LR interactions) from the universe of all cells. This unique characteristic of ABMs enables
effective and flexible ligand/receptor perturbations.

<u>Cell Agent Communication Behavior Rules</u>

In this study, the cellular communications between agents (cells) depend on the states of the cells,
which includes the gene expression of the cell agent, the spatial position of the cell agent (optional)
and the LR universe. To be more specific, the cellular interaction strengths (scores) between cell
agents are quantified based on the set of defined rules as discussed below.

(1) Ligand Diffusion Rate ($\gamma_{i,j}^l$)

The diffusion rate $\gamma_{i,j}^l$ estimates the rate of the diffusion of a ligand l in the microenvironment
from sender cell i to receiver cell j. The diffusion rate depends on a few factors. First, the diffusion
rate is proportional to the number of ligands a sender cell secretes into the environment. We can
estimate this from the number of RNA molecules proportional to its gene expression.

Second, the diffusion rate is inversely proportional to the distance (of the receiver agent) from the
sender cell agent. We calculate the Euclidean distance between two cells based on the spatial
coordinates of all the cell agents. The hyperparameter τ, is a dimensionality parameter representing
the degrees of spatial freedom for spatial data. τ is typically set to two for spatial transcriptomics
data derived from 2-dimensional slices. However, with the advancement of technology and the
increasing prevalence of 3-dimensional slices, we can adjust the τ parameter accordingly ($\tau = 3$)[77].
When spatial information is unavailable (e.g., in single-cell RNA-seq data), τ is set to 0 so that the
distance between all cells in the data will degenerate to 1 ($d^{\tau=0} = 1$).

In the absence of precise information about the rates of ligand travel or the operational distance
range of LR pairs, we employ the delta (δ) parameter as an approximation for the distance range
of LR pairs and model the decay rate of ligands. A user-adjustable slider δ (between *0-10*, default
= *1*) is available to control the decay rate of the diffusion along with the increase of distance, which
allows for inferring long-range or short-range interactions. Keeping δ at the default value of 1
provides no bias toward long-range or short-range interactions. We modeled short-range
interactions with a $\delta = 10$ and long-range interactions with $\delta = 0.1$. By investigating various δ
values and the corresponding agent communication behaviors defined by these values, we can also
determine a specific δ for each LR pair. This approach avoids the assumption that all ligands share
the same diffusion rate. The distance between sender cell i and receiver cell j is calculated as the
Euclidean Distance $d(i, j)$. The expression of ligand l in sender cell i is denoted as x_i^l. Then, the
diffusion rate $\gamma_{i,j}^l$ can be calculated as below:

$$\gamma_{i,j}^l = \frac{x_i^l}{d(i,j)^{\tau \times \delta}}$$

751   (1)

(2) Receptor receiving rate (α_j^r)

The receiving rate α_j^r associated with a receptor r represents the probability that an interaction is
received by the receptor in cell j, and it is proportional to the expression of the receptor. The
receptor receiving rate will be a value in the range (0,1) where 0 represents the receptor cannot

"receive" any ligands that reach it, while 1 denotes that reached ligands will all be "received" by
the receptor. Here, we employed the non-parametric min-max normalization to calculate the
receptor receiving rate for each receptor. Min-max normalization has been used in other methods
for cell interaction inference[78-81], lending support to our choice. For each cell, the expression value
of each receptor is subtracted by the minimum expression value in that cell and then divided by
the difference between the maximum and minimum expression values in that cell (min-max
normalization), enabling the identification of highly expressed receptors within each cell, thereby
establishing a cell-specific metric.
764
(3) Receptor transcriptome conversion rate (β^r)

CCIs play a crucial role in the regulation of gene expression, impacting the transcriptome, and
ultimately influencing cellular function and fate. In order to account for the effect of cellular
interaction on downstream gene expression, we employed a neural network regressor to estimate
the downstream gene expression impact associated with each receptor (here we term it as the
receptor conversion rate β^r). The neural network takes as input two vectors, x_j^R, representing
expression of all receptors in cell j and $\frac{1}{N}\sum_{i=1} x_i^L$, representing the average expression of all
ligands across the set of all N cells. These two vectors are then fed into the neural network ($\mathcal{F}$) to
predict the expression $\hat{x}_j$ of all genes in cell j (regression task). The neural network structure
comprises the following components: two input layers, two hidden layers, and one output layer.
The size of the two input layers corresponds to the number of ligands and receptors, respectively.
The first hidden layer is dedicated to representing specific LR pairs. Each node in this layer
receives two connections, one corresponding to a ligand and the other to a receptor, thus
encapsulating the unique LR pair interactions. The second hidden layer represents transcription
factors involved in downstream signaling pathways. We leverage scSeqComm[26], a database that
incorporates insights from KEGG[82] and REACTOME[83] pathways to delineate the association
between a receptor and its downstream transcription factors (TFs), to inform the connections
between this layer and the previous one. The output layer is responsible for outputting downstream
gene expression predictions. To establish connections between known TFs and their downstream
gene targets, we utilize prior knowledge from databases such as TRRUST v2[84], HTRIdb[85], and
RegNetwork. If specific information regarding the target genes of certain TFs is unavailable, we
utilize dense connections. We use a stochastic gradient descent optimizer to train the network $\mathcal{F}$
that minimizes the mean squared error loss L (between the true observed expression for the set of
all cells X and the predicted expression):

789
$$L = \frac{1}{N}\sum_{j=1}\left(x_j - \mathcal{F}\left(x_j^R, \frac{1}{N}\sum_{i=1} x_i^L\right)\right)^2$$

790   (2)

With the trained neural network regressor, we then perform a sensitivity analysis to score each
input receptor based on its impact on the downstream gene expression. Specifically, we conduct a
perturbation-based feature calculation[86], by permuting feature values for each receptor,
independent of the output. Next, we adjust the expression of the permuted receptor to reflect a
~50% reduction, scaling it by the median expression of all receptors. This represents the perturbed

input receptor vector $(x_j^{R\backslash r})$ simulating a ~50% repression of receptor r in expression. While this
50% repression serves as the default setting, we also offer the option to simulate a 100% repression,
equivalent to a complete knockout, catering to users who prefer a full knockout scenario. Then,
we quantify the importance of the receptor r (I_r) based on the loss increase caused by the
perturbation:

802
$$I_r = L_r - L$$
803

804
$$= \frac{1}{N}\sum_{j=1}\left(x_j - \mathcal{F}\left(x_j^{R\backslash r}, \frac{1}{N}\sum_{i=1} x_i^L\right)\right)^2 - \frac{1}{N}\sum_{j=1}\left(x_j - \mathcal{F}\left(x_j^R, \frac{1}{N}\sum_{i=1} x_i^L\right)\right)^2$$

805   (3)

Employing a similar approach as the receiving rate, we normalize the receptor importance into
range (0,1) via min-max normalization. Here, we set a minimum value for the conversion rate to
be 0.4 based on previous studies that measure the effect of cellular interaction on gene
expression[87]. We confirmed that this approach does not significantly alter the resulting gene
expression of the cells of the training grid (Supplementary Fig. 21 and 22).

(4) Cell-cell interaction score between cells

The interaction score between a ligand-receptor pair (l, r) in a sender-receiver pair (i, j) is
calculated based on the rates calculated above $(\gamma^l, \alpha^r, \beta^r)$:

815
$$LR(i_l, j_r) = \gamma_{i,j}^l \times \alpha_j^r \times \beta^r = \frac{x_i^l}{d(i,j)^{\tau \times \delta}} \times \alpha_j^r \times \beta^r$$

816   (4)

817 The cell receiving score for a given cell i is calculated as the summation over all ligand-receptor
818 pairs (l,r) across all sending cells:

819
$$CRS(i) = \sum_{j \in C} \sum_{(l,r) \in (L,R)} LR(j_l, i_r)$$

820   (5)

821 The combined interaction score between a sender-receiver cell pair (i, j) is calculated as the
822 summation over all ligand-receptor pairs (l,r):

823
$$CPS(i,j) = \sum_{(l,r) \in (L,R)} LR(i_l, j_r)$$

824   (6)

The overall interaction score $IS(C1, C2)$ between two clusters C1 (sender) and C2 (receiver) for a
ligand-receptor pair (l,r) is quantified as the average interaction score of all possible sender-
receptor cell agent pairs between the sender C1 and receiver C2:

$$IS(C1_l, C2_r) \;=\; \frac{1}{|C1| \times |C2|} \sum_{i \in C1} \sum_{j \in C2} LR(i_l, j_r)$$

$$= \frac{1}{|C1| \times |C2|} \sum_{i \in C1} \sum_{j \in C2} \frac{x_i^l}{d(i,j)^{\tau \times \delta}} \times \alpha_j^r \times \beta^r$$

(7)

We also introduce a method to optionally integrate the utilization of pseudotime ordering of cells
into the calculation of CCI scores. Please refer to the supplementary text for further details.

Finding significant ligand-receptor pairs between cell populations

Once we get the interaction score for an LR pair between two cell clusters, we will examine
whether such an interaction score is statistically significant (stronger compared to the random
background). Specifically, we employed a random permutation test to calculate the statistical
significance P-value associated with each LR interaction score. Significant interactions are
identified between two cell groups using a permutation test. It randomly permutes the group labels
of cells and then recalculates the interaction score for each LR pair. The LR score is stored after
each permutation regardless of the cell type pairing. We perform enough permutations so that each
LR pair, has on average 10,000 score values. We then obtained the maximum likelihood estimates
of alpha and scale of Gamma distribution from the random scores for each LR pair. Using the
background distribution, we can compute a P-value for a given interaction score (for the specific
LR pair) by evaluating its position in the distribution's right tail. The P-value is then corrected
using FDR (Benjamini/Hochberg) correction. Interaction Scores (IS) with a P-value above the
threshold (0.05) are considered Significant Interaction Scores (SIS). Using the SIS, we can
calculate the cell type pair scores (CTPS) between two clusters, C1 (sender) and C2 (receiver):

$$CTPS(C1, C2) = \sum_{(l,r) \in (L,R)} SIS(C1_l, C2_r)$$

(8)

Due to the computational expense of calculating the distances between all cells in the dataset, the
permutation test can be quite costly when considering spatial information. To address this, we have
devised a method to match the background gamma distribution of non-spatial calculations with
that of spatial ones. This is achieved by dividing the scale parameter by the average distance
between all cells. Additionally, we halve both the alpha and the modified scale values. Although
this does not equate to perfect scaling, we present very similar results using the scaling method
significantly more efficiently (Supplementary Fig. 23b, c).

In silico receptor perturbation

859 After identifying the top interactions between cell types, we conducted in silico perturbation using
860 our trained deep learning model $\mathcal{F}$. To illustrate the process of in silico receptor blocking, we
861 centered our attention on the receptors involved in the top 25 LR interactions, spanning all cell
862 type pairs. We then subjected these receptors to a perturbation analysis.

863
We permutated feature values for each receptor, independent of the output and simulated receptor
blocking and scaled the expression of the permutated receptor by the median expression of all
receptors. This manipulation of the input receptor vector, $x_j^{R \setminus r}$, allowed us to gauge the ensuing

influence on the gene expression downstream. After each receptor's perturbation, we singled out
the most impacted genes on average across all cells. Specifically, we looked at the top 50 most
impacted genes. This approach enabled us to assess the effects of receptor blocking at the
transcriptome level. To further demonstrate the potency of our model in executing in silico
perturbation, we utilized the DisGeNET[57] database as the potential validation. We verified whether
the perturbed genes are associated with the specific diseases that pique our interest – Breast
Cancer/PDAC (i.e., the disease associated with the dataset). To quantify the strength of the
association between each receptor and the disease, we applied a binomial test to compute the
statistical significance of the overlap between the perturbed genes and the genes associated with
each disease (breast cancer, breast carcinoma, PDAC, pancreatic carcinoma). We further refined
our results by adjusting the P-values over all disease gene sets, utilizing the FDR-Bonferroni
Correction method, ensuring the robustness of our findings. Additionally, we confirmed the
effectiveness of these receptor-inhibitor therapeutics using Kaplan-Meier (KM) survival plots[88].
880

Agent Based Dynamic Simulations and Animation of CCIs

CellAgentChat offers dynamic simulation capabilities for analyzing the impact of CCIs on cell
states over short durations. Users have the flexibility to either execute the model once, emulating
other state-of-the-art methods and inferring CCIs at the current moment, or they can run it over
multiple short time steps to simulate temporal cell state dynamics. At each time point, we compute
CCIs as described earlier. Subsequently, our trained neural network predicts gene expression levels
for each cell post-CCI for the next time step. Although we cannot directly substitute existing gene
expression values for the next time step with predicted ones due to uncertainties since many factors
other than CCIs could have an influence on the expression, we place trust in the directional changes
for each gene —whether they increase or decrease[70]. For each cell we then identify genes with
statistically significant changes and adjust their expression for that cell within the ABM, either
increasing or decreasing them by a small, specified amount (0.01) to simulate minor expression
changes[70]. This iterative process is repeated across all time steps, reflecting the dynamic
modulation of gene expression by CCIs.

CellAgentChat offers a unique feature by allowing the inference of cellular interactions between
individual cells rather than just cell types. Its real-time animation platform displays the CRS of
each cell, with size representing the strength; larger cells indicate stronger CRS (Supplementary
Fig. 23a). The platform also includes customizable hyperparameters, such as the maximum number
of steps (default of 1), δ for controlling long-range and short-range interactions (default of 1) and
the degrees of spatial freedom τ (default of 2) enabling effective in silico permutations. See
supplementary text for more information on how to effectively set up the hyperparameters to
facilitate simulations.

In addition, CellAgentChat offers hyperparameters for controlling the LR universe. Users can
block a specific LR pair or a specific receptor to observe downstream gene expression changes,
which can aid in identifying potential therapeutic targets. The animation platform also allows for
visualizing specific cell types/clusters, with modifications only applied to those selected.
Furthermore, users have the option to visualize specific receptor or LR pairs. This visualization
tool enables researchers to understand cell communication patterns at a granular level.

Functional Evaluation of Ligand-Receptor pairs inferred by CellAgentChat

We conducted pathway enrichment analysis to validate the interactions inferred by CellAgentChat functionally. We utilized both ligands and receptors from the top 25 highest scoring LR pairs among all cell type pairs as the input gene set. To perform pathway enrichment analysis, we employed g: Profiler[89] and selected all terms from the Reactome pathway with a minimum adjusted P-value below 0.01. This approach was consistent across all datasets, regardless of spatial or non-spatial. In the cases where we compared the enrichments for the Reactome pathways between two groups (e.g. long-range group vs. short-range group), we used signed-rank Wilcoxon test.

Benchmarking

We conducted a comparative analysis of CellAgentChat with five other state-of-the-art methods, namely CellChat, CellPhoneDB, NICHES, COMMOT, and Scriabin. Among these, CellChat, CellPhoneDB, and NICHES all facilitate the inference of cell-cell interactions (CCIs) using both spatial and non-spatial data. Conversely, COMMOT exclusively supports spatial data, while Scriabin exclusively supports non-spatial data. Additionally, Scriabin allows for the computation of a cell-cell interaction matrix (CCIM); however, it is recommended only for small datasets due to scalability limitations. Consequently, we could only evaluate Scriabin's performance for the PDAC and the SeqFISH+ mouse somatosensory cortex datasets. We used the LR universe derived from CellTalkDB for all methods.

We conducted systematic benchmarking with and without spatial data to validate CellAgentChat's ability to accurately infer interactions prior to integrating the spatial tendencies of CCIs. To assess the accuracies of the methods, we measured the Pearson Correlation to evaluate the communication pattern across cell types of each of the six methods. In the absence of a ground-truth, we employed an ensemble approach. We normalized the interaction matrix computed by each method and took the average to create an ensemble interaction matrix as our new "ground-truth". We then calculated the Pearson correlation between the ensemble interaction matrix and the inferred communication patterns of each method. Additionally, we compared the interaction matrix of each method with the overall distance matrix of the cells to verify whether CellAgentChat's CCI predictions were influenced by the spatial proximity between cells. Specifically, we calculated the Pearson Correlation between the cell interaction matrix with the cell-cell inverse distance ($1/d$) matrix.

As described above, we also conducted pathway enrichment analysis using the genes from the top 100 LR pairs to evaluate the functional validation of the LR interactions inferred by all six methods. We identified pathways through Reactome pathway enrichment analysis of the inferred interactions by at least two methods, to capture the most relevant pathways, and compared the -$\log_{10}$ adjusted P-values across all methods. To differentiate the functional validity of the LR interactions across the methods we used the Wilcoxon test (one-sided).

Finally, to compare the similarity in the inferred interactions we determined the number of shared LR interactions from the top 100 pairs identified by each method.

Data availability

CellTalkDB is available at (https://github.com/ZJUFanLab/CellTalkDB). The SCC and PDAC datasets analyzed in this study are available from the Gene Expression Omnibus (GEO) repository

under the following accession numbers: GSE144240 (GSM4565826) and GSE111672 (GSM3036911), respectively. The data for the SeqFISH+ mouse somatosensory cortex dataset can be found at the following GitHub link: https://github.com/CaiGroup/seqFISH-PLUS. The Xenium human breast cancer dataset (sample 2) can be found at https://www.10xgenomics.com/products/xenium-in-situ/preview-dataset-human-breast. The stereo-seq developmental mouse hippocampus dataset can be found at STOmicsDB[90] https://db.cngb.org/stomics/mosta/. We used the h5ad file titled *"Dorsal_midbrain_cell_bin.h5ad"*.

Code availability

The CellAgentChat software package and source code are available at our GitHub repository (https://github.com/mcgilldinglab/CellAgentChat). All preprocessed .h5ad files used in this study are also available in the same GitHub repository.

Acknowledgements

This work is supported by grants from the Canadian Institutes of Health Research (CIHR) [PJT-180505 to J.D]; the Fonds de recherche du Québec - Santé (FRQS) [295298 to J.D., 295299 to J.D.]; the Natural Sciences and Engineering Research Council of Canada (NSERC) [RGPIN2022-04399 to J.D.; RGPIN-2016-05174 to Y.L.]; and the Meakins-Christie Chair in Respiratory Research [to J.D.]. This publication is part of the Human Cell Atlas (HCA) publication bundle (HCA-10).

REFERENCES

Hu, M. & Polyak, K. Microenvironmental regulation of cancer development. *Current opinion in genetics & development* **18**, 27-34 (2008).

Shao, X., Lu, X., Liao, J., Chen, H. & Fan, X. New avenues for systematically inferring cell-cell communication: through single-cell transcriptomics data. *Protein & cell* **11**, 866-880 (2020).

Armingol, E., Officer, A., Harismendy, O. & Lewis, N. E. Deciphering cell–cell interactions and communication from gene expression. *Nature Reviews Genetics* **22**, 71-88 (2021).

Zhou, X. *et al.* Circuit design features of a stable two-cell system. *Cell* **172**, 744-757. e717 (2018).

Rouault, H. & Hakim, V. Different cell fates from cell-cell interactions: core architectures of two-cell bistable networks. *Biophysical Journal* **102**, 417-426 (2012).

Bonnans, C., Chou, J. & Werb, Z. Remodelling the extracellular matrix in development and disease. *Nature reviews Molecular cell biology* **15**, 786-801 (2014).

Vento-Tormo, R. *et al.* Single-cell reconstruction of the early maternal–fetal interface in humans. *Nature* **563**, 347-353 (2018).

Stubbington, M. J., Rozenblatt-Rosen, O., Regev, A. & Teichmann, S. A. Single-cell transcriptomics to explore the immune system in health and disease. *Science* **358**, 58-63 (2017).

Giladi, A. *et al.* Dissecting cellular crosstalk by sequencing physically interacting cells. *Nature Biotechnology* **38**, 629-637 (2020).

10 Yuan, G.-C. *et al.* Challenges and emerging directions in single-cell analysis. *Genome*
biology **18**, 1-8 (2017).
11 Marx, V. Method of the Year: spatially resolved transcriptomics. *Nature methods* **18**, 9-14
(2021).
12 Rodriques, S. G. *et al.* Slide-seq: A scalable technology for measuring genome-wide
expression at high spatial resolution. *Science* **363**, 1463-1467 (2019).
13 Xia, C., Babcock, H. P., Moffitt, J. R. & Zhuang, X. Multiplexed detection of RNA using
MERFISH and branched DNA amplification. *Scientific reports* **9**, 7721 (2019).
14 Eng, C.-H. L. *et al.* Transcriptome-scale super-resolved imaging in tissues by RNA
seqFISH+. *Nature* **568**, 235-239 (2019).
15 Chen, A. *et al.* Spatiotemporal transcriptomic atlas of mouse organogenesis using DNA
nanoball-patterned arrays. *Cell* **185**, 1777-1792. e1721 (2022).
16 Janesick, A. *et al.* High resolution mapping of the tumor microenvironment using
integrated single-cell, spatial and in situ analysis. *Nature Communications* **14**, 8353
(2023).
17 Efremova, M., Vento-Tormo, M., Teichmann, S. A. & Vento-Tormo, R. CellPhoneDB:
inferring cell–cell communication from combined expression of multi-subunit ligand–
receptor complexes. *Nature protocols* **15**, 1484-1506 (2020).
18 Jin, S. *et al.* Inference and analysis of cell-cell communication using CellChat. *Nature*
communications **12**, 1088 (2021).
19 Cabello-Aguilar, S. *et al.* SingleCellSignalR: inference of intercellular networks from
single-cell transcriptomics. *Nucleic Acids Research* **48**, e55-e55 (2020).
20 Asp, M., Bergenstråhle, J. & Lundeberg, J. Spatially resolved transcriptomes—next
generation tools for tissue exploration. *BioEssays* **42**, 1900221 (2020).
21 Cang, Z. *et al.* Screening cell–cell communication in spatial transcriptomics via collective
optimal transport. *Nature Methods* **20**, 218-228 (2023).
22 Raredon, M. S. B. *et al.* Comprehensive visualization of cell–cell interactions in single-
cell and spatial transcriptomics with NICHES. *Bioinformatics* **39**, btac775 (2023).
23 Jin, S., Plikus, M. V. & Nie, Q. CellChat for systematic analysis of cell-cell
communication from single-cell and spatially resolved transcriptomics. *BioRxiv*,
2023.2011. 2005.565674 (2023).
24 Wilk, A. J., Shalek, A. K., Holmes, S. & Blish, C. A. Comparative analysis of cell–cell
communication at single-cell resolution. *Nature Biotechnology*, 1-14 (2023).
25 Brady-Kalnay, S. M. Vol. 6 344-345 (Taylor & Francis, 2012).
26 Baruzzo, G., Cesaro, G. & Di Camillo, B. Identify, quantify and characterize cellular
communication from single-cell RNA sequencing data with scSeqComm. *Bioinformatics*
38, 1920-1929 (2022).
27 Bonabeau, E. Agent-based modeling: Methods and techniques for simulating human
systems. *Proceedings of the national academy of sciences* **99**, 7280-7287 (2002).
28 Yu, J. S. & Bagheri, N. Agent-based models predict emergent behavior of heterogeneous
cell populations in dynamic microenvironments. *Frontiers in Bioengineering and*
Biotechnology **8**, 249 (2020).
29 Mendelsohn, J. & Baselga, J. Status of epidermal growth factor receptor antagonists in
the biology and treatment of cancer. *Journal of clinical oncology* **21**, 2787-2799 (2003).
30 Tong, C. K. & Vidyadaran, S. Role of microglia in embryonic neurogenesis.
Experimental Biology and Medicine **241**, 1669-1675 (2016).

Li, W. *et al.* Endothelial cells regulate astrocyte to neural progenitor cell trans-differentiation in a mouse model of stroke. *Nature communications* **13**, 7812 (2022).

Vogenstahl, J., Parrilla, M., Acker-Palmer, A. & Segarra, M. Vascular regulation of developmental neurogenesis. *Frontiers in Cell and Developmental Biology* **10**, 890852 (2022).

Tensaouti, Y., Stephanz, E. P., Yu, T.-S. & Kernie, S. G. ApoE regulates the development of adult newborn hippocampal neurons. *ENeuro* **5** (2018).

Yin, Z. *et al.* APOE4 impairs the microglial response in Alzheimer's disease by inducing TGFβ-mediated checkpoints. *Nature immunology* **24**, 1839-1853 (2023).

Kakoi, C., Udo, H., Matsukawa, T. & Ohnuki, K. Collagen peptides enhance hippocampal neurogenesis and reduce anxietyrelated behavior in mice. *Biomedical Research* **33**, 273-279 (2012).

Swonger, J. M., Liu, J. S., Ivey, M. J. & Tallquist, M. D. Genetic tools for identifying and manipulating fibroblasts in the mouse. *Differentiation* **92**, 66-83 (2016).

Leiter, O. *et al.* Platelet-derived exerkine CXCL4/platelet factor 4 rejuvenates hippocampal neurogenesis and restores cognitive function in aged mice. *Nature Communications* **14**, 4375 (2023).

Malatesta, P., Hartfuss, E. & Götz, M. Isolation of radial glial cells by fluorescent-activated cell sorting reveals a neuronal lineage. *Development* **127**, 5253-5263 (2000).

Qian, X. *et al.* Timing of CNS cell generation: a programmed sequence of neuron and glial cell production from isolated murine cortical stem cells. *Neuron* **28**, 69-80 (2000).

Cope, E. C. & Gould, E. Adult neurogenesis, glia, and the extracellular matrix. *Cell Stem Cell* **24**, 690-705 (2019).

Fromme, J. E. & Zigrino, P. The role of extracellular matrix remodeling in skin tumor progression and therapeutic resistance. *Frontiers in Molecular Biosciences*, 370 (2022).

Wareham, L. K., Baratta, R. O., Del Buono, B. J., Schlumpf, E. & Calkins, D. J. Collagen in the central nervous system: contributions to neurodegeneration and promise as a therapeutic target. *Molecular Neurodegeneration* **19**, 11 (2024).

Kania, A. & Klein, R. Mechanisms of ephrin–Eph signalling in development, physiology and disease. *Nature reviews Molecular cell biology* **17**, 240-256 (2016).

Boschert, V., Teusch, J., Müller-Richter, U. D., Brands, R. C. & Hartmann, S. PKM2 modulation in head and neck squamous cell carcinoma. *International Journal of Molecular Sciences* **23**, 775 (2022).

Ludwig, N. *et al.* CD44 (+) tumor cells promote early angiogenesis in head and neck squamous cell carcinoma. *Cancer Letters* **467**, 85-95 (2019).

Zhang, G. *et al.* Identity of MMP1 and its effects on tumor progression in head and neck squamous cell carcinoma. *Cancer medicine* **11**, 2516-2530 (2022).

Schaafsma, E., Fugle, C. M., Wang, X. & Cheng, C. Pan-cancer association of HLA gene expression with cancer prognosis and immunotherapy efficacy. *British Journal of Cancer* **125**, 422-432 (2021).

Atasoy, M., Anadolu-Braise, R., Pirim, I., Dogan, H. & Ikbal, M. HLA antigen profile differences in patients with SCC (squamous cell carcinoma) in-situ/actinic keratosis and invasive SCC: Is there a genetic succeptibility for invasive SCC development? *The Eurasian Journal of Medicine* **41**, 162 (2009).

49 Pontén, F., Jirström, K. & Uhlen, M. The Human Protein Atlas—a tool for pathology. *The*
Journal of Pathology: A Journal of the Pathological Society of Great Britain and Ireland
216, 387-393 (2008).

50 Parker, A. L. *et al.* Extracellular matrix profiles determine risk and prognosis of the
squamous cell carcinoma subtype of non-small cell lung carcinoma. *Genome Medicine*
14, 1-29 (2022).

51 Ahrends, T. & Borst, J. The opposing roles of CD 4+ T cells in anti-tumour immunity.
Immunology **154**, 582-592 (2018).

52 Centuori, S. M. & Bauman, J. E. c-Met signaling as a therapeutic target in head and neck
cancer. *The Cancer Journal* **28**, 346-353 (2022).

53 Szturz, P. *et al.* Understanding c-MET signalling in squamous cell carcinoma of the head
& neck. *Critical reviews in oncology/hematology* **111**, 39-51 (2017).

54 Meireles Da Costa, N. *et al.* Potential therapeutic significance of laminin in head and
neck squamous carcinomas. *Cancers* **13**, 1890 (2021).

55 Jourquin, J., Tripathi, M., Guess, C. & Quaranta, V. Laminins and cancer progression.
Cell-Extracellular Matrix Interactions in Cancer, 87-109 (2010).

56 Ramovs, V., Te Molder, L. & Sonnenberg, A. The opposing roles of laminin-binding
integrins in cancer. *Matrix Biology* **57**, 213-243 (2017).

57 Piñero, J. *et al.* DisGeNET: a discovery platform for the dynamical exploration of human
diseases and their genes. *Database* **2015** (2015).

58 Masuda, H. *et al.* Role of epidermal growth factor receptor in breast cancer. *Breast*
cancer research and treatment **136**, 331-345 (2012).

59 Ralser, D. J. *et al.* Molecular and immune correlates of PDCD1 (PD-1), PD-L1 (CD274),
and PD-L2 (PDCD1LG2) DNA methylation in triple negative breast cancer. *Journal of*
Immunotherapy **44**, 319-324 (2021).

60 Ghahremani Dehbokri, S. *et al.* CTLA-4: as an immunosuppressive immune checkpoint
in breast cancer. *Current Molecular Medicine* **23**, 521-526 (2023).

61 Ren, J., Chen, Y., Kong, W., Li, Y. & Lu, F. Tumor protein D52 promotes breast cancer
proliferation and migration via the long non-coding RNA NEAT1/microRNA-218-5p
axis. *Annals of Translational Medicine* **9** (2021).

62 Corso, G., Veronesi, P., Sacchini, V. & Galimberti, V. Prognosis and outcome in CDH1-
mutant lobular breast cancer. *European Journal of Cancer Prevention* **27**, 237-238
(2018).

63 Soysal, S. D. *et al.* EpCAM expression varies significantly and is differentially associated
with prognosis in the luminal B HER2+, basal-like, and HER2 intrinsic subtypes of
breast cancer. *British journal of cancer* **108**, 1480-1487 (2013).

64 Moncada, R. *et al.* Integrating microarray-based spatial transcriptomics and single-cell
RNA-seq reveals tissue architecture in pancreatic ductal adenocarcinomas. *Nature*
biotechnology **38**, 333-342 (2020).

65 Friedl, P. & Alexander, S. Cancer invasion and the microenvironment: plasticity and
reciprocity. *Cell* **147**, 992-1009 (2011).

66 Chen, P., Cescon, M. & Bonaldo, P. Collagen VI in cancer and its biological mechanisms.
Trends in molecular medicine **19**, 410-417 (2013).

67 Perez, V. M., Kearney, J. F. & Yeh, J. J. The PDAC extracellular matrix: a review of the
ECM protein composition, tumor cell interaction, and therapeutic strategies. *Frontiers in*
Oncology **11**, 751311 (2021).

68 Liu, Z., Sun, D. & Wang, C. Evaluation of cell-cell interaction methods by integrating
single-cell RNA sequencing data with spatial information. *Genome Biology* **23**, 1-38
(2022).
69 Neufeld, G. *et al.* The role of the semaphorins in cancer. *Cell Adhesion & Migration* **10**,
652-674 (2016).
70 Ding, J., Hagood, J. S., Ambalavanan, N., Kaminski, N. & Bar-Joseph, Z. iDREM:
Interactive visualization of dynamic regulatory networks. *PLoS computational biology*
14, e1006019 (2018).
71 Francis, K. & Palsson, B. O. Effective intercellular communication distances are
determined by the relative time constants for cyto/chemokine secretion and diffusion.
Proceedings of the National Academy of Sciences **94**, 12258-12262 (1997).
72 Wan, X. *et al.* Integrating spatial and single-cell transcriptomics data using deep
generative models with SpatialScope. *Nature Communications* **14**, 7848 (2023).
73 Shao, X. *et al.* CellTalkDB: a manually curated database of ligand–receptor interactions
in humans and mice. *Briefings in bioinformatics* **22**, bbaa269 (2021).
74 Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene expression
data analysis. *Genome biology* **19**, 1-5 (2018).
75 Islam, S. *et al.* Quantitative single-cell RNA-seq with unique molecular identifiers.
Nature methods **11**, 163-166 (2014).
76 Ilicic, T. *et al.* Classification of low quality cells from single-cell RNA-seq data. *Genome*
biology **17**, 1-15 (2016).
77 Rao, A., Barkley, D., França, G. S. & Yanai, I. Exploring tissue architecture using spatial
transcriptomics. *Nature* **596**, 211-220 (2021).
78 Tran, M. *et al.* A robust experimental and computational analysis framework at multiple
resolutions, modalities and coverages. *Frontiers in immunology* **13**, 911873 (2022).
79 Feng, J. *et al.* sc2MeNetDrug: A computational tool to uncover inter-cell signaling targets
and identify relevant drugs based on single cell RNA-seq data. *PLoS computational*
biology **20**, e1011785 (2024).
80 Peng, L. *et al.* CellEnBoost: A boosting-based ligand-receptor interaction identification
model for cell-to-cell communication inference. *IEEE Transactions on NanoBioscience*
(2023).
81 Peng, L., Xiong, W., Han, C., Li, Z. & Chen, X. CellDialog: A Computational Framework
for Ligand-receptor-mediated Cell-cell Communication Analysis III. *IEEE Journal of*
Biomedical and Health Informatics (2023).
82 Kanehisa, M. & Goto, S. KEGG: kyoto encyclopedia of genes and genomes. *Nucleic*
acids research **28**, 27-30 (2000).
83 Jassal, B. *et al.* The reactome pathway knowledgebase. *Nucleic acids research* **48**, D498-
D503 (2020).
84 Han, H. *et al.* TRRUST v2: an expanded reference database of human and mouse
transcriptional regulatory interactions. *Nucleic acids research* **46**, D380-D386 (2018).
85 Bovolenta, L., Acencio, M. & Lemke, N. HTRIdb: an open-access database for
experimentally verified human transcriptional regulation interactions. *Nature Precedings*,
1-1 (2012).
86 Mi, X., Zou, B., Zou, F. & Hu, J. Permutation-based identification of important
biomarkers for complex diseases via machine learning models. *Nature communications*
12, 3008 (2021).

87   Arnol, D., Schapiro, D., Bodenmiller, B., Saez-Rodriguez, J. & Stegle, O. Modeling cell-
1180   cell interactions from spatial molecular data with spatial variance component analysis.
1181   *Cell reports* **29**, 202-211. e206 (2019).
1182   88   Győrffy, B. Survival analysis across the entire transcriptome identifies biomarkers with
1183   the highest prognostic power in breast cancer. *Computational and structural*
1184   *biotechnology journal* **19**, 4101-4109 (2021).
1185   89   Raudvere, U. *et al.* g: Profiler: a web server for functional enrichment analysis and
1186   conversions of gene lists (2019 update). *Nucleic acids research* **47**, W191-W198 (2019).
1187   90   Xu, Z. *et al.* STOmicsDB: a database of Spatial Transcriptomic data. *bioRxiv*, 2022.2003.
1188   2011.481421 (2022).
1189