Gut microbiome remains stable following COVID-19 vaccination in healthy and immuno-compromised individuals

Rebecca H. Boston[1], Rui Guan[1], Lajos Kalmar[1], Sina Beier[1], Emily C. Horner[1], Nonantzin Beristain-Covarrubias[1], Juan Carlos Yam-Puc[1], Pehuén Pereyra Gerber[2,3], Luisa Faria[1], Anna Kuroshchenkova[1], Anna E. Lindell[1], Sonja Blasche[1], Andrea Correa-Noguera[4], Anne Elmer[5], Caroline Saunders[5], Areti Bermperi[5], Sherly Jose[5], Nathalie Kingston[6], CITIID-NIHR COVID-19 BioResource Collaboration, Sofia Grigoriadou[7], Emily Staples[1], Matthew S. Buckland[7,8], Sara Lear[4], Nicholas J. Matheson[2,3,4,9], Vladimir Benes[10], Christine Parkinson[4], James E. D. Thaventhiran[1,4,*] and Kiran R. Patil[1,*]

[1] Medical Research Council Toxicology Unit, University of Cambridge, Cambridge, UK.

[2] Cambridge Institute of Therapeutic Immunology and Infectious Disease (CITIID), University of Cambridge, Cambridge, UK

[3] Department of Medicine, University of Cambridge, Cambridge, UK

[4] Department of Clinical Immunology, Cambridge University NHS Hospitals Foundation Trust, Cambridge, UK

[5] NIHR Cambridge Clinical Research Facility, Cambridge, UK

[6] NIHR BioResource, Cambridge University Hospitals NHS Foundation Trust, Cambridge, UK

[7] Department of Clinical Immunology, Barts Health, London, UK

[8] UCL GOSH Institute of Child Health Division of Infection and Immunity, Section of Cellular and Molecular Immunology, London, UK

[9] NHS Blood and Transplant, Cambridge, UK

[10] European Molecular Biology Laboratory, Heidelberg, Germany

*Correspondence to: JEDT, jedt2@mrc-tox.cam.ac.uk; KRP, kp533@mrc-tox.cam.ac.uk.

Abstract

The bidirectional interaction between the immune system and the gut microbiota is a key contributor to various host physiological functions. Immune-associated diseases such as cancer and autoimmunity, as well as the efficacy of immunomodulatory therapies, have been linked to microbiome variation. While COVID-19 infection has been shown to cause microbial dysbiosis, it remains understudied whether the inflammatory response associated with vaccination also impacts the microbiota. Here, we investigate the temporal impact of COVID-19 vaccination on the gut microbiome in healthy and immuno-compromised individuals; the latter included patients with primary immunodeficiency and cancer patients on immunomodulating therapies. We find that the gut microbiome remained remarkably stable post-vaccination irrespective of diverse immune status, vaccine response, and microbial composition spanned by the cohort. The stability is evident at all evaluated levels including diversity, phylum, species, and functional capacity. Our results indicate the resilience of the gut microbiome to host immune changes triggered by COVID-19 vaccination and suggest minimal, if any, impact on microbiome-mediated processes. These findings encourage vaccine acceptance, particularly when contrasted with the significant microbiome shifts observed during COVID-19 infection.

Introduction

In the first 30 months of the pandemic, there have been reported to be almost 800 million PCR confirmed cases of COVID-19 infection and approaching 7 million related deaths globally[1]. To reduce this severity, vaccines were deployed with the aim of promoting anti-SARS-CoV-2 immunity, with almost 13.5 billion vaccine doses administered globally, 150 million of which were administered in the United Kingdom[1]. Yet, continued COVID-19 transmission remains of concern[2] with one of the reasons being vaccine hesitancy[3]. Thus, data helping to understand holistic effects of vaccination will have a profound impact on the public health management.

The SARS-CoV-2 mRNA and viral vector vaccines induce a strong immune response through the promotion of both innate and adaptive immunity against the spike protein[4]. Of note, there has been reported to be a promotion of inflammatory cytokines IFNy, IL-15 and IL-6 secretion in response to vaccination[5] each known to impact intestinal epithelial architecture and mucosal immunity[6–8]. The sequestration of the spike mRNA by antigen-presenting cells aims to limit the spread into systemic circulation, however the SARS-CoV-2 spike protein was also found in blood plasma following vaccination[9], potentially leading to inflammation at different sites of the body other than the site of vaccination, including that of the gut. This led us to hypothesize that the systemic immune response to the SARS-CoV-2 vaccination may impact the gut microbiota.

Numerous studies have analysed the gut microbiome during COVID-19 infection[10], finding notable depletion in both commensal bacterial, such as *Bacteroides* and *Bifidobacterium spp.* and beneficial *Lachnospiraceae,* coupled with increased abundances of opportunistic pathogens such as *Streptococcus* and *Clostridium hathawayi*; this indicates a marked dybiosis induced by COVID-19 infection. Yet since the initiation of the vaccination programme against COVID-19, very few studies have addressed the impact of the vaccination on the gut microbiome[11–14]. Previous work has addressed the link between the gut microbiome and vaccine immunogenicity in which, baseline abundances of certain bacterial species before the

first vaccine dose have been correlated with a defined end point of vaccine efficacy, typically

a vaccine-related readout, such as virus neutralisation or spike-specific antibody titres.

Yet, two open questions remain: how the gut microbiome is affected by COVID-19 vaccination

in the days following vaccination when the inflammatory response is at its highest, and are

any immediate changes in the gut microbiome maintained or resolved once humoral immunity

has been initiated. Addressing this knowledge-gap could help understand the extent and the

nature of reciprocal links between the gut microbiome and systemic immunity in the context of

vaccination. We therefore sought to analyse the gut microbiome of patients receiving doses

of the COVID-19 vaccines to decipher whether there were any notable, characteristic changes

in the gut microbiome in either healthy or immunocompromised individuals [Table 1]. The

immunocompromised patients we recruited were cancer patients receiving immune

checkpoint modulating therapies and patients with primary immunodeficiencies. These

cohorts consist of patients with impaired immunity resulting from either therapeutic

intervention or monogenic defects in immunoregulatory genes (CTLA4, NFKB1, CD40L),

respectively.

This presented us with the unique opportunity to elucidate whether the COVID-19 vaccines

alter the gut microbiome in the absence of complete, functional immunity and subsequent

impairments of the regulation of the gut microbiome.

Methods

Study recruitment and ethics

Participants volunteered for an observational study and were enrolled to one of 3 cohorts,

healthy controls, cancer patients which presented with either melanoma or renal malignancies,

or primary immunodeficient patients with defined mutations in key immunoregulatory genes,

as well as patients with clinical presentation aligning to that of defined primary

immunodeficiency, including antibody deficiency [Table 1]. Almost all participants in each

cohort received BNT162b2 Pfizer vaccine, aside from one patient at second dose receiving

AstraZeneca, and 6 patients receiving Moderna at the third [Table 1]. Patients were excluded

if presenting with positive COVID-19 serology or if presenting in hospital with clinal

symptoms/features related to their disease which may influence the physiological response to

the COVID-19 vaccination. The research was conducted in accordance with the principles of

Good Clinical Practice and following approved protocols of the NIHR National Bioresource.

Samples were collected at NIHR Cambridge Clinical Research Facility, Addenbrookes

Hospital, Cambridge, UK. Samples were provided with the written informed consent of all

study participants under the NIHR National BioResource - Research Tissue Bank (NBR-RTB)

ethics (research ethics committee (REC):17/EE/0025). All participants were consented under

the East of England Cambridge South national consent (REC:13/EE/0325).

Participant sampling

Participant samples were anonymised by clinical staff prior to sample delivery to the research

lab. Peripheral blood and fecal samples were collected longitudinally over the course of up to

three doses of the vaccines against COVID-19 (sample coverage varied across doses

[Supplementary Figure 1]). Samples were taken from 9[th] February 2021 to the 22[nd] February

2022. Peripheral blood mononuclear cells (PBMCs) were extracted from blood samples using

density gradient centrifugation, stored temporarily at -80°C, before being transferred to long-

term storage in liquid nitrogen. Serum was isolated from peripheral blood via centrifugation

and stored at -80°C until required. Fecal samples were collected at 3 timepoints around each

vaccine dose: pre-dose (94% of samples taken within 3 days prior to vaccination, the

remaining 3 samples taken up to 14 days prior), acute (day 2 or 3 after vaccination) or late

(day 16–28 after vaccination). Fecal matter was collected in OMNIgene•GUT kits (DNA

Genotek, Canada) whereby samples are stored in a stabilizing, inactivating solution. Samples

were transported to the lab and homogenized upon arrival before being stored at -80°C until

required.

Fecal DNA extraction and sequencing

DNA was extracted from fecal samples using QIAamp® PowerFecal® Pro DNA kits (Qiagen).

Samples were thawed and approximately 250mg of fecal sample was lysed via bead beating.

According to the kit protocol, the sample was then cleaned of non-DNA organic and inorganic

material, then washed using ethanol. DNA was eluted into 10mM Tris and quantified using the

Qubit™ 1X dsDNA High Sensitivity (HS) Assay Kit (Thermo Fisher Scientific, UK) using the

Qubit™ fluorometer (Thermo Fisher Scientific, UK). DNA at a concentration of 10mg/uL was

sent for sequencing. Samples were sent in bulk for sequencing blinded to sample timepoints

of each patient. Shotgun metagenomic sequencing was performed with Illumina NextSeq 2000

sequencing platform using paired-end reads of 150bp in length, resulting in a median of

approximately 28 million reads per sample.

Shotgun metagenomic analysis

Raw sequencing data was pre-processed with PRINSEQ++[15] (v.1.2.4) in paired read mode,

quality trimming to a minimal Phred score of 30 in a window of 15 bases and removing reads

of less than 75 bp length after trimming. Additionally, host contamination was removed by

mapping against the GRCh38 reference human genome using Bowtie2[16] (v.2.4.5) and

removing any mapped reads from the dataset.

Raw, trimmed and filtered reads were checked for quality using FastQC (v.0.11.9)

(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/). From the remaining read pairs,

taxonomic profiling was determined using mOTUs3[17] (v.3.0.1) profiler. For functional profiling,

the remaining read pairs after filtering were assembled using metaSPAdes[18] (v.3.15.4) with a

k-mer size of 55. The resulting scaffolds were filtered for at least 200 bp length and weighted

by their average coverage of the filtered reads to enable quantitative analysis. The remaining

scaffolds were aligned to the EggNOG database[19](downloaded on 2022/04/08) using

DIAMOND[20] (v.2.0.13). Microbiome analysis was performed in R (v.4.2.2) using relative

abundances, within the phyloseq[21] (v.1.42.0), microbiome (v.1.20.0)[22] and vegan[23] (v.2.6-4)

packages. Differential abundance analysis was performed using DESeq2[24] (v.1.38.3).

Serological assessment of immune response to COVID-19 vaccines

Serum samples were thawed, heat-inactivated at 56°C for 30 mins and measured for the

dilution of serum that reduces viral activity by 50% (NT_{50}); the method for which has been

previously described[25]. For anti-SARS-CoV-2 specific IgG antibodies, we utilised a previously

described method[26,27], in which Luminex bead sets are covalently coupled to the recombinant

SARS-CoV-2 proteins nucleocapsid protein, spike and receptor-binding domain (RBD) to

quantify antibody levels.

Statistical analysis

Wilcoxon tests with multiple testing correction using the bonferroni were deployed throughout,

using pairwise comparison where appropriate. Mixed effect linear modelling was performed

using lmer4[28] in R. Correlation was determined using Spearman's Rank coefficient.

Results

The composition of gut microbiome is not altered by vaccination against COVID-19

To investigate the impact of the COVID-19 vaccines on the gut microbiome, shotgun metagenomic sequencing was performed on a total of 239 fecal samples from 59 patients from our three cohorts (43 healthy control, 160 cancer and 36 primary immunodeficient patient samples) [Supplementary Figure 1]. Samples were taken over the course of 3 vaccine doses, at one of three vaccine timepoints: pre-dose (before vaccination), acute (2–3 days after vaccination) or late (16–28 days after vaccination) for each vaccine dose [Fig1A]. Pre-dose sampling provides a baseline assessment of the gut microbiome prior to vaccination, whilst acute samples provide the opportunity to assess the effect of vaccination on the gut microbiome at the height of the initial inflammatory response, and late samples allow a determination of any resolution or maintenance of acute effects.

When first observing the alpha diversities of all samples taken from each of our cohorts, we noted significant differences between the cohorts [Supplementary Figure 2A], however when assessing samples taken at different vaccine timepoints from within each cohort, we did not observe any significantly differences [Figure 1B]. In agreement with this, when assessing samples from the same patient at different vaccine timepoints using a paired sample analysis (tracking individual patient samples across multiple timepoints, e.g. pre-dose & acute) approach, we also did not see any significant differences in the alpha diversities of our patient samples [Supplementary Figure 2B][Supplementary Table 1]. This indicates that the COVID-19 vaccine is not affecting the diversity of the gut microbiome, despite the distinct microbial diversity between the cohorts and individuals.

We next applied principal component analysis (PCA) to visualize the beta-diversity of our microbiome composition data using the abundance of all detected operational taxonomic units (OTUs) [Fig1C]. The principal components (PCs) describe the largest variation components in the dataset, representing shifts in microbiome composition and potentially reflecting to the abundance changes of bacterial species between the samples. The first 5 principal

components were responsible for the 2.9%, 2.4%, 2.1%, 2.0% and 1.9% of variation in the

data respectively, and were further analysed using mixed effect linear models with multiple

input variables from our available metadata. In our linear models we asked whether the

vaccination timepoint of the samples in each cohort could improve the explained variance of

the PCs when compared to a baseline model describing the explained variance using patient

samples as the grouping variable. We found that there was no significant improvement on the

baseline model [Supplementary Figure 2C]. To further, a PERMANOVA analysis did not find

the covariate of the timepoints from which the samples were taken to be significantly affecting

the microbial composition [Supplementary Table 2]. These analyses together support that the

variance we see in our samples is not a signature of the COVID-19 vaccines, rather of the

patients presenting with different microbiome compositions themselves.

Next, we asked whether the COVID-19 vaccines induce any changes in the phylum-level

composition of the gut microbiome and profiled the relative abundance of phyla across all

samples, [Fig1D] observing variation in our patient samples. Moreover, when comparing the

top six most prevalent phyla, no significant differences were observed between vaccine

timepoints [Fig1E] despite significant differences in these phyla between cohorts and

individuals [Supplementary Figure 2D]. When using a paired analysis approach, we observed

only marginal, small effect size, differences (after multiple testing correction) in seven (out of

231) timepoint comparisons in the cancer cohort [Supplementary Figure 2E] [Supplementary

Table 3]. Of those 7, however, only one had a log2FC greater than 1 in either direction (+1 or

-1) (Supplementary Figure 2F), indicative of notable change in median relative abundance of

Lentisphaerae.

This demonstrates that the COVID-19 vaccines do not appreciably alter the composition of

the gut microbiome, irrespective of the unique compositions found in our cohort samples.

COVID-19 vaccination does not induce species level changes in the gut microbiome

We next sought to analyse differentially abundant microbial species between vaccine

timepoints, i.e. pre-dose, acute, and late, using DESeq2[24]. All cohorts were analysed

independently for the abundance changes in samples taken at each timepoint with the most

differentially abundant species in a representative heatmap. For the cancer cohort, when

assessing the abundance of these top differential responding bacterial species between

samples taken pre-dose and acutely, unsupervised clustering does not demonstrate evident

grouping of timepoints [Fig2A]. Among all the species, only two were significantly increased in

acute samples compared to pre-dose samples, *Klebsiella pneumoniae* and *Butyrivibrio*

crossotus found in 15 [p = 1.01e-24] and seven samples [p = 7.63e-12] out of the 97 cancer

patient samples respectively [Fig2B]. The former is only representative in a quarter of the

cohort (11 patients), only melanoma patients, and within those has an average relative

abundance of 0.7% [Fig2C]; the latter in two renal cancer patients, representing on average

3% of the relative abundance [Fig2D]. Considering approximately 2500 species are

represented across all patient samples, change in two low-abundant and sparsely represented

species signifies negligible changes. Similar findings were seen for our other two cohorts,

healthy controls and primary immunodeficient patients [Supplementary Figure 3A and 3C, and

3B and 3D respectively]. When performing paired sample differential abundance analysis in

all three cohorts using DESeq2, we find that no significantly altered species between pre-dose

and acute, or pre-dose and late sample in the healthy control and cancer cohorts. Samples

from only one primary immunodeficient patient showed a significant reduction in *Enterobacter*

sp. in a late sample compared to pre-dose [Supplementary Figure 3E]. Thus, our findings

demonstrate that on a species level there is no unified, biologically relevant change in

abundance of microbial species induced by the COVID-19 vaccines.

As the differential abundance analysis considers the change in abundance of all species

irrespective of their relative abundance within each sample, we were curious whether there

were any noticeable changes in the most abundant species found within each patient cohort

that could be attributed to the vaccine timepoints. There was no significant difference in any

of the most abundant 15 species in samples taken at any of the three vaccine timepoints

[Fig2E], representing on average 47% of the relative abundance of the species within patient

samples in the cancer cohort, 50% within the healthy controls and 53% within the primary

immunodeficient patient cohort. This indicates that we see no effect of the vaccine on the

species occupying the highest proportion of the microbial niche.

There was considerable concern both at the time of the initial vaccine programme, and to this

day, on the safety of COVID-19 vaccines. Given that, we sought to observe if there was any

differential outgrowth of bacterial species that have been associated to various immune-

related diseases, including gastric cancer and autoimmunity, as well as metabolic and

neurological diseases. Although our study cannot address long-term outgrowth of bacterial

associated to these diseases, we are able to highlight if there is any temporal, acute changes

in these species which is still of physiological relevance. *Faecalibacterium prausnitzii,* which

is reported to be reduced in both gastric cancers, autoimmunity, and Crohn's disease[29],

showed no significant abundance changes at the vaccine timepoints in our cohorts [Fig2F].

We also found no presence of *Helicobacter pylori*, which is strongly associated with the

initiation and development of gastric cancers[30]. *Akkermansia muciniphila,* found to be

increased in obesity[31] and correlated with response rates to immune checkpoint blockade

therapies in various malignancies[32], was not altered by vaccination in all three cohorts [Fig2G].

In studies of Alzheimer's, *Escherichia coli* has been demonstrated to promote

neurodegeneration[33], in our samples we did not see significant difference induced by the

vaccine in any of our cohorts [Fig2H]. This supports that the COVID-19 vaccine does not

promote the change in abundance of microbes that are associated with various immune-

related diseases within any of the three cohorts and is indicative of no greater risk of the

aforementioned diseases as a result of COVID-19 vaccination.

Gut microbiome diversity is not correlated with the magnitude of the response to the COVID-19 vaccines

Within the current literature, a few studies have reported changes in the gut microbiome that

correlate with vaccine efficacy[12–14], so we sought to determine whether the gut microbiome

composition was related to the magnitude of the COVID-19 vaccine response. We performed

an assessment of vaccine efficacy using a live-virus neutralisation assay, as a predictive

measure of vaccine protection[32], to assess whether vaccine efficacy had correlation with

microbial diversity.

When taking the neutralising capacity of patient serum at both second dose [Fig3A] and third

dose [Fig3B], we asked whether the Shannon diversity of gut microbiome at different vaccine

timepoints, was affected by or correlated with neutralisation. We did not see any correlation

between diversity and vaccine efficacy in any of our patient cohorts; the same is true for the

quantity of anti-spike IgG antibodies [Supplementary Figure 4]. This indicates that the gut

microbiome diversity was not correlated with the magnitude of the immune response in our

patient cohorts, thus suggesting that improved efficacy of the vaccine does not come at a cost

of microbial disturbance.

The gut microbiome functional capacity was not affected by COVID-19 vaccines

Having investigated the composition and relative abundance of the microbial species that

constitute the gut microbiome, we next sought to investigate whether the functional capacity

of the microbial species was altered by the COVID-19 vaccines. Using the EggNOG database,

we assigned functional annotations to the sequenced metagenomes. The highest level of

functional annotation depicts three functional groups, cellular processes and signalling,

information storage and processing, and metabolism. In these, we did not see any significant

differences between the vaccine timepoints within our cohorts [Fig4A]; similar to taxonomic

data presented earlier, when assessing cohort samples separately , there are significant

changes [Supplementary Figure 5A].

We next observed the abundance of the 22 defined functional groups in the next functional

annotation level down in the separate vaccine timepoints within each patient cohort [Fig4B].

Representative graphs of the most abundant functional annotations within each of the highest

three level functional levels remained unchanged after vaccination within our patient cohorts

[Fig4C], the same is true for the remaining 19 (Supplementary Figure 5B).

At the lowest functional annotation level, we interrogated the abundance of cluster of

orthologous genes (COGs) at separate vaccine timepoints within our cohorts. Remarkably, no

COGs out of a possible 2142 presented in our patient samples, were significantly altered

(when corrected for multiple testing) as a result of the COVID-19 vaccines in any of the three

cohorts. This demonstrates that the functional annotations of the gut microbiome are not

altered by the administration of the COVID-19 vaccines.

Discussion

To our knowledge, this study is the first to assess the gut microbiome composition in response

to the COVID-19 vaccines across multiple doses and at multiple timepoints with samples taken

pre-dose, acutely and late following vaccination. The study is also the first to assess the effect

of the COVID-19 vaccines on the gut microbiome in cancer patients and in patients with inborn

errors of immunity associated with severe immune dysregulation. As sampling across the

cohorts varied throughout, we opted to combine the three vaccine doses and assess vaccine

timepoints or samples from within each cohort. This allowed us to better observe the influence

of the COVID-19 vaccines in these contexts.

The relative abundance of microbes within the gut microbiome has more recently been

assessed with vaccine immunogenicity including that of vaccines against SARS-CoV-2 virus.

The first reported study of the gut microbiome in COVID-19 vaccinated patients correlated

vaccine immunogenicity of the inactivated virus, CoronaVac, and the mRNA vaccine encoding

spike protein, BNT162b2 vaccine, with the baseline abundance of gut *Bifidobacterium*

adolescentis and *Roseburia faecis* respectively[12], as well as noting shifts in microbiome

composition. We did not observe changes in these bacterial species, nor the composition, as

our study design was to look at change over vaccine doses individually rather all together. A

previous study investigated the variability of the gut microbiome response to the COVID-19

vaccine by correlating RNAseq data with microbial abundance using 16S rRNA gene amplicon

sequencing[14]. They identified several differentially abundant taxa between high- and low-

antibody   responders   and   high-   and   low-T-cell   responders.   In   the   context   of

immunocompromised cohorts, a previous study assessed patients with inflammatory bowel

disease[13] well known to be characterised by gut microbiome dysbiosis[34], who were receiving

anti-TNF immunomodulators. Their study did not demonstrate changes in diversity in above

geometric mean vaccine responders but found *Bilophila* abundance correlated to an improved

response. What these studies have in common is associating microbiome composition to

vaccine immunogenicity, however these studies analysed single post-vaccination time point

with potential for confounding effects of various lifestyle and other factors known to profoundly

affect microbiome composition such as diet, medications, sampling time of the day etc.[35–38].

Our multi-time-point study around individual vaccine dose allowed us to more robustly assess

the microbiome changes (or the lack thereof). As the patients enrolled in our study were not

all healthy, our cohort samples span a much wider and higher diversity compared to the other

study, and all our samples were taken in UK, it remains a possibility that country-specific

differences explain the variance in the findings across the studies.

In our patient cohorts, we did not find any significant effect on the diversity of the gut

microbiome following COVID-19 vaccination despite considerable differences between the

cohorts. In line with the studies investigating the gut microbiome of patients with primary

immunodeficiencies[39], we also observed decreased diversity in our cohort compared to control

samples. These patients have been demonstrated to exhibit increased gut permeability with

higher rates of bacterial translocation[40], perhaps indicative that there is bidirectional

permeability of systemic immunity affecting the gut microbiome. In our study, we find no

influence on microbiome variation after vaccination in patients at genetically determined

persistent state of immune dysregulation.

In melanoma patients, the presence of species bacteria species from the *Actinobacteria* and

Firmicutes phyla have been associated with better responses to immune checkpoint blockade

therapies[41–43]. An *Akkermansia muciniphilia* signature was also found in renal cancer patients

responding better to immune checkpoint blockade therapy[44,45]. While cancer progression is

reported to be linked to gut microbiome composition and its derived metabolites, these

associations vary between cancer types[46]. We saw wide compositional variation within our

cancer cohort samples, perhaps due to the wide range of disease presentation and treatment

included in our patients. Our study did not consider factors such diet and medications known

to affect the gut microbiome[47]. Nevertheless, this is not critical as our analysis suggests

stability as opposed to specific changes and post-hoc power calculation indicates sufficient

power against false negatives when taking paired samples (Pre-dose-Acute, Pre-dose-Late

and Acute-Late) (for effect size Cohen's d = 0.55, estimated power = 0.8). Cancer patients'

response to vaccination depend significantly on cancer type, for example antibody-related

immune responses in solid cancers are better than in haematological cancers[48]. It is therefore

notable that in an immunologically diverse cohort of individuals, with varied vaccine responses,

we did not observe any effect of the COVID-19 vaccination on the gut microbiome, indicating

stability irrespective of pre-dose composition.

While we did not observe changes at any taxonomic level or functional capacity, we cannot

rule out genetic changes at mutational levels that may alter the microbiota function. An

independent functional validation such as metabolomics to look for bacterial derived short-

chain fatty acids, tryptophan and bile acid metabolites known to mediate microbiome-host

interactions[49] could be utilised to assess this.

Sampling from the PID cohort was limited reflecting the rarity of the individuals within the

general population. Despite this, we were able to recruit patients characterised by monogenic

defects in both intrinsic and extrinsic B-cell aetiologies, thus representing multiple facets within

the rare PID population. Further, we were unable to sample all patients at all vaccine

timepoints across all vaccine doses. Nevertheless, our findings still bare relevance as we

assess patient cohorts individually and where possible the paired sample data analysis aligns

with the overall findings.

Although the global vaccination efforts have controlled the spread of the SARS-CoV-2 virus,

there were still reported to be 2.6 million new cases within the past month[1], highlighting that

prevention of disease through vaccination is still relevant for public health. Considering the

measurable impact of common life factors such as alcohol consumption, meat intake and

commonly used medications on the microbiota[49], our study finds that the vaccination has

negligible, if any, impact on microbiome-mediated processes. The contrast is even starker

when considering large microbiome changes have been reported for COVID-19 infection[10].

Our findings indicates that the gut microbiome remains stable post-vaccination and provides

an additional reassurance towards promoting vaccine uptake.

Data availability

Sequencing data will be uploaded to the European Genome-phenome Archive, whilst code

will be available on GitHub: https://github.com/RHBoston/COVID-19_Vaccination_GM.

Acknowledgements

We are grateful to the clinical staff at Addenbrookes hospital who facilitated patient recruitment

and sample collection, and the members of the Thaventhiran and Patil lab for help in

processing the samples.

Author Information

Consortia CITIID-NIHR COVID−19 BioResource Collaboration

Juan Carlos Yam-Puc, Zhaleh Hosseini, Emily C. Horner, Nonantzin Beristain-Covarrubias,

Robert Hughes, Maria Rust, Rebecca H. Boston, Lucy H. Booth, Edward Simmons-Rosello,

Magda Ali, Anne Elmer, Caroline Saunders, Areti Bermperi, Sherly Jose, Nathalie Kingston,

Thomas E. Mulroney, Sarah Spencer, Nicholas J. Matheson & James E. D. Thaventhiran.

Contributions

R.H.B performed sample collection and performed data analysis and interpretation of results.

R.G., S.B., and L.K. performed data analysis and interpretation of results. E.C.H., N.B-C.,

J.C.Y-C., P.P.G., L.F., A.K., A.E.L., and S.B., performed sample collection and experimental

assays, as well as providing technical support. A.C.-N., A.E., C.S., A.B., S.J., N.K., S.G., E.S.

recruitment of participants of the study and sample collection. M.S.B., C.P., S.L., N.J.M.,

J.E.D.T. provided oversight on recruitment of participants, sample collection and experiments.

V.B. provided technical support. R.H.B., R.G., J.E.D.T., and K.R.P prepared the figures and

wrote the manuscript. R.H.B., J.E.D.T., and K.R.P designed and supervised the study.

J.E.D.T., and K.R.P contributed equally.

Competing interests

The authors declare no competing interests.

Funding declaration

This work was funded by the UK Medical Research Council (project numbers

MC_UU_00025/11 and MC_UU_00025/12).

Table 1. Characteristics of the participants in this study.

Cohort	Participants	Age	Vaccine	Condition	Treatment
Healthy Controls	6F 9M	28–59 Mean = 43.7	Vaccine doses, n = 20 90% Pfizer doses 10% Moderna doses	NA	NA
Immune checkpoint treated cancer patients (ICP)	9F 26M	39–86 Mean = 61.7	Vaccine doses, n = 70 97% Pfizer doses 3% Moderna doses	11 Metastatic Melanoma, 10 Adjuvant Melanoma, 5 Melanoma controls, 6 Metastatic Renal, 3 Renal controls	3 Nivolumab, 13 Pembrolizumab, 10 Ipilimumab + Nivolumab, 1 Ipilimumab + Pembrolizumab
Primary Immunodeficient patients (PID)	4F 5M	19–61 Mean = 41.1	Vaccine doses, n = 19 95% Pfizer doses 5% AstraZeneca doses	1 CD40L deficiency, 2 CTLA4 deficiency, 4 NFKB1 deficiency, 2 Undiagnosed condition	5 intravenous immunoglobulin, 3 Antibiotics

Figure 1. *The composition of the gut microbiome remains unaltered following COVID-19 vaccination.*

(A) 59 patients were recruited for longitudinal analysis of the effect of the vaccines against COVID-19. Samples were assigned to one of three patient cohorts, healthy control (HC), immune-checkpoint therapy treated cancer patients (ICP), or patients with primary immunodeficiencies (PID). Blood samples were analysed for their live-virus neutralisation capacity and quantifying the amount of anti-spike IgG antibodies, whilst fecal samples were analysed with shotgun metagenomics for taxonomic and functional annotations. (B) Diversity measures of chao1 and Shannon assessed in fecal samples taken from different vaccine timepoints, from within healthy control (HC), immune-checkpoint therapy treated cancer patients (ICP) and patients with primary immunodeficiencies (PID). Statistical testing performed using Wilcoxon test and adjusted for multiple testing using FDR correction. (C) Principal component (PC) analysis at the operational taxonomic unit level. Each dot represents a unique sample from within each cohort (shapes) taken at unique timepoints following vaccination (colours). (D) Relative abundance at the phyla taxonomic level   depicted by colours of each of the bars, from samples taken from each of the cohorts (HC, ICP and PID), separated by the vaccine timepoints from which the sample was taken; PD = Pre-Dose, Acute and Late. (E) Relative abundance of the 6 most prevalent phyla in patient samples from within each of the cohorts and separated by the vaccine timepoint from which the sample was taken. Statistical testing performed using Wilcoxon test and adjusted for multiple testing using FDR correction. (N = 43 HC, 160 ICP, and 36 PID).

***Figure 2.** Bacterial species demonstrate minimal change attributable to the COVID-19 vaccines.*

(A) Differential abundance analysis using DESeq2 of relative abundance of the top 35 differential

species between samples taken at pre-dose (N = 29) and acutely (N = 69) after vaccination. (B) Log2

fold-change of the significant differential abundant species taken from the DESeq2 analysis. (C)

Relative abundance of Klebsiella pneumoniae in ICP cohort samples. *(D) Relative abundance of*

Butyrivibrio crossotus in ICP cohort samples. (E) *Relative abundance of the top 15 abundant species*

within the ICP cohort taken at each of the vaccine timepoints (N = 43 HC, 160 ICP, and 36 PID). Relative

abundance of various bacterial species correlated with immune-related diseases: Faecalibacterium

prausnitzii (F), Akkermansia muciniphila (G) and Escherichia coli (H) within patient samples taken at

each vaccine timepoint (N = 43 HC, 160 ICP, and 36 PID).

 Figure 3. *Vaccine efficacy is not correlated with the gut microbiome diversity.*

 Live-virus neutralisation (NT) capacity assessed against Shannon diversity of fecal samples, each point

 represents a different sample taken at one of the three vaccine timepoints. Colours represent cohorts,

 within healthy control (HC), immune-checkpoint therapy treated cancer patients (ICP) and patients with

 primary immunodeficiencies (PID). Correlated vaccine response through NT capacity of patient serum

 taken at second dose (N = 9 HC, 57 ICP, and 15 PID) (A) or third dose (N = 33 HC, 54 ICP, and 5 PID)

 (B). rho and p values from Spearman's Rank correlation testing displayed.

Figure 4. *Functional capacity of microbiome samples are not altered by the COVID-19 vaccines.*

(A) The relative abundance of the highest functional annotation level using the EGGNOG database

within patient samples at different vaccine timepoints in each of our patient cohorts. (B) Functional

composition depicted by colours of each of the bars, from samples taken from each of the cohorts (HC,

ICP and PID), separated by the vaccine timepoints from which the sample was taken; PD = Pre-Dose,

Acute and Late. (C) Relative abundance of the 3 most abundant functional annotations in our patient

samples from within each of our patient cohorts and separated by the vaccine timepoint from which the

sample was taken. Statistical testing performed using Wilcoxon test and adjusted for multiple testing

using FDR (N = 43 HC, 160 ICP, and 36 PID).

References

References

1.   WHO Coronavirus (COVID-19) dashboard. https://covid19.who.int.

2.   Lind, M. L. *et al.* Evidence of leaky protection following COVID-19 vaccination and SARS-
CoV-2 infection in an incarcerated population. *Nat. Commun.* **14**, 5055 (2023).

3.   Jennings, W. *et al.* Trust and vaccine hesitancy during the COVID-19 pandemic: A cross-
national analysis. *Vaccine X* **14**, 100299 (2023).

4.   Bellamkonda, N. *et al.* Immune response to SARS-CoV-2 vaccines. *Biomedicines* **10**,
1464 (2022).

5.   Bergamaschi, C. *et al.* Systemic IL-15, IFN-γ, and IP-10/CXCL10 signature associated
with effective immune response to SARS-CoV-2 in BNT162b2 mRNA vaccine recipients.
Cell Rep. **36**, 109504 (2021).

6.   Nava, P. *et al.* Interferon-gamma regulates intestinal epithelial homeostasis through
converging beta-catenin signaling pathways. *Immunity* **32**, 392–402 (2010).

7.   Pagliari, D. *et al.* The role of IL-15 in gastrointestinal diseases: a bridge between innate
and adaptive immune response. *Cytokine Growth Factor Rev.* **24**, 455–466 (2013).

8.   Wu, S. *et al.* Interleukin-6 absence triggers intestinal microbiota dysbiosis and mucosal
immunity in mice. *Cytokine* **153**, 155841 (2022).

9.   Röltgen, K. *et al.* Immune imprinting, breadth of variant recognition, and germinal center
response in human SARS-CoV-2 infection and vaccination. *Cell* **185**, 1025-1040.e14
(2022).

10. Rocchi, G. *et al.* Gut Microbiota and COVID-19: Potential implications for disease severity.
Pathogens **11**, (2022).

11. Leung, J. S. M. Interaction between gut microbiota and COVID-19 and its vaccines. *World*
J. Gastroenterol. **28**, 5801–5806 (2022).

12. Ng, S. C. *et al.* Gut microbiota composition is associated with SARS-CoV-2 vaccine
immunogenicity and adverse events. *Gut* **71**, 1106–1116 (2022).

13. Alexander, J. L. *et al.* The gut microbiota and metabolome are associated with diminished

COVID-19 vaccine-induced antibody responses in immunosuppressed inflammatory

bowel disease patients. *EBioMedicine* **88**, 104430 (2023).

14. Hirota, M. *et al.* Human immune and gut microbial parameters associated with inter-

individual variations in COVID-19 mRNA vaccine-induced immunity. *Commun. Biol.* **6**, 368

(2023).

15. Cantu, V. A., Sadural, J. & Edwards, R. PRINSEQ++, a multi-threaded tool for fast and

efficient quality control and preprocessing of sequencing datasets. *PeerJ* (2019)

doi:10.7287/peerj.preprints.27553.

16. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods*

9, 357–359 (2012).

17. Ruscheweyh, H.-J. *et al.* Cultivation-independent genomes greatly expand taxonomic-

profiling capabilities of mOTUs across various environments. *Microbiome* **10**, 212 (2022).

18. Nurk, S., Meleshko, D., Korobeynikov, A. & Pevzner, P. A. metaSPAdes: a new versatile

metagenomic assembler. *Genome Res.* **27**, 824–834 (2017).

19. Cantalapiedra, C. P., Hernández-Plaza, A., Letunic, I., Bork, P. & Huerta-Cepas, J.

eggNOG-mapper v2: Functional Annotation, Orthology Assignments, and Domain

Prediction at the Metagenomic Scale. *Mol. Biol. Evol.* **38**, 5825–5829 (2021).

20. Buchfink, B., Xie, C. & Huson, D. H. Fast and sensitive protein alignment using

DIAMOND. *Nat. Methods* **12**, 59–60 (2015).

21. McMurdie, P. J. & Holmes, S. phyloseq: an R package for reproducible interactive analysis

and graphics of microbiome census data. *PLoS One* **8**, e61217 (2013).

22. Microbiome@GitHub. http://microbiome.github.io.

23. *vegan: R package for community ecologists: popular ordination methods, ecological null*

models & diversity analysis. (Github).

24. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

25. Gerber, P. P. *et al.* A protease-activatable luminescent biosensor and reporter cell line for

authentic SARS-CoV-2 infection. *PLoS Pathog.* **18**, e1010265 (2022).

26. Xiong, X. *et al.* A thermostable, closed SARS-CoV-2 spike protein trimer. *Nat. Struct. Mol.*

Biol. **27**, 934–941 (2020).

27. Collier, D. A. *et al.* Age-related immune response heterogeneity to SARS-CoV-2 vaccine

BNT162b2. *Nature* **596**, 417–422 (2021).

28. Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting linear mixed-effects models

Usinglme4. *J. Stat. Softw.* **67**, (2015).

29. Islam, M. Z. *et al.* Reproducible and opposing gut microbiome signatures distinguish

autoimmune diseases and cancers: a systematic review and meta-analysis. *Microbiome*

10, 218 (2022).

30. Polk, D. B. & Peek, R. M., Jr. Helicobacter pylori: gastric cancer and beyond. *Nat. Rev.*

Cancer **10**, 403–414 (2010).

31. Aoun, A., Darwish, F. & Hamod, N. The influence of the gut microbiome on obesity in

adults and the role of probiotics, prebiotics, and synbiotics for weight loss. *Prev. Nutr.*

Food Sci. **25**, 113–123 (2020).

32. Shi, Z. *et al.* Emerging roles of the gut microbiota in cancer immunotherapy. *Front.*

Immunol. **14**, 1139821 (2023).

33. Bairamian, D. *et al.* Microbiota in neuroinflammation and synaptic dysfunction: a focus on

Alzheimer's disease. *Mol. Neurodegener.* **17**, 19 (2022).

34. Qiu, P. *et al.* The gut Microbiota in inflammatory bowel disease. *Front. Cell. Infect.*

Microbiol. **12**, 733992 (2022).

35. Johnson, A. J. *et al.* Daily sampling reveals personalized diet-microbiome associations in

humans. *Cell Host Microbe* **25**, 789-802.e5 (2019).

36. Hasan, N. & Yang, H. Factors affecting the composition of the gut microbiota, and its

modulation. *PeerJ* **7**, e7502 (2019).

37. Vujkovic-Cvijin, I. *et al.* Host variables confound gut microbiota studies of human disease.

Nature **587**, 448–454 (2020).

38. Kurilshikov, A. *et al.* Large-scale association analyses identify host factors influencing
human gut microbiome composition. *Nat. Genet.* **53**, 156–165 (2021).

39. Castagnoli, R. *et al.* Gut Microbiota-host interactions in inborn errors of immunity. *Int. J.*
Mol. Sci. **22**, 1416 (2021).

40. Ho, H.-E., Radigan, L., Bongers, G., El-Shamy, A. & Cunningham-Rundles, C. Circulating
bioactive bacterial DNA is associated with immune activation and complications in
common variable immunodeficiency. *JCI Insight* **6**, (2021).

41. Routy, B. *et al.* Gut microbiome influences efficacy of PD-1–based immunotherapy
against epithelial tumors. *Science* **359**, 91–97 (2018).

42. Matson, V. *et al.* The commensal microbiome is associated with anti–PD-1 efficacy in
metastatic melanoma patients. *Science* **359**, 104–108 (2018).

43. Frankel, A. E. *et al.* Metagenomic shotgun sequencing and unbiased metabolomic
profiling identify specific human gut Microbiota and metabolites associated with immune
checkpoint therapy efficacy in melanoma patients. *Neoplasia* **19**, 848–855 (2017).

44. Derosa, L. *et al.* Gut bacteria composition drives primary resistance to cancer
immunotherapy in renal cell carcinoma patients. *Eur. Urol.* **78**, 195–206 (2020).

45. Salgia, N. J. *et al.* Stool microbiome profiling of patients with metastatic renal cell
carcinoma receiving anti-PD-1 immune checkpoint inhibitors. *Eur. Urol.* **78**, 498–502
(2020).

46. Yang, Q. *et al.* A review of gut Microbiota-derived metabolites in tumor progression and
cancer therapy. *Adv. Sci. (Weinh.)* **10**, e2207366 (2023).

47. Maier, L. *et al.* Extensive impact of non-antibiotic drugs on human gut bacteria. *Nature*
555, 623–628 (2018).

48. Hiam-Galvez, K. J., Allen, B. M. & Spitzer, M. H. Systemic immunity in cancer. *Nat. Rev.*
Cancer **21**, 345–359 (2021).

49. Lindell, A. E., Zimmermann-Kogadeeva, M. & Patil, K. R. Multimodal interactions of drugs,
natural compounds and pollutants with the gut microbiota. *Nat. Rev. Microbiol.* (2022)
doi:10.1038/s41579-022-00681-5.