Gut microbiome remains stable following COVID-19 vaccination in healthy and immuno-compromised individuals

Rebecca H. Boston[1], Rui Guan[1], Sina Beier[1], Lajos Kalmar[1], Emily C. Horner[1], Nonantzin Beristain-Covarrubias[1], Juan Carlos Yam-Puc[1], Pehuén Pereyra Gerber[2,3], Luisa Faria[1], Anna Kuroshchenkova[1], Anna E. Lindell[1], Sonja Blasche[1], Andrea Correa-Noguera[4], Anne Elmer[5], Caroline Saunders[5], Areti Bermperi[5], Sherly Jose[5], Nathalie Kingston[6], CITIID-NIHR COVID-19 BioResource Collaboration, Sofia Grigoriadou[7], Emily Staples[1], Matthew S. Buckland[7,8], Christine Parkinson[4], Sara Lear[4], Nicholas J. Matheson[2,3,4,9], Vladimir Benes[10], James E. D. Thaventhiran[1,4,*] and Kiran R. Patil[1,*].

[1] Medical Research Council Toxicology Unit, University of Cambridge, Cambridge, UK.
[2] Cambridge Institute of Therapeutic Immunology and Infectious Disease (CITIID), University of Cambridge, Cambridge, UK
[3] Department of Medicine, University of Cambridge, Cambridge, UK
[4] Department of Clinical Immunology, Cambridge University NHS Hospitals Foundation Trust, Cambridge, UK
[5] NIHR Cambridge Clinical Research Facility, Cambridge, UK
[6] NIHR BioResource, Cambridge University Hospitals NHS Foundation Trust, Cambridge, UK
[7] Department of Clinical Immunology, Barts Health, London, UK
[8] UCL GOSH Institute of Child Health Division of Infection and Immunity, Section of Cellular and Molecular Immunology, London, UK
[9] NHS Blood and Transplant, Cambridge, UK
[10] European Molecular Biology Laboratory, Heidelberg, Germany
*Correspondence to: JEDT, jedt2@mrc-tox.cam.ac.uk; KRP, kp533@mrc-tox.cam.ac.uk.

Abstract

The bidirectional interaction between the immune system and the gut microbiota is a key contributor to various host physiological functions. Immune-associated diseases such as cancer and autoimmunity, as well as the efficacy of immunomodulatory therapies, have been linked to microbiome variation. Here, we investigate the temporal impact of COVID-19 vaccination on the gut microbiome in healthy and immuno-compromised individuals; the latter included patients with primary immunodeficiency and cancer patients on immunosuppressive therapy. We find that the gut microbiome, assessed using shotgun metagenomics, remained stable post-vaccination irrespective of diverse immune status, vaccine response, and microbial composition spanned by the cohort. The stability is evident at all tested levels including phylum, species, and functional capacity. Our results show the resilience of the gut microbiome to host immune changes triggered by COVID-19 vaccination and suggest minimal, if any, impact on microbiome-mediated processes.

Introduction

As of 9th July 2023, there have been reported almost 800 million cases of COVID-19 infection and approaching 7 million related deaths globally[1]. To reduce this severity, vaccines were deployed with the aim of promoting anti-SARS-CoV-2 immunity, with almost 13.5 billion vaccine doses administered globally, 150million of which administered in the United Kingdom[1].

The SARS-CoV-2 mRNA and viral vector vaccines induce a strong immune response through
the promotion of spike mRNA translation in host cells, resulting in the expression of spike
protein on their cell surface. The spike protein is recognised by either antigen-presenting cells
(APCs), which recognise spike antigens and are activated, inducing the migration to draining
lymph nodes where these cells can present spike antigens to T- and B-cells resulting in
antibody production, or by B-cells themselves with the cognate receptor for the spike protein,
ultimately both resulting in antibody production[2]. Although sequestration of the spike mRNA
by antigen-presenting cells aims to limit the spread into systemic circulation, SARS-CoV-2
spike protein can also be found in blood plasma[3], potentially leading to inflammation at
different sites of the body other than the site of vaccination.

Although numerous studies have analysed the gut microbiome during COVID-19 infection,
since the initiation of the vaccination programme against COVID-19, comparably few studies
have addressed the impact of the vaccination on the gut microbiome[5]. The previous work has
addressed the link between the gut microbiome and vaccine immunogenicity[6,7,8]. Primarily,
baseline abundances of species before the first vaccine dose have been correlated with a
defined end point of vaccine efficacy, typically a vaccine-related readout, such as virus
neutralisation or spike-specific antibody titres.

Yet, an open question is how the gut microbiome is affected by COVID-19 vaccination in the
first few days following vaccination, as well as elucidating whether any immediate changes in
the gut microbiome are maintained or resolved over time. Addressing this knowledge-gap
could help understanding the extent and the nature of reciprocal links between the gut
microbiome and systemic immunity in the context of vaccination. We therefore sought to
analyse the gut microbiome of patients receiving doses of the COVID-19 mRNA vaccine to
decipher whether there were any notable, characteristic changes in the gut microbiome in
either healthy or immunocompromised individuals [Table 1]; the gut microbiomes in these
cohorts are often characterised by distinct compositions when compared to healthy controls.

The immunocompromised patients recruited were either cancer patients receiving immune
checkpoint modulating therapies (ICP) or patients with primary immunodeficiencies (PID).
These cohorts thus consist of patients with impaired immunity resulting from either therapeutic
intervention for the ICP cohort, or patients with inborn errors of immunity with monogenic
defects in immunoregulatory genes for the PID cohort. This therefore also presented us with
the unique opportunity to elucidate whether the COVD-19 vaccines alter the gut microbiome
in the absence of complete, functional immunity and subsequent impairments of the regulation
of the gut microbiome.

Methods

Patient recruitment and ethics

Patients were enrolled and placed into one of 3 cohorts, healthy controls (HCs), cancer receiving either monotherapy or combination therapy of immune checkpoint inhibitors (ICP), or PID patients with defined mutations in key immunoregulatory genes (CTLA4, NFKB1, CD40L) as well as patients with clinical presentation aligning to that of defined PID, including antibody deficiency, termed as undefined PID [Table 1]. The research was conducted in accordance with the principles of Good Clinical Practice and following approved protocols of the NIHR National Bioresource. Samples were collected with the written informed consent of all study participants under the NIHR National BioResource - Research Tissue Bank (NBR-RTB) ethics (REC:17/EE/0025) or under the Barts and the London Immunology Registry (REC:361 11/LO/1689). The patients and healthy controls were consented under the East of England Cambridge South national research ethics committee (REC) reference 13/EE/0325 or Barts and the London Immunology Registry (REC: 11/LO/1689).

Patient sampling

Peripheral blood and fecal samples for were collected longitudinally over the course of 3 doses of the vaccines against COVID-19 (sample and patient coverage varied across doses). Peripheral blood mononuclear cells (PBMCs) were extracted from blood samples using density gradient centrifugation, stored temporarily at -80°C, before being transferred to long-term storage in liquid nitrogen. Serum was isolated from peripheral blood via centrifugation and stored at -80°C until required. Fecal samples were collected at 3 timepoints around each vaccine dose: pre-dose, acute (day 2 or 3 after vaccination) or late (day 16–28 after vaccination). Fecal matter was collected in OMNIgene•GUT kits (DNA Genotek, Canada) whereby samples are stored in a stabilizing, inactivating solution. Samples were transported to the lab and homogenized upon arrival before being stored at -80°C until required.

Serological assessment of immune response to COVID-19 vaccines

Serum samples were thawed, heat-inactivated at 56°C for 30 mins and measured for the dilution of serum that reduces viral activity by 50% (NT_{50}); the method for which has been previously described[9]. For anti-SARS-CoV-2 specific IgG antibodies, we utilised a previously described method[10,11], in which Luminex bead sets are covalently coupled to the recombinant SARS-CoV-2 proteins nucleocapsid protein (NCP), spike (S) and receptor-binding domain (RBD) to quantify antibody levels.

Fecal DNA extraction and sequencing

DNA was extracted from fecal samples using QIAamp® PowerFecal® Pro DNA kits (Qiagen). Samples were thawed and approximately 250mg of fecal sample was lysed via bead beating. According to the kit protocol, the sample was then cleaned of non-DNA organic and inorganic

material, then washed using ethanol. DNA is eluted into 10mM Tris and quantified using the Qubit™ 1X dsDNA High Sensitivity (HS) Assay Kit (Thermo Fisher Scientific, UK) using the Qubit™ fluorometer (Thermo Fisher Scientific, UK). DNA at a concentration of 10mg/uL was sent for sequencing. Shotgun metagenomic sequencing was performed using Illumina NextSeq 2000 sequencing technology using paired end reads of 150bp in length.

Shotgun metagenomic analysis

Raw sequencing data was pre-processed with PRINSEQ++[12] in paired read mode, quality trimming to a minimal PHRED score of 30 in a window of 15 bases and removing reads of less than 75 bp length after trimming. Additionally, host contamination was removed by mapping against the HG38 human database using Bowtie2[13] v2.4.5 and removing any mapped reads from the dataset. Raw, trimmed and filtered reads were checked for quality using FastQC[14]. From the remaining read pairs, taxonomic profiling was determined using mOTUs3[15] profiler, whilst for functional profiling, the remaining read pairs after filtering were assembled using metaSPAdeS[16] with a k-mer size of 55. The resulting scaffolds were filtered for at least 200 bp length and weighted by their average coverage of the filtered reads to enable quantitative analysis. The remaining scaffolds were aligned to the EGGNOG database[17] (downloaded on 2022/04/08) using DIAMOND version 2.0.13. Microbiome analysis was performed in R using phyloseq[18] and vegan[19] packages. Differential abundance analysis was performed using DESeq2[20].

Statistical analysis

Wilcoxon tests with multiple testing correction using the false discovery rate were deployed throughout. Correlation was determined using Spearman rank.

Results

The composition of gut microbiome is not altered by vaccination against COVID-19

To investigate the impact of the COVID-19 vaccines on the gut microbiome, shotgun metagenomic sequencing was performed on a total of 239 fecal samples from our three patient cohorts (43 HC, 160 ICP and 36 PID samples) over the course of 3 vaccine doses, at one of three vaccine timepoints pre-dose (before vaccination), acute (2–3 days after vaccination) or late (16–28 days after vaccination) for each vaccine dose [Fig1A].

Samples taken at different vaccine timepoints from within each cohort were not significantly
different [Figure 1B], however we noted significant diversity differences between the cohorts
[Supplementary figure 1A]. This indicates that despite the distinct microbial diversity between
the patient cohorts, the COVID-19 vaccine is not affecting the diversity of the gut microbiome.

We next used principal component analysis to reduce the dimensionality of our microbiome
composition data (starting with a very high dimension dataset including the abundance of all
detected operational taxonomic units (OTUs)) [Fig1C]. The first 5 principal components that
were responsible for the 2.9%, 2.4%, 2.1%, 2.0% and 1.9% of variation in the data
respectively, were further analysed by using linear models with multiple input variables from
our available metadata. The principal components describing the largest variation components
in the dataset representing shifts in microbiome composition, potentially reflecting to the
parallel abundance changes of multiple bacterial species between the samples. In our linear
models we asked if either the cohort and/or the vaccination timepoint association of the
samples have effect(s) on the value of the principal component. We found that the cohort
grouping has significant effect on PC1, PC2, PC3 and PC5 while none of these major principal
components were dependent on the timepoints in which the samples were taken. This
suggests that the variance we see in our samples is not a signature of the COVID-19 vaccines,
rather those of the unique cohorts presenting with different microbiome compositions.

Next, we asked whether the COVID-19 vaccines induce any changes in the phylum-level
composition of the gut microbiome, we observed the relative abundance of taxa at a phylum
level across all samples [Fig1D]. There were no significant differences observed when
comparing samples at the three vaccine timepoints within each of our three cohorts in the top
6 most prevalent phyla [Fig1E], this is despite significant differences in these phyla between
cohorts [Supplementary figure 1B]. This demonstrates that the COVID-19 vaccines do not
alter the composition of the gut microbiome irrespective of the unique compositions found in
our cohort samples.

COVID-19 vaccination does not induce species level changes in the gut microbiome

We next sought to analyse differentially abundant microbial species between vaccine
timepoints using DESEq2. All cohorts were analysed independently for the abundance
changes in samples taken at each vaccine timepoint with the top 35 differentially abundant
species presented in a representative heatmap of log2 fold-change in abundance. The ICP
cohort demonstrates that when assessing the change in species abundance between samples
taken pre-dose and acutely, unsupervised clustering does not demonstrate clustering based
on the top 35 bacterial species [Fig2A]. Two species that were significantly increased in acute

samples compared to pre-dose samples [Fig2B], *Klebsiella pneumoniae* and *Butyrivibrio*
crossotus found in 11 and 5 samples [p = 1.01e-24 and 7.63e-12 respectively]. The former is
only representative in a quarter of the cohort, only melanoma patients, and within those has
an average relative abundance of 0.7% [Figure 2C]; the latter in 2 renal cancer patients,
representing on average 3% of the relative abundance. This indicates that we see minimal
changes in the relative abundance of bacterial species within the patient samples considering
over 2600 species are represented across all our patient samples, and that the changes we
do see are attributable to species that are not represented by the entire cohort and that
represent a fraction of the microbial niche. Similar findings were seen for our other two cohorts,
HC and PID [Supplementary figure 2], indicating that on a species level there is no unified,
biologically relevant change in abundance of microbial species induced by the COVID-19
vaccines.

As the differential abundance analysis tool DESEq2 considers the change in abundance of all
species irrespective of their relative abundance within each sample, we were curious whether
there were any noticeable changes, falling below DESEq2 threshold, in the abundance of the
top 15 most abundant species found within each patient cohort that could be attributed to the
vaccine timepoints. There was no significant difference in any of the 15 species in samples
taken at any of the three vaccine timepoints [Figure 2D]. This indicates that we see no effect
of the vaccine on the species occupying the highest proportion of the abundant microbial
species. Thus, on a species level there is no unified, biologically relevant change in abundance
of microbial species induced by the COVID-19 vaccines.

There was considerable concern both at the time of the vaccine programme, and to this day,
on the safety of COVID-19 vaccines. Given that, we sought to observe if there was any
differential outgrowth of species that have been associated to various immune-related
diseases, including gastric cancer and autoimmunity, as well as metabolic and neurological
diseases. When observing the abundance of *Faecalibacterium prausnitzii* found to be reduced
in both gastric cancers, autoimmunity, and Crohn's disease[21], we found no significant
abundance changes at the vaccine timepoints in our cohorts [Fig2E]. We also found no
presence of *Helicobacter pylori*, which is strongly associated with the initiation and
development of gastric cancers[22]. *Akkermansia muciniphila,* found to be increased in
obesity[23], was not altered [Fig2F]. In studies of Alzheimer's, *Escherichia coli* has been
demonstrated to promote neurodegeneration[24], in our samples we did not see significant
difference induced by the vaccine in any of our cohorts [Fig2G]. This supports that the COVID-
19 vaccines does not promote the change in abundance of microbes that are negatively
associated with various immune-related diseases from within our patient cohorts.

The magnitude of the response to the BNT162b2 vaccine is not correlated with microbiome diversity

Within the current literature, a few studies have reported changes in the gut microbiome that correlate with vaccine efficacy[5,6,7], so we sought to determine whether the gut microbiome composition was influenced by the magnitude of the BNT162b2 vaccine response. We performed a gold-tier assessment of the vaccine response using a live-virus neutralisation assay to assess whether patients with great vaccine efficacy had any correlation with the microbial diversity. When taking the neutralising capacity (NT50) of patient serum at both second dose [Fig3A] and third dose [Fig3B], we asked whether the Shannon diversity at different vaccine timepoints, correlated with neutralisation. We did not see any correlation of diversity with vaccine efficacy in any of our patient cohorts; the same is true for the quantity of anti-spike IgG antibodies [Supplementary figure 3]. This indicates that the magnitude of the immune response was not correlated with the gut microbiome diversity in our patient cohorts.

The microbiome gene functional annotations were not affected by COVID-19 vaccines

Having investigated the composition and relative abundance of the microbial species that constitute the gut microbiome, we next sought to investigate whether the functional capacity of the microbial species was altered by the COVID-19 vaccines. Using the EGGNOG database, we assigned functional annotations to the sequenced metagenomes. The highest level of functional annotation depicts three functional groups, cellular processes and signalling, information storage and processing, and metabolism. In these, we did not see any significant differences between the vaccine timepoints within our cohorts [Fig4A]; similarly to taxonomic data presented earlier, when combining samples from within the same cohorts, there are significant changes.

We next observed the abundance of the 22 defined functional groups in the next functional annotation level down in the separate vaccine timepoints within each patient cohort [Fig4B]. Representative graphs of the most abundant functional annotations within each of the highest three level functional levels remained unchanged after vaccination within our patient cohorts [Fig4C], the same is true for the remaining 19.

At the lowest functional annotation level, we interrogated the abundance of cluster of orthologous genes (COGs) at separate vaccine timepoints within our cohorts. Remarkably, only two COGs were significantly different as a result of the COVID-19 vaccines, in HC samples COG2243 Precorrin-2 c20-methyltransferase between acute and late vaccine samples (padj value = 0.04 [FDR]), and COG4750 cytidylyltransferase choline kinase between pre-dose and late samples in ICP patients (padj value = 0.03 [FDR]); not a single COG was found to be significantly altered in PID patient samples as a result of the vaccine. This

demonstrates that the functional annotations of the gut microbiome are not practically
unaltered by the administration of the COVID-19 vaccines.

Discussion

To our knowledge, this study is the first to assess the gut microbiome composition in response
to the COVID-19 vaccines across multiple doses and at multiple timepoints after each vaccine
dose, with samples take pre-dose, acutely and late following vaccination. The study is also
the first to assess the effect of the COVID-19 vaccines on the gut microbiome in cancer
patients and the rare population of individuals born with inborn errors of immunity. As sampling
across the cohorts varied throughout, we opted to combine the three vaccine doses and
assess vaccine timepoints or samples from within each cohort. This allowed us to better
observe the influence of the COVID-19 vaccines in these contexts.

The relative abundance of microbes within the gut microbiome has more recently been
assessed with vaccine immunogenicity including that of vaccines against SARS-CoV-2 virus.
The first reported study of the gut microbiome in COVID-19 vaccinated patients correlated
vaccine immunogenicity of the inactivated virus, CoronaVac, and the messenger RNA vaccine
encoding spike protein, BNT162b2 vaccine, with the baseline abundance of gut
Bifidobacterium adolescentis and *Roseburia faecis* respectively[5]. Another study investigated
the variability of the COVID-19 vaccine response to the gut microbiome by correlating RNAseq
data with microbial abundance using 16S amplicon sequencing[7]. They identified several
differentially abundant taxa between high- and low-antibody responders and high- and low-T-
cell responders. In the context of immunocompromised cohorts, a previous study assessed
patients with inflammatory bowel disease[6] well known to be characterised by gut microbiome
dysbiosis[25], who were receiving anti-TNF immunomodulators. This study did not demonstrate
changes in diversity in above geometric mean vaccine responders but found *Bilophia*
abundance correlated to an improved response. They also assessed the fecal metabolome
showing that various metabolites were associated with a better vaccine response; several
metabolites were also associated with a reduced response. What these studies have in
common is associating microbiome composition to vaccine immunogenicity, however these
studies have not addressed temporal changes that can be seen in the gut microbiome,
whereas our longitudinal analysis affords us the opportunity to do so.

In our patient cohorts, we did not find any significant effect on the diversity of the gut
microbiome following COVID-19 vaccination despite considerable differences between the
cohorts. In line with the studies investigating the gut microbiome of PID patients[26], we also
observed decreased diversity in our cohort compared to HC samples. These patients have
been demonstrated to exhibit increased gut permeability with higher rates of bacterial

translocation[27], perhaps indicative that there is bidirectional permeability of systemic immunity
affecting the gut microbiome. With regards to vaccination, PID patients are reported to respond
poorly due to their antibody deficiencies; while able to mount some form of protective antibody
responses, it is often not durable [28,29,30]. Further, in live vaccines there is evidence of induction
of severe disease[26]. In our study, we find no influence on microbiome variation after
vaccination in our patient samples.

In melanoma patients, various bacteria species from the *Actinobacteria* and *Firmicutes* phylum
have been associated with better responses to immune checkpoint blockade therapies[27,28,29]
An *Akkermansia muciniphilia* signature was also found in renal cancer patients responding
better to immune checkpoint inhibitors (ICIs)[35,36]. While cancer progression is reported to be
linked to gut microbiome composition and its derived metabolites, these associations vary
between cancer types[37]. We saw wide compositional variation within our ICP cohort samples,
perhaps due to the wide range of disease presentation and treatment included in our patients.
Our study cannot address the influence of the non-immune modulating drugs taken by the ICP
patients. The patients who undergo both chemotherapy and immunotherapy will typically be
administered a multitude of immunomodulators to aid their prevention of disease and infection.
We are not able to rule out the influence of these drugs at this time, known to affect the gut
microbiome[38]. Cancer patients' response to vaccination depend significantly on cancer type,
for example antibody-related immune responses in solid cancers are better than in
haematological cancers[39], perhaps indicative of their gut microbiome compositional
differences[40]. It is therefore notable that in such a varied cohort of individuals, we did not
observe any effect of the COVID-19 vaccination, showing stability of the gut microbiome
irrespective of pre-dose composition.

While we did not observe any changes also in the functional capacity at the metagenomic
level, further experimental validation using a metabolomics approach may be useful. Bacterial-
derived short-chain fatty acids (SCFAs), tryptophan and bile acid (BAs) metabolites have the
capacity to activate or suppress the immune system[41]. For example, SCFAs have been shown
to reduce APC activation[42], whilst BA metabolites prevent APC maturation[43] and induce B-cell
apoptosis[44] and prevent antibody production[45], both essential immune cell types within the
promotion of a vaccine response. It would therefore be of importance to correlate the predicted
functional annotations of the bacterial species found within our patient samples, with the
metabolites found with the patients' fecal samples to draw a more definitive conclusion on
whether the COVID-19 vaccines induced a biologically relevant changes in the gut
microbiome.

Considering limitations of our study, sampling from the PID cohort was limited reflecting the
rarity of the individuals within the general population. Despite this, we were able to recruit

patients from characterised by monogenic defects in both intrinsic and extrinsic B-cell
aetiologies, thus representing multiple facets within the rare PID population. We could not
account for known confounding factors like diet due to the lack of the metadata; however, this
is likely to be less critical since we observe considerable stability over an extended duration.
Further, we were unable to sample all patients at all vaccine timepoints across all vaccine
doses. Nevertheless, our findings still bare relevance as we assess patient cohorts individually
and where possible the paired sample data analysis aligns with the overall findings.

Although the global vaccination efforts have controlled the spread of the SARS-CoV-2 virus,
there were still reported to be 2.6 million new cases within the past month[1], highlighting that
management of disease through vaccination is still relevant for public health. Our findings
provide evidence that that the gut microbiome remains stable post-vaccination suggesting
minimal if any impact on microbiome-mediated processes.

Acknowledgements

335 We are grateful to the clinical staff at Addenbrookes hospital who facilitated patient
336 recruitment and sample collection, and the members of the Thaventhiran and Patil lab for
337 help in processing the samples.

Competing interests

340 The authors declare no competing interests.

References

[1] World Health Organization. Available: https://covid19.who.int [Accessed 10 July 2023].

[2] Bellamkonda N, et al. Immune Response to SARS-CoV-2 Vaccines. *Biomedicines*. 2022 Jun
21;10(7):1464.

[3] Röltgen K, et al. Immune imprinting, breadth of variant recognition, and germinal center response in
human SARS-CoV-2 infection and vaccination. *Cell*. 2022 Mar 17;185(6):1025-1040.e14.

[4] Baradaran A, et al. COVID-19 Associated Multisystem Inflammatory Syndrome: A Systematic Review
and Meta-analysis. *Iran J Allergy Asthma Immunol*. 2020 Dec 19;19(6):570-588.

[5] Leung JSM. Interaction between gut microbiota and COVID-19 and its vaccines. *World J*
Gastroenterol. 2022 Oct 28;28(40):5801-5806.

[6] Ng SC, et al. Gut microbiota composition is associated with SARS-CoV-2 vaccine immunogenicity and
adverse events. Gut. 2022 Jun;71(6):1106-1116.

[7] Alexander JL, et al. The gut microbiota and metabolome are associated with diminished COVID-19
vaccine-induced antibody responses in immunosuppressed inflammatory bowel disease patients.
EBioMedicine. 2023 Feb;88:104430.

[8] Hirota M, et al. Human immune and gut microbial parameters associated with inter-individual
variations in COVID-19 mRNA vaccine-induced immunity. *Commun Biol*. 2023 Apr 20;6(1):368.

[9] Gerber PP, et al. A protease-activatable luminescent biosensor and reporter cell line for authentic
SARS-CoV-2 infection. PLoS Pathog. 2022 Feb 10;18(2):e1010265.

[10] Xiong X, et al. A thermostable, closed SARS-CoV-2 spike protein trimer. Nat Struct Mol Biol. 2020
Oct;27(10):934-941.

[11] Collier DA, et al. Age-related immune response heterogeneity to SARS-CoV-2 vaccine BNT162b2.
Nature. 2021 Aug;596(7872):417-422.

[12] Cantu VA, Sadural J, Edwards R. PRINSEQ++, a multi-threaded tool for fast and efficient quality
control and preprocessing of sequencing datasets. PeerJ Preprints. 2019. 7:e27553v1

[13] Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods. 2012 Mar
4;9(4):357-9.

[14] Andrews, S. (2010). FastQC: A Quality Control Tool for High Throughput Sequence Data [Online].
Available online at: http://www.bioinformatics.babraham.ac.uk/projects/fastqc/

[15] Ruscheweyh, HJ., Milanese, A., Paoli, L. et al. Cultivation-independent genomes greatly expand
taxonomic-profiling capabilities of mOTUs across various environments. Microbiome 10, 212 (2022).

[16] Nurk S, et al. metaSPAdes: a new versatile metagenomic assembler. Genome Res. 2017
May;27(5):824-834.

[17] Huerta-Cepas J, et al. eggNOG 5.0: a hierarchical, functionally and phylogenetically annotated
orthology resource based on 5090 organisms and 2502 viruses. Nucleic Acids Res. 2019 Jan
8;47(D1):D309-D314.

[18] McMurdie PJ, Holmes S. phyloseq: An R Package for Reproducible Interactive Analysis and Graphics
of Microbiome Census Data. PLOS ONE (2013). 8(4): e61217.

[19] Oksanen J, et al. vegan: Community Ecology Package_. R package version 2.6-4, (2022).
<https://CRAN.R-project.org/package=vegan>.

[20] Love, M.I., Huber, W., Anders, S. Moderated estimation of fold change and dispersion for RNA-seq
data with DESeq2 Genome Biology (2014). 15(12):550

[21] Islam MZ, et al. Reproducible and opposing gut microbiome signatures distinguish autoimmune
diseases and cancers: a systematic review and meta-analysis. Microbiome. 2022 Dec 9;10(1):218.

[22] Polk DB, Peek RM Jr. Helicobacter pylori: gastric cancer and beyond. Nat Rev Cancer. 2010
Jun;10(6):403-14.

[23] Aoun A, Darwish F, Hamod N. The Influence of the Gut Microbiome on Obesity in Adults and the Role
of Probiotics, Prebiotics, and Synbiotics for Weight Loss. Prev Nutr Food Sci. 2020 Jun 30;25(2):113-
123.

[24] Bairamian D, et al. Microbiota in neuroinflammation and synaptic dysfunction: a focus on Alzheimer's
disease. Mol Neurodegener. 2022 Mar 5;17(1):19.

[25] Qiu P, et al. The Gut Microbiota in Inflammatory Bowel Disease. Front Cell Infect Microbiol. 2022 Feb
22;12:733992.

[26] Castagnoli R, et al. Gut Microbiota-Host Interactions in Inborn Errors of Immunity. Int J Mol Sci. 2021
Jan 31;22(3):1416.

[27] Ho HE, et al. Circulating bioactive bacterial DNA is associated with immune activation and
complications in common variable immunodeficiency. JCI Insight. 2021 Oct 8;6(19):e144777.

[28] Guo J, Bet al. Immunodeficiency-related vaccine-derived poliovirus (iVDPV) cases: a systematic
review and implications for polio eradication. Vaccine. 2015 Mar 3;33(10):1235-42.

[29] Yam-Puc JC, et al. Age-associated B cells predict impaired humoral immunity after COVID-19
vaccination in patients receiving immune checkpoint blockade. Nat Commun. 2023 Jun 27;14(1):3292

[30] Pham MN, et al. Immunogenicity and tolerability of COVID-19 messenger RNA vaccines in primary
immunodeficiency patients with functional B-cell defects. J Allergy Clin Immunol. 2022 Mar;149(3):907-
911.e3.

[31] Sobh A, Bonilla FA. Vaccination in Primary Immunodeficiency Disorders. J Allergy Clin Immunol
Pract. 2016 Nov-Dec;4(6):1066-1075.

[32] Routy B, et al. Gut microbiome influences efficacy of PD-1-based immunotherapy against epithelial
tumors. Science. 2018 Jan 5;359(6371):91-97.

[33] Matson V, et al. The commensal microbiome is associated with anti-PD-1 efficacy in metastatic
melanoma patients. Science. 2018 Jan 5;359(6371):104-108.

[34] Frankel AE, et al. Metagenomic Shotgun Sequencing and Unbiased Metabolomic Profiling Identify
Specific Human Gut Microbiota and Metabolites Associated with Immune Checkpoint Therapy Efficacy
in Melanoma Patients. Neoplasia. 2017 Oct;19(10):848-855.

[35] Derosa L, et al. Gut Bacteria Composition Drives Primary Resistance to Cancer Immunotherapy in
Renal Cell Carcinoma Patients. Eur Urol. 2020 Aug;78(2):195-206.

[36] Salgia NJ, et al. Stool Microbiome Profiling of Patients with Metastatic Renal Cell Carcinoma
Receiving Anti-PD-1 Immune Checkpoint Inhibitors. Eur Urol. 2020 Oct;78(4):498-502.

[37] Yang Q, et al. A Review of Gut Microbiota-Derived Metabolites in Tumor Progression and Cancer
Therapy. Adv Sci (Weinh). 2023 May;10(15):e2207366.

[38] Maier L, et al. Extensive impact of non-antibiotic drugs on human gut bacteria. *Nature*.
2018;555(7698):623-628.

[39] Hiam-Galvez KJ, Allen BM, Spitzer MH. Systemic immunity in cancer. Nat Rev Cancer. 2021
Jun;21(6):345-359. doi: 10.1038/s41568-021-00347-z.

[40] Oliva M, et al. Tumor-Associated Microbiome: Where Do We Stand? Int J Mol Sci. 2021 Feb
1;22(3):1446.

[41] Wang J, et al.. Gut-Microbiota-Derived Metabolites Maintain Gut and Systemic Immune Homeostasis.
Cells. 2023 Mar 2;12(5):793.

[42] Nastasi C,et al. The effect of short-chain fatty acids on human monocyte-derived dendritic cells. *Sci*
Rep. 2015;5:16148.

[43] Széles L, et al. 1,25-dihydroxyvitamin D3 is an autonomous regulator of the transcriptional changes
leading to a tolerogenic dendritic cell phenotype. *J Immunol*. 2009;182(4):2074-2083

[44] Lemire JM, et al. . 1 alpha,25-dihydroxyvitamin D3 suppresses proliferation and immunoglobulin
production by normal human peripheral blood mononuclear cells. *J Clin Invest*. 1984;74(2):657-661.

[45] Kim M, et al. Gut Microbial Metabolites Fuel Host Antibody Responses. *Cell Host Microbe*.
2016;20(2):202-214.

Table 1. Characteristics of the participants in this study.

Healthy Controls (HC)	
Participants	*Age*
6F 9M	28–59

Immune checkpoint treated cancer patients (ICP)			
Participants	*Age*	*Disease*	*Treatment*
9F 26M	39–86	11 Metastatic Melanoma	3 Nivolumab
		10 Adjuvant Melanoma	13 Pembrolizumab
		5 Melanoma controls	10 Ipilimumab + Nivolumab
		6 Metastic Renal	1 Ipulumumab + Pembrolizumab
		3 Renal controls	

Primary Immunodeficient patients (PID)			
Participants	*Age*	*In-born error of immunity*	*Treatment*
4F 5M	19–61	1 CD40L	5 IvIg
		2 CTLA4	3 Antibiotics
		4 NFKB1	
		2 Undiagnosed	

Figure 1. (A) Study design. 60 patients were recruited for longitudinal analysis of the effect of the
vaccines against COVID-19. Samples were assigned to one of three patient cohorts, healthy control
(HC), immune-checkpoint therapy treated cancer patients (ICP), or patients with primary
immunodeficiencies (PID). Blood samples were analysed for their live-virus neutralisation capacity and
quantifying the amount of anti-spike IgG antibodies, whilst fecal sample were analysed with shotgun
metagenomics for taxonomic and functional annotations. (B) Diversity measures of chao1 and Shannon

assessed in fecal samples taken from different vaccine timepoints, from within healthy control (HC),

immune-checkpoint therapy treated cancer patients (ICP) and patients with primary immunodeficiencies

(PID). Statistical testing performed using Wilcoxon test and adjusted for multiple testing using FDR

correction. (C) Principal component (PC) analysis at the operational taxonomic unit level. Each dot

represents a unique sample from within each cohort (shapes) taken at unique timepoints following

vaccination (colours). (D) Phylum composition depicted by colours of each of the bars, from samples

taken from each of the cohorts (HC, ICP and PID), separated by the vaccine timepoints from which the

sample was taken; PD = Pre-Dose, Acute and Late. (E) Relative abundance of the 6 most prevalent

phyla in patient samples from within each of the cohorts and separated by the vaccine timepoint from

which the sample was taken. Statistical testing performed using Wilcoxon test and adjusted for multiple

testing using FDR correction.

Figure 2. *Representative analysis of the ICP cohort demonstrates minimal change attributable to the*
COVID-19 vaccines. (A) Differential abundance analysis using DESeq2 of the log2 fold-change of the
top 35 differential species between samples taken at pre-dose and acutely after vaccination. (B) Log2
fold-change of the significant differential abundant species taken from the DESeq2 analysis. (C)
Relative abundance of Klebsiella pneumoniae in ICP cohort samples. *(D) Relative abundance of the*
top 15 abundant species within the ICP cohort taken at each of the vaccine timepoints. Relative
abundance of various bacterial species correlated with immune-related diseases: Faecalibacterium
prausnitzii (E), Akkermansia muciniphila (F) and Escherichia coli (G) within patient samples taken at
each vaccine timepoint.

Figure 3. *Vaccine efficacy is not correlated with the gut microbiome composition. Live-virus*
neutralisation (NT) capacity assessed against Shannon diversity of fecal samples, each point
represents a different sample taken at one of the three vaccine timepoints. Colours represent cohorts,
within healthy control (HC), immune-checkpoint therapy treated cancer patients (ICP) and patients with
primary immunodeficiencies (PID). Correlated vaccine response through NT capacity of patient serum
taken at second dose (A) or third dose (B). rho and p values from Spearman correlation testing
displayed.

Figure 4. *Functional annotations of metagenomics samples using EGGNOG database are not altered by the COVID-19 vaccines. (A) The relative abundance of the highest functional annotation level within patient samples at different vaccine timepoints in each of our patient cohorts. (B) Functional composition depicted by colours of each of the bars, from samples taken from each of the cohorts (HC, ICP and PID), separated by the vaccine timepoints from which the sample was taken; PD = Pre-Dose, Acute and Late. (C) Relative abundance of the 3 most abundant functional annotations in our patient samples from within each of our patient cohorts and separated by the vaccine timepoint from which the sample was taken. Statistical testing performed using Wilcoxon test and adjusted for multiple testing using FDR.*

Supplementary Figure 1. Gut microbiome compositional differences are evident between patient cohorts. (A) Alpha diversity measures of chao1 and Shannon diversity in samples from different patient cohorts*, healthy control (HC), immune-checkpoint therapy treated cancer patients (ICP), or patients with primary immunodeficiencies (PID).* Statistical testing performed using Wilcoxon test and adjusted for multiple testing using FDR. (B) Relative abundance of the 6 most prevalent phyla in our patient samples from within each of our patient cohorts and separated by the vaccine timepoint from which the sample was taken. Statistical testing performed using Wilcoxon test and adjusted for multiple testing using FDR.

Supplementary Figure 2. Differential abundance analysis of HC and PID cohort samples taken at pre-dose and acutely after COVID-19 vaccination. (A) DESeq analysis of HC cohort and PID cohort (B), along with corresponding log2foldchange analysis of significantly different bacterial species between pre-dose and acute samples in HC samples (C) and PID samples (D), colours represent different phyla.

552 **Supplementary Figure 3.** Second dose *anti-spike IgG antibody levels assessed against Shannon*

553 *diversity of fecal samples; each point represents a different sample taken at one of the three vaccine*

554 *timepoints. Colours represent cohorts, within healthy control (HC), immune-checkpoint therapy treated*

555 *cancer patients (ICP) and patients with primary immunodeficiencies (PID. rho and p values from*

556 *Spearman correlation testing displayed.*

558