The HSF–DREB–MYB transcriptional regulatory module regulates flavonol

biosynthesis and flavonoid B-ring hydroxylation in banana (*Musa acuminata*)

Jogindra Naik[1], Ruchika Rajput[1], Samar Singh[1], Ralf Stracke[2], Ashutosh Pandey[1*]

[1]National Institute of Plant Genome Research, Aruna Asaf Ali Marg, New Delhi 110067,

India

[2]Bielefeld University, Faculty of Biology, Genetics and Genomics of Plants, 33615 Bielefeld,

Germany

Short title: HSF–DREB–MYB regulates flavonol biosynthesis

[*]**Corresponding author:**

Ashutosh Pandey

National Institute of Plant Genome Research

New Delhi 110067, India

Email: ashutosh@nipgr.ac.in

Significance statement

Our study reveals that the three stress-responsive SG7 MYB factors, MaMYBFA1-3, are modest regulators of flavonol biosynthesis in banana. In contrast, the drought- and heat-responsive MaHSF11 plays a strong, positive role in this pathway, enhancing flavonol biosynthesis and flavonoid B-ring hydroxylation by upregulating key biosynthetic genes, such as *MaFLS1* and *MaF3'5'H1*, as well as *MaDREB1*, another regulator of flavonol biosynthesis in banana.

Abstract

Plant flavonols act primarily as ultraviolet radiation absorbers, reactive oxygen species scavengers, and phytoalexins, and they contribute to biotic and abiotic stress tolerance in plants. Banana (*Musa acuminata*), an herbaceous monocot and important fruit crop, accumulates flavonol derivatives in different organs, including the edible fruit pulp. Although flavonol content varies greatly in different organs, the molecular mechanisms involving transcriptional regulation of flavonol synthesis in banana are not known. Here, we characterized three SG7-R2R3 MYB transcription factors (MaMYBFA1, MaMYBFA2, and MaMYBFA3) and their upstream regulators, heat shock transcription factor (MaHSF11) and dehydration responsive element binding factor (MaDREB1), to elucidate the molecular mechanism involved in transcriptional regulation of flavonol biosynthesis in banana. MaMYBFA positively regulate *flavonol synthase* 2 (*MaFLS2)* and downregulates *MaFLS1*. We show these transcription factors to be weak regulators of flavonol synthesis. Overexpression of *MaHSF11* enhances flavonol contents, particularly that of myricetin, and promotes flavonol B-ring hydroxylation, which contributes to the diversity of flavonol derivatives. MaHSF11 directly interacts with the *MaFLS1* and *flavonoid 3', 5'-hydroxylase*1 (*MaF3'5'H1)* promoters, both *in vitro* and *in vivo*. MaHSF11 activates the expression of *MaDREB1* directly, which in turn regulates the expression of *MaMYBFA3*. Overall, our study elucidates a novel regulatory mechanism for flavonol synthesis in banana and suggests possible targets for genetic optimization to enhance nutritional value and stress responses in this globally important fruit crop.

Keywords: Flavonol biosynthesis, MYB transcription factors, ABA, banana, HSF transcription factors, transcriptional regulatory module

Introduction

Plants produce a multitude of chemical compounds in response and adaptation to external and environmental stimuli, many of which are specialized metabolites (Oh *et al.,* 2009; Mikkelsen *et al.,* 2015; Murcia *et al.,* 2017). Among these, flavonoids are ubiquitous and play vital roles in both regulation of developmental processes and defense against pathogens (Halbwirth *et al.,* 2010; Maloney *et al.,* 2014; Chen *et al.,* 2019; Yang *et al.,* 2021). Based on their structures, flavonoids are classified into flavonols, flavones, flavanones, isoflavones, anthocyanins, and proanthocyanidins (Lago *et al.,* 2014). Flavonols have a basic 16-carbon, triple-ring structure containing a 3-hydroxyflavone backbone, and the different modifications and substituents in this ring give rise to their diversity (Ko *et al.,* 2014). Besides acting as co-pigments with anthocyanins in flower petals and as reactive oxygen species (ROS) scavengers, flavonols have numerous other important functions in plants (Kitamura *et al.,* 2016; Watkins *et al.,* 2017). They absorb ultraviolet radiation (UV), protecting plants from UV-induced oxidative damage (Liang *et al.,* 2020) and also regulate root growth and lateral root development by inhibiting polar auxin transport via stabilization of PIN-FORMED (PIN) protein dimers (Tan *et al.,* 2019; Chapman *et al.,* 2021; Teale *et al.,* 2021). In addition, flavonols regulate pollen development and germination, nodule development, and symbiosis (Zhang *et al.,* 2009; Muhlemann *et al.,* 2018). Flavonols are synthesized by the flavonoid biosynthesis pathway involving the enzymes chalcone synthase (CHS), chalcone isomerase (CHI), flavanone 3-hydroxylase (F3H), flavonoid 3′ hydroxylase (F3′H), flavonoid 3′5′ hydroxylase (F3′5′H), and flavonol synthase (FLS), which are well characterized in many plant species (Nagamatsu *et al.,* 2007; Guo *et al.,* 2019; Jiang *et al.,* 2020).

The flavonol biosynthesis pathway is regulated at several levels, and its transcriptional regulation has been extensively studied (Matus *et al.,* 2009). A group of MYB transcription factors (TFs), known as subgroup 7 (SG7) R2R3-MYBs, play a major role in the regulation of flavonol biosynthesis (Schwinn *et al.,* 2016); R2 and R3 are conserved regions. MYB12, MYB111, and MYB11, also known as PRODUCTION OF FLAVONOL GLYCOSIDES1–3 (PFG1–3), respectively, modulate flavonol biosynthesis in a tissue-specific manner in Arabidopsis (*Arabidopsis thaliana*) (Mehrtens *et al.,* 2005; Stracke *et al*., 2007; Stracke *et al.,* 2010), tobacco (*Nicotiana tabacum*) (Misra *et al.,* 2010; Pandey *et al.,* 2014a, b; Pandey *et al.,* 2015a), and tomato (*Solanum lycopersicum*) (Pandey *et al.,* 2015b). They regulate the

expression of several rate-limiting biosynthesis genes, including *CHS*, *F3H*, and *FLS*. *FLS* is

also coordinately regulated by MYB99, MYB21, and MYB24 in Arabidopsis stamens and

anthers independently of the PFGs (Battat *et al.,* 2019; Zhang *et al.,* 2021). Many MYB TFs

belonging to SG7 have been reported to regulate flavonol synthesis in various species

namely VvMYBF1 (*Vitis vinifera*), CmMYB3 (*Chrysanthemum morifolium*), MtMYB134

(*Medicago truncatula*), CaMYB39 (*Cicer arietinum*), MdMYB22 (*Malus domestica*),

NtMYB12 (*Nicotiana tabacum*) are among the several factors characterized so far to regulate

the expression of *FLS* and flavonol synthesis (Czemmel *et al.,* 2009; Wang *et al.,* 2017; Song

et al., 2020; Naik *et al.,* 2021; Saxena *et al.,* 2023; Yang *et al.,* 2023). Apart from these,

AtHY5, AtARF2, AtWRKY23, NtWRKY11b, VvWRKY70, GmNAC2, MaABI5-like,

MaC2H2, PaNAC03, CsNAC086 are also among the few transcriptional regulators of

flavonol synthesis in different species including banana (Grunewald *et al.,* 2012; Dalman *et*

al., 2017; Bian *et al.,* 2020; Song *et al.,* 2020; Bhatia *et al.,* 2021; Jiang *et al.,* 2022; Song *et*

al., 2022; Song *et al.,* 2023; Wang *et al.,* 2021f; Wei *et al.,* 2023b; Song *et al.,* 2024). A

recent study highlighted rather an interesting mechanism, where a SG7 MYB ScMYB12

interacts with ScTT8 to regulate flavonol synthesis in Jojoba (Zheng *et al.,* 2024). DELLA

component (RGA and GAI) directly interacts with MYB12 and MYB111 to promote flavonol

synthesis in Arabidopsis (Tan *et al.,* 2019). Many of these characterized factors regulate

flavonol biosynthesis pathway in hormone- and stress-responsive manner.

Because plants are exposed to numerous biotic and abiotic stresses, such as excessive

light and UV radiation, suboptimal water conditions (drought and waterlogging), high salt,

extreme temperatures, nutrient deficiencies, and pathogen infection (López *et al.,* 2008;

Cramer *et al.,* 2011; Kalaji *et al.,* 2017; Choudhury *et al.,* 2017), they have evolved various

defensive mechanisms that are activated after exposure to stress and involve a complex

interplay of signaling pathways. Other classes of TFs, such as heat shock factors (HSFs) and

dehydration-responsive element-binding (DREB) factors, modulate the plant's stress

response to drought and high heat (Zhao *et al.,* 2010; Lee *et al.,* 2010; Huang *et al.,* 2016;

Zinsmeister *et al.,* 2020). HSFs, in their dimeric form, regulate several stress-responsive

genes by binding to heat shock elements (HSEs; nGAAnnTTCn/nTTCnnGAAn) via their N-

terminal DNA-binding domains (Liu *et al.,* 2008; Wang *et al.,* 2020). HSFs can be classified

into three major classes, A, B, and C, depending upon the type of amino acid residues and the

length of flexible linker length between DBD (DNA-binding domain) and OD

(oligomerization domain) (Scharf *et al.,* 2012; Manna *et al.,* 2021). In addition to heat stress

tolerance, HSFs regulate salt, drought, and cold stress responses in Arabidopsis (Yoshida *et al.,* 2011). MdHSFA8a is a positive regulator of drought stress tolerance and flavonoid synthesis in apple (*Malus domestica*) (Wang *et al.,* 2020). GmHSFB2b positively regulate flavonol synthesis in soybean through up-regulation of *GmCHS* and down-regulation of *GmNAC2*, a repressor of *GmFLS1* and *GmFLS2* under salt stress (Bian *et al.,* 2020). Apart from these, The HSFA2 and HSFA3 heterodimeric complex confers transcriptional thermal memory by promoting epigenetic H3K4 hypermethylation (Friedrich *et al.,* 2021). In banana, MaHSF24 negatively regulates cold stress response via repression of HSP and antioxidant genes, while MaHSF26 alleviates chilling injury in heat shock responsive manner via repressing MaRBOH (Si *et al.,* 2022; Si *et al.,* 2024). DREBs, which belong to the AP2/ERF family of TFs acting both upstream and downstream of HSF, are also involved in several stress signaling pathways, in particular drought, salt, heat, and cold stress (Dubouzet *et al.,* 2003; Schramm *et al.,* 2008; Chen *et al.,* 2016; Huang *et al.,* 2016). CBF/DREB TFs bind to the cold and dehydration-responsive CRT/DREB motif core element (CCGAC) present in the promoters of their target genes via the AP2/ERF DNA-binding domain (Li *et al.,* 2020; Wang *et al.,* 2021a). In context of metabolic pathways regulation, they are known to influence the expression of flavonoid related genes in different plants. For instance, in eggplant, SmCBF1/2/3 interacts with SmMYB113 and exerts additive effect on the SmMYB113 and SmTT8 mediated activation of anthocyanin synthesis (Zhou *et al.,* 2020). CitERF32/33 and CitRAV1 cooperatively promote citrus flavone synthesis (Zhao *et al.,* 2021). MdAP2-34, a DREB, modulates flavonol content in apple (Han *et al.,* 2022). In addition, DREBs regulate fruit ripening via a complex transcriptional regulatory pathway in banana (*Musa acuminata*) (Kuang *et al.,* 2017). MaCBF1 interacts with MaNAC1, a downstream target of MaICE1, to confer cold stress tolerance in banana (Shan *et al.,* 2014). Among all the abiotic stress, high temperature is also an important factor where HSFs are master downstream components in heat stress response (Liu *et al.,* 2012; Guo *et al.,* 2016; Rao *et al.,* 2021). It also poses a threat to the normal physiology of plants and greatly affects crop yield and production (Fahad *et al.,* 2017). High temperature is known to modulate flavonoid accumulation in different plant species to confer heat tress tolerance. For instance, in oriental hybrid lily, LhMYBC2 negatively regulates anthocyanin biosynthesis in high temperature-responsive manner via competition with LhMYBA1 for LhTT8 (Yang *et al.,* 2024). Likewise, an atypical SG7 R2R3 MYB CmMYB012 suppresses flavone and anthocyanin accumulation under high temperature in chrysanthemum (*Chrysanthemum x morifolium*) (Zhou *et al.,* 2021). Whereas in *C. sinensis*, heat induced CsHSFA1b and CsHSFA2 activates CsJAZ6, which interacts

with CsEGL3 and CsTTG1 to inhibit catechin synthesis under high temperature (Zhang *et al.,* 2023). Theses finding indicates decreased accumulation of anthocyanin and proanthocyanidin under heat stress. On the contrary, high temperature promotes flavonol biosynthesis (Martinez *et al.,* 2016; Fragkostefanakis *et al.,* 2016; Paupière *et al.,* 2017). Flavonols scavenge excessive ROS during heat stress to promote pollen viability and pollen tube growth in tomato (Muhlemann *et al.,* 2020). In tomato, heat stress significantly enhanced accumulation of flavonols content in leaves and pollen (Martinez *et al.,* 2016, Paupiere *et al.,* 2017).

In terms of global production and monetary value, banana is the fourth most important food crop in the world, after rice (*Oryza sativa*), wheat (*Triticum aestivum*), and maize (*Zea mays*) (Wang *et al.,* 2021c). Banana is a monocarpic herbaceous crop with climacteric fruits, which is characterized by postharvest ripening because of cellular respiration and increased ethylene production. However, global gross production is often affected by stresses including drought, cold, salinity, and, most importantly, *Fusarium oxysporum* f. sp. *cubense* 4 (Foc4), which causes a devastating wilt disease (Xu *et al.,* 2010). We identified and functionally characterized three SG7 R2R3-MYB TFs (MaMYBFA1, MaMYBFA2, and MaMYBFA3) as regulators of flavonol synthesis in banana. These MaMYBFAs positively regulate *MaFLS2* and downregulate *MaFLS1*. MaHSF11 directly regulates the expression of *MaFLS1*, *MaF3'5'H1*, and *MaDREB1*. MaDREB1, in turn, regulates the expression of *MaMYBFA3*. Additionally, we identified MaHSF11 as a regulator of cytochrome P_{450} (CYP450) B-ring hydroxylase in banana, where it binds directly to the promoter of *MaF3'5'H1*. Taken together, our results elucidate a novel regulatory cascade of flavonol synthesis in the important fruit crop banana.

Results

An MaMYBFA1–3 triad modulates flavonol biosynthesis in banana

Our previous *in silico* study (Pucker *et al.,* 2020) determined that the banana genome contains 285 *R2R3-MYB* genes. Three of the encoded MaMYBs form a clade with previously characterized flavonol regulators in a phylogenetic tree (Supplemental Fig. S1a). We selected these three candidates for further study and named them *MaMYBFA1* (Ma02_g00290), *MaMYBFA2* (Ma05_g23640), and *MaMYBFA3* (Ma08_g10260), where the abbreviation FA stands for flavonol activator. A multiple protein sequence alignment identified the conserved R2R3-MYB domain and the two SG7-specific, conserved motifs: SG7-1 and SG7-2

(Supplemental Fig. S1b). The full-length coding sequences (CDSs) of *MaMYBFA1*, *MaMYBFA2*, and *MaMYBFA3* were PCR-amplified from banana cDNA and used to express the encoded proteins. The subcellular localization of MaMYBFA-YFP fusion proteins was analyzed in *Nicotiana benthamiana* epidermal cells and was restricted to the nucleus, co-localizing with the nuclear marker protein NLS-RFP (Supplemental Fig. S1c), indicating that MaMYBFA1–3 are nuclear proteins.

We used RT-qPCR to analyze the expression of these three *MaMYBFA* genes in different organs/tissues of banana variety 'Grand Naine' (Supplemental Fig. S2a). All three *MaMYBFA* genes were differentially expressed in the analyzed organs. *MaMYBFA1* was most highly expressed in young leaf followed by bract tissue, and ripe peel. *MaMYBFA2* transcript levels were highest in young leaf followed by bract and ripe peel, and *MaMYBFA3* was expressed most highly in young leaf followed by bract and pseudostem. Transcript accumulation of the three genes was lowest in ripe pulp. These expression patterns suggest functional differences between the three MaMYBFAs in terms of upstream regulation and downstream regulatory potential toward biosynthesis genes. We also analyzed the expression level of *MaFLS1* and *MaFLS2*, the two probable targets of MaMYBFAs. The expression level of *MaFLS1* was highest in bract and young leaf, while the pseudostem, unripe and ripe peel, and unripe pulp had lower expression level (Supplemental Fig. S2b). The root and ripe pulp had least expressions among all the tissues. On the other hand, the expression level of *MaFLS2* was considerably higher in bract tissue only (Supplemental Fig. S2b). All other tissues, however, had very lower transcript level of *MaFLS2* with almost negligible expression in unripe peel (Supplemental Fig. S2b). Finally, we quantified different flavonol aglycones to study the differential tissue specific accumulation pattern. We found that kaempferol content was highest in leaf, unripe peel, and pulp. Likewise, quercetin content was highest in bract followed by other tissues, while myricetin content was higher in peel, bract and pseudostem tissue (Supplemental Fig. S2c). Banana ripe pulp tissue, however, accumulate very least amount of all the flavonol aglycones. This varying accumulation of *MYB* genes, *FLS* genes, and flavonol metabolites suggests multilayer regulatory mechanism operating to impact overall flavonol accumulation in banana.

To analyze the regulatory role of MaMYBFA1–3, we transiently overexpressed the *MaMYBFA1–3* CDSs under the control of the *Zea mays ubiquitin1* promoter (*proZmUBI1*) in immature banana fruit discs (Fig. 1a-b). The activity of a GUS reporter simultaneously transformed on the same T-DNA in immature banana fruit discs indicated successful

transformation of the fruit tissues (Fig. 1b). RT-qPCR analysis revealed upregulation of the respective *MaMYBFA* transcripts in transgenic tissue samples (Fig. 1c). The expression of selected flavonoid biosynthesis genes was analyzed in MaMYBFA-overexpressing (OE) fruit discs. The expression of *CHALCONE SYNTHASE* (*MaCHS1–6*), *CHALCONE ISOMERASE* (*MaCHI1–2*), *FLAVANONE 3-HYDROXYLASE* (*MaF3H1–2*), and *FLAVONOL SYNTHASE* (*MaFLS1–2*) was modulated to different degrees in the OE fruit discs (Fig. 1d, Supplemental Figure S3). *MaMYBFA1* overexpression enhanced the expression of *MaCHS1*, whereas *MaMYBFA2* and *MaMYBFA3* overexpression did not (Supplemental Fig. S3). *MaCHS4* expression increased significantly with overexpression of all three *MaMYBFA*s. However, *MaCHS3* and *MaCHS5* showed enhanced expression only with overexpression of *MaMYBFA1* and *MaMYBFA3* (Supplemental Fig. S3), and only *MaMYBFA1* and *MaMYBFA2* overexpression increased the *MaCHS6* transcript level. *MaCHI* and *MaF3H* transcripts were also modulated in the transgenic OE fruit tissues: The highest activation of *MaCHI1* resulted from *MaMYBFA1* overexpression, whereas *MaCHI2* transcripts were increased 2-fold by *MaMYBFA3* overexpression (Supplemental Fig. S3). For *MaF3H1*, a significant increase in the transcript level was observed with *MaMYBFA1* and *MaMYBFA3* overexpression (Supplemental Fig. S3). Genes encoding CYP450 proteins also showed differential modulation; only *MaF3'5'H3* and *MaF3'5'H4* transcripts were upregulated in OE fruit tissues, whereas *MaF3'H* and other *MaF3'5'H* paralogs showed a marked decrease in expression (Supplemental Fig. S3). Finally, we analyzed the expression of *MaFLS1* and *MaFLS2*, the putative primary targets of MaMYBFAs, in transgenic *MaMYBFA* OE fruit discs (Fig. 1d). *MaFLS1* was significantly downregulated, whereas *MaFLS2* transcript levels increased in the *MaMYBFA* OE tissues (Fig. 1d). Overall, we found that overexpression of *MaMYBFA1* and *MaMYBFA3* significantly increased the expression of most structural flavonoid biosynthesis genes, whereas *MaMYBFA2* overexpression either decreased or negligibly affected the transcript levels of most of the genes studied.

To analyze the modulation of flavonol metabolite contents in transgenic *MaMYBFA* OE fruit disc tissues, we performed LC-MS-based, targeted flavonol quantification (Fig. 1e). We found decreased levels of kaempferol and quercetin in all the *MaMYBFA* OE tissues and an increase in myricetin content in *MaMYBFA2* OE compared to the empty vector (EV) control. Although the observed changes were significant for kaempferol, the contents of quercetin and myricetin were not altered significantly by *MaMYBFA1* and *MaMYBFA3* OE. Only *MaMYBFA2* OE significantly increased the accumulation of myricetin (Fig. 1e). Taken

together, these results indicate that MaMYBFA1, MaMYBFA2, and MaMYBFA3 act as a
MYB regulatory triad for modulating the expression of flavonol synthesis genes in banana.
Despite significant up-regulation of *MaMYBFA* genes after their overexpression in banana
fruits, flavonol content remain mostly unaltered or else there was a little change in the overall
flavonol pool (Fig. 1e). This finding can be attributed to the downregulation of *MaFLS1*,
MaF3'H and a few other genes, while increased expression of *MaFLS2* could compensate it.

MaHSF11 promotes flavonol synthesis in banana by transactivating *MaFLS1* and regulating several other biosynthesis genes

None of the banana SG7 R2R3-MYBs we examined upregulated the expression of *MaFLS1*
but instead repressed it. Because *MaFLS1* is highly expressed in leaves and is the major
enzyme for flavonol synthesis in banana (Pandey *et al.,* 2016; Busche *et al.,* 2021), we
attempted to identify a putative regulator of *MaFLS1*. An ~2-kb promoter fragment was
searched for possible TF binding sites, including ACGT-motif, E-box, LTRE (DRE/CRT),
HSE, and W-box elements, which are recognized by bZIP, MYC, DREB/CBF, HSF, and
WRKY factors, respectively (Supplemental File S1). Among these, the HSF TFs are poorly
characterized in the context of flavonoids. The *MaFLS1* promoter contained two HSEs
(nTTCnnGAAn) at positions −310 (HSE1) and −1154 (HSE2) upstream of the transcription
start site (TSS). To identify a HSF as a possible HSE-binding candidate, we searched the
publicly available banana transcriptome datasets related to abiotic stresses because both *HSF*
and *FLS* are well-known stress-responsive genes. We found a transcriptome dataset of the
drought-stressed banana cultivars Saba (ABB, drought tolerant) and Grand Naine (AAA,
drought susceptible) to be the most suitable for this analysis (Muthusamy *et al.*, 2016). From
the transcriptome deep sequencing data, we found five *HSF* genes, *MaHSF9* (*Ma08_g28350*),
MaHSF11 (*Ma06_g33440*), *MaHSF10* (Ma03_g17030), *MaHSF25* (*Ma01_g03200*), and
MaHSF26 (Ma01_g09560), that were differentially expressed before and after drought
treatment in the cultivar Grand Naine (Fig. 2a). *MaHSF25* and *MaHSF26* are class B HSFs,
which are known to repress gene expression (Scharf *et al.*, 2012). However, *MaHSF9*,
MaHSF10, and *MaHSF11* are class A HSFs, which are known to act activate gene expression
due to the AHA activator motif (Scharf *et al.*, 2012).

Using a dual-luciferase assay in *N. benthamiana* leaves, we examined whether
MaHSF9, MaHSF10, and MaHSF11 could regulate the expression of *MaFLS1* by testing
their ability to regulate *proMaFLS1–2556*. We observed a significant increase in the activity

of *proMaFLS1–2556* when co-transfected with either *MaHSF10* or *MaHSF11* compared to the EV (Fig. 2b). Between these two, MaHSF11 showed the higher *in vivo* transactivation ability (>3.6-fold, $P < 0.0001$). Therefore, we focused on MaHSF11 in further studies and tested whether MaHSF11 could directly interact with *proMaFLS1*. To verify direct binding of MaHSF11 to the HSEs in the *MaFLS1* promoter, we performed a ChIP-qPCR assay in which *proMaFLS1* was co-infiltrated with a *pro35S:YFP-MaHSF11* construct into *N. benthamiana* leaves and expressed for 3 days. An anti-GFP antibody was used to immunoprecipitate YFP-MaHSF11 and the associated chromatin. A qPCR assay with purified DNA from the precipitated chromatin showed an approximate 5-fold enrichment of fragment F1 containing HSE1; fragment F2, containing HSE2, was not enriched (Fig. 2c,d). We then performed Y1H assays to validate the direct binding of MaHSF11 to HSEs of *proMaFLS1* using 100-bp, HSE-containing promoter fragments F1 and F2 to test for interaction with the MaHSF11-AD fusion protein. The Y1H growth assay showed strong binding of MaHSF11 to F1 and a weak interaction with F2 (Fig. 2e). Furthermore, we performed an EMSA experiment, which demonstrated the direct binding of MaHSF11-GST with the promoter fragment F1 of the *MaFLS1* promoter *in vitro*. MaHSF11 could bind to the TTCTAGAA motif-containing F1 fragment, but not with F1m, which has a mutated motif (Fig. 2c,f). Taken together, our results show that MaHSF11 binds directly to the HSEs in the promoter of *MaFLS1* and activates its expression.

Phylogenetic analysis revealed clustering of MaHSF11 with AtHSFA4a, which confers salt tolerance in the MPK3/6-dependent pathway in Arabidopsis, and other A-class HSF proteins (Fig. 3a). Furthermore, a subcellular localization assay in *N. benthamiana* leaves using a pro35S:YFP-MaHSF11 construct confirmed nuclear localization of YFP-MaHSF11, which co-localized with a nuclear NLS-RFP marker (Fig. 3b). In addition to nuclear localization, we also observed fluorescence in plasma membrane, suggesting probable post-translational modification (Fig. 3b). In tissue-specific expression pattern analysis, we found that *MaHSF11* is highly expressed in roots and leaves, followed by other vegetative tissues. Meanwhile, ripe peel and ripe pulp have very low transcript levels. (Fig. 3c).

To further test the regulatory role of MaHSF11, we transiently overexpressed *proZmUBI:MaHSF11* in banana fruit discs (Fig. 3d). RT-qPCR analysis confirmed that the expression of *MaHSF11* was significantly higher (8-fold, $P \leq 0.001$) in *proZmUBI:MaHSF11* OE banana fruit disc tissues than in the EV controls (Fig. 3e). Furthermore, the expression of

MaFLS1 was significantly upregulated (3-fold, $P \leq 0.01$), and the expression of *MaFLS2* was

downregulated about 2-fold (Fig. 3e). We also observed differential expression of other

structural flavonol biosynthesis genes in *MaHSF11*-OE banana fruit tissues (Supplemental

Fig. S4a-d) as well as ~3-fold upregulation in MaMYBFA3 expression, but no significant

upregulation of the other two MYB genes (Supplemental Fig. S4e). We quantified the levels

of several different flavonol compounds in *MaHSF11*-OE fruit tissues. LC-MS-based

quantification showed increased levels of kaempferol, quercetin, and myricetin in transgenic

fruit discs (Fig. 3f). Taken together, these results suggest that MaHSF11 positively regulates

flavonol synthesis via regulation of several key biosynthesis genes, including direct

regulation of *MaFLS1*.

To further verify the regulatory role of MaHSF11 in banana, we transient expressed

pANIC8b-EV and *proZmUBI:MaHSF11*-RNAi constructs in pANIC8b in immature banana

fruit discs. GUS staining of transformed fruit discs revealed successful integration of the

expression cassette in banana (Fig. 3g). Transcript analysis revealed decreased expression of

MaHSF11 in *MaHSF11*-silenced banana fruit tissues suggesting successful gene silencing

(Fig. 3h). As expected, the expression level of *MaFLS1* decreased significantly, with the

significant increase in the expression of *MaFLS2* (Fig. 3h). Further, transcript analysis of the

other biosynthesis genes revealed insignificant changes in the expression level of many

biosynthesis genes except a few (Supplemental Fig. S5a-d). Interestingly, we found

significant increase in the transcript level of *MaMYBFA1*, but not in *MaMYBFA2* and

MaMYBFA3 genes, in *MaHSF11*-silenced tissues suggesting either direct or indirect negative

regulatory role of MaHSF11 over *MaMYBFA1* (Supplemental Fig. S5e). Promoter analysis

revealed the absence of any HSE element in the promoter of *MaMYBFA1* indicating no

probable direct interaction between the two (Supplemental File S1). Finally, metabolite

analysis showed decreased accumulation of kaempferol, quercetin and myricetin derivatives

in *MaHSF11* silenced tissues (Fig. 3i). These results further reiterate the regulatory role of

MaHSF11 in flavonol biosynthesis in banana.

MaHSF11 directly regulates *MaF3'5'H1* to promote flavonoid B-ring hydroxylation

and enhances myricetin biosynthesis

MaHSF11 overexpression led to significantly higher up-regulation of all the *MaF3'5'H* gene

family members except *MaF3'5'H7* (Fig. 4a). *MaF3'5'H1* was the most upregulated gene (a

7-fold increase), followed by *MaF3'5'H2* and other homologs; *MaF3'H* expression decreased

negligibly (Fig. 4a, Supplemental Fig. S4d). In *MaHSF11*-silenced tissues, we observed significantly decreased expression in the transcript of *MaF3'H*, *MaF3'5'H4* and *MaF3'5'H5*, while insignificant alteration was observed in the transcript level of other paralogs (Fig. 4b, Supplemental Fig. S5d).

Thorough analysis of the promoter sequences of these genes revealed the presence of one or more HSE elements in the promoters of all these genes except *MaF3'5'H7*, suggesting direct regulation by MaHSF11 (Supplemental File S1). The more than 7-fold up-regulation of *MaF3'5'H1*, (the highest among all the paralogs) prompted us to check whether it can be directly regulated by MaHSF11 at the transcriptional level. Promoter of *MaF3'5'H1* contains a single HSE (nGAAnnTTCn) at the –720-bp position relative to the TSS (Fig. 4c, Supplemental File S1). We performed Y1H assays to test the interaction of MaHSF11 with the HSE of *proMaF3'5'H1–1335* using a 100-bp, HSE-containing promoter fragment (F1), which showed a strong interaction with MaHSF11, as indicated by yeast growth (Fig. 4c,d). The ability of MaHSF11 to transactivate *proMaF3'5'H1–1335* was tested in a dual-luciferase assay in *N. benthamiana* leaves. The increased LUC activity with MaHSF11, compared to the EV control, indicated that MaHSF11 activated the *MaF3'5'H1* promoter (Fig. 4e). To further test the direct binding of MaHSF11 via the HSE, we co-infiltrated a *pro35S:YFP-MaHSF11* construct and the *proMaF3'5'H1–1335* fragment into *N. benthamiana* leaves and performed a ChIP assay using an anti-GFP antibody. This ChIP-qPCR assay revealed enrichment of the HSE-containing 100-bp promoter fragment F1 (Fig. 4f). Overall, these results show that MaHSF11 directly regulates the expression of *MaF3'5'H1*.

MaHSF11 regulates *MaMYBFA3* through MaDREB1

Overexpression and gene silencing of *MaHSF11* in banana fruit discs resulted in significant alteration in the expression level of several other structural genes involved in flavonol biosynthesis (Supplemental Fig. S4a-d, Supplemental Fig. S5a-d). Therefore, we investigated the expression of regulatory *MaMYBFA1–3* genes. As stated earlier, we found a significant upregulation of only *MaMYBFA3* in banana fruit discs overexpressing *MaHSF11* (Supplemental Fig. S4e), while no significant change was observed in *MaHSF11* silenced tissues (Supplemental Fig. S5e). A promoter analysis of the ~2-kb region upstream of the *MaMYBFA3* TSS revealed the absence of any HSEs (Supplemental File S1). Therefore, we hypothesized that MaHSF11 might indirectly regulate the expression of *MaMYBFA3* via an unknown factor. Furthermore, promoter analysis of *MaMYBFA3* revealed the presence of

several DRE/CRT (CCGAC) elements in the ~2.5-kb region upstream of the TSS (Supplemental File S1). Because HSFs bind HSEs in the promoter of *DREB* genes (Yoshida *et al.,* 2011; Huang *et al.,* 2016), we hypothesized that DREB TFs may act as mediators between *MaHSF11* and *MaMYBFA3*. To investigate the likelihood that MaHSF11–MaDREB–MaMYBFA3 is a regulatory module in banana, we measured the expression levels of *DREB* genes using the RNA-Seq data from drought-treated bananas of the Grand Naine cultivar (Muthusamy *et al.*, 2016). We found the expression of a single *DREB* gene (*Ma04_t2630*), which was also upregulated after drought stress (Fig. 5a). This gene was previously characterized as MaCBF1 (C-repeat binding factor1) and described as a critical regulator of cold stress tolerance in banana (Zhao *et al.*, 2013; Shan *et al.*, 2014). We isolated the 777-bp full-length CDS of this gene and named it *MaDREB1* due to nucleotide sequence dissimilarity with MaCBF1 (Shan *et al.*, 2014). Phylogenetic analysis revealed that MaDREB1 formed a clade with rice OsDREB3 and maize ZmDREB (Supplemental Fig. S6a). Further, subcellular localization assay in *N. benthamiana* leaves using a pro35S:YFP-MaDREB1 construct confirmed nuclear localization of YFP-MaDREB1, which co-localized with a nuclear NLS-RFP marker (Supplemental Fig. S6b). In the tissue specific transcript accumulation profiling of *MaDREB1*, we found that it followed the same distribution pattern as *MaHSF11*, with higher expression level in root, leaf, and least expression in ripe peel and pulp tissue (Supplemental Fig. S6c). To examine whether the promoter of *MaDREB1* was regulated by MaHSF11, we analyzed the region ~2 kb upstream of the TSS and identified two HSEs at positions –223 and –576 bp upstream of the TSS (Supplemental File S1). We measured the expression of *MaDREB1* in *MaHSF11* OE fruit discs and determined that it was significantly upregulated (>6-fold) (Fig. 5b). Analysis of *MaDREB1* expression in *MaHSF11* silenced fruit tissues showed insignificant alteration in the expression level of *MaDREB1* (Fig. 5c)

To verify the direct regulation of MaDREB1, we conducted a Y1H assay. We cloned two 100-bp fragments (F1 and F2) harboring HSEs (HSE1 and HSE2, respectively) of the *proMaDREB1* and conducted a Y1H assay with MaHSF11 as the effector (Fig. 5d-e). Promoter fragments carrying the HSEs were cloned into the pAbAi vector and transformed into yeast (*Saccharomyces cerevisiae*) Y1H Gold strain together with pGADT7 as a negative control. MaHSF11 in pGADT7 was transformed with both fragments, and the growth assay was performed on media supplemented with aureobasidin A (AbA). MaHSF11 could bind the HSE1-containing fragment F1 of *proMaDREB1 in vitro* (Fig. 5e). To test the transactivation

activity of MaHSF11 on *proMaDREB1*, we cloned a 998-bp promoter fragment of *MaDREB1* into the p635nRRF vector and performed dual-luciferase assays. The EV as a negative control and *MaHSF11* as the effector were individually infiltrated together with the reporter construct carrying a particular promoter fragment into *N. benthamiana* leaves, and the assay was performed 3 days post infiltration. LUC activity increased significantly when MaHSF11 was co-transfected compared to the EV control (Fig. 5f). Furthermore, *pro35S:YFP-MaHSF11* and the promoter fragment were also transformed into *N. benthamiana* leaves, and a ChIP assay was conducted. An anti-GFP antibody was used to precipitate MaHSF11-bound chromatin, and anti-IgG was used as a negative control. ChIP-qPCR showed enrichment of the F1 fragment in the immunoprecipitated chromatin, demonstrating that MaHSF11 directly bound the F1 fragment and regulated *MaDREB1* (Fig. 5g). In addition, EMSA showed direct binding of MaHSF11-GST with the GAAGCTTC motif containing promoter fragment F1 of *MaDREB1* promoter *in vitro* (Fig. 5h). Taken together, these findings strongly suggests that MaHSF11 directly binds to the promoter of *MaDREB1* to enhance its expression.

MaDREB1 directly regulates the expression of *MaMYBFA3* in banana

To test whether MaDREB1 acted as a direct transcriptional activator of *MaMYBFA3*, we isolated *proMaMYBFA3–1892*, which contains all CRT/DREs (CCGAC/GTCGG), and cloned it into pAbAi. We performed a Y1H assay with MaDREB1 as the effector in pGADT7 and EV as a control. Promoter fragments cloned into the pAbAi vector were transformed into the Y1H strain, and a growth assay was performed on AbA-supplemented media. Growth was observed in the MaDREB1 transformed strain, suggesting that it bound to the promoter fragment (Fig. 6a). To study the transactivation activity of MaDREB1, we cloned a 2508-bp fragment of *proMaMYBFA3* into p635nRRF to perform a dual-luciferase assay with MaDREB1 in pBTDest as the effector and the EV as a negative control. LUC activity increased when MaDREB1 was used as the effector compared to the control, indicating its ability to transactivate *proMaMYBFA3* (Fig. 6b). Overall, these findings show direct regulation of *MaMYBFA3* by MaDREB1. To validate the direct binding of the MaDREB1 on to the promoter of MaMYBFA3 *in vitro*, we conducted EMSA assay with four different probes (F1-F4) having different DRE/CRT motifs designed from the promoter of MaMYBFA3 (Fig. 6c). The EMSA result demonstrated the binding of MaDREB1-GST with the promoter fragment F3 having GTCGG motif, but not with promoter fragment F1, F2, and F4 of the MaMYBFA3 promoter despite having CCGAC/GTCGG motif (Fig. 6d). These

results suggests that MaDREB1 specifically binds to F3 region of the MaMYBFA3 promoter

to enhance its expression.

To verify the function of *MaDREB1 in planta*, we transiently overexpressed it in

banana fruit discs. GUS staining indicated successful transformation of the GUS expression

cassette (Fig. 7a). RT-qPCR analysis of the transformed fruit disc tissue showed high

expression (~375-fold, *P* < 0.0001) of *MaDREB1* relative to the EV control, demonstrating

successful overexpression (Fig. 7b). Expression analysis of *MaMYBFA3* in the *MaDREB1*

OE tissue showed an ~10-fold increase in the expression level (*P* < 0.05). We also found

higher expression (~21-fold, *P* < 0.001) of *MaMYBFA2*, while *MaMYBFA1* expression was

unaltered (Fig. 7c). *MaFLS1* and *MaFLS2* also showed opposite trends in expression in the

proZmUBI:MaDREB1 OE fruit tissues: *MaFLS1* was downregulated, and *MaFLS2* was

upregulated (Fig. 7c). Differential expression of other structural flavonol biosynthesis–related

genes was also observed in *proZmUBI:MaDREB1* OE banana fruit tissues (Supplemental Fig.

S7a-e). To study the modulation of flavonol content resulting from *MaDREB1*

overexpression, we used LC-MS to quantify the levels of different metabolites and found

significantly higher accumulation of kaempferol, quercetin, and myricetin (Fig. 7d). These

results indicate that MaDREB1 positively regulates flavonol biosynthesis in banana.

To further verify the regulatory role of MaDREB1 in banana, we did gene silencing of

MaDREB1 in immature banana fruit. GUS staining in pANIC8b and *proZmUBI:MaDREB1-*

RNAi expressing banana fruits revealed successful integration of the expression cassette in

banana (Fig. 7e). Expression analysis showed decreased transcript of *MaDREB1* in

MaDREB1-silenced banana disc fruit tissues suggesting successful gene silencing (Fig. 7f).

We did not see any significant changes in the expression level of *MaMYBFA2* and *FA3* (Fig.

7g). Surprisingly, the transcript level of *MaMYBFA1* increased by 4-fold suggesting that the

later might be directly or indirectly regulated by MaDREB1 (Fig. 7g). As expected, the

expression level of *MaFLS1* opposite trends as compared to overexpression, it increased

significantly. We did not observe any significant change in the transcript of *MaFLS2* (Fig.

7g). Transcript analysis of the other biosynthesis genes revealed significant increase in the

expression level of many biosynthesis genes, except few genes where no change was

observed (Supplemental Fig. S8a-e). Finally, metabolite analysis showed decreased

accumulation of kaempferol, quercetin and myricetin derivatives in *MaDREB1* silenced

tissues (Fig. 7h). Spatio-temporal metabolite accumulation is also controlled at post-

translational level apart from transcriptional regulation. These results reveled that MaDREB1

has an overall regulatory role to modulate flavonol biosynthesis in banana.

483 Heat stress positively modulates flavonol biosynthesis in banana

HSFs, as the terminal component of the heat stress response pathway, also regulate the

expression of many thermo-responsive genes (von Koskull-Döring *et al.,* 2007; Liu *et al.,*

2013). Flavonol synthesis is modulated by heat stress conditions, wherein flavonols are

known to scavenge excess ROS (Muhlemann *et al.,* 2018). Since MaHSF11 was induced

under drought stress (Fig. 2a), we wanted to check whether the expression level of *MaHSF11*

and the flavonol biosynthesis pathway are modulated upon exposure to heat stress. To do this,

we exposed three-month-old banana plantlets to 42ºC under optimal photoperiod and light

conditions, while control plants were kept at 26ºC (Fig. 8a). Expression analysis revealed

increased transcript levels of *MaHSF11* at high temperature as compared to the control plants

(Fig. 8b). Similarly, we found increased expression levels of *MaDREB1*, *MaMYBFA1*-3,

MaFLS1, and *MaFLS2* under heat stress conditions compared to control, each at different

time points (Fig. 8b). We further checked the expression of other biosynthesis genes and

found that transcript levels of a few genes up-regulated while a few genes showed down-

regulation upon heat stress (Supplemental Fig. S9). Interestingly, different paralogs of

MaF3'5'H genes showed increased transcript levels under heat stress at different time points

(Supplemental Fig. S9). Finally, metabolite analysis showed modulation in the accumulation

of kaempferol, quercetin, and myricetin derivatives under heat stress (Fig. 8c). Quercetin

content decreased at early time points, while kaempferol content fluctuated at different time

points (Fig. 8c). However, we observed a sharp increase in the accumulation of myricetin

right after the heat treatments, gradually up to 24 hours (Fig. 8c), which may be attributed to

the up-regulation of different *MaF3'5'H* paralogs at different time points of heat stress

treatment. Taken together, our results show that flavonol biosynthesis is responsive to heat

stress conditions in banana, and that *MaHSF11* might also contribute to enhanced flavonol

accumulation under heat stress in banana.

509 Discussion

510 Flavonoid biosynthesis pathways developed during the evolution of land plants. As they

511 became terrestrial, plants developed response mechanisms to adapt to and cope with several

new environmental conditions. Flavonoid biosynthesis is highly responsive to drought stress along with other biotic and abiotic stress conditions (Wang *et al.,* 2020; Wang *et al.,* 2021b; Naik *et al*., 2022; Song *et al.,* 2022; Gautam *et al.,* 2023). The role of stress signalling in the context of flavonoid biosynthesis has been studied recently, strengthening our knowledge regarding transcriptional regulation of flavonoid biosynthesis. Although flavonol biosynthesis, in particular, is positively modulated by abiotic and biotic stress, a conserved response seen throughout the plant kingdom (Albert *et al.,* 2018; Naik *et al.,* 2022; Davies *et al.,* 2023; Lu *et al.,* 2023), the molecular mechanism controlling this response has remained obscure. In this study, we identified a novel transcriptional regulatory network involving different classes of TFs and showed their interaction in regulating SG7 *MaMYBFA* and flavonol biosynthesis genes in the herbaceous fruit crop banana. We deciphered important details of the molecular mechanisms involving the transcriptional regulatory module of how HSFs, DREB, and MYB proteins orchestrate flavonol accumulation in abiotic stress dependent manner.

Initially, three candidate SG7 R2R3-MYB TFs, *MaMYBFA1–3*, were identified as having high sequence conservation in their R2R3 domains and partial conservation in the SG7-1 (GRTxRSxMK, Stracke *et al.*, 2001) and SG7-2 motifs ([W/x] [L/x]LS, Czemmel *et al.,* 2009). Furthermore, the absence of the bHLH-interacting motif ([DE]Lx2[RK]x3Lx6Lx3R, Zimmermann *et al.,* 2004) in the R3 domain suggested that these three candidate genes were independent regulators of flavonol synthesis in banana. Expression analysis of the three *MaMYBFA* genes revealed the highest level of transcript accumulation in leaf and bract tissue. *MaFLS1* transcript level was highest in bract and leaf tissues as compared to other tissues, whereas *MaFLS2* showed highest expression in bract tissue only. Different flavonol derivatives, on the contrary, are accumulated varyingly in different tissues, suggesting multi-layered regulation of the metabolic pathway. This finding indicates the presence of a regulator other than SG7 MYB in banana, acting upstream of *MaFLS1*. All these results prompted us to functionally characterize the candidate *MaMYBFA* genes in banana to determine their possible roles as flavonol-specific TFs. To do this, we transiently overexpressed the *MaMYBFA* genes in banana fruit. As raising stable transgenics in banana is time-consuming process which takes years of laborious efforts, we preferred transgene expression system in immature banana fruit as it is well established and has been used in many studies (Matsumoto *et al.,* 2009; Xiao *et al.,* 2018: Wu *et al.,* 2019; Wei *et al.,* 2023a). Although *MaFLS2* expression was activated by MaMYBFA, the expression of

MaFLS1 was downregulated. SG7 R2R3 MYB TFs directly regulates the expression of *FLS*

genes in other species. Thus, *MaFLS1* also seems a direct target of MaMYBFAs in banana. If

it is true, then later might depend on some associative or interacting partner for strong

repression. Few TFs have been reported to directly repress the expression of *FLS* genes in

other species (Zhang *et al.,* 2018; Bian *et al.,* 2020; Wang *et al.,* 2021d). It is also apparent

that binding is mediated by MaMYBFAs, as we did not find any known repression motifs in

the protein sequences of the concerned. The repression may be affected by the interacting

partner, which might be spatio-temporally regulated in banana. Few instances have also been

documented in previous studies, where *FLS2* genes are strongly repressed by transcription

regulators. In *Fagopyrum tataricum*, the expression level of *FtFLS* and rutin synthesis is

severely affected in *FtMYB13*, *FtMYB14*, *FtMYB15*, *FtMYB16* and *FtJAZ1* overexpressing

hairy roots (Zhang *et al.,* 2018). FtMYB13-16, the four SG4 R2R3 MYBs, directly interacts

with FtJAZ1, suggesting that *FtFLS1* expression is repressed by cooperative action of both

the factors. *MdBEH2*.2 directly binds to the E-box region in the promoter of *MdFLS* to

repress the expression (Wang *et al.,* 2021d). It also physically interacts with MdMYB60 and

overexpression of *MdMYB60* in apple calli leads to down-regulation of *MdFLS*. GmNAC2

directly binds to and represses the expression of *GmFLS1* and *GmFLS2* in soybean (Bian *et*

al., 2020). These reports hints at the role of probable interacting partners of MaMYBFA,

which may influence the transcriptional activity of MaMYBFA strongly. On the other hand,

DELLAs (RGA and GAI) directly interacts with MYB12 and MYB111 to enhance the

transcriptional activation activity of the later over the promoter of *FLS* in Arabidopsis (Tan *et*

al., 2019). In *Camellia sinensis*, CsMYB12 interacts physically with CsbZIP1, and both

directly binds to promoter of *CsFLS* to promote flavonol synthesis (Zhao *et al.,* 2021). In an

antagonistic way, CsMYB4 and CsMYB7 directly repressed the expression of *CsFLS*. They

also interacted with CsMYB12 to attenuate the transcriptional activation of the later (Zhao *et*

al., 2021). All these reports suggest that some strong repressors might be operating in banana

in the same way as CsMYB4 and CsMYB7. This finding could be an exception, because

none of the SG7 MYB activators downregulate flavonol synthase genes, the key genes in the

flavonol biosynthesis pathway. Flavonol content mainly remained unaltered with MaMYBFA

overexpression, a finding that can be attributed to the downregulation of *MaFLS1* and

MaF3'H and increased expression of *MaFLS2*, which could compensate for it. Thus, although

we concluded that MaMYBFAs are weak regulators of flavonol synthesis in banana, it is

difficult to ascertain their positive regulatory role on flavonol synthesis. However, our results

show the division of labor among the SG7 MYB transcriptional regulators within the

regulatory cascade. Such a mechanism has been observed in the dicot freesia (*Freesia hybrida*), in which four SG7 MYB proteins, FhMYBF1–4, differentially regulate the expression of *FhCHS* and *FhCHI* paralogs, as well as *FhFLS1*, but not *FhFLS2* (Shan *et al.,* 2020). SG7 MYB TFs regulate flavonol synthesis by directly binding to MYB binding sites (MBSs, CNGTTR) present in the promoters of flavonol biosynthesis genes, activating their expression. In apple, MdMYB22 binds to the MBS in the promoters of *MdCHS* and *MdFLS* to regulate their expression (Wang *et al.,* 2017). In freesia, FhMYB1–4 bind to the MYB core (CNGTTR) and MYBPLANT motifs (MACCWAMC), which are essential for promoter binding and transactivation (Shan *et al.,* 2022). Overall, these results suggest that MaMYBFA1–3 are the functional flavonol-specific transcriptional regulators in banana. As many early biosynthesis genes were highly up-regulated upon overexpression of MaMYBFAs, it can be speculated that the MaMYBFAs probably enhances flux towards the overall flavonoid biosynthesis pathway, rather specific flavonol branch. A recent study reported a very interesting rather unusual mechanism in Jojoba, where a flavonol regulator ScMYB12 physically interacts with ScTT8 to regulate the expression of *ScFLS* gene (Zheng *et al.,* 2024). SG7 R2R3 MYBs NtMYB12a, apart from their classical role in the regulation of flavonol synthesis, are also reported to regulate fatty acid synthesis in tobacco (Wang *et al.,* 2021e). It would be interesting to explore the impact of MaMYBFA on the regulation of other metabolic or developmental pathways in banana.

HSFs are critical components of the signaling cascades involved in responses to salt, cold, drought, and heat stress (Chauhan *et al.,* 2011; Bian *et al.,* 2020; Wang *et al.,* 2020). Several HSFs and their regulatory networks have been discovered in Arabidopsis, rice, tomato, soybean (*Glycine max*), wheat, and grape (*Vitis vinifera*) (Guo *et al.,* 2016; Wu *et al.,* 2022). In this study, MaHSF11 was identified as a regulator of *MaFLS1*. MdHSFA8a was previously identified as a regulator of *MdFLS* in apple, promoting flavonol biosynthesis and the drought stress response by regulating *MdFLS* expression (Wang *et al.,* 2020). GmHSFB2B confers salt stress tolerance and flavonol synthesis in soybean by repressing *GmNAC2*, a negative regulator of *GmFLS1* and *GmFLS2* (Bian *et al.,* 2020). Apart from these, other non-SG7 MYB regulators from other species include AtMYB21, AtMYB24, and FhMYB21L2. The presence of three HSEs in the promoter of *MaFLS1* suggested it might be regulated by a HSF. We determined the role of MaHSF11 to be a regulator of CYP450 B-ring hydroxylase in banana, where it directly bound to the promoter of *MaF3'5'H1* (Fig. 4d). Very few regulators of *F3'5'H* have been identified in plants. In poplar (*Populus*

tremula×tremuloides), MYB117 specifically regulates *F3'5'H* expression to promote anthocyanin synthesis and flavonoid B-ring hydroxylation (Ma *et al.,* 2021). MYBC1 and WRKY44 both independently and cooperatively regulate *F3'5''H* expression to regulate the anthocyanin level in kiwifruit (*Actinidia purpurea*) (Peng *et al.,* 2020).

In our study, *MaHSF11* overexpression upregulated *MaFLS1* but downregulated *MaFLS2*, indicating the presence of a complex regulatory cascade leading to preference of one paralog over the other. MaHSF11 also promoted the expression of several *MaF3'5'H* gene homologs, while slightly repressing *MaF3'H* (Supplemental Fig. S4d). Since the enzymes compete with each other for the common substrate dihydrokaempferol, MaHSF11 might be specifically promoting the myricetin synthesis branch and tries to suppresses the quercetin synthesis branch catalyzed by MaF3'H, either directly or indirectly. *MaFLS1* was highly expressed in young leaves, and this finding can be correlated with higher *MaHSF11* expression in leaves compared to fruits and roots (Wei *et al.,* 2016). Fruit tissue, particularly peel, had higher expression of *MaF3'H*, leading to a greater accumulation of rutin and quercetin in peel compared to other tissues (Pandey *et al.,* 2016). Of the seven paralogs of *MaF3'5'H* in banana, five were upregulated 2-fold with *MaHSF11* overexpression, suggesting a role for *MaHSF11* as a strong regulator of this gene family (Fig. 4a). These results further suggest that MaHSF11 promotes flavonol biosynthesis via a distinct regulatory module to regulate stress-responsive genes such as flavonol biosynthesis–related genes involved in drought, cold, and salinity stress responses in banana.

Preference of MaHSF11 for the *MaFLS1* and *MaF3'5'H* homologs (relative to *MaFLS2* and *MaF3'H*) suggested a specificity of MaFLS1 for dihydromyricetin as a substrate. In a recent study, MaFLS1 also showed specificities for dihydrokaempferol and dihydroquercetin. In future studies, it will be important to determine whether MaFLS1 has a substrate specificity for dihydromyricetin as well. CYP450 enzymes are involved in the flavonoid metabolon in both rice and soybean (Shih *et al.,* 2008; Waki *et al.,* 2016), where they act as ER membrane–bound nucleation sites for metabolon formation. Interaction among enzymes varies greatly in terms of their stability and complexity. Preferential upregulation of MaF3'5'H and MaFLS1 and downregulation of MaF3'H and MaFLS2 suggested probable metabolon formation in the ER, which can be investigated in future. In snapdragon (*Antirrhinum majus*), F3'H associates with CHI to form a metabolon (Fujino *et al.,* 2018). In Arabidopsis, FLS1 interacts with CHS via competition with DFR (dihydroflavonol 4-reductase), which can also bind to CHS (Crosby *et al.,* 2011). Negative regulation of

MaFLS2 by MaHSF11 also strongly suggests functional promiscuity of the MaFLS2 enzyme

as it has not been functional characterized. The absence of HSE element in the promoter of

MaFLS2 indicates its indirect repression by MaHSF11. In soybean, GmHSFB2b represses

the expression of *GmNAC2*, negative regulators of GmFLS1/2. So, two possible mechanisms

behind the down-regulation of MaFLS2 can be the up-regulation of any of its repressors or

down-regulation of its positive regulators by MaHSF11.

We also observed that MaHSF11 upregulated *MaMYBFA3* indirectly via MaDREB1.

Interactions of DREB and HSF TFs have been reported previously (Yoshida *et al.,*

2011;Chen *et al.,* 2016). In Arabidopsis, DREB2A is transcriptionally regulated by HSFA1a,

HSFA1b, and HSFA1d (Yoshida *et al.,* 2011), and DREB2A also acts upstream of HSFA3 to

regulate its expression (Schramm *et al.,* 2008). In our study, we found ~2-fold increased

expression of MaHSF11 when *MaDREB1* was overexpressed in banana, whereas suggesting

conservation of an interconnected regulatory cascade (Supplemental Fig. S10). We also

determined that MaHSF11 directly bound to the HSE in *proMaDREB1* to regulate its

expression. Overexpression of *MaDREB1* significantly down-regulated several biosynthesis

genes related to flavonol biosynthesis, while few biosynthesis genes were also up-regulated

notably *MaCHS4* and *MaF3'5'H3*. This antagonistic modulation in the expression level of

different biosynthesis genes leads to increased flavonol accumulation, although inadequately

(Supplemental Fig. S7a–e). While, gene silencing led to significantly up-regulation of few

genes (Supplemental Fig. S8a-e). In future work, it will be intriguing to determine whether

MaDREB1 directly represses these genes. HSF and DREB are not reported so far to interact

physically. In our Y2H analysis, we could not see any direct interaction between MaHSF11

and MaDREB1, suggesting rather conserved genetic interaction between the two regulators

(Supplemental Fig. S11). Recently, *MaCBF1* was reported to be a positive regulator of

chlorophyll catabolism: It possesses strong transactivation activity and is localized to the

nucleus (Xiao *et al*., 2023). It is also known to interact with MaNAC1 and involved in

propylene-induced cold tolerance in banana (Shan *et al.,* 2014). Overall, it may be concluded

that MaHSF11 activates the expression of *MaDREB1* to fine tune flavonol synthesis.

MaDREB1 may act, downstream of MaHSF11, to mediate drought, and cold stress response,

as both are induced (Shan *et al.,* 2014; Muthusamy *et al.,* 2016). Prior to this study, several

DREB proteins were reported to regulate different branches of flavonoid biosynthesis.

CitERF32 and CitERF33 act as transcriptional inducers of *CitCHIL1* to promote flavone and

flavanone biosynthesis in citrus (*Citrus reticulata cv. Suavissima*) (Zhao *et al.,* 2021). In

eggplant (*Solanum melongena*), SmCBF1, SmCBF2, and SmCBF3 interact with SmMYB113 to regulate *SmCHS* and *SmDFR* expression and enhance anthocyanin synthesis (Zhou *et al.,* 2020). In our study, we discovered MaDREB1 to be a regulator of flavonol synthesis, modulating the expression of regulatory SG7 MYB genes. In eggplant as well, SmCBF1, SmCBF2, and SmCBF1 are induced by drought, salt, and ABA, showing that MaDREB1 might also be involved in drought stress signaling as a transcriptional regulator of stress-responsive genes such as flavonoid biosynthesis genes under drought and heat stress. Our study provides insights into a regulatory cascade that promotes flavonol biosynthesis in banana under different stress conditions, as shown in our proposed model (Fig. 9). HSF and DREB TFs have earlier been reported to modulate flavonol, and anthocyanin synthesis in few plant species under drought, cold and salt stress (Zhou *et al.,* 2020; Bian *et al.,* 2020; Wang *et al.,* 2020). Previous studies have investigated the heat stress mediated modulation of flavonoids (Martinez *et al.,* 2016; Zhou *et al.,* 2021; Zhang *et al.,* 2023; Yang *et al.,* 2024). In our study, we explored the role of heat stress on flavonol synthesis, as HSFs are master regulators of heat stress signaling (Ohama *et al.,* 2017). We found increased expression of *MaHSF11* and its downstream factors including MaDREB1, regulatory MYB and biosynthesis genes upon heat treatment. Further, increased accumulation of flavonol derivatives, particularly myricetin, upon heat stress verified the positive modulation of flavonol synthesis in banana by heat stress signalling, which corroborates with few previous studies (Martinez *et al.,* 2016; Fragkostefanakis *et al.,* 2016). Flavonol accumulation is promoted by high temperature in vegetative tissues and pollen in tomato (Martinez *et al.,* 2016; Paupière *et al.,* 2017). Banana genome encodes seven homologs of *MaF3'5'H* (Pandey *et al.,* 2016). Among these few paralogs showed significantly increased transcript levels at early time points of 42□ heat treatment, while *MaF3'5'H4/6* transcript sharply increased at later time points. It might cumulatively contribute to increased myricetin accumulation upon heat treatment. In tomato, *HsfA2* mutant accumulates decreased transcripts of *CHI* which leads to reduced pollen viability under heat stress (Fragkostefanakis *et al.,* 2016). In a recent study, a heat shock-induced B class HSF, MaHSF26 represses the expression of *MaRBOHs*, which leads to decreased ROS and MDA (Malondialdehyde) accumulation and alleviation of chilling injury in banana (Si *et al.,* 2022). Our findings suggest rather consistent role of MaHSF11 under heat, drought, and salt stress mediated promotion of flavonol accumulation in banana (Fig. 2a, Fig. 8b, Wei *et al.,* 2016).

The banana cultivar Grand Naine (Cavendish, AAA) is highly susceptible to different abiotic stresses as well as pathogen infection. Flavonols acting as ROS scavengers and pathogen defense molecules are synthesized in increased amounts under a variety of stresses. Here, we identified the HSF–DREB-MYB transcriptional regulatory module and characterized how it works independently or cooperatively to regulate flavonol synthesis in stress-responsive manner particularly, drought and heat conditions in banana. The TFs in this regulatory module can be used for effective engineering of stress resistance in crop plants like banana, as well as flavonol fortification of banana fruit. These candidate genes could be utilized for metabolic engineering of rutin and myricetin synthesis in banana fruit. Monocots are suggested to be monophyletic arising from a common ancestor (Remizowa and Sokoloff, 2023). It suggests conservation of regulatory mechanisms of stress-mediated flavonol synthesis in other monocot crops like, rice, maize, wheat. We have, in our study, explored a novel regulatory module of stress-mediated flavanol biosynthesis in banana. Which can also be applied or studied in other crops, particularly monocots.

Materials and methods

Plant material

Banana (cultivar subgroup Cavendish [AAA] variety 'Grand Naine') plants were grown in the fields of National Institute of Plant Genome Research (NIPGR), New Delhi, India. Various tissues, including young leaf, bract, pseudostem, root, peel, and pulp, of different fruit developmental stages were sampled during winter season, frozen in liquid nitrogen, and stored at −80□ until further use. *Nicotiana benthamiana* plants were grown in a Percival AR-41L3 growth cabinet at 100 µmol m^{-2} s^{-1} light intensity and 22°C under a 16-h-light/8-h-dark photoperiod.

Identification of candidate flavonol-specific TFs from banana

To identify candidate SG7 R2R3-MYB TFs, which are known activators of flavonol biosynthesis, we retrieved sequences of previously characterized flavonol-specific TFs from various plants from NCBI (https://www.ncbi.nlm.nih.gov). These sequences were used to perform a BLASTp search against the banana genome sequence (DH-Pahang version 2) (https://banana-genome-hub.southgreen.fr). For phylogenetic studies, MEGA X (Kumar *et al.,* 2018) was used to construct a maximum-likelihood phylogenetic tree with 1000 bootstrapping rounds. For multiple sequence alignment, MaMYBFA1–3 peptide sequences were aligned with a set of similar MYBs (Stracke *et al.,* 2014) using MAFFT v.7.299b

(Katoh and Standley, 2013). Different domains and motifs were identified manually (Naik *et*

al., 2021).

Cloning of flavonol-specific transcriptional regulators

The full-length CDSs (without the stop codons) of the MaMYB-encoding genes *MaMYBFA1*

(*Ma02_g00290*), *MaMYBFA2* (*Ma05_g23640*), and *MaMYBFA3* (*Ma08_g10260*) were

amplified using first-strand cDNA of banana fruit tissue and a set of primers containing

Gateway attB sites that were designed based on information from the Banana Genome Hub

(Supplemental Table S1). The resulting amplicons were recombined into the Gateway vector

pDONR^{TM}zeo (Invitrogen) and transformed into *Escherichia coli* TOP10 (Invitrogen) cells.

Sanger sequencing of the resulting entry plasmids was performed by the NIPGR sequencing

core facility (New Delhi, India).

Subcellular localization and co-localization assays

CDSs of *MaMYBFA1–3, MaHSF11,* and *MaDREB1* were cloned into the pSITE-3CA binary

vector (Chakrabarty *et al*., 2007) via LR recombination, producing yellow fluorescent protein

(YFP) fusion proteins. Empty vectors were used as negative controls. The binary plasmids

were transformed into *Agrobacterium tumefaciens* strain GV3101-pMP90 (Koncz &

Schell,□1986). Equal volumes of *A. tumefaciens* cultures having the desired gene and the

marker were mixed for agro-infiltration. *N. benthamiana* leaves were infiltrated on the

abaxial side using a syringe, according to the protocol described by Walter *et al.* (2004).

Infiltrated plants were kept in the dark at 25°C for 2 days prior to analysis. Leaf discs were

then cut from the infiltrated sites and observed under an argon laser confocal laser-scanning

microscope (TCS SP5; Leica Microsystems, Wetzlar, Germany) with YFP and RFP filters.

YFP and RFP fluorescence was observed at 514-nm excitation/527-nm emission and 558-nm

excitation/583-nm emission wavelengths, respectively. YFP/RFP gains (intensities %) for

MaMYBFA1, MaMYBFA2, MaMYBFA3, MaDREB1 and MaHSF11 were 760 (39)/158

(39), 818 (39)/158 (39), 818 (30)/158 (39), 820 (45)/168(42) and 718 (42)/154 (39),

respectively.

RT-qPCR analysis

Total RNA was extracted from different banana organs according to a previously described

protocol (Asif *et al.,* 2006) and subsequently treated with RNase-free DNase (TURBO DNA-

freeTM Kit, Invitrogen). Total RNA was reverse transcribed to generate first-strand cDNA

using a RevertAid H Minus First Strand cDNA Synthesis Kit (Thermo Fisher) and oligo(dT)

primers. RT-qPCR-based gene expression analyses were performed on a 7500 Fast Real-time

PCR System (Applied Biosystems) using the 2x SYBR Green PCR Master mix (Applied

Biosystems) and diluted cDNA corresponding to 10 ng total RNA in a final volume of 10

µL. *MaACTIN1* (GenBank AF246288) was used to normalize transcript abundance. Relative

transcript levels were calculated using the cycle threshold (C_t) $2^{-\Delta\Delta CT}$ method (Livak &

Schmittgen, 2001) and are presented as fold changes relative either to the tissue having the

lowest expression or to the control condition, depending on the requirement. Three biological

replicates or three technical replicates of a pool of 4-5 biological replicates were analyzed in

each RT-qPCR analysis.

Metabolite analysis

Targeted metabolite analysis was performed to quantify flavonols according to a protocol

described previously (Pandey *et al.,* 2016; Naik *et al.,* 2021; Rajput *et al.,* 2022a, b). Plant

metabolites were extracted from shredded plant material in 80% methanol at room

temperature. Finely ground tissues were dissolved in 80% methanol, kept at 70 for 15

minutes, and subjected to centrifugation for 10 minutes at 12,000 rpm (18514 g). After

centrifugation, supernatants were mixed with three volumes of 1N HCl for acid hydrolysis

and incubated at 94 for 2 h. After incubation, an equal volume of ethyl acetate was added,

and the upper phase was removed for further purification of the desired metabolites. The

ethyl acetate was evaporated at 37 under lower pressure (90–100 mbar) on a Buchi

rotavapor R-300 (BUCHI, Switzerland), and samples were finally dissolved in 80%

methanol. Liquid chromatography–mass spectrometry (LC-MS) analysis was performed

using an ultra-high-performance liquid chromatography (UHPLC) system (Exion LC Sciex,

Framingham) coupled to a triple quadrupole system (QTRAP6500+; ABSciex) with

electrospray ionization. The voltage was set to 5500 V for positive ionization. Detailed

UHPLC and LC-MS parameters have been described by Naik *et al.* (2021). Identification and

quantitative analyses were done via Analyst software (version 1.5.2).

Cloning of reporter constructs and identification of putative *cis* elements in promoters

Total genomic DNA was extracted from banana leaves using the DNeasy Plant Mini Kit

(Qiagen). Sequences of the putative target gene promoter fragments were retrieved from the

banana genome (https://banana-genome-hub.southgreen.fr) (D'Hont *et al.,* 2016) and

analyzed for the presence of *cis*-regulatory motifs via New PLACE software

(https://www.dna.affrc.go.jp) supplemented by manual curation. Promoter fragments were

amplified from banana genomic DNA using designed primer sequences and Phusion™ High-
Fidelity DNA Polymerase (Thermo Scientific). The amplified fragments were cloned into the
pENTR™/D-TOPO™ vector (ThermoFisher) and sequenced to confirm their integrity. The
entry clones of promoters (*proMaFLS1–2508*, *proMaF3'5'H1–1335*, *proMaDREB1–998*, and
proMaMYBFA3–2508) were recombined into the vector p635nRRF containing *pro35S:REN*
(Kumar *et al.,* 2018) to make the reporter constructs.

Cloning of cDNA from MaHSF and MaDREB transcriptional regulators

Full-length CDSs of selected homologs of MaHSF and MaDREB were amplified by first-
strand cDNA synthesis of banana leaves using Phusion™ High-Fidelity DNA Polymerase
(Thermo Scientific) and a set of primers containing Gateway attB sites designed on the basis
of information from the Banana Genome Hub (Supplemental Table S1). The resulting
amplicons were cloned into the pDONR^{TM} zeo vector (Invitrogen) and transformed into *E.*
coli TOP10 cells, and integrity was confirmed by Sanger sequencing (Sanger *et al.*, 1977).

Dual-luciferase reporter assay

To test the transactivation potential of the transcriptional regulators, MaHSF11, and
MaDREB1 CDSs were recombined into the destination vector pBTdest (Baudry *et al.,* 2004)
to generate CaMV *pro35S*-driven effector constructs. For dual-luciferase reporter assays,
reporter and effector constructs were transformed into *A. tumefaciens* strain GV3101
(pMP90) (GoldBio) and subsequently infiltrated using a syringe into the abaxial side of *N.*
benthamiana leaves according to a previously described protocol (Walter *et al.,* 2004). LUC
and REN activities were measured 48□h later in extracts from infiltrated leaf discs using the
Dual-Luciferase Reporter Assay System (Promega) according to the manufacturer's protocol.
Luminescence in the samples was quantified using the POLARstar Omega multimode plate
reader (BMG Labtech), and LUC activity was normalized to REN activity. Values are given
as means ± SD of the LUC/REN ratio of 4 independent biological replicates.

Y1H assays

The Matchmaker Gold Yeast 1-hybrid system (Clontech) with the yeast one-hybrid (Y1H)
gold yeast strain was used. Promoter fragments were cloned into the pABAi vector (Takara
Bio), and the pABAi plasmids carrying the promoter fragments were linearized and
transformed into the Y1H Gold strain according to the Matchmaker protocol, whereby they
integrated into the yeast genome by homologous recombination. Effector plasmids and empty

pGADT7-GW vector (Gal4-AD fusion, Ura selection) controls were transformed into the

Y1H strains previously transformed with AurR-reporter constructs (*proMaFLS1*-AurR,

proMaF3'5'H1-AurR, *proMaDREB1*-AurR, and *proMaMYBFA3*-AurR). To test for

interaction, the yeast colonies containing effector and reporter constructs were spotted onto

medium containing AbA per the manufacturer's protocol. Growth of different Ura⁻ yeast

colonies was checked using different concentrations of AbA to determine the minimum

inhibitory concentration (MIC) values and to check for possible interactions. The MIC values

for *proMaFLS1*, *proMaF3'5'H1*, *proMaDREB1*, and *proMaMYBFA3* were 100 ng mL^{-1}

AbA, 1000 ng mL^{-1} AbA, 100 ng mL^{-1}, and 900 ng mL^{-1}, respectively.

ChIP-qPCR assay

The ChIP-qPCR assay was performed using the ChIP Kit - Plants kit (Abcam) according to

the manufacturer's instructions (Shi *et al.,* 2019). Promoter fragments containing the HSE *cis*

motif in p635nRRF that was used for dual-luciferase assays were co-infiltrated with YFP-

MaHSF11 fusion proteins into *N. benthamiana* leaves. Three days after infiltration, equally

sized leaf discs were used for ChIP. Next, 2 µL anti-GFP antibody (Catalog #ab290, Abcam)

was used for chromatin precipitation in two biological replicates, and 2 µL of anti-IgG

antibody (Catalog #ab48386, Abcam) was used as a negative control. qPCR was performed

with 5 µL 2x SYBR Green PCR Master mix (Applied Biosystems), 1 µL ChIP DNA, and 0.5

µL each forward and reverse primer, with 40 cycles of amplification as described above. The

relative fold enrichment of the target promoter fragments was calculated using the $2^{-(-\Delta CT)}$

method (Livak & Schmittgen, 2001). The primers used in ChIP-qPCR are listed in

Supplemental Table S1.

Electrophoretic mobility-shift assay

For the EMSA, the fusion construct MaHSF11-GST or MdDREB1-GST with GST-tag were

obtained by cloning the CDS of MaHSF11 or MaDREB1 into pGEX-4T-2 vector, and then

individually introduced into *Escherichia coli* cells BL21 (DE3/Codon+) for protein

production. The culture solution was added with IPTG (final concentration 0.5 mM) at OD of

0.6 and kept on culturing at 28 , for 12 h followed by harvesting. GST-tagged recombinant

protein was purified using Glutathione Resin (catalog no. 786-310, G-Biosciences). The

probes containing intact and mutated cis-motifs were labelled with biotin using Pierce™

Biotin 3' End DNA Labeling Kit (catalog no. 89818, Thermo Fisher Scientific, USA). The

EMSA was carried out using LightShift Chemiluminescent EMSA Kit (catalog no. 20148,

Thermo Fisher Scientific, USA) according to the protocol described in the kit manual. The

primers and probe sequences for EMSA are shown in Supplementary Table S1.

Transient expression in banana fruit

For transient overexpression studies, the CDSs of *MaMYBFA1*, *MaMYBFA2*, *MaMYBFA3*,

MaHSF11, and *MaDREB1* were cloned into pANIC6b (Mann *et. al.*, 2012), a Gateway-

compatible destination vector suitable for overexpression of genes of interest using *A.*

tumefaciens-mediated transformation, particularly useful for monocot species. pANIC6b

encodes hygromycin resistance under the control of a constitutive rice (*Oryza sativa*) *actin1*

promoter (*proOsACT1*) (LOC4338914), a constitutive *Panicum virgatum ubiquitin1+3*

promoter (*proPvUBI*)-driven GUSPlus visual marker, and a Gateway cassette driven by a *Zea*

mays ubiquitin1 promoter and intron for overexpression of target genes. The resulting

overexpression plasmid and the empty vector control were transformed into *A. tumefaciens*

strain EHA105 (Mehra *et al.*, 2019). Resulting *A. tumefaciens* suspensions were then

separately introduced into discs of immature green banana fruits by vacuum infiltration as

previously described (Rajput *et al.,* 2022a). After 3 days of co-cultivation on MS media (MS

basal salt mixture, HiMedia), transformation of banana fruit disc was confirmed by GUS

staining. Tissues were harvested and frozen in liquid nitrogen for subsequent expression (RT-

qPCR) and metabolite (LC-MS) analyses.

For gene silencing through RNA interference, the 250-350 bp CDS fragments of *MaHSF11*

and *MaDREB1* were cloned into pANIC8b vector, a Gateway-compatible destination vector

suitable for silencing of target genes in monocot species (Mehra *et al.,* 2022). The

transformation into *A. tumefaciens* strain EHA105 and immature banana fruits were done as

described previously for overexpression. After GUS staining, tissues were harvested for

expression (RT-qPCR) and metabolite analysis (LC-MS).

Heat stress treatments

We used plant growth chamber (Model: AR-41L3, Percival) to perform heat stress treatment

at 250 µmol m^{-2} s^{-1} light intensity under a 16-h-light/8-h-dark photoperiod and 60% relative

humidity. Three months old *in vitro* grown banana plants (cultivar subgroup Cavendish

[AAA] variety 'Grand Naine') of equal size were kept at 26°C for control and 42°C for heat

treatment. Leaf samples were harvested at 0, 6, 12, 24, 48, and 72 hours (h) post heat stress

treatment followed by freezing in liquid nitrogen. Expression (RT-qPCR) and metabolite
analysis (LC-MS) were carried out as described earlier.

Y2H assay

For Y2H assays, full lengthen CDS of MaHSF11 and MaDREB1 were cloned into pGADT7
(Clontech Laboratories Inc., San Jose, CA, USA) and pGBKT7 vector, respectively. The
resulting AD and BD constructs were co-transformed into Y2HGold (Clontech) and grown at
30°C on synthetic defined (SD) medium devoid of Trp and Leu. For interaction assay, the
colonies having both the bait and prey clones were spotted onto medium lacking Trp, Leu,
Ade, and His (QDO). p53/large T antigen interaction (Pipas & Levine, 2001) was used as
the positive control and empty pGADT7g/pGBKT7g vectors with opposite and appropriate
constructs were used as a negative control.

Statistical tests

911 One-way and two-way analysis of variance (ANOVA) with Tukey's multiple comparison
912 were performed for statistical analysis. Unpaired Student's *t*-tests were used for the specific
913 comparisons. Asterisks indicate the level of significance: *$P \leq 0.05$; **$P \leq 0.01$;
914 ***$P \leq 0.001$, ****$P \leq 0.0001$; ns, not significant.

Data availability

916 All data supporting the findings of this study are available within the paper and within the
917 supplemental data published online.

Supplemental data

Figure S1. Identification of candidate R2R3-MYB flavonol regulators from the banana
genome and their subcellular localization.

Figure S2. Tissue-specific expression of *MaMYBFA1-FA3, MaFLS1-2* genes and flavonol
content in different tissues of banana**.**

Figure S3. Relative expression of structural flavonol biosynthesis gene transcripts in banana
fruit discs transiently overexpressing *MaMYBFA1-FA3***.**

Figure S4. Relative expression of structural and regulatory flavonol biosynthesis genes in
MaHSF11 overexpressing banana fruit discs.

Figure S5. Relative expression of structural and regulatory flavonol biosynthesis genes in *MaHSF11*-silenced banana fruit discs.

Figure S6. Phylogenetic analysis of MaDREB1 with previously characterized DREB proteins from other plant species, its sub-cellular localization and tissue specific expression pattern in different tissues of banana.

Figure S7. Relative expression of structural flavonol biosynthesis genes in overexpressing *MaDREB1* overexpressing banana fruit discs.

Figure S8. Relative expression of structural and regulatory flavonol biosynthesis genes in *MaDREB1*-silenced banana fruit discs.

Figure S9. Relative expression analysis of flavonol biosynthesis genes in control (26 □) and heat stress (42 □).

Figure S10. Relative expression analysis of *MaHSF11* in overexpressing *MaDREB1*- and *MaDREB1*-silenced banana fruit discs.

Figure S11. Interaction between MaHSF11 and MaDREB1 in yeast.

Acknowledgements

This work was supported by a core grant from the National Institute of Plant Genome Research and a Department of Science and Technology-SERB for Core research grant (CRG/2022/001178) to AP. JN, RR and SS acknowledge the Council of Scientific and Industrial Research, Government of India, for Senior Research Fellowships. The authors thank the DBT-eLibrary Consortium (DeLCON) for providing access to e-resources. We also acknowledge the Metabolome facility at NIPGR for phytochemical analysis.

Declarations

The authors declare no conflicts of interest.

Figure legends

Figure 1. MaMYBFA1, MaMYBFA2, and MaMYBFA3 differentially regulate the expression of *MaFLS1* and *MaFLS2* and modulate flavonol biosynthesis in banana. (a) Diagram of the plant expression cassette used for transient overexpression (OE) of *MaMYBFA* in banana fruit discs. The *Zea mays ubiquitin1* promoter (*proZmUBI1*) drives the expression of *MaMYBFA*. The expression of the *GUSPlus* marker gene is driven by the *Phaseolus vulgaris ubiquitin1* promoter (*proPvUBI*). (b) Representative images showing GUS staining of transiently transformed banana fruit discs indicating successful transformation. Images were taken 3 days after transformation. (c-d) Relative expression of *MaMYBFA1–3* (c) and *MaFLS1* and *MaFLS2* (d) OE lines in transiently transformed *MaMYBFA* OE banana fruit disc tissues as determined by RT-qPCR. EV, pANIC6b empty vector control. Values are represented as fold change in three replicates of pool of 4–5 independent biological samples. (e) Flavonol quantification showing differential alteration in the accumulation of kaempferol, quercetin, and myricetin aglycones in the *MaMYBFA* OE banana fruit discs. Values are given in µg per 100 g of dry weight (DW) of fruit discs of three replicates. One-way ANOVA with Tukey's multiple comparison was used for statistical analyses. *$P \leq 0.05$; **$P \leq 0.01$; ***$P \leq 0.001$; ****$P \leq 0.0001$; ns, not significant.

Figure 2. MaHSF11 directly binds to the *MaFLS1* promoter and regulates the expression of *MaFLS1* to modulate flavonol biosynthesis in banana. (a) Heatmap showing expression of five banana candidate *HSF* genes before and after drought treatment (Grand Naine cultivar, AAA genome). FPKM values for the indicated genes were extracted from publicly available RNA-Seq datasets and are represented according to the 0–7 color scale shown. (b) Dual-luciferase assay showing transactivation of *proMaFLS1–2556* by MaHSF. Values are shown as means ± SD of the LUC/REN ratios of four independent biological replicates ($n = 4$). (c) Illustration of the *cis*-motifs in the region located within 2487 bp upstream of the TSS in the *MaFLS1* promoter. F1 and F2 fragments contain one HSE element each having TTTCTAGAAG and TTTCAAGAAA motifs. +69 shows the bp distance between TSS and start codon. (d) ChIP-qPCR assay analyzing the binding of MaHSF11 to fragments F1 and F2 of *proMaFLS1*, both of which contain the HSE consensus sequence (nTTCnnGAAn). Fold enrichment values are given as means ± SD of two biological replicates. AB, antibody. (e) Y1H assay showing interaction of MaHSF11 with the two HSE-containing *proMaFLS1* fragments F1 and F2. The empty pGADT7 vector (EV) was used as a negative control. Interaction was indicated by yeast growth on synthetic Ura– dropout media supplemented with AbA. AD, activation domain. (f) EMSA showing the binding of MaHSF11-GST protein

with the F1 fragment of *MaFLS1* promoter (*MaFLS1*-F1) *in vitro*. Binding cis-motif and the mutated motif in the probes are highlighted in red in F1 and F1m, respectively. MaHSF11-GST protein, labelled and unlabeled probes were separated in polyacrylamide gel, blotted on nylon membrane, and detected by chemiluminescence. + indicates presence and − indicates absence of probe/protein.

Figure 3. MaHSF11 promotes flavonol synthesis in banana fruit. (a) The phylogenetic tree was constructed in MEGA-X32 software using the maximum-likelihood method with 1000 bootstrap values and was visualized with iTOL v.6.4 (http://itol.embl.de/). Accessions used are MdHSFA8a (NP_001281294.1), GmHSFB2b (XP_014619265.1), AtHSFA2 (AT2G26150), AtHSF1 (AT4G17750), AtHSF3 (AT5G16820), AtHSFA8 (AT1G67970), AtHSFC1 (AT3G24520), AtHSFA3 (AT5G03720), AtHSF4 (AT4G36990), AtHSFB2A (AT5G62020), AtHSFB2B (AT4G11660), and AtHSFA4A (AT4G18880). Md, *Malus domestica*; Gm, *Glycine max*; At, *Arabidopsis thaliana.* (b) Subcellular localization of MaHSF11 in *N. benthamiana* leaves. The YFP-MaHSF11 fusion protein localizes to both the nucleus and the cytoplasm and coincides with the signal from the nuclear localization signal-red fluorescent protein (NLS-RFP) fusion in *N. benthamiana* leaves as observed in confocal microscopy. Empty vector pSITE-3CA as used as negative control to visualize free YFP. Scale bar, 20 μm. (c) Relative expression of *MaHSF11* genes in different tissues of banana as determined by RT-qPCR. Values shown are means ± SD of three biological replicates. (d) GUS staining of banana fruit discs indicating successful transformation with the empty vector pANIC6b (EV) and the *proZmUBI:MaHSF11* construct. (e) Relative expression of *MaHSF11*, *MaFLS1*, and *MaFLS2* in transiently transformed banana fruit discs overexpressing (OE) *proZmUBI:MaHSF11* as determined by RT-qPCR. The empty vector was transformed as a control (expression level set to 1). Values shown are means ± SD of three technical replicates of pool of 4-5 independent biological samples. (f) Quantification of the flavonols kaempferol, quercetin, and myricetin in transformed banana fruit discs. (g) GUS staining of banana fruit discs indicating successful transformation with the empty vector EV and the *proZmUBI:MaHSF11*-RNAi construct. (h) Relative expression of *MaHSF11, MaFLS1,* and *MaFLS2* in *MaHSF11*-silenced banana fruit discs transiently expressing *proZmUBI:MaHSF11*-RNAi as determined by RT-qPCR. The empty vector pANIC8b was transformed as a control (expression level set to 1). Values shown are means ± SD of three technical replicates of pool of 4-5 independent biological samples. (i) Quantification of the flavonols kaempferol, quercetin, and myricetin in EV and *proZmUBI:MaHSF11*-RNAi

banana fruit discs. Values are represented as means ± SD of three replicates. Unpaired

student's t-tests were used for statistical analyses. *P $\leq$ 0.05; **P $\leq$ 0.01;

***P $\leq$ 0.001.

Figure 4. MaHSF11 promotes flavonoid B-ring hydroxylation by modulating the expression

of *MaF3'5'H1*. (a) Relative expression of *MaF3'5'H1* genes in transformed banana fruit discs

transiently overexpressing *proZmUBI:MaHSF11*, as determined by RT-qPCR (EV, empty

vector control). Values shown are means ± SD of three technical replicates of pool of 4-5

independent biological samples. (b) Relative expression of *MaF3'5'H1* genes in *MaHSF11*-

silenced banana fruit discs, as determined by RT-qPCR (EV, empty vector control). Values

shown are means ± SD of three technical replicates of pool of 4-5 independent biological

samples. (c) Illustration of the *cis*-motifs in the region located within 1335 bp upstream of the

TSS in the *MaF3'5'H1* promoter. F1 fragment contain one HSE element having

GGAATATTCT motifs. (d) Y1H assay showing the interaction of MaHSF11 with

proMaF3'5'H1–100 fragment F1 carrying HSE (GGAATATTCT). pGADT7 was used as a

negative control. Growth was checked on synthetic dropout medium supplemented with

AbA. (e) Dual-luciferase assay showing transactivation of *proMaF3'5'H1–1335* by

MaHSF11. The promoter sequence driving the expression of *luciferase* (LUC) was co-

infiltrated with the EV or *MaHSF11* constructs in *N. benthamiana* leaves. Values are shown

as means ± SD of the LUC/REN ratio of four independent biological replicates (*n* = 4). (f)

ChIP-qPCR assay showing binding of MaHSF11 to fragment F1 of *proMaF3'5'H1* carrying

an HSE (nGAAnnTTCn). The promoter fragment carrying HSE was co-infiltrated into *N.*

benthamiana leaves with the *pro35S:YFP-MaHSF11* construct. Fold enrichment values are

represented as means ± SD of two biological replicates. An anti-IgG antibody was used as a

negative control, and an anti-GFP antibody was used to precipitate MaHSF11-enriched

chromatin. Unpaired Student's *t*-tests were used for statistical analyses. *P $\leq$ 0.05.

Figure 5. MaHSF11 acts upstream of *MaDREB1* to positively regulate its expression in

banana. (a) Heatmap showing expression of candidate *MaDREB* genes before and after

drought treatment in banana (Grand Naine cultivar, AAA genome). FPKM values for the

indicated genes were extracted from publicly available RNA-Seq datasets and plotted

according to the 0–7 color scale shown. (b) Relative expression of *MaDREB1* in banana fruit

discs transiently transformed with EV or *proZmUBI:MaHSF11* as determined by RT-qPCR.

(c) Relative expression of *MaDREB1* in *MaHSF11*-silenced banana fruit discs and EV

control as determined by RT-qPCR. (d) Illustration of the *cis*-motifs in the region located

within 688 bp upstream of the TSS in the *MaDREB1* promoter. F1 and F2 fragments contain one HSE element each having AGAAGCTTCA and TGAAGCTTCG motifs. +310 shows the bp distance between TSS and start codon. (e) Y1H assay showing interaction of MaHSF11 with the F1 and F2 fragments of *proMaDREB1–100* carrying an HSE (nGAAnnTTCn). pGADT7 was used as a negative control. Growth was checked on synthetic dropout medium supplemented with AbA. (f) Dual-luciferase assay showing transactivation of *proMaDREB1–998* by MaHSF11. Promoter sequence driving the expression of *LUC* was co-infiltrated with empty vector and *MaHSF11* construct in *N. benthamiana* leaves. Values shown are means ± SD of LUC/REN ratio ($n = 4$ independent biological replicates). (g) ChIP-qPCR assay showing binding of MaHSF11 to *proMaDREB1*. A promoter fragment carrying HSE was co-infiltrated into *N. benthamiana* leaf cells with the *pro35S:MaHSF11-YFP* construct. Fold enrichment values are represented as means ± SD of two biological replicates. An anti-IgG antibody was used as a negative control, and an anti-GFP antibody was used to precipitate MaHSF11 enriched chromatin. Unpaired Student's t-tests were used for statistical analyses. *$P \leq 0.05$; ***$P \leq 0.001$; ns, not significant. (h) EMSA showing the binding of MaHSF11-GST protein with the F1 fragment of *MaDREB1* promoter (*MaDREB1*-F1) *in vitro*. Binding cis-motif and the mutated motif in the probes are highlighted in red in F1 and F1m, respectively. MaHSF11-GST protein, labelled and unlabeled probes were separated in a polyacrylamide gel, blotted on nylon membrane and detected by chemiluminescence. + indicates presence and – indicates absence of probe/protein.

Figure 6. MaDREB1 directly binds to the promoter of *MaMYBFA3* to regulate flavonol biosynthesis. (a) Y1H assay showing interaction of MaDREB1 with the *proMaMYBFA3–1892* fragment carrying CRT/DREs (CCGAC/GTCGG). pGADT7 was used as a negative control. Growth was checked on synthetic dropout medium supplemented with AbA. (b) Dual-luciferase assay showing transactivation of *proMaMYBFA3–2508* by MaDREB1. The *proMaMYBFA3–2508* sequence driving the expression of *LUC* was co-infiltrated with the empty vector or *MaDREB1* construct into *N. benthamiana* leaves. Values are shown as means ± SD of the LUC/REN ratio ($n = 4$ independent biological replicates). (c) Illustration of the cis-motifs in the region located within 2367 bp upstream of the TSS in the *MaMYBFA3* promoter. F1, F2, F3 and F4 fragments contain one to two DRE/CRT elements having either CCGAC or GTCGG motifs. +141 shows the bp distance between TSS and start codon. (d) EMSA showing the binding of MaDREB1-GST protein with the F3 fragment of *MaDREB1* promoter (*proMaDREB1-F3*) *in vitro*. Different cis-motifs (CCGAC/GTCGG) present in the

F1, F2, F3 and F4 probes are highlighted in red. MaDREB1-GST protein, labelled and unlabeled probes were separated in a polyacrylamide gel, blotted on nylon membrane and detected by chemiluminescence. + indicates presence and – indicates absence of probe/protein.

Figure 7. MaDREB1 regulates the expression of *MaMYBFAs*, *MaFLS1*, and *MaFLS2* to modulate flavonol biosynthesis in banana. (a) GUS staining of EV and *proZmUBI:MaDREB1* transformed fruit discs. (b) RT-qPCR-based relative expression of *MaDREB1*, and (c) relative expression of *MaMYBFAs*, *MaFLS1*, and *MaFLS2* in transformed banana fruit discs transiently expressing EV and *proZmUBI:MaDREB1*. EV was transformed as a control. Values shown are means ± SD of three technical replicates of a pool of 4-5 biological replicates. (d) Quantification of the flavonols kaempferol, quercetin, and myricetin in transformed fruit discs. (e) GUS staining of EV and *proZmUBI:MaDREB1-RNAi* transformed fruit discs. (f) RT-qPCR-based relative expression of *MaDREB1, and (g)* relative expression of *MaMYBFAs, MaFLS1*, and *MaFLS2* in *MaDREB1*-silenced banana fruit discs transiently expressing *proZmUBI:MaDREB1-RNAi* and EV. EV pANIC8b was transformed as a control. Values shown are means ± SD of three technical replicates of pool of 4-5 independent biological pools. (h) Quantification of the flavonols kaempferol, quercetin, and myricetin in transiently expressing *proZmUBI:MaDREB1*-RNAi banana fruit discs. Values are represented as means ± SD of three replicates. Unpaired Student's t-tests were used for statistical analyses. *P□≤□0.05; **P□≤□0.01; ***P□≤□0.001; ****P ≤ 0.0001.

Figure 8. Heat stress positively modulates flavonol biosynthesis in banana (a) Representative images of three months old banana seedlings exposed heat stress at 42□. Control seedlings were kept at 26□. (b) RT-qPCR-based relative expression of *MaHSF11, MaDREB1, MaMYBFA1-3, MaFLS1* and *MaFLS2* at different time points (0h, 6h, 12h, 24h, 48h, 72h) of heat stress (42□) treatment in banana seedlings, and control (26□) seedlings. (c) Quantification of the flavonols aglycones i.e., kaempferol, quercetin, and myricetin in the heat stress treated and control plants at different time intervals. Values are represented as mean ±SD of three biological replicates.

Figure 9. A working model showing the modulation of hydroxylated flavonol synthesis by the MaHSF11–MaDREB1–MaMYBFA regulatory module. The banana SG7 R2R3 MYB, MaMYBFA, targets *MaFLS2* to promote its expression, while strongly repressing the expression of *MaFLS1*, and modulate flavonol synthesis. Drought stress increases the

transcript level of *MaHSF11*, enhancing the expression of *MaFLS1* and *MaF3'5'H1*. MaHSF11 also directly activates the expression of *MaDREB1*. MaDREB1 binds to the *MaMYBFA3* promoter to regulate its expression. This regulatory module enhances overall accumulation of flavonols under drought stress. MaMYBFA1, MaMYBFA2, and MaMYBFA3 differentially regulate the expression of *MaFLS1* and *MaFLS2*. MRE, MYB recognition element; DRE, dehydration response element; HSE, heat shock element. Heat stress modulates the expression of *MaHSF11, MaDREB1,* and *MaMYBFA1-3,* which subsequently leads to up-regulation of flavonol synthesis genes and increased accumulation of flavonols under heat stress in banana. Flavonols are likely transported into the vacuole by MATE transporters (multidrug and toxic compound extrusion), once the stress subsides.

Supplemental Figures

Supplemental Figure S1. Identification of candidate R2R3-MYB flavonol regulators from the banana genome and their subcellular localization. (a) Various R2R3-MYB proteins were classified into three major groups as flavonol-, anthocyanin-, or proanthocyanidin biosynthesis–specific regulators using landmark flavonoid-specific R2R3-MYBs from different plant species. Numbers on each branch show branch length, and solid dark blue circles represent bootstrap values. (b) Amino acid sequence alignment of candidate SG7 R2R3-MYB transcription factors (MaMYBFA1, MaMYBFA2, and MaMYBFA3, where FA stands for flavonol activator) and selected flavonol regulators from other plant species. The R2 and R3 domains are highlighted in grey. The SG7-1 and SG7-2 motifs, found in subgroup 7 R2R3-MYBs, are highlighted in dark green and yellow, respectively. Accession numbers and two-letter species abbreviations are provided in Supplemental File S2. (c) Subcellular localization of MaMYBFAs. MaMYBFA-YFP fusion proteins were expressed together with the nuclear marker NLS-RFP in *N. benthamiana* leaf epidermal cells. Confocal imaging shows co-localization of YFP and RFP fluorescence, indicating nuclear localization of the three MaMYBFA proteins. Empty vector pSITE-3CA was also taken as negative control, which shows free YFP. Representative images are shown with scale bar of 50 µm.

Supplemental Figure S2. Tissue-specific expression of *MaMYBFA1–3, MaFLS1-2* genes and flavonol content in different tissues of banana. (a) Relative expression of *MaMYBFA1–3* genes in different tissues of banana as determined by RT-qPCR. (b) Relative

expression of *MaFLS1* and *MaFLS2* genes in different tissues of banana. (c) Flavonol aglycones i.e., kaempferol, quercetin and myricetin content in µg per 100g dry weight (DW) in different tissues of banana. Tissues from three independent biological replicates were used for expression and metabolite (LC-MS) analysis. YL, young leaf; BRT, bract; P.STEM, pseudostem; UR.PL, unripe peel; UR.PP, unripe pulp; R.PL, ripe peel; R.PP, ripe pulp. Values shown are means ± SD of three biological replicates.

Supplemental Figure S3. Relative expression of structural flavonol biosynthesis gene transcripts in banana fruit discs transiently overexpressing *MaMYBFA1-FA3*. RT-qPCR-based transcript profiling of structural flavonol biosynthesis genes *MaCHS1–6*, *MaCHI1–2*, *MaF3H1–2*, *MaF3'5'H1–7*, and *MaF3'H* in *MaMYBFA1–3* OE banana fruit discs. EV, empty vector control. Values shown are means ± SD of three technical replicates of pool of 4-5 independent biological samples. One-way ANOVA was used for statistical analysis with Tukey's multiple comparison. $*P \leq 0.05$; $**P \leq 0.01$; $***P \leq 0.001$; $****P \leq 0.0001$; ns, not significant.

Supplemental Figure S4. Relative expression of structural and regulatory flavonol biosynthesis genes in *MaHSF11* overexpressing banana fruit discs. RT-qPCR-based expression analysis of (a) *MaCHS1–6*, (b) *MaCHI1–2*, (c) *MaF3H1–2*, (d) *MaF3'H*, and (e) *MaMYBFA1–3* in *MaHSF11* overexpressing banana fruit discs. Values shown are means ± SD of three technical replicates of a pool of 4-5 independent biological samples. EV, empty vector control. Two-way ANOVA was used with Tukey's multiple comparison. $P \leq 0.05$; $**P \leq 0.01$; ns, not significant. N.D., not determined.

Supplemental Figure S5. Relative expression of structural and regulatory flavonol biosynthesis genes in *MaHSF11*-silenced banana fruit discs. RT-qPCR-based expression analysis of (a) *MaCHS1–6*, (b) *MaCHI1–2*, (c) *MaF3H1–2*, (d) *MaF3'H*, and (e) *MaMYBFA1–3* in EV and *proZmUBI:MaHSF11-RNAi* banana fruit discs. Values shown are means ± SD of three technical replicates of a pool of 4-5 independent biological samples. EV, empty vector control. Two-way ANOVA was used with Tukey's multiple comparison. $*P \leq 0.05$; $**P \leq 0.01$; $****P \leq 0.0001$; ns, not significant. N.D., not determined.

Supplemental Figure S6. Phylogenetic analysis of MaDREB1 with previously characterized DREB proteins from other plant species, its sub-cellular localization and tissue specific expression pattern in different tissues of banana. (a) The phylogenetic tree was constructed in MEGA-X32 software using the maximum-likelihood method with 1000

bootstrap values and visualized with iTOL v.6.4 (http://itol.embl.de/). Accessions used were AtDREB1B (AAV80413.1), AtDREB1C (NP_567719.1), AtDREB1C (AAV80414.1), GmDREB3 (AAZ03388.1), GmDREB2 (NP_001237254.2), GmDREB2A2 (NP_001240942.3), OsDREB1 (AAL40870.1), OsDREB1B (AAN85707.1), OsDREB3 (ADO79620.1), SlDREB3 (NP_001234707.1), HvDREB2 (AFH75401.1), and ZmDREB (NP_001105081.1). At, *Arabidopsis thaliana*; Gm, *Glycine max*; Os, *Oryza sativa*; Sl, *Solanum lycopersicum*; Hv, *Hordeum vulgare*; Zm, *Zea mays*). (b) Subcellular localization of MaDREB1 in *N. benthamiana* leaves. The YFP-MaDREB1 fusion protein localizes to the nucleus and coincides with the signal from the nuclear localization signal-red fluorescent protein (NLS-RFP) fusion in *N. benthamiana* leaves as observed in confocal microscopy. Empty vector pSITE-3CA as used as negative control to visualize free YFP. Scale bar, 50 µm. (c) Relative expression of *MaDREB1* gene in different tissues of banana as determined by RT-qPCR. Values shown are means ± SD of three biological replicates.

Supplemental Figure S7. Relative expression of structural flavonol biosynthesis genes in *MaDREB1* overexpressing banana fruit discs. RT-qPCR-based expression analysis of (a) *MaCHS1–6*, (b) *MaCHI1–2*, (c) *MaF3H1–2*, (d) *MaF3'H* and (e) *MaF3'5'H1–7* in *MaDREB1* overexpressing banana fruit discs. Values shown are means ± SD of three technical replicates of a pool of 4-5 independent biological pools. Two-way ANOVA was used with Tukey's multiple comparison. *$P ≤ 0.05$; **$P ≤ 0.01$; ***$P ≤ 0.001$; ns, not significant.

Supplemental Figure S8. Relative expression of structural flavonol biosynthesis genes in *MaDREB1*-silenced banana fruit discs. RT-qPCR-based expression analysis of (a) *MaCHS1–6*, (b) *MaCHI1–2*, (c) *MaF3'H1–2*, (d) *MaF3'H* and (e) *MaF3'5'H1–7* in *MaDREB1*-silenced banana fruit discs. Values shown are means ± SD of three technical replicates of a pool of 4-5 independent biological pools. Two-way ANOVA was used with Tukey's multiple comparison. *$P ≤ 0.05$; **$P ≤ 0.01$; ****$P ≤ 0.0001$; ns, not significant.

Supplemental Figure S9. Relative expression analysis of other flavonol biosynthesis genes in control (26□) and heat stress (42□) treated banana seedlings. RT-qPCR based relative expression level of *MaCHS1-6, MaCHI1-2, MaF3H1-2, MaF3'H,* and *MaF3'5'H1-7* at different time points (0h, 6h, 12h, 24h, 48h, 72h) of heat stress (42 □) treatment in banana

seedlings, and control (26 ☐) seedlings. Values shown are means ± SD of three biological replicates.

Supplemental Figure S10. Relative expression analysis of *MaHSF11* in *MaDREB1* overexpressing *and MaDREB1*-silenced banana fruit discs. Values shown are means ± SD of three technical replicates of a pool of 4-5 independent biological samples. Unpaired Student's *t*-tests were used with Tukey's multiple comparison for statistical analysis. ****P*☐≤☐0.001, ns. Not significant.

Supplemental Figure S11. Interaction between MaHSF11 and MaDREB1 in yeast. Yeast-two-hybrid assays showing no direct interaction of MaHSF11 with MaDREB1 in yeast. AD, GAL4 activation domain; BD, GAL4 DNA-binding domain; -LT medium, synthetic defined (SD)–Leu–Trp medium; -AHLT, SD–Leu–Trp–Ade–His medium.

Supplemental File S1. Promoter sequences with highlighted *cis* motifs used in this study.

Supplemental File S2. Accession numbers and two-letter species abbreviations used in phylogenetic tree figures.

Supplemental Table S1. List of primers used in this study.

References

Albert, N.W., Thrimawithana, A.H., McGhie, T.K., Clayton, W.A., Deroles, S.C., Schwinn, K.E., Bowman, J.L., Jordan, B.R. and Davies, K.M., 2018. Genetic analysis of the liverwort *Marchantia polymorpha* reveals that R2R3 MYB activation of flavonoid production in response to abiotic stress is an ancient character in land plants. *New Phytologist*, 218(2), pp.554-566.

Asif, M., Trivedi, P., Solomos, T. and Tucker, M., 2006. Isolation of high-quality RNA from apple *(Malus domestica)* fruit. *Journal of agricultural and food chemistry*, 54(15), pp.5227-5229.

Battat, M., Eitan, A., Rogachev, I., Hanhineva, K., Fernie, A., Tohge, T., Beekwilder, J. and Aharoni, A., 2019. A MYB triad controls primary and phenylpropanoid metabolites for pollen coat patterning. *Plant Physiology*, 180(1), pp.87-108.

Baudry, A., Heim, M.A., Dubreucq, B., Caboche, M., Weisshaar, B. and Lepiniec, L., 2004. TT2, TT8, and TTG1 synergistically specify the expression of BANYULS and

proanthocyanidin biosynthesis in *Arabidopsis thaliana*. *The Plant Journal*, 39(3), pp.366-380.

Bhatia, C., Gaddam, S.R., Pandey, A. and Trivedi, P.K., 2021. COP1 mediates light-dependent regulation of flavonol biosynthesis through HY5 in Arabidopsis. *Plant Science*, 303, p.110760.

Bian, X.H., Li, W., Niu, C.F., Wei, W., Hu, Y., Han, J.Q., Lu, X., Tao, J.J., Jin, M., Qin, H. and Zhou, B., 2020. A class B heat shock factor selected for during soybean domestication contributes to salt tolerance by promoting flavonoid biosynthesis. *New Phytologist*, 225(1), pp.268-283.

Busche, M., Acatay, C., Martens, S., Weisshaar, B. and Stracke, R., 2021. Functional characterisation of banana (*Musa spp.*) 2-oxoglutarate-dependent dioxygenases involved in flavonoid biosynthesis. *Frontiers in Plant Science*, 12, p.701780.

Chapman, J.M. and Muday, G.K., 2021. Flavonols modulate lateral root emergence by scavenging reactive oxygen species in Arabidopsis thaliana. *Journal of Biological Chemistry*, 296.

Chauhan, H., Khurana, N., Agarwal, P. and Khurana, P., 2011. Heat shock factors in rice (Oryza sativa L.): genome-wide expression analysis during reproductive development and abiotic stress. *Molecular Genetics and Genomics*, 286(2), pp.171-187.

Chen, H., Liu, L., Wang, L., Wang, S. and Cheng, X., 2016. VrDREB2A, a DREB-binding transcription factor from *Vigna radiata*, increased drought and high-salt tolerance in transgenic *Arabidopsis thaliana*. *Journal of plant research*, 129(2), pp.263-273.

Chen, J., Ullah, C., Reichelt, M., Gershenzon, J. and Hammerbacher, A., 2019. *Sclerotinia sclerotiorum* circumvents flavonoid defenses by catabolizing flavonol glycosides and aglycones. *Plant Physiology*, 180(4), pp.1975-1987.

Choudhury, F.K., Rivero, R.M., Blumwald, E. and Mittler, R., 2017. Reactive oxygen species, abiotic stress and stress combination. *The Plant Journal*, 90(5), pp.856-867.

Cramer, G.R., Urano, K., Delrot, S., Pezzotti, M. and Shinozaki, K., 2011. Effects of abiotic stress on plants: a systems biology perspective. *BMC plant biology*, 11(1), pp.1-14.

Crosby, K.C., Pietraszewska-Bogiel, A., Gadella Jr, T.W. and Winkel, B.S., 2011. Förster resonance energy transfer demonstrates a flavonoid metabolon in living plant cells that displays competitive interactions between enzymes. *FEBS letters*, 585(14), pp.2193-2198.

Czemmel, S., Stracke, R., Weisshaar, B., Cordon, N., Harris, N.N., Walker, A.R., Robinson, S.P. and Bogs, J., 2009. The grapevine R2R3-MYB transcription factor VvMYBF1 regulates flavonol synthesis in developing grape berries. *Plant physiology*, 151(3), pp.1513-1530.

D'hont, A., Denoeud, F., Aury, J.M., Baurens, F.C., Carreel, F., Garsmeur, O., Noel, B., Bocs, S., Droc, G., Rouard, M. and Da Silva, C., 2012. The banana (*Musa acuminata*) genome and the evolution of monocotyledonous plants. *Nature*, 488(7410), pp.213-217.

Dalman, K., Wind, J.J., Nemesio-Gorriz, M., Hammerbacher, A., Lundén, K., Ezcurra, I. and Elfstrand, M., 2017. Overexpression of *PaNAC03*, a stress induced NAC gene family transcription factor in Norway spruce leads to reduced flavonol biosynthesis and aberrant embryo development. *BMC Plant Biology*, 17, pp.1-17.

Davies, K.M., Jibran, R., Albert, N.W., Zhou, Y. and Schwinn, K.E., 2021. Conservation and divergence between bryophytes and angiosperms in the biosynthesis and regulation of flavonoid production. *Recent Advances in Polyphenol Research*, 7, pp.227-263.

Dubouzet, J.G., Sakuma, Y., Ito, Y., Kasuga, M., Dubouzet, E.G., Miura, S., Seki, M., Shinozaki, K. and Yamaguchi-Shinozaki, K., 2003. *OsDREB* genes in rice, *Oryza sativa* L., encode transcription activators that function in drought-, high-salt-and cold-responsive gene expression. *The Plant Journal*, 33(4), pp.751-763.

Fahad, S., Bajwa, A.A., Nazir, U., Anjum, S.A., Farooq, A., Zohaib, A., Sadia, S., Nasim, W., Adkins, S., Saud, S. and Ihsan, M.Z., 2017. Crop production under drought and heat stress: plant responses and management options. *Frontiers in Plant Science*, 8, p.1147. doi.org/10.3389/fpls.2017.01147.

Fragkostefanakis, S., Mesihovic, A., Simm, S., Paupière, M.J., Hu, Y., Paul, P., Mishra, S.K., Tschiersch, B., Theres, K., Bovy, A. and Schleiff, E., 2016. HsfA2 controls the activity of developmentally and stress-regulated heat stress protection mechanisms in tomato male reproductive tissues. *Plant Physiology*, 170(4), pp.2461-2477.

Friedrich, T., Oberkofler, V., Trindade, I., Altmann, S., Brzezinka, K., Lämke, J., Gorka, M., Kappel, C., Sokolowska, E., Skirycz, A. and Graf, A., 2021. Heteromeric HSFA2/HSFA3

complexes drive transcriptional memory after heat stress in Arabidopsis. *Nature*
Communications, 12(1), pp.1-15.

Fujino, N., Tenma, N., Waki, T., Ito, K., Komatsuzaki, Y., Sugiyama, K., Yamazaki, T.,
Yoshida, S., Hatayama, M., Yamashita, S. and Tanaka, Y., 2018. Physical interactions among
flavonoid enzymes in snapdragon and torenia reveal the diversity in the flavonoid metabolon
organization of different plant species. *The Plant Journal*, 94(2), pp.372-392.

Gautam, H., Sharma, A. and Trivedi, P.K., 2023. The role of flavonols in insect resistance
and stress response. *Current Opinion in Plant Biology*, 73, p.102353.

Grunewald, W., De Smet, I., Lewis, D.R., Löfke, C., Jansen, L., Goeminne, G., Vanden
Bossche, R., Karimi, M., De Rybel, B., Vanholme, B. and Teichmann, T., 2012.
Transcription factor WRKY23 assists auxin distribution patterns during Arabidopsis root
development through local control on flavonol biosynthesis. *Proceedings of the National*
Academy of Sciences, 109(5), pp.1554-1559.

Guo, L., Gao, L., Ma, X., Guo, F., Ruan, H., Bao, Y., Xia, T. and Wang, Y., 2019. Functional
analysis of flavonoid 3'-hydroxylase and flavonoid 3', 5'-hydroxylases from tea plant
(*Camellia sinensis*), involved in the B-ring hydroxylation of flavonoids. *Gene*, 717,
p.144046.

Guo, M., Liu, J.H., Ma, X., Luo, D.X., Gong, Z.H. and Lu, M.H., 2016. The plant heat stress
transcription factors (HSFs): structure, regulation, and function in response to abiotic stresses.
Frontiers in Plant Science, 7, p.114.

Halbwirth, H., 2010. The creation and physiological relevance of divergent hydroxylation
patterns in the flavonoid pathway. *International Journal of Molecular Sciences*, 11(2),
pp.595-621.

Han, D., Huang, B., Li, Y., Dang, Q., Fan, L., Nie, J., Wang, Y., Yuan, Y. and Jia, D., 2022.
The MdAP2-34 modulates flavonoid accumulation in apple (*Malus domestica* Borkh.) by
regulating MdF3'H. *Postharvest Biology and Technology*, 192, p.111994.

Huang, Y.C., Niu, C.Y., Yang, C.R. and Jinn, T.L., 2016. The heat stress factor HSFA6b
connects ABA signaling and ABA-mediated heat responses. *Plant Physiology*, 172(2),
pp.1182-1199.

Jiang, W., Xia, Y., Su, X. and Pang, Y., 2022. ARF2 positively regulates flavonols and
proanthocyanidins biosynthesis in *Arabidopsis thaliana*. *Planta*, 256(2), p.44.

Jiang, X., Shi, Y., Fu, Z., Li, W.W., Lai, S., Wu, Y., Wang, Y., Liu, Y., Gao, L. and Xia, T.,
2020. Functional characterization of three flavonol synthase genes from *Camellia sinensis*:
Roles in flavonol accumulation. *Plant Science*, 300, p.110632.

Kalaji, H.M., Dąbrowski, P., Cetner, M.D., Samborska, I.A., Łukasik, I., Brestic, M., Zivcak,
M., Tomasz, H., Mojski, J., Kociel, H. and Panchal, B.M., 2017. A comparison between
different chlorophyll content meters under nutrient deficiency conditions. *Journal of Plant*
Nutrition, 40(7), pp.1024-1034.

Katoh, K. and Standley, D.M., 2013. MAFFT multiple sequence alignment software version
7: improvements in performance and usability. *Molecular Biology and Evolution*, 30(4),
pp.772-780.

Kitamura, S., Oono, Y. and Narumi, I., 2016. Arabidopsis *pab1*, a mutant with reduced
anthocyanins in immature seeds from banyuls, harbors a mutation in the MATE transporter
FFT. *Plant Molecular Biology*, 90(1), pp.7-18.

Ko, M.J., Cheigh, C.I. and Chung, M.S., 2014. Relationship analysis between flavonoids
structure and subcritical water extraction (SWE). *Food Chemistry*, 143, pp.147-155.

Koncz, C. and Schell, J., 1986. The promoter of TL-DNA gene 5 controls the tissue-specific
expression of chimaeric genes carried by a novel type of Agrobacterium binary vector.
Molecular and General Genetics MGG, 204(3), pp.383-396.

Kuang, J.F., Chen, J.Y., Liu, X.C., Han, Y.C., Xiao, Y.Y., Shan, W., Tang, Y., Wu, K.Q., He,
J.X. and Lu, W.J., 2017. The transcriptional regulatory network mediated by banana *(Musa*
acuminata) dehydration□responsive element binding (MaDREB) transcription factors in fruit
ripening. *New Phytologist*, 214(2), pp.762-781.

Kumar, K., Purayannur, S., Kaladhar, V.C., Parida, S.K. and Verma, P.K., 2018. mQTL□seq
and classical mapping implicates the role of an AT□HOOK MOTIF CONTAINING
NUCLEAR LOCALIZED (AHL) family gene in *A. scochyta* blight resistance of chickpea.
Plant, Cell & Environment, 41(9), pp.2128-2140.

Kumar, S., Stecher, G., Li, M., Knyaz, C. and Tamura, K., 2018. MEGA X: molecular evolutionary genetics analysis across computing platforms. *Molecular Biology and Evolution*, 35(6), p.1547.

Lago, J.H.G., Toledo-Arruda, A.C., Mernak, M., Barrosa, K.H., Martins, M.A., Tibério, I.F. and Prado, C.M., 2014. Structure-activity association of flavonoids in lung diseases. *Molecules*, 19(3), pp.3570-3595.

Lee, S.J., Kang, J.Y., Park, H.J., Kim, M.D., Bae, M.S., Choi, H.I. and Kim, S.Y., 2010. DREB2C interacts with ABF2, a bZIP protein regulating abscisic acid-responsive gene expression, and its overexpression affects abscisic acid sensitivity. *Plant Physiology*, 153(2), pp.716-727.

Li, L., Peng, H., Tan, S., Zhou, J., Fang, Z., Hu, Z., Gao, L., Li, T., Zhang, W. and Chen, L., 2020. Effects of early cold stress on gene expression in *Chlamydomonas reinhardtii*. *Genomics*, 112(2), pp.1128-1138.

Liang, T., Shi, C., Peng, Y., Tan, H., Xin, P., Yang, Y., Wang, F., Li, X., Chu, J., Huang, J. and Yin, Y., 2020. Brassinosteroid-activated BRI1-EMS-SUPPRESSOR 1 inhibits flavonoid biosynthesis and coordinates growth and UV-B stress responses in plants. *Plant Cell*, 32(10), pp.3224-3239.

Liu, H.C. and Charng, Y.Y., 2012. Acquired thermotolerance independent of heat shock factor A1 (HsfA1), the master regulator of the heat stress response. *Plant Signaling & Behavior*, 7(5), pp.547-550.

Liu, H.T., Gao, F., Li, G.L., Han, J.L., Liu, D.L., Sun, D.Y. and Zhou, R.G., 2008. The calmodulin binding protein kinase 3 is part of heat shock signal transduction in *Arabidopsis thaliana*. *The Plant Journal*, 55(5), pp.760-773.

Liu, J., Sun, N., Liu, M., Liu, J., Du, B., Wang, X. and Qi, X., 2013. An autoregulatory loop controlling Arabidopsis HsfA2 expression: role of heat shock-induced alternative splicing. *Plant Physiology*, 162(1), pp.512-521.

Livak, K.J. and Schmittgen, T.D., 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2− ΔΔCT method. *Methods*, 25(4), pp.402-408.

López, M.A., Bannenberg, G. and Castresana, C., 2008. Controlling hormone signaling is a plant and pathogen challenge for growth and survival. *Current Opinion in Plant Biology*, 11(4), pp.420-427.

Lu, J., Tong, P., Xu, Y., Liu, S., Jin, B., Cao, F. and Wang, L., 2023. SA-responsive transcription factor GbMYB36 promotes flavonol accumulation in *Ginkgo biloba*. *Forestry Research*, 3(1).

Ma, D., Tang, H., Reichelt, M., Piirtola, E.M., Salminen, J.P., Gershenzon, J. and Constabel, C.P., 2021. Poplar MYB117 promotes anthocyanin synthesis and enhances flavonoid B-ring hydroxylation by up-regulating the flavonoid 3', 5'-hydroxylase gene. *Journal of Experimental Botany*, 72(10), pp.3864-3880.

Maloney, G.S., DiNapoli, K.T. and Muday, G.K., 2014. The anthocyanin reduced tomato mutant demonstrates the role of flavonols in tomato lateral root and root hair development. *Plant Physiology*, 166(2), pp.614-631.

Mann DG, LaFayette PR, Abercrombie LL, King ZR, Mazarei M, Halter MC, Poovaiah CR, Baxter H, Shen H, Dixon RA, Parrott WA. Gateway compatible vectors for high throughput gene functional analysis in switchgrass (*Panicum virgatum* L.) and other monocot species. *Plant Biotechnology Journal*, 10(2), pp. 226-36.

Manna, M., Thakur, T., Chirom, O., Mandlik, R., Deshmukh, R. and Salvi, P., 2021. Transcription factors as key molecular target to strengthen the drought stress tolerance in plants. *Physiologia Plantarum*, 172(2), pp.847-868.

Martinez, V., Mestre, T.C., Rubio, F., Girones-Vilaplana, A., Moreno, D.A., Mittler, R. and Rivero, R.M., 2016. Accumulation of flavonols over hydroxycinnamic acids favors oxidative damage protection under abiotic stress. *Frontiers in Plant Science*, 7, p.838.

Matsumoto, K., Cabral, G., Teixeira, J. and Monte, D., 2009. Agrobacterium-mediated transient expression system in banana immature fruits. *African Journal of Biotechnology*, 8(17), pp. 4039-4042.

Matus, J.T., Loyola, R., Vega, A., Peña-Neira, A., Bordeu, E., Arce-Johnson, P. and Alcalde, J.A., 2009. Post-veraison sunlight exposure induces MYB-mediated transcriptional regulation of anthocyanin and flavonol synthesis in berry skins of *Vitis vinifera*. *Journal of experimental botany*, 60(3), pp.853-867.

Mehra, P., Pandey, B.K., Verma, L. and Giri, J., 2019. A novel glycerophosphodiester phosphodiesterase improves phosphate deficiency tolerance in rice. *Plant, Cell & Environment*, 42(4), pp.1167-1179.

Mehra, P., Pandey, B.K., Verma, L., Prusty, A., Singh, A.P., Sharma, S., Malik, N., Bennett, M.J., Parida, S.K., Giri, J. and Tyagi, A.K., 2022. OsJAZ11 regulates spikelet and seed development in rice. *Plant Direct*, 6(5), p.e401.

Mehrtens, F., Kranz, H., Bednarek, P. and Weisshaar, B., 2005. The Arabidopsis transcription factor MYB12 is a flavonol-specific regulator of phenylpropanoid biosynthesis. *Plant Physiology*, 138(2), pp.1083-1096.

Mikkelsen, B.L., Olsen, C.E. and Lyngkjær, M.F., 2015. Accumulation of secondary metabolites in healthy and diseased barley, grown under future climate levels of CO2, ozone and temperature. *Phytochemistry*, 118, pp.162-173.

Misra, P., Pandey, A., Tiwari, M., Chandrashekar, K., Sidhu, O.P., Asif, M.H., Chakrabarty, D., Singh, P.K., Trivedi, P.K., Nath, P. and Tuli, R., 2010. Modulation of transcriptome and metabolome of tobacco by Arabidopsis transcription factor, *AtMYB12*, leads to insect resistance. *Plant Physiology*, 152(4), pp.2258-2268.

Muhlemann, J.K., Younts, T.L. and Muday, G.K., 2018. Flavonols control pollen tube growth and integrity by regulating ROS homeostasis during high-temperature stress. *Proceedings of the National Academy of Sciences*, 115(47), pp.E11188-E11197.

Murcia, G., Fontana, A., Pontin, M., Baraldi, R., Bertazza, G. and Piccoli, P.N., 2017. ABA and GA3 regulate the synthesis of primary and secondary metabolites related to alleviation from biotic and abiotic stresses in grapevine. *Phytochemistry*, 135, pp.34-52.

Muthusamy, M., Uma, S., Backiyarani, S., Saraswathi, M.S. and Chandrasekar, A., 2016. Transcriptomic changes of drought-tolerant and sensitive banana cultivars exposed to drought stress. *Frontiers in Plant Science*, 7, pp.1609.

Nagamatsu, A., Masuta, C., Senda, M., Matsuura, H., Kasai, A., Hong, J.S., Kitamura, K., Abe, J. and Kanazawa, A., 2007. Functional analysis of soybean genes involved in flavonoid biosynthesis by virus□induced gene silencing. *Plant Biotechnology Journal*, 5(6), pp.778-790.

Naik, J., Misra, P., Trivedi, P.K. and Pandey, A., 2022. Molecular components associated with the regulation of flavonoid biosynthesis. *Plant Science*, 317, p.111196.

Naik, J., Rajput, R., Pucker, B., Stracke, R. and Pandey, A., 2021. The R2R3-MYB transcription factor MtMYB134 orchestrates flavonol biosynthesis in *Medicago truncatula*. *Plant Molecular Biology*, 106(1), pp.157-172.

Oh, M.M., Trick, H.N. and Rajashekar, C.B., 2009. Secondary metabolism and antioxidants are involved in environmental adaptation and stress tolerance in lettuce. *Journal of Plant Physiology*, 166(2), pp.180-191.

Ohama, N., Sato, H., Shinozaki, K. and Yamaguchi-Shinozaki, K., 2017. Transcriptional regulatory network of plant heat stress response. *Trends in Plant Science*, 22(1), pp.53-65.

Pandey, A., Alok, A., Lakhwani, D., Singh, J., Asif, M.H. and Trivedi, P.K., 2016. Genome-wide expression analysis and metabolite profiling elucidate transcriptional regulation of flavonoid biosynthesis and modulation under abiotic stresses in banana. *Scientific Reports*, 6(1), pp.1-13.

Pandey, A., Misra, P. and Trivedi, P.K., 2015. Constitutive expression of Arabidopsis MYB transcription factor, *AtMYB11*, in tobacco modulates flavonoid biosynthesis in favor of flavonol accumulation. *Plant cell reports*, 34(9), pp.1515-1528.

Pandey, A., Misra, P., Bhambhani, S., Bhatia, C. and Trivedi, P.K., 2014b. Expression of Arabidopsis MYB transcription factor, *AtMYB111*, in tobacco requires light to modulate flavonol content. *Scientific Reports*, 4(1), p.5018. https://doi.org/10.1038/srep05018.

Pandey, A., Misra, P., Choudhary, D., Yadav, R., Goel, R., Bhambhani, S., Sanyal, I., Trivedi, R. and Kumar Trivedi, P., 2015. *AtMYB12* expression in tomato leads to large scale differential modulation in transcriptome and flavonoid content in leaf and fruit tissues. *Scientific Reports*, 5(1), pp.1-14.

Pandey, A., Misra, P., Khan, M.P., Swarnkar, G., Tewari, M.C., Bhambhani, S., Trivedi, R., Chattopadhyay, N. and Trivedi, P.K., 2014a. Co☐expression of Arabidopsis transcription factor, *AtMYB12*, and soybean isoflavone synthase, *GmIFS1*, genes in tobacco leads to enhanced biosynthesis of isoflavones and flavonols resulting in osteoprotective activity. *Plant Biotechnology Journal*, 12(1), pp.69-80.

Paupière, M.J., Müller, F., Li, H., Rieu, I., Tikunov, Y.M., Visser, R.G. and Bovy, A.G., 2017. Untargeted metabolomic analysis of tomato pollen development and heat stress response. *Plant Reproduction*, 30, pp.81-94.

Peng, Y., Thrimawithana, A.H., Cooney, J.M., Jensen, D.J., Espley, R.V. and Allan, A.C., 2020. The proanthocyanin-related transcription factors MYBC1 and WRKY44 regulate branch points in the kiwifruit anthocyanin pathway. *Scientific Reports*, 10(1), pp.1-19.

Pipas, J.M. and Levine, A.J., 2001, February. Role of T antigen interactions with p53 in tumorigenesis. *Seminars in Cancer Biology,* 11(1), pp. 23-30.

Pucker, B., Pandey, A., Weisshaar, B. and Stracke, R., 2020. The R2R3-MYB gene family in banana (*Musa acuminata*): Genome-wide identification, classification and expression patterns. *PloS one*, 15(10), pp.e0239275.

Rajput, R., Naik, J., Stracke, R. and Pandey, A., 2022a. Interplay between R2R3 MYB-type activators and repressors regulates proanthocyanidin biosynthesis in banana (*Musa acuminata*). *New Phytologist*, 236, pp.1108-1127.

Rajput, R., Tyagi, S., Naik, J., Pucker, B., Stracke, R. and Pandey, A., 2022b. The R2R3-MYB gene family in *Cicer arietinum*: genome-wide identification and expression analysis leads to functional characterization of proanthocyanidin biosynthesis regulators in the seed coat. *Planta*, 256(4), pp.1-24.

Rao, S., Das, J.R. and Mathur, S., 2021. Exploring the master regulator heat stress transcription factor HSFA1a-mediated transcriptional cascade of HSFs in the heat stress response of tomato. *Journal of Plant Biochemistry and Biotechnology*, 30, pp.878-888.

Remizowa, M.V. and Sokoloff, D.D., 2023. Patterns of Carpel Structure, Development, and Evolution in Monocots. *Plants*, 12(24), p.4138.https://doi.org/10.3390/plants12244138.

Sanger, F., Nicklen, S. and Coulson, A.R., 1977. DNA sequencing with chain-terminating inhibitors. *Proceedings of the National Academy of Sciences*, 74(12), pp.5463-5467.

Saxena, S., Pal, L., Naik, J., Singh, Y., Verma, P.K., Chattopadhyay, D. and Pandey, A., 2023. The R2R3□MYB□SG7 transcription factor CaMYB39 orchestrates surface phenylpropanoid metabolism and pathogen resistance in chickpea. *New Phytologist*, 238(2), pp.798-816.

Scharf, K.D., Berberich, T., Ebersberger, I. and Nover, L., 2012. The plant heat stress
transcription factor (Hsf) family: structure, function and evolution. *Biochimica et Biophysica*
Acta (BBA)-Gene Regulatory Mechanisms, 1819(2), pp.104-119.

Schramm, F., Larkindale, J., Kiehlmann, E., Ganguli, A., Englich, G., Vierling, E. and Von
Koskull□Döring, P., 2008. A cascade of transcription factor DREB2A and heat stress
transcription factor HsfA3 regulates the heat stress response of Arabidopsis. *The Plant*
Journal, 53(2), pp.264-274.

Schwinn, K.E., Ngo, H., Kenel, F., Brummell, D.A., Albert, N.W., McCallum, J.A., Pither-
Joyce, M., Crowhurst, R.N., Eady, C. and Davies, K.M., 2016. The onion *(Allium cepa* L.)
R2R3-MYB gene MYB1 regulates anthocyanin biosynthesis. *Frontiers in Plant Science*, 7,
p.1865.

Shan, W., KUANG, J.F., LU, W.J. and CHEN, J.Y., 2014. Banana fruit NAC transcription
factor MaNAC 1 is a direct target of MaICE1 and involved in cold stress through interacting
with MaCBF1. *Plant, cell & environment*, 37(9), pp.2116-2127.

Shan, X., Li, Y., Yang, S., Yang, Z., Qiu, M., Gao, R., Han, T., Meng, X., Xu, Z., Wang, L.
and Gao, X., 2020. The spatio□temporal biosynthesis of floral flavonols is controlled by
differential phylogenetic MYB regulators in *Freesia hybrida*. *New Phytologist*, 228(6),
pp.1864-1879.

Shi, C., Luo, P., Du, Y.T., Chen, H., Huang, X., Cheng, T.H., Luo, A., Li, H.J., Yang, W.C.,
Zhao, P. and Sun, M.X., 2019. Maternal control of suspensor programmed cell death via
gibberellin signaling. *Nature Communications*, 10(1), pp.1-12.

Shih, C.H., Chu, H., Tang, L.K., Sakamoto, W., Maekawa, M., Chu, I.K., Wang, M. and Lo,
C., 2008. Functional characterization of key structural genes in rice flavonoid biosynthesis.
Planta, 228(6), pp.1043-1054.

Si, J., Fan, Z.Q., Wu, C.J., Yang, Y.Y., Shan, W., Kuang, J.F., Lu, W.J., Wei, W. and Chen,
J.Y., 2024. MaHsf24, a novel negative modulator, regulates cold tolerance in banana fruits by
repressing the expression of HSPs and antioxidant enzyme genes. *Plant Biotechnology*
Journal. https://doi.org/10.1111/pbi.14410.

Si, J., Ye, B.B., Liu, Z.L., Xiao, X.M., Yang, Y.Y., Fan, Z.Q., Shan, W., Kuang, J.F., Lu,
W.J., Su, X.G. and Chen, J.Y., 2022. Transcriptional repression of MaRBOHs by MaHsf26 is

associated with heat shock-alleviated chilling injury in banana fruit. *Postharvest Biology and Technology*, 193, p.112056.

Song, S.S., Ran, W.X., Gao, L.H., Wang, Y.C., Lv, W.Y., Tao, Y., Chen, L. and Li, C.F., 2024. A functional study reveals CsNAC086 regulated the biosynthesis of flavonols in *Camellia sinensis*. Planta, 259(6), p.147.

Song, Z., Lai, X., Chen, H., Wang, L., Pang, X., Hao, Y., Lu, W., Chen, W., Zhu, X. and Li, X., 2022. Role of MaABI5-like in abscisic acid-induced cold tolerance of 'Fenjiao'banana fruit. *Horticulture Research*, 9, p.uhac130.

Song, Z., Lai, X., Chen, H., Wang, L., Yao, Y., Chen, W., Zhu, X. and Li, X., 2023. MaC2H2-like regulates chilling stress response of 'Fenjiao'banana by modulating flavonoid synthesis and fatty acid desaturation. *Food Chemistry*, 419, p.136089.

Song, Z., Luo, Y., Wang, W., Fan, N., Wang, D., Yang, C. and Jia, H., 2020. NtMYB12 positively regulates flavonol biosynthesis and enhances tolerance to low Pi stress in *Nicotiana tabacum. Frontiers in Plant Science*, 10, p.1683.

Stracke, R., Favory, J.J., Gruber, H., Bartelniewoehner, L., Bartels, S., Binkert, M., Funk, M., Weisshaar, B. and Ulm, R., 2010. The Arabidopsis bZIP transcription factor HY5 regulates expression of the PFG1/MYB12 gene in response to light and ultraviolet B radiation. *Plant, Cell & Environment*, 33(1), pp.88-103.

Stracke, R., Holtgräwe, D., Schneider, J., Pucker, B., Rosleff Sörensen, T. and Weisshaar, B., 2014. Genome-wide identification and characterisation of R2R3-MYB genes in sugar beet (*Beta vulgaris*). *BMC Plant Biology*, 14(1), pp.1-17.

Stracke, R., Ishihara, H., Huep, G., Barsch, A., Mehrtens, F., Niehaus, K. and Weisshaar, B., 2007. Differential regulation of closely related R2R3 MYB transcription factors controls flavonol accumulation in different parts of the Arabidopsis thaliana seedling. *The Plant Journal*, 50(4), pp.660-677.

Stracke, R., Jahns, O., Keck, M., Tohge, T., Niehaus, K., Fernie, A.R. and Weisshaar, B., 2010. Analysis of PRODUCTION OF FLAVONOL GLYCOSIDES dependent flavonol glycoside accumulation in *Arabidopsis thaliana* plants reveals MYB11 , MYB12 and MYB111 independent flavonol glycoside accumulation. *New Phytologist*, 188(4), pp.985-1000.

Stracke, R., Werber, M. and Weisshaar, B., 2001. The R2R3-MYB gene family in *Arabidopsis thaliana*. *Current Opinion in Plant Biology*, 4(5), pp.447-456.

Tan, H., Man, C., Xie, Y., Yan, J., Chu, J. and Huang, J., 2019. A crucial role of GA-regulated flavonol biosynthesis in root growth of Arabidopsis. *Molecular Plant*, 12(4), pp.521-537.

Teale, W.D., Pasternak, T., Dal Bosco, C., Dovzhenko, A., Kratzat, K., Bildl, W., Schwörer, M., Falk, T., Ruperti, B., V Schaefer, J. and Shahriari, M., 2021. Flavonol mediated stabilization of PIN efflux complexes regulates polar auxin transport. *The EMBO Journal*, 40(1), p.e104416.

von Koskull-Döring, P., Scharf, K.D. and Nover, L., 2007. The diversity of plant heat stress transcription factors. *Trends in Plant Science*, 12(10), pp.452-457.

Waki, T., Yoo, D., Fujino, N., Mameda, R., Denessiouk, K., Yamashita, S., Motohashi, R., Akashi, T., Aoki, T., Ayabe, S.I. and Takahashi, S., 2016. Identification of protein–protein interactions of isoflavonoid biosynthetic enzymes with 2-hydroxyisoflavanone synthase in soybean (*Glycine max* (L.) Merr.). *Biochemical and biophysical research communications*, 469(3), pp.546-551.

Walter, M., Chaban, C., Schütze, K., Batistic, O., Weckermann, K., Näke, C., Blazevic, D., Grefen, C., Schumacher, K., Oecking, C. and Harter, K., 2004. Visualization of protein interactions in living plant cells using bimolecular fluorescence complementation. *The Plant Journal*, 40(3), pp.428-438.

Wang, J., Ren, Y., Liu, X., Luo, S., Zhang, X., Liu, X., Lin, Q., Zhu, S., Wan, H., Yang, Y. and Zhang, Y., 2021a. Transcriptional activation and phosphorylation of OsCNGC9 confer enhanced chilling tolerance in rice. *Molecular Plant*, 14(2), pp.315-329.

Wang, N., Liu, W., Yu, L., Guo, Z., Chen, Z., Jiang, S., Xu, H., Fang, H., Wang, Y., Zhang, Z. and Chen, X., 2020. HEAT SHOCK FACTOR A8a modulates flavonoid synthesis and drought tolerance. *Plant Physiology*, 184(3), pp.1273-1290.

Wang, N., Xu, H., Jiang, S., Zhang, Z., Lu, N., Qiu, H., Qu, C., Wang, Y., Wu, S. and Chen, X., 2017. MYB12 and MYB22 play essential roles in proanthocyanidin and flavonol synthesis in red fleshed apple (*Malus sieversii* f. niedzwetzkyana). *The Plant Journal*, 90(2), pp.276-292.

Wang, X., Niu, Y. and Zheng, Y., 2021b. Multiple functions of MYB transcription factors in abiotic stress responses. *International Journal of Molecular Sciences,* 22(11), p.6125.

Wang, X., Wang, L., Wang, Y., Liu, H., Hu, D., Zhang, N., Zhang, S., Cao, H., Cao, Q., Zhang, Z. and Tang, S., 2018. Arabidopsis PCaP2 plays an important role in chilling tolerance and ABA response by activating CBF-and SnRK2-mediated transcriptional regulatory network. *Frontiers in Plant Science*, 9, p.215.

Wang, X., Yu, R. and Li, J., 2021c. Using genetic engineering techniques to develop banana cultivars with fusarium wilt resistance and ideal plant architecture. *Frontiers in Plant Science*, 11, p.617528.

Wang, Y., Mao, Z., Jiang, H., Zhang, Z., Wang, N. and Chen, X., 2021d. Brassinolide inhibits flavonoid biosynthesis and red-flesh coloration via the MdBEH2. 2–MdMYB60 complex in apple. Journal of experimental botany, 72(18), pp.6382-6399.

Wang, Z., Huang, S., Jia, C., Liu, J., Zhang, J., Xu, B. and Jin, Z., 2013. Molecular cloning and expression of five glutathione S-transferase (GST) genes from Banana (*Musa acuminata* L. AAA group, cv. Cavendish). *Plant Cell Reports*, 32(9), pp.1373-1380.

Wang, Z., Li, Z.F., Wang, S.S., Xiao, Y.S., Xie, X.D., Wu, M.Z., Yu, J.L., Cheng, L.R., Yang, A.G. and Yang, J., 2021e. NtMYB12a acts downstream of sucrose to inhibit fatty acid accumulation by targeting lipoxygenase and SFAR genes in tobacco. *Plant, Cell & Environment*, 44(3), pp.775-791.

Wang, Z., Luo, Z., Liu, Y., Li, Z., Liu, P., Bai, G., Zhou, Z., Xie, H. and Yang, J., 2021f. Molecular cloning and functional characterization of NtWRKY11b in promoting the biosynthesis of flavonols in *Nicotiana tabacum. Plant Science*, 304, p.110799.

Watkins, J.M., Chapman, J.M. and Muday, G.K., 2017. Abscisic acid-induced reactive oxygen species are modulated by flavonols to control stomata aperture. *Plant Physiology*, 175(4), pp.1807-1825.

Wei, W., Yang, Y.Y., Lakshmanan, P., Kuang, J.F., Lu, W.J., Pang, X.Q., Chen, J.Y. and Shan, W., 2023a. Proteasomal degradation of MaMYB60 mediated by the E3 ligase MaBAH1 causes high temperature-induced repression of chlorophyll catabolism and green ripening in banana. *The Plant Cell*, 35(5), pp.1408-1428.

Wei, Y., Hu, W., Xia, F., Zeng, H., Li, X., Yan, Y., He, C. and Shi, H., 2016. Heat shock transcription factors in banana: genome-wide characterization and expression profile analysis during development and stress response. *Scientific Reports*, 6(1), pp.1-11.

Wei, Y., Meng, N., Wang, Y., Cheng, J., Duan, C. and Pan, Q., 2023b. Transcription factor VvWRKY70 inhibits both norisoprenoid and flavonol biosynthesis in grape. *Plant Physiology*, 193(3), pp.2055-2070.

Wu, C., Shan, W., Liang, S., Zhu, L., Guo, Y., Chen, J., Lu, W., Li, Q., Su, X. and Kuang, J., 2019. MaMPK2 enhances MabZIP93-mediated transcriptional activation of cell wall modifying genes during banana fruit ripening. *Plant Molecular Biology*, 101, pp.113-127.

Wu, T.Y., Hoh, K.L., Boonyaves, K., Krishnamoorthi, S. and Urano, D., 2022. Diversification of heat shock transcription factors expanded thermal stress responses during early plant evolution. *The Plant Cell*, 34(10), pp.3557-3576.

Wu, Y.L., Yi, G.J. and Peng, X.X., 2010. Rapid screening of *Musa* species for resistance to Fusarium wilt in an *in vitro* bioassay. *European Journal of Plant Pathology*, 128, pp.409-415.

Xiao, X.M., Li, L.L., Kuang, J.F., Chen, J.Y., Lu, W.J., Wei, W. and Shan, W., 2023. Cold pretreatment promotes chlorophyll degradation of green ripening banana peel by activating MaCBF1 to *MaCBR* and *MaSGR1*. *Food Chemistry*, 413, p.135575. doi.org/10.1016/j.foodchem.2023.135575.

Xiao, Y.Y., Kuang, J.F., Qi, X.N., Ye, Y.J., Wu, Z.X., Chen, J.Y. and Lu, W.J., 2018. A comprehensive investigation of starch degradation process and identification of a transcriptional activator MabHLH6 during banana fruit ripening. *Plant Biotechnology Journal,* 16(1), pp.151-164.

Xu, F., Li, L., Zhang, W., Cheng, H., Sun, N., Cheng, S. and Wang, Y., 2012. Isolation, characterization, and function analysis of a flavonol synthase gene from *Ginkgo biloba*. *Molecular Biology Reports*, 39(3), pp.2285-2296.

Xu, Y., Hu, W., Liu, J., Song, S., Hou, X., Jia, C., Li, J., Miao, H., Wang, Z., Tie, W. and Xu, B., 2020. An aquaporin gene MaPIP2-7 is involved in tolerance to drought, cold and salt stresses in transgenic banana (*Musa acuminata* L.). *Plant Physiology and Biochemistry*, 147, pp.66-76.

Yang, F., Wang, T., Guo, Q., Zou, Q. and Yu, S., 2023. The CmMYB3 transcription factors

isolated from the *Chrysanthemum morifolium* regulate flavonol biosynthesis in Arabidopsis

thaliana. *Plant Cell Reports*, 42(4), pp.791-803.

Yang, J., Guo, C., Chen, F., Lv, B., Song, J., Ning, G., He, Y., Lin, J., He, H., Yang, Y. and

Xiang, F., 2024. Heat-induced modulation of flavonoid biosynthesis via a LhMYBC2-

Mediated regulatory network in oriental hybrid lily. *Plant Physiology and Biochemistry*,

p.108966. https://doi.org/10.1016/j.plaphy.2024.10896.

Yang, Q., Song, Z., Dong, B., Niu, L., Cao, H., Li, H., Du, T., Liu, T., Yang, W., Meng, D.

and Fu, Y., 2021. Hyperoside regulates its own biosynthesis via MYB30 in promoting

reproductive development and seed set in okra. *Plant Physiology*, 185(3), pp.951-968.

Yoshida, T., Fujita, Y., Maruyama, K., Mogami, J., Todaka, D., Shinozaki, K. and

Yamaguchi□Shinozaki, K.A.Z.U.K.O., 2015. Four Arabidopsis AREB/ABF transcription

factors function predominantly in gene expression downstream of SnRK2 kinases in abscisic

acid signalling in response to osmotic stress. *Plant, Cell & Environment*, 38(1), pp.35-49.

Yoshida, T., Ohama, N., Nakajima, J., Kidokoro, S., Mizoi, J., Nakashima, K., Maruyama,

K., Kim, J.M., Seki, M., Todaka, D. and Osakabe, Y., 2011. Arabidopsis HsfA1 transcription

factors function as the main positive regulators in heat shock-responsive gene expression.

Molecular Genetics and Genomics, 286(5), pp.321-332.

Zhang, J., Subramanian, S., Stacey, G. and Yu, O., 2009. Flavones and flavonols play distinct

critical roles during nodulation of *Medicago truncatula* by *Sinorhizobium meliloti*. *The Plant

Journal*, 57(1), pp.171-183.

Zhang, K., Logacheva, M.D., Meng, Y., Hu, J., Wan, D., Li, L., Janovská, D., Wang, Z.,

Georgiev, M.I., Yu, Z. and Yang, F., 2018. Jasmonate-responsive MYB factors spatially

repress rutin biosynthesis in *Fagopyrum tataricum*. *Journal of Experimental Botany*, 69(8),

pp.1955-1966.

Zhang, L., Cenci, A., Rouard, M., Zhang, D., Wang, Y., Tang, W. and Zheng, S.J., 2019.

Transcriptomic analysis of resistant and susceptible banana corms in response to infection by

Fusarium oxysporum f. sp. cubense tropical race 4. *Scientific Reports*, 9(1), pp.1-14.

Zhang, X., He, Y., Li, L., Liu, H. and Hong, G., 2021. Involvement of the R2R3-MYB transcription factor MYB21 and its homologs in regulating flavonol accumulation in Arabidopsis stamen. *Journal of Experimental Botany*, 72(12), pp.4319-4332.

Zhang, X., Li, L., He, Y., Lang, Z., Zhao, Y., Tao, H., Li, Q. and Hong, G., 2023. The CsHSFA□CsJAZ6 module□mediated high temperature regulates flavonoid metabolism in *Camellia sinensis*. *Plant, Cell & Environment*, 46(8), pp.2401-2418.

Zhao, C., Liu, X., Gong, Q., Cao, J., Shen, W., Yin, X., Grierson, D., Zhang, B., Xu, C., Li, X. and Chen, K., 2021. Three AP2/ERF family members modulate flavonoid synthesis by regulating type IV chalcone isomerase in citrus. *Plant Biotechnology Journal*, 19(4), pp.671-688.

Zhao, L., Hu, Y., Chong, K. and Wang, T., 2010. ARAG1, an ABA-responsive DREB gene, plays a role in seed germination and drought tolerance of rice. *Annals of Botany*, 105(3), pp.401-409.

Zhao, M.L., Wang, J.N., Shan, W.E.I., Fan, J.G., Kuang, J.F., Wu, K.Q., Li, X.P., Chen, W.X., He, F.Y., Chen, J.Y. and Lu, W.J., 2013. Induction of jasmonate signalling regulators MaMYC2s and their physical interactions with MaICE1 in methyl jasmonate□induced chilling tolerance in banana fruit. *Plant, Cell & Environment*, 36(1), pp.30-51.

Zhao, X., Zeng, X., Lin, N., Yu, S., Fernie, A.R. and Zhao, J., 2021. CsbZIP1-CsMYB12 mediates the production of bitter-tasting flavonols in tea plants (*Camellia sinensis*) through a coordinated activator–repressor network. *Horticulture Research*, 8. https://doi.org/10.1038/s41438-021-00545-8.

Zheng, L., Li, B., Zhang, G., Zhou, Y. and Gao, F., 2024. Jasmonate enhances cold acclimation in jojoba by promoting flavonol synthesis. *Horticulture Research*, p.uhae125. https://doi.org/10.1093/hr/uhae125.

Zhou, L., He, Y., Li, J., Liu, Y. and Chen, H., 2020. CBFs function in anthocyanin biosynthesis by interacting with MYB113 in eggplant (*Solanum melongena* L.). *Plant and Cell Physiology*, 61(2), pp.416-426.

Zhou, L.J., Geng, Z., Wang, Y., Wang, Y., Liu, S., Chen, C., Song, A., Jiang, J., Chen, S. and Chen, F., 2021. A novel transcription factor CmMYB012 inhibits flavone and anthocyanin

biosynthesis in response to high temperatures in chrysanthemum. Horticulture research, 8.

https://doi.org/10.1038/s41438-021-00675-z.

Zimmermann, I.M., Heim, M.A., Weisshaar, B. and Uhrig, J.F., 2004. Comprehensive

identification of Arabidopsis thaliana MYB transcription factors interacting with R/B□like

BHLH proteins. *The Plant Journal*, 40(1), pp.22-34.

Zinsmeister, J., Berriri, S., Basso, D.P., Ly□Vu, B., Dang, T.T., Lalanne, D., da Silva,

E.A.A., Leprince, O. and Buitink, J., 2020. The seed□specific heat shock factor A9 regulates

the depth of dormancy in *Medicago truncatula* seeds via ABA signalling. *Plant, Cell &*

Environment, 43(10), pp.2508-2522.

(a)

LB // *pPvUB*I intron *GUSPlus* nosT *pZmUBI1* intron *MaMYBFA* AcV5 tag OCS T RB

(b)

Empty vector (EV)   *MaMYBFA1*-OE   *MaMYBFA2*-OE   *MaMYBFA3*-OE

(c)

MaMYBFA1 — Relative expression — EV, MaMYBFA1-OX (****), MaMYBFA2-OX (ns), MaMYBFA3-OX (ns)

MaMYBFA2 — Relative expression — EV, MaMYBFA1-OX (ns), MaMYBFA2-OX (***), MaMYBFA3-OX (ns)

MaMYBFA3 — Relative expression — EV, MaMYBFA1-OX (ns), MaMYBFA2-OX (ns), MaMYBFA3-OX (****)

(e)

Kaempferol — µg/100g DW — EV, MaMYBFA1-OE (*), MaMYBFA2-OE (*), MaMYBFA3-OE (*)

Quercetin — µg/100g DW — EV, MaMYBFA1-OE (ns), MaMYBFA2-OE (ns), MaMYBFA3-OE (**)

(d)

MaFLS1 — Relative expression — EV, MaMYBFA1-OX (***), MaMYBFA2-OX (***), MaMYBFA3-OX (***)

MaFLS2 — Relative expression — EV, MaMYBFA1-OX (**), MaMYBFA2-OX (*), MaMYBFA3-OX (ns)

Myricetin — µg/100g DW — EV, MaMYBFA1-OE (ns), MaMYBFA2-OE (*), MaMYBFA3-OE (ns)

(a)

(b) CDS-YFP   NLS-RFP   Merged

MaHSF11-YFP

Free YFP

(c) MaHSF11

(d) EV   *ZmUBI:MaHSF11-OE*

(e) EV   *MaHSF11-OE*

(f) Kaempferol   Quercetin   Myricetin

(g) EV   *ZmUBI:MaHSF11-RNAi*

(h) EV   *MaHSF11-RNAi*

(i) Kaempferol   Quercetin   Myrecetin

(a) *MaDREB1*
Before drought
After drought
Log2(FPKM)
0 1 2 3 4 5 6 7

(b)
EV
MaHSF11-OE
Relative expression
***
MaDREB1

(c)
EV
MaHSF11-RNAi
Relative expression
ns
MaDREB1

(d) *proMaDREB1*
+310
ATG
F2   F1
HSE   HSE   -1
(TGAAGCTTCG) (AGAAGCTTCA)
-688

(e)
-AbA   +AbA
proMaDREB1 (F2)   pGADT7 (EV)
MaHSF11-AD
proMaDREB1 (F1)   pGADT7 (EV)
MaHSF11-AD

(f) *proMaDREB1-998*
(n=4)
LUC/REN ratio
*
EV   MaHSF11

(g) *pro35S*:YFP-MaHSF11
anti-IgG AB
anti-GFP AB
proMaDREB1 Relative fold enrichment
ns   ns
F1   F2

(h)
F1 : GGCTTGGGAAGGGTAGAGAAGCTTCAGGGGATGCTTCTTG
F1m : GGCTTGGGAAGGGTAGACCCCCCCCAGGGGATGCTTCTTG

GST   +   -   -   -   -
GST-MaHSF11   -   +   +   +   +
Biotin labelled probe   +   +   +   +   -
Unlabelled cold probe   -   -   +   +   -
Biotin labelled mutated probe   -   -   -   -   +

proMaDREB1-F1 shifted probe →

proMaDREB1-F1 free probe