The HSF–DREB–MYB transcriptional regulatory module regulates flavonol biosynthesis
and flavonoid B-ring hydroxylation in banana (*Musa acuminata*)

Jogindra Naik[1], Ruchika Rajput[1], Ralf Stracke[2], Ashutosh Pandey[1*]

[1]National Institute of Plant Genome Research, Aruna Asaf Ali Marg, New Delhi 110067, India

[2]Bielefeld University, Faculty of Biology, Genetics and Genomics of Plants, 33615 Bielefeld,
Germany

Short title: HSF–DREB–MYB module regulates flavonol biosynthesis in banana

[*]**Corresponding author:**

Ashutosh Pandey

National Institute of Plant Genome Research

New Delhi 110067, India

Email: ashutosh@nipgr.ac.in

Abstract

Plant flavonols act primarily as ultraviolet radiation absorbers, reactive oxygen species
scavengers, and phytoalexins, and they contribute to biotic and abiotic stress tolerance in
plants. Banana (*Musa acuminata*), an herbaceous monocot and important fruit crop,
accumulates flavonol derivatives in different organs, including the edible fruit pulp. Although
flavonol content varies greatly in different organs, the molecular mechanisms involving
transcriptional regulation of flavonol synthesis in banana are not known. Here, we
characterized three SG7-R2R3 MYB transcription factors MaMYBFA1, MaMYBFA2, and
MaMYBFA3) and their upstream regulators, heat shock transcription factor (MaHSF11) and
dehydration responsive element binding factor (MaDREB1), to elucidate the molecular
mechanism involved in transcriptional regulation of flavonol biosynthesis in banana.
MaMYBFA positively regulate *flavonol synthase* 2 (*MaFLS2)* and downregulates *MaFLS1*.
We show these transcription factors to be weak regulators of flavonol synthesis.

Overexpression of *MaHSF11* enhances flavonol contents, particularly that of myricetin, and

promotes flavonol B-ring hydroxylation, which contributes to the diversity of flavonol

derivatives. MaHSF11 directly interacts with the *MaFLS1* and *flavonoid 3′, 5′-hydroxylase*1

(*MaF3′5′H1)* promoters, both *in vitro* and *in vivo*. MaHSF11 activates the expression of

MaDREB1 directly, which in turn regulates the expression of *MaMYBFA3*. Overall, our study

elucidates a key regulatory mechanism for flavonol synthesis in banana and suggests possible

targets for genetic optimization to enhance nutritional value and stress responses in this

globally important fruit crop.

Keywords: Flavonol biosynthesis, MYB transcription factors, ABA, banana, HSF
transcription factors, transcriptional regulatory module, metabolon

Introduction

Plants produce a multitude of chemical compounds in response and adaptation to external and

environmental stimuli, many of which are specialized metabolites (Oh *et al.,* 2009; Mikkelsen

et al., 2015; Murcia *et al.,* 2017). Among these, flavonoids are ubiquitous and play vital roles

in both regulation of developmental processes and defense against pathogens (Halbwirth *et al.,*

2010; Maloney *et al.,* 2014; Chen *et al.,* 2019; Yang *et al.,* 2021). Based on their structures,

flavonoids are classified into flavonols, flavones, flavanones, isoflavones, anthocyanins, and

proanthocyanidins (Lago *et al.,* 2014). Flavonols have a basic 16-carbon, triple-ring structure

containing a 3-hydroxyflavone backbone, and the different modifications and substituents in

this ring give rise to their diversity (Ko *et al.,* 2014). Besides acting as co-pigments with

anthocyanins in flower petals and as reactive oxygen species (ROS) scavengers, flavonols have

numerous other important functions in plants (Kitamura *et al.,* 2016; Watkins *et al.,* 2017).

They absorb ultraviolet radiation (UV), protecting plants from UV-induced oxidative damage

(Liang *et al.,* 2020) and also regulate root growth and lateral root development by inhibiting

polar auxin transport via stabilization of PIN-FORMED (PIN) protein dimers (Tan *et al.,* 2019;

Chapman *et al.,* 2021; Teale *et al.,* 2021). In addition, flavonols regulate pollen development

and germination, nodule development, and symbiosis (Zhang *et al.,* 2009; Muhlemann *et al.,*

2018). Flavonols are synthesized in early steps of the flavonoid biosynthesis pathway involving

the enzymes chalcone synthase (CHS), chalcone isomerase (CHI), flavanone 3-hydroxylase

(F3H), flavonoid 3′ hydroxylase (F3′H), flavonoid 3′5′ hydroxylase (F3′5′H), and flavonol

synthase (FLS), which are well characterized in many plant species (Nagamatsu *et al.,* 2007;

Yu *et al.,* 2015; Guo *et al.,* 2019; Jiang *et al.,* 2020).

The flavonol biosynthesis pathway is regulated at several levels, and its transcriptional

regulation has been extensively studied (Matus *et al.,* 2009). A group of MYB transcription

factors (TFs), known as subgroup 7 (SG7) R2R3-MYBs, play a major role in the regulation of

flavonol biosynthesis (Schwinn *et al.,* 2016); R2 and R3 are conserved regions. MYB12,

MYB111, and MYB11, also known as PRODUCTION OF FLAVONOL GLYCOSIDES1–3

(PFG1–3), respectively, modulate flavonol biosynthesis in a tissue-specific manner in

Arabidopsis (*Arabidopsis thaliana*) (Mehrtens *et al.,* 2005; Stracke *et al*., 2007; Stracke *et al.,*

2010), tobacco (*Nicotiana tabacum*) (Misra *et al.,* 2010; Pandey *et al.,* 2014a, b; Pandey *et al.,*

2015a), and tomato (*Solanum lycopersicum*) (Pandey *et al.,* 2015b). They regulate the

expression of several rate-limiting biosynthesis genes, including *CHS*, *F3H*, and *FLS*. *FLS* is

also coordinately regulated by MYB99, MYB21, and MYB24 in Arabidopsis stamens and

anthers independently of the PFGs (Battat *et al.,* 2019; Zhang *et al.,* 2021).

Because plants are exposed to numerous biotic and abiotic stresses, such as excessive

light and UV radiation, suboptimal water conditions (drought and waterlogging), high salt,

extreme temperatures, nutrient deficiencies, and pathogen infection (López *et al.,* 2008; Cramer

et al., 2011; Kalaji *et al.,* 2017; Choudhury *et al.,* 2017), they have evolved various defensive

mechanisms that are activated after exposure to stress and involve a complex interplay of

signaling pathways. Other classes of TFs, such as heat shock factors (HSFs) and dehydration-

responsive element-binding (DREB) factors, modulate the abscisic acid (ABA)–dependent

stress response to drought and high heat (Zhao *et al.,* 2010; Lee *et al.,* 2010; Huang *et al.,* 2016;

Zinsmeister *et al.,* 2020). HSFs, in their dimeric form, regulate several stress-responsive genes

by binding to heat shock elements (HSEs; nGAAnnTTCn/nTTCnnGAAn) via their N-terminal

DNA-binding domains (Liu *et al.,* 2008; Wang *et al.,* 2020). HSFs can be classified into three

major classes, A, B, and C, depending upon the type of amino acid residues and the length of

flexible linker length between DBD (DNA-binding domain) and OD (oligomerization domain)

(Scharf *et al.*, 2012; Manna *et al.,* 2021). In addition to heat stress tolerance, HSFs regulate

salt, drought, and cold stress responses in Arabidopsis (Yoshida *et al.,* 2011). MdHSFA8a is a

positive regulator of drought stress tolerance and flavonoid synthesis in apple (*Malus*

domestica) (Wang *et al.,* 2020). The HSFA2 and HSFA3 heterodimeric complex confers

transcriptional thermal memory by promoting epigenetic H3K4 hypermethylation (Friedrich *et*

al., 2021). DREBs, which belong to the AP2/ERF family of TFs, are involved in several stress

signaling pathways, in particular drought, salt, and cold stress (Dubouzet *et al.,* 2003; Chen *et al.,* 2016). CBF/DREB TFs bind to the cold and dehydration-responsive CRT/DREB motif core element (CCGAC) present in the promoters of their target genes via the AP2/ERF DNA-binding domain (Li *et al.,* 2020; Wang *et al.,* 2021). In addition, DREBs regulate fruit ripening via a complex transcriptional regulatory pathway in banana (*Musa acuminata*) (Kuang *et al.,* 2017).

In terms of global production and monetary value, banana is the fourth most important food crop in the world, after rice (*Oryza sativa*), wheat (*Triticum aestivum*), and maize (*Zea mays*) (Wang *et al.,* 2021). Banana is a monocarpic herbaceous crop with climacteric fruits, which is characterized by postharvest ripening because of cellular respiration and increased ethylene production. However, global gross production is often affected by stresses including drought, cold, salinity, and, most importantly, *Fusarium oxysporum* f. sp. *cubense* 4 (Foc4), which causes a devastating wilt disease (Xu *et al.,* 2010; Dong *et al.,* 2020). We identified and functionally characterized three SG7 R2R3-MYB TFs (MaMYBFA1, MaMYBFA2, and MaMYBFA3) as regulators of flavonol synthesis in banana. These MaMYBFAs positively regulate *MaFLS2* and downregulate *MaFLS1*. MaHSF11 directly regulates the expression of *MaFLS1*, *MaF3'5'H1*, and *MaDREB1*. MaDREB1, in turn, regulates the expression of *MaMYBFA3*. We also found that MaFLS1 transiently interacts with MaF3′5′H2 to form a metabolon that promotes the synthesis of the hydroxylated flavonol myricetin, which has strong antioxidant, anticancer, antidiabetic, and anti-inflammatory activities. Additionally, we identified MaHSF11 as a regulator of cytochrome P_{450} (CYP450) B-ring hydroxylase in banana, where it binds directly to the promoter of *MaF3'5'H1*. Taken together, our results elucidate a key regulatory cascade of flavonol synthesis in the important fruit crop banana.

Results

An MaMYBFA1–3 triad modulates flavonol biosynthesis in banana

Our previous *in silico* study (Pucker *et al.,* 2022) determined that the banana genome contains 285 *R2R3-MYB* genes. Three of the encoded MaMYBs form a clade with previously characterized flavonol regulators in a phylogenetic tree (Fig. 1a). We selected these three candidates for further study and named them *MaMYBFA1* (Ma02_g00290), *MaMYBFA2* (Ma05_g23640), and *MaMYBFA3* (Ma08_g10260), where the abbreviation FA stands for flavonol activator. A multiple protein sequence alignment identified the conserved R2R3-MYB domain and the two SG7-specific, conserved motifs: SG7-1 and SG7-2 (Fig. 1b).

We used RT-qPCR to analyze the expression of these three *MaMYBFA* genes in different organs/tissues of banana variety 'Grand Naine' (Supplemental Fig. S1). All three *MaMYBFA* genes were differentially expressed in the analyzed organs. *MaMYBFA1* was most highly expressed in bract tissue, followed by unripe peel and root. *MaMYBFA2* transcript levels were highest in unripe peel, followed by bract and unripe pulp, and *MaMYBFA3* was expressed most highly in bract followed by pseudostem, young leaf, and unripe pulp. Transcript accumulation of the three genes was lowest in ripe peel and ripe pulp. Overall, *MaMYBFA3* showed the highest levels of expression in all tissues. These expression patterns suggest functional differences between the three MaMYBFAs in terms of upstream regulation and downstream regulatory potential toward biosynthesis genes.

The full-length coding sequences (CDSs) of *MaMYBFA1*, *MaMYBFA2*, and *MaMYBFA3* were PCR-amplified from banana cDNA and used to express the encoded proteins. The subcellular localization of MaMYBFA-YFP fusion proteins was analyzed in *Nicotiana benthamiana* epidermal cells and was restricted to the nucleus, co-localizing with the nuclear marker protein NLS-RFP (Fig. 1c), indicating that MaMYBFA1–3 are nuclear proteins. To analyze the regulatory role of MaMYBFA1–3, we transiently overexpressed the *MaMYBFA1–3* CDSs under the control of the *Zea mays ubiquitin1* promoter (*proZmUBI1*) in immature banana fruit discs (Fig. 2a-b). The activity of a GUS reporter simultaneously transformed on the same T-DNA indicated successful transformation of the fruit tissues (Fig. 1b). RT-qPCR analysis revealed upregulation of the respective *MaMYBFA* transcripts in transgenic tissue samples (Fig. 2c). The expression of selected flavonoid biosynthesis genes was analyzed in MaMYBFA-overexpressing (OE) fruit discs. The expression of *CHALCONE SYNTHASE* (*MaCHS1–6*), *CHALCONE ISOMERASE* (*MaCHI1–2*), *FLAVANONE 3-HYDROXYLASE* (*MaF3H1–2*), and *FLAVONOL SYNTHASE* (*MaFLS1–2*) was modulated to different degrees in the OE fruit discs (Fig. 2d, Supplemental Figure S2). *MaMYBFA1* overexpression enhanced the expression of *MaCHS1*, whereas *MaMYBFA2* and *MaMYBFA3* overexpression did not (Supplemental Fig. S2). *MaCHS4* expression increased significantly with overexpression of all three *MaMYBFA*s. However, *MaCHS3* and *MaCHS5* showed enhanced expression only with overexpression of *MaMYBFA1* and *MaMYBFA3* (Supplemental Fig. S2), and only *MaMYBFA1* and *MaMYBFA2* overexpression increased the *MaCHS6* transcript level. *MaCHI* and *MaF3H* transcripts were also modulated in the transgenic OE fruits: The highest activation of *MaCHI1* resulted from *MaMYBFA1* overexpression, whereas *MaCHI2* transcripts were increased 2-fold by *MaMYBFA3* overexpression (Supplemental Fig.

S2). For *MaF3H1*, a significant increase in the transcript level was observed with *MaMYBFA1* and *MaMYBFA3* overexpression (Supplemental Fig. S2). Genes encoding CYP450 proteins also showed differential modulation; only *MaF3'5'H3* and *MaF3'5'H4* transcripts were upregulated in OE fruits, whereas *MaF3'H* and other *MaF3'5'H* paralogs showed a marked decrease in expression (Supplemental Fig. S2). Finally, we analyzed the expression of *MaFLS1* and *MaFLS2*, the putative primary targets of MaMYBFAs, in transgenic *MaMYBFA* OE fruit discs (Fig. 2d). *MaFLS1* was significantly downregulated, whereas *MaFLS2* transcript levels increased in the *MaMYBFA* OE tissues (Fig. 2d). Overall, we found that overexpression of *MaMYBFA1* and *MaMYBFA3* significantly increased the expression of most structural flavonoid biosynthesis genes, whereas *MaMYBFA2* overexpression either decreased or negligibly affected the transcript levels of most of the genes studied.

To analyze the modulation of flavonol metabolite contents in transgenic *MaMYBFA* OE fruit slices, we performed LC-MS-based, targeted flavonol quantification (Fig. 2e). We found decreased levels of kaempferol and quercetin in all the *MaMYBFA* OE tissues and an increase in myricetin content in *MaMYBFA2* OE compared to the empty vector (EV) control. Although the observed changes were significant for kaempferol, the contents of quercetin and myricetin were not altered significantly by *MaMYBFA1* and *MaMYBFA3* OE. Only *MaMYBFA2* OE significantly increased the accumulation of myricetin (Fig. 2e). Taken together, these results indicate that MaMYBFA1, MaMYBFA2, and MaMYBFA3 act as a MYB regulatory triad for modulating flavonol synthesis in banana.

To test the opposite regulation of *MaFLS1* and *MaFLS2* expression by MaMYBFAs, we examined the promoters of the *MaFLS* genes to identify possible MYB binding sites (MYB core elements, CNGTTR). Structural analyses of a 2556-bp *MaFLS1* (*proMaFLS1–2556*) promoter fragment and a 1500-bp *MaFLS2* (*proMaFLS2–1500*) promoter fragment revealed several MYB core elements (Fig. 3a). To analyze the transactivation capacity of the MaMYBFAs on these *MaFLS* promoter fragments, we performed dual-luciferase assays in *N. benthamiana* leaves. The reporter constructs in the p635nRRF vector used for this assay contain the promoters to be analyzed positioned upstream of the LUC gene encoding firefly luciferase, with simultaneous expression of REN (Renilla luciferase) driven by the constitutive cauliflower mosaic virus (CaMV) 35S promoter. The LUC/REN ratio therefore represents the level of activation by the particular promoter analyzed. The effectors in the pBTdest vector contain the *MaMYBBFA* CDS under the control of the CaMV35S promoter (Fig. 3b). The results of these dual-luciferase assays indicated a decreased activity of *proMaFLS1–2556* in

193 response to all three MaMYBFAs, with the largest effect on MaMYBFA1 and MaMYBFA2;
194 the activity of *proMaFLS2–1500* was mildly, but not significantly, increased in response to
195 MaMYBFA1–3 (Fig. 3c). These results suggest that all three MaMYBFAs repress *MaFLS1*
196 expression and that MaMYBFA1 slightly activates *MaFLS2* expression.

**197 MaHSF11 promotes flavonol synthesis in banana by transactivating *MaFLS1* and
regulating several other biosynthesis genes**

None of the banana SG7 R2R3-MYBs we examined upregulated the expression of *MaFLS1*
but instead repressed it. Because *MaFLS1* is highly expressed in leaves and is the major enzyme
for flavonol synthesis in banana (Pandey *et al.,* 2016; Busche *et al.,* 2021), we attempted to
identify a putative regulator of *MaFLS1*. An ~2-kb promoter fragment was searched for
possible TF binding sites, including ACGT-motif, E-box, LTRE (DRE/CRT), HSE, and W-
box elements, which are recognized by bZIP, MYC, DREB/CBF, HSF, and WRKY factors,
respectively (Supplemental File S1). Among these, the HSF TFs are poorly characterized in
the context of flavonoids. The *MaFLS1* promoter contained two HSEs (nTTCnnGAAn) at
positions −310 (HSE1) and −1154 (HSE2) upstream of the transcription start site (TSS). To
identify a HSF as a possible HSE-binding candidate, we searched the publicly available banana
transcriptome datasets related to abiotic stresses because both *HSF* and *FLS* are well-known
stress-responsive genes. We found a transcriptome dataset of the drought-stressed banana
cultivars Saba (ABB, drought tolerant) and Grand Naine (AAA, drought susceptible) to be the
most suitable for this analysis (Muthusamy *et al.*, 2016). From the transcriptome deep
sequencing data, we found five *HSF* genes, *MaHSF9* (*GSMUA_Achr8G27750_001*),
MaHSF11 (*GSMUA_Achr6G31750_001*), *MaHSF16* (*GSMUA_Achr3G15640_001*),
MaHSF25 (*GSMUA_Achr1G07880_001*), and *MaHSF26* (*GSMUA_Achr1G16240_001*), that
were differentially expressed before and after drought treatment in the cultivar Grand Naine
(Fig. 4a). *MaHSF25* and *MaHSF26* are class B HSFs, which are known to repress gene
expression (Scharf *et al*., 2012). However, *MaHSF9*, *MaHSF10*, and *MaHSF11* are class A
HSFs, which are known to act activate gene expression due to the AHA activator motif (Scharf
et al., 2012).

Using a dual-luciferase assay in *N. benthamiana* leaves, we examined whether
MaHSF9, MaHSF10, and MaHSF11 could regulate the expression of *MaFLS1* by testing their
ability to regulate *proMaFLS1–2556*. We observed a significant increase in the activity of
proMaFLS1–2556 when co-transfected with either *MaHSF10* or *MaHSF11* compared to the

EV (Fig. 4b). Between these two, MaHSF11 showed the higher *in vivo* transactivation ability (>3.6-fold, $P < 0.0001$). Therefore, we focused on MaHSF11 in further studies and tested whether MaHSF11 could directly interact with *proMaFLS1*. To verify direct binding of MaHSF11 to the HSEs in the *MaFLS1* promoter, we performed a ChIP-qPCR assay in which *proMaFLS1* was co-infiltrated with a *pro35S:YFP-MaHSF11* construct into *N. benthamiana* leaves and expressed for 3 days. An anti-GFP antibody was used to immunoprecipitate YFP-MaHSF11 and the associated chromatin. A qPCR assay with purified DNA from the precipitated chromatin showed an approximate 5-fold enrichment of fragment F1 containing HSE1; fragment F2, containing HSE2, was not enriched (Fig. 4c). We then performed Y1H assays to validate the direct binding of MaHSF11 to HSEs of *proMaFLS1* using 100-bp, HSE-containing promoter fragments F1 and F2 to test for interaction with the MaHSF11-AD fusion protein. The Y1H growth assay showed strong binding of MaHSF11 to F1 and a weak interaction with F2 (Fig. 4d). Taken together, our results show that MaHSF11 binds directly to the HSEs in the promoter of *MaFLS1* and activates its expression. Phylogenetic analysis revealed clustering of MaHSF11 with AtHSFA4a, which confers salt tolerance in the MPK3/6-dependent pathway in Arabidopsis, and other A-class HSF proteins (Supplemental Fig. S3). Furthermore, a subcellular localization assay in *N. benthamiana* leaves using a *pro35S:YFP-MaHSF11* construct confirmed co-localization with a nuclear NLS-RFP marker (Fig. 5a). In addition to nuclear localization, we also observed fluorescence in plasma membrane, suggesting probable post-translational modification.

To further test the regulatory role of MaHSF11, we transiently overexpressed *proZmUBI:MaHSF11* in banana fruit discs (Fig. 5b). RT-qPCR analysis confirmed that the expression of *MaHSF11* was significantly higher (8-fold, $P \leq 0.001$) in *proZmUBI:MaHSF11* OE banana fruits than in the EV controls (Fig. 5c). Furthermore, the expression of *MaFLS1* was significantly upregulated (3-fold, $P \leq 0.01$), and the expression of *MaFLS2* was downregulated about 2-fold (Fig. 5c). We also observed differential expression of other structural flavonol biosynthesis genes in *MaHSF11*-OE banana fruit discs (Supplemental Fig. S4-d) as well as ~3-fold upregulation in MaMYBFA3 expression, but no significant upregulation of the other two MYB genes (Supplemental Fig. S4e).

We quantified the levels of several different flavonol compounds in *MaHSF11*-OE fruit tissues. LC-MS-based quantification showed increased levels of kaempferol, quercetin, and myricetin in transgenic fruit discs (Fig. 5d). Taken together, these results suggest that

257 MaHSF11 positively regulates flavonol synthesis via regulation of several key biosynthesis
258 genes, including direct regulation of *MaFLS1*.

259 MaHSF11 directly regulates *MaF3'5'H1* to promote flavonoid B-ring hydroxylation and
260 enhances myricetin biosynthesis

MaHSF11 overexpression modulated the expression of *MaF3'H* (Supplemental Fig. S4d) and
all the *MaF3'5'H1–7* gene family members (Fig. 6a). *MaF3'5'H1* was the most upregulated
gene (a 7-fold increase), followed by *MaF3'5'H2* and other homologs; *MaF3'H* expression
decreased negligibly (Supplemental Fig. S4d). Analysis of *cis* motifs in the promoter of
MaF3'5'H1 revealed a single HSE (nGAAnnTTCn) at the –720-bp position relative to the TSS
(Supplemental File S1). We performed Y1H assays to test the interaction of MaHSF11 with
the HSE of *proMaF3'5'H1–1335* using a 100-bp, HSE-containing promoter fragment (F1),
which showed a strong interaction with MaHSF11, as indicated by yeast growth (Fig. 6b). The
ability of MaHSF11 to transactivate *proMaF3'5'H1–1335* was tested in a dual-luciferase assay
in *N. benthamiana* leaves. The increased LUC activity with MaHSF11, compared to the EV
control, indicated that MaHSF11 activated the *MaF3'5'H1* promoter (Fig. 6c). To further test
the direct binding of MaHSF11 via the HSE, we co-infiltrated a *pro35S:YFP-MaHSF11*
construct and the *proMaF3'5'H1–1335* fragment into *N. benthamiana* leaves and performed a
ChIP assay using an anti-GFP antibody. This ChIP-qPCR assay revealed enrichment of the
HSE-containing 100-bp promoter fragment F1 (Fig. 6d). Overall, these results show that
MaHSF11 directly regulates the expression of *MaF3'5'H1*.

277 MaHSF11 regulates *MaMYBFA3* through MaDREB1

Overexpression of *MaHSF11* in banana fruit discs resulted in higher expression of several other
structural genes involved in flavonol biosynthesis (Supplemental Fig. S4a-d). Therefore, we
investigated the expression of regulatory *MaMYBFA1–3* genes. We found a significant
upregulation of only *MaMYBFA3* in banana fruit discs overexpressing *MaHSF11*
(Supplemental Fig. S4e). A promoter analysis of the ~2-kb region upstream of the *MaMYBFA3*
TSS revealed the absence of any HSEs (Supplemental File S1). Therefore, we hypothesized
that MaHSF11 might indirectly regulate the expression of *MaMYBFA3* via an unknown factor.
Furthermore, promoter analysis of *MaMYBFA3* revealed the presence of several DRE/CRT
(CCGAC) elements in the ~2.5-kb region upstream of the TSS (Supplemental File S1). Because
HSFs bind HSEs in the promoter of *DREB* genes (Yoshida *et al.,* 2011; Huang *et al.,* 2016),
we hypothesized that DREB TFs may act as mediators between *MaHSF11* and *MaMYBFA3*.

To investigate the likelihood that MaHSF11–MaDREB–MaMYBFA3 is a regulatory module in banana, we measured the expression levels of *DREB* genes using the RNA-Seq data from drought-treated bananas of the Grand Naine cultivar (Muthusamy *et al.*, 2016). We found the expression of a single *DREB* gene (*GSMUA_Achr4G19660_001*), which was also upregulated after drought stress (Fig. 7a). This gene was previously characterized as MaCBF1 (C-repeat binding factor1) and described as a critical regulator of cold stress tolerance in banana (Zhao *et al.*, 2013; Shan *et al.*, 2014). We isolated the 777-bp full-length CDS of this gene and named it *MaDREB1* due to nucleotide sequence dissimilarity with MaCBF1 (Shan *et al.*, 2014). Phylogenetic analysis revealed that MaDREB1 formed a clade with rice OsDREB3 and maize ZmDREB (Supplemental Fig. S5). To examine whether the promoter of *MaDREB1* was regulated by MaHSF11, we analyzed the region ~2 kb upstream of the TSS (*proMaDREB1–998*) and identified two HSEs at positions –223 and –576 bp upstream of the TSS (Supplemental File S1). We measured the expression of *MaDREB1* in *MaHSF11* OE fruit discs and determined that it was significantly upregulated (>6-fold) (Fig. 7b).

To verify the direct regulation of MaDREB1, we conducted a Y1H assay. We cloned two 100-bp fragments (F1 and F2) harboring HSEs (HSE1 and HSE2, respectively) of the *proMaDREB1* and conducted a Y1H assay with MaHSF11 as the effector. Promoter fragments carrying the HSEs were cloned into the pAbAi vector and transformed into yeast (*Saccharomyces cerevisiae*) Y1H Gold strain together with pGADT7 as a negative control. MaHSF11 in pGADT7 was transformed with both fragments, and the growth assay was performed on media supplemented with aureobasidin A (AbA). MaHSF11 could bind the HSE1-containing fragment F1 of *proMaDREB1 in vitro* (Fig. 7c). To test the transactivation activity of MaHSF11 on *proMaDREB1*, we cloned a 998-bp promoter fragment of *MaDREB1* into the p635nRRF vector and performed dual-luciferase assays. The EV as a negative control and *MaHSF11* as the effector were individually infiltrated together with the reporter construct carrying a particular promoter fragment into *N. benthamiana* leaves, and the assay was performed 3 days post infiltration. LUC activity increased significantly when MaHSF11 was co-transfected compared to the EV control (Fig. 7d). Furthermore, *pro35S:YFP-MaHSF11* and the promoter fragment were also transformed into *N. benthamiana* leaves, and a ChIP assay was conducted. An anti-GFP antibody was used to precipitate MaHSF11-bound chromatin, and anti-IgG was used as a negative control. ChIP-qPCR showed enrichment of the F1 fragment in the immunoprecipitated chromatin, demonstrating that MaHSF11 directly bound the F1 fragment and regulated *MaDREB1* (Fig. 7e).

To test whether MaDREB1 acted as a direct transcriptional activator of *MaMYBFA3*, we isolated *proMaMYBFA3–1892*, which contains all three DREs (CCGAC), and cloned it into pAbAi. We performed a Y1H assay with MaDREB1 as the effector in pGADT7 and EV as a control. Promoter fragments cloned into the pAbAi vector were transformed into the Y1H strain, and a growth assay was performed on AbA-supplemented media. Growth was observed in the MaDREB1 transformed strain, suggesting that it bound to the promoter fragment (Fig. 8a). To study the transactivation activity of MaDREB1, we cloned a 2508-bp fragment of *proMaMYBFA3* into p635nRRF to perform a dual-luciferase assay with MaDREB1 in pBTDest as the effector and the EV as a negative control. LUC activity increased when MaDREB1 was used as the effector compared to the control, indicating its ability to transactivate *proMaMYBFA3* (Fig. 8b). Overall, these findings show direct regulation of *MaMYBFA3* by MaDREB1.

To verify the function of *MaDREB1 in planta*, we transiently overexpressed it in banana fruit discs. GUS staining indicated successful transformation of the GUS expression cassette (Fig. 8c). RT-qPCR analysis of the transformed fruit tissue showed high expression (~375-fold, $P < 0.0001$) of *MaDREB1* relative to the EV control, demonstrating successful overexpression (Fig. 8d). Expression analysis of *MaMYBFA3* in the *MaDREB1* OE tissue showed an ~10-fold increase in the expression level ($P < 0.05$). We also found higher expression (~21-fold, $P < 0.001$) of *MaMYBFA2*, while *MaMYBFA1* expression was unaltered (Fig. 8d). *MaFLS1* and *MaFLS2* also showed opposite trends in expression in the *proZmUBI:MaDREB1* OE fruit tissues: *MaFLS1* was downregulated, and *MaFLS2* was upregulated (Fig. 8d). Differential expression of other structural flavonol biosynthesis–related genes was also observed in *proZmUBI:MaDREB1* OE banana fruit tissues (Supplemental Fig. S6). To study the modulation of flavonol content resulting from *MaDREB1* overexpression, we used LC-MS to quantify the levels of different metabolites and found significantly higher accumulation of kaempferol, quercetin, and myricetin (Fig. 8e). These results indicate that MaDREB1 positively regulates flavonol biosynthesis in banana.

Discussion

Flavonoid biosynthesis pathways developed during the evolution of land plants. As they became terrestrial, plants developed response mechanisms to adapt to and cope with several new environmental conditions. Flavonoid biosynthesis is highly responsive to drought stress hormones like ABA (Loreti *et al.,* 2008; Medina-Puche *et al.,* 2014; Song *et al.,* 2022). The

role of ABA in the context of flavonoid biosynthesis has been studied recently, strengthening our knowledge regarding transcriptional regulation of flavonoid biosynthesis. Although flavonol biosynthesis, in particular, is positively modulated by ABA, a conserved response seen throughout the plant kingdom (Xu *et al.,* 2012; Song *et al.,* 2022), the molecular mechanism controlling this response has remained obscure. In this study, we identified a key transcriptional regulatory network involving different classes of TFs and showed their interaction in regulating SG7 *MaMYBFA* and flavonol biosynthesis genes in the herbaceous fruit crop banana. We deciphered important details of the molecular mechanisms involving the transcriptional regulatory module of how HSFs, DREB, and MYB proteins orchestrate flavonol accumulation.

Initially, three candidate SG7 R2R3-MYB TFs, *MaMYBFA1–3*, were identified as having high sequence conservation in their R2R3 domains and partial conservation in the SG7-1 (GRTxRSxMK, Stracke *et al.*, 2001) and SG7-2 motifs ([W/x] [L/x]LS, Czemmel *et al.,* 2009). Furthermore, the absence of the bHLH-interacting motif ([DE]Lx2[RK]x3Lx6Lx3R, Zimmermann *et al.,* 2004) in the R3 domain suggested that these three candidate genes were independent regulators of flavonol synthesis in banana. Expression analysis of the three *MaMYBFA* genes revealed the highest level of transcript accumulation in bract tissue, a result that correlated well with expression of *MaFLS2*, their probable target. It also showed a positive correlation with *MaFLS2* in other tissues. By contrast, expression of *MaFLS1* showed an inverse correlation regarding the three candidate genes in young leaves, where its transcript level was highest. This finding indicates the presence of a regulator other than SG7 MYB in banana, acting upstream of *MaFLS1*. All these results prompted us to functionally characterize the candidate *MaMYBFA* genes in banana to determine their possible roles as flavonol-specific TFs. To do this, we transiently overexpressed the *MaMYBFA* genes in banana fruit. Although *MaFLS2* expression was activated by MaMYBFA, the expression of *MaFLS1* was downregulated. This finding could be an exception, because none of the SG7 MYB activators downregulate flavonol synthase genes, the key genes in the flavonol biosynthesis pathway. Flavonol content mainly remained unaltered with MaMYBFA overexpression, a finding that can be attributed to the downregulation of *MaFLS1* and *MaF3'H* and increased expression of *MaFLS2*, which could compensate for it. Thus, although we concluded that MaMYBFAs are weak regulators of flavonol synthesis in banana, it is difficult to ascertain their positive regulatory role on flavonol synthesis. However, our results show the division of labor among the SG7 MYB transcriptional regulators within the regulatory cascade. Such a mechanism has

been observed in the dicot freesia (*Freesia hybrida*), in which four SG7 MYB proteins,

FhMYBF1–4, differentially regulate the expression of *FhCHS* and *FhCHI* paralogs, as well as

FhFLS1, but not *FhFLS2* (Shan *et al.,* 2020). SG7 MYB TFs regulate flavonol synthesis by

directly binding to MYB binding sites (MBSs, CNGTTR) present in the promoters of flavonol

biosynthesis genes, activating their expression. In apple, MdMYB22 binds to the MBS in the

promoters of *MdCHS* and *MdFLS* to regulate their expression (Wang *et al.,* 2017). In freesia,

FhMYB1–4 bind to the MYB core (CNGTTR) and MYBPLANT motifs (MACCWAMC),

which are essential for promoter binding and transactivation (Shan *et al.,* 2022). Overall, these

results suggest that MaMYBFA1–3 are the key functional flavonol-specific transcriptional

regulators in banana.

HSFs are critical components of the signaling cascades involved in responses to salt,

cold, drought, and heat stress (Chauhan *et al.,* 2011; Bian *et al.,* 2020; Wang *et al.,* 2020).

Several HSFs and their regulatory networks have been discovered in Arabidopsis, rice, tomato,

soybean (*Glycine max*), wheat, and grape (*Vitis vinifera*) (Guo *et al.,* 2016; Wu *et al.,* 2022).

In this study, MaHSF11 was identified as a regulator of *MaFLS1*. MdHSFA8a was previously

identified as a regulator of *MdFLS* in apple, promoting flavonol biosynthesis and the drought

stress response by regulating *MdFLS* expression (Wang *et al.,* 2020). GmHSFB2B confers salt

stress tolerance and flavonol synthesis in soybean by repressing GmNAC2, a negative regulator

of *GmFLS1* and *GmFLS2* (Bian *et al.,* 2020). Apart from these, other non-SG7 MYB regulators

from other species include AtMYB21, AtMYB24, and FhMYB21L2. The presence of three

HSEs in the promoter of *MaFLS1* suggested it might be regulated by a HSF. We determined

the role of MaHSF11 to be a regulator of CYP450 B-ring hydroxylase in banana, where it

directly bound to the promoter of *MaF3'5'H1* (Fig. 6b). Very few regulators of *F3'5'H* have

been identified in plants. In poplar (*Populus tremula×tremuloides*), MYB117 specifically

regulates *F3'5'H* expression to promote anthocyanin synthesis and flavonoid B-ring

hydroxylation (Ma *et al.,* 2021). MYBC1 and WRKY44 both independently and cooperatively

regulate *F3'5''H* expression to regulate the anthocyanin level in kiwifruit (*Actinidia purpurea*)

(Peng *et al.,* 2020).

In our study, *MaHSF11* overexpression upregulated *MaFLS1* but downregulated

MaFLS2, indicating the presence of a complex regulatory cascade leading to preference of one

paralog over the other. MaHSF11 also promoted the expression of several *MaF3'5'H* gene

homologs, while slightly repressing *MaF3'H* (Supplemental Fig. S4d). Since the enzymes

compete with each other for the common substrate dihydrokaempferol, MaHSF11 might be

specifically promoting the myricetin synthesis branch and tries to suppresses the quercetin synthesis branch catalyzed by MaF3′H, either directly or indirectly. *MaFLS1* was highly expressed in young leaves, and this finding can be correlated with higher *MaHSF11* expression in leaves compared to fruits and roots (Wei *et al.,* 2016). Fruit tissue, particularly peel, had higher expression of *MaF3'H*, leading to a greater accumulation of rutin and quercetin in peel compared to other tissues (Pandey *et al.,* 2016). Of the seven paralogs of *MaF3'5'H* in banana, five were upregulated 2-fold with *MaHSF11* overexpression, suggesting a role for *MaHSF11* as a master regulator of this gene family (Fig. 6a). These results further suggest that MaHSF11 promotes flavonol biosynthesis via a distinct regulatory module to regulate stress-responsive genes such as flavonol biosynthesis–related genes involved in drought, cold, and salinity stress responses in banana.

Preference of MaHSF11 for the *MaFLS1* and *MaF3'5'H* homologs (relative to *MaFLS2* and *MaF3'H*) suggested a specificity of MaFLS1 for dihydromyricetin as a substrate. In a recent study, MaFLS1 also showed specificities for dihydrokaempferol and dihydroquercetin. In future studies, it will be important to determine whether MaFLS1 has a substrate specificity for dihydromyricetin as well. CYP450 enzymes are involved in the flavonoid metabolon in both rice and soybean (Shih *et al.,* 2008; Waki *et al.,* 2016), where they act as ER membrane–bound nucleation sites for metabolon formation. Interaction among enzymes varies greatly in terms of their stability and complexity. Preferential upregulation of MaF3′5′H and MaFLS1 and downregulation of MaF3′H and MaFLS2 suggested probable metabolon formation in the ER. In snapdragon (*Antirrhinum majus*), F3′H associates with CHI to form a metabolon (Fujino *et al.,* 2018). In Arabidopsis, FLS1 interacts with CHS via competition with DFR (dihydroflavonol 4-reductase), which can also bind to CHS (Crosby *et al.,* 2011).

We also observed that MaHSF11 upregulated *MaMYBFA3* indirectly via MaDREB1. Interactions of DREB and HSF TFs have been reported previously (Yoshida *et al.,* 2011; Ya-Chen *et al.,* 2016). In Arabidopsis, DREB2A is transcriptionally regulated by HSFA1a, HSFA1b, and HSFA1d (Yoshida *et al.,* 2011), and DREB2A also acts upstream of HSFA3 to regulate its expression (Schramm *et al.,* 2008). In our study, we found ~2-fold increased expression of MaHSF11 when *MaDREB1* was overexpressed in banana, suggesting conservation of an interconnected regulatory cascade (Supplemental Fig. S7). We also determined that MaHSF11 directly bound to the HSE in *proMaDREB1* to regulate its expression. Although overexpression of *MaDREB1* upregulated several biosynthesis genes related to flavonol biosynthesis, leading to increased flavonol accumulation, *MaCHS2* and

MaFLS1 were downregulated (Supplemental Fig. S6a–e). In future work, it will be intriguing to determine whether MaDREB1 directly represses these genes. Recently, *MaCBF1* was reported to be a positive regulator of chlorophyll catabolism: It possesses strong transactivation activity and is localized to the nucleus (Xiao *et al.*, 2023). Prior to this study, several DREB proteins were reported to regulate different branches of flavonoid biosynthesis. CitERF32 and CitERF33 act as transcriptional inducers of *CitCHIL1* to promote flavone and flavanone biosynthesis in citrus (*Citrus reticulata cv. Suavissima*) (Zhao *et al.,* 2021). In eggplant (*Solanum melongena*), SmCBF1, SmCBF2, and SmCBF3 interact with SmMYB113 to regulate *SmCHS* and *SmDFR* expression and enhance anthocyanin synthesis (Zhou *et al.,* 2020). In our study, we discovered MaDREB1 to be a key regulator of flavonol synthesis, modulating the expression of regulatory SG7 MYB genes. In eggplant as well, SmCBF1, SmCBF2, and SmCBF1 are induced by drought, salt, and ABA, showing that MaDREB1 might also be involved in drought stress signaling as a transcriptional regulator of stress-responsive genes such as flavonoid biosynthesis genes. Our study provides insights into a key regulatory cascade that promotes flavonol biosynthesis in banana under different stress conditions, as shown in our proposed model (Fig. 9).

The banana cultivar Grand Naine (Cavendish, AAA) is highly susceptible to different abiotic stresses as well as pathogen infection. Flavonols acting as ROS scavengers and pathogen defense molecules are synthesized in increased amounts under a variety of stresses. Here, we identified the HSF–DREB-MYB transcriptional regulatory module and characterized how it works independently or cooperatively to regulate flavonol synthesis in an ABA-responsive manner. The TFs in this regulatory module can be used for effective engineering of stress resistance in crop plants like banana. These candidate genes could be utilized for metabolic engineering of rutin and myricetin synthesis in banana fruit.

Materials and methods

Plant material

Banana (cultivar subgroup Cavendish [AAA] variety 'Grand Naine') plants were grown in the fields of National Institute of Plant Genome Research (NIPGR), New Delhi, India. Various tissues, including young leaf, bract, pseudostem, root, peel, and pulp, of different fruit developmental stages were sampled during winter season, frozen in liquid nitrogen, and stored at −80°C until further use. *Nicotiana benthamiana* plants were grown in a Percival AR-41L3

484   growth cabinet at 100 µmol m^{-2} s^{-1} light intensity and 22°C under a 16-h-light/8-h-dark

485   photoperiod.

Identification of candidate flavonol-specific TFs from banana

To identify candidate SG7 R2R3-MYB TFs, which are known activators of flavonol

biosynthesis, we retrieved sequences of previously characterized flavonol-specific TFs from

various plants from NCBI (https://www.ncbi.nlm.nih.gov). These sequences were used to

perform a BLASTp search against the banana genome sequence (DH-Pahang version 2)

(https://banana-genome-hub.southgreen.fr). For phylogenetic studies, MEGA X (Kumar *et al.,*

2018) was used to construct a maximum-likelihood phylogenetic tree with 1000 bootstrapping

rounds. For multiple sequence alignment, MaMYBFA1–3 peptide sequences were aligned with

a set of similar MYBs (Stracke *et al.,* 2014) using MAFFT v.7.299b (Katoh and Standley,

2013). Different domains and motifs were identified manually (Naik *et al.,* 2021).

Cloning of flavonol-specific transcriptional regulators

The full-length CDSs (without the stop codons) of the MaMYB-encoding genes *MaMYBFA1*

(*Ma02_g00290*), *MaMYBFA2* (*Ma05_g23640*), and *MaMYBFA3* (*Ma08_g10260*) were

amplified using first-strand cDNA of banana fruit tissue and a set of primers containing

Gateway attB sites that were designed based on information from the Banana Genome Hub

(Supplemental Table S1). The resulting amplicons were recombined into the Gateway vector

pDONR^{TM}zeo (Invitrogen) and transformed into *Escherichia coli* TOP10 (Invitrogen) cells.

Sanger sequencing of the resulting entry plasmids was performed by the NIPGR sequencing

core facility (New Delhi, India).

Subcellular localization and co-localization assays

CDSs of *MaMYBFA1–3* and *MaHSF11* were cloned into the pSITE-3CA binary vector

(Chakrabarty *et al*., 2007) via LR recombination, producing yellow fluorescent protein (YFP)

fusion proteins. Empty vectors were used as negative controls. The binary plasmids were

transformed into *Agrobacterium tumefaciens* strain GV3101-pMP90 (Koncz & Schell, 1986).

Equal volumes of *A. tumefaciens* cultures having the desired gene and the marker were mixed

for agro-infiltration. *N. benthamiana* leaves were infiltrated on the abaxial side using a syringe,

according to the protocol described by Walter *et al.* (2004). Infiltrated plants were kept in the

dark at 25°C for 2 days prior to analysis. Leaf discs were then cut from the infiltrated sites and

observed under an argon laser confocal laser-scanning microscope (TCS SP5; Leica

Microsystems, Wetzlar, Germany) with YFP and RFP filters. YFP and RFP fluorescence was observed at 514-nm excitation/527-nm emission and 558-nm excitation/583-nm emission wavelengths, respectively. YFP/RFP gains (intensities %) for MaMYBFA1, MaMYBFA2, MaMYBFA3, and MaHSF11 were 760 (39)/158 (39), 818 (39)/158 (39), 818 (30)/158 (39), and 718 (42)/154 (39), respectively.

RT-qPCR analysis

Total RNA was extracted from different banana organs according to a previously described protocol (Asif *et al.,* 2006) and subsequently treated with RNase-free DNase (TURBO DNA-freeTM Kit, Invitrogen). Total RNA was reverse transcribed to generate first-strand cDNA using a RevertAid H Minus First Strand cDNA Synthesis Kit (Thermo Fisher) and oligo(dT) primers. RT-qPCR-based gene expression analyses were performed on a 7500 Fast Real-time PCR System (Applied Biosystems) using the 2x SYBR Green PCR Master mix (Applied Biosystems) and diluted cDNA corresponding to 10 ng total RNA in a final volume of 10 µL. *MaACTIN1* (GenBank AF246288) was used to normalize transcript abundance. Relative transcript levels were calculated using the cycle threshold (C_t) $2^{-\Delta\Delta CT}$ method (Livak & Schmittgen, 2001) and are presented as fold changes relative either to the tissue having the lowest expression or to the control condition, depending on the requirement. Three biological replicates were analyzed in each RT-qPCR analysis.

Metabolite analysis

Targeted metabolite analysis was performed to quantify flavonols according to a protocol described previously (Pandey *et al.,* 2016; Naik *et al.,* 2021; Rajput *et al.,* 2022a). Plant metabolites were extracted from shredded plant material in 80% methanol at room temperature. Finely ground tissues were dissolved in 80% methanol, kept at 70℃ for 15 minutes, and subjected to centrifugation for 10 minutes at 12,000 rpm (18514 g). After centrifugation, supernatants were mixed with three volumes of 1 N HCl for acid hydrolysis and incubated at 94℃ for 2 h. After incubation, an equal volume of ethyl acetate was added, and the upper phase was removed for further purification of the desired metabolites. The ethyl acetate was evaporated at 37℃ under lower pressure (90–100 mbar) on a Buchi rotavapor R-300 (BUCHI, Switzerland), and samples were finally dissolved in 80% methanol. Liquid chromatography–mass spectrometry (LC-MS) analysis was performed using an ultra-high-performance liquid chromatography (UHPLC) system (Exion LC Sciex, Framingham) coupled to a triple quadrupole system (QTRAP6500+; ABSciex) with electrospray ionization. The voltage was

547   set to 5500 V for positive ionization. Detailed UHPLC and LC-MS parameters have been

548   described by Naik *et al.* (2021). Identification and quantitative analyses were done via Analyst

549   software (version 1.5.2).

Cloning of reporter constructs and identification of putative *cis* elements in promoters

Total genomic DNA was extracted from banana leaves using the DNeasy Plant Mini Kit

(Qiagen). Sequences of the putative target gene promoter fragments were retrieved from the

banana genome (https://banana-genome-hub.southgreen.fr) (D'Hont *et al.,* 2016) and analyzed

for the presence of *cis*-regulatory motifs via New PLACE software

(https://www.dna.affrc.go.jp) supplemented by manual curation. Promoter fragments were

amplified from banana genomic DNA using designed primer sequences and Phusion™ High-

Fidelity DNA Polymerase (Thermo Scientific). The amplified fragments were cloned into the

pENTR™/D-TOPO™ vector (ThermoFisher) and sequenced to confirm their integrity. The

entry clones of promoters (*proMaFLS1–2508*, *proMaFLS2–1500*, *proMaF3'5'H1–1335*,

proMaDREB1–998, and *proMaMYBFA3–2508*) were recombined into the vector p635nRRF

containing *pro35S:REN* (Kumar *et al.,* 2018) to make the reporter constructs.

Cloning of cDNA from MaHSF and MaDREB transcriptional regulators

Full-length CDSs of selected homologs of MaHSF and MaDREB were amplified by first-

strand cDNA synthesis of banana leaves using Phusion™ High-Fidelity DNA Polymerase

(Thermo Scientific) and a set of primers containing Gateway attB sites designed on the basis

of information from the Banana Genome Hub (Supplemental Table S1). The resulting

amplicons were cloned into the pDONR™ zeo vector (Invitrogen) and transformed into *E. coli*

TOP10 cells, and integrity was confirmed by Sanger sequencing (Sanger *et al.*, 1977).

Dual-luciferase reporter assay

To test the transactivation potential of the different transcriptional regulators, MaMYBFA1-

MaMYBFA3, MaHSF11, and MaDREB1 CDSs were recombined into the destination vector

pBTdest (Baudry *et al.,* 2004) to generate CaMV *pro35S*-driven effector constructs. For dual-

luciferase reporter assays, reporter and effector constructs were transformed into *A.*

tumefaciens strain GV3101 (pMP90) (GoldBio) and subsequently infiltrated using a syringe

into the abaxial side of *N. benthamiana* leaves according to a previously described protocol

(Walter *et al.,* 2004). LUC and REN activities were measured 48 h later in extracts from

infiltrated leaf discs using the Dual-Luciferase Reporter Assay System (Promega) according to

the manufacturer's protocol. Luminescence in the samples was quantified using the

POLARstar Omega multimode plate reader (BMG Labtech), and LUC activity was normalized

to REN activity. Values are given as means ± SD of the LUC/REN ratio.

Y1H assays

The Matchmaker Gold Yeast 1-hybrid system (Clontech) with the yeast one-hybrid (Y1H) gold

yeast strain was used. Promoter fragments were cloned into the pABAi vector (Takara Bio),

and the pABAi plasmids carrying the promoter fragments were linearized and transformed into

the Y1H Gold strain according to the Matchmaker protocol, whereby they integrated into the

yeast genome by homologous recombination. Effector plasmids and empty pGADT7-GW

vector (Gal4-AD fusion, Ura selection) controls were transformed into the Y1H strains

previously transformed with AurR-reporter constructs (*proMaFLS1*-AurR, *proMaF3'5'H1*-

AurR, *proMaDREB1*-AurR, and *proMaMYBFA3*-AurR). To test for interaction, the yeast

colonies containing effector and reporter constructs were spotted onto medium containing AbA

per the manufacturer's protocol. Growth of different Ura$^-$ yeast colonies was checked using

different concentrations of AbA to determine the minimum inhibitory concentration (MIC)

values and to check for possible interactions. The MIC values for *proMaFLS1*, *proMaF3'5'H1*,

proMaDREB1, and *proMaMYBFA3* were 100 ng mL^{-1} AbA, 1000 ng mL^{-1} AbA, 100 ng mL^{-1},

and 900 ng mL^{-1}, respectively.

ChIP-qPCR assay

The ChIP-qPCR assay was performed using the ChIP Kit - Plants kit (Abcam) according to the

manufacturer's instructions (Shi *et al.,* 2019). Promoter fragments containing the HSE *cis* motif

in p635nRRF that was used for dual-luciferase assays were co-infiltrated with YFP-MaHSF11

fusion proteins into *N. benthamiana* leaves. Three days after infiltration, equally sized leaf

discs were used for ChIP. Next, 2 µL anti-GFP antibody (Catalog #ab290, Abcam) was used

for chromatin precipitation in two biological replicates, and 2 µL of anti-IgG antibody (Catalog

#ab48386, Abcam) was used as a negative control. qPCR was performed with 5 µL 2x SYBR

Green PCR Master mix (Applied Biosystems), 1 µL ChIP DNA, and 0.5 µL each forward and

reverse primer, with 40 cycles of amplification as described above. The relative fold

enrichment of the target promoter fragments was calculated using the $2^{-(-\Delta CT)}$ method (Livak

& Schmittgen, 2001). The primers used in ChIP-qPCR are listed in Supplemental Table S1.

Transient overexpression in banana fruit

For transient overexpression studies, the CDSs of *MaMYBFA1*, *MaMYBFA2*, *MaMYBFA3*, *MaHSF11*, and *MaDREB1* were cloned into pANIC6b (Mann *et. al.*, 2012), a Gateway-compatible destination vector suitable for overexpression of genes of interest using *A. tumefaciens*-mediated transformation, particularly useful for monocot species. pANIC6b encodes hygromycin resistance under the control of a constitutive rice (*Oryza sativa*) *actin1* promoter (*proOsACT1*) (LOC4338914), a constitutive *Panicum virgatum ubiquitin1+3* promoter (*proPvUBI*)-driven GUSPlus visual marker, and a Gateway cassette driven by a *Zea mays ubiquitin1* promoter and intron for overexpression of target genes. The resulting overexpression plasmid and the empty vector control were transformed into *A. tumefaciens* strain EHA105 (Mehra *et al.*, 2019). Resulting *A. tumefaciens* suspensions were then separately introduced into slices of immature green banana fruits by vacuum infiltration as previously described (Rajput *et al.,* 2022a). After 3 days of co-cultivation on MS media (MS basal salt mixture, HiMedia), transformation of banana fruit slices was confirmed by GUS staining. Tissues were harvested and frozen in liquid nitrogen for subsequent expression (RT-qPCR) and metabolite (LC-MS) analyses.

Statistical tests

One-way and two-way analysis of variance (ANOVA) with Tukey's multiple comparison were performed for statistical analysis. Unpaired Student's *t*-tests were used for the specific comparisons. Asterisks indicate the level of significance: $*P \leq 0.05$; $**P \leq 0.01$; $***P \leq 0.001$, $****P \leq 0.0001$; ns, not significant.

Data availability

All data supporting the findings of this study are available within the paper and within the supplemental data published online.

Supplemental data

Figure S1. Tissue-specific expression of *MaMYBFA1-FA3* genes*.*

Figure S2. Relative expression of structural flavonol biosynthesis gene transcripts in EV and *MaMYBFA1-FA3* transiently overexpressing lines.

Figure S3. Phylogenetic analysis of MaHSF11 with previously characterized ABF gens from
other plant species.

Figure S4. Relative expression of structural and regulatory flavonol biosynthesis genes in
MaHSF11-overexpressing (OE) banana fruit discs.

Figure S5. Phylogenetic analysis of MaDREB1 with previously characterized ABF genes from
other plant species.

Figure S6. Relative expression of structural flavonol biosynthesis genes in *MaDREB1*-
overexpressing banana fruit tissues.

Figure S7. Relative expression analysis of *MaHSF11* in *MaDREB1*-overexpressing banana
fruits.

Acknowledgements

This work was supported by a core grant from the National Institute of Plant Genome Research
and a Department of Science and Technology-SERB for Core research grant
(CRG/2022/001178) to AP. JN and RR acknowledge the Council of Scientific and Industrial
Research, Government of India, for Senior Research Fellowships. The authors thank the DBT-
eLibrary Consortium (DeLCON) for providing access to e-resources. We also acknowledge
the Metabolome facility at NIPGR for phytochemical analysis.

Declarations

The authors declare no conflicts of interest.

Figure legends
Figure 1. Identification of candidate R2R3-MYB flavonol regulators from the banana genome
and their subcellular localization. (a) Various R2R3-MYB proteins were classified into three
major groups as flavonol-, anthocyanin-, or proanthocyanidin biosynthesis–specific regulators
using landmark flavonoid-specific R2R3-MYBs from different plant species. Numbers on each
branch show branch length, and solid dark blue circles represent bootstrap values. (b) Amino
acid sequence alignment of candidate SG7 R2R3-MYB transcription factors (MaMYBFA1,

MaMYBFA2, and MaMYBFA3, where FA stands for flavonol activator) and selected flavonol regulators from other plant species. The R2 and R3 domains are highlighted in grey. The SG7-1 and SG7-2 motifs, found in subgroup 7 R2R3-MYBs, are highlighted in dark green and yellow, respectively. Accession numbers and two-letter species abbreviations are provided in Supplemental File S2. (c) Subcellular localization of MaMYBFAs. MaMYBFA-YFP fusion proteins were expressed together with the nuclear marker NLS-RFP in *N. benthamiana* leaf epidermal cells. Confocal imaging shows co-localization of YFP and RFP fluorescence, indicating nuclear localization of the three MaMYBFA proteins. Representative images are shown with scale bar of 50 μm.

Figure 2. MaMYBFA1, MaMYBFA2, and MaMYBFA3 differentially regulate the expression of *MaFLS1* and *MaFLS2* and modulate flavonol biosynthesis in banana. (a) Diagram of the plant expression cassette used for transient overexpression (OE) of *MaMYBFA* in banana fruit discs. The *Zea mays ubiquitin1* promoter (*proZmUBI1*) drives the expression of *MaMYBFA*. The expression of the *GUSPlus* marker gene is driven by the *Phaseolus vulgaris ubiquitin1* promoter (*proPvUBI*). (b) Representative images showing GUS staining of transiently transformed banana fruit discs indicating successful transformation. Images were taken 3 days after transformation. (c-d) Relative expression of *MaMYBFA1–3* (c) and *MaFLS1* and *MaFLS2* (d) OE lines in transiently transformed *MaMYBFA* OE banana fruit tissues as determined by RT-qPCR. EV, empty vector control. (e) Flavonol quantification showing differential alteration in the accumulation of kaempferol, quercetin, and myricetin aglycones in the *MaMYBFA* OE banana fruit discs. Values are given in μg per 100 g of dry weight (DW) of fruit discs of three replicates. One-way ANOVA with Tukey's multiple comparison was used for statistical analyses. *$P \leq 0.05$; **$P \leq 0.01$; ***$P \leq 0.001$; ****$P \leq 0.0001$; ns, not significant.

Figure 3. MaMYBFAs regulate the expression of *MaFLS1* and *MaFLS2 in vivo*. (a) Schematic representation of the 2556- and 1500-bp promoter fragments of *MaFLS1* and *MaFLS2*. The locations of the different MYB core (CNGTTR) sequences and their positions upstream of the transcription start sites (−1) are indicated by red ovals. A plus (+) sign with a number indicates the length of the 5′ UTR. (b) Diagram of the pBTdest effector and the p635nRRF reporter constructs. Constitutive cauliflower mosaic virus (CaMV) promoter drives the expression of transcription factor encoding genes and renilla *luciferase* (REN). Desired amplified promoter fragments were placed upstream of firefly *luciferase* (LUC) (c) Dual-luciferase assay showing the altered activity of the promoters *proMaFLS1–2508* and *proMaFLS2–1500* when transiently

co-infiltrated into *N. benthamiana* leaves with the effectors MaMYBFA1, MaMYBFA2, and

MaMYBFA3. Promoter activity is represented as the LUC/REN ratio in four biological

replicates ($n = 4$).

Figure 4. MaHSF11 directly binds to the *MaFLS1* promoter and regulates the expression of

MaFLS1 to modulate flavonol biosynthesis in banana. (a) Heatmap showing expression of five

banana candidate *HSF* genes before and after drought treatment (Grand Naine cultivar, AAA

genome). FPKM values for the indicated genes were extracted from publicly available RNA-

Seq datasets and are represented according to the 0–7 color scale shown. (b) Dual-luciferase

assay showing transactivation of *proMaFLS1–2556* by MaHSF. Values are shown as means ±

SD of the LUC/REN ratios ($n = 4$). (c) ChIP-qPCR assay analyzing the binding of MaHSF11

to fragments F1 and F2 of *proMaFLS1*, both of which contain the HSE consensus sequence

(nTTCnnGAAn). Fold enrichment values are given as means ± SD of two biological replicates.

AB, antibody. (d) Yeast one-hybrid assay showing interaction of MaHSF11 with the two HSE-

containing *proMaFLS1* fragments F1 and F2. The empty pGADT7 vector (EV) was used as a

negative control. Interaction was indicated by yeast growth on synthetic Ura– dropout media

supplemented with aureobasidin (AbA). AD, activation domain.

Figure 5. MaHSF11 promotes flavonol synthesis in banana fruit. (a) Subcellular localization

of MaHSF11 in *N. benthamiana* leaves. The YFP-MaHSF11 fusion protein localizes to both

the nucleus and the cytoplasm and coincides with the signal from the nuclear localization

signal-red fluorescent protein (NLS-RFP) fusion in *N. benthamiana* leaves as observed in

confocal microscopy. Scale bar, 20 μm. (b) GUS staining of banana fruit slices indicating

successful transformation with the empty vector (EV) and the *proZmUBI:MaHSF11* construct.

(c) Relative expression of *MaHSF11*, *MaFLS1*, and *MaFLS2* in transiently transformed banana

fruit slices overexpressing (OE) *proZmUBI:MaHSF11* as determined by RT-qPCR. The empty

vector was transformed as a control (expression level set to 1). Values shown are means ± SD

of three technical replicates. (d) Quantification of the flavonols kaempferol, quercetin, and

myricetin in transformed banana fruit slices. Values are represented as means ± SD of three

replicates. Unpaired student's *t*-tests were used for statistical analyses. *$P ≤ 0.05$; **$P ≤ 0.01$;

***$P ≤ 0.001$.

Figure 6. MaHSF11 promotes flavonoid B-ring hydroxylation by modulating the expression

of *MaF3'5'H1*. (a) Relative expression of *MaF3'5'H1* genes in transformed banana fruit slices

transiently overexpressing *proZmUBI:MaHSF11*, as determined by RT-qPCR (EV, empty

vector control). Values shown are means ± SD of three technical replicates. (b) Yeast one-hybrid assay showing the interaction of MaHSF11 with *proMaF3'5'H1–100* fragment F1 carrying HSE (GGAATATTCT). pGADT7 was used as a negative control. Growth was checked on synthetic dropout medium supplemented with aureobasidin. (c) Dual-luciferase assay showing transactivation of *proMaF3'5'H1–1335* by MaHSF11. The promoter sequence driving the expression of luciferase (LUC) was co-infiltrated with the EV or *MaHSF11* constructs in *N. benthamiana* leaves. Values are shown as means ± SD of the LUC/REN ratio ($n = 4$). (d) ChIP-qPCR assay showing binding of MaHSF11 to fragment F1 of *proMaF3'5'H1* carrying an HSE (nGAAnnTTCn). The promoter fragment carrying HSE was co-infiltrated into *N. benthamiana* leaves with the *pro35S:YFP-MaHSF11* construct. Fold enrichment values are represented as means ± SD of two biological replicates. An anti-IgG antibody was used as a negative control, and an anti-GFP antibody was used to precipitate MaHSF11-enriched chromatin. Unpaired Student's *t*-tests were used for statistical analyses. *$P \leq 0.05$.

Figure 7. MaHSF11 acts upstream of *MaDREB1* to positively regulate its expression in banana. (a) Heatmap showing expression of candidate DREB genes before and after drought treatment in banana (Grand Naine cultivar, AAA genome). FPKM values for the indicated genes were extracted from publicly available RNA-Seq datasets and plotted according to the 0–7 color scale shown. (b) Relative expression of *MaDREB1* in banana fruit tissues transiently transformed with EV or *proZmUBI:MaHSF11* as determined by RT-qPCR. (c) Yeast one-hybrid assay showing interaction of MaHSF11 with the F1 and F2 fragments of *proMaDREB1–100* carrying an HSE (nGAAnnTTCn). pGADT7 was used as a negative control. Growth was checked on synthetic dropout medium supplemented with aureobasidin. (d) Dual-luciferase assay showing transactivation of *proMaDREB1–998* by MaHSF11. Promoter sequence driving the expression of LUC was co-infiltrated with empty vector and *MaHSF11* construct in *N. benthamiana* leaves. Values shown are means ± SD of LUC/REN ratio ($n = 4$). (e) ChIP-qPCR assay showing binding of MaHSF11 to *proMaDREB1*. A promoter fragment carrying an HSE was co-infiltrated into *N. benthamiana* leaf cells with the *pro35S:MaHSF11-YFP* construct. Fold enrichment values are represented as means ± SD of two biological replicates. An anti-IgG antibody was used as a negative control, and an anti-GFP antibody was used to precipitate MaHSF11 enriched chromatin. Unpaired Student's t-tests were used for statistical analyses. *$P \leq 0.05$; ***$P \leq 0.001$; ns, not significant.

Figure 8. MaDREB1 directly binds to the promoter of *MaMYBFA3* to regulate flavonol biosynthesis. (a) Yeast one-hybrid assay showing interaction of MaDREB1 with the

proMaMYBFA3–1892 fragment carrying three DREs (CCGAC). pGADT7 was used as a negative control. Growth was checked on synthetic dropout medium supplemented with aureobasidin. (b) Dual-luciferase assay showing transactivation of *proMaMYBFA3–2508* by MaDREB1. The *proMaMYBFA3–2508* sequence driving the expression of LUC was co-infiltrated with the empty vector or *MaDREB1* construct into *N. benthamiana* leaves. Values are shown as means ± SD of the LUC/REN ratio ($n = 4$). (c) GUS staining of transformed fruit. (d) RT-qPCR-based relative expression of *MaDREB1*, *MaMYBFAs*, *MaFLS1*, and *MaFLS2* in transformed banana fruit tissues transiently expressing EV and *proZmUBI:MaDREB1*. EV was transformed as a control. Values shown are means ± SD of three technical replicates. (e) Quantification of the flavonols kaempferol, quercetin, and myricetin in transformed fruit tissues. Values are represented as means ± SD of three replicates. Unpaired Student's *t*-tests were used for statistical analyses. *$P \leq 0.05$; **$P \leq 0.01$; ***$P \leq 0.001$; ****$P \leq 0.0001$.

Figure 9. A working model showing the modulation of hydroxylated flavonol synthesis by the MaHSF11–MaDREB1–MaMYBFA regulatory module. The banana SG7 R2R3 MYB, MaMYBFA, targets *MaFLS2* to promote its expression and promote flavonol synthesis. Drought stress increases the transcript level of *MaHSF11*, enhancing the expression of *MaFLS1* and *MaF3′5′H1*. MaFLS1 and MaF3′5′H might interact to form a metabolon at the ER membrane to promote myricetin biosynthesis. MaHSF11 also directly activates the expression of *MaDREB1*. MaDREB1 binds to the *MaMYBFA3* promoter to regulate its expression. MaMYBFA1, MaMYBFA2, and MaMYBFA3 differentially regulate the expression of *MaFLS1* and *MaFLS2*. MRE, MYB recognition element; DRE, dehydration response element; ABRE, ABA response element; HSE, heat shock element. CHS (chalcone synthase) catalyzes the condensation of one molecule of *p*-coumaryl-CoA and three molecules of malonyl-CoA into naringenin chalcone, which is further converted into naringenin by CHI (chalcone isomerase). F3H (flavanone 3-hydroxylase) converts naringenin into DHK (dihydrokaempferol). F3′H (flavonoid 3′-hydroxylase) and F3′5′H (flavonoid 3′5′-hydroxylase) compete for the substrate DHK to synthesize DHQ (dihydroquercetin) and DHM (dihydromyricetin), respectively. FLS (flavonol synthase) converts DHK, DHQ, and DHM into kaempferol, quercetin, and myricetin, respectively. F3′5′H also physically interacts with FLS1 to form a metabolon to accelerate myricetin synthesis. Flavonols are likely transported into the vacuole by MATE transporters (multidrug and toxic compound extrusion).

Supplemental Figures

Supplemental Figure S1. Tissue-specific expression of *MaMYBFA1–3* genes. Relative expression of *MaMYBFA1–3* genes in different tissues of banana as determined by RT-qPCR. Tissues from three independent biological replicates were pooled and used for expression analysis. YL, young leaf; BRT, bract; P.STEM, pseudostem; UR.PL, unripe peel; UR.PP, unripe pulp; R.PL, ripe peel; R.PP, ripe pulp. Values shown are means ± SD of three technical replicates.

Supplemental Figure S2. Relative expression of structural flavonol biosynthesis gene transcripts in banana fruit discs transiently overexpressing *MaMYBFA1-FA3*. RT-qPCR-based transcript profiling of structural flavonol biosynthesis genes *MaCHS1–6*, *MaCHI1–2*, *MaF3H1–2*, *MaF3'5'H1–7*, and *MaF3'H* in banana fruit tissues of *MaMYBFA1–3* OE tissues. EV, empty vector control. Values shown are means ± SD of three technical replicates. One-way ANOVA was used for statistical analysis with Tukey's multiple comparison. **P* ≤ 0.05; ***P* ≤ 0.01; ****P* ≤ 0.001; *****P* ≤ 0.0001; ns, not significant.

Supplemental Figure S3. Phylogenetic analysis of MaHSF11 and previously characterized HSF genes from other plant species. The phylogenetic tree was constructed in MEGA-X32 software using the maximum-likelihood method with 1000 bootstrap values and was visualized with iTOL v.6.4 (http://itol.embl.de/). Accessions used are MdHSFA8a (NP_001281294.1), GmHSFB2b (XP_014619265.1), AtHSFA2 (AT2G26150), AtHSF1 (AT4G17750), AtHSF3 (AT5G16820), AtHSFA8 (AT1G67970), AtHSFC1 (AT3G24520), AtHSFA3 (AT5G03720), AtHSF4 (AT4G36990), AtHSFB2A (AT5G62020), AtHSFB2B (AT4G11660), and AtHSFA4A (AT4G18880). Md, *Malus domestica*; Gm, *Glycine max*; At, *Arabidopsis thaliana.*

Supplemental Figure S4. Relative expression of structural and regulatory flavonol biosynthesis genes in banana fruit discs overexpressing *MaHSF11*. RT-qPCR-based expression analysis of (a) *MaCHS1–6*, (b) *MaCHI1–2*, (c) *MaF3H1–2*, (d) *MaF3'H*, and (e) *MaMYBFA1–3* in *MaHSF11* OE banana fruit slices. Values shown are means ± SD of three technical replicates. EV, empty vector control. Two-way ANOVA was used with Tukey's multiple comparison. *P* ≤ 0.05; ***P* ≤ 0.01; ns, not significant. N.D., not determined.

Supplemental Figure S5. Phylogenetic analysis of MaDREB1 with previously characterized DREB genes from other plant species. The phylogenetic tree was constructed

in MEGA-X32 software using the maximum-likelihood method with 1000 bootstrap values
and visualized with iTOL v.6.4 (http://itol.embl.de/). Accessions used were AtDREB1B
(AAV80413.1), AtDREB1C (NP_567719.1), AtDREB1C (AAV80414.1), GmDREB3
(AAZ03388.1), GmDREB2 (NP_001237254.2), GmDREB2A2 (NP_001240942.3),
OsDREB1 (AAL40870.1), OsDREB1B (AAN85707.1), OsDREB3 (ADO79620.1), SlDREB3
(NP_001234707.1), HvDREB2 (AFH75401.1), and ZmDREB (NP_001105081.1). At,
Arabidopsis thaliana; Gm, *Glycine max*; Os, *Oryza sativa*; Sl, *Solanum lycopersicum*; Hv,
Hordeum vulgare; Zm, *Zea mays*)

Supplemental Figure S6. Relative expression of structural flavonol biosynthesis genes in
banana fruit tissues overexpressing *MaDREB1*. RT-qPCR-based expression analysis of (a)
MaCHS1–6, (b) *MaCHI1–2*, (c) *MaF3H1–2*, (d) *MaF3'5'H1–7*, and (e) *MaF3'H* in *MaDREB1*-
overexpressing tissues. Values shown are means ± SD of three technical replicates. Two-way
ANOVA was used with Tukey's multiple comparison. $P \leq 0.05$; **$P \leq 0.01$; ***$P \leq 0.001$; ns,
not significant.

Supplemental Figure S7. Relative expression analysis of *MaHSF11* in banana fruits
overexpressing *MaDREB*. Values shown are means ± SD of three technical replicates.
Unpaired Student's *t*-tests were used with Tukey's multiple comparison for statistical analysis.
***$P \leq 0.001$

Supplemental File S1. Promoter sequences with highlighted *cis* motifs used in this study.

Supplemental File S2. Accession numbers and two-letter species abbreviations used in
phylogenetic tree figures.

Supplemental Table S1. List of primers used in this study.

References

An, J.P., Zhang, X.W., Liu, Y.J., Wang, X.F., You, C.X. and Hao, Y.J., 2021. ABI5 regulates
ABA-induced anthocyanin biosynthesis by modulating the MYB1-bHLH3 complex in apple.
Journal of Experimental Botany, 72(4), pp.1460-1472.

Asif, M., Trivedi, P., Solomos, T. and Tucker, M., 2006. Isolation of high-quality RNA from
apple (Malus domestica) fruit. *Journal of agricultural and food chemistry*, 54(15), pp.5227-
5229.

Battat, M., Eitan, A., Rogachev, I., Hanhineva, K., Fernie, A., Tohge, T., Beekwilder, J. and

Aharoni, A., 2019. A MYB triad controls primary and phenylpropanoid metabolites for pollen

coat patterning. *Plant Physiology*, 180(1), pp.87-108.

Baudry, A., Heim, M.A., Dubreucq, B., Caboche, M., Weisshaar, B. and Lepiniec, L., 2004.

TT2, TT8, and TTG1 synergistically specify the expression of BANYULS and

proanthocyanidin biosynthesis in Arabidopsis thaliana. *The Plant Journal*, 39(3), pp.366-380.

Bhatia, C., Gaddam, S.R., Pandey, A. and Trivedi, P.K., 2021. COP1 mediates light-dependent

regulation of flavonol biosynthesis through HY5 in Arabidopsis. *Plant Science*, 303, p.110760.

Bian, X.H., Li, W., Niu, C.F., Wei, W., Hu, Y., Han, J.Q., Lu, X., Tao, J.J., Jin, M., Qin, H.

and Zhou, B., 2020. A class B heat shock factor selected for during soybean domestication

contributes to salt tolerance by promoting flavonoid biosynthesis. *New Phytologist*, 225(1),

pp.268-283.

Busche, M., Acatay, C., Martens, S., Weisshaar, B. and Stracke, R., 2021. Functional

characterisation of banana (Musa spp.) 2-oxoglutarate-dependent dioxygenases involved in

flavonoid biosynthesis. *Frontiers in Plant Science*, 12, p.701780.

Chapman, J.M. and Muday, G.K., 2021. Flavonols modulate lateral root emergence by

scavenging reactive oxygen species in Arabidopsis thaliana. *Journal of Biological Chemistry*,

296.

Chauhan, H., Khurana, N., Agarwal, P. and Khurana, P., 2011. Heat shock factors in rice

(Oryza sativa L.): genome-wide expression analysis during reproductive development and

abiotic stress. *Molecular Genetics and Genomics*, 286(2), pp.171-187.

Chen, H., Liu, L., Wang, L., Wang, S. and Cheng, X., 2016. VrDREB2A, a DREB-binding

transcription factor from Vigna radiata, increased drought and high-salt tolerance in transgenic

Arabidopsis thaliana. *Journal of plant research*, 129(2), pp.263-273.

Chen, H.H., Qu, L., Xu, Z.H., Zhu, J.K. and Xue, H.W., 2018. EL1-like casein kinases suppress

ABA signaling and responses by phosphorylating and destabilizing the ABA receptors

PYR/PYLs in Arabidopsis. *Molecular Plant*, 11(5), pp.706-719.

Chen, H.H., Qu, L., Xu, Z.H., Zhu, J.K. and Xue, H.W., 2018. EL1-like casein kinases suppress ABA signaling and responses by phosphorylating and destabilizing the ABA receptors PYR/PYLs in Arabidopsis. *Molecular Plant*, 11(5), pp.706-719.

Chen, J., Ullah, C., Reichelt, M., Gershenzon, J. and Hammerbacher, A., 2019. Sclerotinia sclerotiorum circumvents flavonoid defenses by catabolizing flavonol glycosides and aglycones. *Plant physiology*, 180(4), pp.1975-1987.

Choudhury, F.K., Rivero, R.M., Blumwald, E. and Mittler, R., 2017. Reactive oxygen species, abiotic stress and stress combination. *The Plant Journal*, 90(5), pp.856-867.

Cramer, G.R., Urano, K., Delrot, S., Pezzotti, M. and Shinozaki, K., 2011. Effects of abiotic stress on plants: a systems biology perspective. *BMC plant biology*, 11(1), pp.1-14.

Crosby, K.C., Pietraszewska-Bogiel, A., Gadella Jr, T.W. and Winkel, B.S., 2011. Förster resonance energy transfer demonstrates a flavonoid metabolon in living plant cells that displays competitive interactions between enzymes. *FEBS letters*, 585(14), pp.2193-2198.

Czemmel, S., Stracke, R., Weisshaar, B., Cordon, N., Harris, N.N., Walker, A.R., Robinson, S.P. and Bogs, J., 2009. The grapevine R2R3-MYB transcription factor VvMYBF1 regulates flavonol synthesis in developing grape berries. *Plant physiology*, 151(3), pp.1513-1530.

D'hont, A., Denoeud, F., Aury, J.M., Baurens, F.C., Carreel, F., Garsmeur, O., Noel, B., Bocs, S., Droc, G., Rouard, M. and Da Silva, C., 2012. The banana (Musa acuminata) genome and the evolution of monocotyledonous plants. *Nature*, 488(7410), pp.213-217.

Dubouzet, J.G., Sakuma, Y., Ito, Y., Kasuga, M., Dubouzet, E.G., Miura, S., Seki, M., Shinozaki, K. and Yamaguchi-Shinozaki, K., 2003. OsDREB genes in rice, Oryza sativa L., encode transcription activators that function in drought-, high-salt-and cold-responsive gene expression. *The Plant Journal*, 33(4), pp.751-763.

Friedrich, T., Oberkofler, V., Trindade, I., Altmann, S., Brzezinka, K., Lämke, J., Gorka, M., Kappel, C., Sokolowska, E., Skirycz, A. and Graf, A., 2021. Heteromeric HSFA2/HSFA3 complexes drive transcriptional memory after heat stress in Arabidopsis. *Nature Communications*, 12(1), pp.1-15.

Fujii, H., Verslues, P.E. and Zhu, J.K., 2007. Identification of two protein kinases required for
abscisic acid regulation of seed germination, root growth, and gene expression in Arabidopsis.
The Plant Cell, 19(2), pp.485-494.

Fujino, N., Tenma, N., Waki, T., Ito, K., Komatsuzaki, Y., Sugiyama, K., Yamazaki, T.,
Yoshida, S., Hatayama, M., Yamashita, S. and Tanaka, Y., 2018. Physical interactions among
flavonoid enzymes in snapdragon and torenia reveal the diversity in the flavonoid metabolon
organization of different plant species. *The Plant Journal*, 94(2), pp.372-392.

Giri, J., Vij, S., Dansana, P.K. and Tyagi, A.K., 2011. Rice A20/AN1 zinc-finger containing
stress-associated proteins (SAP1/11) and a receptor-like cytoplasmic kinase (OsRLCK253)
interact via A20 zinc-finger and confer abiotic stress tolerance in transgenic Arabidopsis plants.
New Phytologist, 191(3), pp.721-732.

Gong, S., Ding, Y., Hu, S., Ding, L., Chen, Z. and Zhu, C., 2019. The role of HD-Zip class I
transcription factors in plant response to abiotic stresses. *Physiologia plantarum*, 167(4),
pp.516-525.

Grunewald, W., De Smet, I., Lewis, D.R., Löfke, C., Jansen, L., Goeminne, G., Vanden
Bossche, R., Karimi, M., De Rybel, B., Vanholme, B. and Teichmann, T., 2012. Transcription
factor WRKY23 assists auxin distribution patterns during Arabidopsis root development
through local control on flavonol biosynthesis. *Proceedings of the National Academy of*
Sciences, 109(5), pp.1554-1559.

Guo, L., Gao, L., Ma, X., Guo, F., Ruan, H., Bao, Y., Xia, T. and Wang, Y., 2019. Functional
analysis of flavonoid 3′-hydroxylase and flavonoid 3′, 5′-hydroxylases from tea plant (Camellia
sinensis), involved in the B-ring hydroxylation of flavonoids. *Gene*, 717, p.144046.

Guo, M., Liu, J.H., Ma, X., Luo, D.X., Gong, Z.H. and Lu, M.H., 2016. The plant heat stress
transcription factors (HSFs): structure, regulation, and function in response to abiotic stresses.
Frontiers in plant science, 7, p.114.

Halbwirth, H., 2010. The creation and physiological relevance of divergent hydroxylation
patterns in the flavonoid pathway. *International journal of molecular sciences*, 11(2), pp.595-
621.

He, Y., Zhang, X., Li, L., Sun, Z., Li, J., Chen, X. and Hong, G., 2021. SPX4 interacts with
both PHR1 and PAP1 to regulate critical steps in phosphorus-status-dependent anthocyanin
biosynthesis. *New Phytologist*, 230(1), pp.205-217.

Huang, Y.C., Niu, C.Y., Yang, C.R. and Jinn, T.L., 2016. The heat stress factor HSFA6b
connects ABA signaling and ABA-mediated heat responses. *Plant physiology*, 172(2),
pp.1182-1199.

Huang, Y.C., Niu, C.Y., Yang, C.R. and Jinn, T.L., 2016. The heat stress factor HSFA6b
connects ABA signaling and ABA-mediated heat responses. *Plant physiology*, 172(2),
pp.1182-1199.

Jiang, X., Shi, Y., Fu, Z., Li, W.W., Lai, S., Wu, Y., Wang, Y., Liu, Y., Gao, L. and Xia, T.,
2020. Functional characterization of three flavonol synthase genes from Camellia sinensis:
Roles in flavonol accumulation. *Plant Science*, 300, p.110632.

Kalaji, H.M., Dąbrowski, P., Cetner, M.D., Samborska, I.A., Łukasik, I., Brestic, M., Zivcak,
M., Tomasz, H., Mojski, J., Kociel, H. and Panchal, B.M., 2017. A comparison between
different chlorophyll content meters under nutrient deficiency conditions. *Journal of Plant
Nutrition*, 40(7), pp.1024-1034.

Katoh, K. and Standley, D.M., 2013. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Molecular biology and evolution*, 30(4), pp.772-
780.

Kitamura, S., Oono, Y. and Narumi, I., 2016. Arabidopsis pab1, a mutant with reduced
anthocyanins in immature seeds from banyuls, harbors a mutation in the MATE transporter
FFT. *Plant molecular biology*, 90(1), pp.7-18.

Ko, M.J., Cheigh, C.I. and Chung, M.S., 2014. Relationship analysis between flavonoids
structure and subcritical water extraction (SWE). *Food chemistry*, 143, pp.147-155.

Koncz, C. and Schell, J., 1986. The promoter of TL-DNA gene 5 controls the tissue-specific
expression of chimaeric genes carried by a novel type of Agrobacterium binary vector.
Molecular and General Genetics MGG, 204(3), pp.383-396.

Kuang, J.F., Chen, J.Y., Liu, X.C., Han, Y.C., Xiao, Y.Y., Shan, W., Tang, Y., Wu, K.Q., He,
J.X. and Lu, W.J., 2017. The transcriptional regulatory network mediated by banana (Musa

acuminata) dehydration-responsive element binding (MaDREB) transcription factors in fruit

ripening. *New Phytologist*, 214(2), pp.762-781.

Kumar, K., Purayannur, S., Kaladhar, V.C., Parida, S.K. and Verma, P.K., 2018. mQTL-seq

and classical mapping implicates the role of an AT-HOOK MOTIF CONTAINING

NUCLEAR LOCALIZED (AHL) family gene in A scochyta blight resistance of chickpea.

Plant, Cell & Environment, 41(9), pp.2128-2140.

Kumar, S., Stecher, G., Li, M., Knyaz, C. and Tamura, K., 2018. MEGA X: molecular

evolutionary genetics analysis across computing platforms. *Molecular biology and evolution*,

35(6), p.1547.

Lago, J.H.G., Toledo-Arruda, A.C., Mernak, M., Barrosa, K.H., Martins, M.A., Tibério, I.F.

and Prado, C.M., 2014. *Structure-activity association of flavonoids in lung diseases.*

Molecules, 19(3), pp.3570-3595.

Lee, S.J., Kang, J.Y., Park, H.J., Kim, M.D., Bae, M.S., Choi, H.I. and Kim, S.Y., 2010.

DREB2C interacts with ABF2, a bZIP protein regulating abscisic acid-responsive gene

expression, and its overexpression affects abscisic acid sensitivity. *Plant physiology*, 153(2),

pp.716-727.

Lewis, D.R., Ramirez, M.V., Miller, N.D., Vallabhaneni, P., Ray, W.K., Helm, R.F., Winkel,

B.S. and Muday, G.K., 2011. Auxin and ethylene induce flavonol accumulation through

distinct transcriptional networks. *Plant physiology*, 156(1), pp.144-164.

Li, L., Peng, H., Tan, S., Zhou, J., Fang, Z., Hu, Z., Gao, L., Li, T., Zhang, W. and Chen, L.,

2020. Effects of early cold stress on gene expression in Chlamydomonas reinhardtii. *Genomics*,

112(2), pp.1128-1138.

Liang, T., Shi, C., Peng, Y., Tan, H., Xin, P., Yang, Y., Wang, F., Li, X., Chu, J., Huang, J.

and Yin, Y., 2020. Brassinosteroid-activated BRI1-EMS-SUPPRESSOR 1 inhibits flavonoid

biosynthesis and coordinates growth and UV-B stress responses in plants. *Plant Cell*, 32(10),

pp.3224-3239.

Lin, Z., Li, Y., Zhang, Z., Liu, X., Hsu, C.C., Du, Y., Sang, T., Zhu, C., Wang, Y., Satheesh,

V. and Pratibha, P., 2020. A RAF-SnRK2 kinase cascade mediates early osmotic stress

signaling in higher plants. *Nature communications*, 11(1), pp.1-10.

Liu, H.T., Gao, F., Li, G.L., Han, J.L., Liu, D.L., Sun, D.Y. and Zhou, R.G., 2008. The
calmodulin-binding protein kinase 3 is part of heat-shock signal transduction in Arabidopsis
thaliana. *The Plant Journal*, 55(5), pp.760-773.

López, M.A., Bannenberg, G. and Castresana, C., 2008. Controlling hormone signaling is a
plant and pathogen challenge for growth and survival. *Current opinion in plant biology*, 11(4),
pp.420-427.

Lu, Q., Tang, X., Tian, G., Wang, F., Liu, K., Nguyen, V., Kohalmi, S.E., Keller, W.A., Tsang,
E.W., Harada, J.J. and Rothstein, S.J., 2010. Arabidopsis homolog of the yeast TREX-2 mRNA
export complex: components and anchoring nucleoporin. *The Plant Journal*, 61(2), pp.259-
270.

Ma, D., Tang, H., Reichelt, M., Piirtola, E.M., Salminen, J.P., Gershenzon, J. and Constabel,
C.P., 2021. Poplar MYB117 promotes anthocyanin synthesis and enhances flavonoid B-ring
hydroxylation by up-regulating the flavonoid 3′, 5′-hydroxylase gene. *Journal of Experimental*
Botany, 72(10), pp.3864-3880.

Ma, Y., Szostkiewicz, I., Korte, A., Moes, D., Yang, Y., Christmann, A. and Grill, E., 2009.
Regulators of PP2C phosphatase activity function as abscisic acid sensors. *Science*, 324(5930),
pp.1064-1068.

Maloney, G.S., DiNapoli, K.T. and Muday, G.K., 2014. The anthocyanin reduced tomato
mutant demonstrates the role of flavonols in tomato lateral root and root hair development.
Plant Physiology, 166(2), pp.614-631.

Mann DG, LaFayette PR, Abercrombie LL, King ZR, Mazarei M, Halter MC, Poovaiah CR,
Baxter H, Shen H, Dixon RA, Parrott WA. Gateway-compatible vectors for high-throughput
gene functional analysis in switchgrass (Panicum virgatum L.) and other monocot species.
Plant biotechnology journal, 10(2), pp. 226-36.

Manna, M., Thakur, T., Chirom, O., Mandlik, R., Deshmukh, R. and Salvi, P., 2021.
Transcription factors as key molecular target to strengthen the drought stress tolerance in
plants. *Physiologia Plantarum*, 172(2), pp.847-868.

Martin, K., Kopperud, K., Chakrabarty, R., Banerjee, R., Brooks, R. and Goodin, M.M., 2009.
Transient expression in Nicotiana benthamiana fluorescent marker lines provides enhanced

definition of protein localization, movement and interactions in planta. *The Plant Journal*,
59(1), pp.150-162.

Matus, J.T., Loyola, R., Vega, A., Peña-Neira, A., Bordeu, E., Arce-Johnson, P. and Alcalde,
J.A., 2009. Post-veraison sunlight exposure induces MYB-mediated transcriptional regulation
of anthocyanin and flavonol synthesis in berry skins of Vitis vinifera. *Journal of experimental*
botany, 60(3), pp.853-867.

Medina-Puche, L., Cumplido-Laso, G., Amil-Ruiz, F., Hoffmann, T., Ring, L., Rodríguez-
Franco, A., Caballero, J.L., Schwab, W., Muñoz-Blanco, J. and Blanco-Portales, R., 2014.
MYB10 plays a major role in the regulation of flavonoid/phenylpropanoid metabolism during
ripening of Fragaria× ananassa fruits. *Journal of experimental botany*, 65(2), pp.401-417.

Mehra, P., Pandey, B.K., Verma, L. and Giri, J., 2019. A novel glycerophosphodiester
phosphodiesterase improves phosphate deficiency tolerance in rice. *Plant, Cell &*
Environment, 42(4), pp.1167-1179.

Mehrtens, F., Kranz, H., Bednarek, P. and Weisshaar, B., 2005. The Arabidopsis transcription
factor MYB12 is a flavonol-specific regulator of phenylpropanoid biosynthesis. *Plant*
physiology, 138(2), pp.1083-1096.

Mikkelsen, B.L., Olsen, C.E. and Lyngkjær, M.F., 2015. Accumulation of secondary
metabolites in healthy and diseased barley, grown under future climate levels of CO2, ozone
and temperature. *Phytochemistry*, 118, pp.162-173.

Misra, P., Pandey, A., Tiwari, M., Chandrashekar, K., Sidhu, O.P., Asif, M.H., Chakrabarty,
D., Singh, P.K., Trivedi, P.K., Nath, P. and Tuli, R., 2010. Modulation of transcriptome and
metabolome of tobacco by Arabidopsis transcription factor, AtMYB12, leads to insect
resistance. *Plant Physiology*, 152(4), pp.2258-2268.

Muhlemann, J.K., Younts, T.L. and Muday, G.K., 2018. Flavonols control pollen tube growth
and integrity by regulating ROS homeostasis during high-temperature stress. *Proceedings of*
the National Academy of Sciences, 115(47), pp.E11188-E11197.

Murcia, G., Fontana, A., Pontin, M., Baraldi, R., Bertazza, G. and Piccoli, P.N., 2017. ABA
and GA3 regulate the synthesis of primary and secondary metabolites related to alleviation
from biotic and abiotic stresses in grapevine. *Phytochemistry*, 135, pp.34-52.

Muthusamy, M., Uma, S., Backiyarani, S., Saraswathi, M.S. and Chandrasekar, A., 2016. Transcriptomic changes of drought-tolerant and sensitive banana cultivars exposed to drought stress. *Frontiers in plant science*, 7, pp.1609.

Nagamatsu, A., Masuta, C., Senda, M., Matsuura, H., Kasai, A., Hong, J.S., Kitamura, K., Abe, J. and Kanazawa, A., 2007. Functional analysis of soybean genes involved in flavonoid biosynthesis by virus-induced gene silencing. *Plant biotechnology journal*, 5(6), pp.778-790.

Naik, J., Rajput, R., Pucker, B., Stracke, R. and Pandey, A., 2021. The R2R3-MYB transcription factor MtMYB134 orchestrates flavonol biosynthesis in Medicago truncatula. *Plant Molecular Biology*, 106(1), pp.157-172.

Née, G., Kramer, K., Nakabayashi, K., Yuan, B., Xiang, Y., Miatton, E., Finkemeier, I. and Soppe, W.J., 2017. DELAY OF GERMINATION1 requires PP2C phosphatases of the ABA signalling pathway to control seed dormancy. *Nature communications*, 8(1), pp.1-9.

Oh, M.M., Trick, H.N. and Rajashekar, C.B., 2009. Secondary metabolism and antioxidants are involved in environmental adaptation and stress tolerance in lettuce. *Journal of plant physiology*, 166(2), pp.180-191.

Pandey, A., Alok, A., Lakhwani, D., Singh, J., Asif, M.H. and Trivedi, P.K., 2016. Genome-wide expression analysis and metabolite profiling elucidate transcriptional regulation of flavonoid biosynthesis and modulation under abiotic stresses in banana. *Scientific reports*, 6(1), pp.1-13.

Pandey, A., Misra, P. and Trivedi, P.K., 2015. Constitutive expression of Arabidopsis MYB transcription factor, AtMYB11, in tobacco modulates flavonoid biosynthesis in favor of flavonol accumulation. *Plant cell reports*, 34(9), pp.1515-1528.

Pandey, A., Misra, P., Choudhary, D., Yadav, R., Goel, R., Bhambhani, S., Sanyal, I., Trivedi, R. and Kumar Trivedi, P., 2015. AtMYB12 expression in tomato leads to large scale differential modulation in transcriptome and flavonoid content in leaf and fruit tissues. *Scientific reports*, 5(1), pp.1-14.

Pandey, A., Misra, P., Khan, M.P., Swarnkar, G., Tewari, M.C., Bhambhani, S., Trivedi, R., Chattopadhyay, N. and Trivedi, P.K., 2014. Co-expression of Arabidopsis transcription factor, At MYB 12, and soybean isoflavone synthase, Gm IFS 1, genes in tobacco leads to enhanced

biosynthesis of isoflavones and flavonols resulting in osteoprotective activity. *Plant biotechnology journal*, 12(1), pp.69-80.

Peng, Y., Thrimawithana, A.H., Cooney, J.M., Jensen, D.J., Espley, R.V. and Allan, A.C., 2020. The proanthocyanin-related transcription factors MYBC1 and WRKY44 regulate branch points in the kiwifruit anthocyanin pathway. *Scientific Reports*, 10(1), pp.1-19.

Pucker, B., Pandey, A., Weisshaar, B. and Stracke, R., 2020. The R2R3-MYB gene family in banana (Musa acuminata): Genome-wide identification, classification and expression patterns. *PloS one*, 15(10), pp.e0239275.

Qian, C., Chen, Z., Liu, Q., Mao, W., Chen, Y., Tian, W., Liu, Y., Han, J., Ouyang, X. and Huang, X., 2020. Coordinated transcriptional regulation by the UV-B photoreceptor and multiple transcription factors for plant UV-B responses. *Molecular plant*, 13(5), pp.777-792.

Rajput, R., Naik, J., Stracke, R. and Pandey, A., 2022. Interplay between R2R3 MYB-type activators and repressors regulates proanthocyanidin biosynthesis in banana (Musa acuminata). *New Phytologist*, 236, pp.1108-1127.

Rajput, R., Tyagi, S., Naik, J., Pucker, B., Stracke, R. and Pandey, A., 2022. The R2R3-MYB gene family in Cicer arietinum: genome-wide identification and expression analysis leads to functional characterization of proanthocyanidin biosynthesis regulators in the seed coat. *Planta*, 256(4), pp.1-24.

Scharf, K.D., Berberich, T., Ebersberger, I. and Nover, L., 2012. The plant heat stress transcription factor (Hsf) family: structure, function and evolution. *Biochimica et Biophysica Acta (BBA)-Gene Regulatory Mechanisms*, 1819(2), pp.104-119.

Schramm, F., Larkindale, J., Kiehlmann, E., Ganguli, A., Englich, G., Vierling, E. and Von Koskull-Döring, P., 2008. A cascade of transcription factor DREB2A and heat stress transcription factor HsfA3 regulates the heat stress response of Arabidopsis. *The Plant Journal*, 53(2), pp.264-274.

Schwinn, K.E., Ngo, H., Kenel, F., Brummell, D.A., Albert, N.W., McCallum, J.A., Pither-Joyce, M., Crowhurst, R.N., Eady, C. and Davies, K.M., 2016. The onion (Allium cepa L.) R2R3-MYB gene MYB1 regulates anthocyanin biosynthesis. *Frontiers in Plant Science*, 7, p.1865.

Shan, W., KUANG, J.F., LU, W.J. and CHEN, J.Y., 2014. Banana fruit NAC transcription factor MaNAC 1 is a direct target of MaICE 1 and involved in cold stress through interacting with MaCBF 1. *Plant, cell & environment*, 37(9), pp.2116-2127.

Shan, X., Li, Y., Yang, S., Yang, Z., Qiu, M., Gao, R., Han, T., Meng, X., Xu, Z., Wang, L. and Gao, X., 2020. The spatio-temporal biosynthesis of floral flavonols is controlled by differential phylogenetic MYB regulators in Freesia hybrida. *New Phytologist*, 228(6), pp.1864-1879.

Shi, C., Luo, P., Du, Y.T., Chen, H., Huang, X., Cheng, T.H., Luo, A., Li, H.J., Yang, W.C., Zhao, P. and Sun, M.X., 2019. Maternal control of suspensor programmed cell death via gibberellin signaling. *Nature communications*, 10(1), pp.1-12.

Shih, C.H., Chu, H., Tang, L.K., Sakamoto, W., Maekawa, M., Chu, I.K., Wang, M. and Lo, C., 2008. Functional characterization of key structural genes in rice flavonoid biosynthesis. *Planta*, 228(6), pp.1043-1054.

Si, J., Ye, B.B., Liu, Z.L., Xiao, X.M., Yang, Y.Y., Fan, Z.Q., Shan, W., Kuang, J.F., Lu, W.J., Su, X.G. and Chen, J.Y., 2022. Transcriptional repression of MaRBOHs by MaHsf26 is associated with heat shock-alleviated chilling injury in banana fruit. *Postharvest Biology and Technology*, 193, p.112056.

Song, Z., Lai, X., Chen, H., Wang, L., Pang, X., Hao, Y., Lu, W., Chen, W., Zhu, X. and Li, X., 2022. Role of MaABI5-like in abscisic acid-induced cold tolerance of 'Fenjiao'banana fruit. *Horticulture Research*, 9, p.uhac130.

Stracke, R., Favory, J.J., Gruber, H., Bartelniewoehner, L., Bartels, S., Binkert, M., Funk, M., Weisshaar, B. and Ulm, R., 2010. The Arabidopsis bZIP transcription factor HY5 regulates expression of the PFG1/MYB12 gene in response to light and ultraviolet-B radiation. *Plant, cell & environment*, 33(1), pp.88-103.

Stracke, R., FAVORY, J.J., Gruber, H., Bartelniewoehner, L., Bartels, S., Binkert, M., Funk, M., Weisshaar, B. and Ulm, R., 2010. The Arabidopsis bZIP transcription factor HY5 regulates expression of the PFG1/MYB12 gene in response to light and ultraviolet-B radiation. *Plant, cell & environment*, 33(1), pp.88-103.

Stracke, R., Holtgräwe, D., Schneider, J., Pucker, B., Rosleff Sörensen, T. and Weisshaar, B.,
2014. Genome-wide identification and characterisation of R2R3-MYB genes in sugar beet
(Beta vulgaris). *BMC plant biology*, 14(1), pp.1-17.

Stracke, R., Jahns, O., Keck, M., Tohge, T., Niehaus, K., Fernie, A.R. and Weisshaar, B., 2010.
Analysis of PRODUCTION OF FLAVONOL GLYCOSIDES-dependent flavonol glycoside
accumulation in Arabidopsis thaliana plants reveals MYB11-, MYB12-and MYB111-
independent flavonol glycoside accumulation. *New Phytologist*, 188(4), pp.985-1000.

Stracke, R., Werber, M. and Weisshaar, B., 2001. The R2R3-MYB gene family in Arabidopsis
thaliana. *Current opinion in plant biology*, 4(5), pp.447-456.

Tan, H., Man, C., Xie, Y., Yan, J., Chu, J. and Huang, J., 2019. A crucial role of GA-regulated
flavonol biosynthesis in root growth of Arabidopsis. *Molecular plant*, 12(4), pp.521-537.

Teale, W.D., Pasternak, T., Dal Bosco, C., Dovzhenko, A., Kratzat, K., Bildl, W., Schwörer,
M., Falk, T., Ruperti, B., V Schaefer, J. and Shahriari, M., 2021. Flavonol-mediated
stabilization of PIN efflux complexes regulates polar auxin transport. *The EMBO Journal*,
40(1), p.e104416.

Umezawa, T., Sugiyama, N., Mizoguchi, M., Hayashi, S., Myouga, F., Yamaguchi-Shinozaki,
K., Ishihama, Y., Hirayama, T. and Shinozaki, K., 2009. Type 2C protein phosphatases directly
regulate abscisic acid-activated protein kinases in Arabidopsis. *Proceedings of the National
Academy of sciences*, 106(41), pp.17588-17593.

Upendra, K., Shekhawat, S., Srinivas, L. and Ganapathi, T.R., 2011. MusaDHN-1, a novel
multiple stress-inducible SK^ sub 3^-type dehydrin gene, contributes affirmatively to drought-
and salt-stress tolerance in banana. *Planta*, 234(5), p.915.

Waki, T., Yoo, D., Fujino, N., Mameda, R., Denessiouk, K., Yamashita, S., Motohashi, R.,
Akashi, T., Aoki, T., Ayabe, S.I. and Takahashi, S., 2016. Identification of protein–protein
interactions of isoflavonoid biosynthetic enzymes with 2-hydroxyisoflavanone synthase in
soybean (Glycine max (L.) Merr.). *Biochemical and biophysical research communications*,
469(3), pp.546-551.

Walter, M., Chaban, C., Schütze, K., Batistic, O., Weckermann, K., Näke, C., Blazevic, D.,
Grefen, C., Schumacher, K., Oecking, C. and Harter, K., 2004. Visualization of protein

interactions in living plant cells using bimolecular fluorescence complementation. *The Plant Journal*, 40(3), pp.428-438.

Wang, J., Ren, Y., Liu, X., Luo, S., Zhang, X., Liu, X., Lin, Q., Zhu, S., Wan, H., Yang, Y. and Zhang, Y., 2021. Transcriptional activation and phosphorylation of OsCNGC9 confer enhanced chilling tolerance in rice. *Molecular plant*, 14(2), pp.315-329.

Wang, N., Liu, W., Yu, L., Guo, Z., Chen, Z., Jiang, S., Xu, H., Fang, H., Wang, Y., Zhang, Z. and Chen, X., 2020. HEAT SHOCK FACTOR A8a modulates flavonoid synthesis and drought tolerance. *Plant Physiology*, 184(3), pp.1273-1290.

Wang, N., Xu, H., Jiang, S., Zhang, Z., Lu, N., Qiu, H., Qu, C., Wang, Y., Wu, S. and Chen, X., 2017. MYB12 and MYB22 play essential roles in proanthocyanidin and flavonol synthesis in red-fleshed apple (Malus sieversii f. niedzwetzkyana). *The Plant Journal*, 90(2), pp.276-292.

Wang, X., Wang, L., Wang, Y., Liu, H., Hu, D., Zhang, N., Zhang, S., Cao, H., Cao, Q., Zhang, Z. and Tang, S., 2018. Arabidopsis PCaP2 plays an important role in chilling tolerance and ABA response by activating CBF-and SnRK2-mediated transcriptional regulatory network. *Frontiers in plant science*, 9, p.215.

Wang, X., Yu, R. and Li, J., 2021. Using genetic engineering techniques to develop banana cultivars with fusarium wilt resistance and ideal plant architecture. *Frontiers in Plant Science*, 11, p.617528.

Wang, Z., Huang, S., Jia, C., Liu, J., Zhang, J., Xu, B. and Jin, Z., 2013. Molecular cloning and expression of five glutathione S-transferase (GST) genes from Banana (Musa acuminata L. AAA group, cv. Cavendish). *Plant cell reports*, 32(9), pp.1373-1380.

Watkins, J.M., Chapman, J.M. and Muday, G.K., 2017. Abscisic acid-induced reactive oxygen species are modulated by flavonols to control stomata aperture. *Plant Physiology*, 175(4), pp.1807-1825.

Wei, Y., Hu, W., Xia, F., Zeng, H., Li, X., Yan, Y., He, C. and Shi, H., 2016. Heat shock transcription factors in banana: genome-wide characterization and expression profile analysis during development and stress response. *Scientific Reports*, 6(1), pp.1-11.

Wu, T.Y., Hoh, K.L., Boonyaves, K., Krishnamoorthi, S. and Urano, D., 2022. Diversification of heat shock transcription factors expanded thermal stress responses during early plant evolution. *The Plant Cell*, 34(10), pp.3557-3576.

Xu, F., Li, L., Zhang, W., Cheng, H., Sun, N., Cheng, S. and Wang, Y., 2012. Isolation, characterization, and function analysis of a flavonol synthase gene from Ginkgo biloba. *Molecular Biology Reports*, 39(3), pp.2285-2296.

Xu, Y., Hu, W., Liu, J., Song, S., Hou, X., Jia, C., Li, J., Miao, H., Wang, Z., Tie, W. and Xu, B., 2020. An aquaporin gene MaPIP2-7 is involved in tolerance to drought, cold and salt stresses in transgenic banana (Musa acuminata L.). *Plant Physiology and Biochemistry*, 147, pp.66-76.

Yang, M., Han, X., Yang, J., Jiang, Y. and Hu, Y., 2021. The Arabidopsis circadian clock protein PRR5 interacts with and stimulates ABI5 to modulate abscisic acid signaling during seed germination. *The Plant Cell*, 33(9), pp.3022-3041.

Yang, Q., Song, Z., Dong, B., Niu, L., Cao, H., Li, H., Du, T., Liu, T., Yang, W., Meng, D. and Fu, Y., 2021. Hyperoside regulates its own biosynthesis via MYB30 in promoting reproductive development and seed set in okra. *Plant Physiology*, 185(3), pp.951-968.

Yoshida, T., Fujita, Y., Maruyama, K., Mogami, J., Todaka, D., Shinozaki, K. and Yamaguchi-Shinozaki, K.A.Z.U.K.O., 2015. Four A rabidopsis AREB/ABF transcription factors function predominantly in gene expression downstream of SnRK2 kinases in abscisic acid signalling in response to osmotic stress. *Plant, cell & environment*, 38(1), pp.35-49.

Yoshida, T., Ohama, N., Nakajima, J., Kidokoro, S., Mizoi, J., Nakashima, K., Maruyama, K., Kim, J.M., Seki, M., Todaka, D. and Osakabe, Y., 2011. Arabidopsis HsfA1 transcription factors function as the main positive regulators in heat shock-responsive gene expression. *Molecular Genetics and Genomics*, 286(5), pp.321-332.

Zhang, J., Subramanian, S., Stacey, G. and Yu, O., 2009. Flavones and flavonols play distinct critical roles during nodulation of Medicago truncatula by Sinorhizobium meliloti. *The Plant Journal*, 57(1), pp.171-183.

Zhang, K., Logacheva, M.D., Meng, Y., Hu, J., Wan, D., Li, L., Janovská, D., Wang, Z., Georgiev, M.I., Yu, Z. and Yang, F., 2018. Jasmonate-responsive MYB factors spatially

repress rutin biosynthesis in Fagopyrum tataricum. *Journal of Experimental Botany*, 69(8), pp.1955-1966.

Zhang, L., Cenci, A., Rouard, M., Zhang, D., Wang, Y., Tang, W. and Zheng, S.J., 2019. Transcriptomic analysis of resistant and susceptible banana corms in response to infection by Fusarium oxysporum f. sp. cubense tropical race 4. *Scientific reports*, 9(1), pp.1-14.

Zhang, X., He, Y., Li, L., Liu, H. and Hong, G., 2021. Involvement of the R2R3-MYB transcription factor MYB21 and its homologs in regulating flavonol accumulation in Arabidopsis stamen. *Journal of experimental botany*, 72(12), pp.4319-4332.

ZHAO, M.L., WANG, J.N., Shan, W., FAN, J.G., KUANG, J.F., WU, K.Q., LI, X.P., CHEN, W.X., HE, F.Y., CHEN, J.Y. and LU, W.J., 2013. Induction of jasmonate signalling regulators MaMYC2s and their physical interactions with MaICE1 in methyl jasmonate-induced chilling tolerance in banana fruit. *Plant, cell & environment*, 36(1), pp.30-51.

Zhao, C., Liu, X., Gong, Q., Cao, J., Shen, W., Yin, X., Grierson, D., Zhang, B., Xu, C., Li, X. and Chen, K., 2021. Three AP2/ERF family members modulate flavonoid synthesis by regulating type IV chalcone isomerase in citrus. *Plant biotechnology journal*, 19(4), pp.671-688.

Zhao, L., Hu, Y., Chong, K. and Wang, T., 2010. ARAG1, an ABA-responsive DREB gene, plays a role in seed germination and drought tolerance of rice. *Annals of Botany*, 105(3), pp.401-409.

Zimmermann, I.M., Heim, M.A., Weisshaar, B. and Uhrig, J.F., 2004. Comprehensive identification of Arabidopsis thaliana MYB transcription factors interacting with R/B-like BHLH proteins. *The Plant Journal*, 40(1), pp.22-34.

Zinsmeister, J., Berriri, S., Basso, D.P., Ly-Vu, B., Dang, T.T., Lalanne, D., da Silva, E.A.A., Leprince, O. and Buitink, J., 2020. The seed-specific heat shock factor A9 regulates the depth of dormancy in Medicago truncatula seeds via ABA signalling. *Plant, Cell & Environment*, 43(10), pp.2508-2522.

Figure 1. Identification of candidate R2R3-MYB flavonol regulators from the banana genome and their subcellular localization. (a) Various R2R3-MYB proteins were classified into three major groups as flavonol-, anthocyanin-, or proanthocyanidin biosynthesis–specific regulators using landmark flavonoid-specific R2R3-MYBs from different plant species. Numbers on each branch show branch length, and solid dark blue circles represent bootstrap values. (b) Amino acid sequence alignment of candidate SG7 R2R3-MYB transcription factors (MaMYBFA1, MaMYBFA2, and MaMYBFA3, where FA stands for flavonol activator) and selected flavonol regulators from other plant species. The R2 and R3 domains are highlighted in grey. The SG7-1 and SG7-2 motifs, found in subgroup 7 R2R3-MYBs, are highlighted in dark green and yellow, respectively. Accession numbers and two-letter species abbreviations are provided in Supplemental File S2. (c) Subcellular localization of MaMYBFAs. MaMYBFA-YFP fusion proteins were expressed together with the nuclear marker NLS-RFP in *N. benthamiana* leaf epidermal cells. Confocal imaging shows co-localization of YFP and RFP fluorescence, indicating nuclear localization of the three MaMYBFA proteins. Representative images are shown with scale bar of 50 μm.

Figure 2. MaMYBFA1, MaMYBFA2, and MaMYBFA3 differentially regulate the expression of *MaFLS1* and *MaFLS2* and modulate flavonol biosynthesis in banana. (a) Diagram of the plant expression cassette used for transient overexpression (OE) of *MaMYBFA* in banana fruit discs. The *Zea mays ubiquitin1* promoter (*proZmUBI1*) drives the expression of *MaMYBFA*. The expression of the *GUSPlus* marker gene is driven by the *Phaseolus vulgaris ubiquitin1* promoter (*proPvUBI*). (b) Representative images showing GUS staining of transiently transformed banana fruit discs indicating successful transformation. Images were taken 3 days after transformation. (c-d) Relative expression of *MaMYBFA1–3* (c) and *MaFLS1* and *MaFLS2* (d) OE lines in transiently transformed *MaMYBFA* OE banana fruit tissues as determined by RT-qPCR. EV, empty vector control. (e) Flavonol quantification showing differential alteration in the accumulation of kaempferol, quercetin, and myricetin aglycones in the *MaMYBFA* OE banana fruit discs. Values are given in µg per 100 g of dry weight (DW) of fruit discs of three replicates. One-way ANOVA with Tukey's multiple comparison was used for statistical analyses. *$P \leq 0.05$; **$P \leq 0.01$; ***$P \leq 0.001$; ****$P \leq 0.0001$; ns, not significant.

Figure 3. MaMYBFAs regulate the expression of *MaFLS1* and *MaFLS2 in vivo*. (a) Schematic representation of the 2556- and 1500-bp promoter fragments of *MaFLS1* and *MaFLS2*. The locations of the different MYB core (CNGTTR) sequences and their positions upstream of the transcription start sites (−1) are indicated by red ovals. A plus (+) sign with a number indicates the length of the 5′ UTR. (b) Diagram of the pBTdest effector and the p635nRRF reporter constructs. Constitutive cauliflower mosaic virus (CaMV) promoter drives the expression of transcription factor encoding genes and renilla *luciferase* (REN). Desired amplified promoter fragments were placed upstream of firefly *luciferase* (LUC) (c) Dual-luciferase assay showing the altered activity of the promoters *proMaFLS1–2508* and *proMaFLS2–1500* when transiently co-infiltrated into *N. benthamiana* leaves with the effectors MaMYBFA1, MaMYBFA2, and MaMYBFA3. Promoter activity is represented as the LUC/REN ratio in four biological replicates (*n* = 4).

Figure 4. MaHSF11 directly binds to the *MaFLS1* promoter and regulates the expression of *MaFLS1* to modulate flavonol biosynthesis in banana. (a) Heatmap showing expression of five banana candidate *HSF* genes before and after drought treatment (Grand Naine cultivar, AAA genome). FPKM values for the indicated genes were extracted from publicly available RNA-Seq datasets and are represented according to the 0–7 color scale shown. (b) Dual-luciferase assay showing transactivation of *proMaFLS1–2556* by MaHSF. Values are shown as means ± SD of the LUC/REN ratios (*n* = 4). (c) ChIP-qPCR assay analyzing the binding of MaHSF11 to fragments F1 and F2 of *proMaFLS1*, both of which contain the HSE consensus sequence (nTTCnnGAAn). Fold enrichment values are given as means ± SD of two biological replicates. AB, antibody. (d) Yeast one-hybrid assay showing interaction of MaHSF11 with the two HSE-containing *proMaFLS1* fragments F1 and F2. The empty pGADT7 vector (EV) was used as a negative control. Interaction was indicated by yeast growth on synthetic Ura– dropout media supplemented with aureobasidin (AbA). AD, activation domain.

Figure 5. MaHSF11 promotes flavonol synthesis in banana fruit. (a) Subcellular localization of MaHSF11 in *N. benthamiana* leaves. The YFP-MaHSF11 fusion protein localizes to both the nucleus and the cytoplasm and coincides with the signal from the nuclear localization signal-red fluorescent protein (NLS-RFP) fusion in *N. benthamiana* leaves as observed in confocal microscopy. Scale bar, 20 μm. (b) GUS staining of banana fruit slices indicating successful transformation with the empty vector (EV) and the *proZmUBI:MaHSF11* construct. (c) Relative expression of *MaHSF11*, *MaFLS1*, and *MaFLS2* in transiently transformed banana fruit slices overexpressing (OE) *proZmUBI:MaHSF11* as determined by RT-qPCR. The empty vector was transformed as a control (expression level set to 1). Values shown are means ± SD of three technical replicates. (d) Quantification of the flavonols kaempferol, quercetin, and myricetin in transformed banana fruit slices. Values are represented as means ± SD of three replicates. Unpaired student's *t*-tests were used for statistical analyses. *$P \leq 0.05$; **$P \leq 0.01$; ***$P \leq 0.001$.

(a)

(d)

proMaF3'5'H1 F1 ATG

−1335 HSE
(GGAATATTCT) TSS
(−1)

(b)

-AbA   +AbA

proMaF3'5'H1(F1)

pGADT7 (EV)

MaHSF11-AD

pro35S:YFP-MaHSF11

■ anti-IgG AB
■ anti-GFP AB

proMaF3'5'H1   F1

(c)

proMaF3'5'H1-1335
(n=4)

EV   *MaHSF11*

Figure 6. MaHSF11 promotes flavonoid B-ring hydroxylation by modulating the expression of *MaF3'5'H1*. (a) Relative expression of *MaF3'5'H1* genes in transformed banana fruit slices transiently overexpressing *proZmUBI:MaHSF11*, as determined by RT-qPCR (EV, empty vector control). Values shown are means ± SD of three technical replicates. (b) Yeast one-hybrid assay showing the interaction of MaHSF11 with *proMaF3'5'H1–100* fragment F1 carrying HSE (GGAATATTCT). pGADT7 was used as a negative control. Growth was checked on synthetic dropout medium supplemented with aureobasidin. (c) Dual-luciferase assay showing transactivation of *proMaF3'5'H1–1335* by MaHSF11. The promoter sequence driving the expression of luciferase (LUC) was co-infiltrated with the EV or *MaHSF11* constructs in *N. benthamiana* leaves. Values are shown as means ± SD of the LUC/REN ratio (*n* = 4). (d) ChIP-qPCR assay showing binding of MaHSF11 to fragment F1 of *proMaF3'5'H1* carrying an HSE (nGAAnnTTCn). The promoter fragment carrying HSE was co-infiltrated into *N. benthamiana* leaves with the *pro35S:YFP-MaHSF11* construct. Fold enrichment values are represented as means ± SD of two biological replicates. An anti-IgG antibody was used as a negative control, and an anti-GFP antibody was used to precipitate MaHSF11-enriched chromatin. Unpaired Student's *t*-tests were used for statistical analyses. *$*P \leq 0.05$.

Figure 7. MaHSF11 acts upstream of *MaDREB1* to positively regulate its expression in banana. (a) Heatmap showing expression of candidate DREB genes before and after drought treatment in banana (Grand Naine cultivar, AAA genome). FPKM values for the indicated genes were extracted from publicly available RNA-Seq datasets and plotted according to the 0–7 color scale shown. (b) Relative expression of *MaDREB1* in banana fruit tissues transiently transformed with EV or *proZmUBI:MaHSF11* as determined by RT-qPCR. (c) Yeast one-hybrid assay showing interaction of MaHSF11 with the F1 and F2 fragments of *proMaDREB1–100* carrying an HSE (nGAAnnTTCn). pGADT7 was used as a negative control. Growth was checked on synthetic dropout medium supplemented with aureobasidin. (d) Dual-luciferase assay showing transactivation of *proMaDREB1–998* by MaHSF11. Promoter sequence driving the expression of LUC was co-infiltrated with empty vector and *MaHSF11* construct in *N. benthamiana* leaves. Values shown are means ± SD of LUC/REN ratio (*n* = 4). (e) ChIP-qPCR assay showing binding of MaHSF11 to *proMaDREB1*. A promoter fragment carrying an HSE was co-infiltrated into *N. benthamiana* leaf cells with the *pro35S:MaHSF11-YFP* construct. Fold enrichment values are represented as means ± SD of two biological replicates. An anti-IgG antibody was used as a negative control, and an anti-GFP antibody was used to precipitate MaHSF11 enriched chromatin. Unpaired Student's t-tests were used for statistical analyses. *$P \leq 0.05$; ***$P \leq 0.001$; ns, not significant.

Figure 8. MaDREB1 directly binds to the promoter of *MaMYBFA3* to regulate flavonol biosynthesis. (a) Yeast one-hybrid assay showing interaction of MaDREB1 with the *proMaMYBFA3–1892* fragment carrying three DREs (CCGAC). pGADT7 was used as a negative control. Growth was checked on synthetic dropout medium supplemented with aureobasidin. (b) Dual-luciferase assay showing transactivation of *proMaMYBFA3–2508* by MaDREB1. The *proMaMYBFA3–2508* sequence driving the expression of LUC was co-infiltrated with the empty vector or *MaDREB1* construct into *N. benthamiana* leaves. Values are shown as means ± SD of the LUC/REN ratio (*n* = 4). (c) GUS staining of transformed fruit. (d) RT-qPCR-based relative expression of *MaDREB1*, *MaMYBFAs*, *MaFLS1*, and *MaFLS2* in transformed banana fruit tissues transiently expressing EV and *proZmUBI:MaDREB1*. EV was transformed as a control. Values shown are means ± SD of three technical replicates. (e) Quantification of the flavonols kaempferol, quercetin, and myricetin in transformed fruit tissues. Values are represented as means ± SD of three replicates. Unpaired Student's *t*-tests were used for statistical analyses. *$P \leq 0.05$; **$P \leq 0.01$; ***$P \leq 0.001$; ****$P \leq 0.0001$.

Figure 9. A working model showing the modulation of hydroxylated flavonol synthesis by the MaHSF11–MaDREB1–MaMYBFA regulatory module. The banana SG7 R2R3 MYB, MaMYBFA, targets *MaFLS2* to promote its expression and promote flavonol synthesis. Drought stress increases the transcript level of *MaHSF11*, enhancing the expression of *MaFLS1* and *MaF3′5′H1*. MaFLS1 and MaF3′5′H might interact to form a metabolon at the ER membrane to promote myricetin biosynthesis. MaHSF11 also directly activates the expression of *MaDREB1*. MaDREB1 binds to the *MaMYBFA3* promoter to regulate its expression. MaMYBFA1, MaMYBFA2, and MaMYBFA3 differentially regulate the expression of *MaFLS1* and *MaFLS2*. MRE, MYB recognition element; DRE, dehydration response element; ABRE, ABA response element; HSE, heat shock element. CHS (chalcone synthase) catalyzes the condensation of one molecule of *p*-coumaryl-CoA and three molecules of malonyl-CoA into naringenin chalcone, which is further converted into naringenin by CHI (chalcone isomerase). F3H (flavanone 3-hydroxylase) converts naringenin into DHK (dihydrokaempferol). F3′H (flavonoid 3′-hydroxylase) and F3′5′H (flavonoid 3′5′-hydroxylase) compete for the substrate DHK to synthesize DHQ (dihydroquercetin) and DHM (dihydromyricetin), respectively. FLS (flavonol synthase) converts DHK, DHQ, and DHM into kaempferol, quercetin, and myricetin, respectively. F3′5′H also physically interacts with FLS1 to form a metabolon to accelerate myricetin synthesis. Flavonols are likely transported into the vacuole by MATE transporters (multidrug and toxic compound extrusion).

Parsed Citations

An, J.P., Zhang, X.W., Liu, Y.J., Wang, X.F., You, C.X. and Hao, Y.J., 2021. ABI5 regulates ABA-induced anthocyanin biosynthesis by modulating the MYB1-bHLH3 complex in apple. Journal of Experimental Botany, 72(4), pp.1460-1472.
  Google Scholar: Author Only Title Only Author and Title

Asif, M., Trivedi, P., Solomos, T. and Tucker, M., 2006. Isolation of high-quality RNA from apple (Malus domestica) fruit. Journal of agricultural and food chemistry, 54(15), pp.5227-5229.
  Google Scholar: Author Only Title Only Author and Title

Battat, M., Eitan, A., Rogachev, I., Hanhineva, K., Fernie, A., Tohge, T., Beekwilder, J. and Aharoni, A., 2019. A MYB triad controls primary and phenylpropanoid metabolites for pollen coat patterning. Plant Physiology, 180(1), pp.87-108.
  Google Scholar: Author Only Title Only Author and Title

Baudry, A., Heim, M.A., Dubreucq, B., Caboche, M., Weisshaar, B. and Lepiniec, L., 2004. TT2, TT8, and TTG1 synergistically specify the expression of BANYULS and proanthocyanidin biosynthesis in Arabidopsis thaliana. The Plant Journal, 39(3), pp.366-380.
  Google Scholar: Author Only Title Only Author and Title

Bhatia, C., Gaddam, S.R., Pandey, A. and Trivedi, P.K., 2021. COP1 mediates light-dependent regulation of flavonol biosynthesis through HY5 in Arabidopsis. Plant Science, 303, p.110760.
  Google Scholar: Author Only Title Only Author and Title

Bian, X.H., Li, W., Niu, C.F., Wei, W., Hu, Y., Han, J.Q., Lu, X., Tao, J.J., Jin, M., Qin, H. and Zhou, B., 2020. A class B heat shock factor selected for during soybean domestication contributes to salt tolerance by promoting flavonoid biosynthesis. New Phytologist, 225(1), pp.268-283.
  Google Scholar: Author Only Title Only Author and Title

Busche, M., Acatay, C., Martens, S., Weisshaar, B. and Stracke, R., 2021. Functional characterisation of banana (Musa spp.) 2-oxoglutarate-dependent dioxygenases involved in flavonoid biosynthesis. Frontiers in Plant Science, 12, p.701780.
  Google Scholar: Author Only Title Only Author and Title

Chapman, J.M. and Muday, G.K., 2021. Flavonols modulate lateral root emergence by scavenging reactive oxygen species in Arabidopsis thaliana. Journal of Biological Chemistry, 296.
  Google Scholar: Author Only Title Only Author and Title

Chauhan, H., Khurana, N., Agarwal, P. and Khurana, P., 2011. Heat shock factors in rice (Oryza sativa L.): genome-wide expression analysis during reproductive development and abiotic stress. Molecular Genetics and Genomics, 286(2), pp.171-187.
  Google Scholar: Author Only Title Only Author and Title

Chen, H., Liu, L., Wang, L., Wang, S. and Cheng, X., 2016. VrDREB2A, a DREB-binding transcription factor from Vigna radiata, increased drought and high-salt tolerance in transgenic Arabidopsis thaliana. Journal of plant research, 129(2), pp.263-273.
  Google Scholar: Author Only Title Only Author and Title

Chen, H.H., Qu, L., Xu, Z.H., Zhu, J.K. and Xue, H.W., 2018. EL1-like casein kinases suppress ABA signaling and responses by phosphorylating and destabilizing the ABA receptors PYR/PYLs in Arabidopsis. Molecular Plant, 11(5), pp.706-719.
  Google Scholar: Author Only Title Only Author and Title

Chen, H.H., Qu, L., Xu, Z.H., Zhu, J.K. and Xue, H.W., 2018. EL1-like casein kinases suppress ABA signaling and responses by phosphorylating and destabilizing the ABA receptors PYR/PYLs in Arabidopsis. Molecular Plant, 11(5), pp.706-719.
  Google Scholar: Author Only Title Only Author and Title

Chen, J., Ullah, C., Reichelt, M., Gershenzon, J. and Hammerbacher, A., 2019. Sclerotinia sclerotiorum circumvents flavonoid defenses by catabolizing flavonol glycosides and aglycones. Plant physiology, 180(4), pp.1975-1987.
  Google Scholar: Author Only Title Only Author and Title

Choudhury, F.K., Rivero, R.M., Blumwald, E. and Mittler, R., 2017. Reactive oxygen species, abiotic stress and stress combination. The Plant Journal, 90(5), pp.856-867.
  Google Scholar: Author Only Title Only Author and Title

Cramer, G.R., Urano, K., Delrot, S., Pezzotti, M. and Shinozaki, K., 2011. Effects of abiotic stress on plants: a systems biology perspective. BMC plant biology, 11(1), pp.1-14.
  Google Scholar: Author Only Title Only Author and Title

Crosby, K.C., Pietraszewska-Bogiel, A., Gadella Jr, T.W. and Winkel, B.S., 2011. Förster resonance energy transfer demonstrates a flavonoid metabolon in living plant cells that displays competitive interactions between enzymes. FEBS letters, 585(14), pp.2193-2198.
  Google Scholar: Author Only Title Only Author and Title

Czemmel, S., Stracke, R., Weisshaar, B., Cordon, N., Harris, N.N., Walker, A.R., Robinson, S.P. and Bogs, J., 2009. The grapevine

R2R3-MYB transcription factor VvMYBF1 regulates flavonol synthesis in developing grape berries. Plant physiology, 151(3), pp.1513-1530.
Google Scholar: Author Only Title Only Author and Title

D'hont, A, Denoeud, F., Aury, J.M., Baurens, F.C., Carreel, F., Garsmeur, O., Noel, B., Bocs, S., Droc, G., Rouard, M. and Da Silva, C., 2012. The banana (Musa acuminata) genome and the evolution of monocotyledonous plants. Nature, 488(7410), pp.213-217.

Dubouzet, J.G., Sakuma, Y., Ito, Y., Kasuga, M., Dubouzet, E.G., Miura, S., Seki, M., Shinozaki, K. and Yamaguchi-Shinozaki, K., 2003. OsDREB genes in rice, Oryza sativa L., encode transcription activators that function in drought-, high-salt-and cold-responsive gene expression. The Plant Journal, 33(4), pp.751-763.
Google Scholar: Author Only Title Only Author and Title

Friedrich, T., Oberkofler, V., Trindade, I., Altmann, S., Brzezinka, K., Lämke, J., Gorka, M., Kappel, C., Sokolowska, E., Skirycz, A. and Graf, A, 2021. Heteromeric HSFA2/HSFA3 complexes drive transcriptional memory after heat stress in Arabidopsis. Nature Communications, 12(1), pp.1-15.
Google Scholar: Author Only Title Only Author and Title

Fujii, H., Verslues, P.E. and Zhu, J.K., 2007. Identification of two protein kinases required for abscisic acid regulation of seed germination, root growth, and gene expression in Arabidopsis. The Plant Cell, 19(2), pp.485-494.
Google Scholar: Author Only Title Only Author and Title

Fujino, N., Tenma, N., Waki, T., Ito, K., Komatsuzaki, Y., Sugiyama, K., Yamazaki, T., Yoshida, S., Hatayama, M., Yamashita, S. and Tanaka, Y., 2018. Physical interactions among flavonoid enzymes in snapdragon and torenia reveal the diversity in the flavonoid metabolon organization of different plant species. The Plant Journal, 94(2), pp.372-392.
Google Scholar: Author Only Title Only Author and Title

Giri, J., Vij, S., Dansana, P.K. and Tyagi, A.K., 2011. Rice A20/AN1 zinc-finger containing stress-associated proteins (SAP1/11) and a receptor-like cytoplasmic kinase (OsRLCK253) interact via A20 zinc-finger and confer abiotic stress tolerance in transgenic Arabidopsis plants. New Phytologist, 191(3), pp.721-732.
Google Scholar: Author Only Title Only Author and Title

Gong, S., Ding, Y., Hu, S., Ding, L., Chen, Z. and Zhu, C., 2019. The role of HD-Zip class I transcription factors in plant response to abiotic stresses. Physiologia plantarum, 167(4), pp.516-525.
Google Scholar: Author Only Title Only Author and Title

Grunewald, W., De Smet, I., Lewis, D.R., Löfke, C., Jansen, L., Goeminne, G., Vanden Bossche, R., Karimi, M., De Rybel, B., Vanholme, B. and Teichmann, T., 2012. Transcription factor WRKY23 assists auxin distribution patterns during Arabidopsis root development through local control on flavonol biosynthesis. Proceedings of the National Academy of Sciences, 109(5), pp.1554-1559.
Google Scholar: Author Only Title Only Author and Title

Guo, L., Gao, L., Ma, X., Guo, F., Ruan, H., Bao, Y., Xia, T. and Wang, Y., 2019. Functional analysis of flavonoid 3′-hydroxylase and flavonoid 3′, 5′-hydroxylases from tea plant (Camellia sinensis), involved in the B-ring hydroxylation of flavonoids. Gene, 717, p.144046.
Google Scholar: Author Only Title Only Author and Title

Guo, M., Liu, J.H., Ma, X., Luo, D.X., Gong, Z.H. and Lu, M.H., 2016. The plant heat stress transcription factors (HSFs): structure, regulation, and function in response to abiotic stresses. Frontiers in plant science, 7, p.114.
Google Scholar: Author Only Title Only Author and Title

Halbwirth, H., 2010. The creation and physiological relevance of divergent hydroxylation patterns in the flavonoid pathway. International journal of molecular sciences, 11(2), pp.595-621.
Google Scholar: Author Only Title Only Author and Title

He, Y., Zhang, X., Li, L., Sun, Z, Li, J., Chen, X. and Hong, G., 2021. SPX4 interacts with both PHR1 and PAP1 to regulate critical steps in phosphorus-status-dependent anthocyanin biosynthesis. New Phytologist, 230(1), pp.205-217.
Google Scholar: Author Only Title Only Author and Title

Huang, Y.C., Niu, C.Y., Yang, C.R. and Jinn, T.L., 2016. The heat stress factor HSFA6b connects ABA signaling and ABA-mediated heat responses. Plant physiology, 172(2), pp.1182-1199.
Google Scholar: Author Only Title Only Author and Title

Huang, Y.C., Niu, C.Y., Yang, C.R. and Jinn, T.L., 2016. The heat stress factor HSFA6b connects ABA signaling and ABA-mediated heat responses. Plant physiology, 172(2), pp.1182-1199.
Google Scholar: Author Only Title Only Author and Title

Jiang, X., Shi, Y., Fu, Z, Li, W.W., Lai, S., Wu, Y., Wang, Y., Liu, Y., Gao, L. and Xia, T., 2020. Functional characterization of three flavonol synthase genes from Camellia sinensis: Roles in flavonol accumulation. Plant Science, 300, p.110632.
Google Scholar: Author Only Title Only Author and Title

Kalaji, H.M., Dąbrowski, P., Cetner, M.D., Samborska, I.A., Łukasik, I., Brestic, M., Zivcak, M., Tomasz, H., Mojski, J., Kociel, H. and Panchal, B.M., 2017. A comparison between different chlorophyll content meters under nutrient deficiency conditions. Journal of Plant Nutrition, 40(7), pp.1024-1034.
  Google Scholar: Author Only Title Only Author and Title

Katoh, K. and Standley, D.M., 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. Molecular biology and evolution, 30(4), pp.772-780.
  Google Scholar: Author Only Title Only Author and Title

Kitamura, S., Oono, Y. and Narumi, I., 2016. Arabidopsis pab1, a mutant with reduced anthocyanins in immature seeds from banyuls, harbors a mutation in the MATE transporter FFT. Plant molecular biology, 90(1), pp.7-18.
  Google Scholar: Author Only Title Only Author and Title

Ko, M.J., Cheigh, C.I. and Chung, M.S., 2014. Relationship analysis between flavonoids structure and subcritical water extraction (SWE). Food chemistry, 143, pp.147-155.
  Google Scholar: Author Only Title Only Author and Title

Koncz, C. and Schell, J., 1986. The promoter of TL-DNA gene 5 controls the tissue-specific expression of chimaeric genes carried by a novel type of Agrobacterium binary vector. Molecular and General Genetics MGG, 204(3), pp.383-396.
  Google Scholar: Author Only Title Only Author and Title

Kuang, J.F., Chen, J.Y., Liu, X.C., Han, Y.C., Xiao, Y.Y., Shan, W., Tang, Y., Wu, K.Q., He, J.X. and Lu, W.J., 2017. The transcriptional regulatory network mediated by banana (Musa acuminata) dehydration-responsive element binding (MaDREB) transcription factors in fruit ripening. New Phytologist, 214(2), pp.762-781.
  Google Scholar: Author Only Title Only Author and Title

Kumar, K., Purayannur, S., Kaladhar, V.C., Parida, S.K. and Verma, P.K., 2018. mQTL-seq and classical mapping implicates the role of an AT-HOOK MOTIF CONTAINING NUCLEAR LOCALIZED (AHL) family gene in Ascochyta blight resistance of chickpea. Plant, Cell & Environment, 41(9), pp.2128-2140.
  Google Scholar: Author Only Title Only Author and Title

Kumar, S., Stecher, G., Li, M., Knyaz, C. and Tamura, K., 2018. MEGA X: molecular evolutionary genetics analysis across computing platforms. Molecular biology and evolution, 35(6), p.1547.
  Google Scholar: Author Only Title Only Author and Title

Lago, J.H.G., Toledo-Arruda, A.C., Mernak, M., Barrosa, K.H., Martins, M.A., Tibério, I.F. and Prado, C.M., 2014. Structure-activity association of flavonoids in lung diseases. Molecules, 19(3), pp.3570-3595.
  Google Scholar: Author Only Title Only Author and Title

Lee, S.J., Kang, J.Y., Park, H.J., Kim, M.D., Bae, M.S., Choi, H.I. and Kim, S.Y., 2010. DREB2C interacts with ABF2, a bZIP protein regulating abscisic acid-responsive gene expression, and its overexpression affects abscisic acid sensitivity. Plant physiology, 153(2), pp.716-727.
  Google Scholar: Author Only Title Only Author and Title

Lewis, D.R., Ramirez, M.V., Miller, N.D., Vallabhaneni, P., Ray, W.K., Helm, R.F., Winkel, B.S. and Muday, G.K., 2011. Auxin and ethylene induce flavonol accumulation through distinct transcriptional networks. Plant physiology, 156(1), pp.144-164.
  Google Scholar: Author Only Title Only Author and Title

Li, L., Peng, H., Tan, S., Zhou, J., Fang, Z., Hu, Z, Gao, L., Li, T., Zhang, W. and Chen, L., 2020. Effects of early cold stress on gene expression in Chlamydomonas reinhardtii. Genomics, 112(2), pp.1128-1138.
  Google Scholar: Author Only Title Only Author and Title

Liang, T., Shi, C., Peng, Y., Tan, H., Xin, P., Yang, Y., Wang, F., Li, X., Chu, J., Huang, J. and Yin, Y., 2020. Brassinosteroid-activated BRI1-EMS-SUPPRESSOR 1 inhibits flavonoid biosynthesis and coordinates growth and UV-B stress responses in plants. Plant Cell, 32(10), pp.3224-3239.
  Google Scholar: Author Only Title Only Author and Title

Lin, Z., Li, Y., Zhang, Z., Liu, X., Hsu, C.C., Du, Y., Sang, T., Zhu, C., Wang, Y., Satheesh, V. and Pratibha, P., 2020. A RAF-SnRK2 kinase cascade mediates early osmotic stress signaling in higher plants. Nature communications, 11(1), pp.1-10.
  Google Scholar: Author Only Title Only Author and Title

Liu, H.T., Gao, F., Li, G.L., Han, J.L., Liu, D.L., Sun, D.Y. and Zhou, R.G., 2008. The calmodulin-binding protein kinase 3 is part of heat-shock signal transduction in Arabidopsis thaliana. The Plant Journal, 55(5), pp.760-773.
  Google Scholar: Author Only Title Only Author and Title

López, M.A., Bannenberg, G. and Castresana, C., 2008. Controlling hormone signaling is a plant and pathogen challenge for growth and survival. Current opinion in plant biology, 11(4), pp.420-427.
  Google Scholar: Author Only Title Only Author and Title

Lu, Q., Tang, X., Tian, G., Wang, F., Liu, K., Nguyen, V., Kohalmi, S.E., Keller, W.A., Tsang, E.W., Harada, J.J. and Rothstein, S.J.,

2010. Arabidopsis homolog of the yeast TREX-2 mRNA export complex: components and anchoring nucleoporin. The Plant Journal, 61(2), pp.259-270.
  Google Scholar: Author Only Title Only Author and Title

Ma, D., Tang, H., Reichelt, M., Piirtola, E.M., Salminen, J.P., Gershenzon, J. and Constabel, C.P., 2021. Poplar MYB117 promotes anthocyanin synthesis and enhances flavonoid B-ring hydroxylation by up-regulating the flavonoid 3′, 5′-hydroxylase gene. Journal of Experimental Botany, 72(10), pp.3864-3880.
  Google Scholar: Author Only Title Only Author and Title

Ma, Y., Szostkiewicz, I., Korte, A., Moes, D., Yang, Y., Christmann, A. and Grill, E., 2009. Regulators of PP2C phosphatase activity function as abscisic acid sensors. Science, 324(5930), pp.1064-1068.
  Google Scholar: Author Only Title Only Author and Title

Maloney, G.S., DiNapoli, K.T. and Muday, G.K., 2014. The anthocyanin reduced tomato mutant demonstrates the role of flavonols in tomato lateral root and root hair development. Plant Physiology, 166(2), pp.614-631.
  Google Scholar: Author Only Title Only Author and Title

Mann DG, LaFayette PR, Abercrombie LL, King ZR, Mazarei M, Halter MC, Poovaiah CR, Baxter H, Shen H, Dixon RA, Parrott WA. Gateway-compatible vectors for high-throughput gene functional analysis in switchgrass (Panicum virgatum L.) and other monocot species. Plant biotechnology journal, 10(2), pp. 226-36.

Manna, M., Thakur, T., Chirom, O., Mandlik, R., Deshmukh, R. and Salvi, P., 2021. Transcription factors as key molecular target to strengthen the drought stress tolerance in plants. Physiologia Plantarum, 172(2), pp.847-868.
  Google Scholar: Author Only Title Only Author and Title

Martin, K., Kopperud, K., Chakrabarty, R., Banerjee, R., Brooks, R. and Goodin, M.M., 2009. Transient expression in Nicotiana benthamiana fluorescent marker lines provides enhanced definition of protein localization, movement and interactions in planta. The Plant Journal, 59(1), pp.150-162.
  Google Scholar: Author Only Title Only Author and Title

Matus, J.T., Loyola, R., Vega, A., Peña-Neira, A., Bordeu, E., Arce-Johnson, P. and Alcalde, J.A., 2009. Post-veraison sunlight exposure induces MYB-mediated transcriptional regulation of anthocyanin and flavonol synthesis in berry skins of Vitis vinifera. Journal of experimental botany, 60(3), pp.853-867.
  Google Scholar: Author Only Title Only Author and Title

Medina-Puche, L., Cumplido-Laso, G., Amil-Ruiz, F., Hoffmann, T., Ring, L., Rodríguez-Franco, A., Caballero, J.L., Schwab, W., Muñoz-Blanco, J. and Blanco-Portales, R., 2014. MYB10 plays a major role in the regulation of flavonoid/phenylpropanoid metabolism during ripening of Fragaria× ananassa fruits. Journal of experimental botany, 65(2), pp.401-417.
  Google Scholar: Author Only Title Only Author and Title

Mehra, P., Pandey, B.K., Verma, L. and Giri, J., 2019. A novel glycerophosphodiester phosphodiesterase improves phosphate deficiency tolerance in rice. Plant, Cell & Environment, 42(4), pp.1167-1179.
  Google Scholar: Author Only Title Only Author and Title

Mehrtens, F., Kranz, H., Bednarek, P. and Weisshaar, B., 2005. The Arabidopsis transcription factor MYB12 is a flavonol-specific regulator of phenylpropanoid biosynthesis. Plant physiology, 138(2), pp.1083-1096.
  Google Scholar: Author Only Title Only Author and Title

Mikkelsen, B.L., Olsen, C.E. and Lyngkjær, M.F., 2015. Accumulation of secondary metabolites in healthy and diseased barley, grown under future climate levels of CO2, ozone and temperature. Phytochemistry, 118, pp.162-173.
  Google Scholar: Author Only Title Only Author and Title

Misra, P., Pandey, A., Tiwari, M., Chandrashekar, K., Sidhu, O.P., Asif, M.H., Chakrabarty, D., Singh, P.K., Trivedi, P.K., Nath, P. and Tuli, R., 2010. Modulation of transcriptome and metabolome of tobacco by Arabidopsis transcription factor, AtMYB12, leads to insect resistance. Plant Physiology, 152(4), pp.2258-2268.
  Google Scholar: Author Only Title Only Author and Title

Muhlemann, J.K., Younts, T.L. and Muday, G.K., 2018. Flavonols control pollen tube growth and integrity by regulating ROS homeostasis during high-temperature stress. Proceedings of the National Academy of Sciences, 115(47), pp.E11188-E11197.
  Google Scholar: Author Only Title Only Author and Title

Murcia, G., Fontana, A., Pontin, M., Baraldi, R., Bertazza, G. and Piccoli, P.N., 2017. ABA and GA3 regulate the synthesis of primary and secondary metabolites related to alleviation from biotic and abiotic stresses in grapevine. Phytochemistry, 135, pp.34-52.
  Google Scholar: Author Only Title Only Author and Title

Muthusamy, M., Uma, S., Backiyarani, S., Saraswathi, M.S. and Chandrasekar, A., 2016. Transcriptomic changes of drought-tolerant and sensitive banana cultivars exposed to drought stress. Frontiers in plant science, 7, pp.1609.
  Google Scholar: Author Only Title Only Author and Title

Nagamatsu, A., Masuta, C., Senda, M., Matsuura, H., Kasai, A., Hong, J.S., Kitamura, K., Abe, J. and Kanazawa, A., 2007. Functional

analysis of soybean genes involved in flavonoid biosynthesis by virus-induced gene silencing. Plant biotechnology journal, 5(6), pp.778-790.
Google Scholar: Author Only Title Only Author and Title

Naik, J., Rajput, R., Pucker, B., Stracke, R. and Pandey, A, 2021. The R2R3-MYB transcription factor MtMYB134 orchestrates flavonol biosynthesis in Medicago truncatula. Plant Molecular Biology, 106(1), pp.157-172.
Google Scholar: Author Only Title Only Author and Title

Née, G., Kramer, K., Nakabayashi, K., Yuan, B., Xiang, Y., Miatton, E., Finkemeier, I. and Soppe, W.J., 2017. DELAY OF GERMINATION1 requires PP2C phosphatases of the ABA signalling pathway to control seed dormancy. Nature communications, 8(1), pp.1-9.
Google Scholar: Author Only Title Only Author and Title

Oh, M.M., Trick, H.N. and Rajashekar, C.B., 2009. Secondary metabolism and antioxidants are involved in environmental adaptation and stress tolerance in lettuce. Journal of plant physiology, 166(2), pp.180-191.
Google Scholar: Author Only Title Only Author and Title

Pandey, A, Alok, A, Lakhwani, D., Singh, J., Asif, M.H. and Trivedi, P.K., 2016. Genome-wide expression analysis and metabolite profiling elucidate transcriptional regulation of flavonoid biosynthesis and modulation under abiotic stresses in banana. Scientific reports, 6(1), pp.1-13.
Google Scholar: Author Only Title Only Author and Title

Pandey, A, Misra, P. and Trivedi, P.K., 2015. Constitutive expression of Arabidopsis MYB transcription factor, AtMYB11, in tobacco modulates flavonoid biosynthesis in favor of flavonol accumulation. Plant cell reports, 34(9), pp.1515-1528.
Google Scholar: Author Only Title Only Author and Title

Pandey, A, Misra, P., Choudhary, D., Yadav, R., Goel, R., Bhambhani, S., Sanyal, I., Trivedi, R. and Kumar Trivedi, P., 2015. AtMYB12 expression in tomato leads to large scale differential modulation in transcriptome and flavonoid content in leaf and fruit tissues. Scientific reports, 5(1), pp.1-14.
Google Scholar: Author Only Title Only Author and Title

Pandey, A, Misra, P., Khan, M.P., Swarnkar, G., Tewari, M.C., Bhambhani, S., Trivedi, R., Chattopadhyay, N. and Trivedi, P.K., 2014. Co-expression of Arabidopsis transcription factor, At MYB 12, and soybean isoflavone synthase, GmIFS 1, genes in tobacco leads to enhanced biosynthesis of isoflavones and flavonols resulting in osteoprotective activity. Plant biotechnology journal, 12(1), pp.69-80.
Google Scholar: Author Only Title Only Author and Title

Peng, Y., Thrimawithana, A.H., Cooney, J.M., Jensen, D.J., Espley, R.V. and Allan, A.C., 2020. The proanthocyanin-related transcription factors MYBC1 and WRKY44 regulate branch points in the kiwifruit anthocyanin pathway. Scientific Reports, 10(1), pp.1-19.
Google Scholar: Author Only Title Only Author and Title

Pucker, B., Pandey, A, WeissShaar, B. and Stracke, R., 2020. The R2R3-MYB gene family in banana (Musa acuminata): Genome-wide identification, classification and expression patterns. PloS one, 15(10), pp.e0239275.
Google Scholar: Author Only Title Only Author and Title

Qian, C., Chen, Z, Liu, Q., Mao, W., Chen, Y., Tian, W., Liu, Y., Han, J., Ouyang, X. and Huang, X., 2020. Coordinated transcriptional regulation by the UV-B photoreceptor and multiple transcription factors for plant UV-B responses. Molecular plant, 13(5), pp.777-792.
Google Scholar: Author Only Title Only Author and Title

Rajput, R., Naik, J., Stracke, R. and Pandey, A, 2022. Interplay between R2R3 MYB-type activators and repressors regulates proanthocyanidin biosynthesis in banana (Musa acuminata). New Phytologist, 236, pp.1108-1127.
Google Scholar: Author Only Title Only Author and Title

Rajput, R., Tyagi, S., Naik, J., Pucker, B., Stracke, R. and Pandey, A, 2022. The R2R3-MYB gene family in Cicer arietinum: genome-wide identification and expression analysis leads to functional characterization of proanthocyanidin biosynthesis regulators in the seed coat. Planta, 256(4), pp.1-24.
Google Scholar: Author Only Title Only Author and Title

Scharf, K.D., Berberich, T., Ebersberger, I. and Nover, L., 2012. The plant heat stress transcription factor (Hsf) family: structure, function and evolution. Biochimica et Biophysica Acta (BBA)-Gene Regulatory Mechanisms, 1819(2), pp.104-119.
Google Scholar: Author Only Title Only Author and Title

Schramm, F., Larkindale, J., Kiehlmann, E., Ganguli, A, Englich, G., Vierling, E. and Von Koskull-Döring, P., 2008. A cascade of transcription factor DREB2A and heat stress transcription factor HsfA3 regulates the heat stress response of Arabidopsis. The Plant Journal, 53(2), pp.264-274.
Google Scholar: Author Only Title Only Author and Title

Schwinn, K.E., Ngo, H., Kenel, F., Brummell, D.A, Albert, N.W., McCallum, J.A, Pither-Joyce, M., Crowhurst, R.N., Eady, C. and

Davies, K.M., 2016. The onion (Allium cepa L.) R2R3-MYB gene MYB1 regulates anthocyanin biosynthesis. Frontiers in Plant Science, 7, p.1865.

Google Scholar: Author Only Title Only Author and Title

Shan, W., KUANG, J.F., LU, W.J. and CHEN, J.Y., 2014. Banana fruit NAC transcription factor MaNAC 1 is a direct target of MaICE 1 and involved in cold stress through interacting with MaCBF 1. Plant, cell & environment, 37(9), pp.2116-2127.

Google Scholar: Author Only Title Only Author and Title

Shan, X., Li, Y., Yang, S., Yang, Z., Qiu, M., Gao, R., Han, T., Meng, X., Xu, Z., Wang, L. and Gao, X., 2020. The spatio-temporal biosynthesis of floral flavonols is controlled by differential phylogenetic MYB regulators in Freesia hybrida. New Phytologist, 228(6), pp.1864-1879.

Google Scholar: Author Only Title Only Author and Title

Shi, C., Luo, P., Du, Y.T., Chen, H., Huang, X., Cheng, T.H., Luo, A., Li, H.J., Yang, W.C., Zhao, P. and Sun, M.X., 2019. Maternal control of suspensor programmed cell death via gibberellin signaling. Nature communications, 10(1), pp.1-12.

Google Scholar: Author Only Title Only Author and Title

Shih, C.H., Chu, H., Tang, L.K., Sakamoto, W., Maekawa, M., Chu, I.K., Wang, M. and Lo, C., 2008. Functional characterization of key structural genes in rice flavonoid biosynthesis. Planta, 228(6), pp.1043-1054.

Google Scholar: Author Only Title Only Author and Title

Si, J., Ye, B.B., Liu, Z.L., Xiao, X.M., Yang, Y.Y., Fan, Z.Q., Shan, W., Kuang, J.F., Lu, W.J., Su, X.G. and Chen, J.Y., 2022. Transcriptional repression of MaRBOHs by MaHsf26 is associated with heat shock-alleviated chilling injury in banana fruit. Postharvest Biology and Technology, 193, p.112056.

Google Scholar: Author Only Title Only Author and Title

Song, Z., Lai, X., Chen, H., Wang, L., Pang, X., Hao, Y., Lu, W., Chen, W., Zhu, X. and Li, X., 2022. Role of MaABI5-like in abscisic acid-induced cold tolerance of 'Fenjiao'banana fruit. Horticulture Research, 9, p.uhac130.

Google Scholar: Author Only Title Only Author and Title

Stracke, R., Favory, J.J., Gruber, H., Bartelniewoehner, L., Bartels, S., Binkert, M., Funk, M., Weisshaar, B. and Ulm, R., 2010. The Arabidopsis bZIP transcription factor HY5 regulates expression of the PFG1/MYB12 gene in response to light and ultraviolet-B radiation. Plant, cell & environment, 33(1), pp.88-103.

Google Scholar: Author Only Title Only Author and Title

Stracke, R., FAVORY, J.J., Gruber, H., Bartelniewoehner, L., Bartels, S., Binkert, M., Funk, M., Weisshaar, B. and Ulm, R., 2010. The Arabidopsis bZIP transcription factor HY5 regulates expression of the PFG1/MYB12 gene in response to light and ultraviolet-B radiation. Plant, cell & environment, 33(1), pp.88-103.

Google Scholar: Author Only Title Only Author and Title

Stracke, R., Holtgräwe, D., Schneider, J., Pucker, B., Rosleff Sörensen, T. and Weisshaar, B., 2014. Genome-wide identification and characterisation of R2R3-MYB genes in sugar beet (Beta vulgaris). BMC plant biology, 14(1), pp.1-17.

Google Scholar: Author Only Title Only Author and Title

Stracke, R., Jahns, O., Keck, M., Tohge, T., Niehaus, K., Fernie, A.R. and Weisshaar, B., 2010. Analysis of PRODUCTION OF FLAVONOL GLYCOSIDES-dependent flavonol glycoside accumulation in Arabidopsis thaliana plants reveals MYB11-, MYB12- and MYB111-independent flavonol glycoside accumulation. New Phytologist, 188(4), pp.985-1000.

Google Scholar: Author Only Title Only Author and Title

Stracke, R., Werber, M. and Weisshaar, B., 2001. The R2R3-MYB gene family in Arabidopsis thaliana. Current opinion in plant biology, 4(5), pp.447-456.

Google Scholar: Author Only Title Only Author and Title

Tan, H., Man, C., Xie, Y., Yan, J., Chu, J. and Huang, J., 2019. A crucial role of GA-regulated flavonol biosynthesis in root growth of Arabidopsis. Molecular plant, 12(4), pp.521-537.

Google Scholar: Author Only Title Only Author and Title

Teale, W.D., Pasternak, T., Dal Bosco, C., Dovzhenko, A., Kratzat, K., Bildl, W., Schwörer, M., Falk, T., Ruperti, B., V Schaefer, J. and Shahriari, M., 2021. Flavonol-mediated stabilization of PIN efflux complexes regulates polar auxin transport. The EMBO Journal, 40(1), p.e104416.

Google Scholar: Author Only Title Only Author and Title

Umezawa, T., Sugiyama, N., Mizoguchi, M., Hayashi, S., Myouga, F., Yamaguchi-Shinozaki, K., Ishihama, Y., Hirayama, T. and Shinozaki, K., 2009. Type 2C protein phosphatases directly regulate abscisic acid-activated protein kinases in Arabidopsis. Proceedings of the National Academy of sciences, 106(41), pp.17588-17593.

Google Scholar: Author Only Title Only Author and Title

Upendra, K., Shekhawat, S., Srinivas, L. and Ganapathi, T.R., 2011. MusaDHN-1, a novel multiple stress-inducible SK^ sub 3^-type dehydrin gene, contributes affirmatively to drought-and salt-stress tolerance in banana. Planta, 234(5), p.915.

Google Scholar: Author Only Title Only Author and Title

Waki, T., Yoo, D., Fujino, N., Mameda, R., Denessiouk, K., Yamashita, S., Motohashi, R., Akashi, T., Aoki, T., Ayabe, S.I. and Takahashi, S., 2016. Identification of protein–protein interactions of isoflavonoid biosynthetic enzymes with 2-hydroxyisoflavanone synthase in soybean (Glycine max (L.) Merr.). Biochemical and biophysical research communications, 469(3), pp.546-551.
    Google Scholar: Author Only Title Only Author and Title

Walter, M., Chaban, C., Schütze, K., Batistic, O., Weckermann, K., Näke, C., Blazevic, D., Grefen, C., Schumacher, K., Oecking, C. and Harter, K., 2004. Visualization of protein interactions in living plant cells using bimolecular fluorescence complementation. The Plant Journal, 40(3), pp.428-438.
    Google Scholar: Author Only Title Only Author and Title

Wang, J., Ren, Y., Liu, X., Luo, S., Zhang, X., Liu, X., Lin, Q., Zhu, S., Wan, H., Yang, Y. and Zhang, Y., 2021. Transcriptional activation and phosphorylation of OsCNGC9 confer enhanced chilling tolerance in rice. Molecular plant, 14(2), pp.315-329.
    Google Scholar: Author Only Title Only Author and Title

Wang, N., Liu, W., Yu, L., Guo, Z., Chen, Z., Jiang, S., Xu, H., Fang, H., Wang, Y., Zhang, Z. and Chen, X., 2020. HEAT SHOCK FACTOR A8a modulates flavonoid synthesis and drought tolerance. Plant Physiology, 184(3), pp.1273-1290.
    Google Scholar: Author Only Title Only Author and Title

Wang, N., Xu, H., Jiang, S., Zhang, Z., Lu, N., Qiu, H., Qu, C., Wang, Y., Wu, S. and Chen, X., 2017. MYB12 and MYB22 play essential roles in proanthocyanidin and flavonol synthesis in red-fleshed apple (Malus sieversii f. niedzwetzkyana). The Plant Journal, 90(2), pp.276-292.
    Google Scholar: Author Only Title Only Author and Title

Wang, X., Wang, L., Wang, Y., Liu, H., Hu, D., Zhang, N., Zhang, S., Cao, H., Cao, Q., Zhang, Z. and Tang, S., 2018. Arabidopsis PCaP2 plays an important role in chilling tolerance and ABA response by activating CBF-and SnRK2-mediated transcriptional regulatory network. Frontiers in plant science, 9, p.215.
    Google Scholar: Author Only Title Only Author and Title

Wang, X., Yu, R. and Li, J., 2021. Using genetic engineering techniques to develop banana cultivars with fusarium wilt resistance and ideal plant architecture. Frontiers in Plant Science, 11, p.617528.
    Google Scholar: Author Only Title Only Author and Title

Wang, Z., Huang, S., Jia, C., Liu, J., Zhang, J., Xu, B. and Jin, Z., 2013. Molecular cloning and expression of five glutathione S-transferase (GST) genes from Banana (Musa acuminata L. AAA group, cv. Cavendish). Plant cell reports, 32(9), pp.1373-1380.
    Google Scholar: Author Only Title Only Author and Title

Watkins, J.M., Chapman, J.M. and Muday, G.K., 2017. Abscisic acid-induced reactive oxygen species are modulated by flavonols to control stomata aperture. Plant Physiology, 175(4), pp.1807-1825.
    Google Scholar: Author Only Title Only Author and Title

Wei, Y., Hu, W., Xia, F., Zeng, H., Li, X., Yan, Y., He, C. and Shi, H., 2016. Heat shock transcription factors in banana: genome-wide characterization and expression profile analysis during development and stress response. Scientific Reports, 6(1), pp.1-11.
    Google Scholar: Author Only Title Only Author and Title

Wu, T.Y., Hoh, K.L., Boonyaves, K., Krishnamoorthi, S. and Urano, D., 2022. Diversification of heat shock transcription factors expanded thermal stress responses during early plant evolution. The Plant Cell, 34(10), pp.3557-3576.
    Google Scholar: Author Only Title Only Author and Title

Xu, F., Li, L., Zhang, W., Cheng, H., Sun, N., Cheng, S. and Wang, Y., 2012. Isolation, characterization, and function analysis of a flavonol synthase gene from Ginkgo biloba. Molecular Biology Reports, 39(3), pp.2285-2296.
    Google Scholar: Author Only Title Only Author and Title

Xu, Y., Hu, W., Liu, J., Song, S., Hou, X., Jia, C., Li, J., Miao, H., Wang, Z., Tie, W. and Xu, B., 2020. An aquaporin gene MaPIP2-7 is involved in tolerance to drought, cold and salt stresses in transgenic banana (Musa acuminata L.). Plant Physiology and Biochemistry, 147, pp.66-76.
    Google Scholar: Author Only Title Only Author and Title

Yang, M., Han, X., Yang, J., Jiang, Y. and Hu, Y., 2021. The Arabidopsis circadian clock protein PRR5 interacts with and stimulates ABI5 to modulate abscisic acid signaling during seed germination. The Plant Cell, 33(9), pp.3022-3041.
    Google Scholar: Author Only Title Only Author and Title

Yang, Q., Song, Z., Dong, B., Niu, L., Cao, H., Li, H., Du, T., Liu, T., Yang, W., Meng, D. and Fu, Y., 2021. Hyperoside regulates its own biosynthesis via MYB30 in promoting reproductive development and seed set in okra. Plant Physiology, 185(3), pp.951-968.
    Google Scholar: Author Only Title Only Author and Title

Yoshida, T., Fujita, Y., Maruyama, K., Mogami, J., Todaka, D., Shinozaki, K. and Yamaguchi-Shinozaki, K.A.Z.U.K.O., 2015. Four A rabidopsis AREB/ABF transcription factors function predominantly in gene expression downstream of SnRK2 kinases in abscisic acid signalling in response to osmotic stress. Plant, cell & environment, 38(1), pp.35-49.
    Google Scholar: Author Only Title Only Author and Title

Yoshida, T., Ohama, N., Nakajima, J., Kidokoro, S., Mizoi, J., Nakashima, K., Maruyama, K., Kim, J.M., Seki, M., Todaka, D. and Osakabe, Y., 2011. Arabidopsis HsfA1 transcription factors function as the main positive regulators in heat shock-responsive gene expression. Molecular Genetics and Genomics, 286(5), pp.321-332.

Google Scholar: Author Only Title Only Author and Title

Zhang, J., Subramanian, S., Stacey, G. and Yu, O., 2009. Flavones and flavonols play distinct critical roles during nodulation of Medicago truncatula by Sinorhizobium meliloti. The Plant Journal, 57(1), pp.171-183.

Google Scholar: Author Only Title Only Author and Title

Zhang, K., Logacheva, M.D., Meng, Y., Hu, J., Wan, D., Li, L., Janovská, D., Wang, Z., Georgiev, M.I., Yu, Z. and Yang, F., 2018. Jasmonate-responsive MYB factors spatially repress rutin biosynthesis in Fagopyrum tataricum. Journal of Experimental Botany, 69(8), pp.1955-1966.

Google Scholar: Author Only Title Only Author and Title

Zhang, L., Cenci, A., Rouard, M., Zhang, D., Wang, Y., Tang, W. and Zheng, S.J., 2019. Transcriptomic analysis of resistant and susceptible banana corms in response to infection by Fusarium oxysporum f. sp. cubense tropical race 4. Scientific reports, 9(1), pp.1-14.

Google Scholar: Author Only Title Only Author and Title

Zhang, X., He, Y., Li, L., Liu, H. and Hong, G., 2021. Involvement of the R2R3-MYB transcription factor MYB21 and its homologs in regulating flavonol accumulation in Arabidopsis stamen. Journal of experimental botany, 72(12), pp.4319-4332.

Google Scholar: Author Only Title Only Author and Title

ZHAO, M.L., WANG, J.N., Shan, W., FAN, J.G., KUANG, J.F., WU, K.Q., LI, X.P., CHEN, W.X., HE, F.Y., CHEN, J.Y. and LU, W.J., 2013. Induction of jasmonate signalling regulators MaMYC2s and their physical interactions with MaICE1 in methyl jasmonate-induced chilling tolerance in banana fruit. Plant, cell & environment, 36(1), pp.30-51.

Google Scholar: Author Only Title Only Author and Title

Zhao, C., Liu, X., Gong, Q., Cao, J., Shen, W., Yin, X., Grierson, D., Zhang, B., Xu, C., Li, X. and Chen, K., 2021. Three AP2/ERF family members modulate flavonoid synthesis by regulating type IV chalcone isomerase in citrus. Plant biotechnology journal, 19(4), pp.671-688.

Google Scholar: Author Only Title Only Author and Title

Zhao, L., Hu, Y., Chong, K. and Wang, T., 2010. ARAG1, an ABA-responsive DREB gene, plays a role in seed germination and drought tolerance of rice. Annals of Botany, 105(3), pp.401-409.

Google Scholar: Author Only Title Only Author and Title

Zimmermann, I.M., Heim, M.A, Weisshaar, B. and Uhrig, J.F., 2004. Comprehensive identification of Arabidopsis thaliana MYB transcription factors interacting with R/B-like BHLH proteins. The Plant Journal, 40(1), pp.22-34.

Google Scholar: Author Only Title Only Author and Title

Zinsmeister, J., Berriri, S., Basso, D.P., Ly-Vu, B., Dang, T.T., Lalanne, D., da Silva, E.A.A., Leprince, O. and Buitink, J., 2020. The seed-specific heat shock factor A9 regulates the depth of dormancy in Medicago truncatula seeds via ABA signalling. Plant, Cell & Environment, 43(10), pp.2508-2522.

Google Scholar: Author Only Title Only Author and Title