Title: Predictive evolutionary modelling for influenza virus by site-based
dynamics of mutations

Authors: Jingzhi Lou[1,2]†, Weiwen Liang[3]†, Lirong Cao[1,4]†, Shi Zhao[1,4], Zigui Chen[5], Renee
Wan Yi Chan[6,7], Peter Pak Hang Cheung[8], Hong Zheng[1], Caiqi Liu[1], Qi Li[1], Marc Ka Chun
Chong[1,4], Yexian Zhang[1,4], Eng-kiong Yeoh[1,9], Paul Kay-Sheung Chan[5,10], Benny Chung Ying
Zee[1,4], Chris Ka Pun Mok[1,11]*, Maggie Haitian Wang[1,4]*

Affiliations:

[1] JC School of Public Health and Primary Care (JCSPHPC), the Chinese University of Hong
Kong (CUHK); Hong Kong SAR, China.

[2] Beth Bioinformatics Co. Ltd; Hong Kong SAR, China.

[3] HKU-Pasteur Research Pole, School of Public Health, Li Ka Shing Faculty of Medicine,
The University of Hong Kong; Hong Kong SAR, China.

[4] CUHK Shenzhen Research Institute, Shenzhen, China.

[5] Department of Microbiology, CUHK; Hong Kong SAR, China.

[6] Department of Paediatrics, CUHK; Hong Kong SAR, China.

[7] Hong Kong Hub of Paediatric Excellence, CUHK; Hong Kong SAR, China.

[8] Department of Chemical Pathology, CUHK; Hong Kong SAR, China.

[9] Centre for Health Systems and Policy Research, CUHK; Hong Kong SAR, China.

[10] Stanley Ho Centre for Emerging Infectious Diseases, CUHK; Hong Kong SAR, China.

[11] Li Ka Shing Institute of Health Sciences, Faculty of Medicine, CUHK, Hong Kong SAR,
China.

† These authors contributed equally to this work.

* Corresponding authors. Emails: kapunmok@cuhk.edu.hk, maggiew@cuhk.edu.hk

Abstract: A predictive evolutionary model was developed to forecast representative influenza viral strains for upcoming epidemic seasons. Influenza virus continuously evolves to escape human adaptive immunity and generates seasonal epidemics. A computational approach beth-1 that models site-wise mutation dynamics demonstrated remarkable matching of predicted strains to the circulating viruses in subsequent seasons for the influenza A(H1N1)pdm09 and A(H3N2) viruses in both retrospective and prospective validations. The method offers a promising and ready-to-use tool to facilitate vaccine strain selection for the influenza virus, achieved by capturing heterogeneous evolutionary dynamics over genome space-time and linking molecular variants to population immune response.

One-Sentence Summary: A computational model predicts virus evolution and facilitates vaccine strain selection for the influenza virus

Main Text:

The major driver of recurrent influenza epidemics is fast virus evolution that enables the influenza virus to escape from human immunity acquired from prior vaccination or infection[1]. In response, influenza vaccines need to be updated annually to match the circulating virus population[2]. Prediction of virus evolution has a critical role in ensuring the protective effect of influenza vaccines, which aids the selection of candidate vaccine strains nearly a year ahead before the arrival of next epidemic season[2,3]. We present a computational approach to predict influenza virus evolution through modelling the dynamic process of mutation adaptation at individual sites and locating the optimal wild-type strains for vaccine candidates by a combined evaluation of multiple gene segments and population immune profiles.

Virus evolution is shaped by a complex interplay of genetic mutations, host immune response, and epidemiology[1,4]. Thus, although mutation is stochastic[5,6], the evolutionary process could be traced with the observed viral genetic and antigenic profiling in the host population[6]. The influenza virus evolves through antigenic drifts on the two surface proteins, hemagglutinin (HA) and neuraminidase (NA), the primary immuno-active components of influenza vaccines. Previous studies showed that mutations at epitope sites of the HA played a dominant role in characterizing virus antigenic change and were under higher immune selection pressure[5,7]. In addition to the major antigenic substitutions, the virus evolution is also critically influenced by epistatic mutations or mutation interference effects[8-10], making predictions challenging as the evolutionary dynamics is non-uniform across genomic regions and time.

Currently, influenza vaccine strain determination involves extensive surveillance and characterization of the virus in terms of genetic, antigenic evolution, and epidemiological profiles, a global effort coordinated by the World Health Organization (WHO)[11]. While antigenic evolution of influenza virus can be mapped by the antigenic cartography based on hemagglutinin inhibition (HAI) assay data[12], genetic evolution is mostly delineated with the phylogenetic trees[13]. Current prediction methods of virus evolution mainly focus on modelling the fitness of tree parts[14-16], using sequences of the HA or HA1 segment, with epitope or antigenic data incorporated as component information[14,15,17]. These methods projected the most likely predominant future lineages by tracing the fitness of clusters of strains. However, as the virus evolution is driven by major antigenic substitutions, a substantial amount of information is contained in the dynamic process of the fitness of mutations, characterizing which might provide critical information for predicting genetic evolution.

A site-based dynamic model for mutation forecasting. Our prediction method is

primarily based on modelling the time-resolved frequency pattern of mutations for individual

sites across virus genome segments (**Fig. 1**). The selective advantage of a mutation could be

reflected in their growing prevalence in the host population. To make a projection into the

future, the velocity of mutation frequency growth can be estimated by solving the first-order

derivative of a frequency function over a period of mutation adaption in the host population,

which is captured by the mutation transition time (**Methods**). Since the transition time is

specific to substituted residues, the dynamic model is site-based and time-dependent. Scanning

over the genome restores a global picture of mutation-selection. Such design shares the

advantage of the agent-based model that agglomerates individual agents carrying simple rules

and avoids fitting a complicated system with copious parameters or unrealistic assumptions[18].

The potential relationship between the genome-wide dynamics of mutation prevalence and

population epidemics is illustrated in **Extended Data Fig. 1**. The site-based model is also

highly computationally efficient in analyzing large genomic datasets.

Specifically, transition time in this study is defined as the duration for a mutation to

emerge until it reaches an influential frequency in the population. This quantity is related to

but different from the conventional fixation time, which covers a period from the first

observation time of a mutation to the point when it reaches 0.99 prevalence in the population.

The fixation time was reported to span over a long period of 4-32 years for the influenza virus

A(H3N2)[8]. In contrast, the transition time we identified for the same viral subtype had a median

length of only ~17 months and ranged between 0-7 years (**Extended Data Fig. 2**). As the

transition time calibrates the initial period of mutation adaptation, it may have the advantage

of informing the emerging genetic variants on a short-term time horizon. The transition time is

determined with a frequency threshold (θ) indicating fitness strength, at which the overall

mutation activities are detected to influence population epidemics significantly (**Extended

Data Fig. 1**); and it can be estimated using the virus epidemic-genetic association model we

previously developed[19] (**Methods**).

This site-based mutation dynamic model enables the prediction for fitness of competing

residues at individual sites, thereby the construction of a genome-wide fitness landscape of the

virus population in future time (**Fig. 1**). Thus, a pseudo consensus strain can be shaped

containing all mutations showing selective advantage relative to their precedent or competing

alleles in the upcoming epidemic season. It serves as a prototype to search for the optimal wild-

type virus for consideration as vaccine strain.

Identification of optimal wild-type virus The second part of our method is

identification of optimal wild-type virus based on the predicted virus population from a pool

of available strains (**Fig. 1**). The aim is to minimize the genetic distance of HA and NA proteins

to the forecasted representative strain of future virus population simultaneously. This could be

achieved by referencing the estimated relationship between vaccine effectiveness (VE) and

genetic distance (GD)[20-22], with a goal of maximizing the projected VE in the future virus

population (**Methods**). In particular, the VE-GD model is powered by a subset of epitopes, the

predictor codon sites, and their relative contribution to explaining VE (**Extended Data Table**

1). Therefore, the step of vaccine strain selection is essentially informed by the antigenic

relevance of mutations estimated over the seasons. Although only HA concentration is

standardized in the current vaccine production[23], both genes are major components of the

influenza vaccines; the integrative evaluation would provide a useful tool for identifying wild-

type virus for potential vaccine strains.

This two-step evolution prediction and wild-type virus selection method are referred to

as the "beth-1" for easy reference.

Genetic matching of retroactively predicted vaccine strains. We applied the beth-

1 to predict vaccine strains of the influenza A (H1N1)pdm09 (pH1N1) and A (H3N2) viruses.

Data was collected from the Global Initiative on Sharing All Influenza Data (GISAID)[24]

between 1999/2000 and 2021/22, involving a total number of 11,243 HA and 9,396 NA

sequences of pH1N1, and 31,501 HA and 28,967 NA sequences of H3N2 from ten

geographical regions in the Northern Hemisphere, covering North America (New York State,

California State, Canada), Europe (United Kingdom, Germany, France), and Asia (Hong Kong

SAR, South China provinces, Japan, Singapore) (**Extended Data Table 2**). The prediction was

performed using data up to February in season T, targeting the subsequent epidemic season

$T+1$ (October-April next year). We calculate the average amino acid (AA) mismatch between

the predicted strain and sequences of circulating viruses in the target season to determine

prediction accuracy. To fully understand the performance of beth-1 in the context of existing

vaccines, we calculated the following comparison groups: (1) a naïve approach using a

randomly selected circulating strain in season T, repeated ten times; (2) WHO-recommended

vaccine strains (the "current-system") for season $T+1$; (3) beth-1 predicted strains for season

T+1; (4) the "answer": the observed representative strains in season *T*+1 (**Methods**) (**Extended**
Data Table 3).

In the retroactive season-to-season prediction, beth-1 demonstrated substantially
improved prediction accuracy compared to the current-system on full-length HA and NA gene,
their epitopes, and for both pH1N1 and H3N2 subtypes (**Fig. 2, Extended Data Table 4**).
Three-year data was burnt-in for model building. For pH1N1, prediction was conducted for
seasons from 2012/13 to 2018/19. Averaging over the seven seasonal prediction outcomes,
beth-1 reduced the genetic mismatch of the current-system by 7.7 AAs (SD 6.3, paired *t*-test
p-value < 0.001) on full-length HA protein and reduced the HA epitope mismatch by 2.3 AAs
(SD 1.2, paired *t*-test *p*-value < 0.001) (**Fig. 2**). This accuracy was close to the best possible
outcome of a 0.7 AA mismatch given by the answer strain (**Fig. 2, Extended Data Table 4**).
Similarly, the NA proteins achieved significant improvements in vaccine matching by beth-1
on both the full-length NA sequence and its epitope regions. We also observed that the current-
system performed worse than the naïve approach, likely due to the lack of updates on vaccine
strains in multiple of the influenza seasons and a number of practical constraints of WHO
vaccine strain selection, such as the availability of egg-isolated virus and sufficiency of viral
growth in production platform[25]. Overall, in the retrospective validation data, beth-1 prediction
is expected to improve the current VE against influenza pH1N1 by an absolute 11.5% (95%
CI: 9.4-13.5), estimated by the VE-GD model.

For the H3N2 virus, between seasons 2002/03 to 2018/19, beth-1 prediction
significantly reduced genetic mismatch of the current-system by 4.2 AAs (SD 5.3) on full-
length HA and by 2.6 AAs (SD 3.6) on the epitope sites (**Fig. 2, Extended Data Table 4**).
These improvements corresponded to a VE increase by absolute 8.5% (95% CI: 6.8-10.2)
relative to the current vaccines against the H3N2. Among the seasons when the predicted
strains differ between the two-protein model and the single-protein model, the two-protein
model enhanced VE by absolute 12.88% (95% CI: 9.96 – 15.82, *t*-test *p*-value < 0.001) relative
to the HA-only model on average in three seasons.

Seasonal and geographical breakdown of prediction accuracies were also analyzed.
Year-by-year comparison with the current-system showed that the beth-1 gave vastly lower
genetic mismatch throughout the epidemic seasons (**Fig. 3, Extended Data Table 5-8**).
However, discrepancy between the two predictions narrowed in recent years as a result of both
annual vaccine updates and improved predictions of the current-system. A separate analysis by

geographical regions showed no systematic difference in vaccine mismatch across continents
by both prediction methods and for the two influenza A subtypes (**Extended Data Fig. 3,**
Extended Data Table 9-12).

We performed two predictions for the 2019/20 epidemic season: a prospective
prediction conducted in October 2019 using training data up to June 2019, from which the
predicted strains were recorded on blockchain **(Extended Data Table 13)**; and a retroactive
prediction conducted in March 2022 using data up to March 2019 to better match the current-
system's timeline. Genetic mismatch was calculated by all prediction experiments **(Extended**
Data Table 14). Between the 2019-03 and 2019-06 predictions by beth-1, the latter one
recorded 0.1 less HA epitope mismatches for pH1N1 and 1.1 less HA epitope mismatch for
H3N2. Both predictions showed significant superior performance comparing to the current-
system for pH1N1 on all protein segments and NA of H3N2, and showed non-inferior
performance in other segments **(Extended Data Table 14)**. We further evaluated the
immunogenicity of predicted strains for this particular season using clinical samples collected
in Hong Kong (**Methods, Extended Data Fig. 4**). Against the viral isolates of H1 and H3
positive samples, beth-1 strains induced significantly higher neutralizing antibodies in $PRNT_{50}$
for pH1N1 and non-inferior antibodies for H3N2, compared to the current vaccines (**Extended**
Data Fig. 5).

Prospective predictions and validations for the 2020/21 and 2021/22 seasons. To
further validate our prediction method, we conducted prospective predictions and validations
for two more seasons in 2020/21 and 2021/22. The predictions were blockchain recorded and
sent to the WHO before vaccine composition meetings (**Extended Data Table 13**). In 2020/21,
on the pH1N1 HA epitopes, beth-1 gave 2.7 AAs (SD 1.1) mismatch and the current-system
gave 3.8 AAs (SD 1.2) mismatch, while on the H3N2 HA epitopes, both methods resulted in
~13 AAs mismatch (**Fig. 2** dashed lines, **Extended Data Table 14**). This season was under the
impact of the COVID-19 pandemic since early 2020 that recorded sharply reduced sequence
samples of influenza[26,27]. Nevertheless, activities of influenza virus A resurged with relaxed
pandemic control measures in 2021/22[28].

In the 2021/22 season (**Fig. 2** dotted lines, **Extended Data Table 14**), for the pH1N1,
the HA epitope mismatch are ~3.5 AAs by both beth-1 and the current-system; however, for
H3N2, the HA epitope mismatches were 4.2 AAs (SD 1.6) by beth-1 and 10.8 AAs (SD 1.9)
by the current-system (p-value < 0.001). This genetic matching corresponds to expected VE

improvement resulted from beth-1 by absolute 31.7% (95% CI: 31.5-31.9) for the H3N2, compared to the current vaccines in the 2021/22 epidemic season.

To further understand the relationship between the beth-1 strains and the current vaccine viruses, we mapped them on phylogenetic trees for the three prospective seasons, 2019/20 – 2021/22 (**Fig. 4**). The phylogenies of pH1N1 delineated two major clades through the seasons, of which one is predominant with strains from Europe and the other is predominant with strains from Asia; while strains from North America are distributed in both clades. Interestingly, beth-1 and the current-system each captures one of the two clades in these seasons. For the H3N2, the epidemic seasons are dominated by a single major trunk. In 2019/20, beth-1 captured the correct trunk while the current-system did not; in 2020/21, the two methods identified almost the identical strain; and in the latest 2021/22 season, both methods successfully forecasted the correct clade, while beth-1 predicted strains extended slightly more into the future clusters.

How predictable are the pH1N1 and H3N2 viruses? Our predictions for the two influenza pH1N1 and H3N2 viruses allowed us to compare the predictability for the two subtypes by the genetic mismatch of the HA epitopes in the retrospective data. Ceiling of prediction can be approximated by the answer vaccine strain, that is, the observed representative strain in respective seasons. For the pH1N1 virus, the lowest possible mismatch of HA epitope was 0.7 AAs (SD 0.4) achieved by the answer strain, and the beth-1's mismatch was 1.2 AAs (SD 0.6) (**Fig. 2, Extended Data Table 4**). This suggests that a highly precise hit by predictive modelling is achievable for the pH1N1. However, for the H3N2 virus, the lowest possible HA epitope mismatch was as large as 3.5 AAs (SD 1.1), that is 2.8 more epitope mismatches compared to the pH1N1. Although the 5.1 AAs (SD 1.7) mismatch by beth-1 was only 1.6 residue away from the answer, the prediction for H3N2 would be ultimately bounded by the ceiling of genetic matching. The large genetic mismatch of the H3N2 vaccine strain might be attributed to the high genetic diversity of this virus[29,30], making the selection of a single representative wild-type strain challenging. New vaccinology strategies, such as developing broadly reactive vaccines, designing antigens containing multiple H3N2 strains, or preparing region-specific vaccines, may provide solutions from other dimensions to enhance vaccine protection against this subtype.

Comparing prediction performances of beth-1 and the existing prediction models for influenza evolution. Existing predictive models for influenza evolution mostly adopt

strain-based analyses and extract information presented by the phylogenetic structure. Neher et al. used the local branching index (LBI) to rank tree-nodes in a given phylogeny to identify lineage with the highest fitness as a progenitor of strains in an upcoming influenza season, where fitness is estimated by integrating exponentially discounted branch-length surrounding a node[16]. Luksza and Lassig calculated frequency of tree-clades and predicted the future predominant clade by an exponential function (the Malthusian model) with a certain growth rate[14]. Steinbruck and McHardy used allele dynamics plots to identify the top three alleles characterizing antigenic novelty of tree-clades in a given season, pre-screened based on the epitope information, HI data, and phylogeny[31,32]. Also based on the Malthusian model, Huddleston et al regarded strain rather than clade as a primary unit of analysis, and investigated different combination of factors to calculate growth rate for strain, including epitope mutations, non-epitope mutations (mutational load), LBI, HI antigenic novelty, and deep mutational scanning[15]. The best performing model of Huddleston et al incorporated mutational load and strain-specific HI data, trained for the H3N2. Herein, we compared beth-1 to the current state-of-the-art methods, the LBI and Huddleston's best model on H3N2 in the retrospective data from 2002/03 to 2018/19 (**Methods**). The results showed that beth-1 outperformed the existing models significantly (**Fig 5**). For instance, over the ten geographical regions, the mean mismatch obtained by beth-1 on full-length HA of H3N2 is 7.5 AAs (SD 2.2), lower than the 9.4 AAs (SD 2.8) by the method of Huddleston et al (p-value < 0.001) and 9.5 AAs (SD 4.7) by the LBI (p-value $= 0.008$). For predicting NA of H3N2 and pH1N1, we compare beth-1 to the LBI, as strain-specific HI data is not fully available to train the model of Huddleston et al. Results shows that the beth-1 gives significantly improved genetic matching compared to the LBI on full-length and epitopes of HA of pH1N1 and NA of H3N2, and full-length NA of pH1N1 (**Extended Data Fig. 6**).

Advantages of site-based dynamic model for evolution prediction. Strain-based dynamic model often requires the estimation of a single-valued growth rate for a given genome to project future fitness, whereas a site-based dynamic model is not constraint by a constant growth rate over genome space-time. This property leads to the following three advantages in predicting virus evolution. First, key mutations distributed over multiple clades could be captured as they arise, while the virus is trialing various epistatic combinations before shaping a stable lineage. Second, a site-based model avoids making assumption for directional mutation effects on fitness, which is self-evident in mutation frequency. In contrast, strain-based models

often need to assume negative effect for the non-epitope mutations, to offset the genetic

distance obtained from gene-based analysis in modelling evolutionary pathway[14-16], although

studies suggested that the alternative might be true[33,34]. Third, the site-based dynamic model is

adaptive to the altered residue fitness from epistasis and environmental factors[10,35], by sampling

and re-estimating the dynamic function at each time period. When such adaptive framework is

not in place, study showed that the fitness advantage estimated in the initial stage of mutation

emergence cannot predict their ultimate fixation[36]. However, the proposed site-based dynamic

model avoids making assumptions on the constant effect of mutations. Through adaptive

estimation of fitness by genomic site and time, it projects a probable fitness landscape including

all the trackable in-advantage mutations to the near future.

"Representativeness" of viral strains in the perspective of strain-based and site-

based analysis. Under the perspective of the existing strain-based analysis, a representative

virus is naturally indicated by the majority vote of strain or clades in a host population. While

under the perspective of site-based analysis, a genome-wide fitness landscape can be

constructed for a virus population, without missing a single mutation showing selective

advantage. Based on this fitness landscape, a mode estimator can be operated at individual sites

and generate a consensus sequence. Therefore, the consensus sequence is a natural

representative strain for a virus population through site-wise analysis of viral fitness. Beth-1 in

its primary objective establishes the theoretical framework to model mutation dynamics site-

wise and projects the fitness landscape into the future.

Limitation of the study. One major limitation in our analysis is the sparse time interval,

constrained by the sequence sample size in the earlier years, which prevented us from using

more complex trajectory functions for prediction. In the future, this sampling gap may be

gradually closed with increasing surveillance strength and global collaborations, and non-linear

functions can be explored. Second, the epitope mismatch could have been underestimated as it

was subjected to the known epitope sites, especially in the immune-subdominant protein

segments. Nevertheless, the comparison across prediction methods was fair with the same

evaluation criteria.

Current WHO vaccine virus selection considerations. Currently, WHO selects

vaccine strains by considering the emergence of virus with distinct genetic and antigenic

characteristics, their geographical spread, and the potential loss of effective binding of

antibodies from antisera of previously vaccinated subjects against the current circulating

viruses[25]. These factors depict a picture of the current global virus population and their
antigenic relationship to the previous representative viruses and vaccine strains, based on which
recommendations of vaccine strains are made. One major consideration of vaccine strain
selection is the HAI titre for antigenic characterization[37]. Nevertheless, the HAI only evaluates
one specific type of reaction, that is the prevention of HA binding to sialic acid receptor on
host cells inhibited by the anti-HA head antibodies[38], while other types of immune responses
are undetected, such as the antibodies against HA-stem and T-cell responses that also play
important roles in protection[39,40]. Developing improved assays with broader range or finer
specificity for vaccine protection may facilitate the assessment and selection of vaccine strain[25].
Another major constraint of the current vaccine strain selection process is the limited
availability of high-yield virus in embryonated eggs, with which more than 95% of the current
influenza vaccines are produced[41]. These candidate vaccine viruses (CVVs) are prepared from
representative strains by the WHO Collaborating Centres (CC) and Essential Regulatory Labs
(ERLs) before the annual WHO consultation meetings[25]. Thus, earlier prediction, even year-
round projection of future representative strains may facilitate the preparation of CVVs and the
subsequent vaccine strain selection, while the development of alternative vaccine technologies
for influenza virus would sidestep the constraints of egg-based platform. The potential of
computational optimized vaccine strains may be fully explored coupling with the availability
of new production platforms, as "new wine in new wineskins".

In summary, we have introduced a new computational method, the beth-1, for
predicting influenza evolution through the site-based fitness dynamic modelling, and enables
simultaneously considering multiple proteins to locate optimal wild-type virus. The model
demonstrated promising prediction performances in both retrospective and prospective real
data applications. The framework has potentially wide applications by virology labs, vaccine
manufacturers, health authorities and the WHO for indicating virus evolution and preparing
vaccine virus, to facilitate influenza vaccine strain selection towards more effective vaccines.

References

1   Petrova, V. N. & Russell, C. A. The evolution of seasonal influenza viruses. *Nat Rev Microbiol* **16**, 47-
60, doi:10.1038/nrmicro.2017.118 (2018).
2   Ampofo, W. K. *et al.* Improving influenza vaccine virus selection: report of a WHO informal consultation
held at WHO headquarters, Geneva, Switzerland, 14-16 June 2010. *Influenza and other respiratory*
viruses **7 Suppl 2**, 52-53, doi:10.1111/irv.12081 (2013).
3   Morris, D. H. *et al.* Predictive Modeling of Influenza Shows the Promise of Applied Evolutionary
Biology. *Trends in microbiology* **26**, 102-118, doi:10.1016/j.tim.2017.09.004 (2018).

4   Nelson, M. I. & Holmes, E. C. The evolution of epidemic influenza. *Nat Rev Genet* **8**, 196-205,
doi:10.1038/nrg2053 (2007).
5   Bush, R. M., Fitch, W. M., Bender, C. A. & Cox, N. J. Positive selection on the H3 hemagglutinin gene
of human influenza virus A. *Molecular biology and evolution* **16**, 1457-1465,
doi:10.1093/oxfordjournals.molbev.a026057 (1999).
6   Lässig, M., Mustonen, V. & Walczak, A. M. Predicting evolution. *Nature ecology & evolution* **1**, 77,
doi:10.1038/s41559-017-0077 (2017).
7   Koel, B. F. *et al.* Substitutions near the receptor binding site determine major antigenic change during
influenza virus evolution. *Science* **342**, 976-979, doi:10.1126/science.1244730 (2013).
8   Shih, A. C., Hsiao, T. C., Ho, M. S. & Li, W. H. Simultaneous amino acid substitutions at antigenic sites
drive influenza A hemagglutinin evolution. *Proc Natl Acad Sci U S A* **104**, 6283-6288,
doi:10.1073/pnas.0701396104 (2007).
9   Kryazhimskiy, S., Dushoff, J., Bazykin, G. A. & Plotkin, J. B. Prevalence of epistasis in the evolution of
influenza A surface proteins. *PLoS Genet* **7**, e1001301, doi:10.1371/journal.pgen.1001301 (2011).
10   Illingworth, C. J. & Mustonen, V. Components of selection in the evolution of the influenza virus: linkage
effects beat inherent selection. *PLoS Pathog* **8**, e1003091, doi:10.1371/journal.ppat.1003091 (2012).
11   Ampofo, W. K. *et al.* Improving influenza vaccine virus selection: report of a WHO informal consultation
held at WHO headquarters, Geneva, Switzerland, 14-16 June 2010. *Influenza and other respiratory*
viruses **6**, 142-152, e141-145, doi:10.1111/j.1750-2659.2011.00277.x (2012).
12   Smith, D. J. *et al.* Mapping the antigenic and genetic evolution of influenza virus. *Science* **305**, 371-376,
doi:10.1126/science.1097211 (2004).
13   Hadfield, J. *et al.* Nextstrain: real-time tracking of pathogen evolution. *Bioinformatics (Oxford, England)*
34, 4121-4123, doi:10.1093/bioinformatics/bty407 (2018).
14   Luksza, M. & Lassig, M. A predictive fitness model for influenza. *Nature* **507**, 57-61,
doi:10.1038/nature13087 (2014).
15   Huddleston, J. *et al.* Integrating genotypes and phenotypes improves long-term forecasts of seasonal
influenza A/H3N2 evolution. *eLife* **9**, doi:10.7554/eLife.60067 (2020).
16   Neher, R. A., Russell, C. A. & Shraiman, B. I. Predicting evolution from the shape of genealogical trees.
eLife **3**, doi:10.7554/eLife.03568 (2014).
17   Steinbrück, L., Klingen, T. R. & McHardy, A. C. Computational prediction of vaccine strains for human
influenza A (H3N2) viruses. *Journal of virology* **88**, 12123-12132, doi:10.1128/jvi.01861-14 (2014).
18   Kennedy, J. & Eberhart, R. in *Proceedings of ICNN'95-international conference on neural networks.*
1942-1948 (IEEE).
19   Wang, M. H. *et al.* Characterization of key amino acid substitutions and dynamics of the influenza virus
H3N2 hemagglutinin. *The Journal of infection* **83**, 671-677, doi:10.1016/j.jinf.2021.09.026 (2021).
20   Cao, L. *et al.* Rapid evaluation of COVID-19 vaccine effectiveness against symptomatic infection with
SARS-CoV-2 variants by analysis of genetic distance. *Nat Med*, doi:10.1038/s41591-022-01877-1
(2022).
21   Cao, L. *et al.* In silico prediction of influenza vaccine effectiveness by sequence analysis. *Vaccine* **39**,
1030-1034, doi:10.1016/j.vaccine.2021.01.006 (2021).
22   Cao, L. *et al.* Improving the prediction of influenza vaccine effectiveness by refined genetic distance
measure. *medRxiv*, 2023.2002. 2014.23285900 (2023).
23   Krammer, F. *et al.* NAction! How Can Neuraminidase-Based Immunity Contribute to Better Influenza
Virus Vaccines? *mBio* **9**, doi:10.1128/mBio.02332-17 (2018).
24   Shu, Y. & McCauley, J. GISAID: Global initiative on sharing all influenza data - from vision to reality.
Euro surveillance : bulletin Europeen sur les maladies transmissibles = European communicable
disease bulletin **22**, doi:10.2807/1560-7917.Es.2017.22.13.30494 (2017).
25   Group, W. H. O. W. *et al.* Improving influenza vaccine virus selection: report of a WHO informal
consultation held at WHO headquarters, Geneva, Switzerland, 14-16 June 2010. *Influenza Other Respir*
Viruses **6**, 142-152, e141-145, doi:10.1111/j.1750-2659.2011.00277.x (2012).
26   Uyeki, T. M., Wentworth, D. E. & Jernigan, D. B. Influenza Activity in the US During the 2020-2021
Season. *JAMA* **325**, 2247-2248, doi:10.1001/jama.2021.6125 (2021).
27   Subbarao, K. What influenza activity can we anticipate in 2022? *Med J Aust* **216**, 239-241,
doi:10.5694/mja2.51437 (2022).
28   Larrauri, A. & Prosenc Trilar, K. Preparing for an influenza season 2021/22 with a likely co-circulation
of influenza virus and SARS-CoV-2. *Euro surveillance : bulletin Europeen sur les maladies*
transmissibles = European communicable disease bulletin **26**, doi:10.2807/1560-
7917.ES.2021.26.41.2100975 (2021).

29 Su, Y. C. F. *et al.* Phylodynamics of H1N1/2009 influenza reveals the transition from host adaptation to
immune-driven selection. *Nat Commun* **6**, 7952, doi:10.1038/ncomms8952 (2015).
30 Ma, Y. *et al.* The Phylodynamics of Seasonal Influenza A/H1N1pdm Virus in China Between 2009 and
2019. *Front Microbiol* **11**, 735, doi:10.3389/fmicb.2020.00735 (2020).
31 Steinbruk, L., Klingen, T. R. & McHardy, A. C. Computational Prediction of Vaccine Strains for Human
Influenza A (H3N2) Viruses. *Journal of virology* **88**, 12123-12132, doi:10.1128/Jvi.01861-14 (2014).
32 Steinbruck, L. & McHardy, A. C. Allele dynamics plots for the study of evolutionary dynamics in viral
populations. *Nucleic Acids Res* **39**, doi:ARTN e410.1093/nar/gkq909 (2011).
33 Bershtein, S., Segal, M., Bekerman, R., Tokuriki, N. & Tawfik, D. S. Robustness-epistasis link shapes
the fitness landscape of a randomly drifting protein. *Nature* **444**, 929-932, doi:10.1038/nature05385
(2006).
34 Li, C., Qian, W. F., Maclean, C. J. & Zhang, J. Z. The fitness landscape of a tRNA gene. *Science* **352**,
837-840, doi:10.1126/science.aae0568 (2016).
35 Strelkowa, N. & Lassig, M. Clonal interference in the evolution of influenza. *Genetics* **192**, 671-682,
doi:10.1534/genetics.112.143396 (2012).
36 Barrat-Charlaix, P., Huddleston, J., Bedford, T. & Neher, R. A. Limited Predictability of Amino Acid
Substitutions in Seasonal Influenza Viruses. *Molecular biology and evolution* **38**, 2767-2777,
doi:10.1093/molbev/msab065 (2021).
37 Houser, K. & Subbarao, K. Influenza vaccines: challenges and solutions. *Cell host & microbe* **17**, 295-
300, doi:10.1016/j.chom.2015.02.012 (2015).
38 Stephenson, I., Wood, J. M., Nicholson, K. G. & Zambon, M. C. Sialic acid receptor specificity on
erythrocytes affects detection of antibody to avian influenza haemagglutinin. *J Med Virol* **70**, 391-398,
doi:10.1002/jmv.10408 (2003).
39 Reber, A. & Katz, J. Immunological assessment of influenza vaccines and immune correlates of
protection. *Expert Rev Vaccines* **12**, 519-536, doi:10.1586/erv.13.35 (2013).
40 Doherty, P. C. *et al.* Effector CD4+ and CD8+ T-cell mechanisms in the control of respiratory virus
infections. *Immunol Rev* **159**, 105-117, doi:10.1111/j.1600-065x.1997.tb01010.x (1997).
41 Perdue, M. L. *et al.* The future of cell culture-based influenza vaccine production. *Expert Rev Vaccines*
10, 1183-1194, doi:10.1586/erv.11.82 (2011).

Fig. 1 | Overview of the computational model and study design. Site-wise fitness dynamics are modelled and projected to the next season (T+1). Based on the predicted genome-wide fitness landscape of future virus population, an optimal wild-type virus can be selected to maximize expected vaccine effectiveness integrating information of both hemagglutinin and neuraminidase genes. The prediction is validated by genetic mismatch against observed circulating viruses in the Northern Hemisphere retroactively and prospectively for pH1N1 and H3N2. In addition, mouse model is used to evaluate the antibodies elicited by predicted vaccine strain in neutralizing the clinical isolates in 2019/20 season.

Fig. 2 | Overall prediction performance in retrospective and prospective data for influenza pH1N1 and H3N2. a, Full-length protein. **b**, Epitope. The boxplots display distribution of yearly genetic mismatch (amino acids, AA) of vaccine strains in retrospective data during seasons 2012/13-2018/19 for pH1N1 and 2002/03-2018/19 for H3N2. Panel **b** left Y-axis: average genetic mismatch of pH1N1; right Y-axis: average genetic mismatch of H3N2. Four prediction comparison groups include the current-system (green), random selection (yellow), beth-1 (pink), and the actual representative strain (blue). The asterisks represent mismatches in prospective prediction in the 2019/20, 2020/21 and 2021/22 flu seasons. In both the retrospective and prospective validations, beth-1 shows prevailingly lower genetic mismatch on the protein segments of the two influenza subtypes, compared to the current-system.

Fig. 3 | Genetic mismatch of predicted vaccine strains of pH1N1 and H3N2 against circulating viruses in retrospective data. a, pH1N1, HA full sequence (566 codons). **b**, pH1N1, NA full sequence (469 codons). **c**, pH1N1, HA epitopes (50 sites). **d**, pH1N1, NA epitopes (30 sites). **e**, H3N2, HA full sequence (566 codons). **f**, H3N2, NA full sequence (469 codons). **g**, H3N2, HA epitopes (131 sites). **h**, H3N2, NA epitopes (12 sites). Vaccine strains were predicted by the proposed method (beth-1) and were assessed by the average amino acids mismatch (Y-axis) against circulating viruses in ten geographical regions in respective epidemic seasons (X-axis). In the retrospective epidemic seasons and the genomic segments, the genetic mismatch of beth-1 was prevailingly lower than the current-system.

Fig. 4 | Prospectively predicted vaccine strains on phylogenetic tree from 2019/20 to 2021/22. a, 2019/20, pH1N1. **b**, 2019/20, H3N2. **c**, 2020/21, pH1N1. **d**, 2020/21, H3N2. **e**, 2021/22, pH1N1. **f**, 2021/22, H3N2. Predicted vaccine strains of the proposed method (beth-1) and current-system are marked in phylogenetic trees. Predictions were made prospectively for the coming epidemic season in the Northern Hemisphere (highlighted by yellow band in the background). The phylogeny relationship shows that beth-1's predictions are generally more advanced into the future. For the pH1N1, each prediction captures one of the two major clades in the epidemic seasons. For the H3N2, beth-1 captures the correct clades in 2019/20 while the current system does not; both systems locate the predominant clades for 2020/21 and 2021/22, while beth-1 strains are slightly more advanced into future.

Fig. 5 | Comparison of beth-1 to alternative computational models for influenza virus evolution on HA of H3N2. a, HA full-length. **b**, HA epitopes. Retroactive season-to-season prediction is performed between 2002/03 and 2018/19 by alternative methods, including beth-1 (this study), Local Branching Index (LBI) (Neher et al 2014), the model of Huddleston et al (2020) and the current-system. Prediction accuracy is assessed by genetic mismatch of the predicted strains to the circulating virus in the upcoming epidemic, averaged over the 17 epidemic seasons. The beth-1 shows significantly lower genetic mismatch compared to the existing models on both HA full-length and epitopes of H3N2.

Methods

Dataset   We downloaded genetic sequences of the influenza virus from the Global

Initiative on Sharing All Influenza Data (GISAID)[24]. All samples were retrieved and analyzed

if the strain has complete sequence and were isolated from the target epidemic seasons and

geographical regions (**Extended Data Table 2**). The data of genetic sequences for pH1N1

spanned from 2009/10 to 2021/22, and for the H3N2 from 1999/2000 to 2021/22. Ten

geographical regions in the Northern Hemisphere (NH) were considered, including North

America (New York State, California State, Canada), Europe (United Kingdom, Germany,

France), and Asia (Hong Kong SAR, South China provinces, Japan, Singapore). In 2020/21

and 2021/22, due to sharp decline of available samples in these regions, all sequences in NH

were considered. Overall, the total number of genetic sequences used in analysis was 42,744

for the HA and 38,093 for the NA. Sequence alignment was performed by MEGA (7.0.26)[42].

All statistical analysis were conducted in R version 4.1.3[43]. All *p*-values reported are two-sided.

The vaccine strain prediction and selection model: beth-1   In our prior work[19], we used

the *threshold θ* to detect the change-point of mutation prevalence in relation to a significant

rise in epidemic level, and the *extended effective mutation period* (*h*) to measure the duration

of a mutation to remain in selective advantage in the host population after reaching θ. Let

$x_{jk}(t)$ denote amino acid residue or nucleotide type at site *j* in time *t*. $j \in J$, where *J* is the

sequence length. $k \in K_j$, and K_j is the total number of unique substitutions observed at site *j*.

$K_j \leq 4$ for nucleotide sequence, and $K_j \leq 20$ for amino acid sequence. Amino acid sequences

are used in the prediction modelling of this study. Denote the prevalence of $x_{jk}(t)$ in host

population as $p_{jk}(t)$. For a mutation $x_{jk}(t)$, let t_{jk}^0 denote the time point when $p_{jk}(t^0)$=0

and $p_{jk}(t^0 + 1) > 0$; and t_{jk}^θ denote the time when $p_{jk}(t^\theta) = \theta$. The *transition time* (τ) for

a mutation $x_{jk}(t)$ is defined as,

$$\tau_{jk}(\theta, t) = t_{jk}^\theta - t_{jk}^0.$$ **Eq. 1**

τ describes the duration that a mutation takes from its first emergence to reaching an influential

prevalence in population; $\tau > 0$. The parameters (θ, h) are estimated from fitting the

epidemic trend and a summary statistic of genetic activities, the *g*-measure[19], towards

optimizing a goodness-of-fit statistic $S(\cdot)$. We have,

$$(\hat{\theta}, \hat{h}) = \arg \max_{\theta \in \Theta, h \in H} S[y(t), g(\theta, h, t)],$$ **Eq. 2**

where $y(t)$ is a measure of epidemic level through time. $\Theta \in (0,1)$, and $H = \{0,1,2,...\}$. The

g-measure is written as,

$$g(\theta, h, t) = \mathbf{m}^t \cdot \mathbf{p}^t = \sum_{j,k} m_{jk}(\theta, h, t) p_{jk}(t), \qquad \textbf{Eq. 3}$$

in which $m_{jk}(\theta, h, t)$ is a time-dependent indicator function of influential mutations (or

effective mutations, EM)[19], such that $m_{jk}(\theta, h, t) \triangleq I\{t_{jk}^0 \le t \le t_{jk}^\theta + h, j \in J, k \in K_j\}$. Thus,

the subset of influential mutations varies across epidemic seasons (**Extended Data Table 1**).

We showed in other works that the *g*-measure is a good predictor for epidemic cycles of the

A(H3N2)[19], A(H1N1)pdm09[44] and COVID-19[45,46]. In this study, $S(\cdot)$ is the R-square of a linear

model, of which the response variable $y(t)$ is the annual sero-positivity rate, and independent

variables include the *g*-measure, mean temperature, absolute humidity and season, fitted

separately by geographical regions in the training set (**Extended Data Table 15**). When

sufficient sequences and epidemic data are available, the time-varying reproduction number R_t

is a preferred quantity for $y(t)$[45,46].

Prediction of future mutation prevalence can be made by solving a classical initial value

problem of the differential equation,

$$\begin{cases} f'(t) = F[t, p_{jk}(t), \tau_{jk}(\theta, t)], & t > T \\ \quad\quad f(T) = \eta \end{cases}, \qquad \textbf{Eq. 4}$$

where $f(t)$ is a function describing mutation prevalence through time, $F[t, p_{jk}(t), \tau_{jk}(\theta, t)]$ is

the velocity of the prevalence change, η is the initial value, and T is the latest observation

time. Using Euler's method of approximation, the projected mutation prevalence at time $T+1$

for substitution k at position j is,

$$\hat{p}_{jk}(T + 1) = \{p_{jk}(T) + F[T, p_{jk}(T), \hat{\tau}_{jk}(\theta, T)]\} \cdot C, \qquad \textbf{Eq. 5}$$

in which $k \in K_j$, and $j \in J$. $C = 1/\sum_{k \in K_j} \hat{p}_{jk}(T + 1)$ is a normalization constant. $\hat{\tau}_{jk}(\theta, T)$

is estimated with history transition event at site j by the time of making prediction. When no

history of transition event is available for the site, its transition time is estimated by the mean

transition time of mutations from the same protein in the training data (**Extended Data Table**

15).

Eq. 4 and **5** describe the prediction step. In practice, the model form of $F(t)$ shall be

chosen such that the training data can support a robust estimation. In the earlier seasons, only

a dozen influenza sequence samples were available each year in many regions. Thus, yearly

interval is chosen and the simplest form of $F(t)$ as a rate is adopted. Specifically,

530
$$F\left[T, p_{jk}(T), \hat{\tau}_{jk}(\theta, T)\right] = \frac{\Delta p_{jk}(T)}{\Delta t} = \frac{p_{jk}(T) - p_{jk}[T - \hat{\tau}_{jk}(\theta, T)]}{\hat{\tau}_{jk}(\theta, T)} \quad . \qquad \textbf{Eq. 6}$$

531   For viruses with more abundant samples at denser time interval, other model form of $F(t)$ can

532   be incorporated.

533   Based on the projected genome-wide fitness landscape given by **Eq. 5**, we can thus

534   select the most advantageous mutations per site in $T+1$ using the mode estimator,

535
$$k_j^* = \arg\max_{k \in K_j} \hat{p}_{jk}(T + 1).$$

536   This gives us a consensus strain to represent the projected future virus population,

537
$$\{x_j^c(T + 1) \mid c = k_j^*\}. \qquad \textbf{Eq. 7}$$

Since a high mutation prevalence infers selective advantage of the mutation in population, the

mode sequence carries all the in-advantage mutations at time $T+1$. The prediction model is

trained using sample sequences from the South-east Asia region that is the source of influenza

A epidemics[47,48]; for seasons after 2019/20, all Northern Hemisphere samples were used.

According to the guideline of vaccine preparation[49], the vaccine strains shall be selected

among the available wildtypes. Thus, we locate the optimal wildtype virus for vaccine strains

based on the predicted HA and NA sequences. We can use the prediction model for vaccine

effectiveness by genetic distance (VE-GD)[20,21] to select the available present wildtype virus

that maximizes the estimated VE against the predicted virus population.

A brief account of the VE-GD model is described here. The model captures the

statistical relationship between VE and the genetic mismatch of vaccine strains against the

circulating viruses, fitted with VE data collected from published observational studies over

multiple seasons and genetic sequences matched to the same period and geographical region

of the VE surveys[20-22]. Let $\{x_j^i\}$ denote a vaccine sequence indexed by i. The genetic distance

d_s^i was measured by the average hamming distance on gene s between a vaccine strain i and

the circulating viruses, $s \in S = \{HA, NA\}$. The effect of genetic mismatch was first estimated

for individual genes by $VE \sim d_s^i$ using a linear model, resulting the estimated effect size

$\hat{\beta}_s$. Next, the effect of multiple genes is combined through the polygenic risk score (PRS)[50,51]

to form an integrated predictor $\sum_{s \in S} \hat{\beta}_s d_s^i$. We showed that the VE-GD relationship for

influenza virus was best fitted when the genetic distance was evaluated on a subset of EMs

residing on epitope A or B for H3N2 and the EMs on antigenic sites for pH1N1[22], referred to

as the predictor codon sites $W(t)$ (**Extended Data Table 1**). For a future season, using the

projected $\hat{p}_j^k(T + 1)$ in **Eq. 5** and fitted (θ, h) (**Extended Data Table 15**), we can thus

obtain the predicted EMs and corresponding $W(T+1)$.

The genetic distance between a candidate vaccine virus i and $x_j^c(T + 1)$ is,

$$d_s^i(T + 1) = \sum_{j \in W(t), t \in [1, T+1]} I\{x_j^i \neq x_j^c(T + 1)\}, \qquad \text{Eq. 8}$$

Thus, VE of a candidate vaccine strain i in season $T+1$ can be estimated by,

$$\widehat{VE}_i(T + 1) \sim \sum_{s \in S} \hat{\beta}_s d_s^i(T + 1). \qquad \text{Eq. 9}$$

With **Eq. 9**, the optimal wild-type virus $\{x_j^v\}$ can be selected by searching through all

available strains for the one that maximizes the VE in $T+1$, integrating mutation information

from both HA and NA. Hence, the estimated optimal wild-type virus is given by,

$$\{x_j^v\} = \arg \max_{i \in N}\{ x_j^i \mid \widehat{VE}_i(T + 1)\}, \qquad \text{Eq. 10}$$

where N is the total number of available wildtype viruses on and before time T.

Vaccine strains predicted by alternative models.   The list of predicted strains by the beth-

1 and the current-system can be found in **Extended Data Table 3**. Random selection is

repeated ten times and the averaged genetic mismatch is reported. The answer strain is set as

the closest wildtype to the consensus sequence in season $T+1$ obtained using the observed data.

The predicted strains of LBI[16] are obtained by running the model's source code in python

(https://github.com/rneher/FitnessInference) on the same training data as the beth-1. The

predicted strains of Huddleston et al's model[15] (HI + mutational load) are directly retrieved

from Github estimated by the original authors (https://github.com/elifesciences-

publications/flu-forecasting). Two-sample t-test is used to compare log mismatch against

circulating viruses between alternative prediction models in a given season. When comparing

model performances over multiple seasons, paired t-test of two methods by season is used to

control seasonal variation.

Phylogenetic tree   To better visualize the predicted strains, we present them in the time-

scaled maximum-likelihood phylogenies generated from TreeTime[52]. The input tree was built

in MEGA (7.0.26) using the neighbor-joining (NJ) approach with Jukes-Cantor gamma

distance of shape parameter 4[42]. Genetic sequences for building the tree were sampled in

proportion to continent distributions in the corresponding epidemic season, targeting a total

number of 400 sequences per season for each of the pH1N1 and H3N2 subtype per season. In

2020/21, all available sequences in the NH were used.

Clinical samples for animal experiment   Between October 2019 to February 2020, a total

number of 117 positive H1 and 43 H3 nasopharyngeal swab samples were collected in the

Prince of Wales Hospital, Hong Kong SAR. Ten samples from each of the H1 and H3 subtypes

were randomly selected for virus isolation in hMDCK cells. The sequences of the HA gene

from each virus was then determined by sanger sequencing. Phylogenetic analysis showed that

the viruses belonged to the same clusters of their respective subtype in these isolates, of which

HA ectodomain shared >98% amino acids similarity for the pH1N1 and >96% for H3N2. Thus,

one pH1N1 and one H3N2 virus from these isolates were randomly picked for animal

experiments. Human study ethics approval has been obtained from the Joint Chinese University

of Hong Kong – New Territories East Cluster Clinical Research Ethics Committee.

Cell culture   Humanized Madin-Darby canine kidney (hMDCK) cells[53] and human

embryonic kidney (HEK) 293T cells were used in this study. Both cells were maintained in

minimum essential medium (MEM) supplemented with 10% fetal bovine serum (FBS; Gibco),

25 mM HEPES, and 100 U/mL penicillin-streptomycin (PS; Gibco).

Virus rescuing by reverse genetics   The H1N1 A/Puerto Rico/8/1934 (PR8) eight-plasmid

reverse genetic system was used to produce the recombinant influenza virus with the predicted

HA and NA genes[54]. Chimeric 6:2 recombinant viruses with six internal genes from PR8 (PB2,

PB1, PA, NP, M, and NS) and 2 genes (HA and NA) from strain of interest were synthesized

by Sangon Technology (Guangzhou, China) and were cloned into the pHW2000 vector[55]. The

HEK 293T cells and hMDCK cells were then mixed at a ratio of 2:1 one-day before transfection

and co-cultured in a 6-well plate until they reached 70% confluence. For each recombinant

virus, 16 μL TransIT®-LT1 (Mirus) and 1 μg for each of the 8 plasmids encoding the

corresponding virus fragments were mixed thoroughly for transfection. The medium was

replaced with 1mL of MEM supplemented with 25 mM HEPES and 100 U/mL PS at 6-hour

post transfection. At 24-hour post-transfection, another 1mL of MEM was added,

supplemented 25 mM HEPES, 100 U/mL PS, and 1 μg/mL tosylphenylalanyl chloride methyl

ketone (TPCK) trypsin (Sigma). At 72-hour post-transfection, cell supernatants were

inoculated into hMDCK cells maintained in MEM containing 25mM HEPES, 100 U/mL PS,

and 1 μg/mL TPCK trypsin. The viruses were harvested at 72 hours after infection and titrated

by plaque assay.

Immunization of the influenza vaccine virus in a mouse model   The research protocol

of animal experiment was carried out in strict accordance with the recommendations and was

approved by the Teaching and Research Committee on the Use of Live Animals (CULATR

5598-20) of the University of Hong Kong. Batches of ten 8-10 weeks old female BALB/c mice

were inoculated intraperitoneally (i.p) with 10^5 PFU of Addavax-adjuvant recombinant virus

in 200 μL of phosphate-buffered saline (PBS; Gibco). At twenty-one days post immunization,

peripheral blood was drawn from the immunized mice by cardiac puncture, and the serum was

collected by centrifugation. All sera were treated with receptor destroying enzyme (RDE;

Sigma) for 16-18 hours to reduce non-specific antibodies, then heated at 56 degrees for 30

minutes to inactivate the RDE residue. The serum was stored at -80 degrees for further

experiments.

Immunogenicity assessment based on plaque reduction neutralization test   hMDCK

cells were used for virus titration by plaque assay and plaque reduction neutralization test

(PRNT). hMDCK cells were prepared in a 12-well plate at least 24 hours before PRNT and

cultured until obtaining a 100% confluent monolayer in each well. Two-fold serial dilutions of

antiserum from 1:10 to 1:1280 were prepared, where each 100 μL was mixed with 100 PFU of

viruses and incubated at 37°C for 1 hour. Cells were washed once with PBS and inoculated

with corresponding virus-antiserum mixture for further incubation at 37°C for one hour. After

that, the supernatant in each well was replaced with 3 mL of MEM supplied with a final

concentration of 25 mM HEPES, 100 U/mL PS, 1% agarose gel, and 1 μg/mL TPCK trypsin.

Against the live viruses, the highest dilution of the antiserum that inhibited at least 50% of the

viral plaques ($PRNT_{50}$ titer) was recorded. The plate was inverted and incubated at 37°C for

48 hours. Finally, the cells were fixed with 4% formaldehyde and stained with 0.5% crystal

violet for virus plaque counting. We recorded the highest dilution of the antiserum that

inhibited at least 50% of the viral plaques as the $PRNT_{50}$ titer of the antiserum. Detailed

information of the reagents and cell lines used in this study were listed in **Extended Data**

Table 16.

42   Kumar, S., Stecher, G. & Tamura, K. MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0
for Bigger Datasets. *Molecular biology and evolution* **33**, 1870-1874, doi:10.1093/molbev/msw054
(2016).

43   Team, R. C. R: A language and environment for statistical computing.   (2013).

44   Lou, J. *et al.* Predicting the dominant influenza A serotype by quantifying mutation activities.
International journal of infectious diseases : IJID : official publication of the International Society for
Infectious Diseases **100**, 255-257, doi:10.1016/j.ijid.2020.08.053 (2020).

45   Lou, J. *et al.* Quantifying the effect of government interventions and virus mutations on transmission
advantage during COVID-19 pandemic. *Journal of infection and public health* **15**, 338-342,
doi:10.1016/j.jiph.2022.01.020 (2022).

46   Zhao, S. *et al.* Modelling the association between COVID-19 transmissibility and D614G substitution in
SARS-CoV-2 spike protein: using the surveillance data in California as an example. *Theoretical biology*
& medical modelling **18**, 10, doi:10.1186/s12976-021-00140-3 (2021).

47   Rambaut, A. *et al.* The genomic and epidemiological dynamics of human influenza A virus. *Nature* **453**,
615-619, doi:10.1038/nature06945 (2008).

48   Bedford, T. *et al.* Global circulation patterns of seasonal influenza viruses vary with antigenic drift.
Nature **523**, 217-220, doi:10.1038/nature14460 (2015).

49   Weir, J. P. & Gruber, M. F. An overview of the regulation of influenza vaccines in the United States.
Influenza Other Respir Viruses **10**, 354-360, doi:10.1111/irv.12383 (2016).

50   International Schizophrenia, C. *et al.* Common polygenic variation contributes to risk of schizophrenia
and bipolar disorder. *Nature* **460**, 748-752, doi:10.1038/nature08185 (2009).

51   Torkamani, A., Wineinger, N. E. & Topol, E. J. The personal and clinical utility of polygenic risk scores.
Nat Rev Genet **19**, 581-590, doi:10.1038/s41576-018-0018-x (2018).

52   Sagulenko, P., Puller, V. & Neher, R. A. TreeTime: Maximum-likelihood phylodynamic analysis. *Virus*
evolution **4**, vex042, doi:10.1093/ve/vex042 (2018).

53   Takada, K. *et al.* A humanized MDCK cell line for the efficient isolation and propagation of human
influenza viruses. *Nature microbiology* **4**, 1268-1273, doi:10.1038/s41564-019-0433-6 (2019).

54   Hoffmann, E., Neumann, G., Kawaoka, Y., Hobom, G. & Webster, R. G. A DNA transfection system
for generation of influenza A virus from eight plasmids. *Proc Natl Acad Sci U S A* **97**, 6108-6113,
doi:10.1073/pnas.100133697 (2000).

55   Liang, W. *et al.* Egg-adaptive mutations of human influenza H3N2 virus are contingent on natural
evolution. *PLoS Pathog* **18**, e1010875, doi:10.1371/journal.ppat.1010875 (2022).

Data availability

All data used in this study is publicly available. Viral sequence data were downloaded from the global initiative on sharing all influenza data (GISAID) at *http://platform.gisaid.org/* and the accession numbers were provided in the online supplementary acknowledgment table.

Code availability

The code is available at *https://github.com/JingzhiLou/beth-1*. Access and use of the code is subject to a revocable, non-transferable, and limited right for the exclusive purpose of undertaking academic or not-for-profit research. Use of the Code or any part thereof for commercial purposes requires a Commercial License Agreement from Beth Bioinformatics (info@bethbio.com).

Acknowledgments

We thank the GISAID Initiative for critical surveillance efforts and open data sharing, and we greatly acknowledge the authors' contributions, submitting and originating laboratories.

Funding

Health and Medical Research Fund, the Food and Health Bureau, the Government of the Hong Kong Special Administrative Region: INF-CUHK-1 (E.K.Y.), COVID190103 (M.H.W.) and 19180932 (C.K.P.M.)

National Natural Science Foundation of China: 31871340 (M.H.W.) and 71974165 (M.K.C.C.) the Chinese University of Hong Kong Direct Grant: PIEF/Ph2/COVID/06 and 4054600 (M.H.W.)

Guangdong-Hong Kong-Macau Joint Laboratory of Respiratory Infectious Disease: 20191205 (C.K.P.M.)

Visiting scientist scheme from Lee Kong Chian School of Medicine, Nanyang Technological University, Singapore (C.K.P.M.)

Research Grants Committee, General Research Fund: 16301319 (P.P.H.C.)

Author contributions

M.H.W. conceived the study. M.H.W. and J.L. developed the prediction method. J.L. analyzed the data and prepared the results. C.K.P.M. designed the animal experiment. P.K.C. and Z.C. provided clinical samples. W.L. and C.K.P.M. conducted the immunology experiments. L.C., S. Z., H.Z., Y.Z., C.L. and Q.L. contributed to data collection and interpretation. E.Y., M.H.W.

711   and C.K.P.M. acquired funding. M.H.W. and J.L. wrote the original manuscript. C.K.P.M. and

712   W.L. revised the manuscript. Z.C., R.W.Y.C., P.P.H.C., and M.K.C.C. reviewed and edited the

713   manuscript. B.C.Y.Z. and E.K.Y. supervised the project.

Competing interests

715   M.H.W., J.L., M.K.C.C. and B.C.Y.Z. filed a related patent PCT/CN2019/091652. M.H.W.

716   and B.C.Y.Z. are shareholders of Beth Bioinformatics Co., Ltd. B.C.Y.Z. is a shareholder of

717   Health View Bioanalytics Ltd. The remaining authors declare no competing interests.

Additional information

719   Extended data is available for this paper.

720   Correspondence and requests for materials should be addressed to M.H.W. and C.K.P.M.