TCCIA: A Comprehensive Resource for Exploring CircRNA in Cancer Immunotherapy

Shixiang Wang[1,2,#], Yi Xiong[3,4,5,#], Yihao Zhang[3,4,5,#], Haitao Wang[6,^], Minjun Chen[1], Jianfeng Li[7], Peng Luo[8], Yung-Hung Luo[9,10], Markus Hecht[11], Benjamin Frey[12], Udo S Gaipl[12], Xuejun Li[3,4,5,*], Qi Zhao[2,*], Hu Ma[1,*], Jian-Guo Zhou[1,*]

Affiliations of authors:

[1] Department of Oncology, The Second Affiliated Hospital of Zunyi Medical University, Zunyi, 563000, P. R. China.

[2] Department of Experimental Research, Sun Yat-sen University Cancer Center, State Key Laboratory of Oncology in South China, Collaborative Innovation Center for Cancer Medicine, Sun Yat-sen University, Guangzhou, 510060, P. R. China.

[3] Xiangya School of Medicine, Central South University, Changsha, 410013, P. R. China.

[4] Hunan International Scientific and Technological Cooperation Base of Brain Tumor Research, Xiangya Hospital, Central South University, Changsha, 410008, P. R. China.

5 Department of Neurosurgery, Xiangya Hospital, Central South University, Changsha, 410008, P. R. China.

[6] Center for Precision Medicine Research and Training, Faculty of Health Sciences, University of Macau, Macau SAR, 999087, P. R. China.

[7] State Key Laboratory of Medical Genomics, Shanghai Institute of Hematology, National Research Center for Translational Medicine, Rui-Jin Hospital, Shanghai Jiao Tong University, School of Medicine, Shanghai, 200025, P. R. China.

[8] Department of Oncology, Zhujiang Hospital, Southern Medical University, Guangzhou, 510091, P. R. China.

[9] Department of Chest Medicine, Taipei Veterans General Hospital, Taipei, Taiwan.

[10] School of Medicine, College of Medicine, National Yang Ming Chiao Tung University, Taipei, Taiwan.

[11] Department of Radiotherapy and Radiation Oncology, Saarland University Medical Center,

Homburg, 66421, Germany.

[12] Department of Radiation Oncology, Universitätsklinikum Erlangen, Erlangen, 91054, Germany.

^ Current address: Thoracic Surgery Branch, Center for Cancer Research, NCI, NIH, Bethesda, MD 20892, United States

Co-first author.

* Corresponding author.

Correspondence: Jian-Guo Zhou (jianguo.zhou@zmu.edu.cn), Hu Ma (mahuab@163.com), Qi Zhao (zhaoqi@sysucc.org.cn), Xuejun Li (lxjneuro@csu.edu.cn)

Lead contact: Jian-Guo Zhou

Keywords: Circular RNA, Database, Checkpoint immunotherapy, Biomarker

Abstract

Background Immunotherapies targeting immune checkpoints have gained increasing attention in cancer treatment, emphasizing the need for predictive biomarkers. Circular RNAs (circRNAs) have emerged as critical regulators of tumor immunity, particularly in the PD-1/PD-L1 pathway, and have shown potential in predicting the efficacy of immunotherapies. However, the precise roles of circRNAs in cancer immunotherapy remain incompletely understood. While existing databases focus on either circRNA profiles or immunotherapy cohorts, there is currently no platform that enables the exploration of the intricate interplay between circRNAs and anti-tumor immunotherapy. Therefore, the development of a comprehensive resource that integrates circRNA profiles, immunotherapy response data, and clinical benefits is crucial for advancing our understanding of circRNA-mediated tumor-immune interactions and developing effective immunotherapy biomarkers.

Methods To address these gaps, we constructed the Cancer CircRNA Immunome Atlas (TCCIA), the first database that combines circRNA profiles, immunotherapy response data, and clinical outcomes across multi-cancer types. The construction of TCCIA involved applying standardized preprocessing to the raw sequencing FASTQ files, characterizing circRNA profiles using CIRCexplorer2, analyzing tumor immunophenotypes through IOBR, and compiling immunotherapy response data from diverse cohorts treated with immune-checkpoint blockades (ICBs).

Results TCCIA encompasses over 3,700 clinical samples obtained from 18 cohorts treated with ICBs, including PD-1/PD-L1 and CTLA-4 inhibitors, along with other treatment modalities. The database provides researchers and clinicians with a cloud-based platform that enables interactive exploration of circRNA data in the context of ICB. The platform offers a range of analytical tools, including visualization of circRNA abundance and correlation, association analysis between circRNAs and clinical variables, assessment of the tumor immune microenvironment, exploration of tumor molecular signatures, evaluation of treatment response or prognosis, and identification of altered circRNAs in immunotherapy-sensitive and resistant tumors. To illustrate the utility of TCCIA, we performed a re-analysis on a melanoma cohort with TCCIA, and found that an isoform of circTMTC3, TMTC3:+:chr12:88148287:88176319, played a significant role in predicting unfavorable

survival outcomes and treatment nonresponse.

Conclusions TCCIA represents a significant advancement over existing resources, providing

a comprehensive platform to investigate the role of circRNAs in immune oncology.

What is already known on this topic

Prior knowledge indicated that circRNAs are involved in tumor immunity and have potential

as predictive biomarkers for immunotherapy efficacy. However, there lacked a

comprehensive database that integrated circRNA profiles and immunotherapy response data,

necessitating this study.

What this study adds

This study introduces TCCIA, a database that combines circRNA profiles, immunotherapy

response data, and clinical outcomes. It provides a diverse collection of clinical samples and

an interactive platform, enabling in-depth exploration of circRNAs in the context of

checkpoint-blockade immunotherapy.

How this study might affect research, practice or policy

The findings of this study offer valuable insights into the roles of circRNAs in tumor-immune

interactions and provide a resource for researchers and clinicians in the field of immune-

oncology. TCCIA has the potential to guide personalized immunotherapeutic strategies and

contribute to future research, clinical practice, and policy decisions in checkpoint-blockade

immunotherapy and biomarker development.

Introduction

Immunotherapy has revolutionized the treatment of cancer over the past decade, emerging as a groundbreaking approach that harnesses the patient's own immune system to fight cancer. Therapies like immune checkpoint inhibitors, chimeric antigen receptor (CAR) T-cell therapy, and therapeutic vaccines aim to reinvigorate anti-tumor immunity against malignant cells [1–3]. Checkpoint inhibitors targeting programmed cell death protein-1 (PD-1), PD-1 ligand (PD-L1), and cytotoxic T-lymphocyte-associated protein 4 (CTLA-4) in particular have demonstrated remarkable clinical efficacy across diverse cancer types, highlighting immunotherapy's potential for a durable and curative response [4]. Adoptive cell transfer using engineered T cells expressing CARs has also shown great promise for blood cancers [5]. However, significant challenges remain in extending immunotherapies to larger patient populations and solid tumors. Heterogeneity in response is a major limitation – while some patients achieve long-term remission, others exhibit intrinsic resistance or relapse after an initial response [6]. This variable efficacy likely stems from immunosuppressive mechanisms within the tumor microenvironment (TME) that enable cancer cells to evade immune attack [7]. Elucidating the complex cellular and molecular interactions underlying immunotherapy resistance will be critical to unlock the full potential of immune-based cancer treatments. Reliable predictive biomarkers are also imperative to guide patient selection and combination immunotherapies tailored to each patient's TME [8].

Circular RNAs (circRNAs) have recently emerged as fascinating non-coding RNA regulators with unique covalently closed loop structures. Initially considered splicing byproducts, circRNAs are now recognized as important gene expression modulators with diverse functions [9,10]. In cancer, circRNAs have been implicated in proliferation, metastasis, and malignancy hallmarks [11]. Moreover, circRNAs are now recognized as critical regulators and potential biomarker of tumor immunity and immunotherapy response [7,12,13]. Accumulating evidence indicates circRNAs modulate TME and immunotherapy outcomes through various mechanisms in cancers like lung cancer, melanoma, colorectal cancer, and pancreatic cancer [14,15]. For example, circRNAs such as circFGFR1, circ-CPA4, and circ_0000284 facilitate immune evasion by modulating PD-L1 via sponging tumor-suppressive microRNAs [16–18]. Additionally, circRNAs including hsa_circ_0000190 [19],

circ_0020710 [20], CDR1-AS [16], and circ-UBAP2 [21] upregulate immune checkpoint proteins like PD-L1, CTLA-4 and PD-1, hampering T cell function and promoting immune evasion. Furthermore, cancer cell-derived circRNAs can reprogram intratumoral immune cells via exosomal transfer or cytokine signaling, thereby impacting facets like angiogenesis that affect immunotherapy efficacy [22–24]. CircRNAs influence various aspects of the TME, including vascularization [25], metabolism [26], hypoxia [27], macrophage polarization [28], natural killer cell cytotoxicity [17], and T cell exhaustion/apoptosis [29]. These factors can impede the efficacy of immunotherapy [15,30]. Dysregulation of circRNAs promotes immune destruction evasion and reduced immunotherapy efficacy. CircRNAs employ diverse regulatory mechanisms—from sponging miRNAs and proteins to scaffolding proteins and translating peptides [31]. While many circRNAs originate in tumors, others come from stromal and immune cells, underscoring complex multicellular regulation [32]. Exploring circRNA networks will be critical to unraveling this intricate cancer-immunity interplay. With emerging roles in tumor immunity, prognostic potential, and biomarker utility, circRNAs represent a promising new frontier in cancer immunotherapy.

Despite growing interest in circRNAs and their potential relevance in cancer immunotherapy, a comprehensive understanding of their precise functions and clinical implications remains incomplete. Existing databases have limitations in either profiling circRNAs, such as riboCIRC [33], CSCD [34] and CircNet [35] offering circRNA profiles across tissues or cancers, or curating immunotherapy cohorts, like ICBatlas [36] and TCIA [37] compiling immune infiltration and immunotherapy data across tumor types. Crucially, no resources systematically integrate comprehensive circRNA expression with multi-omics datasets including immune cell fractions, ICB types, and clinical outcomes for systematic exploration of the circRNA-immunotherapy interplay.

To address this unmet need, we developed the first-of-its-kind database, The Cancer CircRNA Immunome Atlas (TCCIA), a comprehensive database that integrates circRNA profiles, immunotherapy response data, and clinical outcomes for multiple cancer types, with the objective of providing a valuable resource for systematic exploration of the circRNA-immune axis, advancing our understanding of their functions and to facilitates discovery of potential biomarkers, therapeutic targets and clinical implications in cancer immunotherapy.

Results

Integrating circular RNAs in cancer immunotherapy

The development of the TCCIA database encompassed a comprehensive process involving data collection, preprocessing, and integration (Figure 1). In terms of data collection, we carefully curated research articles detailing cohorts treated with immune-checkpoint blockades (ICBs), utilizing the PubMed database (https://www.ncbi.nlm.nih.gov/pubmed/) for selection. We acquired raw RNA-seq datasets from several genome sequence archive repositories, including dbGaP, EGA, EMBL-EBI and GSA. We applied standardized preprocessing techniques to identify circRNAs, quantify TME/gene signatures, and incorporate clinical annotations and outcomes. This approach was aimed at improving the consistency and comparability of data across different datasets. Notably, TCCIA addresses a crucial gap by providing a unified platform with multiple tools that facilitates the exploration of circRNAs' impact on immunotherapy outcomes (Figure 1). This sets it apart from many existing databases [33–47] that predominantly focus on circRNA profiles, circRNA annotations, or immunotherapy cohorts (Table 1).

Figure 1. Overview of TCCIA.

Table 1. Comparison between TCCIA and other similar database resources [33–47].

Type	Name	Publication	Last update	Support species	Sample types (N)	Number of samples	Data sources	Characteristics
Mixed	TCCIA	Submitted	2023	Homo sapiens	cancer (5)	> 3700 cancer samples	dbGaP, EMBL-EBI, EGA, GSA	The first online interface for exploring circRNA expression and analysis in 18 immunotherapy cohorts, supporting systematic comparisons between circRNAs, clinical phenotypes, and immune signatures/infiltration at both the cohort and molecular levels, and offering the ability to explore differential gene expression between responsive and non-responsive groups for biomarker discovery.
CircRNA related	CSCD2	NAR Database Issue (2022)	2022	Homo sapiens	cancer (23)	825 tissues + 288 cell lines	ENCODE, SRA	Includes a large number of circRNAs, predicts potential miRNA–circRNA and RBP–circRNA interactions, and the potential full-length and open reading frame sequence.
	circMine	NAR Database Issue (2022)	2022	Homo sapiens	disease (87)	1107 samples	GEO	Provides online analytical functions to comprehensively evaluate the clinical and biological significance of circRNA and discover the circRNA–miRNA interaction and circRNA translatability.
	CircR2Disease2	Genomics, Proteomics & Bioinformatics (2021)	2022	5 species	disease (313)	2449 studies	PubMed	Serves as a platform to systematically investigate the roles of dysregulated circRNAs in various diseases and further explore the posttranscriptional regulatory function in diseases.
	CircNet2	NAR Database Issue (2021)	2021	Homo sapiens	cancer (37)	2732 cancer samples	TCGA, GEO, CircAtlas, MiOncoCirc	Cancer tissue-specific circRNA expression profiles and circRNA-miRNA-gene regulatory network.
	riboCIRC	Genome Biology (2021)	2021	21 species	tissue/cell line (314)	1970 samples	GEO	Provides computationally predicted ribosome-associated circRNAs and experimentally verified translated circRNAs.
	CircAtlas2	Genome Biology (2020)	2020	7 species	tissue (20)	1070 samples	SRA, NGDC, GeneBank	Integrating the most comprehensive circRNAs and their expression and functional profiles in vertebrates, which provides a foundation for investigating their biological significance.
	CircInteractome	RNA Biology (2016)	2020	Homo sapiens	tissue/cell line (34)	34 samples	circBase	Predicts the interactions between miRNAs and circRNAs with 109 RBPs, and the database also focus on IRESs and ORFs.
	TransCirc	NAR Database Issue (2020)	2020	Homo sapiens	tissues (17)	17 tissues	CircAtlas	Provides comprehensive evidence supporting the translation potential of circRNAs.
	circRic	Genome Medicine (2019)	2019	Homo sapiens	cancer (22)	935 cancer celllines	CCLE	Characterizes circRNA expression profiles; analyzes the circRNA biogenesis regulators, the effect of circRNAs on drug response, the association of circRNAs with mRNAs, proteins, and mutations, etc.
	CIRCpedia2	Genomics, Proteomics & Bioinformatics (2018)	2018	6 species	tissue (13)	185 samples	GEO, ENCODE, EMBL-EBI	Comprehensive circRNA annotation from over 180 RNA-seq datasets across six different species. Conservation analysis of circRNAs between humans and mice.
	circBase	RNA (2014)	2014	6 species	tissues/cell line (77)	77 samples	GEO, ENCODE, EMBL-EBI	Explores merged and unified circRNA data sets and the evidence supporting their expression. Provides scripts to identify known and novel circRNAs in sequencing data.
Immunotherapy related	TISMO	NAR Database Issue (2022)	2022	Mouse	cancer (19)	1518 mouse samples	GEO and In-house data	Interactive interfaces for exploring gene expression and immune infiltration, and allowing systematic comparisons between different model characteristics, and treatment and response groups.
	ICBatlas	Cancer Immunol Res (2022)	2022	Homo sapiens	cancer (9)	1515 cancer samples	GEO,ArrayExpress, TCGA, dbGaP	Transcriptome features of ICB therapy through the analysis of 1,515 ICB-treated samples from 25 studies across nine cancer types.
	ImmuCellAI	Advanced Science (2020)	2020	Homo sapiens	cancer (37)	NA	GEO, TCGA, dbGaP	The abundance of 24 immune cell types including 18 T-cell subsets, from gene expression data from self-designed approach Immune Cell Abundance Identifier.
	TCIA	Cell Reports (2017)	2017	Homo sapiens	cancer (20)	9562 cancer samples	TCGA and two immunotherapy studies	Exploration of comprehensive immunogenomic analyses of next generation sequencing data for 20 solid cancers from TCGA and other datasources.

Figure 2. Content of TCCIA. (**A**) Inclusion of cancer types, cohort and publication years in this study. Abbr. BLCA, Bladder urothelial carcinoma; HNSC, Head and neck squamous cell carcinoma; KIRC, Kidney renal clear cell carcinoma; NSCLC, non-small lung cancer; SKCM, Skin cutaneous melanoma. (**B-C**) The number of detected circRNAs, length of circRNAs, mean BSJ (back-splicing junction count) and mean CPM (counts per million) in different cancer types (**B**) or cohort (**C**). (**D**) UMAP plot of all samples, colored by cancer types.

Data summary of TCCIA

In this study, a comprehensive compilation was made, involving approximately 3700 patients from 18 immune-checkpoint blockade (ICB) cohorts [48–63] with raw RNA-seq datasets published between 2012 and 2022, encompassing 5 distinct cancer types (Figure 2A). The circRNA profiling revealed the identification of an impressive total of 281,556 circRNAs. Among these, kidney renal clear cell carcinoma (KIRC) exhibited the highest count, encompassing 159,577 circRNAs (representing 56.7% of the total). Conversely, head and neck squamous cell carcinoma (HNSC) demonstrated the lowest count, with only 680 circRNAs, constituting a mere 0.2% of the total (Figure 2B). The average lengths of these circRNAs demonstrated notable consistency across various cancer types, ranging from 21,739.64 to 23,582.26, as well as within individual cohorts, ranging from 17,041.13 to

23,112.20 (Figure 2B, C). Intriguingly, skin cutaneous melanoma (SKCM) exhibited the highest mean back-splice junction (BSJ) reads at 0.13, whereas non-small cell lung cancer (NSCLC) had the lowest mean at 0.04. Additionally, HNSC displayed the highest mean counts per million (CPM) at 1,470.6, while KIRC had the lowest mean CPM at 9.0. A comprehensive analysis encompassing all the circRNAs from the sampled datasets was visualized using a UMAP plot (Figure 2D). This visualization revealed discernible circRNA clustering patterns specific to various cancer types, highlighting the nuanced circRNA heterogeneity within human cancers and emphasizing the need for independent circRNA analysis considerations.

Web functionality of TCCIA

TCCIA introduces an array of advanced analytical tools, encompasses multifaceted functionalities to aid researchers in uncovering intricate connections and insights (Figure 1). These functionalities empower the exploration of circRNA abundance, correlation, associations with clinical variables, the tumor immune microenvironment, molecular signatures, treatment responses, and prognosis predictions, along with identifying circRNAs implicated in immunotherapy-sensitive and resistant tumor scenarios (Figure 3). The well-established exploration and analysis pipeline within the TCCIA framework is described in Figure 4. This schematic outlines the typical path that researchers follow when engaging with the platform. A more comprehensive elucidation of all fundamental modules is provided below.

Cohort Selection and Data Access. The TCCIA interface offers an intuitive approach for cohort selection and data access. The Repository Page serves as a gateway, enabling users to filter datasets based on crucial parameters such as cancer type, treatment modalities, drugs administered, and cohort sizes. Essential details pertaining to each dataset are presented in a comprehensive cohort table, facilitating informed decision-making regarding cohort selection.

Cohort/Molecule-Centered Analysis Modules. At the heart of TCCIA's capabilities lie the cohort-centered analysis modules and molecule-centred analysis modules (for analyzing

circRNAs across multiple cohorts), providing a profound lens into circRNA dynamics within specific immunotherapy cohorts. These modules encompass:

(1) *Scatter-Correlation and Heatmap-Correlation*: Researchers gain insights into circRNA correlations through scatter plots and heatmaps. These visualizations are pivotal in elucidating potential connections between circRNAs and other variables within the chosen cohort.

(2) *Group-Comparison (including simplified and comprehensive versions)*: TCCIA facilitates nuanced analysis of numeric differences in circRNA expression across multiple groups within a cohort. The dual modes of simplified and comprehensive group comparison empower researchers to unravel intricate circRNA expression patterns.

(3) *KM-Analysis and Cox-Analysis*: Survival analysis is made accessible through the KM-Analysis module, which generates Kaplan Meier survival curves among distinct variable groups. Additionally, the Cox-Analysis module allows for an in-depth examination of survival outcomes of any circRNA expression, opening avenues to prognostic evaluations.

Signature and DEG Analysis. TCCIA introduces dedicated modules for signature analysis and differential expression circRNAs (DEG) assessment. The Signature Page facilitates the investigation of associations between circRNAs and tumor microenvironment metrics using eight prominent deconvolution methods. It also allows for the examination of connections between circRNAs and 255 cancer signatures categorized into three distinct groups: TME-associated, tumor-metabolism, and tumor-intrinsic signatures. These analyses encompass a wide range of cohorts, ensuring comprehensive exploration of these relationships. The DEG Page empowers researchers to pinpoint differentially expressed circRNAs between patients who respond and those who do not respond to immunotherapy, thus unraveling the intricate web of circRNA involvement in treatment outcomes.

User Customized Configurations. Global settings within TCCIA add a layer of refinement to the user experience, allowing for customized exploration. These settings grant users control over data access and enable tailoring analyses to align with their specific research objectives. For example, by default, the platform prioritizes immunotherapy-related sample

data by filtering out samples without checkpoint immunotherapy treatment, streamlining analyses for coherent research goals. As users become acclimated to the platform, customization options foster enhanced flexibility, enabling researchers to uncover novel insights.

In essence, the web functionality of TCCIA embodies an advanced and user-centric avenue for investigating the complex roles of circRNAs in cancer immunotherapy. The integration of diverse analysis modules, coupled with a cohort-centered approach and adaptable settings, positions TCCIA as an indispensable tool for advancing our comprehension of circRNA-mediated immune responses and guiding the formulation of personalized immunotherapeutic strategies. This interactive platform stands poised to reshape the landscape of circRNA-immunotherapy research.

Figure 3. The core modules of TCCIA.

Figure 4. A standard exploration and analysis pipeline within TCCIA. (A) Users need to navigate to the Repository

page and select the clinical trial dataset of their interest. On the leftmost panel, users can filter the dataset based on

various parameters. **(B)** Users can choose an appropriate analysis strategy according to their needs (core analysis steps

are listed on the rightmost panel). The analysis consists of four steps: I. Select the dataset. II. Choose the CircRNA ID or

Host Gene of interest. III. Adjust various parameters such as Test Type, Color Selection, etc. IV. Click the Submit button to

obtain the analysis results. **(C)** For users aiming for in-depth analysis, a more personalized clinical data filtering suite is

provided. Users can perform patient selection based on different features such as Sex, Response, etc., and redraw the

plots. **(D)** Users have the option to save the result images in PDF or PNG format, with the desired dimensions.

Case study: validating circTMTC3 prediction efficacy in Gide et al. melanoma cohort

ICB therapies targeting PD-1 and CTLA-4 have significantly transformed the field of oncology,

particularly in the treatment of metastatic melanoma. However, it is important to note that only

a limited number of melanoma patients experience positive outcomes from these

immunotherapies. Consequently, there is a pressing need to identify predictive biomarkers

that can guide precision oncology.

In a study conducted by Dong et al. [13], it was observed that melanoma patients from

the study (cohort ID: PRJEB23709) by Gide et al. [54], who exhibited high expression of

circTMTC3, experienced poorer survival outcomes and demonstrated reduced treatment

responsiveness compared to those with low circTMTC3 expression. To illustrate the

functionality of the TCCIA, here, we performed a re-analysis on the same cohort. Figure 5A

provides an overview of the analysis panel used to assess the association between a circRNA and patient survival. Additionally, Figure 5B clearly demonstrates that high levels of circTMTC3 are predictive of poor overall survival ($HR_{low\ vs\ high}$=0.47, 95% CI: 0.24-0.92, P=0.02). Upon closer examination of the different isoforms of circRNAs derived from TMTC3, it was found that only TMTC3:+:chr12:88148287:88176319 was abundant in this patient cohort and played a significant role in predicting unfavorable survival outcomes (Figure 5C, $HR_{low\ vs\ high}$=0.42, 95% CI: 0.2-0.9, P=0.008). The remaining isoforms, such as TMTC3:+:chr12:88160113:88190622, did not exhibit the same predictive ability (Figure 5D). Furthermore, employing three different approaches to assess differential expression of circRNAs between checkpoint immunotherapy responders and non-responders, it was consistently discovered that TMTC3:+:chr12:88148287:88176319 was the sole isoform of circTMTC3 that displayed significant upregulation in non-responding patients (Figure 5E-G).

Discussion

Circular RNAs, emerging as pivotal gene expression regulators, exert diverse functions across biological processes, with substantial clinical research potential [64]. However, the existing landscape of circRNA-related tumor-immune checkpoint research falls short of satisfying the burgeoning need for insights. This limitation is compounded by constraints stemming from sample quantity and diversity, which subsequently curtail the generalizability of research findings due to geographical, racial, and tumor-specific factors. Consequently, given this evolving landscape, the urgency and significance of developing a CircRNA tool for preliminary data mining have become increasingly pronounced. TCCIA (the Cancer CircRNA Immunome Atlas) an integrated online platform, building upon datasets from four genome sequence archive repositories, encompassing around 3,700 cancer samples, spanning five cancer types and 18 checkpoint-blockade immunotherapy cohorts, incorporating over 280,000 circRNA expression profiles, 255 established cancer signatures, and TME decomposition results from eight immune infiltration algorithms. TCCIA emphasizes user-friendly visual presentations, eliminating the need for intricate programming skills. Moreover, the platform offers a range of customizable visualization options, ensuring adaptability to user needs. Notably, TCCIA is readily accessible without mandatory registration or login,

potentially rendering it an economical and efficient solution for both researchers and clinical

practitioners.

Figure 5. circTMTC3 predicts poor survival and non-response to checkpoint immunotherapy in the melanoma

cohort of Gide et al. (**A**) Analysis panel for analyzing and visualizing associations between circRNAs and patient survival.

Here, we present the data and analysis settings used to generate the plot shown in (**C**). (**B**) CircTMTC3 predicts poor survival.

(**C**) A circRNA isoform of circTMTC3, TMTC3:+:chr12:88148287:88176319, predicts poor survival. (**D**) A circRNA isoform

of circTMTC3, TMTC3:+:chr12:88160113:88190622, predicts poor survival. (**E-G**) Volcano plots showing differential

expressed circRNAs between checkpoint immunotherapy response and nonresponse patients using the approaches: (**E**)

Limma Voom, (**F**) DESeq2, and (**G**) edgeR.

In this manuscript, we delineate the data sources, collection and standardization

processes, TCCIA's functionalities, website analysis modules, and provide a step-by-step

guide to its operation. To further illustrate TCCIA, we present a concrete example of the

circTMTC3 as a molecular marker for melanoma and confirm its specific isoform, TMTC3:+:chr12:88148287:88176319, in playing a major role in prognosticating unfavorable survival outcomes and non-response to treatment.

While TCCIA has various advantages and uniqueness, there are still some limitations. In terms of data, although TCCIA includes data from five different types of cancer, the sample size for head and neck cancer remains sparse. Recently, the application of immunotherapy has been expanding to more cancer types, such as digestive system tumors. However, it is important to note that our TCCIA does not currently cover these types of tumors. The main reason for this is that current clinical genomics research of such cancer types primarily focuses on the DNA level, specifically on whole exome sequencing and targeted sequencing [65]. As a result, there are few RNA-seq datasets to infer the presence of circRNAs. Another reason is that some raw RNA-seq datasets are difficult to access due to restricted availability, e.g., Carroll et al. study [66]. In terms of circRNA abundance, there are significant differences between different cohorts, batch effects may exist in different cohorts, and there is also strong heterogeneity within the same tumor type, so users need to be cautious when performing cross-dataset analysis and comparison. In terms of functionality, some databases provide the characteristics of studying circRNA-miRNA-gene regulatory networks (such as CSCD2 [34], circMine [38], and CircNet2 [35]), which have not been considered in TCCIA with two reasons: First, further experimental validation is generally required to confirm the authenticity of detected circRNAs. In the recent large-scale circRNA detection benchmark study [67], Vromman et al. recommended using qPCR+ Ribonuclease R or qPCR+amplicon sequencing for circRNA validation. Therefore, it is recommended to incorporate these experimental validation methods to ensure the accuracy of circRNA detection. Second, the focus of our study is on integrating circRNA profiles and clinical outcomes in cancer patients treated with immunotherapy. However, instead of duplicating efforts, users can leverage already well-established and high-quality circRNA databases to complete other circRNA annotation and analysis explorations. By linking to these databases (a widget is provided in the footer of the TCCIA website), users can access comprehensive circRNA information and utilize existing tools for further analysis, optimizing the accuracy and efficiency of circRNA annotation and deep investigation.

Looking ahead, we aim to continually update TCCIA by incorporating more circRNA data from diverse immunotherapy cohorts and introducing new functionalities based on user feedback. In summary, the distinctive features, analytical capabilities, and potential for growth position it as a pivotal tool in advancing our understanding of circRNAs in tumor immunity and in shaping development of personalized immunotherapy strategies guided by circRNA.

Methods

Data collection

To conduct a systematic search, we utilized PubMed (https://www.ncbi.nlm.nih.gov/pubmed/) to search for articles related to Bulk RNAseq data from solid cancer patients treated with immune checkpoint blockers (ICB). The search expression used was "(ICB [Title/Abstract] OR PD-1[Title/Abstract] OR PD-L1[Title/Abstract] OR CTLA-4[Title/Abstract]) AND (rnaseq[Title/Abstract] OR rna-seq[Title/Abstract] OR rna-sequencing[Title/Abstract] OR rna seq[Title/Abstract] OR rna sequencing[Title/Abstract])". No filters were applied, and there were no restrictions on language or geographic region. Peer-reviewed publications, preprints, and press releases were considered for inclusion. To obtain raw RNA sequencing data, we submitted requests to the Database of Genotypes and Phenotypes (dbGaP) (https://dbgap.ncbi.nlm.nih.gov/), the European Genome-phenome Archive (EGA) (https://ega-archive.org/), and the Genome Sequence Archive (GSA) (https://ngdc.cncb.ac.cn/gsa/) of the National Genomics Data Center (NGDC) after receiving approval from the Data Access Committee (DAC). However, it is important to note that not all raw RNAseq datasets were accessible and available for use. In total, we collected 16 studies [48–63] related to checkpoint immunotherapy that provided raw RNAseq datasets. We gathered relevant clinical data from publications and clinical meta documents associated with these RNAseq datasets. Additionally, we extracted information on the cohorts' fundamental characteristics, such as sample size, treatment methods, and specific drugs used, based on the abstracts. It should be noted that, for Patil et al. study, two clinical cohorts were included; and for the study identified as PHS000452, the two patient subgroups had distinct drug treatments and clinical annotations. Hence, we treated them as two separate cohorts during the analysis. For the TCCIA project, CircRNA profiles, immunotherapy response, and clinical

benefits were analyzed for five cancer types. This analysis included over 3,700 clinical samples from 18 cohorts treated with immune-checkpoint blockades (ICBs) such as PD-1/PD-L1 and CTLA-4 inhibitors, as well as other treatments. The analysis considered both pre-treatment and on-treatment responses.

CircRNA identification and differential expression analysis

We aligned the raw RNA sequencing data to the human genome hg38 using STAR [68]. Next, we utilized CIRCexplorer2 [69] to identify, parse, and annotate circRNA junctions within each sample. These identified junctions were then analyzed for differential expression using Limma Voom [70], edgeR [71], and DESeq2 [72], enabling a comparison between patients who responded to checkpoint immunotherapy and those who did not.

TME decomposition and cancer gene signature estimation

We employed IOBR [73] for TME decomposition and the scoring of cancer gene signatures. IOBR seamlessly integrates eight widely-used open-source deconvolution methods, including CIBERSORT [74], ESTIMATE [75], quanTIseq [76], TIMER [77], IPS [37], MCPCounter [78], xCell [79], and EPIC [80]. Furthermore, IOBR incorporates a comprehensive compilation of 255 established cancer signatures. These diverse signatures are organized into three distinct categories: TME-associated, tumor-metabolism, and tumor-intrinsic signatures.

TCCIA implementation

The TCCIA database is developed as a Web application leveraging R Shiny (https://shiny.posit.co/) and built using the golem framework (https://github.com/ThinkR-open/golem) to achieve optimization. TCCIA, is developed solely for research purposes and does not utilize any cookies or collect any personal identifiable information. TCCIA is free available in https://tccia.zmu-zhoulab.com/ and https://shiny.hiplot.cn/TCCIA.

Statistical analysis

We performed Kaplan-Meier survival analysis to generate and compare survival curves. The

log-rank test was used for comparison. We also conducted multivariate survival analysis using the Cox regression model. All reported *P*-values are two-tailed, and a significance level of p≤0.05 was used unless otherwise specified. All statistical analyses and visualization were conducted using R v4.2.0.

Patient consent for publication

Not applicable.

Data availability

All relevant data reported in the study can be found in the article or on the TCCIA website. Please note that access to the raw RNA-seq datasets is not provided. For any other data requests, please contact the leader of this project, Jian-Guo Zhou.

Funding

This work was supported by the National Natural Science Foundation of China (Grant No. 81660512, 81472594, 81770781, 82270825), Chunhui program of the Chinese Ministry of Education (Grant No. HZKY20220231), the Natural Science Foundation of Guizhou Province (Grant No. ZK2021-YB435), Guangdong Basic and Applied Basic Research Foundation (Grant No. 2021A1515011743), Youth Talent Project of Guizhou Provincial Department of Education (Grant No. QJJ2022-224), China Postdoctoral Science Foundation (Grant No. 2021M703733), China Lung Cancer Immunotherapy Research Project, and Special funds for innovation in Hunan Province (Grant No. 2020SK2062).

Contributions

SW: Conceptualization, software, methodology, formal analysis, writing original draft, review and editing. YX: Software, methodology, formal analysis, visualization, writing original draft. YZ: Software, formal analysis, visualization, writing original draft. HW: Conceptualization, writing original draft, review and editing. JL and USG: Resources, review and editing. MC: Visualization, review and editing. PL, YHL, MH and BF: Review and editing. XL, QZ and HM: Supervision, resources, funding acquisition. JGZ: Conceptualization, methodology,

resources, supervision, funding acquisition, project administration, writing original draft,

writing–review and editing.

Conflict of interest

None were declared.

Acknowledgments

We thank Dr. Jianming Zeng (University of Macau), and all the members of his bioinformatics

team, biotrainee, for generously sharing their experience and codes. We thank Juan Zhang

(Echo Biotech Co., Ltd., Beijing, China) for her help in the data pre-processing.

References

Sharma P, Allison JP. Immune Checkpoint Targeting in Cancer Therapy: Toward Combination Strategies with Curative Potential. *Cell* 2015;**161**:205–14. doi:10.1016/j.cell.2015.03.030

Maude SL, Laetsch TW, Buechner J, *et al.* Tisagenlecleucel in Children and Young Adults with B-Cell Lymphoblastic Leukemia. *N Engl J Med* 2018;**378**:439–48. doi:10.1056/NEJMoa1709866

Guo C, Manjili MH, Subjeck JR, *et al.* Chapter Seven - Therapeutic Cancer Vaccines: Past, Present, and Future. In: Tew KD, Fisher PB, eds. *Advances in Cancer Research*. Academic Press 2013. 421–75. doi:10.1016/B978-0-12-407190-2.00007-1

Topalian SL, Hodi FS, Brahmer JR, *et al.* Safety, Activity, and Immune Correlates of Anti–PD-1 Antibody in Cancer. *N Engl J Med* 2012;**366**:2443–54. doi:10.1056/NEJMoa1200690

Neelapu SS, Locke FL, Bartlett NL, *et al.* Axicabtagene Ciloleucel CAR T-Cell Therapy in Refractory Large B-Cell Lymphoma. *N Engl J Med* 2017;**377**:2531–44. doi:10.1056/NEJMoa1707447

Sharma P, Hu-Lieskovan S, Wargo JA, *et al.* Primary, Adaptive, and Acquired Resistance to Cancer Immunotherapy. *Cell* 2017;**168**:707–23. doi:10.1016/j.cell.2017.01.017

Vinay DS, Ryan EP, Pawelec G, *et al.* Immune evasion in cancer: Mechanistic basis and therapeutic strategies. *Semin Cancer Biol* 2015;**35**:S185–98. doi:10.1016/j.semcancer.2015.03.004

Gibney GT, Weiner LM, Atkins MB. Predictive biomarkers for checkpoint inhibitor-based immunotherapy. *Lancet Oncol* 2016;**17**:e542–51. doi:10.1016/S1470-2045(16)30406-5

Salzman J. Circular RNA Expression: Its Potential Regulation and Function. *Trends Genet* 2016;**32**:309–16. doi:10.1016/j.tig.2016.03.002

Liu C-X, Chen L-L. Circular RNAs: Characterization, cellular roles, and applications. *Cell* 2022;**185**:2016–34. doi:10.1016/j.cell.2022.04.021

Kristensen LS, Hansen TB, Venø MT, *et al.* Circular RNAs in cancer: opportunities and challenges in the field. *Oncogene* 2018;**37**:555–65. doi:10.1038/onc.2017.361

Zhou J-G, Liang R, Wang H-T, *et al.* Identification and characterization of circular RNAs as novel putative biomarkers to predict anti-PD-1 monotherapy response in metastatic melanoma patients – Knowledge from two independent international studies. *Neoplasia* 2023;**37**:100877. doi:10.1016/j.neo.2023.100877

Dong Y, Gao Q, Chen Y, *et al.* Identification of CircRNA signature associated with tumor immune infiltration to predict therapeutic efficacy of immunotherapy. *Nat Commun* 2023;**14**:2540. doi:10.1038/s41467-023-38232-y

Pandey PR, Young KH, Kumar D, *et al.* RNA-mediated immunotherapy regulating tumor immune microenvironment: next wave of cancer therapeutics. *Mol Cancer* 2022;**21**:58. doi:10.1186/s12943-022-01528-6

Guan L, Hao Q, Shi F, *et al.* Regulation of the tumor immune microenvironment by cancer-derived circular RNAs. *Cell Death Dis* 2023;**14**:1–12. doi:10.1038/s41419-023-05647-w

Tanaka E, Miyakawa Y, Kishikawa T, *et al.* Expression of circular RNA CDR1-AS in colon cancer cells increases cell surface PD-L1 protein levels. *Oncol Rep* 2019;**42**:1459–66. doi:10.3892/or.2019.7244

Zhang P-F, Gao C, Huang X-Y, *et al.* Cancer cell-derived exosomal circUHRF1 induces natural killer cell exhaustion and may cause resistance to anti-PD1 therapy in hepatocellular carcinoma.

Mol Cancer 2020;**19**:110. doi:10.1186/s12943-020-01222-5

18   Hong W, Xue M, Jiang J, *et al.* Circular RNA circ-CPA4/ let-7 miRNA/PD-L1 axis regulates
cell growth, stemness, drug resistance and immune evasion in non-small cell lung cancer
(NSCLC). *J Exp Clin Cancer Res* 2020;**39**:149. doi:10.1186/s13046-020-01648-1

19   Luo Y-H, Yang Y-P, Chien C-S, *et al.* Circular RNA hsa_circ_0000190 Facilitates the
Tumorigenesis and Immune Evasion by Upregulating the Expression of Soluble PD-L1 in Non-
Small-Cell Lung Cancer. *Int J Mol Sci* 2021;**23**:64. doi:10.3390/ijms23010064

20   Wei C-Y, Zhu M-X, Lu N-H, *et al.* Circular RNA circ_0020710 drives tumor progression and
immune evasion by regulating the miR-370-3p/CXCL12 axis in melanoma. *Mol Cancer*
2020;**19**:84. doi:10.1186/s12943-020-01191-9

21   Zhao R, Ni J, Lu S, *et al.* CircUBAP2-mediated competing endogenous RNA network
modulates tumorigenesis in pancreatic adenocarcinoma. *Aging* 2019;**11**:8484–501.
doi:10.18632/aging.102334

22   Ou Z-L, Luo Z, Wei W, *et al.* Hypoxia-induced shedding of MICA and HIF1A-mediated
immune escape of pancreatic cancer cells from NK cells: role of circ_0000977/miR-153 axis.
RNA Biol 2019;**16**:1592–603. doi:10.1080/15476286.2019.1649585

23   Wang X, Sheng W, Xu T, *et al.* CircRNA hsa_circ_0110102 inhibited macrophage activation
and hepatocellular carcinoma progression via miR-580-5p/PPARα/CCL2 pathway. *Aging*
2021;**13**:11969–87. doi:10.18632/aging.202900

24   Li Z, Huang C, Bao C, *et al.* Exon-intron circular RNAs regulate transcription in the nucleus.
Nat Struct Mol Biol 2015;**22**:256–64. doi:10.1038/nsmb.2959

25   Liu C, Yao M-D, Li C-P, *et al.* Silencing Of Circular RNA-ZNF609 Ameliorates Vascular
Endothelial Dysfunction. *Theranostics* 2017;**7**:2863–77. doi:10.7150/thno.19353

26   Di Timoteo G, Dattilo D, Centrón-Broco A, *et al.* Modulation of circRNA Metabolism by m6A
Modification. *Cell Rep* 2020;**31**:107641. doi:10.1016/j.celrep.2020.107641

27   Jiao B, Liu S, Zhao H, *et al.* Hypoxia-responsive circRNAs: A novel but important participant
in non-coding RNAs ushered toward tumor hypoxia. *Cell Death Dis* 2022;**13**:666.
doi:10.1038/s41419-022-05114-y

28   Pan Z, Zhao R, Li B, *et al.* EWSR1-induced circNEIL3 promotes glioma progression and
exosome-mediated macrophage immunosuppressive polarization via stabilizing IGF2BP3. *Mol*
Cancer 2022;**21**:16. doi:10.1186/s12943-021-01485-6

29   Yang C, Wu S, Mou Z, *et al.* Exosome-derived circTRPS1 promotes malignant phenotype and
CD8+ T cell exhaustion in bladder cancer microenvironments. *Mol Ther J Am Soc Gene Ther*
2022;**30**:1054–70. doi:10.1016/j.ymthe.2022.01.022

30   Zheng Y, Ren S, Zhang Y, *et al.* Circular RNA circWWC3 augments breast cancer progression
through promoting M2 macrophage polarization and tumor immune escape via regulating the
expression and secretion of IL-4. *Cancer Cell Int* 2022;**22**:264. doi:10.1186/s12935-022-02686-
9

31   Pamudurti NR, Bartok O, Jens M, *et al.* Translation of CircRNAs. *Mol Cell* 2017;**66**:9-21.e7.
doi:10.1016/j.molcel.2017.02.021

32   Fang Z, Jiang C, Li S. The Potential Regulatory Roles of Circular RNAs in Tumor Immunology
and Immunotherapy. *Front Immunol*
2021;**11**.https://www.frontiersin.org/articles/10.3389/fimmu.2020.617583 (accessed 11 Aug
2023).

33 Li H, Xie M, Wang Y, *et al.* riboCIRC: a comprehensive database of translatable circRNAs.
Genome Biol 2021;**22**:79. doi:10.1186/s13059-021-02300-7

34 Feng J, Chen W, Dong X, *et al.* CSCD2: an integrated interactional database of cancer-specific
circular RNAs. *Nucleic Acids Res* 2022;**50**:D1179–83. doi:10.1093/nar/gkab830

35 Chen Y, Yao L, Tang Y, *et al.* CircNet 2.0: an updated database for exploring circular RNA
regulatory networks in cancers. *Nucleic Acids Res* 2022;**50**:D93–101.
doi:10.1093/nar/gkab1036

36 Yang M, Miao Y-R, Xie G-Y, *et al.* ICBatlas: A comprehensive resource for depicting immune
checkpoint blockade therapy characteristics from transcriptome profiles. *Cancer Immunol Res*
2022;:CIR-22-0249. doi:10.1158/2326-6066.CIR-22-0249

37 Charoentong P, Finotello F, Angelova M, *et al.* Pan-cancer Immunogenomic Analyses Reveal
Genotype-Immunophenotype Relationships and Predictors of Response to Checkpoint
Blockade. *Cell Rep* 2017;**18**:248–62. doi:10.1016/j.celrep.2016.12.019

38 Zhang W, Liu Y, Min Z, *et al.* circMine: a comprehensive database to integrate, analyze and
visualize human disease–related circRNA transcriptome. *Nucleic Acids Res* 2022;**50**:D83–92.
doi:10.1093/nar/gkab809

39 Fan C, Lei X, Tie J, *et al.* CircR2Disease v2.0: An Updated Web Server for Experimentally
Validated circRNA–disease Associations and Its Application. *Genomics Proteomics*
Bioinformatics 2022;**20**:435–45. doi:10.1016/j.gpb.2021.10.002

40 Wu W, Ji P, Zhao F. CircAtlas: an integrated resource of one million highly accurate circular
RNAs from 1070 vertebrate transcriptomes. *Genome Biol* 2020;**21**:101. doi:10.1186/s13059-
020-02018-y

41 Dudekula DB, Panda AC, Grammatikakis I, *et al.* CircInteractome: A web tool for exploring
circular RNAs and their interacting proteins and microRNAs. *RNA Biol* 2016;**13**:34–42.
doi:10.1080/15476286.2015.1128065

42 Huang W, Ling Y, Zhang S, *et al.* TransCirc: an interactive database for translatable circular
RNAs based on multi-omics evidence. *Nucleic Acids Res* 2021;**49**:D236–42.
doi:10.1093/nar/gkaa823

43 Ruan H, Xiang Y, Ko J, *et al.* Comprehensive characterization of circular RNAs in ~ 1000
human cancer cell lines. *Genome Med* 2019;**11**:55. doi:10.1186/s13073-019-0663-5

44 Dong R, Ma X-K, Li G-W, *et al.* CIRCpedia v2: An Updated Database for Comprehensive
Circular RNA Annotation and Expression Comparison. *Genomics Proteomics Bioinformatics*
2018;**16**:226–33. doi:10.1016/j.gpb.2018.08.001

45 Glažar P, Papavasileiou P, Rajewsky N. circBase: a database for circular RNAs. *RNA*
2014;**20**:1666–70. doi:10.1261/rna.043687.113

46 Zeng Z, Wong CJ, Yang L, *et al.* TISMO: syngeneic mouse tumor database to model tumor
immunity and immunotherapy response. *Nucleic Acids Res* 2022;**50**:D1391–7.
doi:10.1093/nar/gkab804

47 Miao Y, Zhang Q, Lei Q, *et al.* ImmuCellAI: A Unique Method for Comprehensive T-Cell
Subsets Abundance Prediction and its Application in Cancer Immunotherapy. *Adv Sci*
2020;**7**:1902880. doi:10.1002/advs.201902880

48 Patil NS, Nabet BY, Müller S, *et al.* Intratumoral plasma cells predict outcomes to PD-L1
blockade in non-small cell lung cancer. *Cancer Cell* 2022;**40**:289-300.e4.
doi:10.1016/j.ccell.2022.02.002

49 Bellmunt J, Hussain M, Gschwend JE, *et al.* Adjuvant atezolizumab versus observation in
muscle-invasive urothelial carcinoma (IMvigor010): a multicentre, open-label, randomised,
phase 3 trial. *Lancet Oncol* 2021;**22**:525–37. doi:10.1016/S1470-2045(21)00004-8

50 Cui C, Xu C, Yang W, *et al.* Ratio of the interferon-γ signature to the immunosuppression
signature predicts anti-PD-1 therapy response in melanoma. *Npj Genomic Med* 2021;**6**:7.
doi:10.1038/s41525-021-00169-w

51 Liu S, Knochelmann HM, Lomeli SH, *et al.* Response and recurrence correlates in individuals
treated with neoadjuvant anti-PD-1 therapy for resectable oral cavity squamous cell carcinoma.
Cell Rep Med 2021;**2**:100411. doi:10.1016/j.xcrm.2021.100411

52 Grasso CS, Tsoi J, Onyshchenko M, *et al.* Conserved Interferon-γ Signaling Drives Clinical
Response to Immune Checkpoint Blockade Therapy in Melanoma. *Cancer Cell* 2020;**38**:500-
515.e3. doi:10.1016/j.ccell.2020.08.005

53 Motzer RJ, Banchereau R, Hamidi H, *et al.* Molecular Subsets in Renal Cancer Determine
Outcome to Checkpoint and Angiogenesis Blockade. *Cancer Cell* 2020;**38**:803-817.e4.
doi:10.1016/j.ccell.2020.10.011

54 Gide TN, Quek C, Menzies AM, *et al.* Distinct Immune Cell Populations Define Response to
Anti-PD-1 Monotherapy and Anti-PD-1/Anti-CTLA-4 Combined Therapy. *Cancer Cell*
2019;**35**:238-255.e6. doi:10.1016/j.ccell.2019.01.003

55 Mariathasan S, Turley SJ, Nickles D, *et al.* TGFβ attenuates tumour response to PD-L1
blockade by contributing to exclusion of T cells. *Nature* 2018;**554**:544–8.
doi:10.1038/nature25501

56 McDermott DF, Huseni MA, Atkins MB, *et al.* Clinical activity and molecular correlates of
response to atezolizumab alone or in combination with bevacizumab versus sunitinib in renal
cell carcinoma. *Nat Med* 2018;**24**:749–57. doi:10.1038/s41591-018-0053-3

57 Miao D, Margolis CA, Gao W, *et al.* Genomic correlates of response to immune checkpoint
therapies in clear cell renal cell carcinoma. *Science* 2018;**359**:801–6.
doi:10.1126/science.aan5951

58 Auslander N, Zhang G, Lee JS, *et al.* Robust prediction of response to immune checkpoint
blockade therapy in metastatic melanoma. *Nat Med* 2018;**24**:1545–9. doi:10.1038/s41591-018-
0157-9

59 Riaz N, Havel JJ, Makarov V, *et al.* Tumor and Microenvironment Evolution during
Immunotherapy with Nivolumab. *Cell* 2017;**171**:934-949.e16. doi:10.1016/j.cell.2017.09.028

60 Hugo W, Zaretsky JM, Sun L, *et al.* Genomic and Transcriptomic Features of Response to Anti-
PD-1 Therapy in Metastatic Melanoma. *Cell* 2016;**165**:35–44. doi:10.1016/j.cell.2016.02.065

61 Snyder A, Makarov V, Merghoub T, *et al.* Genetic Basis for Clinical Response to CTLA-4
Blockade in Melanoma. *N Engl J Med* 2014;**371**:2189–99. doi:10.1056/NEJMoa1406498

62 Tumeh PC, Harview CL, Yearley JH, *et al.* PD-1 blockade induces responses by inhibiting
adaptive immune resistance. *Nature* 2014;**515**:568–71. doi:10.1038/nature13954

63 Berger MF, Hodis E, Heffernan TP, *et al.* Melanoma genome sequencing reveals frequent
PREX2 mutations. *Nature* 2012;**485**:502–6. doi:10.1038/nature11071

64 Pisignano G, Michael DC, Visal TH, *et al.* Going circular: history, present, and future of
circRNAs in cancer. *Oncogene* Published Online First: 16 August 2023. doi:10.1038/s41388-
023-02780-w

65 Horak P, Fröhling S, Glimm H. Integrating next-generation sequencing into clinical oncology:

strategies, promises and pitfalls. *ESMO Open* 2016;**1**:e000094. doi:10.1136/esmoopen-2016-
000094

66 Carroll TM, Chadwick JA, Owen RP, *et al.* Tumor monocyte content predicts
immunochemotherapy outcomes in esophageal adenocarcinoma. *Cancer Cell* 2023;**41**:1222-
1241.e7. doi:10.1016/j.ccell.2023.06.006

67 Vromman M, Anckaert J, Bortoluzzi S, *et al.* Large-scale benchmarking of circRNA detection
tools reveals large differences in sensitivity but not in precision. *Nat Methods* 2023;**20**:1159–69.
doi:10.1038/s41592-023-01944-6

68 Dobin A, Davis CA, Schlesinger F, *et al.* STAR: ultrafast universal RNA-seq aligner.
Bioinformatics 2013;**29**:15–21. doi:10.1093/bioinformatics/bts635

69 Zhang X-O, Dong R, Zhang Y, *et al.* Diverse alternative back-splicing and alternative splicing
landscape of circular RNAs. *Genome Res* 2016;**26**:1277–87. doi:10.1101/gr.202895.115

70 Law CW, Chen Y, Shi W, *et al.* voom: precision weights unlock linear model analysis tools for
RNA-seq read counts. *Genome Biol* 2014;**15**:R29. doi:10.1186/gb-2014-15-2-r29

71 Robinson MD, McCarthy DJ, Smyth GK. edgeR : a Bioconductor package for differential
expression analysis of digital gene expression data. *Bioinformatics* 2010;**26**:139–40.
doi:10.1093/bioinformatics/btp616

72 Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. *Genome Biol* 2014;**15**:550. doi:10.1186/s13059-014-0550-8

73 Zeng D, Ye Z, Shen R, *et al.* IOBR: Multi-Omics Immuno-Oncology Biological Research to
Decode Tumor Microenvironment and Signatures. *Front Immunol*
2021;**12**.https://www.frontiersin.org/articles/10.3389/fimmu.2021.687975 (accessed 20 Aug
2023).

74 Newman AM, Liu CL, Green MR, *et al.* Robust enumeration of cell subsets from tissue
expression profiles. *Nat Methods* 2015;**12**:453–7. doi:10.1038/nmeth.3337

75 Yoshihara K, Shahmoradgoli M, Martínez E, *et al.* Inferring tumour purity and stromal and
immune cell admixture from expression data. *Nat Commun* 2013;**4**:2612.
doi:10.1038/ncomms3612

76 Finotello F, Mayer C, Plattner C, *et al.* Molecular and pharmacological modulators of the tumor
immune contexture revealed by deconvolution of RNA-seq data. *Genome Med* 2019;**11**:34.
doi:10.1186/s13073-019-0638-6

77 Li B, Liu JS, Liu XS. Revisit linear regression-based deconvolution methods for tumor gene
expression data. *Genome Biol* 2017;**18**:127. doi:10.1186/s13059-017-1256-5

78 Becht E, Giraldo NA, Lacroix L, *et al.* Estimating the population abundance of tissue-
infiltrating immune and stromal cell populations using gene expression. *Genome Biol*
2016;**17**:218. doi:10.1186/s13059-016-1070-5

79 Aran D, Hu Z, Butte AJ. xCell: digitally portraying the tissue cellular heterogeneity landscape.
Genome Biol 2017;**18**:220. doi:10.1186/s13059-017-1349-1

80 Racle J, de Jonge K, Baumgaertner P, *et al.* Simultaneous enumeration of cancer and immune
cell types from bulk tumor gene expression data. *eLife* 2017;**6**:e26476. doi:10.7554/eLife.26476

Database Construction

Data Collection

db GaP genotypes and phenotypes
EMBL-EBI
European Genome-Phenome Archive
GSA Genome Sequence Archive

Integration of Data

Standardized preprocessing

fastq → CircRNA expression profiles ⟷ Immunotherapy response & Clinical outcomes

GRCh38

Exploration and Analysis Tools

ⓘ **Repository**
▤ **Signature**
⧗ **Molecule-centred Analysis**
👥 **Correlation**

👥 **Grouping**

◈ **Cox-Analysis**

📈 **Distribution**

🔥 **DEG**

CircRNA profiles

TCCIA
The Cancer CircRNA Immunome Atlas

Immunotherapy response data/Clinical outcomes across five cancer types

TME/gene signatures

A Repository Selection

Filter Cohort → Select Cohort → Click

B Analysis Controls

I.   **Select datasets**

II. **Select circRNA ID or Host Gene**

III. **Adjust parameters for analysis**

IV. **Click to analyze**

Notion:

1. **Homepage Overview**
2. **Repository Selection**
3. **Cohort-centred Analysis**
   - Scatter-Correlation
   - Heatmap-Correlation
   - Group-Comparison
   - KM-Analysis
   - Cox-Analysis
   - Cohort Filter
4. **Molecular-centred Analysis**
   - Distribution of circRNA
   - Cox analysis for circRNA
5. **Signature Analysis**
6. **DEG Analysis**
7. **Demo Output**

C Filter (Optional)

Filter patients and redraw

D Save Plots

I.   **Change sizes**

II. **Choose format**

III. **Click**