TCCIA: A Comprehensive Resource for Exploring CircRNA in Cancer Immunotherapy

Shixiang Wang[1,2,#], Yi Xiong[3,4,5,#], Yihao Zhang[3,4,5,#], Haitao Wang[6,^], Minjun Chen[1], Jianfeng Li[7], Peng Luo[8], Yung-Hung Luo[9,10], Markus Hecht[11], Benjamin Frey[12], Udo S Gaipl[12], Xuejun Li[3,4,5,*], Qi Zhao[2,*], Hu Ma[1,*], Jian-Guo Zhou[1,12*]

Affiliations of authors:

[1] Department of Oncology, The Second Affiliated Hospital of Zunyi Medical University, Zunyi, 563000, P. R. China.

[2] State Key Laboratory of Oncology in South China, Guangdong Key Laboratory of Nasopharyngeal Carcinoma Diagnosis and Therapy, Guangdong Provincial Clinical Research Center for Cancer, Sun Yat-sen University Cancer Center, Guangzhou 510060, P. R. China.

[3] Xiangya School of Medicine, Central South University, Changsha, 410013, P. R. China.

[4] Hunan International Scientific and Technological Cooperation Base of Brain Tumor Research, Xiangya Hospital, Central South University, Changsha, 410008, P. R. China.

5 Department of Neurosurgery, Xiangya Hospital, Central South University, Changsha, 410008, P. R. China.

[6] Center for Precision Medicine Research and Training, Faculty of Health Sciences, University of Macau, Macau SAR, 999087, P. R. China.

[7] State Key Laboratory of Medical Genomics, Shanghai Institute of Hematology, National Research Center for Translational Medicine, Rui-Jin Hospital, Shanghai Jiao Tong University, School of Medicine, Shanghai, 200025, P. R. China.

[8] Department of Oncology, Zhujiang Hospital, Southern Medical University, Guangzhou, 510091, P. R. China.

[9] Department of Chest Medicine, Taipei Veterans General Hospital, Taipei, 11217, Taiwan (R.O.C.).

[10] School of Medicine, College of Medicine, National Yang Ming Chiao Tung University, Taipei, 112304, Taiwan (R.O.C.)

[11] Department of Radiotherapy and Radiation Oncology, Saarland University Medical Center, Homburg, 66421, Germany.

[12] Translational Radiobiology, Department of Radiation Oncology, Universitätsklinikum

34   Erlangen, Friedrich-Alexander-Universität Erlangen-Nürnberg, Erlangen, 91054, Germany.

35   ^ Current address: Thoracic Surgery Branch, Center for Cancer Research, NCI, NIH,
36   Bethesda, MD 20892, United States
37

Co-first author.

* Corresponding author.

Correspondence: Jian-Guo Zhou (jianguo.zhou@zmu.edu.cn), Hu Ma (mahuab@163.com),

Qi Zhao (zhaoqi@sysucc.org.cn), Xuejun Li (lxjneuro@csu.edu.cn)

Lead contact: Jian-Guo Zhou

Keywords: Circular RNA, Database, Checkpoint immunotherapy, Biomarker

Abstract

Background Immunotherapies targeting immune checkpoints have gained increasing
attention in cancer treatment, emphasizing the need for predictive biomarkers. Circular
RNAs (circRNAs) have emerged as critical regulators of tumor immunity, particularly in the
PD-1/PD-L1 pathway, and have shown potential in predicting immunotherapy efficacy. Yet,
the detailed roles of circRNAs in cancer immunotherapy are not fully understood. While
existing databases focus on either circRNA profiles or immunotherapy cohorts, there is
currently no platform that enables the exploration of the intricate interplay between
circRNAs and anti-tumor immunotherapy. A comprehensive resource combining circRNA
profiles, immunotherapy responses, and clinical outcomes is essential to advance our
understanding of circRNA-mediated tumor-immune interactions and to develop effective
biomarkers.
Methods To address these gaps, we constructed the Cancer CircRNA Immunome Atlas
(TCCIA), the first database that combines circRNA profiles, immunotherapy response data,
and clinical outcomes across multi-cancer types. The construction of TCCIA involved
applying standardized preprocessing to the raw sequencing FASTQ files, characterizing
circRNA profiles using an ensemble approach based on four established circRNA detection
tools, analyzing tumor immunophenotypes, and compiling immunotherapy response data
from diverse cohorts treated with immune-checkpoint blockades (ICBs).
Results TCCIA encompasses over 4,000 clinical samples obtained from 25 cohorts treated
with ICBs along with other treatment modalities. The database provides researchers and
clinicians with a cloud-based platform that enables interactive exploration of circRNA data

in the context of ICB. The platform offers a range of analytical tools, including browse of identified circRNAs, visualization of circRNA abundance and correlation, association analysis between circRNAs and clinical variables, assessment of the tumor immune microenvironment, exploration of tumor molecular signatures, evaluation of treatment response or prognosis, and identification of altered circRNAs in immunotherapy-sensitive and resistant tumors. To illustrate the utility of TCCIA, we showcase two examples, including circTMTC3 and circMGA, by employing analysis of large-scale melanoma and bladder cancer cohorts, which unveil distinct impacts and clinical implications of different circRNA expression in cancer immunotherapy.

Conclusions TCCIA represents a significant advancement over existing resources, providing a comprehensive platform to investigate the role of circRNAs in immuno-oncology.

What is already known on this topic

Prior knowledge indicated that circRNAs are involved in tumor immunity and have potential as predictive biomarkers for immunotherapy efficacy. However, there lacked a comprehensive database that integrated circRNA profiles and immunotherapy response data, necessitating this study.

What this study adds

This study introduces TCCIA, a database that combines circRNA profiles, immunotherapy response data, and clinical outcomes. It provides a diverse collection of clinical samples and an interactive platform, enabling in-depth exploration of circRNAs in the context of checkpoint-blockade immunotherapy.

How this study might affect research, practice or policy

The findings of this study offer valuable insights into the roles of circRNAs in tumor-immune interactions and provide a resource for researchers and clinicians in the field of immune-oncology. TCCIA has the potential to guide personalized immunotherapeutic strategies and contribute to future research, clinical practice, and policy decisions in checkpoint-blockade immunotherapy and biomarker development.

Introduction

Immunotherapy has revolutionized the treatment of cancer over the past decade, emerging as a groundbreaking approach that harnesses the patient's own immune system to fight cancer. Therapies like immune checkpoint blockades (ICBs), chimeric antigen receptor (CAR) T-cell therapy, and therapeutic vaccines aim to reinvigorate anti-tumor immunity against malignant cells [1–3]. ICBs targeting programmed cell death protein-1 (PD-1), PD-1 ligand (PD-L1), and cytotoxic T-lymphocyte-associated protein 4 (CTLA-4) in particular have demonstrated remarkable clinical efficacy across diverse cancer types, highlighting immunotherapy's potential for a durable and curative response [4]. Adoptive cell transfer using engineered T cells expressing CARs has also shown great promise for blood cancers [5]. However, significant challenges remain in extending immunotherapies to larger patient populations and solid tumors. Heterogeneity in response is a major limitation – while some patients achieve long-term remission, others exhibit intrinsic resistance or relapse after an initial response [6]. This variable efficacy likely stems from immunosuppressive mechanisms within the tumor microenvironment (TME) that enable cancer cells to evade immune attack [7]. Elucidating the complex cellular and molecular interactions underlying immunotherapy resistance will be critical to unlock the full potential of immune-based cancer treatments. Reliable predictive biomarkers are also imperative to guide patient selection and combination immunotherapies tailored to each patient's TME [8].

Circular RNAs (circRNAs) have recently emerged as fascinating non-coding RNA regulators with unique covalently closed loop structures. Initially considered splicing byproducts, circRNAs are now recognized as important gene expression modulators with diverse functions [9,10]. In cancer, circRNAs have been implicated in proliferation, metastasis, and malignancy hallmarks [11]. Moreover, circRNAs are now recognized as critical regulators and potential biomarker of tumor immunity and immunotherapy response [7,12,13]. Accumulating evidence indicates circRNAs modulate TME and immunotherapy outcomes through various mechanisms in cancers like lung cancer, melanoma, colorectal cancer, and pancreatic cancer [14,15]. For example, circRNAs such as circFGFR1, circ-CPA4, and circ_0000284 facilitate immune evasion by modulating PD-L1 via sponging tumor-suppressive microRNAs [16–18]. Additionally, circRNAs including hsa_circ_0000190 [19], circ_0020710 [20], CDR1-AS [16], and circ-UBAP2 [21] upregulate immune checkpoint proteins like PD-L1, CTLA-4 and PD-1, hampering T cell function and promoting immune evasion. Furthermore, cancer cell-derived circRNAs can reprogram intratumoral immune

cells via exosomal transfer or cytokine signaling, thereby impacting facets like angiogenesis that affect immunotherapy efficacy [22–24]. CircRNAs influence various aspects of the TME, including vascularization [25], metabolism [26], hypoxia [27], macrophage polarization [28], natural killer cell cytotoxicity [17], and T cell exhaustion/apoptosis [29]. These factors can impede the efficacy of immunotherapy [15,30]. Dysregulation of circRNAs promotes immune destruction evasion and reduced immunotherapy efficacy. CircRNAs employ diverse regulatory mechanisms—from sponging miRNAs and proteins to scaffolding proteins and translating peptides [31]. While many circRNAs originate in tumors, others come from stromal and immune cells, underscoring complex multicellular regulation [32]. Exploring circRNA networks will be critical to unraveling this intricate cancer-immunity interplay. With emerging roles in tumor immunity, prognostic potential, and biomarker utility, circRNAs represent a promising new frontier in cancer immunotherapy.

Despite growing interest in circRNAs and their potential relevance in cancer immunotherapy, a comprehensive understanding of their precise functions and clinical implications remains incomplete. Existing databases have limitations in either profiling circRNAs, such as riboCIRC [33], CSCD [34] and CircNet [35] offering circRNA profiles across tissues or cancers, or curating immunotherapy cohorts, like ICBatlas [36] and TCIA [37] compiling immune infiltration and immunotherapy data across tumor types. Crucially, no resources systematically integrate comprehensive circRNA expression with multi-omics datasets including immune cell fractions, ICB types, and clinical outcomes for systematic exploration of the circRNA-immunotherapy interplay.

To address this unmet need, we developed the first-of-its-kind database, The Cancer CircRNA Immunome Atlas (TCCIA), a comprehensive database that integrates circRNA profiles, immunotherapy response data, and clinical outcomes for multiple cancer types, with the objective of providing a valuable resource for systematic exploration of the circRNA-immune axis, advancing our understanding of their functions and to facilitates discovery of potential biomarkers, therapeutic targets and clinical implications in cancer immunotherapy.

Results

Integrating circular RNAs in cancer immunotherapy

The development of the TCCIA database encompassed a comprehensive process involving data collection, preprocessing, and integration (Figure 1). In terms of data collection, we carefully curated research articles detailing cohorts treated with ICBs, utilizing the PubMed

database (https://www.ncbi.nlm.nih.gov/pubmed/) for selection. We acquired raw RNA-seq datasets from several genome sequence archive repositories, including dbGaP, EGA, EMBL-EBI and GSA. We applied standardized preprocessing techniques to identify circRNAs, quantify TME/gene signatures, and incorporate clinical annotations and outcomes. This approach was aimed at improving the consistency and comparability of data across different datasets. Notably, TCCIA addresses a crucial gap by providing a unified platform with multiple tools that facilitates the exploration of circRNAs' impact on immunotherapy outcomes (Figure 1). This sets it apart from many existing databases [33–47] that predominantly focus on circRNA profiles, circRNA annotations, or immunotherapy cohorts (Table 1).

Figure 1. Overview of TCCIA.

Table 1. Comparison between TCCIA and other similar database resources [33–47].

Type	Name	Publication (Year)	Last update	Support species	Sample types (N)	Number of samples	Data sources	Characteristics
Mixed	TCCIA	Submitted	2023	Homo sapiens	cancer (10)	> 4000 cancer samples	dbGaP, EMBL-EBI, EGA, GSA	Characterization of circRNA expression profiles within and across checkpoint bloackade immunotherapy clinical datasets; Comparison of circRNA between clinical phenotypes (including response vs. nonresponse in pre-treatment or on-treatment); circRNA based survival analyses; the association of circRNA with 255 cancer signatures and 8 types of TME metrics; analyzing multiple circRNAs simultaneously
CircRNA related	CSCD2	NAR Database Issue (2022)	2022	Homo sapiens	cancer (23)	825 tissues + 288 cell lines	ENCODE, SRA	Prediction of potential miRNA–circRNA and RBP–circRNA interactions; prediction of potential circRNA full-length and open reading frame sequence
	circMine	NAR Database Issue (2022)	2022	Homo sapiens	disease (87)	1107 samples	GEO	Provide 13 online analytical functions to comprehensively investigate circRNAs, including search, browse, differential expresion analysis, co-expression analysis, circRNA-miRNA prediction, circRNA IRES prediction, ribo-circRNA location, etc.
	CircR2Disease2	Genomics, Proteomics & Bioinformatics (2021)	2022	5 species	disease (313)	2449 studies	PubMed	Investigation of dysregulated circRNAs and their posttranscriptional regulatory function
	CircNet2	NAR Database Issue (2021)	2021	Homo sapiens	cancer (37)	2732 cancer samples	TCGA, GEO, CircAtlas, MiOncoCirc	Characterization of tissue-specific circRNA expression profiles and circRNA-miRNA-gene regulatory network
	riboCIRC	Genome Biology (2021)	2021	21 species	tissue/cell line (314)	1970 samples	GEO	Exploration of computationally predicted ribosome-associated circRNAs, experimentally verified translated circRNAs, and cross-species conserved ribo-cirRNAs
	CircAtlas2	Genome Biology (2020)	2020	7 species	tissue (20)	1070 samples	SRA, NGDC, GeneBank	Comprehensive profiling of circRNAs, including their searching, orthologous, converter, expression, sequences, translations, and functions
	CircInteractome	RNA Biology (2016)	2020	Homo sapiens	tissue/cell line (34)	34 samples	circBase	Exploration of the interactions between miRNAs and circRNAs with 109 RBPs
	TransCirc	NAR Database Issue (2020)	2020	Homo sapiens	tissues (17)	17 tissues	CircAtlas	Search and browse six types of evidence supporting the translation potential of circRNAs
	circRic	Genome Medicine (2019)	2019	Homo sapiens	cancer (22)	935 cancer celllines	CCLE	Characterization of circRNA expression profiles; analysis of the circRNA biogenesis regulators, the effect of circRNAs on drug response, the association of circRNAs with mRNAs, proteins, mutations, and RBPs and miRNA binding sites.
	CIRCpedia2	Genomics, Proteomics & Bioinformatics (2018)	2018	6 species	tissue (13)	185 samples	GEO, ENCODE, EMBL-EBI	Search, browse and download CIRCexplorer2 identified circRNAs; simple visualization of circRNA expression
	circBase	RNA (2014)	2014	6 species	tissues/cell line (77)	77 samples	GEO, ENCODE, EMBL-EBI	Public (evidence supporting) circRNA search; annotation preview and export
Immunotherapy related	TISMO	NAR Database Issue (2022)	2022	Mouse	cancer (19)	1518 mouse samples	GEO and In-house data	Summary data browser; Boxplots comparing expression of a gene, pathway, or immune cell type either in vivo or in vitro samples across treatment and response conditions
	ICBatlas	Cancer Immunol Res (2022)	2022	Homo sapiens	cancer (9)	1515 cancer samples	GEO,ArrayExpress, TCGA, dbGaP	Gene-, pathway-, and immune cell-level search, summary, and visualization; Comparison between response vs. nonresponse, pre-treatment vs. on-treatment of checkpoint bloackade immunotherapy; Provide a response score system
	ImmuCellAI	Advanced Science (2020)	2020	Homo sapiens	cancer (37)	NA	GEO, TCGA, dbGaP	Estimation of the abundance of 24 immune cells; survival analysis
	TCIA	Cell Reports (2017)	2017	Homo sapiens	cancer (20)	9562 cancer samples	TCGA and two immunotherapy studies	Advanced data selection; gene expression analysis; cell type abundance analysis; GSEA; survival analysis; tumor heterogeneity summary; neoantigen summary and analysis

186
Figure 2. Content of TCCIA. (**A**) Inclusion of cancer types, cohort, and publication years in this study.
Abbr. BLCA, Bladder urothelial carcinoma; HNSC, Head and neck squamous cell carcinoma; KIRC,
Kidney renal clear cell carcinoma; NSCLC, Non-small lung cancer; SCLC, Small lung cancer; SKCM,
Skin cutaneous melanoma; SGC, Salivary gland carcinoma; BRCA, Breast cancer; SARC, Sarcoma;
STAD, Stomach Adenocarcinoma. (**B-C**) The number of detected circRNAs, length of circRNAs, mean
(back-splicing junction count (BSJ) and mean counts per million (CPM) in different cancer types (**B**) or
cohort (**C**). (**D**) UMAP plot of all samples, colored by cancer types.
194

Data summary of TCCIA

In this study, a comprehensive compilation was made, involving approximately 4000 clinical

patient samples from 25 immune-checkpoint blockade (ICB) cohorts [48–70] with raw RNA-

seq datasets published between 2012 and 2023, encompassing 10 distinct cancer types

(Figure 2A, B and Supplementary Table 1). We employed an ensemble strategy that

combined the outcomes of four different circRNA detection tools: CIRCexplorer2 [71],

CIRIquant [72], find_circ [73], and circRNA_finder [74]. This approach is akin to the one

utilized in the study conducted by Dong et al. [13] (refer to the Methods section for more

details). In total, we detected 793,316 circRNAs using at least two circRNA detection

methods. By applying the ensemble strategy, we successfully identified and included a total

of 555,098 unique circRNAs in the TCCIA database (Figure S1). A significant proportion of these circRNAs (23.2% to CIRCpedia [44] and 40.3% to circAltas [40]) could be confidently mapped to well-known circRNA databases. Among included cancer types, Bladder urothelial carcinoma (BLCA) exhibited the highest level of circRNA abundance, encompassing 141,085 circRNAs (representing 25.4% of the total). Conversely, Sarcoma (SARC) demonstrated the lowest level of circRNA abundance, with only 4109 circRNAs, constituting a mere 0.7% of the total (Figure 2B). The average lengths of these circRNAs demonstrated notable consistency across various cancer types, ranging from 11,176.81 to 19,970.70, as well as within individual cohorts, ranging from 8,899.24 to 21,357.33 (Figure 2C). Intriguingly, Head and neck squamous cell carcinoma (HNSC) exhibited the highest mean back-splice junction (BSJ) reads at 5.10, whereas BLCA had the lowest mean at 0.07 (Figure 2C). A comprehensive analysis encompassing all the circRNAs from the sampled datasets was visualized using t-SNE (Figure 2D). This visualization revealed discernible circRNA clustering patterns specific to various cancer types, highlighting the nuanced circRNA heterogeneity within human cancers and emphasizing the need for independent circRNA analysis considerations.

Web functionality of TCCIA

TCCIA offers a diverse array of exploratory and advanced analytical tools, providing a comprehensive platform for researchers to uncover intricate connections and gain valuable insights (Figure 1). The modules are thoughtfully introduced, organized, and accessible across nine web pages: "Home", "Browse", "Repository", "Cohort-centred Analysis", "Molecule-centred Analysis", "Signature", "DE circRNAs", "Setting", and "Help" These functionalities enable users to explore circRNA information, abundance, correlations with clinical variables, the tumor immune microenvironment, molecular signatures, treatment responses, and prognosis predictions. Researchers can also identify circRNAs associated with both immunotherapy-sensitive and resistant tumor scenarios. The well-established exploration and analysis workflow within the TCCIA framework is illustrated in Figure 3, outlining the typical path that researchers follow when engaging with the platform. A more comprehensive explanation of each fundamental module is provided below.

Figure 3. A standard exploration and analysis pipeline within TCCIA.
(A) Users need to navigate to the Repository page and select the clinical cancer dataset(s) of their interest. On the leftmost panel, users can filter datasets based on various parameters. **(B)** Users can choose an appropriate analysis strategy according to their needs (core analysis steps are listed on the rightmost panel). The analysis typically consists of four steps: I. Select the dataset(s). II. Choose the CircRNA ID or Host Gene of interest. III. Adjust various parameters such as Test Type, Color Selection, etc. IV. Click the Submit button to obtain the analysis and visualization results. **(C)** Users can browse various information about all CircRNAs included in TCCIA, and at the same time, they can use the ID Converter to transform their interested CircRNAs into other CircRNA Database IDs. **(D)** Users can use the provided options to save result images in PDF or PNG format, with the desired dimensions.

CircRNA Browser. TCCIA's circRNA browser offers a distinctive and valuable resource for the exploration and retrieval of identified circRNAs within cancer cohorts. The browser provides essential information such as gene symbols, genomic locations (based on the hg38 genome build), cohort origins, statistical metrics reflecting circRNA expression distribution (including mean, median, maximum, standard deviation, and sample size - N), and cross-references to identifiers from CIRCpedia and circAtlas. Users can conveniently pinpoint circRNAs of interest by applying filters based on gene symbols, CIRCpedia identifiers, and/or circAtlas identifiers.

Cohort Selection and Data Access. The TCCIA interface offers an intuitive approach for cohort selection and data access. The Repository Page serves as a gateway, enabling users to filter datasets based on crucial parameters such as cancer type, treatment modalities, drugs administered, and cohort sizes. Essential details pertaining to each

dataset are presented in a comprehensive cohort table, facilitating informed decision-
making regarding cohort selection.

Cohort/Molecule-Centered Analysis Modules. At the heart of TCCIA's capabilities lie the
cohort-centered analysis modules and molecule-centered analysis modules (for analyzing
circRNAs across multiple cohorts), providing a profound lens into circRNA dynamics within
specific immunotherapy cohorts. These modules encompass:
(1) *Scatter-Correlation and Heatmap-Correlation*: Researchers gain insights into circRNA
correlations through scatter plots and heatmaps. These visualizations are pivotal in
elucidating potential connections between circRNAs and other variables within the
chosen cohort.
(2) *Group-Comparison (including simplified and comprehensive versions)*: TCCIA facilitates
nuanced analysis of numeric differences in circRNA expression across multiple groups
within a cohort. The dual modes of simplified and comprehensive group comparison
empower researchers to unravel intricate circRNA expression patterns.
(3) *Kaplan-Meier(KM)-Analysis and Cox-Analysis*: Survival analysis is made accessible
through the KM-Analysis module, which generates Kaplan Meier survival curves among
distinct variable groups. Additionally, the Cox-Analysis module allows for an in-depth
examination of survival outcomes of any circRNA expression, opening avenues to
prognostic evaluations.

Signature and DEG Analysis. TCCIA introduces dedicated modules for signature analysis
and differential expression circRNAs (DEG) assessment. The Signature Page facilitates the
investigation of associations between circRNAs and tumor microenvironment metrics using
eight prominent deconvolution methods. It also allows for the examination of connections
between circRNAs and 255 cancer signatures categorized into three distinct groups: TME-
associated, tumor-metabolism, and tumor-intrinsic signatures. These analyses encompass
a wide range of cohorts, ensuring comprehensive exploration of these relationships. The
DEG Page empowers researchers to pinpoint differentially expressed circRNAs between
patients who respond and those who do not respond to immunotherapy, thus unraveling the
intricate web of circRNA involvement in treatment outcomes.

User Customized Configurations. Global settings within TCCIA add a layer of refinement
to the user experience, allowing for customized exploration. These settings grant users'

control over data access and enable tailoring analyses to align with their specific research objectives. For example, by default, the platform prioritizes immunotherapy-related sample data by filtering out samples without checkpoint immunotherapy treatment, streamlining analyses for coherent research goals. Besides, TCCIA supports gene signature in a nutshell, i.e., the addition, subtraction, multiplication, and division are support for aggregating circRNAs of interest as a whole. As users become acclimated to the platform, customization options foster enhanced flexibility, enabling researchers to uncover novel insights.

In essence, the web functionality of TCCIA embodies an advanced and user-centric avenue for investigating the complex roles of circRNAs in cancer immunotherapy. The integration of diverse analysis modules, coupled with a cohort-centered approach and adaptable settings, positions TCCIA as an indispensable tool for advancing our comprehension of circRNA-mediated immune responses and guiding the formulation of personalized immunotherapeutic strategies. This interactive platform stands poised to reshape the landscape of circRNA-immunotherapy research.

Case studies validating and extending data of two reported CircRNAs

ICB therapies targeting PD-1 and CTLA-4 have significantly transformed the field of oncology, particularly in the treatment of metastatic melanoma. However, it is important to note that only a limited number of melanoma patients experience positive outcomes from these immunotherapies. Consequently, there is a pressing need to identify predictive biomarkers that can guide precision oncology.

In the study conducted by Dong et al. [13], it was observed in metastatic melanoma patients from the Gide et al. cohort [54] and independently validated in metastatic melanoma patients from the Berger et al. cohort [63], who exhibited high expression of circTMTC3, experienced poorer survival outcomes and demonstrated reduced treatment responsiveness compared to those with low circTMTC3 expression. To illustrate the functionality of the TCCIA, here, we performed a reproducible analysis on the same cohorts (Figure 4).

325
Figure 4. circTMTC3 predicts poor survival and non-response to checkpoint immunotherapy in
the melanoma cohorts of Gide et al. and Berger et al.
(**A**) Analysis panel for analyzing and visualizing associations between circRNAs and patient survival.
Here, we present the data and analysis settings used to generate the plot shown in (**D**). (**B-C**)
CircTMTC3 predicts poor survival in (**B**) Gide et al. cohort and (**C**) Berger et al. cohort. (**D-E**) A circRNA
isoform of circTMTC3, TMTC3:+:chr12:88148287:88176319, predicts poor survival in (**D**) Gide et al.
cohort and (**E**) Berger et al. cohort. Red, group of high circRNA expression. Blue, group of low circRNA
expression. (**F-G**) Volcano plots showing differential expressed circRNAs between checkpoint
immunotherapy response and nonresponse patients using the approaches Limma Voom, DESeq2, and
edgeR in (**F**) Gide et al. cohort and (**G**) Berger et al. cohort.

TCCIA provides an overview of the analysis panel used to assess the association between a circRNA and patient survival (Figure 4). Here, using TCCIA clearly demonstrate that high levels of circTMTC3 are predictive of poor overall survival in the Gide et al. cohort ($HR_{low\ vs\ high}$=0.48, 95% CI: 0.24-0.93, P=0.023) and the Berger et al. cohort ($HR_{low\ vs\ high}$=0.5, 95% CI: 0.28-0.89, P=0.006) (Figure 4B-C). Upon closer examination of the different isoforms of circRNAs derived from TMTC3, it was found that only TMTC3:+:chr12:88148287:88176319 was abundant and played a significant role in predicting unfavorable survival outcomes in the Gide et al. cohort (Figure 4D, $HR_{low\ vs\ high}$=0.42, 95% CI: 0.21-0.85, P=0.007) and the Berger et al. cohort (Figure 4E, $HR_{low\ vs\ high}$=0.5, 95% CI: 0.26-0.95, P=0.01). Furthermore, we utilized three commonly employed methods to evaluate the differential expression of circRNAs between checkpoint immunotherapy responders and non-responders. Consistent with the survival outcome, we observed that TMTC3:+:chr12:88148287:88176319 was the only isoform of circTMTC3 to exhibit significant upregulation in non-responding patients from both metastatic melanoma cohorts (Figure 4F-G). Mechanistically, Dong et al. [13] demonstrated that the overexpression of circTMTC3 could elevate PD-L1 expression through the miR-142-5p/PD-L1 axis, thereby reducing T cell activity and promoting immune evasion. Our identification of the specific functional isoform of circTMTC3 may stimulate further investigation into the dynamics and role of circRNA isoforms in immuno-oncology.

In the recent study by Sun et al. [75], it was reported that circMGA, a circRNA with tumor-suppressing properties, functions as a chemoattractant for CD8+ T cells. The study indicated a positive correlation between elevated levels of circMGA and improved response to immunotherapy in bladder cancer. However, the analysis conducted by Sun et al. solely encompassed a small cohort of 73 patients, without considering the effects of checkpoint blockade immunotherapy. To validate their findings and assess the clinical relevance of circMGA, we performed an expanded analysis employing the Bellmunt et al. BLCA cohort [49] (IMvigor010, NCT02450331), which comprised 728 clinical patients encompassing two treatment modalities (374 patients received adjuvant immunotherapy, while 354 were assigned to observation arm). Our analysis demonstrated that heightened levels of circMGA were associated with a favorable prognosis in patients, regardless of their receipt of immunotherapy (Supplementary Figure 2A-B). Notably, we observed that increased circMGA expression presented as a particularly strong predictor of improved overall survival (OS) in early-stage tumors (≤ pT2) received immunotherapy ($HR_{low\ vs\ high}$ = 3.77, 95% CI:

1.43-9.94, *P* = 0.0042) (Supplementary Figure 2A). Furthermore, patients with elevated circMGA expression in the immunotherapy group exhibited a higher CD8 T cell signature (Supplementary Figure 2C), potentially clarifying the augmented effectiveness of immunotherapy in these individuals. These findings provide additional substantiation for the conclusions drawn by Sun et al.

Together, the two aforementioned case studies, employing analysis of large-scale clinical cancer cohorts, have unveiled distinct impacts and clinical implications of different circRNA expression in cancer immunotherapy across cancer types. These studies showcase the exploratory and validating value of TCCIA in investigating the role of circRNA in tumor immunity.

Discussion

Circular RNAs, emerging as pivotal gene expression regulators, exert diverse functions across biological processes, with substantial clinical research potential [76]. However, the existing landscape of circRNA-related tumor-immune checkpoint research falls short of satisfying the burgeoning need for insights. This limitation is compounded by constraints stemming from sample quantity and diversity, which subsequently curtail the generalizability of research findings due to geographical, racial, and tumor-specific factors. Consequently, given this evolving landscape, the urgency and significance of developing a CircRNA tool for preliminary data mining have become increasingly pronounced. TCCIA (the Cancer CircRNA Immunome Atlas) an integrated online platform, building upon datasets from four genome sequence archive repositories, encompassing around 4,000 cancer samples, spanning five cancer types and 25 checkpoint-blockade immunotherapy cohorts, incorporating over 0.55 million circRNA expression profiles, 255 established cancer signatures, and TME decomposition results from eight immune infiltration algorithms. TCCIA emphasizes user-friendly visual presentations, eliminating the need for intricate programming skills. Moreover, the platform offers a range of customizable visualization options, ensuring adaptability to user needs. Notably, TCCIA is readily accessible without mandatory registration or login, potentially rendering it an economical and efficient solution for both researchers and clinical practitioners.

In this manuscript, we delineate the data sources, collection and standardization processes, TCCIA's functionalities, website analysis modules, and provide a step-by-step guide to its operation. To further illustrate TCCIA, we present two concrete examples of the circTMTC3 and circMGA as molecular markers in melanoma and bladder cancer,

respectively.

While TCCIA has various advantages and uniqueness, there are still some
limitations. In terms of data, although TCCIA includes data from 10 types of cancer, the
sample size for some cancer types (e.g., head and neck cancer) remains sparse. Recently,
the application of immunotherapy has been expanding to more cancer types, such as
colorectal cancer. The main reason for this is that current clinical genomics research of
such cancer types primarily focuses on the DNA level, specifically on whole exome
sequencing and targeted sequencing [77]. As a result, there are few RNA-seq datasets to
infer the presence of circRNAs. Another reason is that some raw RNA-seq datasets are
difficult to access due to restricted availability, e.g., Carroll et al. study [78]. In terms of data
integration, we acknowledge that while TCCIA supports the integration and display of
multiple cohorts, there are inherent challenges when it comes to providing uniform
functionality for multi-cohort analyses. These challenges arise from the heterogeneity in
circRNA profiles and phenotypic data across different cohorts. For users with specific,
detailed analysis needs that span multiple cohorts, we recommend leveraging the flexibility
of the platform to perform combined analyses based on the results obtained from single-
cohort analyses. This approach allows for a more tailored and comprehensive exploration
of individual cohort data while accommodating the inherent variations in circRNA profiles
and phenotypes across different cohorts. In terms of functionality, some databases provide
the characteristics of studying circRNA-miRNA-gene regulatory networks (such as CSCD2
[34], circMine [38], and CircNet2 [35]), which have not been considered in TCCIA with two
reasons: First, further experimental validation is generally required to confirm the
authenticity of detected circRNAs. In the recent large-scale circRNA detection benchmark
study [79], Vromman et al. recommended using qPCR+ Ribonuclease R or qPCR+amplicon
sequencing for circRNA validation. Therefore, it is recommended to incorporate these
experimental validation methods to ensure the accuracy of circRNA detection. Second, the
focus of our study is on integrating circRNA profiles and clinical outcomes in cancer
patients treated with immunotherapy. However, instead of duplicating efforts, users can
leverage already well-established and high-quality circRNA databases to complete other
circRNA annotation and analysis explorations. By linking to these databases (a widget is
provided in the footer of the TCCIA website), users can access comprehensive circRNA
information and utilize existing tools for further analysis, optimizing the accuracy and
efficiency of circRNA annotation and deep investigation.

Looking ahead, we aim to continually update TCCIA by incorporating more circRNA

data from diverse and updating cancer immunotherapy cohorts and introducing new
functionalities based on user feedback. In summary, the distinctive features, analytical
capabilities, and potential for growth position it as a pivotal tool in advancing our
understanding of circRNAs in tumor immunity and in shaping development of personalized
immunotherapy strategies guided by circRNA.

Methods

Data collection

To conduct a systematic search, we utilized PubMed
(https://www.ncbi.nlm.nih.gov/pubmed/) to search for articles related to Bulk RNA-seq data
from solid cancer patients treated with immune checkpoint blockers (ICB). The search
expression used was "(ICB [Title/Abstract] OR PD-1[Title/Abstract] OR PD-L1[Title/Abstract]
OR CTLA-4[Title/Abstract]) AND (rnaseq[Title/Abstract] OR rna-seq[Title/Abstract] OR rna-
sequencing[Title/Abstract] OR rna seq[Title/Abstract] OR rna sequencing[Title/Abstract])".
No filters were applied, and there were no restrictions on language or geographic region.
Peer-reviewed publications, preprints, and press releases were considered for inclusion. To
obtain raw RNA sequencing data, we submitted requests to the Database of Genotypes
and Phenotypes (dbGaP) (https://dbgap.ncbi.nlm.nih.gov/), the European Genome-
phenome Archive (EGA) (https://ega-archive.org/), and the Genome Sequence Archive
(GSA) (https://ngdc.cncb.ac.cn/gsa/) of the National Genomics Data Center (NGDC) after
receiving approval from the Data Access Committee (DAC). However, it is important to note
that not all raw RNA-seq datasets were accessible and available for use. In total, we
collected 23 studies [48–70] related to checkpoint immunotherapy that provided raw RNA-
seq datasets (Supplementary Table 1). We gathered relevant clinical data from publications
and clinical meta documents associated with these RNA-seq datasets. Additionally, we
extracted information on the cohorts' fundamental characteristics, such as sample size,
treatment methods, and specific drugs used, based on the abstracts. It should be noted that,
for Patil et al. study, two clinical cohorts were included; and for the study identified as
PHS000452, the two patient subgroups had distinct drug treatments and clinical
annotations. Hence, we treated them as two separate cohorts during the analysis. For the
TCCIA project, CircRNA profiles, immunotherapy response, and clinical benefits were
analyzed for 10 cancer types. This analysis included over 4,000 clinical samples from 25
cohorts treated with immune-checkpoint blockades (ICBs) such as PD-1/PD-L1 and CTLA-
4 inhibitors alone or in combination with conventional therapeutics. The analysis considered

both pre-treatment and on-treatment responses.

CircRNA identification and differential expression analysis

We employed multiple circRNA detection tools, including CIRCexplorer2 [71], CIRIquant

[72], find_circ [73], and circRNA_finder [74], to identify, parse, and annotate circRNA

junctions within each sample, using the human genome version hg38 as the reference. In

our pursuit of biologically robust circRNA identification within each cohort, we undertook an

approach that amalgamated results from various circRNA detection tools and implemented

a rigorous ensemble strategy characterized by the following stringent criteria. Firstly, we

exclusively retained circRNAs located on chromosomes 1 to 22, as well as the X and Y

chromosomes. Secondly, circRNAs were mandated to exhibit genomic overlap with gene

regions   as   defined   in   the   reference   genome   annotation   file,   specifically

"gencode.v34.annotation.gtf." Lastly, to ensure heightened robustness, circRNAs had to be

concurrently identified by a minimum of two out of the four employed detection tools, with at

least one of these tools detecting no fewer than two back-splicing junction counts. This

meticulous methodological approach was instrumental in ensuring the acquisition of

biologically sound and reliable circRNA datasets, thereby fortifying the integrity of our

analysis within each cohort. Subsequently, the back-splicing junction counts underwent

differential expression analysis using established approaches, namely Limma Voom [80],

edgeR [81], and DESeq2 [82]. This analysis facilitated comparisons between patient groups,

such as those who responded to checkpoint immunotherapy (responders) versus non-

responders, as well as between samples collected before and during ICB treatment.

TME decomposition and cancer gene signature estimation

We employed IOBR [83] for TME decomposition and the scoring of cancer gene signatures.

IOBR seamlessly integrates eight widely-used open-source deconvolution methods,

including CIBERSORT [84], ESTIMATE [85], quanTIseq [86], TIMER [87], IPS [37],

MCPCounter [88], xCell [89], and EPIC [90]. Furthermore, IOBR incorporates a

comprehensive compilation of 255 established cancer signatures. These diverse signatures

are organized into three distinct categories: TME-associated, tumor-metabolism, and tumor-

intrinsic signatures.

TCCIA implementation

The TCCIA database is developed as a Web application leveraging R Shiny

(https://shiny.posit.co/) and built using the golem framework (https://github.com/ThinkR-open/golem) to achieve optimization. TCCIA, is developed solely for research purposes and does not utilize any cookies or collect any personal identifiable information. We independently deployed TCCIA on both the reliable hiplot cloud platform (https://shiny.hiplot.cn/TCCIA/) and our team's servers (http://tccia.zmu-zhoulab.com/ or http://biotrainee.vip:18888/TCCIA/) for free access. We're dedicated to promptly addressing and resolving any issues impacting user experience. We will continually expand the TCCIA database, incorporate additional cancer types, and strive to gain access to more data resources every six months.

Statistical analysis

We performed Kaplan-Meier survival analysis to generate and compare survival curves. The log-rank test was used for comparison. We also conducted multivariate survival analysis using the Cox regression model. All reported *P*-values are two-tailed, and a significance level of p≤0.05 was used unless otherwise specified. All statistical analyses and visualization were conducted using R v4.2.0.

Patient consent for publication

Not applicable.

Ethics

In this study, we conducted a secondary analysis of trial datasets, which was determined to carry minimal risk. The Institutional Review Board of the Second Affiliated Hospital, Zunyi Medical University (No. YXLL(KY-R)-2021-010) approved the study protocol. As per national legislation and institutional guidelines, written informed consent for participation was not deemed necessary for this particular study.

Data availability

All the processed data can be accessed on the TCCIA website or in the article. We collected ICB-related cancer data sets with tumor gene expression profiles from the dbGaP, EGA, EMBL-EBI and GSA databases. Following the accession instruction described in published ICB studies (Table S1), we downloaded ICB patients' RNA-Seq raw sequencing data, clinical information, and response outcome information from ICB studies (if available).

Curated circRNA identification results from 4 detection methods and our ensemble approach are available at Zenodo (https://zenodo.org/doi/10.5281/zenodo.7969298). For requesting processed circRNA expression data from the TCCIA, please contact the project leader, Jian-Guo Zhou.

Code availability

The cancer circRNA identification pipeline with an ensemble approach and corresponding scripts and logs for handling TCCIA cohorts are available at https://github.com/ShixiangWang/circrna-pipeline (commit id: 1115905).

Funding

This work was supported by the National Natural Science Foundation of China (Grant No. 81660512, 81472594, 81770781, 82270825), Chunhui program of the Chinese Ministry of Education (Grant No. HZKY20220231), the Natural Science Foundation of Guizhou Province (Grant No. ZK2021-YB435), Guangdong Basic and Applied Basic Research Foundation (Grant No. 2021A1515011743), Youth Talent Project of Guizhou Provincial Department of Education (Grant No. QJJ2022-224), China Postdoctoral Science Foundation (Grant No. 2021M703733), China Lung Cancer Immunotherapy Research Project, Special funds for innovation in Hunan Province (Grant No. 2020SK2062), the Science and Technology Programs of Zunyi City (Grant No. ZSKH-HZ2023-142), and Collaborative Innovation Center of Chinese Ministry of Education (Grant No. 2020-39).

Contributions

SW: Conceptualization, software, methodology, formal analysis, writing original draft, review and editing. YX: Software, methodology, formal analysis, visualization, writing original draft. YZ: Software, formal analysis, visualization, writing original draft. HW: Conceptualization, writing original draft, review and editing. JL and USG: Resources, review and editing. MC: Visualization, review and editing. PL, YHL, MH and BF: Review and editing. XL, QZ and HM: Supervision, resources, funding acquisition. JGZ: Conceptualization, methodology, resources, supervision, funding acquisition, project administration, writing original draft, writing–review and editing.

Conflict of interest

None were declared.

Acknowledgments

We thank Dr. Jianming Zeng (University of Macau), and all the members of his
bioinformatics team, biotrainee, for generously sharing their experience and codes. We
thank Juan Zhang (Echo Biotech Co., Ltd., Beijing, China) for her help in the data pre-
processing. We thank Jieyin Tan (University of Sydney) for test TCCIA and report bugs.
The present work was performed in (partial) fulfillment of the requirements for obtaining the
degree "Dr. rer. biol. hum."at the Friedrich-Alexander-Universität Erlangen-Nürnberg (FAU).

References

Sharma P, Allison JP. Immune Checkpoint Targeting in Cancer Therapy: Toward Combination Strategies with Curative Potential. *Cell*. 2015;161:205–14.

Maude SL, Laetsch TW, Buechner J, *et al.* Tisagenlecleucel in Children and Young Adults with B-Cell Lymphoblastic Leukemia. *N Engl J Med*. 2018;378:439–48.

Guo C, Manjili MH, Subjeck JR, *et al.* Chapter Seven - Therapeutic Cancer Vaccines: Past, Present, and Future. In: Tew KD, Fisher PB, eds. *Advances in Cancer Research*. Academic Press 2013:421–75. https://doi.org/10.1016/B978-0-12-407190-2.00007-1

Topalian SL, Hodi FS, Brahmer JR, *et al.* Safety, Activity, and Immune Correlates of Anti–PD-1 Antibody in Cancer. *N Engl J Med*. 2012;366:2443–54.

Neelapu SS, Locke FL, Bartlett NL, *et al.* Axicabtagene Ciloleucel CAR T-Cell Therapy in Refractory Large B-Cell Lymphoma. *N Engl J Med*. 2017;377:2531–44.

Sharma P, Hu-Lieskovan S, Wargo JA, *et al.* Primary, Adaptive, and Acquired Resistance to Cancer Immunotherapy. *Cell*. 2017;168:707–23.

Vinay DS, Ryan EP, Pawelec G, *et al.* Immune evasion in cancer: Mechanistic basis and therapeutic strategies. *Semin Cancer Biol*. 2015;35:S185–98.

Gibney GT, Weiner LM, Atkins MB. Predictive biomarkers for checkpoint inhibitor-based immunotherapy. *Lancet Oncol*. 2016;17:e542–51.

Salzman J. Circular RNA Expression: Its Potential Regulation and Function. *Trends Genet*. 2016;32:309–16.

Liu C-X, Chen L-L. Circular RNAs: Characterization, cellular roles, and applications. *Cell*. 2022;185:2016–34.

Kristensen LS, Hansen TB, Venø MT, *et al.* Circular RNAs in cancer: opportunities and challenges in the field. *Oncogene*. 2018;37:555–65. doi: 10.1038/onc.2017.361

Zhou J-G, Liang R, Wang H-T, *et al.* Identification and characterization of circular RNAs as novel putative biomarkers to predict anti-PD-1 monotherapy response in metastatic melanoma patients – Knowledge from two independent international studies. *Neoplasia*. 2023;37:100877.

Dong Y, Gao Q, Chen Y, *et al.* Identification of CircRNA signature associated with tumor immune infiltration to predict therapeutic efficacy of immunotherapy. *Nat Commun*. 2023;14:2540. doi: 10.1038/s41467-023-38232-y

Pandey PR, Young KH, Kumar D, *et al.* RNA-mediated immunotherapy regulating tumor immune microenvironment: next wave of cancer therapeutics. *Mol Cancer*. 2022;21:58.

Guan L, Hao Q, Shi F, *et al.* Regulation of the tumor immune microenvironment by cancer-derived circular RNAs. *Cell Death Dis*. 2023;14:1–12. doi: 10.1038/s41419-023-05647-w

Tanaka E, Miyakawa Y, Kishikawa T, *et al.* Expression of circular RNA CDR1⸺AS in colon cancer cells increases cell surface PD⸺L1 protein levels. *Oncol Rep*. 2019;42:1459–66.

Zhang P-F, Gao C, Huang X-Y, *et al.* Cancer cell-derived exosomal circUHRF1 induces natural killer cell exhaustion and may cause resistance to anti-PD1 therapy in hepatocellular carcinoma. *Mol Cancer*. 2020;19:110.

Hong W, Xue M, Jiang J, *et al.* Circular RNA circ-CPA4/ let-7 miRNA/PD-L1 axis regulates cell growth, stemness, drug resistance and immune evasion in non-small cell lung cancer (NSCLC). *J Exp Clin Cancer Res*. 2020;39:149.

Luo Y-H, Yang Y-P, Chien C-S, *et al.* Circular RNA hsa_circ_0000190 Facilitates the Tumorigenesis and Immune Evasion by Upregulating the Expression of Soluble PD-L1 in Non-Small-Cell Lung Cancer. *Int J Mol Sci*. 2021;23:64.

Wei C-Y, Zhu M-X, Lu N-H, *et al.* Circular RNA circ_0020710 drives tumor progression

631 and immune evasion by regulating the miR-370-3p/CXCL12 axis in melanoma. *Mol*
632 *Cancer*. 2020;19:84.

21 Zhao R, Ni J, Lu S, *et al.* CircUBAP2-mediated competing endogenous RNA network
634 modulates tumorigenesis in pancreatic adenocarcinoma. *Aging*. 2019;11:8484–501.

22 Ou Z-L, Luo Z, Wei W, *et al.* Hypoxia-induced shedding of MICA and HIF1A-mediated
636 immune escape of pancreatic cancer cells from NK cells: role of circ_0000977/miR-153
637 axis. *RNA Biol*. 2019;16:1592–603.

23 Wang X, Sheng W, Xu T, *et al.* CircRNA hsa_circ_0110102 inhibited macrophage
639 activation and hepatocellular carcinoma progression via miR-580-5p/PPARα/CCL2
640 pathway. *Aging*. 2021;13:11969–87.

24 Li Z, Huang C, Bao C, *et al.* Exon-intron circular RNAs regulate transcription in the
642 nucleus. *Nat Struct Mol Biol*. 2015;22:256–64. doi: 10.1038/nsmb.2959

25 Liu C, Yao M-D, Li C-P, *et al.* Silencing Of Circular RNA-ZNF609 Ameliorates Vascular
644 Endothelial Dysfunction. *Theranostics*. 2017;7:2863–77.

26 Di Timoteo G, Dattilo D, Centrón-Broco A, *et al.* Modulation of circRNA Metabolism by
646 m6A Modification. *Cell Rep*. 2020;31:107641.

27 Jiao B, Liu S, Zhao H, *et al.* Hypoxia-responsive circRNAs: A novel but important
648 participant in non-coding RNAs ushered toward tumor hypoxia. *Cell Death Dis*.
649 2022;13:666.

28 Pan Z, Zhao R, Li B, *et al.* EWSR1-induced circNEIL3 promotes glioma progression
651 and exosome-mediated macrophage immunosuppressive polarization via stabilizing
652 IGF2BP3. *Mol Cancer*. 2022;21:16.

29 Yang C, Wu S, Mou Z, *et al.* Exosome-derived circTRPS1 promotes malignant
654 phenotype and CD8+ T cell exhaustion in bladder cancer microenvironments. *Mol Ther*
655 *J Am Soc Gene Ther*. 2022;30:1054–70.

30 Zheng Y, Ren S, Zhang Y, *et al.* Circular RNA circWWC3 augments breast cancer
657 progression through promoting M2 macrophage polarization and tumor immune
658 escape via regulating the expression and secretion of IL-4. *Cancer Cell Int*.
659 2022;22:264.

31 Pamudurti NR, Bartok O, Jens M, *et al.* Translation of CircRNAs. *Mol Cell*. 2017;66:9-
661 21.e7.

32 Fang Z, Jiang C, Li S. The Potential Regulatory Roles of Circular RNAs in Tumor
663 Immunology and Immunotherapy. *Front Immunol*. 2021;11.

33 Li H, Xie M, Wang Y, *et al.* riboCIRC: a comprehensive database of translatable
665 circRNAs. *Genome Biol*. 2021;22:79.

34 Feng J, Chen W, Dong X, *et al.* CSCD2: an integrated interactional database of
667 cancer-specific circular RNAs. *Nucleic Acids Res*. 2022;50:D1179–83.

35 Chen Y, Yao L, Tang Y, *et al.* CircNet 2.0: an updated database for exploring circular
669 RNA regulatory networks in cancers. *Nucleic Acids Res*. 2022;50:D93–101.

36 Yang M, Miao Y-R, Xie G-Y, *et al.* ICBatlas: A comprehensive resource for depicting
671 immune checkpoint blockade therapy characteristics from transcriptome profiles.
672 *Cancer Immunol Res*. 2022;CIR-22-0249.

37 Charoentong P, Finotello F, Angelova M, *et al.* Pan-cancer Immunogenomic Analyses
674 Reveal Genotype-Immunophenotype Relationships and Predictors of Response to
675 Checkpoint Blockade. *Cell Rep*. 2017;18:248–62.

38 Zhang W, Liu Y, Min Z, *et al.* circMine: a comprehensive database to integrate, analyze
677 and visualize human disease–related circRNA transcriptome. *Nucleic Acids Res*.
678 2022;50:D83–92.

39 Fan C, Lei X, Tie J, *et al.* CircR2Disease v2.0: An Updated Web Server for
Experimentally Validated circRNA–disease Associations and Its Application. *Genomics*

Proteomics Bioinformatics. 2022;20:435–45.

40   Wu W, Ji P, Zhao F. CircAtlas: an integrated resource of one million highly accurate
circular RNAs from 1070 vertebrate transcriptomes. *Genome Biol*. 2020;21:101.

41   Dudekula DB, Panda AC, Grammatikakis I, *et al.* CircInteractome: A web tool for
exploring circular RNAs and their interacting proteins and microRNAs. *RNA Biol*.
2016;13:34–42.

42   Huang W, Ling Y, Zhang S, *et al.* TransCirc: an interactive database for translatable
circular RNAs based on multi-omics evidence. *Nucleic Acids Res*. 2021;49:D236–42.

43   Ruan H, Xiang Y, Ko J, *et al.* Comprehensive characterization of circular RNAs in ~
1000 human cancer cell lines. *Genome Med*. 2019;11:55.

44   Dong R, Ma X-K, Li G-W, *et al.* CIRCpedia v2: An Updated Database for
Comprehensive Circular RNA Annotation and Expression Comparison. *Genomics*
Proteomics Bioinformatics. 2018;16:226–33.

45   Glažar P, Papavasileiou P, Rajewsky N. circBase: a database for circular RNAs. *RNA*.
2014;20:1666–70.

46   Zeng Z, Wong CJ, Yang L, *et al.* TISMO: syngeneic mouse tumor database to model
tumor immunity and immunotherapy response. *Nucleic Acids Res*. 2022;50:D1391–7.

47   Miao Y, Zhang Q, Lei Q, *et al.* ImmuCellAI: A Unique Method for Comprehensive T-
Cell Subsets Abundance Prediction and its Application in Cancer Immunotherapy. *Adv*
Sci. 2020;7:1902880.

48   Patil NS, Nabet BY, Müller S, *et al.* Intratumoral plasma cells predict outcomes to PD-
L1 blockade in non-small cell lung cancer. *Cancer Cell*. 2022;40:289-300.e4.

49   Bellmunt J, Hussain M, Gschwend JE, *et al.* Adjuvant atezolizumab versus observation
in muscle-invasive urothelial carcinoma (IMvigor010): a multicentre, open-label,
randomised, phase 3 trial. *Lancet Oncol*. 2021;22:525–37.

50   Cui C, Xu C, Yang W, *et al.* Ratio of the interferon-γ signature to the
immunosuppression signature predicts anti-PD-1 therapy response in melanoma. *Npj*
Genomic Med. 2021;6:7.

51   Liu S, Knochelmann HM, Lomeli SH, *et al.* Response and recurrence correlates in
individuals treated with neoadjuvant anti-PD-1 therapy for resectable oral cavity
squamous cell carcinoma. *Cell Rep Med*. 2021;2:100411.

52   Grasso CS, Tsoi J, Onyshchenko M, *et al.* Conserved Interferon-γ Signaling Drives
Clinical Response to Immune Checkpoint Blockade Therapy in Melanoma. *Cancer Cell*.
2020;38:500-515.e3.

53   Motzer RJ, Banchereau R, Hamidi H, *et al.* Molecular Subsets in Renal Cancer
Determine Outcome to Checkpoint and Angiogenesis Blockade. *Cancer Cell*.
2020;38:803-817.e4.

54   Gide TN, Quek C, Menzies AM, *et al.* Distinct Immune Cell Populations Define
Response to Anti-PD-1 Monotherapy and Anti-PD-1/Anti-CTLA-4 Combined Therapy.
Cancer Cell. 2019;35:238-255.e6.

55   Mariathasan S, Turley SJ, Nickles D, *et al.* TGFβ attenuates tumour response to PD-L1
blockade by contributing to exclusion of T cells. *Nature*. 2018;554:544–8.

56   McDermott DF, Huseni MA, Atkins MB, *et al.* Clinical activity and molecular correlates
of response to atezolizumab alone or in combination with bevacizumab versus sunitinib
in renal cell carcinoma. *Nat Med*. 2018;24:749–57.

57   Miao D, Margolis CA, Gao W, *et al.* Genomic correlates of response to immune
checkpoint therapies in clear cell renal cell carcinoma. *Science*. 2018;359:801–6.

58   Auslander N, Zhang G, Lee JS, *et al.* Robust prediction of response to immune
checkpoint blockade therapy in metastatic melanoma. *Nat Med*. 2018;24:1545–9.

59   Riaz N, Havel JJ, Makarov V, *et al.* Tumor and Microenvironment Evolution during

Immunotherapy with Nivolumab. *Cell*. 2017;171:934-949.e16.

60 Hugo W, Zaretsky JM, Sun L, *et al.* Genomic and Transcriptomic Features of
Response to Anti-PD-1 Therapy in Metastatic Melanoma. *Cell*. 2016;165:35–44.

61 Snyder A, Makarov V, Merghoub T, *et al.* Genetic Basis for Clinical Response to CTLA-
4 Blockade in Melanoma. *N Engl J Med*. 2014;371:2189–99.

62 Tumeh PC, Harview CL, Yearley JH, *et al.* PD-1 blockade induces responses by
inhibiting adaptive immune resistance. *Nature*. 2014;515:568–71.

63 Berger MF, Hodis E, Heffernan TP, *et al.* Melanoma genome sequencing reveals
frequent PREX2 mutations. *Nature*. 2012;485:502–6.

64 Vos JL, Burman B, Jain S, *et al.* Nivolumab plus ipilimumab in advanced salivary gland
cancer: a phase 2 trial. *Nat Med*. 2023;1–13.

65 Kim ST, Cristescu R, Bass AJ, *et al.* Comprehensive molecular characterization of
clinical responses to PD-1 inhibition in metastatic gastric cancer. *Nat Med*.
2018;24:1449–58.

66 Nguyen VP, Campbell KM, Nowicki TS, *et al.* A Pilot Study of Neoadjuvant Nivolumab,
Ipilimumab, and Intralesional Oncolytic Virotherapy for HER2-negative Breast Cancer.
Cancer Res Commun. 2023;3:1628–37.

67 Rosenbaum E, Seier K, Bradic M, *et al.* Immune-related Adverse Events after Immune
Checkpoint Blockade–based Therapy Are Associated with Improved Survival in
Advanced Sarcomas. *Cancer Res Commun*. 2023;3:2118–25.

68 Roper N, Velez MJ, Chiappori A, *et al.* Notch signaling and efficacy of PD-1/PD-L1
blockade in relapsed small cell lung cancer. *Nat Commun*. 2021;12:3880.

69 Gettinger S, Choi J, Hastings K, *et al.* Impaired HLA Class I Antigen Processing and
Presentation as a Mechanism of Acquired Resistance to Immune Checkpoint Inhibitors
in Lung Cancer. *Cancer Discov*. 2017;7:1420–35.

70 Liu D, Lin J-R, Robitschek EJ, *et al.* Evolution of delayed resistance to immunotherapy
in a melanoma responder. *Nat Med*. 2021;27:985–92.

71 Zhang X-O, Dong R, Zhang Y, *et al.* Diverse alternative back-splicing and alternative
splicing landscape of circular RNAs. *Genome Res*. 2016;26:1277–87.

72 Zhang J, Chen S, Yang J, *et al.* Accurate quantification of circular RNAs identifies
extensive circular isoform switching events. *Nat Commun*. 2020;11:90.

73 Memczak S, Jens M, Elefsinioti A, *et al.* Circular RNAs are a large class of animal
RNAs with regulatory potency. *Nature*. 2013;495:333–8.

74 Westholm JO, Miura P, Olson S, *et al.* Genome-wide Analysis of Drosophila Circular
RNAs Reveals Their Structural and Sequence Properties and Age-Dependent Neural
Accumulation. *Cell Rep*. 2014;9:1966–80.

75 Sun J, Zhang H, Wei W, *et al.* Regulation of CD8+ T cells infiltration and
immunotherapy by circMGA/HNRNPL complex in bladder cancer. *Oncogene*.
2023;42:1247–62. doi: 10.1038/s41388-023-02637-2

76 Pisignano G, Michael DC, Visal TH, *et al.* Going circular: history, present, and future of
circRNAs in cancer. *Oncogene*. Published Online First: 16 August 2023. doi:
10.1038/s41388-023-02780-w

77 Horak P, Fröhling S, Glimm H. Integrating next-generation sequencing into clinical
oncology: strategies, promises and pitfalls. *ESMO Open*. 2016;1:e000094.

78 Carroll TM, Chadwick JA, Owen RP, *et al.* Tumor monocyte content predicts
immunochemotherapy outcomes in esophageal adenocarcinoma. *Cancer Cell*.
2023;41:1222-1241.e7.

79 Vromman M, Anckaert J, Bortoluzzi S, *et al.* Large-scale benchmarking of circRNA
detection tools reveals large differences in sensitivity but not in precision. *Nat Methods*.
2023;20:1159–69. doi: 10.1038/s41592-023-01944-6

80 Law CW, Chen Y, Shi W, *et al.* voom: precision weights unlock linear model analysis
tools for RNA-seq read counts. *Genome Biol.* 2014;15:R29.
81 Robinson MD, McCarthy DJ, Smyth GK. edgeR□: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics.*
2010;26:139–40.
82 Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol.* 2014;15:550.
83 Zeng D, Ye Z, Shen R, *et al.* IOBR: Multi-Omics Immuno-Oncology Biological
Research to Decode Tumor Microenvironment and Signatures. *Front Immunol.*
2021;12.
84 Newman AM, Liu CL, Green MR, *et al.* Robust enumeration of cell subsets from tissue
expression profiles. *Nat Methods.* 2015;12:453–7.
85 Yoshihara K, Shahmoradgoli M, Martínez E, *et al.* Inferring tumour purity and stromal
and immune cell admixture from expression data. *Nat Commun.* 2013;4:2612.
86 Finotello F, Mayer C, Plattner C, *et al.* Molecular and pharmacological modulators of
the tumor immune contexture revealed by deconvolution of RNA-seq data. *Genome*
Med. 2019;11:34.
87 Li B, Liu JS, Liu XS. Revisit linear regression-based deconvolution methods for tumor
gene expression data. *Genome Biol.* 2017;18:127.
88 Becht E, Giraldo NA, Lacroix L, *et al.* Estimating the population abundance of tissue-
infiltrating immune and stromal cell populations using gene expression. *Genome Biol.*
2016;17:218.
89 Aran D, Hu Z, Butte AJ. xCell: digitally portraying the tissue cellular heterogeneity
landscape. *Genome Biol.* 2017;18:220.
90 Racle J, de Jonge K, Baumgaertner P, *et al.* Simultaneous enumeration of cancer and
immune cell types from bulk tumor gene expression data. *eLife.* 2017;6:e26476.

807
808
809 **Table Legends**
810
811 **Table 1. Comparison between TCCIA and other similar database resources.**
812
813 **Supplementary Table1. TCCIA cohort summary.**
814 **Abbr.** BLCA, Bladder urothelial carcinoma; HNSC, Head and neck squamous cell
815   carcinoma; KIRC, Kidney renal clear cell carcinoma; NSCLC, Non-small lung cancer;
816   SCLC, Small lung cancer; SKCM, Skin cutaneous melanoma; SGC, Salivary gland
817   carcinoma; BRCA, Breast cancer; SARC, Sarcoma; STAD, Stomach Adenocarcinoma.
818
819 **Figure Legends**
820
821 **Figure 1. Overview of TCCIA.**
822
Figure 2. Content of TCCIA. (**A**) Inclusion of cancer types, cohort and publication years in
this study. **Abbr.** BLCA, Bladder urothelial carcinoma; HNSC, Head and neck squamous
cell carcinoma; KIRC, Kidney renal clear cell carcinoma; NSCLC, Non-small lung cancer;
SCLC, Small lung cancer; SKCM, Skin cutaneous melanoma; SGC, Salivary gland
carcinoma; BRCA, Breast cancer; SARC, Sarcoma; STAD, Stomach Adenocarcinoma. (**B-**
C) The number of detected circRNAs, length of circRNAs, mean (back-splicing junction
count) BSJ and mean (counts per million) CPM in different cancer types (**B**) or cohort (**C**).
(**D**) UMAP plot of all samples, colored by cancer types.
831

Figure 3. A standard exploration and analysis pipeline within TCCIA.
(A) Users need to navigate to the Repository page and select the clinical cancer dataset(s) of their interest. On the leftmost panel, users can filter datasets based on various parameters. **(B)** Users can choose an appropriate analysis strategy according to their needs (core analysis steps are listed on the rightmost panel). The analysis typically consists of four steps: I. Select the dataset(s). II. Choose the CircRNA ID or Host Gene of interest. III. Adjust various parameters such as Test Type, Color Selection, etc. IV. Click the Submit button to obtain the analysis and visualization results. **(C)** Users can browse various information about all CircRNAs included in TCCIA, and at the same time, they can use the ID Converter to transform their interested CircRNAs into other CircRNA Database IDs. **(D)** Users can use the provided options to save result images in PDF or PNG format, with the desired dimensions.

Figure 4. circTMTC3 predicts poor survival and non-response to checkpoint immunotherapy in the melanoma cohorts of Gide et al. and Berger et al. (**A**) Analysis panel for analyzing and visualizing associations between circRNAs and patient survival. Here, we present the data and analysis settings used to generate the plot shown in (**D**). (**B-C**) CircTMTC3 predicts poor survival in (**B**) Gide et al. cohort and (**C**) Berger et al. cohort. (**D-E**) A circRNA isoform of circTMTC3, TMTC3:+:chr12:88148287:88176319, predicts poor survival in (**D**) Gide et al. cohort and (**E**) Berger et al. cohort. Red, group of high circRNA expression. Blue, group of low circRNA expression. (**F-G**) Volcano plots showing differential expressed circRNAs between checkpoint immunotherapy response and nonresponse patients using the approaches Limma Voom, DESeq2, and edgeR in (**F**) Gide et al. cohort and (**G**) Berger et al. cohort.

Supplementary Figure 1. CircRNA identification and overlap analysis of TCCIA cohorts. (**A**)The number of detected circRNAs in each cohort by different method. (**B**) t-SNE plot of all samples, colored by cohorts.
Abbr. BLCA, Bladder urothelial carcinoma; HNSC, Head and neck squamous cell carcinoma; KIRC, Kidney renal clear cell carcinoma; NSCLC, Non-small lung cancer; SCLC, Small lung cancer; SKCM, Skin cutaneous melanoma; SGC, Salivary gland carcinoma; BRCA, Breast cancer; SARC, Sarcoma; STAD, Stomach Adenocarcinoma.

Supplementary Figure 2. circMGA as a predictive biomarker for adjuvant immunotherapy in bladder cancer. (**A**) Overall survival (OS) of bladder cancer patients (all patients, pathological tumor stage ≤pT2, pT3/4) treated with adjuvant immunotherapy was analyzed by Kaplan–Meier curves. Red, group of high circRNA expression. Blue, group of low circRNA expression. We used the maximally selected rank statistics method to determine the cutoff. The minimal proportion for optimal cutoff was set as 0.4. P value was calculated using a log-rank test. (B) Overall survival (OS) of bladder cancer patients (all patients, early-stage tumors(≤pT2) and advanced-stage tumors (pT3/4)) treated with adjuvant immunotherapy was analyzed by Kaplan–Meier curves. Red, group of high circRNA expression. Blue, group of low circRNA expression. We used the maximally selected rank statistics method to determine the cutoff. The minimal proportion for optimal cutoff was set as 0.4. P value was calculated using a log-rank test. (C) Boxplot visualizing the association of circMGA subgroups for patients receiving adjuvant immunotherapy with CD8 T cells signature.