MiDAS 5: Global diversity of bacteria and archaea in anaerobic digesters

Authors: Morten Kam Dahl Dueholm[1,*], Kasper Skytte Andersen[1], Anne-Kirstine C.
Petersen[1], Vibeke Rudkjøbing[1], MiDAS Global Consortium for Anaerobic Digesters[**]
& Per Halkjær Nielsen[1,*]

Affiliations:

[1]Center for Microbial Communities, Department of Chemistry and Bioscience,
Aalborg University, Aalborg, Denmark. [**]A list of authors and their affiliations
appears at the end of the paper.

[*]Correspondence to: Per Halkjær Nielsen, Center for Microbial Communities,
Department of Chemistry and Bioscience, Aalborg University, Fredrik Bajers Vej 7H,
9220 Aalborg, Denmark; Phone: +45 2173 5089; Fax: Not available; E-mail:
phn@bio.aau.dk or Morten Kam Dahl Dueholm, Center for Microbial Communities,
Department of Chemistry and Bioscience, Aalborg University, Fredrik Bajers Vej 7H,
9220 Aalborg, Denmark; Phone: (+45) 9940 8508; Fax: Not available; E-mail:
md@bio.aau.dk

19
20   **Running title:** Global microbiota of anaerobic digesters

Abstract

Anaerobic digestion represents a key biotechnology for the transformation of organic waste into renewable energy (biogas) and relies on complex microbial communities that work in concert to degrade the complex substrates into methane and carbon dioxide. Here, we sequenced more than half a million high-quality, full-length 16S rRNA gene sequences from 285 full-scale anaerobic digesters (ADs) across the world to expand our knowledge about diversity and function of the bacteria and archaea in ADs. The sequences were processed into full-length 16S rRNA amplicon sequence variants (FL-ASVs), which were added to the MiDAS 4 database for bacteria and archaea in wastewater treatment systems to create 'MiDAS 5'. The expansion of the MiDAS database significantly increased the coverage for bacteria and archaea in ADs worldwide, leading to an improved rate of genus and species-level classification. Using MiDAS 5, we carried out an amplicon-based, global-scale microbial community profiling of the sampled ADs using three common sets of primers targeting different regions of the 16S rRNA gene in bacteria and/or archaea. We revealed how environmental conditions and biogeography shape the AD microbiota. We also identify core and conditionally rare or abundant taxa, encompassing less than 1000 genera and species. These represent 83-99% and 10-64% of the accumulated read abundance respectively, across samples depending on the amplicon primers used. Finally, we examined the global diversity of functional groups with known importance for the anaerobic digestion process. Our online global MiDAS Field Guide presents the data generated in this study and summarizes present knowledge about all taxa.

Introduction:

Anaerobic digestion has gained attention as an important, sustainable biotechnology as it provides a number of benefits that align with the goals of sustainability. It can help to produce renewable energy (biogas) from organic waste such as manure, food waste, and sludge from wastewater treatment plants (WWTPs) [1,2]. The anaerobic digestion process also reduces pathogens and the amount of organic waste that is sent to landfills, thereby reducing methane emissions and supporting sustainable waste management practices [1]. Finally, the fertilizer that is produced as a byproduct of anaerobic digestion can be used to support sustainable agriculture, reducing the need for synthetic fertilizers that can have negative environmental impacts [3,4].

The anaerobic digestion process relies on the microbial degradation and conversion of organic matter, which requires a complex interplay between several functional guilds. These include hydrolyzing, acidogenic, and acetogenic syntrophic bacteria as well as methanogenic archaea [5]. The taxonomy is poorly characterized for many of the microorganisms in anaerobic digesters (ADs), and even among the most abundant taxa many lack genus- or species-level classifications [6]. To optimize performance, a comprehensive knowledge about microbial immigration/competition, environmental/operational conditions, and taxonomy is essential [7–9]. Recent microbial surveys have increased our knowledge about the anaerobic digestion process [7,10–14]. However, sharing knowledge across studies is still hindered by the absence of standardized protocols and a common reference database with a unifying taxonomy [15,16]. To facilitate collaboration and knowledge sharing, it is essential to establish these standard protocols and resources.

The Microbial Database for Activated Sludge and Anaerobic Digesters (MiDAS) project was established as an open-source platform for sharing updated knowledge about the physiology and ecology of the important microorganisms present in engineered ecosystems of activated sludge plants, ADs, and related WWTPs [15–18]. MiDAS provides standardized protocols for microbial profiling of microbes in wastewater treatment systems [19], an ecosystem-specific full-length 16S rRNA gene reference database [16,18], and a field guide where knowledge about the specific genera are stored and shared (https://www.midasfieldguide.org).

The MiDAS 16S rRNA gene reference database was created based on millions of high-quality, chimera-free, full-length 16S rRNA genes resolved into amplicon sequence variants (ASVs) and classified using automated taxonomy assignment (AutoTax) [6,16,18]. AutoTax provides a comprehensive seven-rank taxonomy (kingdom to species-level) to all reference sequences based on the most recent version of the SILVA SSURef 99 NR taxonomy, and adds a robust placeholder taxonomy for lineages without an official taxonomy [6]. The placeholder taxa can easily be distinguished by their names (midas_x_y, where x indicates the taxonomic rank and y is a numerical identifier) and

86   provides an important opportunity to study unclassified taxa along classified taxa at
87   different taxonomic ranks. The placeholder taxonomy should not be seen as a
88   replacement for proper taxonomic classifications but can pinpoint important lineages
89   that should be studied in depth using phylogenomics [20–24].
90
91   The MiDAS 16S rRNA gene reference database (MiDAS 4.8.1) currently contains
92   reference sequences from anaerobic digesters located at WWTPs in Denmark and
93   WWTPs worldwide [18]. However, it may not provide comprehensive coverage for all
94   important microbes found in ADs treating other types of waste or in other locations.
95
In this study, we introduce MiDAS 5, an updated version of MiDAS 4 expanded with
more than half a million high-quality, full-length archaeal and bacterial 16S rRNA gene
sequences from 285 ADs worldwide treating different types of biowaste. We carried
out a global survey of ADs using three commonly used short-read amplicon primer sets
targeting bacteria (V1-V3), archaea (V3-V5), and both (V4). This data was used in
combination with MiDAS 5 to i) link the global diversity of bacteria and archaea to
biogeography and environmental factors, ii) identify important core taxa, and iii)
uncover the global diversity within selected functional guilds. The results provide a
solid foundation for future research on AD microbiology.
105
Results and Discussion:
The MiDAS Global Consortium for Anaerobic Digesters was established in 2018 to
coordinate the sampling and collection of metadata from ADs worldwide
(Supplementary Data 1). Samples were obtained in duplicates from 285 ADs in 188
cities in 19 countries on five continents (Fig. 1a). The majority of the ADs treated
surplus sludge from wastewater treatment plants (60.0%) (Fig. 1b). However, ADs
treating food waste (6.7%), industrial waste (7.4%), and manure (4.9%) were also
included in the survey. Most of the ADs were mesophilic (85.6%), few were
thermophilic (6.0%), and some missed temperature data (8.4%). The main digester
technology used was completely mixed reactors (67.7%) followed by two-stage
reactors (12.6%). A few upflow anaerobic sludge blanket (UASB) and plug-flow
reactors were also sampled to expand the diversity of digester types.
118
Expanding the MiDAS database with reference from global ADs
To expand the MiDAS database with sequences from ADs across the globe, we applied
high-fidelity, full-length 16S rRNA gene sequencing on all samples collected in this
study [25]. More than half a million full-length 16S rRNA gene sequences, representing
both bacteria and archaea, were obtained after quality filtering and primer trimming.
The sequences were processed with AutoTax to produce full-length 16S rRNA gene
ASVs (FL-ASVs), and these were added to the MiDAS 4.8.1 database leading to an
expanded database (MiDAS 5) with a total of 120,408 unique, chimera-free FL-ASV
reference sequences.

Fig. 1: Sampling of anaerobic digesters (ADs) across the world. a) Geographical distribution of ADs
included. b) Distribution of primary substrates. c) Distribution of digester technologies. CSTR:
Continuous stirred-tank reactor; TSAD: Two-stage anaerobic digestion; UASB: Upflow anaerobic
sludge blanket: d) Distribution of digester temperatures. The values next to the bars are the number of
ADs in each group.

Table 1: Sequence novelty of FL-ASVs obtained in this study. Sequence novelty was determined based
on the percentage identity between each of the 30,246 new FL-ASV and their closest relative in the
databases indicated and identity thresholds for each taxonomic rank proposed by Yarza *et al.* [26] shown
in the parentheses.

	SILVA 138.1 SSURef NR99		MiDAS 4.8.1	
	Sequences	Percentage	Sequences	Percentage
Phylum (<75.0%)	28	0.09%	67	0.22%
Class (<78.5%)	55	0.18%	186	0.61%
Order (<82.0%)	112	0.37%	462	1.53%
Family (<86.5%)	354	1.17%	1,483	4.90%
Genus (<94.5%)	5,240	17.32%	9,419	31.14%
Species (<98.7%)	15,806	52.26%	16,863	55.75%

The novelty of the 30,246 new FL-ASVs were determined based on the percent identity
shared with their closest relatives in the SILVA 138.1 SSURef NR99 and MiDAS 4.8.1
database using the threshold for each taxonomic rank proposed by Yarza *et al.* [26] (Table
1). 17% and 31% of the new FL-ASV lacked genus-level homologs (≥94.5% identity)
and 52% and 56% were without species-level homologs (≥98.7% identity) in SILVA
138.1 and MiDAS 4, respectively. This suggests a substantial increase in the diversity
within the MiDAS database.

MiDAS 5 introduces many new taxa

To investigate how the new FL-ASVs affected the taxonomic diversity in the MiDAS database, we determined the number of additional taxa introduced at different taxonomic ranks (Table 2). A substantial increase in diversity was observed with the addition of 2,770 new genera (29.2% increase) and 8,858 new species (28.3% increase). However, many additional taxa were also introduced at higher taxonomic ranks as highlighted with the introduction of six more bacterial and five more archaeal phyla previously known from the SILVA taxonomy. In addition, we observed nine deep branching lineages classified as novo phyla, however phylogenetic analysis revealed that these likely represent mitochondrial sequences. The largest percentage of the new FL-ASVs (42.8%) were found within the Firmicutes (Supplementary Fig. 1a). Firmicutes often occur in high abundance in ADs, where they are involved in fermentation and thereby directly stimulate biogas yields [7,10,13,25]. A closer look into the expanded diversity within the Firmicutes revealed that new FL-ASVs were associated with several families (Supplementary Fig. 1b), including Hungateiclostridiaceae (1,324 FL-ASVs), Lachnospiraceae (788 FL-ASVs), Peptostreptococcales-Tissierellales Family_XI (763 FL-ASVs), Christensenellaceae (754 FL-ASVs), Caldicoprobacteraceae (620 FL-ASVs), and Syntrophomonadaceae (555 FL-ASVs). The Syntrophomonadaceae are of special relevance, as this family includes several syntrophic fatty acid degrading bacteria, which are often the metabolic bottleneck in the overall ADs process [27,28].

Table 2: New taxa introduced with MiDAS 5. The number of new taxa represent unique taxa at the different taxonomic ranks that were not part of MiDAS 4.8.1 [18] and includes both official taxonomic names and de novo placeholder names provided by AutoTax [6].

	Total taxa	New taxa	Increase (%)
Phylum	105	20	23.5%
Class	259	37	16.7%
Order	727	104	16.7%
Family	2,212	360	19.4%
Genus	12,254	2,770	29.2%
Species	40,207	8,858	28.3%

MiDAS 5 provides improved coverage and classifications for AD microbiota

The performance of the MiDAS 5 database was evaluated based on three ASV-resolved, short-read, 16S rRNA gene amplicon datasets generated from the AD samples collected in this study (Fig. 2). The V1-V3 amplicons include only bacteria and provide high phylogenetic resolution. However, the primers targeting this region have a lower coverage for the known bacterial diversity according to in silico evaluations [6,29]. The V4 amplicons include both bacteria and some archaeal lineages and are commonly used due to a very good coverage of the known bacterial diversity. However, the amplicons have a weaker phylogenetic resolution compared to V1-V3, which in many cases prevent species-level classifications [6,29]. The V3-V5 amplicons cover mainly archaea and have previously been used to describe their diversity in ADs [7,10].

Our first evaluation was based on stringent mapping of the short-read ASVs to MiDAS 5 and other commonly applied reference databases. This allowed us to determine the percent identity shared between each ASV and their best hit in each reference database, which was subsequently used to calculate the percentage of ASVs with high-identity hits (≥99% identity) per sample and database after removing the rare biosphere (ASV with <0.01% relative abundance), known to be enriched in non-growing organisms and environmental DNA [10]. MiDAS 5 performed exceptionally well for bacteria with high-identity hits of 94.5%±4.6% (mean±SD) for V1-V3 and 96.4%±2.8% for V4 ASVs, compared to 69.9%±19.7% and 73.6%±16.1% for MiDAS 4, and 61.9%±9.2% and 78.8%±7.0% for SILVA v.138.1 (Fig. 2). For the V3-V5 archaeal dataset, an increase in coverage was observed from 34.1%±8.7% with MiDAS 4 to 56.0%±11.8% with MiDAS 5. However, the SILVA database provides even better coverage (66.4%±12.0%). The 16S rRNA gene reference database derived from the genome taxonomy database (GTDB) performed poorly, which relates to the fact that many metagenome-assembled genomes (MAGs) created from short-read data lack 16S rRNA genes due to assembly issues [30,31]. The lower coverage seen for archaea is likely related to reduced sequencing efforts and difficulties designing universal primers for archaeal full-length 16S rRNA gene sequencing [32,33]. Because the sampling of ADs was directed towards mesophilic digesters treating surplus sludge from WWTPs, we also evaluated the MiDAS 5 coverage for ADs treating different primary substrates and temperatures (Supplementary Fig. 2). MiDAS 5 gave very good coverage for all sample types supporting the general applicability of the reference database for ADs.

Fig. 2: Database evaluation based on short-amplicon data from this study. The ASVs for each of the samples were filtered based on their relative abundance (only ASVs with ≥0.01% relative abundance were kept) before the analyses. The percentage of the microbial community represented by the remaining ASVs after the filtering was 95.43% ± 2.38% (mean±SD) for V1-V3 amplicons (only bacteria), 99.69% ± 0.13% for V3-V5 amplicons (mainly archaea), and 97.68% ± 1.97% for V4 amplicons (bacteria and archaea) across samples. High-identity (≥99%) hits were determined by stringent mapping of ASVs to each reference database. Classification of ASVs was done using the SINTAX classifier. The violin and box plot represent the distribution of percent of ASVs with high-identity hits or genus/species-level classifications for each database across n=469 (V1-V3), n=273 (V3-V5), and n=470 (V4) duplicate, biologically independent samples duplicates. Box plots indicate median (middle line), 25th, 75th percentile (box), and the min and max values after removing outliers based on 1.5x interquartile range (whiskers). Outliers have been removed from the box plots to ease visualization. Different colors are used to distinguish the different databases: GTDB SSU_r202, SILVA 138.1 SSURef NR99, MiDAS 4.8.1, and MiDAS 5.0.

Our second database evaluation was based on the classification of ASVs from each amplicon dataset using the SINTAX classifier (Fig. 2). We found that MiDAS 5 greatly improved the rates of genus-level classification (96.1%±1.9% for V1-V3, 91.2%±2.9% for V4, and 80.6%±6.0% for V3-V5) compared to MiDAS 4 (81.7%±14.2 for V1-V3, 78.6%±9.7% for V4, and 75.0%±8.1% for V3-V5), and that these rates of classification were much higher than those obtained with SILVA (30.7%±6.0 for V1-V3, 43.8%±7.8% for V4, and 42.7%±9.2% for V3-V5) and GTDB (29.0%±6.9 for V1-V3, 31.7%±6.9% for V4, and 62.7%±7.2% for V3-V5). Analysis of species-level classifications revealed similar improvements with MiDAS 5 for bacteria (Fig. 2). However, a decrease in species-level classifications was observed between MiDAS 4 and 5 for the archaeal V3-V5 dataset. We hypothesize that this effect relates to over-classifications with MiDAS 4 due to the lack of appropriate reference sequences in MiDAS 4.

Finally, we investigated if the additional reference sequences introduced in MiDAS 5 could improve classification of amplicon data from WWTPs based on data from the MiDAS global sampling of WWTPs [18] and the Global Water Microbiome Consortium project [34] (Supplementary Fig. 3). The improvements were insignificant, highlighting that most of the added references originated from AD-specific taxa.

Fig. 3: Ecosystem-specific primer coverage for all archaea and bacteria and the 20 most diverse bacterial and all archaeal phyla based on unique FL-ASVs. The coverage was determined as the percentage of FL-ASVs in the MiDAS 5 database with perfect hits for both forward and reverse primers. The primer pairs marked in red were used in the current study. Detailed information of all primer pairs and coverage information for all taxa in MiDAS 5 are provided in Supplementary Data 2. *Only the reverse primer was evaluated for these primer pairs because the forward primer was used to create the MiDAS 5 reference sequences. The coverage might therefore be overestimated.

Evaluation of 16S rRNA gene amplicon primers for community profiling of ADs

The comprehensive ASV-resolved MiDAS 5 database provides a unique opportunity to determine the theoretical coverage of commonly applied 16S rRNA gene amplicon primer pairs for bacteria and archaea in ADs (Fig. 3). This information is highly valuable when designing experiments, especially if targeting specific taxa. Accordingly, we have determined the theoretical coverage for several commonly applied primer pairs for all kingdom to species-level taxa in MiDAS 5 (Supplementary Data 2). We found a fairly low coverage of the V1-V3 primer pair (perfect hits for ≤79% of the bacterial FL-ASVs), which we commonly use due to its high phylogenetic resolution [6,18]. We should therefore expect a bias when using this primer pair, although it has previously shown good correlation with quantitative FISH results [19]. The V4c primer used here and in the Earth Microbiome project [35] showed good coverage for both bacteria (perfect hits for 87% of the FL-ASVs) and archaea (perfect hits for 98% of the FL-ASVs). It may therefore be applied for profiling the entire prokaryotic diversity in ADs. However, it should be noted that the V4 amplicon has a lower phylogenetic resolution which prevents species-level classification for many genera [6,29]. Interestingly, we observed an even higher coverage for both bacteria (perfect hits for 90% of the FL-ASVs) and archaea (perfect hits for 99% of the FL-ASVs) with the V4-V5a primer pair. Because this longer amplicon also carries a stronger phylogenetic resolution, we would recommend this primer pair for general profiling of ADs in future studies. The V3-V5b primer pair, used here to target archaea, had a good coverage for archaea, although not as good as the V4c and V4-V5a universal primers. In most cases, we would therefore recommend the application of universal primers.

Effect of process and environmental factors on the AD microbiota

Alpha diversity analyses showed that the rarefied (10,000 read per sample) ASV richness and inverse Simpsons diversity in ADs were affected mainly by the primary substrate type and the temperature in the ADs (Supplementary Fig. 4). Significantly higher bacterial richness and diversity were observed for ADs treating surplus sludge from WWTPs compared to the other types of substrates. This effect likely reflects the extensive immigration of bacteria into the ADs with the surplus sludge [7,10,36]. A higher richness and diversity were observed for bacteria in mesophilic ADs compared to thermophilic ADs. A similar trend has previously been observed for full-scale ADs treating manure [37,38], household waste [39], and surplus sludge from WWTPs [7].

Genus-level taxonomic beta-diversity was used to investigate the effect of process conditions and geography on the overall microbiota in ADs using principal coordinate analysis (PCoA) and permutational multivariate analysis of variance (PERMANOVA) (Fig. 4). We used this approach because many of the important traits are categorical (yes/no) and only conserved at lower taxonomic ranks (genus/species). Furthermore, MiDAS 5 enabled us to classify almost all our ASVs at the genus-level, thereby providing a comprehensive description of the microbiota. We found that the overall

microbial community could be explained mainly by the primary substrate and to a lesser extent temperature, continent, and digester technology (Fig. 4). This trend was observed for both bacteria and archaea. The percentage of total variation explained by each parameter was, except for the primary substrate, low, suggesting that the global AD microbiota represents a continuous distribution rather than distinct states, as also observed for the human gut microbiota [40] and WWTPs [18]. The pronounced effect of the primary substrates highlights that these substrates are distinct and abundant in microbes, particularly in the case of manure and wastewater sludge.

Fig. 4: Effects of process and environmental factors on the anaerobic digester microbiota. Principal coordinate analyses of Bray-Curtis beta-diversity for genera based on V1-V3 (bacteria), V4 (archaea and bacteria), and V3-V5 (archaea) amplicon data. Samples are colored based on metadata. The fraction of variation in the microbial community explained by each variable in isolation was determined by PERMANOVA (Adonis R^2-values). Exact P-values <0.001 could not be confidently determined due to the permutational nature of the test. CSTR: Continuous stirred-tank reactor; TSAD: Two-stage anaerobic digestion; UASB: Upflow anaerobic sludge blanket.

Core and conditional rare or abundant taxa in the global AD microbiota

The global AD microbiota represents a huge microbial diversity. However, most organisms only occur in very low abundance and are therefore unlikely to have any quantitative impact on the overall metabolism and the process performance in ADs. Analysis of core and conditionally rare or abundant taxa (CRAT) is a powerful approach to identify the most important genera and species within a specific ecosystem [18,25,41]. Core taxa are commonly defined based on how frequently specific taxa are observed in samples from a well-defined habitat [42], whereas CRAT are typically present in low abundance but can occasionally become very abundant [43,44]. The CRAT may include taxa related to process disturbances, such as filamentous microbes associated with foam formation, or taxa associated with the degradation of special substrates found in, e.g., industrial waste.

We recently introduced and applied the following core and CRAT definitions in our survey of the global microbiota of wastewater treatment plants: strict core (>0.1% relative abundance in >80% of samples), general core (>0.1% relative abundance in >50% of samples), loose core (>0.1% relative abundance in >20% of samples), and CRAT (not part of the core, but present in at least one sample with a relative abundance >1%) [18]. Here, we applied the same criteria to identify core and CRAT genera and species in our global AD dataset. Because the primary substrate showed a strong effect on the overall microbial community (Fig. 4), we determined the core and CRAT for each individual substrate separately (Supplementary Data 3). Only mesophilic ADs were examined for ADs treating food waste, industrial waste, and manure due to the low number of thermophilic ADs sampled. Both mesophilic and thermophilic digesters were examined for ADs treating wastewater sludge. All three amplicon datasets were examined, and the results combined to reduce the effect of primer bias.

The core analysis revealed that most core genera were uniquely associated with specific primary substrates and temperature range (Fig. 5a). However, there were also a significant number of core genera shared across substrates (Fig. 5a). In contrast, very few core species were shared between ADs treating different primary substrates (Fig. 5b). This fits well with similar results from a study of ADs in Belgium and Luxemburg [13]. To define a most wanted list for bacteria and archaea in ADs globally, we linked each core and CRAT with the best classification across primary substrates and process temperature (Supplementary Data 3). The resulting list contained 506 core genera (71 strict, 122 general, and 313 loose) and 152 CRAT genera. The strict core genera included 11 known methanogens and four known syntrophs (*Smithella*, *Syntrophaceticus*, *Syntrophomonas*, *Syntrophorhabdus*). At the species-level, we identified 597 core (27 strict, 127 general, and 443 loose) and 339 CRAT species. The strict core species included three methanogens (*Methanobrevibacter smithii*, *Methanosphaera stadtmanae* and *Methanothermobacter* midas_s_3958) and one syntroph (*Syntrophomonas* midas_s_90707). It is worth noting that a large fraction

350 of the taxa observed in ADs is unable to grow, but only occurs because the taxa are in
351 high abundance in the feed [7,10,36]. Data from Danish ADs treating wastewater sludge [7]
352 showed that 140 (28%) of the core genera and 21 (3.5%) of the core species were non-
353 growing (<20% of ASVs belonging to the specific taxa were classified as growing), but
354 it is not known whether this also translates to global ADs.
355

Many core and CRAT are only identified due to the MiDAS placeholder taxonomy

A large proportion of core and CRAT identified was classified as MiDAS *de novo* taxa.
At the genus-level, 281/506 (56%) of the core genera and 100/152 (66%) of the CRAT
genera had MiDAS placeholder names, and at the species-level, the proportion was
even higher. Here placeholder names were assigned to 552/597 (92%) of the core
species and 322/339 (95%) CRAT species. These proportions are similar to those
observed for the global microbiota in activated sludge [18] and reveals the importance of
a taxonomic framework that can handle uncultured taxa which have not yet been
officially classified.
365

366
Fig. 5: Core and conditionally rare or abundant taxa (CRAT) in anaerobic digesters globally. a) and b)
UpSet plots displaying the number of shared core genera and species, respectively, across ADs treating
different primary substrates and operating at different temperatures. c) and d) Number of observed genera
and species, respectively, and their abundance in mesophilic ADs treating different primary substrates
based on V4 amplicon data (bacteria and archaea). Values for genera and species are divided into strict
core, general core, loose core, CRAT, other taxa, and unclassified ASVs based on the most wanted list
(Supplementary Data 3). The relative abundance of different groups was calculated based on the mean
relative abundance of individual genera or species across samples. Similar figures for V1-V3 (bacteria
only) and V3-V5 (archaea only) amplicons data can be found in Supplementary Fig. 5.

The global AD microbiota is dominated by core and CRAT taxa

Despite only accounting for a minor fraction of the total diversity in the global AD microbiota, the core and CRAT represented most of the microbes in the ecosystem according to relative amplicon read abundance (Fig. 5c, 5d, Supplementary Fig. 5). The core and CRAT genera accounted for 86-93% (V1-V3), 83-90% (V4), and 95-99% (V3-V5) of the accumulated read abundance in mesophilic ADs depending on primary substrates. The remaining fractions consisted mainly of ASVs unclassified at the genus level, and genera present in very low abundance, presumably with minor importance for the AD performance.

For the species level, the core and CRAT represented 54-64% (V1-V3), 38-47% (V4), and 10-58% (V3-V5) accumulated read abundance depending on the primary substrate. The remaining fractions were mainly composed of ASVs, which could not be classified at the species level, probably due to insufficient phylogenetic resolution in the short-read amplicons [6,29]. The lack of species-level classification was especially pronounced for the archaeal V3-V5 ASVs in ADs treating industrial waste, manure, and wastewater sludge (Fig. 5).

The large relative abundance of core and CRAT in the global AD microbiota suggests that we can explain a substantial part of the metabolic processes in ADs, if we understand the physiology and metabolic potential of these taxa.

Global diversity of archaea reveals new potential methanogens

As methanogenic archaea are ultimately responsible for the generation of methane in ADs, we examined the global diversity of archaea in all samples based on V4 (Fig. 6) and V3-V5 amplicon data (Supplementary Fig. 6). The V4 amplicon data, encompassing both archaea and bacteria, showed that the archaeal reads constituted 3.9%±3.0% for ADs treating food waste, 4.6%±2.8% for manure, 6.0%±2.4% for wastewater sludge, and 14.2%±12.2% for industrial waste. Many of the abundant archaea represented well-known methanogens. However, we also observed several abundant genera, only classified based on the MiDAS placeholder taxonomy, affiliated to orders and families of known methanogens. These include midas_g_90791 and midas_g_8154, which represent new families within the orders Methanomicrobiales and Methanofastidiosales, respectively, and midas_g_90473 and midas_g_93310, representing new genera within Methanomassiliicoccaceae and Methanospirillaceae, respectively. In addition, we observed two abundant MiDAS placeholder genera (midas_g_90791 and midas_g_97217) that represent a new order within the class *Ca.* Bathyarchaeia. Members of his class can have a versatile metabolism, and some encode the key methanogenic enzyme methyl-coenzyme M reductase (MCR) [45,46]. Targeted metagenomics and assembly of MAGs should be applied to confirm the methanogenic potential of these new potential methanogens, and our amplicon datasets provide insight into where these taxa occur in high abundance.

418
Fig. 6: Top 25 archaeal genera based on V4 amplicon data. The percent relative abundance represents
the mean abundance relative to all archaea across a) different primary substrates and temperature range,
and b) different countries considering only mesophilic ADs treating mainly wastewater sludge.

The methanogenic community composition was clearly affected by the primary
substrate and whether the digestion was performed at mesophilic or thermophilic
conditions (Fig. 6a, Supplementary Fig. 6a). The most common methanogens across
substrates and temperatures were *Methanoculleus*, *Methanosarcina*, *Methanothrix*, and
Methanothermobacter, although the latter was only abundant in thermophilic ADs
treating wastewater sludge and food waste and in a single mesophilic reactor treating
food waste, indicating that this digester might use a two-stage thermophilic-mesophilic
configuration.

Because most of our samples originated from mesophilic reactors treating wastewater
sludge, we examined the diversity of methanogens across countries in these ADs (Fig.
6b, Supplementary Fig. 6b). This analysis revealed that the same genera were
dominating across the world. The most common methanogens in these ADs were
Methanothrix, *Methanolinea*, *Methanospirillum*, *Methanoculleus*, and the recently
discovered *Ca*. Methanofastidiosum [47] (Fig. 6b, Supplementary Fig. 6b). Next, we
examined if the methanogens were also conserved at higher phylogenetic resolution.
As many archaeal ASVs could not be classified at the species-level, we examined the
global diversity at the ASV-level (Supplementary Fig.7). We found that the vast
majority of the abundant ASVs occurred globally. This high similarity among
methanogens across the world suggests a great potential for knowledge transfer
between countries.

Among the highly abundant archaea, we also observed an ammoniaoxidizing archaeon
(AOA) from the genus *Ca*. Nitrosocosmicus [48], which was especially abundant in
thermophilic ADs treating food waste. This is very surprising and may indicate that
they also have an anaerobic physiology which should be investigated further. Another
abundant archaeon was the *Ca*. Diapherotrites ADurb.bin253 belonging to the order

Woesearchaeales which are characterized by ultra-small genomes and an anaerobic and
parasitic/fermentation-based lifestyle [49].
452

Global diversity of syntrophic bacteria
Syntrophic bacteria play a vital role in ADs by converting substrates, such as short-
chain fatty acids to acetate, H_2, and formate, which act as substrates or reducing
equivalents for methanogens to produce methane and CO_2 [27,50,51]. This obligately
mutualistic metabolism between the syntrophs and methanogens is critical because
substrate oxidation can only occur and sustain growth of the syntrophs under anaerobic
conditions if the products are kept at a very low concentration by rapid consumption by
the methanogens [50,52]. Due to the fastidious metabolism, syntrophs are usually present
in low abundance, and can easily become the bottleneck in the anaerobic digestion
process [7,8]. Accordingly, we investigated the global diversity of this functional guild in
the ADs sampled (Fig. 7, Supplementary Fig. 8).
464

A clear effect of the primary substrates and digester temperature was observed on the
composition and abundance of syntrophic genera in the digesters (Fig. 7a,
Supplementary Fig. 8a). The most abundant genus across substrates and temperature
was *Syntrophaceticus*, despite being barely detected in ADs treating wastewater sludge.
The type strain of this genus, *S. schinkii* $Sp3^T$, is an acetate-oxidizing syntroph that
thrives, and has a competitive advantage, under high ammonium concentrations (up to
8,400 mgN/L) [53,54]. The lack of *Syntrophaceticus* in ADs treating wastewater sludge
may therefore be explained by lower ammonium concentrations in these ADs
(1,617±4,312 mgN/L, n=145) compared to those treating food waste (2,913±1,681
mgN/L, n=33), and manure (3,449±933 mgN/L, n=18).
475

476
Fig. 7: Global diversity of syntrophs based on V1-V3 amplicon data. The percent relative abundance
represents the mean for genera across a) different primary substrates and temperature range, and b)
different countries considering only mesophilic ADs treating mainly wastewater sludge. Colored circles
next to the genus labels indicate whether the genera have previously been identified as growing in ADs
at Danish WWTPs according to Jiang *et al.* [7]. Blue: >50% of ASVs classified as growing; Yellow: 20-
50% of ASVs classified as growing. Red: <20% of ASVs classified as growing. Gray: No information
available for the specific genus.

Syntrophomonas, the second most abundant genus, was common in all AD types
investigated, indicating a broader ecological niche. Isolated representatives from this
genus can grow syntrophically via β-oxidation of saturated fatty acids of various lengths
(C4-C18, depending on strain) [55–58], and they are therefore likely important for the
conversion of long-chain fatty acids in ADs. Among the abundant syntrophs,
Tepidimicrobium, a member of the order Clostridiales, was also observed in all AD
types except mesophilic ADs treating wastewater sludge. The exact metabolism of
Tepidimicrobium in ADs remains to be determined, however all isolated representatives
can degrade proteinaceous compounds and some species can also use carbohydrates [59].
Furthermore, *Tepidimicrobium* has been proposed to grow syntrophically by direct
interspecies electron transfer (DIET) with *Methanothermobacter* in a process like that
observed for *Geobacter* [60]. Accordingly, it is likely that the *Tepidimicrobium* acts as a
syntrophic primary degrader in the ADs targeting mainly proteins, carbohydrates, and
derivatives.
498
Finally, we observed a high abundance of the genus *Smithella* in mesophilic ADs
treating industrial waste, manure, and wastewater sludge. The type strain *S. propionica*
LYP^T is a propionate oxidizing syntroph, which uses a unique dismutation pathway in
which propionate is first converted to acetate and butyrate, and the latter is hereafter β-
oxidized syntrophically to acetate and hydrogen [61,62]. Calculations of Gibbs free energy
for this special propionate metabolism indicates a higher tolerance toward elevated
hydrogen concentrations [63], which could explain why some *Smithella* prevail in certain
ADs. However, *Smithella* has also been implicated in the syntrophic degradation of
long-chain alkanes [64,65], which could reflect a more versatile metabolism.
508
When investigating geographical diversity of syntrophic fatty acid oxidizing bacteria
in mesophilic ADs treating wastewater sludge, a similar pattern was observed across
countries (Fig. 7b, Supplementary Fig. 8b). *Smithella*, was generally the dominating
syntroph. However, *Syntrophomonas*, *Syntrophorhabdus*, *Ca*. Phosphitivorax, and
Syntrophus also occurred at a fairly high relative abundance in almost all countries.
Isolates of *Syntrophorhabdus,* including the type strain *S. aromaticus* UI^T, are
syntrophic fermenters of aromatic compounds and may accordingly play an important
role in the detoxification of these substrates in ADs [66,67]. *Ca*. Phosphitivorax was
recently discovered as a butyrate degrading syntroph by genome-resolved meta-
transcriptomics in a digester treating wastewater sludge [51], and *Syntrophus* participates
in the degradation of fatty acids and aromatics [68,69]. Overall, the results suggest a
complex syntrophic degradation process, which involves multiple genera with different
substrate specificities.
522
523   To gain additional insight into the global diversity of syntrophs, we also investigated
524   the species-level diversity across mesophilic digesters treating wastewater sludge
525   (Supplementary Fig. 9). We observed a large species diversity among most of the

abundant syntrophic genera. Furthermore, we found that the most abundant species in the ADs were often distinct from the isolated representatives, which prompts for further investigations into the metabolic potential of syntrophs *in situ*.

Global diversity of filamentous bacteria

Foaming is a common operational problem in ADs and has a strong negative impact on process performance resulting in considerable costs. Both abiotic and biotic factors are involved in foaming [70]. The abiotic factors include high loading rates of surfactants (oil, grease, fatty acids, detergent, proteins, and particulate matter) and biosurfactants produced by microbes in the digester [71]. The biotic factors cover increased abundance of hydrophobic, filamentous microorganisms that can interact with, and stabilize, gas bubbles in the foam [70,72]. To gain further insight into potential foam forming microbes, we examined the global diversity of known filamentous bacteria in ADs (Fig. 8, Supplementary Fig. 10).

The diversity and mean relative abundance of known filamentous organisms were generally low in the ADs examined except for those treating wastewater sludge (Fig. 8a, Supplementary Fig. 10). However, the increased diversity and abundance in the latter are to a large extent the result of passive immigration from the fed surplus sludge. However, most of these are unable to grow in the ADs [7] (Fig. 8a, Supplementary Fig. 10). *Anaerolinea*, *Ca*. Brevefilum, and *Trichococcus* were common across ADs treating all primary substrates (Fig. 8a, Supplementary Fig. 10), whereas *Ca*. Microthrix and *Ca*. Promineofilum were mainly observed in ADs treating wastewater sludge. Many of the Chloroflexi genera found here were also observed in a recent meta-analysis of amplicon data from 17 studies representing 62 ADs [73]. Several of the abundant filamentous genera, including *Ca*. Microthrix and *Ca*. Brevefilum, were previously found to correlate with the foaming potential of full-scale digester sludge from mesophilic ADs at WWTPs [72]. *Ca*. Brevefilum seems especially interesting as it grows well in ADs [7,74].

The species-level diversity was generally low for the filamentous bacteria (Supplementary Fig. 11). *Ca*. Brevefilum was dominated by *Ca*. B. fermentans, *Trichococcus* by midas_s_4, *Ca*. Microthrix by *Ca.* M. parvicella and *Ca.* M. subdominans, and *Gordonia* by *G. amarae* and *G. defluvii*. *Ca.* Promineofilum was more diverse with five common species. The low species-level diversity of potential foam-forming bacteria suggests that it may be feasible to develop and implement universal mitigation strategies for these bacteria in ADs worldwide.

563
Fig. 8: Global diversity of known filamentous organisms based on V1-V3 amplicon data. The percent
relative abundance represents the mean for genera across a) different primary substrates and temperature
range, and b) different countries considering only mesophilic ADs treating mainly wastewater sludge.
Colored circles next to the genus labels indicate whether the genera have previously been identified as
growing in ADs at Danish WWTPs according to Jiang *et al.* [7]. Blue: >50% of ASVs classified as growing;
Yellow: 20-50% of ASVs classified as growing. Red: <20% of ASVs classified as growing. Gray: No
information available for the specific genus.

Conclusion and perspectives
MiDAS 5 was made possible thanks to a huge collaborative effort from experts
worldwide, who contributed to the project by sampling and providing metadata for ADs
in their respective countries. Building on the success of its predecessor, MiDAS 4, this
latest expansion covers ASV-resolved, full-length 16S rRNA gene references from
numerous ADs from all parts of the globe covering different operations parameters and
different substrates. This expanded database provides greatly improved coverage for
AD-specific taxa and a strongly needed taxonomy for uncultured lineages, which lacks
official taxonomic classification. As such, it will be an invaluable resource for
researchers and AD professionals, providing them with a common point of reference to
facilitate knowledge sharing and pave the way for a comprehensive understanding of
the AD microbiome.

Our in silico 16S rRNA gene primer evaluation based on the MiDAS 5 database
revealed that the coverage of commonly applied primer pairs varies significantly, with
some having low coverage and potential bias towards certain taxa. Because the primer
coverage was evaluated for all taxa in the MiDAS 5 database at all taxonomic rank, it
provides a solid foundation for designing experiments and targeting specific taxa in
future studies. For general microbial profiling of ADs, we would recommend the use
of the V4a or the V4-V5a primer pairs, as they both show excellent coverage for archaea
and bacteria in the ecosystem.

Although the total microbial diversity in ADs is huge, importantly, we showed that less
than 1000 genera and species accounted for most of the microbes in the AD ecosystem.
By focusing on these abundant and common microbes, we will be able to explain most
of the microbial processes that occur in the anaerobic digestion process. This list, the
"Most wanted" organisms, contain species that should be prime targets for future in situ
studies and genome reconstruction projects.
599
600   The global survey of the ADs microbiota using three different primer pairs provided a
601   unique insight into the global diversity of individual AD taxa and clues into the
602   environmental and operational factors that define their ecological niches. This
603   information will be invaluable in the development of future microbiome management
604   strategies and improved sustainability of the field of anaerobic digestion.
605
To enhance knowledge dissemination, we have updated the MiDAS Field Guide
available at www.midasfieldguide.org. This dynamic resource allows users to delve
into specifics related to the physiology, morphology, and ecology of genera listed in
the MiDAS database. Additionally, it offers country-specific data on the prevalence of
all MiDAS genera and species in WWTPs and ADs. Finally, it provides information on
the availability of fluorescence in situ hybridization probes and reference genomes,
paving the way for subsequent research endeavors.

Methods:

Sampling and metadata collection

To facilitate sampling of ADs worldwide, we established the MiDAS Global Consortium for Anaerobic Digesters, which consists of 25 anaerobic digestion experts in 19 countries. Members of the consortium acted as national sampling coordinators and were in direct contact with the ADs. Two samples were obtained from each AD and shipped on ice to the sampling coordinators. For each replicate, 2 mL sample was preserved in 2 mL RNAlater (Invitrogen), stored at 4°C until all national samples were collected (usually within a few days), and then shipped to Aalborg University with cooling elements. Upon arrival, the samples were separated into aliquots that were prepared for nucleic acid purification. Metadata associated with each AD was also obtained by the sampling coordinators and is provided as Supplementary Data 1. Minimum information from all ADs included continent, country, GPS coordinates, sampling date, temperature in the digester ("Mesophilic" (≤45°C) or "Thermophilic" (50-60°C)), primary substrate ("Wastewater sludge", "Industrial", "Food waste", "Manure", or "Other"), and digester technology ("Two-stage digester (TSAD)", "Continuous Stirred Tank Reactors (CSTR)", "Plug-flow", "Upflow anaerobic sludge blanket (UASB)", or "Other").

General molecular methods

All commercial kits were used according to the protocols provided by the manufacturer unless otherwise stated. The concentration and quality of nucleic acids were determined using a Qubit 3.0 fluorometer (Thermo Fisher Scientific) and an Agilent 2200 Tapestation (Agilent Technologies), respectively.

Nucleic acid purification

DNA was purified using a custom plate-based extraction protocol based on the FastDNA spin kit for soil (MP Biomedicals). The protocol is available at www.midasfieldguide.org (aau_ad_dna_v 2.0). RNAlater preserved samples were thawed and homogenized using a Heidolph RZR 2020 laboratory stirrer. 20 µL of sample was resuspended in 300 µL PBS and transferred to Lysing Matrix E barcoded tubes (MP Biomedicals). 40 µL of MT buffer was added and lysis was performed by bead beating in a FastPrep-96 bead beater (MP Biomedicals) (3x 120 s, 1800 rpm with 2 min incubation on ice between cycles). The samples were centrifuged (3,486 ×g, 10 min) and 200 µL supernatant was transferred to a 96-well PCR-plate. 50 µL Protein Precipitation Solution (PPS) was mixed with each sample, which was then centrifuged again. 150 µL supernatant was cleaned-up using 100 µL CleanNGS beads with elution into 60 µL of nuclease-free water. 40 µL of the purified DNA was transferred to a new 96-well plate and stored at -80°C.

Full-length 16S rRNA gene library preparation, sequencing, and processing

Full-length 16S rRNA gene sequencing was carried out using high-accuracy, long-read amplicon sequencing using UMIs and PacBio CCS [75]. Oligonucleotides used can be found in Supplementary Table 1. Bacterial and archaeal 16S rRNA genes were UMI-tagged using overhang primers based on the 27F and 1391R [76] and SSU1ArF and SSU1000ArR [32] primer pairs, respectively. These primers have shown excellent coverage for the known bacterial and archaeal diversity in silico [32,76].

Addition of UMI-tags by overhang PCR: Adaptors containing UMIs, and defined primer binding sites were added to each end of the bacterial and archaeal 16S rRNA genes by PCR. The reaction contained 20 µL of 5x SuperFi Buffer (Invitrogen), 2 µL of 10 mM dNTP mix, 5 µL of 10 µM f16S_pcr1_fw, 5 µL of 10 µM f16S_pcr1_rv, 1 µL of 2 U/µL Platinum SuperFi DNA polymerase (Invitrogen), 100 ng of pooled template DNA (from all ADs), and nuclease-free water to 100 µL. The reaction was incubated with an initial denaturation at 98°C for 30 s followed by 2 cycles of denaturation at 98°C for 20 s, annealing at 55°C for 30 s, and extension at 72°C for 45 s, and then a final extension at 72°C for 5 min. The sample was purified using 0.6x CleanNGS beads and eluted in 20 µL nuclease-free water.

Primary library amplification: The tagged 16S rRNA gene amplicons were amplified using PCR to obtain enough product for quantification. The reaction contained 19 µL of UMI-tagged sample, 20 µL 5x SuperFi buffer (Invitrogen), 2 µL of 10 mM dNTP, 5 µL of 10 µM f16S_pcr2_fw, 5 µL of 10 µM f16S_pcr2_rv, 48 µL nuclease-free water, and 1 µL 2U/µL Platinum SuperFi DNA polymerase (Invitrogen). The reaction was incubated with an initial denaturation at 98°C for 30 s followed by 15 cycles of denaturation at 98°C for 20 s, annealing at 60°C for 30 s, and extension at 72°C for 45 s and then a final extension at 72°C for 5 min. The PCR product was purified using 0.6x CleanNGS beads and eluted in 11 µL nuclease-free water. The amplicons were validated on a Genomic screentape and quantified with the Qubit dsDNA HS assay kit.

Clonal library amplification: Tagged amplicon libraries were diluted to approximately 250,000 molecules/µL and amplified by PCR to obtain clonal copies of each uniquely tagged amplicon molecule. Three libraries were made for the bacterial 16S rRNA genes and one for archaea. The PCR reactions contained 1 µL diluted primary library, 20 µL 5x SuperFi buffer (Invitrogen), 2 µL of 10 mM dNTP, 5 µL of 10 µM f16S_pcr2_fw, 5 µL of 10 µM f16S_pcr2_rv, 66 µL nuclease-free water, and 1 µL 2U/µL Platinum SuperFi DNA polymerase (Invitrogen). The reaction was incubated with an initial denaturation at 98°C for 30 s followed by 25 cycles of denaturation at 98°C for 20 s, annealing at 60°C for 30 s, and extension at 72°C for 45 s and then a final extension at 72°C for 5 min. The PCR product was purified using 0.6x CleanNGS beads and eluted in 20 µL nuclease-free water. The amplicons were validated on a Genomic screentape and quantified with the Qubit dsDNA HS assay kit.

PacBio CCS sequencing: The four clonal libraries were sent to Admera Health
(Plainfield, NJ, USA) for PacBio library preparation and sequencing. Here amplicons
were incubated with T4 polynucleotide kinase (New England Biolabs) following the
manufacturer's instructions, and sequencing library prepared using SMRTbell Express
Template Preparation kit 1.0 following the standard protocol. Sequencing was
performed using 4x SMRT cells on a Sequel II using a Sequel II Sequencing kit 1.0,
Sequel II Binding and Int Ctrl kit 1.0 and Sequel II SMRT Cell 8M, following the
standard protocol with 1 h pre-extension and 15 h collection time (Pacific Biosciences).

Bioinformatic processing: CCS reads were generated from raw PacBio data using CCS
v.3.4.1 (https://github.com/PacificBiosciences/ccs) with default settings. UMI
consensus sequences (consensus_raconx3.fa) were obtained using the longread_umi
script (https://github.com/SorenKarst/longread_umi) [75] using the following options:
pacbio_pipeline, -v 3, -m 1000, -M 2000, -s 60, -e 60, -f
CAAGCAGAAGACGGCATACGAGAT, -F AGRGTTYGATYMTGGCTCAG
(bacteria) or TCCGGTTGATCCYGCBRG (archaea), -r
AATGATACGGCGACCACCGAGATC, -R GACGGGCGGTGWGTRCA (bacteria)
or GGCCATGCAMYWCCTCTC (archaea), and -c 3. The UMI-consensus reads were
oriented based on the SILVA 138.1 SSURef NR99 database using the usearch
v.11.0.667 -orient command and trimmed between the 27f and 1391r (bacteria) or
SSU1ArF and SSU1000ArR (archaea) primer binding sites using the trimming function
in CLC genomics workbench v. 20.0. Sequences without both primer binding sites were
discarded. The trimmed high-fidelity reads were processed with AutoTax v. 1.7.4 [6] to
create full-length 16S rRNA gene amplicon sequence variants (FL-ASV) and these
were added to the MiDAS 4.8.1 reference database [18] to create MiDAS 5.0.

Short-read amplicon sequencing
V1-V3 amplicons were made using the 27F (5'-AGAGTTTGATCCTGGCTCAG-3')
[77] and 534R (5'-ATTACCGCGGCTGCTGG-3') [78] primers with barcodes and Illumina
adaptors (IDT) [79]. 25 μL PCR reactions in duplicate were run for each sample using 1X
PCRBIO Ultra Mix (PCR Biosystems), 400 nM of both forward and reverse primer,
and 10 ng template DNA. PCR conditions were 95°C, for 2 min followed by 20 cycles
of 95°C for 20 s, 56°C for 30 s, and 72°C for 60 s, followed by a final elongation at
72°C for 5 min. PCR products were purified using 0.8x CleanNGS beads and eluted in
25 μL nuclease-free water.

V3-V5 amplicons were made using the Arch-340F (5'-CCCTAHGGGGYGCASCA-
3') and Arch-915R (5'-GWGCYCCCCGYCAATTC-3') primers [80]. 25 μL PCR
reactions in duplicate were run for each sample using 1X PCRBIO Ultra Mix (PCR
Biosystems), 400 nM of both forward and reverse primer, and 10 ng template DNA.
PCR conditions were 95°C, for 2 min followed by 30 cycles of 95°C for 15 s, 55°C for
15 s, and 72°C for 50 s, followed by a final elongation at 72°C for 5 min. PCR products
were purified using 0.8x CleanNGS beads and eluted in 25 μL nuclease-free water. 2

µL of purified PCR product from above was used as template for a 25 µL Illumina
barcoding PCR reaction containing 1x PCRBIO Reaction buffer, 1 U PCRBIO HiFi
Polymerase (PCR Biosystems) and 10 µL of Nextera adaptor mix (Illumina). PCR
conditions were 95°C, for 2 min, 8 cycles of 95°C for 20 s, 55°C for 30 s, and 72°C for
60 s, followed by a final elongation at 72°C for 5 min. PCR products were purified
using 0.8x CleanNGS beads and eluted in 25 µL nuclease-free water.
739
V4 amplicons were made using the 515F (5'-GTGYCAGCMGCCGCGGTAA-3') [78]
and 806R (5'-GGACTACNVGGGTWTCTAAT-3') [81] primers. 25 µL PCR reactions
in duplicate were run for each sample using 1X PCRBIO Ultra Mix (PCR Biosystems),
400 nM of both forward and reverse primer, and 10 ng template DNA. PCR conditions
were 95°C, for 2 min followed by 30 cycles of 95°C for 15 s, 55°C for 15 s, and 72°C
for 50 s, followed by a final elongation at 72°C for 5 min. PCR products were purified
using 0.8x CleanNGS beads and eluted in 25 µL nuclease-free water. 2 µL of purified
PCR product from above was used as template for a 25 µL Illumina barcoding PCR
reaction as described for the V3-V5 amplicons.
749
750 16S rRNA gene V1-V3, V3-V5, and V4 amplicon libraries were pooled separately in
751 equimolar concentrations and diluted to 4 nM. The amplicon libraries were paired-end
752 sequenced (2 × 300 bp) on the Illumina MiSeq using v3 chemistry (Illumina, USA). 10
753 to 20% PhiX control library was added to mitigate low diversity library effects.
754
Processing of short-read amplicon data
Usearch v.11.0.667 [82] was used for processing of 16S rRNA gene amplicon data and
for read mapping. V1-V3 forward and reverse reads were merged using the usearch -
fastq_mergepairs command, filtered to remove phiX sequences using usearch -
filter_phix, and quality filtered using usearch -fastq_filter with -fastq_maxee 1.0.
Dereplication was performed using -fastx_uniques with -sizeout, and amplicon
sequence variants (ASVs) were resolved using the usearch -unoise3 command. An
ASV-table was created by mapping the quality filtered reads to the ASVs using the
usearch -otutab command with the -zotus and -strand plus options. Taxonomy was
assigned to ASVs using the usearch -sintax command with -strand both and -
sintax_cutoff 0.8 options. Mapping of ASVs to reference databases was done with the
usearch -usearch_global command and the -id 0, -maxaccepts 0, -maxrejects 0, -
top_hit_only, and -strand plus options.
768
16S rRNA gene V3-V5 forward reads (reverse reads in relation the 16S rRNA gene)
were filtered to remove phiX sequences using usearch -filter_phix, trimmed to remove
primers and obtain a fixed length of 250 bp using -fastx_truncate with -stripleft -17 and
trunclen 250, reverse complemented with usearch -fastx_revcomp, and quality filtered
using usearch -fastq_filter with -fastq_maxee 1.0. Subsequent processing was like that
for the V1-V3 amplicons.

16S rRNA gene V4 forward reads (reverse reads in relation the 16S rRNA gene) were
trimmed with cutadapt v.2.8 [83] based on the V4 primers with the -g
^GGACTACHVGGGTWTCTAAT...TTACCGCGGCKGCTGGCAC and --discard-
untrimmed options. The trimmed reads, which span the entire V4 amplicon, were
reverse complemented with usearch -fastx_revcomp, and quality filtered using usearch
-fastq_filter with -fastq_maxee 1.0. Subsequent processing was like that for the V1-V3
amplicons.
782

In silico *primer evaluation*
The specificity of commonly used amplicon primers was determined for each FL-ASV
using the analyze_primers.py script from Primer Prospector v. 1.0.1 [84]. The specificity
of primer sets was defined based on the overall weighted scores (OWS) for the primer
with the highest score as follows: Perfect hit (OWS = 0), partial hit (OWS > 0, and ≤
1), poor hit (OWS > 1). The percentage of perfect hits were calculated in R for all taxa
in MiDAS 5.
790

791   *Microbial community analyses*
792   Short-read amplicon data was analyzed with R v.4.2.0 [85] through RStudio IDE [86], with
793   the tidyverse v.1.3.1 (https://www.tidyverse.org/), vegan v.2.6-4 [87], maps v.3.4.0 [88],
794   data.table v.1.14.4 [89], FSA v.0.9.3 [90], rcompanion v. 2.4.18 [91], patchwork v.1.1.1 [92],
795   ggupset v.0.3.0 [93] and Ampvis2 v.2.7.34 [94] packages.
796

The microbial community analyses were performed based on all three 16S rRNA gene
short-read amplicon dataset (V1-V3, V3-V5, and V4). Samples with less than 10,000
reads and those lacking information about digester technology, primary substrate, and
temperature in the digester were discarded from the analyses. After filtration, 365 V1-
V3, 200 V3-V5, and 269 V4 samples remained.

Associations between the AD microbiota and the following process-related or
environmental variables were investigated: Digester technology, primary substrate,
temperature in the digester, and continent (see definitions above). All variables were
treated as factors.

For alpha diversity analyses, samples were rarefied to 10,000 reads, and alpha diversity
(observed ASVs and inverse Simpson) was calculated using the ampvis2 package. The
Kruskal-Wallis with Dunn's post-hoc test (Bonferroni correction with α=0.01 before
correction) was used to determine statistically significant differences in alpha diversity
between samples grouped by process and environmental variables.

Beta diversity distances based on Bray-Curtis (abundance-based) for genera was
calculated using the vegdist function in the vegan R package and visualized by PCoA
plots with the ampvis2 package. To determine how much individual parameters
affected the structure of the microbial community across the ADs, a permutational

multivariate analysis of variance (PERMANOVA) test was performed on the beta-
diversity matrices using the adonis function in the vegan package with 999
permutations.

Core taxa (genera and species) were determined separately for ADs treating different
primary substrates and operating at different temperatures (mesophilic and
thermophilic) based on their relative abundances in individual ADs according to the
three short-read amplicon datasets. Core taxa definitions were identical to those applied
in the MiDAS global survey of WWTPs [18]. Taxa were considered abundant when
present at >0.1% relative read abundance in individual ADs. Based on how frequently
taxa were observed to be abundant, we defined the following core communities: loose
core (>20% of ADs), general core (>50% of ADs), and strict core (>80% of ADs).
Additionally, we defined conditionally rare or abundant taxa (CRAT) [42] composed of
taxa present in one or more ADs at >1% relative abundance, but not belonging to the
core taxa.
829

Data Availability:
The raw and assembled sequencing data generated in this study have been deposited in
the NCBI SRA database and will be made available when the final article is accepted
for peer-reviewed publication. The MiDAS 5 reference database in SINTAX, QIIME
and Dada2 format is available at the MiDAS fieldguide website
[https://www.midasfieldguide.org/guide/downloads].
836

837 **Code Availability:**
838 R scripts and raw data used for data analyses and figures will be made available at
839 GitHub and Figshare when the final article is accepted for peer-reviewed publication.

References:

1. Tiwary, A., Williams, I. D., Pant, D. C. & Kishore, V. V. N. Emerging perspectives on environmental burden minimisation initiatives from anaerobic digestion technologies for community scale biomass valorisation. *Renewable and Sustainable Energy Reviews* **42**, 883–901 (2015).

2. Achinas, S., Achinas, V. & Euverink, G. J. W. A technological overview of biogas production from biowaste. *Engineering* **3**, 299–307 (2017).

3. Samoraj, M. *et al.* The challenges and perspectives for anaerobic digestion of animal waste and fertilizer application of the digestate. *Chemosphere* **295**, 133799 (2022).

4. Czekała, W., Jasiński, T., Grzelak, M., Witaszek, K. & Dach, J. Biogas plant operation: Digestate as the valuable product. *Energies* **15**, 8275 (2022).

5. Briones, A. & Raskin, L. Diversity and dynamics of microbial communities in engineered environments and their implications for process stability. *Current Opinion in Biotechnology* **14**, 270–276 (2003).

6. Dueholm, M. S. *et al.* Generation of comprehensive ecosystem-specific reference databases with species-level resolution by high-throughput full-length 16S rRNA gene sequencing and automated taxonomy assignment (AutoTax). *mBio* **11**, e01557-20 (2020).

7. Jiang, C. *et al.* Characterizing the growing microorganisms at species level in 46 anaerobic digesters at Danish wastewater treatment plants: A six-year survey on microbial community structure and key drivers. *Water Research* **193**, 116871 (2021).

8. Vanwonterghem, I. *et al.* Deterministic processes guide long-term synchronised population dynamics in replicate anaerobic digesters. *ISME Journal* **8**, 2015–2028 (2014).

9. Ofiţeru, I. D. *et al.* Combined niche and neutral effects in a microbial wastewater treatment community. *Proceedings of the National Academy of Sciences* **107**, 15345–15350 (2010).

10. Kirkegaard, R. H. *et al.* The impact of immigration on microbial community composition in full-scale anaerobic digesters. *Scientific Reports* **7**, 9343 (2017).

11. Werner, J. J. *et al.* Bacterial community structures are unique and resilient in full-scale bioenergy systems. *Proceedings of the National Academy of Sciences of the United States of America* **108**, 4158–4163 (2011).

12. Campanaro, S. *et al.* Metagenomic analysis and functional characterization of the biogas microbiome using high throughput shotgun sequencing and a novel binning strategy. *Biotechnology for biofuels* **9**, 26 (2016).

13. Calusinska, M. *et al.* A year of monitoring 20 mesophilic full-scale bioreactors reveals the existence of stable but different core microbiomes in bio-waste and wastewater anaerobic digestion systems. *Biotechnology for Biofuels* **11**, 196 (2018).

14. Mei, R. *et al.* Operation-driven heterogeneity and overlooked feed-associated
populations in global anaerobic digester microbiome. *Water Research* **124**, 77–84
(2017).
15. McIlroy, S. J. *et al.* MiDAS 2.0: An ecosystem-specific taxonomy and online
database for the organisms of wastewater treatment systems expanded for anaerobic
digester groups. *Database* **2017**, bax016 (2017).
16. Nierychlo, M. *et al.* MiDAS 3: An ecosystem-specific reference database,
taxonomy and knowledge platform for activated sludge and anaerobic digesters
reveals species-level microbiome composition of activated sludge. *Water Research*
182, 115955 (2020).
17. McIlroy, S. J. *et al.* MiDAS: The field guide to the microbes of activated sludge.
Database : the journal of biological databases and curation **2015**, bav062 (2015).
18. Dueholm, M. K. D. *et al.* MiDAS 4: A global catalogue of full-length 16S rRNA
gene sequences and taxonomy for studies of bacterial communities in wastewater
treatment plants. *Nat Commun* **13**, 1908 (2022).
19. Albertsen, M., Karst, S. M., Ziegler, A. S., Kirkegaard, R. H. & Nielsen, P. H. Back
to basics - the influence of DNA extraction and primer choice on phylogenetic
analysis of activated sludge communities. *PloS one* **10**, e0132783 (2015).
20. Kristensen, J. M., Singleton, C., Clegg, L.-A., Petriglieri, F. & Nielsen, P. H. High
diversity and functional potential of undescribed "Acidobacteriota" in Danish
wastewater treatment plants. *Front. Microbiol.* **12**, 906 (2021).
21. Nierychlo, M. *et al.* Low global diversity of *Candidatus* Microthrix, a troublesome
filamentous organism in full-scale WWTPs. *Front Microbiol* **12**, 690251 (2021).
22. Petriglieri, F. *et al.* "*Candidatus* Dechloromonas phosphoritropha" and "*Ca.* D.
phosphorivorans", novel polyphosphate accumulating organisms abundant in
wastewater treatment systems. *ISME J* **15**, 3605–3614 (2021).
23. Petriglieri, F. *et al.* Reevaluation of the phylogenetic diversity and global
distribution of the genus "*Candidatus* Accumulibacter". *mSystems* **7**, e00016-22
(2022).
24. Singleton, C. M. *et al.* The novel genus, '*Candidatus* Phosphoribacter', previously
identified as *Tetrasphaera*, is the dominant polyphosphate accumulating lineage in
EBPR wastewater treatment plants worldwide. *ISME J* **16**, 1605–1616 (2022).
25. Rivière, D. *et al.* Towards the definition of a core of microorganisms involved in
anaerobic digestion of sludge. *ISME Journal* **3**, 700–714 (2009).
26. Yarza, P. *et al.* Uniting the classification of cultured and uncultured bacteria and
archaea using 16S rRNA gene sequences. *Nature Reviews Microbiology* **12**, 635–
645 (2014).
27. Fujimoto, M., Carey, D. E., Zitomer, D. H. & McNamara, P. J. Syntroph diversity
and abundance in anaerobic digestion revealed through a comparative core
microbiome approach. *Appl Microbiol Biotechnol* **103**, 6353–6367 (2019).

28. Westerholm, M., Calusinska, M. & Dolfing, J. Syntrophic propionate-oxidizing
bacteria in methanogenic systems. *FEMS Microbiology Reviews* **46**, fuab057
(2022).
29. Johnson, J. S. *et al.* Evaluation of 16S rRNA gene sequencing for species and strain-
level microbiome analysis. *Nature Communications* **10**, 1–11 (2019).
30. Yuan, C., Lei, J., Cole, J. & Sun, Y. Reconstructing 16S rRNA genes in
metagenomic data. *Bioinformatics* **31**, i35–i43 (2015).
31. Singleton, C. M. *et al.* Connecting structure to function with the recovery of over
1000 high-quality metagenome-assembled genomes from activated sludge using
long-read sequencing. *Nature Communications* **12**, 2009 (2021).
32. Bahram, M., Anslan, S., Hildebrand, F., Bork, P. & Tedersoo, L. Newly designed
16S rRNA metabarcoding primers amplify diverse and novel archaeal taxa from
the environment. *Environmental Microbiology Reports* **11**, 487–494 (2018).
33. Pausan, M. R. *et al.* Exploring the archaeome: Detection of archaeal signatures in
the human body. *Frontiers in Microbiology* **10**, 2796 (2019).
34. Wu, L. *et al.* Global diversity and biogeography of bacterial communities in
wastewater treatment plants. *Nature Microbiology* **4**, 1183–1195 (2019).
35. Thompson, L. R. *et al.* A communal catalogue reveals Earth's multiscale microbial
diversity. *Nature* **551**, 457–463 (2017).
36. Mei, R., Narihiro, T., Nobu, M. K., Kuroda, K. & Liu, W.-T. Evaluating digestion
efficiency in full-scale anaerobic digesters by identifying active microbial
populations through the lens of microbial activity. *Scientific Reports* **6**, 34090
(2016).
37. Sun, L., Pope, P. B., Eijsink, V. G. H. & Schnürer, A. Characterization of microbial
community structure during continuous anaerobic digestion of straw and cow
manure. *Microbial Biotechnology* **8**, 815–827 (2015).
38. Karakashev, D., Batstone, D. J. & Angelidaki, I. Influence of environmental
conditions on methanogenic compositions in anaerobic biogas reactors. *Applied
and Environmental Microbiology* **71**, 331–338 (2005).
39. Levén, L., Eriksson, A. R. B. & Schnürer, A. Effect of process temperature on
bacterial and archaeal communities in two methanogenic bioreactors treating
organic household waste. *FEMS Microbiology Ecology* **59**, 683–693 (2007).
40. Knights, D. *et al.* Rethinking "enterotypes". *Cell Host & Microbe* **16**, 433–437
(2014).
41. Saunders, A. M., Albertsen, M., Vollertsen, J. & Nielsen, P. H. The activated sludge
ecosystem contains a core community of abundant organisms. *The ISME journal*
10, 11–20 (2016).
42. Dai, T. *et al.* Identifying the key taxonomic categories that characterize microbial
community diversity using full-scale classification: a case study of microbial
communities in the sediments of Hangzhou Bay. *FEMS Microbiology Ecology* **92**,
fiw150 (2016).

43. Astudillo-García, C. *et al.* Evaluating the core microbiota in complex communities: A systematic investigation. *Environmental Microbiology* **19**, 1450–1462 (2017).

44. Shade, A. *et al.* Conditionally rare taxa disproportionately contribute to temporal changes in microbial diversity. *mBio* **5**, e01371-14 (2014).

45. Schorn, S. *et al.* Diverse methylotrophic methanogenic archaea cause high methane emissions from seagrass meadows. *Proceedings of the National Academy of Sciences* **119**, e2106628119 (2022).

46. Evans, P. N. *et al.* Methane metabolism in the archaeal phylum Bathyarchaeota revealed by genome-centric metagenomics. *Science* **350**, 434–438 (2015).

47. Nobu, M. K., Narihiro, T., Kuroda, K., Mei, R. & Liu, W. T. Chasing the elusive Euryarchaeota class WSA2: genomes reveal a uniquely fastidious methyl-reducing methanogen. *The ISME Journal* **10**, 2478–2487 (2016).

48. Jung, M.-Y. *et al.* A hydrophobic ammonia-oxidizing archaeon of the *Nitrosocosmicus* clade isolated from coal tar-contaminated sediment. *Environmental Microbiology Reports* **8**, 983–992 (2016).

49. Huang, W.-C. *et al.* Comparative genomic analysis reveals metabolic flexibility of Woesearchaeota. *Nat Commun* **12**, 5281 (2021).

50. Morris, B. E. L., Henneberger, R., Huber, H. & Moissl-Eichinger, C. Microbial syntrophy: Interaction for the common good. *FEMS Microbiology Reviews* **37**, 384–406 (2013).

51. Hao, L. *et al.* Novel syntrophic bacteria in full-scale anaerobic digesters revealed by genome-centric metatranscriptomics. *ISME J* **14**, 906–918 (2020).

52. Sieber, J. R., McInerney, M. J. & Gunsalus, R. P. Genomic insights into syntrophy: the paradigm for anaerobic metabolic cooperation. *Annual Review of Microbiology* **66**, 429–452 (2012).

53. Westerholm, M., Roos, S. & Schnürer, A. *Syntrophaceticus schinkiigen* nov., sp. nov., an anaerobic, syntrophic acetate-oxidizing bacterium isolated from a mesophilic anaerobic filter. *FEMS Microbiology Letters* **309**, 100–104 (2010).

54. Schnürer, A., Müller, B. & Westerholm, M. *Syntrophaceticus*. in *Bergey's Manual of Systematics of Archaea and Bacteria* 1–10 (John Wiley & Sons, Ltd, 2018). doi:10.1002/9781118960608.gbm01452.

55. McInerney, M. J., Bryant, M. P., Hespell, R. B. & Costerton, J. W. *Syntrophomonas wolfei* gen. nov. sp. nov., an anaerobic, syntrophic, fatty acid-oxidizing bacterium. *Applied and Environmental Microbiology* **41**, 1029–1039 (1981).

56. Wu, C., Liu, X. & Dong, X. *Syntrophomonas cellicola* sp. nov., a spore-forming syntrophic bacterium isolated from a distilled-spirit-fermenting cellar, and assignment of *Syntrophospora bryantii* to *Syntrophomonas bryantii* comb. nov. *Int J Syst Evol Microbiol* **56**, 2331–2335 (2006).

57. Sousa, D. Z., Smidt, H., Alves, M. M. & Stams, A. J. M. *Syntrophomonas zehnderi* sp. nov., an anaerobe that degrades long-chain fatty acids in co-culture with *Methanobacterium formicicum*. *Int J Syst Evol Microbiol* **57**, 609–615 (2007).

58. Hatamoto, M., Imachi, H., Fukayo, S., Ohashi, A. & Harada, H. *Syntrophomonas*
palmitatica sp. nov., an anaerobic, syntrophic, long-chain fatty-acid-oxidizing
bacterium isolated from methanogenic sludge. *International Journal of Systematic*
and Evolutionary Microbiology **57**, 2137–2142 (2007).

59. Niu, L., Song, L., Liu, X. & Dong, X. *Tepidimicrobium xylanilyticum* sp. nov., an
anaerobic xylanolytic bacterium, and emended description of the genus
Tepidimicrobium. International Journal of Systematic and Evolutionary
Microbiology **59**, 2698–2701 (2009).

60. Wang, G., Li, Q., Gao, X. & Wang, X. C. Sawdust-derived biochar much mitigates
VFAs accumulation and improves microbial activities to enhance methane
production in thermophilic anaerobic digestion. *ACS Sustainable Chem. Eng.* **7**,
2141–2150 (2019).

61. Liu, Y., Balkwill, D. L., Aldrich, H. C., Drake, G. R. & Boone, D. R.
Characterization of the anaerobic propionate-degrading syntrophs *Smithella*
propionica gen. nov., sp. nov. and *Syntrophobacter wolinii. International Journal*
of Systematic and Evolutionary Microbiology **49**, 545–556 (1999).

62. de Bok, F. A. M., Stams, A. J. M., Dijkema, C. & Boone, D. R. Pathway of
propionate oxidation by a syntrophic culture of *Smithella propionica* and
Methanospirillum hungatei. Applied and Environmental Microbiology **67**, 1800–
1804 (2001).

63. Dolfing, J. Syntrophic propionate oxidation via butyrate: a novel window of
opportunity under methanogenic conditions. *Applied and Environmental*
Microbiology **79**, 4515–4516 (2013).

64. Embree, M., Nagarajan, H., Movahedi, N., Chitsaz, H. & Zengler, K. Single-cell
genome and metatranscriptome sequencing reveal metabolic interactions of an
alkane-degrading methanogenic community. *ISME J* **8**, 757–767 (2014).

65. Tan, B., Nesbø, C. & Foght, J. Re-analysis of omics data indicates *Smithella* may
degrade alkanes by addition to fumarate under methanogenic conditions. *ISME*
Journal **8**, 2353–2356 (2014).

66. Nobu, M. K. *et al.* The genome of *Syntrophorhabdus aromaticivorans* strain UI
provides new insights for syntrophic aromatic compound metabolism and electron
flow. *Environmental Microbiology* **17**, 4861–4872 (2015).

67. Qiu, Y.-L. *et al. Syntrophorhabdus aromaticivorans* gen. nov., sp. nov., the first
cultured anaerobe capable of degrading phenol to acetate in obligate syntrophic
associations with a hydrogenotrophic methanogen. *Appl Environ Microbiol* **74**,
2051–2058 (2008).

68. McInerney, M. J. *et al.* The genome of *Syntrophus aciditrophicus*: Life at the
thermodynamic limit of microbial growth. *Proceedings of the National Academy of*
Sciences **104**, 7600–7605 (2007).

69. Jackson, B. E., Bhupathiraju, V. K., Tanner, R. S., Woese, C. R. & McInerney, M.
J. *Syntrophus aciditrophicus* sp. nov., a new anaerobic bacterium that degrades fatty

acids and benzoate in syntrophic association with hydrogen-using microorganisms. *Arch Microbiol* **171**, 107–114 (1999).

70. Ganidi, N., Tyrrel, S. & Cartmell, E. Anaerobic digestion foaming causes – A review. *Bioresource Technology* **100**, 5546–5554 (2009).

71. Duan, J.-L. *et al.* Unraveling anaerobic digestion foaming via association between bacterial metabolism and variations in microbiota. *ACS EST Eng.* **1**, 978–988 (2021).

72. Jiang, C. *et al.* Identification of microorganisms responsible for foam formation in mesophilic anaerobic digesters treating surplus activated sludge. *Water Research* **191**, 116779 (2021).

73. Bovio-Winkler, P., Cabezas, A. & Etchebehere, C. Database mining to unravel the ecology of the phylum Chloroflexi in methanogenic full scale bioreactors. *Frontiers in Microbiology* **11**, 603234 (2021).

74. McIlroy, S. J. *et al.* Culture-independent analyses reveal novel Anaerolineaceae as abundant primary fermenters in anaerobic digesters treating waste activated sludge. *Frontiers in Microbiology* **8**, 1134 (2017).

75. Karst, S. M. *et al.* High-accuracy long-read amplicon sequences using unique molecular identifiers with Nanopore or PacBio sequencing. *Nat Methods* **18**, 165–169 (2021).

76. Klindworth, A. *et al.* Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies. *Nucleic Acids Research* **41**, e1 (2013).

77. Lane, D. J. 16S/23S rRNA sequencing. in *Nucleic Acid Techniques in Bacterial Systematics* (eds. Stackebrandt, E. & Goodfellow, M.) 115–175 (John Wiley and Sons, 1991). doi:10.1007/s00227-012-2133-0.

78. Parada, A. E., Needham, D. M. & Fuhrman, J. A. Every base matters: Assessing small subunit rRNA primers for marine microbiomes with mock communities, time series and global field samples. *Environmental Microbiology* **18**, 1403–1414 (2016).

79. Caporaso, J. G. *et al.* Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. *The ISME Journal* **6**, 1621–1624 (2012).

80. Pinto, A. J. & Raskin, L. PCR biases distort bacterial and archaeal community structure in pyrosequencing datasets. *PLoS One* **7**, e43093 (2012).

81. Apprill, A., Mcnally, S., Parsons, R. & Weber, L. Minor revision to V4 region SSU rRNA 806R gene primer greatly increases detection of SAR11 bacterioplankton. *Aquatic Microbial Ecology* **75**, 129–137 (2015).

82. Edgar, R. C. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* **26**, 2460–2461 (2010).

83. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal* **17**, 10–12 (2011).

84. Walters, W. A. *et al.* PrimerProspector: De novo design and taxonomic analysis of
barcoded polymerase chain reaction primers. *Bioinformatics* **27**, 1159–1161
(2011).
85. R Development Core Team, R. F. F. S. C. & R Core Team. *R: A Language and*
Environment for Statistical Computing. Vienna Austria R Foundation for Statistical
Computing vol. 1 (R Foundation for Statistical Computing, Vienna, 2008).
86. RStudio Team. *RStudio: Integrated Development Environment for R*. (RStudio,
PBC., 2020).
87. Oksanen, J. *et al. vegan: Community Ecology Package*. (2019).
88. Richard A. Becker, Allan R. Wilks, Ray Brownrigg, Thomas P Minka, & Alex
Deckmyn. maps: Draw Geographical Maps. (2021).
89. Dowle, M. & Srinivasan, A. data.table: Extension of `data.frame`. (2019).
90. Ogle, D. H., Doll, J. C., Wheeler, A. P. & Dinno, A. FSA: Simple fisheries stock
assessment methods. (2023).
91. Mangiafico, S. rcompanion: Functions to support extension education program
evaluation. (2023).
92. Pedersen, T. L. Patchwork: the composer of plots. (2020).
93. Constantin Ahlmann-Eltze. ggupset: Combination matrix axis for 'ggplot2' to
create 'UpSet' plots. (2020).
94. Andersen, K. S. S., Kirkegaard, R. H., Karst, S. M. & Albertsen, M. Ampvis2: an
R package to analyse and visualise 16S rRNA amplicon data. *bioRxiv* 299537
(2018) doi:10.1101/299537.
1106

Acknowledgements:

The project has been funded by the Danish Research Council (grant 6111-00617A, P.H.N.) and the Villum Foundation (Dark Matter and grant 13351, P.H.N.). We thank all the involved anaerobic digester plants for providing samples and plant metadata.

Author Contributions:

P.H.N. and M.K.D.D. designed the study. M.K.D.D. and P.H.N. wrote the manuscript and all authors reviewed and approved the final manuscript. The MiDAS Global Consortium for Anaerobic Digesters provided samples and metadata. V.R. handled sampling, DNA extraction and library preparation for DNA sequencing. K.S.A. and M.K.D.D. performed the bioinformatics analyses. M.K.D.D., A-K.C.P. and K.S.A. curated metadata and carried out statistical analyses.

MiDAS Global Consortium for Anaerobic Digesters:

Madalena Alves[2], Yadira Bajón-Fernández[3], Damien Batstone[4], Caitlyn Butler[5], Mercedes Cecilia Cruz[6], Åsa Davidsson[7], Leonardo Erijman[8], Christof Holliger[9], Konrad Koch[10], Norbert Kreuzinger[11], Changsoo Lee[12], Gerasimos Lyberatos[13], Srikanth Mutnuri[14], Vincent O'Flaherty[15], Piotr Oleskowicz-Popiel[16], Dana Pokorna[17], Veronica Rajal[6], Michael Recktenwald[18], Jorge Rodríguez[19], Pascal E. Saikaly[20], Nick Tooker[5], Julia Vierheilig[11], Jo De Vrieze[21], Christian Wurzbacher[10a10] & Per Halkjær Nielsen[1]

[2]Centre of Biological Engineering, University of Minho, Portugal. [3]School of Water, Energy and Environment, Cranfield University. Cranfield, United Kingdom. [4]Australian Centre for Water and Environmental Biotechnology (ACWEB), The University of Queensland, Australia. [5]Department of Civil and Environmental Engineering, University of Massachusetts Amherst, MA, USA. [6]National University of Salta, Salta, Argentina. [7]Department of Chemical Engineering, Lund University, Sweden. [8]INGEBI-CONICET, University of Buenos Aires, Argentina. [9]Laboratory for Environmental Biotechnology, Ecole Polytechnique Fédérale de Lausanne (EPFL), Switzerland. [10]Chair of Urban Water Systems Engineering, Technical University of Munich (TUM), Garching, Germany. [11]Institute of Water Quality and Resource Management, Vienna University of Technology, Austria. [12]Department of Urban and Environmental Engineering & Graduate School of Carbon Neutrality, Ulsan National Institute of Science and Technology (UNIST), South Korea. [13]School of Chemical Engineering, National Technical University of Athens, Greece. [14]Applied Environmental Biotechnology Laboratory, Birla Institute of Technology and Science (BITS-Pilani), India. [15]School of Biological and Chemical Sciences and Ryan Institute, University of Galway, Ireland. [16]Water Supply and Bioeconomy Division, Faculty of Environmental Engineering and Energy, Poznan University of Technology, Poland. [17]Department of Water Technology and Environmental Engineering, University of Chemistry and Technology Prague, Czech Republic. [18]Research Scientist at Kemira

Oyj, Espoo R&D Center, Finland. [19]Chemical Engineering Department, Khalifa University, United Arab Emirates. [20]Environmental Science and Engineering Program, Biological and Environmental Science and Engineering Division, King Abdullah University of Science and Technology (KAUST), Saudi Arabia. [21]Center for Microbial Ecology and Technology (CMET), Ghent University, Belgium.

Competing Interests:

The authors declare no competing interests.