Genome replication in asynchronously growing microbial populations

Florian Pflug[1], Deepak Bhat[1,2], Simone Pigolotti[1*]
[1]*Biological Complexity Unit,*
Okinawa Institute of Science and Technology Graduate University,
Onna, Okinawa 904-0495, Japan and
[2]*Department of Physics, School of Advanced Sciences,*
Vellore Institute of Technology, Vellore, Tamil Nadu, India

Biological cells adopt specific programs to replicate their genomes. Information about the replication program of an organism can be obtained by sequencing an exponentially growing cell culture and studying the frequency of DNA fragments as a function of genomic position. However, a quantitative interpretation of this data has been challenging for asynchronously growing cultures. In this paper, we introduce a general theory to predict the abundance of DNA fragments in asynchronously growing cultures from any given stochastic model of the DNA replication program. As key examples, we present stochastic models of DNA replication in *Escherichia coli* and in budding yeast. In both cases, our approach leads to analytical predictions that are in excellent agreement with experimental data and permit to infer biophysically relevant parameters. In particular, our method is able to infer the locations of known replication origins in budding yeast with high accuracy. These examples demonstrate that our method can provide insight into a broad range of organisms, from bacteria to eukaryotes.

I. INTRODUCTION

The genome of an organism contains precious information about its functioning. Genomes need to be reliably and quickly replicated for cells to pass biological information to the next generation. Replication of a genome proceeds according to a certain plan, termed the "replication program" [1–4]. For example, most bacteria have a circular genome, where two replisomes initiate replication by binding at the same origin site [5]. They replicate the genome in opposite directions. Replication is completed when they meet, after each of them has copied approximately half of the genome. The replication program is carefully orchestrated, but not completely deterministic. Stochasticity is particularly relevant in eukaryotes and archaea, where many origin sites are present in each chromosome [6, 7]. Replication can initiate at these origin sites at different times. These times are characterized by some degree of randomness [8, 9] and, depending on the circumstances, some origin sites may not be activated at all [1].

Replication programs must be coordinated with the cell cycle in some way [10]. This idea has led to experimental efforts to determine the replication program. Many of these experiments focus on cultures in which all cells are approximately at the same stage of the cell cycle. This can be achieved by either arresting growth of a synchronized culture at a certain stage of the cell cycle, or by sorting cells depending on their stage in the cell cycle [4]. The fraction of these synchronized cells that have copied each genome location is then measured using deep sequencing. This approach has been extensively used to study the eukaryotic replication program [4, 11, 12].

An alternative method is to measure the abundance of DNA fragments in asynchronous, exponentially growing populations. This approach, originally called marker frequency analysis [13], has been applied to bacterial replication, for example to study *Escherichia coli* mutant lacking genes that assist DNA replication [14–16] and artificially engineered *E.coli* strains with multiple replication origins [17]. The asynchronous approach is experimentally much simpler and avoids potential artifacts caused by the cell cycle arrest [4]. Progress in DNA sequencing have made these experiments high-throughput and relatively inexpensive. However, asynchronous measurements sample cells at different stages in their life cycle, rendering the theoretical interpretation more problematic than for synchronous measurements [18].

Theoretical approaches have attempted to describe measurements in asynchronous populations. However, these approaches either neglect stochasticity, or are limited to specific model systems. For example, we recently proposed a stochastic model describing DNA replication in growing *E.coli* populations [19]. A broader range of bacterial systems have been studied assuming a deterministic replication program [20, 21]. Finally, a model was used to infer replication origins in yeast from an asynchronous growing population, assuming that cells in the population originate from random, uniformly distributed stages in their cell cycle [3].

In this paper, we develop a general theory to infer the replication program from sequencing of an asynchronously growing population of cells. Our theory builds upon classic results on age-structured populations [13, 22–25], that we extend to populations of stochastically replicating genomes. Our approach requires minimal assumptions on the replication dynamics. In particular, it allows for a stochastic replication program and equally applies to bacteria and eukaryotes. We apply our method to experimental data from *E. coli* and budding

* simone.pigolotti@oist.jp

yeast. In the case of *E. coli*, we propose and exactly solve a model of DNA replication that incorporates stochastic stalling. The fit of the solution to experimental data sheds light on recently observed oscillations of bacterial replisome speed. In the case of yeast, our approach permits to reliably infer the location of replication origins.

The paper is organized as follows. In Section II, we present our general theory. In Section III, we apply our framework to DNA replication in *E. coli*. In Section IV, we consider eukaryotic DNA replication, with budding yeast as an example. In Section V, we present possible extensions of our theory. Section VI is devoted to conclusions and future perspectives.

II. DNA REPLICATION IN ASYNCHRONOUSLY GROWING POPULATIONS

A. General theory

We consider a large, growing population of cells. We call $N_c(t)$ the number of cells that are present in the population at time t. Each cell may contain multiple genomes, some complete and other undergoing synthesis (incomplete). We denote by $N_g(t)$ the the total number of complete and incomplete genomes in the population.

Our theory is based on the following assumptions. The number of cells grows exponentially: $N_c(t) \propto \exp(\Lambda t)$, where Λ is the exponential growth rate. The population grows in a steady, asynchronous manner. This assumption implies that the average number of genomes per cell must remain constant, and therefore $N_g(t) \propto \exp(\Lambda t)$. Genomes in the population are immortal, i.e. we neglect the rate at which they might be degraded. All genomes in the population are statistically identical, in the sense that they are all characterized by the same stochastic replication program. These assumptions are realistic in common experimental situations. Moreover, as we shall discuss in Section V, some of them can be readily relaxed, if necessary.

We now assign an age to each genome in the population. To this aim, we conventionally set the birth time a genome as the time at which the replication process that generates it from a parent genome is initiated, see Fig. 1a. We note that the distinction between a "parent" and a "daughter" genome is somewhat arbitrary, as each of them is made up of a preexisting strand and a newly copied complementary one. Since genomes are immortal, the probability density of new genomes is proportional to $\dot{N}_g(t)$, which is proportional to $\exp(\Lambda t)$ as well. It follows that the distribution of ages τ of genomes in the population must be proportional to $\dot{N}_g(t - \tau)$. From this fact, we conclude that the distribution of genome ages in the population is

$$P(\tau) = \Lambda e^{-\Lambda \tau}. \tag{1}$$

We now look into the DNA replication program in more details. In bacteria, replication is carried out by a pair of replisomes, that bind at a specific genome site (the origin) and proceed in opposite directions, each replicating both strands. DNA replication is substantially more complex in eukaryotes, where a large number of replisomes replicate the same chromosome, and initiation site might be activated stochastically. We summarize the effect of all these complex processes into the probability $f(x, \tau)$ that the genome location x has already been copied in a genome of age τ. By definition, $f(x, \tau)$ is a non-decreasing function of τ. Our assumption that genomes are statistically identical means that all genomes are characterized by the same $f(x, \tau)$. Examples of deterministic and stochastic replication programs are represented in Fig. 1b and 1c, respectively.

We now define the probability $\mathcal{P}(x)$ that a randomly chosen genome in the population contains the genome location x. We remark that $\mathcal{P}(x)$ is not normalized to one when integrated over the entire genome, in general. Its normalized counterpart represents the probability density that a randomly chosen genome fragment in the population originates from genomic location x. For this reason, we call $\mathcal{P}(x)$ the DNA abundance distribution. The DNA abundance distribution can be experimentally measured using deep sequencing.

By using Eq. (1) and integrating over all genome ages, we find that $\mathcal{P}(x)$ is related with $f(x, \tau)$ by

$$\mathcal{P}(x) = \int_0^\infty d\tau \, \Lambda e^{-\Lambda \tau} f(x, \tau), \tag{2}$$

see Fig. 1d. To find a more transparent expression, we introduce the probability density $\psi(x, \tau) = \partial_\tau f(x, \tau)$. Substituting this definition into Eq. (2) and integrating by parts we obtain

$$\mathcal{P}(x) = \left\langle e^{-\Lambda \tau} \right\rangle_x, \tag{3}$$

where $\langle \dots \rangle_x = \int_0^\infty d\tau \dots \psi(x, \tau)$ is the average over the distribution of replication times. Equation (2), or equivalently Eq. (3), is the basis of our approach.

In our derivation, we intentionally avoided modeling the specific dynamics of the cell cycle, how it is regulated, and how it is coordinated with the DNA replication program. Our theory is rigorously valid independently of these aspects, provided that our initial assumptions hold.

B. Deterministic limit

In the simple case where replication proceeds deterministically and the replisome speed is a function of its position on the DNA, one has

$$\psi(x, \tau) = \delta \left(\tau - \tau_0 - \int_{x_0}^x \frac{dx'}{v(x')} \right) \tag{4}$$

where $v(x)$ is the replisome speed at position x, x_0 is the replication origin for the replisome that copied position x and τ_0 is the firing time relative to the beginning of

FIG. 1. DNA abundance in an asynchronous, exponentially growing microbial population. **(a)**. The total number of genomes (black line) and the template-copy relationship between individual genomes (tree). Individual genomes are indicated by different colors. Nodes (colored circles) represent replication initiations. Such events leave the template unchanged (same-colored descendant) and create a new genome with initial age $\tau = 0$ (differently-colored descendant). **(b)**. Example of a deterministic replication program $f(x,\tau)$ on a linear genome in which replication is initiated at two origins, one firing at time $\tau = 0$ and one at time $\tau = \tau_0$, and proceeds deterministically with constant speed. **(c)**. A stochastic version of the replication program in (b), in which replication at the second origin is initiated randomly at τ_0, τ_1 or τ_2. **(d)**. The predicted DNA abundance profile arising from replication programs (b) and (c) via Eq. (2).

replication. The speed v might take positive or negative values depending on whether the replisome proceeds in the positive genome direction. It follows from Eqs. (3) and (4) that

$$\mathcal{P}(x) = \exp\left(-\Lambda\tau_0 - \Lambda \int_{x_0}^{x} \frac{dx'}{v(x')}\right). \tag{5}$$

Solving for the speed, we obtain

$$v(x) = -\Lambda \left[\frac{d}{dx}\ln\mathcal{P}(x)\right]^{-1}, \tag{6}$$

i.e., the replication speed is inversely proportional to the logarithmic slope of $\mathcal{P}(x)$ [20, 21]. An advantage of the deterministic assumption is that it leads to a one-to-one correspondence between DNA abundance and local replisome speed thanks to Eq. (6). However, in many realistic cases, neglecting stochasticity might lead to inaccurate predictions.

Unfortunately, in the general stochastic case, different replication programs might give rise to the same DNA abundance distribution. This implies that one can not directly invert Eq. (3) to express $f(x,\tau)$ in terms of $\mathcal{P}(x)$. In these situations, one has to complement the information contained in $\mathcal{P}(x)$ with modeling assumptions, as exemplified in the following.

III. BACTERIAL DNA REPLICATION

A. Model

In this Section, we introduce a stochastic model for the DNA replication of the bacterium *E. coli*. The model includes the possibility for speed variations, as recently observed [19]. In addition, replisomes in the model can stochastically stall, as observed in single molecule experiments [26, 27].

Most bacteria, including *E. coli*, have a single circular chromosome that is replicated by two replisomes. The two replisomes start from the same origin site, one proceeding clockwise and the other counterclockwise. Replication is completed when the two replisomes meet. We assume that the two replisomes do not backtrack and that they act independently from one another until they meet. A base is therefore replicated by whichever replisome reaches it first. The joint replication program of two replisomes is therefore $f(x,\tau) = 1 - [1 - f_1(x,\tau)][1 - f_2(x,\tau)]$, where $f_1(x,\tau)$ and $f_2(x,\tau)$ are their individual replication programs, i.e. the probabilities that position x has been replicated at time τ by the respective replisome. Substituting this expression in Eq. (2) we obtain

$$\mathcal{P}(x) = 1 - \int_0^{\infty} d\tau\,\Lambda e^{-\Lambda\tau}\left(1 - f_1(x,\tau)\right)\left(1 - f_2(x,\tau)\right). \tag{7}$$

We define the genome coordinate $x \in [0, L]$ where L is the genome length and 0 (and equivalently L, since the genome is circular) is the coordinate of the replication origin. We call $x_1(t)$ and $x_2(t)$ the positions of the two replisomes along this coordinate as a function of time. The first replisome starts at $x(0) = 0$ and moves in the direction of increasing x, whereas the second starts at $x(0) = L$ and moves in the direction of decreasing x.

We express the replisome dynamics in terms of two Langevin equations

$$\frac{d}{d\tau}y_1 = v_0 h(\tau) + \sqrt{2Dh(\tau)}\,\xi_1(\tau)$$
$$\frac{d}{d\tau}y_2 = -v_0 h(\tau) + \sqrt{2Dh(\tau)}\,\xi_2(\tau)$$
$$x_1(\tau) = \max_{\tau' \leq \tau} y_1(\tau')$$
$$x_2(\tau) = \min_{\tau' \leq \tau} y_2(\tau')\,, \tag{8}$$

where $\xi_1(\tau)$ and $\xi_2(\tau)$ are Gaussian white noise sources satisfying $\langle\xi_i(\tau)\xi_j(\tau')\rangle = \delta_{ij}\delta(\tau-\tau')$. In Eq. (8), we have

FIG. 2. Bacterial DNA replication model. Parameters are: genome length $L = 5 \cdot 10^6$, growth rate $\Lambda = 2\mathrm{h}^{-1}$, baseline speed $v_0 = 10^3$ bp/s. **(a)**. Trajectories x_1, x_2 of the model (black lines) and auxiliary processes y_1, y_2 (blue lines). Backwards movements of y_1, y_2 correspond to stochastic stalling of x_1, x_2. Replication concludes at time τ_x when x_1 and x_2 first meet at a random meeting point $Z = x_1(\tau_x) = x_2(\tau_x)$. **(b)**. Bacterial replication program $f(x,\tau) = 1 - (1-f_1)(1-f_2)$ with f_1, f_2 from Eq. (9) for diffusivities $D = 10^5, 10^7, 10^8$ bp^2/s and genome ages $\tau = 100, 1200, 2400$ s. **(c)**. DNA abundance distribution for constant and oscillating replisome speed and different values of D. In the constant speed case, $h(\tau) = 1$, whereas in the oscillating speed case $h(\tau) = 1 + \delta \cos(\omega\tau + \phi)$ with $\delta = 0.5$, $\omega = 2\pi/1800$, and $\phi = 0$. **(d)**. Curvature $\mathcal{P}''(x)/\mathcal{P}(x)$ at the terminus $x = L/2$ predicted by Eq. (11) vs. numerical simulations. **(e)**. Uncertainty of the meeting point Z predicted by Eq. (12) vs. numerical simulations.

assumed that the instantaneous average speeds of the two replisomes are opposite in sign. These instantaneous speeds are modulated in time by a function $h(\tau)$, as result of varying conditions during the cell cycle (see [19]). We have assumed that the noise amplitude $\sqrt{2Dh(\tau)}$ is modulated in time by same the function $h(\tau)$ as the instantaneous average speed. This assumption is also convenient to mathematically derive the exact replication program.

The physics of Eq. (8) can be interpreted as follows. Whenever $y_i(\tau)$ attains a new maximal distance from its origin, then $x_i(\tau)$ is moving forward and it coincides with $y_i(\tau)$. If, instead, $y_i(\tau)$ is making a negative excursion from its past maximal distance, then $x_i(\tau)$ remains frozen at the value of the last maximal distance of $y_i(\tau)$. We interpret these stochastic pausing as replisome stalling events. The amplitude of the noise terms in Eqs. (8) controls the commonness of long stalling events. Stochastic trajectories generated by the model are qualitatively similar of those observed in single-molecule experiments with DNA polymerases [26, 27], see Fig. 2a.

The balance between replisome progress and stalling is governed by the dimensionless Peclet number Pe $= Lv_0/4D$. For large Pe, the dynamics are nearly deter-

ministic and long stalling events are infrequent.

A consequence of Eq. (8) is that the individual replication program $f_i(x,\tau)$ is equal to the first-passage probability of the associated process y_i through x. This first-passage probability is expressed by

$$f_i(x,\tau) = \int_0^{H(\tau)} \sqrt{\frac{\mu_i^3}{2\pi\sigma_i^2 u^3}} \exp\left(-\frac{1}{2}\frac{(u-\mu_i)^2}{\sigma_i^2}\frac{\mu_i}{u}\right) du\,,$$

$$(9)$$

where $H(\tau) = \int_0^\tau h(u)du$ and

$$\mu_1 = x/v_0, \qquad \mu_2 = (L-x)/v_0,$$
$$\sigma_1^2 = 2Dx/v_0^3, \qquad \sigma_2^2 = 2D(L-x)/v_0^3\,. \qquad (10)$$

Eqs. (9) and (10) are derived in Appendix A. We compute the DNA abundance $\mathcal{P}(x)$ by substituting Eq. (9) into Eq. (7), see Fig. 2b. We numerically evaluate the final integral over τ appearing in Eq. (7).

The main effect of diffusivity is to smoothen the DNA abundance distribution predicted by the model, see Fig. 2c. This effect is particularly pronounced at the expected meeting point $x = L/2$ of the two replisomes,

FIG. 3. Bacterial model fitted to the data of Ref. [19] for speed oscillations as given in Eq. (13). We fitted the parameters v_0, D, δ, ω, and ϕ from the measured DNA abundances, growth rate Λ was measured separately (see Appendix E). **(a)**. Observed DNA abundance and model predictions for E. coli cultures growing exponentially at temperatures $T = 17^\circ C, 22^\circ C, 27^\circ C, 32^\circ C, 37^\circ C$. **(b)**. Relative decrease in residuals $(SQ(\Theta) = \sum_{i=1}^{N}(a_i - \lambda \mathcal{P}(x_i|\Theta))^2/\sigma_i^2$, see Eq. (E2)) of oscillatory model (Θ_{osc}) vs. constant-speed model (Θ_c). **(c)** Average instantaneous speed $v(\tau) = v_0 h(\tau)$ as a function of the fraction τ/τ_2 of the doubling time $\tau_2 = \log(2)/\Lambda$. **(d)** Average instantaneous speed $v(\tau) = v_0 h(\tau)$ as the function of the time τ since replication initiation.**(e)** Average instantaneous speed $v(\tau) = v_0 h(\tau)$ as a function of the replication progress, i.e. of the fraction $2\int_0^\tau v(u)du/L$ of the genome replicated. In (c-d) we omitted $17^\circ C$ since the effect of speed fluctuations on $\mathcal{P}(x)$ is negligible at that temperature (see panel b).

where the DNA abundance exhibits a cusp for $D = 0$ but is smooth for positive D, see Figure 2c. In contrast, far from the meeting point diffusivity does not affect the DNA abundance much, see Appendix B. By exploiting Eq. (9), we now quantify the effect of diffusivity. In the case of constant speed $v(\tau) = v_0$, we find that the relative curvature of $\mathcal{P}(x)$ at the expected meeting point is approximated by

$$\frac{\mathcal{P}''(L/2)}{\mathcal{P}(L/2)} \approx \frac{2\Lambda}{\sqrt{\pi D L v_0}} , \qquad (11)$$

see Fig. 2d and Appendix B. In principle, Eq. (11) can be used to estimate the diffusion coefficient from the curvature in the terminus region, without having to fit the entire distribution.

From the point of view of trajectories, the smoothing of $\mathcal{P}(x)$ for $D > 0$ occurs because the two replisomes no longer necessarily meet exactly at $x = L/2$. In particular, we find that the uncertainty on the location Z of the meeting point is approximately equal to

$$\langle (Z - L/2)^2 \rangle \approx \frac{DL}{2v_0} , \qquad (12)$$

see Fig. 2e. Equation (12) is derived in Appendix C.

B. Comparison with E.coli sequencing data

In this Section, we fit our model of replisome dynamics in Eq. (8) to previously measured DNA abundance data [19] from wild type *E. coli* grown at different temperatures. We assume that the speed and diffusion coefficient are modulated in time by a function

$$h(\tau) = 1 + \delta \cos(\omega\tau + \phi), \qquad (13)$$

see Fig. 2. We fit the model to the experimental data with v_0, D, δ, ω, and ϕ as free parameters (see Appendix E).

The model fits the experimental DNA abundances very well (Fig. 3a) and yields highly consistent estimates of v_0 among replicates across all temperatures, see Table I. At temperatures above $17^\circ C$ we find robust evidence of speed fluctuations. The oscillatory model provides consistent estimates of δ, ω and ϕ in these cases, with an improvement of the quality of fit ranging between 20% to 40% percent compared to a constant-speed model (Fig. 3b). At $17^\circ C$, the effect of the speed fluctuations on $\mathcal{P}(x)$ is small and, consequently, the uncertainty in the associated parameters ω and ϕ is high. Regardless of temperature, model selection appears to prefer a vanishing value of D for some replicates. The reason is likely that the estimates of D correspond to Peclet numbers

T [°C]	v_0 [bp/s]	D [kbp^2/s]	δ	ω [rad/h]	ϕ [rad]
17	230 ± 30	1.6 ± 1.4	0.29 ± 0.1	5.8 ± 4.2	1.3 ± 1.5
22	350 ± 21	0.5 ± 0.8	0.19 ± 0.07	3.9 ± 0.6	3.1 ± 0.5
27	530 ± 20	1.0 ± 1.2	0.19 ± 0.03	8.7 ± 0.4	2.0 ± 0.2
32	810 ± 57	1.7 ± 1.8	0.12 ± 0.04	16 ± 1.3	1.5 ± 0.2
37	950 ± 24	1.2 ± 2.3	0.19 ± 0.03	15 ± 0.4	2.9 ± 0.2

TABLE I. Parameter estimates for time-dependent speed $v(t) = v_0\big(1 + \delta \cos(\omega t + \phi)\big)$. The reported uncertainties represent the variability over replicates.

approximately in the range from Pe = 200 to Pe = 1000, which lies close to the detection threshold; see the discussion in Appendix E.

Comparing the speed oscillations across different temperatures shows that their frequency appears linked with the population doubling time. The oscillations for 22°C to 37°C align best when time is rescaled according to the duration of a cell cycle, see Fig. 3c. The speed consistently attains a minimum after one doubling time $\tau_2 = \log(2)/\Lambda$ when the next replication cycle starts and the number of active forks is thus increases. When plotted for comparison against absolute time since replication initiation, the speed fluctuations do not align (Fig. 3d). When plotted against replication progress (i.e. replisome position), temperatures 27°C to 37°C do align, but 22°C does not (Fig. 3e).

Ref. [19] first observed the speed oscillations and explained them using a computational model in which speed is modulated in space, rather than in time. Our approach permits to analytically solve also the model of Ref. [19] in the small-noise limit. The resulting parameter values match those of Ref. [19] well, and are consistent with our temporally modulated model as far as that can be expected; see Appedix D for details. The model with temporal speed oscillations yields a better fit for the majority of samples; the improvement in likelihood is, however, small (Fig. 7b in Appendix D).

IV. EUKARYOTIC DNA REPLICATION

A. Model

In this Section, we apply our approach to eukaryotic DNA replication. The eukaryotic case is substantially more complex than the bacterial one, because of the presence of multiple, randomly activated replication origins. In this case, theoretical progress [28, 29] has made use of an analogy with the physics of freezing/crystallization kinetics, as described by the so-called Kolmogorov-Johnson-Mehl-Avrami model [30–32], see also [33, 34].

We briefly summarize this idea and then extend it to asynchronously growing populations. When an origin fires, two replisomes start moving from that origin in opposite directions. We consider a given location x in a eukaryotic genome at a time τ after the beginning of the replication program. The probability $f(x,\tau)$ that this genomic location has been already replicated can then be seen as the probability that the past "light-cone" $V_{x,\tau}$ of the space-time point x, τ contains at least one origin firing event, see Fig. 4a. This elegant argument circumvents the problem of determining which origin replicated a given genomic location.

Assuming that replisomes progress deterministically with constant speed v_0, the past light-cone is the set of space-time points x', τ' from which x is reachable by a replisome within a time τ, i.e.,

$$V_{x,\tau} = \left\{ (x', \tau') \,\big|\, |x - x'| \leq v_0\,(\tau - \tau') \right\}, \qquad (14)$$

see Fig. 4a. Following [1, 35], we now assume that origins fire independently from one another in space and time with position- and time-dependent rates $I(x,\tau)$. This assumption implies that the space-time points at which origins fire form a two-dimensional Poisson point process.

The probability that location x has been replicated by time τ is then expressed by

$$f(x,\tau) = 1 - \exp\left(-\iint_{V_{x,\tau}} dx' d\tau'\, I(x', \tau') \right). \qquad (15)$$

We note that Eq. (15) would remain valid if we chose a different replisome dynamics, corresponding to a form of the light-cone $V_{x,\tau}$ other than that expressed by Eq. (14). Using Eq. (2), we formally express the DNA abundance distribution as

$$\mathcal{P}(x) = 1 - \Lambda \int_0^\infty d\tau \exp\left(-\Lambda\tau - \iint_{V_{x,\tau}} dx' d\tau'\, I(x', \tau') \right). \qquad (16)$$

We now focus on budding yeast, where origins are characterized by a specific DNA sequence [6]. We therefore assume that origins are fired at well-defined locations $x_1, \ldots, x_K$ on the chromosome, meaning $I(x,\tau) = \sum_{j=1}^K I_j(\tau)\delta(x - x_j)$. We further assume that firing rates are constant in time, so that

$$I(x,\tau) = \sum_{j=1}^K I_j^\star \delta(x - x_j). \qquad (17)$$

In budding yeast, the origin firing rate was observed to be time-dependent [8], so that this second assumption is not fully accurate. However, it is still a useful approximation that considerably simplifies the model.

To express the DNA abundance $\mathcal{P}(x)$ for firing rates as in Eq. 17, we consider the travel time τ_k from origin x_k to a given position x,

$$\tau_k = \frac{|x - x_k|}{v_0}. \qquad (18)$$

At fixed x, we reorder the origins so that $0 < \tau_1 < \cdots < \tau_K$. We then define the effective replication times

$$\mathcal{T}_k = \tau_k + \sum_{j=1}^k (\tau_k - \tau_j)\, I_j^\star/\Lambda \qquad (19)$$

FIG. 4. Eukaryotic model. **(a)**. Past light-cone $V_{x,\tau}$ of space-time points within which at least one origin must have fired for location x to be replicated by time τ assuming replisome speed v. **(b)**. Inference of origin location and intensity from simulated DNA abundance. Top plot shows the model (black line) fitted to simulated abundances (green line), bottom plot shows the inferred 26 origins (black bars) and 39 true origins (green bars). Intensities are plotted on a log-scale as $I_i^\star/v$ [1/bp]. Parameters are: $v = 27$ bp/s, $\Lambda = 9.6 \cdot 10^{-5}$ 1/s (doubling time 120 minutes). The true origin locations are known origins on *S. cerevisiae* W303 chromosome IV. True intensities are log-uniformly distributed on $[10^{-5}, 2 \cdot 10^{-3}]$. The resulting true abundances are perturbed by multiplicative Gamma-distributed noise with mean 1.0 and coefficient of variation 0.04 to mimic measurement errors.

which collectively take into account the effect of all origins. We also introduce the effective cumulative weights

$$W_k = 1 + \sum_{j=1}^{k} I_j^\star / \Lambda, \qquad (20)$$

with the convention that $W_0 = 1$. Using these definitions, the DNA abundance is expressed by

$$\mathcal{P}(x) = \sum_{k=1}^{K} \left(\frac{1}{W_{k-1}} - \frac{1}{W_k} \right) e^{-\Lambda \mathcal{T}_k}. \qquad (21)$$

Equation (21) is derived in Appendix F.

We implemented a simulated annealing algorithm to infer the origin number, location, and intensities based on DNA abundance data via Eq. (21). Our inference procedure treats Λ/v_0, the number of origins K, the origin positions $x_1, \ldots, x_K$, and the re-scaled firing rates $I_1^\star/v_0, \ldots, I_K^\star/v_0$ as free parameters. We call the compound parameter $I_j^\star/v_0$ the intensity of origin j. Our algorithm uses the Akaike information criterion (AIC) as the cost function to avoid over-fitting (details in Appendix G).

We first tested this inference procedure on artificial data, in which the genome length (1.6Mbp) and origins distribution are comparable to those of the longest chromosome of budding yeast (chromosome IV). Our inference algorithm detected the location and intensity of 26 out of 39 true origins with high accuracy (Fig. 4b; median distance to true origin 1.5kb, median relative error of intensity 40% if non-resolved clusters of true origins are merged; true intensities range over two orders of magnitude). The 13 non-recovered origins either have low intensity or were merged with another origin in close proximity.

B. Application to experimental data

We now apply our model and inference procedure to experimentally measured DNA abundance in an asyn-

chronous populations of budding yeast (*S. cerevisiae* W303) [36]. Our algorithm infers $K = 234$ origins across the 16 yeast chromosomes. The predicted DNA abundance well matches the experimental one, see Fig. 5a.

The inferred origins locations correlate with known origins of *S. cerevisiae* when binned on a length scale of about 7kb, see Fig. 5b. A second correlation peak at 100kb suggests a large-scale pattern in the distribution of origins. As expected, the peak at 7kb vanishes when known origin locations are randomly shifted by ± 35kbp, or shuffled by randomly reordering chromosomes. The results of the correlation analysis are corroborated by matching each inferred origin to the closed known origin (figure 5c; median distance 3.9kb) respectively each known origin to the closest inferred origin (figure 5d; median distance 6.6kb). In both cases, the average distances are significantly increased if the known origins are shifted or shuffled.

Despite the simplifying assumptions, our method can thus successfully recover origins of replication with a resolution on the order of a few kilobases.

V. EXTENSIONS OF THE THEORY

A. Non-exponential growth

Our approach can be generalized to populations that do not grow exponentially. We here assume that the number of genomes grow with time as an arbitrary function $N_g(t)$. In this case, following the same logic of Section II, we find that Eq. (2) becomes

$$\mathcal{P}(x, t) = \int_0^\infty d\tau P(\tau; t) f(x, \tau) \qquad (22)$$

where $P(\tau; t) \propto N_g'(t - \tau)$. The proportionality constant can be determined by imposing normalization of $P(\tau)$. After integrating Eq. 22 by parts, we obtain

$$\mathcal{P}(x, t) = \langle N_g(t - \tau) \rangle. \qquad (23)$$

FIG. 5. Eukaryotic model fitted to *S. cerevisiae* data from [36]. Fitted parameters are: Λ/v, origin positions $x_1, \ldots, x_n$ and intensities $I_1^\star/v, \ldots, I_n^\star/v$. The fitted number of origins is $K = 234$. Known origins used for validation were lifted from the annotated genome for strain 288C (RefSeq assembly *R64-3-1*, accession *GCF_000146045.2*) using *liftoff* [37]. We excluded the mitochondrial genome. **(a)**. Observed (green circles) and predicted (black line) DNA abundances (top) and inferred origin positions and intensities (bottom). Known origin positions (blue) shown for comparison. **(b)**. Correlation of estimated ($K = 234$) and known ($K = 354$) origin densities at different smoothing length scales (yellow). For comparison, we also show the correlation to the density of shuffled (green) and randomly shifted (blue) known origins. **(c)**. Distribution of distances between estimated and closest known origins. **(d)**. Distribution of distances between known and closest estimated origins. In (c) and (d), the y-axis is non-linearly scaled to enlarge the region of interest. Horizontal lines indicate the medians. Reported p-values are computed using Mann–Whitney U tests.

This generalization can be applied, for example, to bacterial or eukaryotic populations that grow in a partially synchronous manner. In such cases, $N_g(t)$ might exhibit temporal peaks.

We note that, in Eqs. (22) and (23) we implicitly made the assumption that the replication program $f(x, \tau)$ does not depend on time. The validity of this assumption should be verified for populations that are not growing in steady conditions. Moreover, in practical application, the limiting factor can be the accuracy with which $N_g(t)$ can be estimated.

B. Non-immortal genomes

For the purpose of computing the DNA abundance distribution, our assumption of immortal genomes is justified as long as typical genome lifetimes $\tau_{\max}$ are much longer than the doubling time $\tau_2 = \log(2)/\Lambda$. This is

usually a realistic assumption, but in some cases one may want to generalize our approach to include genome degradation. In such case, the exponential distribution in Eq. (2) should be replaced with the age distribution $P(\tau)$ of surviving genomes. In particular, if genomes survive until age τ with probability $s(\tau)$, the age distribution is expressed by

$$P(\tau) \propto \Lambda e^{-\Lambda\tau} s(\tau) \qquad (24)$$

up to a normalization constant.

C. Multiple genome types

The theory we developed so far applies to statistically identical genomes. This assumption is sometimes referred to as "identity at birth" in the literature on renewal processes [38]: all genomes are born identical, and the differences that they might develop as they synthe-

size can be seen as outcomes of independent realizations of the same stochastic process.

There might be cases in which this assumption does not hold. For example, certain knockout mutants of *E. coli* are known to lead to defective genome types that are inheritable [39]. To address these scenarios, we extend our theory to a case in which different population types coexist in the population, each characterized by a different replication program.

We call $N_g^{(i)}(t)$ the number of genomes of type i at time t in the population, with $i = 1 \ldots k$. The total number of genomes is $N_g(t) = \sum_i N_g^{(i)}(t)$. Each genome of type i can be replicated into a genome of type i' according a certain stochastic process. We assume that this stochastic process satisfies the Perron-Frobenius theorem, so that at long time the number of genomes are given by

$$N_g^{(i)}(t) \propto r_i \, e^{\Lambda t}. \tag{25}$$

Here Λ is the leading eigenvalue of the dynamics, corresponding to the exponential growth rate of the population. The vector $(r_1 \ldots r_k)$ is the leading eigenvector, having non-negative real entries according to the Perron-Frobenius theorem, and normalized such that $\sum_i r_i = 1$. The resulting replication program can then be obtained as an average over the genome types:

$$f(x, \tau) = \sum_{i=1}^{k} r_i f^{(i)}(x, \tau). \tag{26}$$

This apparent replication program can be substituted into Eq. (2) to determine the DNA abundance distribution.

D. Numerical simulations and analogy with stochastic resetting

The models we considered in Sections III and IV are analytically solvable. However, in more complex cases, analytical solutions might not be available. In this Section, we briefly present an interpretation of the DNA abundance distribution $\mathcal{P}(x)$ as the steady-state of a dynamical process implementing stochastic resetting [40, 41]. This analogy can be used for efficient numerical simulations.

The models introduced in Sections III and IV are described by linear equations. Formally, we assume that we can always write the stochastic process associated with a certain DNA replication program in terms of linear evolution operator $\hat{\mathcal{L}}$, so that

$$\partial_\tau f(x, \tau) = \hat{\mathcal{L}} f(x, \tau). \tag{27}$$

We now introduce a modified version of this equation:

$$\partial_\tau f(x, \tau) = \hat{\mathcal{L}} f(x, \tau) - \Lambda f(x, \tau). \tag{28}$$

Equation 28 can be interpreted as describing a replication program equivalent to that of Eq. (27), but that is

stochastically reset to its beginning at rate Λ. Stochastic simulations of this process can be easily implemented numerically.

We note that $\mathcal{P}(x)$ is a steady solution of Eq. (28), i.e., it satisfies

$$\left(\hat{\mathcal{L}} - \Lambda \right) \mathcal{P} = 0. \tag{29}$$

This means that we can determine $\mathcal{P}(x)$, up to an appropriate normalization constant, by sampling the stochastic dynamics described Eq. (28) at steady state. Such stochastic simulations can be efficiently implemented numerically [19].

VI. CONCLUSIONS

In this paper, we introduced a general theory that connects the DNA replication program with the abundance of DNA fragments that one should expect in an asynchronously growing population of cells. Our theory builds up on previous approaches [3, 13, 19, 20, 23] and has the advantage of being based on a minimal set of realistic assumptions and allowing for stochastic DNA replication programs. As we have demonstrated, these key properties make our theory applicable to a broad range of organisms, from bacteria to eukaryotes.

In the case of bacterial DNA replication, we proposed a model in which the replisome speed is modulated in time and replisomes can stochastically stall. We solved the model exactly and fitted its prediction against sequencing data of *E.coli* growing at different temperatures [19]. The fits show that the period of speed oscillations matches the population doubling time, or equivalently the time interval between consecutive fork firing. Our model that assumes time-periodic variations leads to slightly better fits than the space-periodic variations postulated in [19]. Taken together, these observations support that the cause of oscillations must be synchronized with the cell cycle, or alternatively with the fork firing rate. A possible candidate would be competition among multiple forks on the same genome [19]. An important advancement compared to the theory proposed in [19] is that the approach introduced in this paper leads to an analytical expression for the DNA abundance distribution, which considerably simplifies the fitting procedure.

The fits of the *E.coli* data reveal that the Peclet number characterizing the replisome dynamics is rather large. On the one hand, this observation confirms that simpler approaches that neglect stochasticity [20, 21] provide reliable results, at least in the case of wild type *E.coli*. On the other hand, we have argued that the diffusion constant, albeit small, provides important information about the uncertainty of the replisome meeting point. In *E.coli*, the Tus-Ter system is know to sets bounds on the region in which replisomes can meet [42], thereby likely affecting this accuracy. It will be interesting for future studies to apply our approach to mutant strains, to see

whether they are characterized by a different degree of uncertainty.

Stochasticity is definitely crucial in the eukaryotic case. We have used our approach to estimate the origins location and intensities for budding yeast. For simplicity, we assumed that replisome speed is constant; that origins are placed at well defined sites and fire at an origin-dependent rate that is constant in time. Despite these simplifying assumptions, the predictions of our algorithm recover the majority of known origins in budding yeast with an accuracy on the order of kilobases. The accuracy can likely be further increased by exploiting advances in sequencing technology, in particular increased sequencing depth and read lengths; and by employing an optimization algorithm tailored more specifically to our model. This makes the combination of deep sequencing of asynchronous populations and our inference approach a cost-effective way of locating origins of replication in any single-cell eukaryotic species which can be cultured and sequenced.

Given the flexibility and generality of our approach, it will be interesting in the future to relax some of the assumptions that we currently make, most notably that origins fire at a constant rate. This would allow our approach to not only accurately recover origin locations but also their relative timing behaviour, and thus to provide a more complete picture of the replication program.

ACKNOWLEDGMENTS

We thank S. Hauf, C. Plessy, and Y. Yokobayashi for fruitful discussions. We thank A. Sassi for feedback on a preliminary manuscript. This work was supported by JSPS KAKENHI Grant No. 23H01146 (to SP). DB thanks Vellore Institute of Technology, Vellore for providing VIT SEED Grant-RGEMS Fund (SG20220060) for carrying out this research work.

Appendix A: Bacterial DNA replication with time-dependent speed

We consider the Langevin Eqs. (8). We write the associated Fokker-Planck equation for the first replisome:

$$\frac{\partial}{\partial \tau} p = h(\tau) \left(-v_0 \frac{\partial}{\partial y} p + D \frac{\partial^2}{\partial y^2} p \right). \tag{A1}$$

Equation (A1) shows that the function $h(\tau)$ acts as global time rescaling. Therefore, if $\hat{p}$ is a solution of the Fokker-Planck equation with constant drift and diffusion terms

$$\frac{\partial}{\partial \tau} \hat{p} = -v_0 \frac{\partial}{\partial y} \hat{p} + D \frac{\partial^2}{\partial y^2} \hat{p} \tag{A2}$$

under certain boundary and initial conditions, then

$$p(y, \tau) = \hat{p}(y, H(\tau)) \tag{A3}$$

solves the original Fokker-Planck Eq. (A1) with the same boundary and initial conditions, where $H(\tau) = \int_0^\tau h(u)$. We impose, in particular, an absorbing boundary condition at x, $\hat{p}(x, \tau) = 0 \; \forall \tau$ and an initial condition $p(y, \tau = 0) = \delta(y)$. The flux $\hat{J} = v_0 p - D \partial_y p$ through the absorbing boundary is given by an inverse Gaussian distribution:

$$\hat{J}(x, \tau) = \frac{x}{\sqrt{4\pi D \tau^3}} e^{-\frac{(x - v_0 \tau)^2}{4D\tau}}. \tag{A4}$$

Since the first-passage density of Eq. (A1) equals the flux $J(x, \tau) = h(\tau)\hat{J}(x, H(\tau))$, we have

$$\frac{\partial}{\partial \tau} f_1(x, \tau) = h(\tau)\hat{J}(x, H(\tau)) = \frac{h(\tau)x}{\sqrt{4\pi D H^3(\tau)}} e^{-\frac{(x - v_0 H(\tau))^2}{4D H(\tau)}}. \tag{A5}$$

The same approach can be used for the second replisome after a suitable change of coordinates and initial conditions. The individual replication programs of the two replisomes are therefore given by Eqs. (9) and (10), where we reparametrized the distributions in terms of their means and variances.

Appendix B: Effect of diffusivity on the shape of the bacterial DNA abundance distribution

We here study the effect of increasing D on the shape of the DNA abundance distribution $\mathcal{P}(x)$. We first focus on the the meeting point region, and then consider the region far from the expected meeting point.

We call the expected meeting point $x = L/2$ the "replication terminus". Since this is the last location on the genome to be replicated, $\mathcal{P}$ attains its global minimum there. For diffusivity $D = 0$, $\mathcal{P}$ exhibits a cusp at that minimum, and thus infinite curvature. For positive diffusivity D, the cusp vanishes, the curvature is finite and decreases with D (Fig. 2c-d). To quantitatively link D with this curvature, we first use that Eq. (8) is symmetric under the swap of replisomes and mirroring of the genomic coordinate: $f_1(x, \tau) = f_2(L - x, \tau)$. As a consequence, $f'(L/2)$ vanishes for all time τ and therefore $\mathcal{P}'(L/2) = 0$ according to Eq. 2. The curvature at $x = L/2$ is thus simply $\mathcal{P}''(L/2)$. We focus on the relative curvature expressed by

$$\frac{\mathcal{P}''(L/2)}{\mathcal{P}(L/2)} = \frac{\int_0^\infty d\tau \Lambda e^{-\Lambda \tau} f''(L/2, \tau)}{\int_0^\infty d\tau \Lambda e^{-\Lambda \tau} f(L/2, \tau)}. \tag{B1}$$

Direct evaluation of the two integrals in Eq. (B1) does not look tractable for the replication program from Eqs. (7) and (9). To make progress, we approximate the replication time distribution with a Gaussian:

$$\psi(x, \tau) = \frac{\partial}{\partial \tau} f(x, \tau) \approx \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(t - \mu(x))^2}{2\sigma^2(x)}}, \tag{B2}$$

where the mean $\mu(x)$ and the variance $\sigma^2(x)$ have to be determined. It then follows from Eq. (3) that

$$\mathcal{P}(x) = \int_0^\infty d\tau e^{-\Lambda\tau}\psi(x,\tau) \approx e^{-\mu(x)\Lambda+\sigma^2(x)\Lambda^2/2}. \quad (B3)$$

Here, we have approximated the integral by extending the lower extreme to $-\infty$. We expect the error caused by this approximation to be negligible, since the Gaussian distribution that approximates $\psi(x,\tau)$ must be concentrated on the positive real numbers. By using that $f'(L/2,\tau) \equiv 0$ and thus $\mu'(x) = 0$, and assuming that the dependence of σ on x is small enough to be ignored, we obtain

$$\frac{\mathcal{P}''(L/2)}{\mathcal{P}(L/2)} \approx -\Lambda\mu''(L/2). \quad (B4)$$

Since $\mu(x)$ was left unspecified above, we have to evaluate the integral $\mu''(L/2) = \partial_x^2 \int \tau\partial_\tau f(x,\tau)d\tau|_{x=L/2}$. To make this tractable, we again replace f with an approximation. Instead of the replacing $\psi = \partial_\tau f$ with a Gaussian density for which $\mu(x)$ and $\sigma(x)$ are unknown as we did above, we now replace the individual inverse Gaussian laws of f_1, f_2 from Eq. (9) with Gaussian distributions. For these individual laws, the means and variances are given by Eq. (10) which yields

$$\hat{f}(x,\tau) = 1 - \left[1 - \Phi\left(\frac{\tau - x/v_0}{\sqrt{2Dx/v_0^3}}\right)\right] \times$$
$$\left[1 - \Phi\left(\frac{\tau - (L-x)/v_0}{\sqrt{2D(L-x)/v_0^3}}\right)\right]. \quad (B5)$$

where we have for simplicity assumed that the replisome speed is constant, i.e. $h(\tau) = 1$. With $\hat{f}$ in place of f, we now proceed to find

$$\mu''(L/2) \approx \frac{\partial^2}{\partial x^2}\int_{-\infty}^\infty \tau\frac{\partial}{\partial\tau}\hat{f}(x,\tau)d\tau\bigg|_{x=L/2}$$
$$= \frac{\partial^2}{\partial x^2}\left(\int_0^\infty d\tau\left(1-\hat{f}(x,\tau)\right) - \int_{-\infty}^0 d\tau\hat{f}(x,\tau)\right)\bigg|_{x=L/2}$$
$$= -\int_{-\infty}^\infty \hat{f}''(L/2,\tau)d\tau$$
$$= -\frac{2}{\sqrt{\pi DLv_0}} \quad (B6)$$

and by inserting into Eq. (B4) we find the expression for the normalized curvature of the DNA abundance at the terminus stated in Eq. (11).

Eq. B3 also provides an approximation for the value of $\mathcal{P}(x)$ itself at the terminus. Similarly to $\mu''(L/2)$, we find that

$$\mu(L/2) \approx \int_{-\infty}^\infty \tau\frac{\partial}{\partial\tau}\hat{f}(x,\tau)d\tau$$
$$= \int_0^\infty \left(1 - \hat{f}(x,\tau)\right)d\tau - \int_{-\infty}^0 \hat{f}(x,\tau)d\tau$$
$$= \frac{L}{2v_0} - \sqrt{\frac{DL}{\pi v_0^3}} \quad (B7)$$

and

$$\sigma^2(L/2) \approx \int_{-\infty}^\infty \tau^2\frac{\partial}{\partial\tau}\hat{f}(x,\tau)d\tau - \mu^2(L/2)$$
$$= 2\int_0^\infty \tau\left(1 - \hat{f}(x,\tau)\right)d\tau$$
$$- 2\int_{-\infty}^0 \tau\hat{f}(x,\tau)d\tau - \mu^2(L/2)$$
$$= \frac{\pi-1}{\pi}\frac{DL}{v_0^3}. \quad (B8)$$

Approximating $\mu(L/2 + \epsilon) \approx \mu(L/2) + \epsilon^2\mu''(L/2)/2$ and $\sigma^2(L/2+\epsilon) \approx \sigma^2(L/2)$ in Eq. (B3) yields the approximation

$$\mathcal{P}(L/2 + \epsilon) \approx \exp\left[-\frac{\Lambda L}{2v_0} + \Lambda\sqrt{\frac{DL}{\pi v_0^3}}\right.$$
$$\left.+ \frac{\pi-1}{\pi}\frac{\Lambda^2 DL}{2v_0^3} + \epsilon^2\frac{\Lambda}{\sqrt{\pi DLv_0}}\right], \quad (B9)$$

close to the replication terminus. The curvature at $x = L/2$ is therefore

$$\mathcal{P}''(L/2) \approx \frac{2\Lambda}{\sqrt{\pi DLv_0}}\exp\left[-\frac{\Lambda L}{2v_0} + \Lambda\sqrt{\frac{DL}{\pi v_0^3}}\right.$$
$$\left.+ \frac{\pi-1}{\pi}\frac{\Lambda^2 DL}{2v_0^3}\right]. \quad (B10)$$

We now discuss the shape of the DNA abundance far from the terminus region. Within these regions, only one of the two replisomes plays a relevant role. For simplicity, we focus on the region replicated by replisome 1. We also assume constant replisome speed, i.e. $h(\tau) = 1$. Since replisome 2 has virtually no chance of reaching the area of interest before replisome 1, we assume $f(x,\tau) \approx f_1(x,\tau)$. The DNA abundance $\mathcal{P}(x)$ can then be explicitly found from the characteristic function of the inverse Gaussian distribution; the expansion $\sqrt{1+x} \approx 1 + x/2 - x^2/8$ yields for small D

$$\mathcal{P}(x) \approx \exp\left[\left(-\frac{\Lambda}{v_0} + D\frac{\Lambda^2}{v_0^3}\right)x\right]. \quad (B11)$$

The change in rate due to D is therefore of order $D\Lambda^2/v^3$.

T	$\overline{v}$ [bp/s]	D [kbp^2/s]	δ	ω [rad/Mbp]	ϕ [rad]
17	230 ± 22	1.0 ± 1.2	0.18 ± 0.07	6.2 ± 6.7	2.3 ± 1.6
22	350 ± 37	1.0 ± 1.6	0.23 ± 0.15	2.8 ± 0.73	3.4 ± 0.72
27	540 ± 17	0.4 ± 0.5	0.18 ± 0.04	4.6 ± 0.11	2.1 ± 0.11
32	820 ± 57	1.4 ± 2.3	0.12 ± 0.04	5.5 ± 0.13	1.6 ± 0.16
37	970 ± 25	2.9 ± 2.3	0.18 ± 0.02	4.3 ± 0.15	3.0 ± 0.14

TABLE II. Parameter estimates for position-dependent speed $v(x) = \overline{v}(1 + \delta \cos(\omega x + \phi))$ model of Bhat *et al.*[19].

Appendix C: Uncertainty of the meeting point

We now consider the random point Z at which the two replisomes meet and its uncertainty $\langle (Z - \langle Z \rangle)^2 \rangle$. Let $\tau_1(x)$ and $\tau_2(x)$ be the times at which replisomes 1 and 2 reach location x, respectively. Then, $\tau_1(z) < \tau_2(z)$ for all points z to the left of the meeting point Z, and $\tau_1(z) > \tau_2(z)$ for all points to the right of Z. Therefore

$$\mathbb{P}(Z < z) = \mathbb{P}(\tau_1(z) > \tau_2(z)) . \tag{C1}$$

We note that the means and variances in Eq. (10) are the means and variances of $H(\tau_i(x))$. Since $H(t)$ increases monotonically, we substitute it into Eq. (C1):

$$\mathbb{P}(Z < z) = \mathbb{P}(H(\tau_2(z)) - H(\tau_1(z)) < 0). \tag{C2}$$

By using that $\tau_1(z)$ and $\tau_2(z)$ are independent, and substituting the means and variances from Eq. (10), we find the mean and variance of $H(\tau_2(z)) - H(\tau_1(z))$ to be $(L - 2z)/v_0$ and $2DL/v_0^3$, respectively. By approximating the distribution of the difference between the two replication times with a Gaussian, we obtain

$$\mathbb{P}(Z < z) \approx \Phi\left(\frac{-(L - 2z)/v_0}{\sqrt{2DL/v_0^3}} \right) = \Phi\left(\frac{z - L/2}{\sqrt{DL/2v_0}} \right) \tag{C3}$$

where $\Phi(x) = 1/2 + \text{erf}(x/\sqrt{2})/2$ is the cumulative function of the Gaussian distribution. The uncertainly about the meeting point is therefore given by Eq. (12).

Appendix D: Bacterial DNA replication with position-dependent speed

In this Appendix, we solve a model with position-dependent speeds $v_1(x)$ and $v_2(x)$ similar to that in Ref. [19] in the limit of small diffusivity. This will allow us to directly compare time-dependent and position-dependent speed models. We allow the diffusion constants $D_1(x)$, $D_2(x)$ to vary in space as well. The Langevin equations are:

$$\frac{d}{d\tau} x_1 = v_1(x) + \sqrt{2D_1(x)}\, \xi_1(\tau)$$

$$\frac{d}{d\tau} x_2 = v_2(x) + \sqrt{2D_2(x)}\, \xi_2(\tau). \tag{D1}$$

where $v_1(x) > 0$ and $v_2(x) < 0$. As in the case of Eq. (8), the initial conditions are $x_1(0) = 0$, $x_2(0) = L$ and we define replication programs $f_1(x, \tau)$, $f_2(x, \tau)$ via the first-passage times of $x_1(\tau)$ and $x_2(\tau)$, respectively, through position x.

We now assume that (i) $v_1(x)$, $v_2(x)$ change only slowly, (ii) $D_1(x)$, $D_2(x)$ are small in comparison, and (iii) x is not too close to the replication origin. Under these assumptions, replication times are approximately additive, meaning that if $\tau_{x_0 \to x_1}$ is the (random) time a replisome originating from x_0 takes to reach x_1, then $\tau_{x_0 \to x_1}$ and $\tau_{x_1 \to x_2}$ are independent, and $\tau_{x_0 \to x_2} = \tau_{x_0 \to x_1} + \tau_{x_1 \to x_2}$. For large enough x, we invoke the central limit theorem for the sum $\tau_x = \tau_{x_0 \to x_1} + \cdots + \tau_{x_k \to x}$ and conclude that ψ_1 and ψ_2 are approximately Gaussian, implying

$$f_i(x, \tau) \approx \frac{1}{2} + \frac{1}{2} \text{erf}\left(\frac{\tau - \mu_i(x)}{\sqrt{2}\sigma_i(x)} \right) \tag{D2}$$

To find the position-dependent mean $\mu_i(x)$ and variance $\sigma_i^2(x)$, we rely on $v_i(x)$ changing sufficiently slowly relative to $D_i(x)$ so that we can find a subdivision $x_0 \leq \cdots \leq x_k \leq x$ where $v_i(x)$ is approximately constant in each interval. By approximating the mean and variance of $\tau_{x_l \to x_{l+1}}$ with the constant-speed solution, we obtain

$$\mu_i(x) = \int_{x_i(0)}^{x} \frac{dx'}{v_i(x')}, \quad \sigma_i^2(x) = \int_{x_i(0)}^{x} \frac{2D_i(x')}{v_i^3(x')} dx'. \tag{D3}$$

We now compute μ_i and σ_i^2 for the position-dependent speed fluctuations from Ref. [19] of the form

$$v(x) = \overline{v}(1 + \delta \cos(\omega x + \phi)) \tag{D4}$$

where x is the distance of the replisome from the origin. We note that because we express $v(x)$ in terms of distance travelled, the speed of the two replisomes when they traverse the same physical location will in general differ. Since integration of $1/v(x)$ and $1/v^3(x)$ is cumbersome, we first compute the following integrals of powers of $g(u) = 1 + \delta \cos u$, which are valid for $z \in (-\pi, \pi)$:

$$\hat{G}_1(z) = \int_0^z \frac{du}{g(u)} = \frac{2 \tanh^{-1}\left(\frac{(1-\delta)\tan\frac{z}{2}}{\sqrt{1-\delta^2}} \right)}{\sqrt{1 - \delta^2}}$$

$$\hat{G}_3(z) = \int_0^z \frac{du}{g^3(u)} = \frac{(\delta^2 + 2) \tanh^{-1}\left(\frac{(1-\delta)\tan\frac{z}{2}}{\sqrt{1-\delta^2}} \right)}{(1 - \delta^2)^{5/2}}$$
$$+ \frac{\delta \sin(z)(\delta^2 - 3\delta\cos(z) - 4)}{2(1 - \delta^2)^2(\delta\cos(z) + 1)^2} \tag{D5}$$

We also introduce the definite integrals

$$\overline{G}_1 = \int_{-\pi}^{\pi} \frac{du}{g(u)} = \frac{2\pi}{\sqrt{1 - \delta^2}},$$

$$\overline{G}_3 = \int_{-\pi}^{\pi} \frac{dz}{g^3(z)} = \frac{\pi(\delta^2 + 2)}{(1 - \delta^2)^{5/2}}. \tag{D6}$$

Writing $[[\cdot]]$ for the nearest integer we get

$$G_p(z) = \int_0^z \frac{du}{g^p(u)} = \hat{G}_p(z) + \overline{G}_p \left[\left[\frac{z}{2\pi} \right]\right] \tag{D7}$$

FIG. 6. Comparison of our parameter estimates with those of Ref.[19]. The plots shows data points for all 9 combinations of biological replicate (3 per temperature) and stationary reference sample (3 in total, taken at temperature 17°C, 27°C and 37°C). **(a)**. Position-dependent speed $v(x) = \overline{v}\big(1 + \delta\cos(\omega x + \phi)\big)$. **(b)**. Time-dependent speed $v(\tau) = v_0\big(1 + \delta\cos(\omega\tau + \phi)\big)$.

FIG. 7. Time-dependent vs. position-dependent speed oscillations. **(a)**. Time-dependent speed oscillations $v(\tau)$ from figure 3c vs. the position-dependent speed oscillations from Bhat *et al.*[19]. **(b)**. Likelihoods L of the best-fitting time-dependent speed model vs. $L_{v(x)}$ of the best-fitting position-dependent model (computed as described in Appendix E).

and in terms of G_p we finally have

$$V_p(x) = \int_0^x \frac{du}{v^p(u)} = \frac{v_0}{\omega}\big(G_p(\omega x + \phi) - G_p(\phi)\big). \quad \text{(D8)}$$

By setting $v_1(x) = v(x)$, $v_2(x) = -v(L - x)$ and $D_1(t) = D_2(t) = D$ in Eq. (D3) and inserting the means and variances into Eq. (D2), we arrive at the replication program

$$f(x,\tau) = 1 - \frac{1}{4}\operatorname{erfc}\left(\frac{t - V_1(x)}{2^{\frac{3}{2}}DV_3(x)}\right)\operatorname{erfc}\left(\frac{t - V_1(L - x)}{2^{\frac{3}{2}}DV_3(L - x)}\right). \quad \text{(D9)}$$

Consistency of parameter estimates

Most parameter estimates for the position-dependent speed model obtained using the procedure outlined in Appendix E agree well with the values reported in Ref. [19], see Fig. 6a. Estimates that deviate also exhibit large uncertainties between replicates, see Table II. These large uncertainties can be explained in a similar way as for the time-dependent model presented in Section III.

Parameter estimates also agree well between time- and

position-modulated models, see Fig.6b. The oscillation frequency ω is not directly comparable between the two models, but when plotted against replication progress both models yield similar oscillations, see Fig. 7a. When the two models are compared across samples, we find that for a large majority of samples, the time-dependent speed model yields a slightly higher likelihood than the position-dependent speed model, see Fig. 7b.

Appendix E: Parameter estimation

We compute the position-wise bias-corrected abundances and their uncertainties as

$$a_i = \frac{n_i^{\text{exp}}}{n_i^{\text{stat}}}, \qquad \sigma_i = a_i \sqrt{\frac{1}{n_i^{\text{exp}}} + \frac{1}{n_i^{\text{stat}}}} \qquad \text{(E1)}$$

given the experimentally observed read counts at genomic locations $x_1, \ldots, x_n$ in an exponentially growing (n_i^{exp}) and stationary (n_i^{stat}) populations, see [19]. Since the counts n_i^{exp} and n_i^{stat} are typically large, we approximate the error distributions as Gaussian. Defining the parameter vector $\Theta = (v_0, D, \delta, \omega, \phi)$, we express the log likelihood by

$$\log L(\Theta) = -\frac{N}{2} \log(2\pi) - \sum_{i=1}^{N} \log(\sigma_i)$$
$$- \frac{1}{2} \sum_{i=1}^{N} \frac{\left(a_i - \lambda \mathcal{P}(x_i|\Theta)\right)^2}{\sigma_i^2}. \qquad \text{(E2)}$$

where $\lambda = \sum_i a_i / \sum_i \mathcal{P}(x_i|\Theta)$ is a scaling factor.

We evaluate $\mathcal{P}(x|\Theta)$ by a numerical integration of Eq.(3). This procedure requires a limited computational effort compared with Ref. [19], that evaluated the likelihood using stochastic simulations. We fit two different models corresponding to two different replication programs with are both instances of the general bacterial program from Eq. (7): (1) the time-dependent speed model in Eq. (9) with the modulating function $h(\tau)$ given in Eq.(13). (2) the position-dependent speed model of Ref. [19] with replication program Eq. (D9), see Appendix D (parameters are $\Theta = (\bar{v}, D, \delta, \omega, \phi)$ in this case, the parameter $\bar{v}$ takes the place of v_0).

The likelihood surfaces of these models are quite rough, in particular with respect to ω and ϕ. This causes combined gradient-ascent optimization of all five parameters to remain stuck in local maxima, unless the initial estimate is close to the global optimum. On the other hand, exhaustive exploration of the parameter space for all 5 parameters at sufficient resolution is computationally expensive. We thus used the following procedure:

1. Fit *constant-speed, no-diffusion* ($\delta = 0$, $D = 0$) regime by optimizing v_0 while keeping $D = 0$ and $\delta = 0$. Initial value is $v_0 = 500$bp/s.

2. Fit *oscillatory no-diffusion* ($\delta, \omega, \phi \geq 0$, $D = 0$) regime by optimizing v_0, δ, ω and ϕ while keeping $D = 0$. Initial values lie on a grid formed by the constant-speed, no-diffusion estimate of v_0, $\delta \in \{0, 1/2\}$, 25 logarithmically spaced values for ω from $\pi L/2$ to $32\pi L$ (L being the genome length), and 5 uniformly spaced values for ϕ from 0 to 2π.

3. Fit *constant-speed diffusive* ($\delta = 0$, $D \geq 0$) regime by optimizing v_0 and D while keeping $\delta = 0$. Initial values lie on a grid formed by the constant-speed, no-diffusion estimate of v_0, and 21 logarithmically spaced values for D from 10 to 10^{10}.

4. Find initial parameter for the *oscillatory diffusive* ($\delta, \omega, \phi \geq 0$, $D \geq 0$) regime by optimizing v_0, δ, ω and ϕ while fixing D to the *constant-speed diffusive* estimate. Initial values are the same as in (2), except that for v_0 we now use the *constant-speed diffusive* estimate.

5. Fit *oscillatory diffusive* regime by optimizing v_0, D, δ, ω and ϕ, using the estimates found in (1)-(4) as initial values.

6. Select the parameter regime with the lowest AIC score.

Identifiability

To study the identifiability of parameters v_0, D, δ, ω, ϕ of the bacterial replication program Eq. 9 with $h(t) = v_0\left(1 + \cos(\omega t + \phi)\right)$ we simulated abundance profiles and attempted to recover the parameter values using the estimation procedure outlined in Appendix E. We randomly selected 2000 sets of parameters (Fig. 8a), and computed their predicted abundance profiles by simulating 10^6 genomes using a fixed growth rate of $\Lambda = 1\text{h}^{-1}$). We then generated Poisson-distributed read counts n_i^{exp} (and also n_i^{stat}, using a flat abundance profile) from these abundance profiles, and ran the estimation procedure described in appendix E. All parameters except D were recovered well (Fig. 8b). Whether or not the diffusion was recovered correctly strongly depended on the value of the diffusion constant, specifically on the Peclet number $\text{Pe} = Lv_0/4D$ at the replication terminus (Fig. 8c). For Peclet numbers up to about 300, the estimation procedure typically correctly infers that $D \neq 0$ after AIC-based model selection (Fig. 8d). For larger Peclet numbers, the effect of diffusion is negligible (Fig. 2), and the AIC-based model selection step thus tends to prefer models with $D = 0$.

Appendix F: Eukaryotic DNA replication

We here derive Eq. (21) for the DNA abundance in the case of spatially discrete origins with time-homogeneous

FIG. 8. Parameter recovery. **(a)**. Distributions of true parameters θ. **(b)**. Estimation errors for simulated abundance profiles with $\Lambda = 1\,h^{-1}$ and 40k reads / 10kbp window on average. **(c)**. True value D vs. recovered value $\hat{D}$ for different ranges of the Peclet number $\mathrm{Pe} = Lv_0/4D$. **(d)**. Distribution of the relative error of $\hat{D}$ for different ranges of Pe.

firing rates as per Eq. (17). The general eukaryotic replication program [1, 35] is given by

$$f(x,\tau) = 1 - e^{-\int\int dx'\,d\tau'\,I(x',\tau')\theta\left(\tau-\tau'-\frac{|x-x'|}{v}\right)} \quad \text{(F1)}$$

which is equivalent to Eq. (15), but with an explicit expression for the light cone. We note that

$$\frac{d}{d\tau}\int\int dx'd\tau' I(x',\tau')\theta\left(\tau-\tau'-\frac{|x-x'|}{v}\right)$$
$$= \int dx' I\left(x',\tau-\frac{|x-x'|}{v}\right). \quad \text{(F2)}$$

We use Eq. (2) to obtain

$$\mathcal{P}(x) = \int_0^\infty dt \left[\int dx'' I\left(x'',\tau-\frac{|x-x''|}{v}\right)\right] \times$$
$$\left[e^{-\Lambda\tau-\int\int dx'dt' I(x',\tau')\theta\left(\tau-\tau'-\frac{|x-x'|}{v}\right)}\right]. \quad \text{(F3)}$$

Setting $I(x,\tau) = \sum_{j=1}^K I_j(\tau)\delta(x-x_j)$, i.e. assuming discrete origins $x_1,\ldots,x_K$, yields

$$\mathcal{P}(x) = \int_0^\infty d\tau \left[\sum_j I_j\left(\tau-\frac{|x-x_j|}{v}\right)\right] \times$$
$$\left[e^{-\Lambda\tau-\sum_j\int_0^{\tau-|x-x_j|/v} d\tau' I_j(\tau')}\right], \quad \text{(F4)}$$

and in the time homogeneous case $I_j(\tau) = I_j^*\theta(\tau)$ we find

$$\mathcal{P}(x) = 1-\Lambda\int_0^\infty dt\, e^{-\Lambda\tau-\sum_j I_j^*\left(\tau-\frac{|x-x_j|}{v}\right)\theta\left(\tau-\frac{|x-x_j|}{v}\right)}. \quad \text{(F5)}$$

The integral can be then done by breaking it into different time intervals in which a certain set of the theta functions are non-zero. This yields

$$\int_0^\infty dt\, e^{-\Lambda\tau-\sum_j I_j^*(\tau-|x-x_j|/v)\theta(\tau-|x-x_j|/v)}$$
$$= \frac{1-e^{-\Lambda\tau_1}}{\Lambda}$$
$$+ \sum_{k=1}^{K-1} \frac{e^{-\Lambda\tau_k-\sum_{j=1}^k I_j^*(\tau_k-\tau_j)} - e^{-\Lambda\tau_{k+1}-\sum_{j=1}^k I_j^*(\tau_{k+1}-\tau_j)}}{\Lambda+\sum_{j=1}^k I_j^*}$$
$$+ \frac{e^{-\Lambda\tau_K-\sum_{j=1}^K I_j^*(\tau_K-\tau_j)}}{\Lambda+\sum_{j=1}^K I_j^*} \quad \text{(F6)}$$

where $\tau_k = \frac{|x-x_k|}{v}$ are the times it takes a replisome formed at the k-th origin to reach position x, and are ordered for each x in such a way that $0 \le \tau_1 \le \cdots \le \tau_K$. Therefore, we have

$$\mathcal{P}(x) = e^{-\Lambda\tau_1}$$
$$-\Lambda\sum_{k=1}^{K-1} \frac{e^{-\Lambda\tau_k-\sum_{j=1}^k I_j^*(\tau_k-\tau_j)} - e^{-\Lambda t_{k+1}-\sum_{j=1}^k I_j^*(\tau_{k+1}-\tau_j)}}{\Lambda+\sum_{j=1}^k I_j^*}$$
$$-\Lambda\frac{e^{-\Lambda\tau_K-\sum_{j=1}^K I_j^*(\tau_K-\tau_j)}}{\Lambda+\sum_{j=1}^K I_j^*}. \quad \text{(F7)}$$

Equation (21) is obtained from Eq. (F7) by substituting T_k and W_k as defined in Eqs. 19 and 20 and grouping terms with the same exponent.

Appendix G: Eukaryotic origin inference by simulated annealing

We fit the solution of our eukaryotic model, Eq. (21), to experimentally measured DNA abundances in an exponentially growing asynchronous population of budding yest (*S. cerevisiae* W303)[36]. Also in this case, we correct for sequencing bias using the DNA abundance measured in a stationary population. Our input data $a_1, \ldots, a_N$ are thus the ratios of number of reads in the exponential vs. stationary population found within each 1kb window along the W303 genome (GenBank accession *CM007964.1*). To estimate number and positions of origins and their fork firing rates, we use simulated annealing [43].

Our free parameters are the number of origins K; their positions x_j, $j = 1 \ldots K$; their intensities (i.e. ratio of each fork firing rates to speed) I_j^*/v_0; and the ratio of growth rate to speed Λ/v_0. We collectively refer to the vector of $2K + 1$ free parameters (excluding K itself) as Θ. To prevent over fitting, we use the Akaike information criterion (AIC) as the cost function, defined as

$$\text{AIC}(\Theta) = 2(2K + 1) - 2 \log \text{L}(\Theta). \quad \text{(G1)}$$

The first term on the right hand side of Eq. (G1) penalizes a large number of origins, while the second term penalizes small log-likelihoods. The quantity $\log L(\Theta)$ is computed as in Eq. (E2). Since the data is over-dispersed compared to a Poissonian distribution, we use the empirical estimate $\sigma_i = \sigma = \sqrt{\frac{1}{N-1} \sum_i (a_{i+1} - a_i)^2 / 2} \approx 3.3 \cdot 10^{-9}$ instead of the Poissonian error estimates from Eq. (E1). The validity of this estimate is confirmed post-hoc by comparison with the same estimate computed with a_i replaced by the fitted model predictions. The resulting value $2 \cdot 10^{-10}$ is an order of magnitude smaller; σ therefore indeed mainly represents measurement errors and not expected differences between a_{i+1} and a_i. Our implementation of simulated annealing comprises the following steps:

1. Set the temperature to $T = 200$ and generate an initial parameter vector Θ. We draw Λ/v_0 uniformly from $[7.7 \times 10^{-8}, 2.3 \times 10^{-5}]$ bp^{-1} and place two origins randomly on each chromosomes with intensities $I_j^\star/v_0$ drawn uniformly from $[1.1 \times 10^{-7}, 1.6 \times 10^{-3}]$ bp^{-1}.

2. Generated a parameter proposal Θ' by executing one of the following moves with the indicated probability:

   - *Relocating an origin (prob. 24%)*. We choose one among the K origins randomly and relocate it to a uniformly drawn position on the same chromosome. The origin firing rate remains unchanged.
   - *Altering one origin intensity (prob. 24%)*. We randomly choose one of the origins and replace its intensity rate $I_j^\star/v_0$ with a value drawn uniformly from $[1.1 \times 10^{-7}, 1.6 \times 10^{-3}]$ bp^{-1}.
   - *Removing an origin (prob. 24%)*. We choose one among the K origins and remove it unless it is the only origin on the chromosome.
   - *Adding an origin (prob. 24%)*. We add an origin at a uniformly drawn location on the genome with an intensity uniformly drawn from $[1.1 \times 10^{-7}, 1.6 \times 10^{-3}]$ bp^{-1}.
   - *Altering the growth rate (prob. 4%)*. We draw a new value of Λ/v uniformly from $[7.7 \times 10^{-8}, 2.3 \times 10^{-5}]$ bp^{-1}.

   The ranges of free parameters are chosen to cover replisome speeds (v) from 10bps^{-1} to 100bps^{-1}, doubling times $\ln 2/\Lambda$ between 50 minutes to 150 minutes, and fork firing delays $1/I_j^*$ between 1 minute and 150 minutes.

3. Accept Θ' with Metropolis probability

$$q = \min \left\{ 1, \ \exp \left(\frac{\text{AIC}(\Theta) - \text{AIC}(\Theta')}{T} \right) \right\} \quad \text{(G2)}$$

   where T is the current temperature.

4. Reduce the temperature by a factor $1 - 1.336 \cdot 10^{-5}$ and reiterate from step (2). The total number of iterations is $2 \cdot 10^6$. The factor is chosen such that the final temperature is equal to $5 \cdot 10^{-10}$.

After half the number of iterations (10^6), the number of origins is already close to the final value (222 vs. the final number of 234), and repeating the procedure 10 times yield similar final numbers of origins (225 ± 6). We take this as evidence that the algorithm has sufficiently converged after $2 \cdot 10^6$ iterations, and that the solutions at that point are close to the global optimum.

[1] J. Bechhoefer and N. Rhind, Replication timing and its emergence from stochastic processes, Trends in Genetics **28**, 374 (2012).

[2] A. Baker, B. Audit, S.-H. Yang, J. Bechhoefer, A. Arneodo, *et al.*, Inferring where and when replication initiates from genome-wide replication timing data, Physical review letters **108**, 268101 (2012).

[3] A. Gispan, M. Carmi, and N. Barkai, Model-based analysis of DNA replication profiles: predicting replication fork velocity and initiation rate by profiling free-cycling cells, Genome Research **27**, 310 (2017).

[4] M. L. Hulke, D. J. Massey, and A. Koren, Genomic meth-

ods for measuring DNA replication dynamics, Chromosome Research **28**, 49 (2020).

[5] Z.-Q. Xu and N. E. Dixon, Bacterial replisomes, Current opinion in structural biology **53**, 159 (2018).

[6] D. M. Gilbert, Making sense of eukaryotic DNA replication origins, Science **294**, 96 (2001).

[7] A. Costa and J. F. Diffley, The initiation of eukaryotic DNA replication, Annual Review of Biochemistry **91**, 107 (2022).

[8] S. C.-H. Yang, N. Rhind, and J. Bechhoefer, Modeling genome-wide replication kinetics reveals a mechanism for regulation of replication timing, Molecular systems biology **6**, 404 (2010).

[9] N. Rhind, DNA replication timing: Biochemical mechanisms and biological significance, BioEssays **44**, 2200097 (2022).

[10] S. Cooper and C. E. Helmstetter, Chromosome replication and the division cycle of Escherichia coli, Journal of Molecular Biology **31**, 519 (1968).

[11] M. K. Raghuraman, E. A. Winzeler, D. Collingwood, S. Hunt, L. Wodicka, A. Conway, D. J. Lockhart, R. W. Davis, B. J. Brewer, and W. L. Fangman, Replication dynamics of the yeast genome, science **294**, 115 (2001).

[12] Y. Daigaku, A. Keszthelyi, C. A. Müller, I. Miyabe, T. Brooks, R. Retkute, M. Hubank, C. A. Nieduszynski, and A. M. Carr, A global profile of replicative polymerase usage, Nature structural & molecular biology **22**, 192 (2015).

[13] N. Sueoka and H. Yoshikawa, The chromosome of Bacillus subtilis. I. Theory of marker frequency analysis, Genetics **52**, 747 (1965).

[14] B. M. Wendel, C. T. Courcelle, and J. Courcelle, Completion of DNA replication in Escherichia coli, Proceedings of the National Academy of Sciences **111**, 16454 (2014).

[15] B. M. Wendel, J. M. Cole, C. T. Courcelle, and J. Courcelle, SbcC-SbcD and ExoI process convergent forks to complete chromosome replication, Proceedings of the National Academy of Sciences **115**, 349 (2018).

[16] S. L. Midgley-Smith, J. U. Dimude, T. Taylor, N. M. Forrester, A. L. Upton, R. G. Lloyd, and C. J. Rudolph, Chromosomal over-replication in Escherichia coli recG cells is triggered by replication fork fusion and amplified if replichore symmetry is disturbed, Nucleic acids research **46**, 7701 (2018).

[17] J. U. Dimude, M. Stein, E. E. Andrzejewska, M. S. Khalifa, A. Gajdosova, R. Retkute, O. Skovgaard, and C. J. Rudolph, Origins left, right, and centre: increasing the number of initiation sites in the Escherichia coli chromosome, Genes **9**, 376 (2018).

[18] N. Rhind, S. C.-H. Yang, and J. Bechhoefer, Reconciling stochastic origin firing with defined replication timing, Chromosome Research **18**, 35 (2010).

[19] D. Bhat, S. Hauf, C. Plessy, Y. Yokobayashi, and S. Pigolotti, Speed variations of bacterial replisomes, Elife **11**, e75884 (2022).

[20] D. Huang, A. E. Johnson, B. S. Sim, T. Lo, H. Merrikh, and P. A. Wiggins, The high-resolution in vivo measurement of replication fork velocity and pausing, Nature Communications **14**, 1762 (2022).

[21] D. Huang, T. Lo, H. Merrikh, and P. A. Wiggins, Characterizing stochastic cell-cycle dynamics in exponential growth, Physical Review E **105**, 014420 (2022).

[22] E. Powell, Growth rate and generation time of bacteria, with special reference to continuous culture, Microbiol-

ogy **15**, 492 (1956).

[23] H. Yoshikawa and N. Sueoka, Sequential replication of Bacillus subtilis chromosome, I. Comparison of marker frequencies in exponential and stationary growth phases, Proceedings of the National Academy of Sciences **49**, 559 (1963).

[24] M. Chandler and R. Pritchard, The effect of gene concentration and relative gene dosage on gene output in Escherichia coli, Molecular and General Genetics MGG **138**, 127 (1975).

[25] F. Jafarpour, C. S. Wright, H. Gudjonson, J. Riebling, E. Dawson, K. Lo, A. Fiebig, S. Crosson, A. R. Dinner, and S. Iyer-Biswas, Bridging the timescales of single-cell and population dynamics, Physical Review X **8**, 021007 (2018).

[26] J. A. Morin, F. J. Cao, J. M. Lázaro, J. R. Arias-Gonzalez, J. M. Valpuesta, J. L. Carrascosa, M. Salas, and B. Ibarra, Active DNA unwinding dynamics during processive DNA replication, Proceedings of the National Academy of Sciences **109**, 8115 (2012).

[27] J. A. Morin, F. J. Cao, J. M. Lázaro, J. R. Arias-Gonzalez, J. M. Valpuesta, J. L. Carrascosa, M. Salas, and B. Ibarra, Mechano-chemical kinetics of DNA replication: identification of the translocation step of a replicative DNA polymerase, Nucleic acids research **43**, 3643 (2015).

[28] S. Jun, H. Zhang, and J. Bechhoefer, Nucleation and growth in one dimension. I. The generalized Kolmogorov-Johnson-Mehl-Avrami model, Physical Review E **71**, 011908 (2005).

[29] S. Jun and J. Bechhoefer, Nucleation and growth in one dimension. II. Application to DNA replication kinetics, Physical Review E **71**, 011909 (2005).

[30] A. Kolmogorov, Statistical theory of nucleation processes., Izu. Akad. Nauk SSSR **3**, 355 (1937).

[31] J. William and R. Mehl, Reaction kinetics in processes of nucleation and growth, Trans. Metall. Soc. AIME **135**, 416 (1939).

[32] M. Avrami, Kinetics of phase change. I General theory, The Journal of chemical physics **7**, 1103 (1939).

[33] K. Sekimoto, Evolution of the domain structure during the nucleation-and-growth process with non-conserved order parameter, Physica A: Statistical Mechanics and its Applications **135**, 328 (1986).

[34] K. Sekimoto, Evolution of the domain structure during the nucleation-and-growth process with non-conserved order parameter, International Journal of Modern Physics B **5**, 1843 (1991).

[35] A. Baker and J. Bechhoefer, Inferring the spatiotemporal DNA replication program from noisy data, Physical Review E **89**, 032703 (2014).

[36] C. A. Müller, M. Hawkins, R. Retkute, S. Malla, R. Wilson, M. J. Blythe, R. Nakato, M. Komata, K. Shirahige, A. P. de Moura, and C. A. Nieduszynski, The dynamics of genome replication using deep sequencing, Nucleic Acids Research **42**, e3 (2013).

[37] A. Shumate and S. L. Salzberg, Liftoff: accurate mapping of gene annotations, Bioinformatics **37**, 1639 (2021).

[38] B. Boldin, O. Diekmann, and J. Metz, Population growth in discrete time: a renewal equation oriented survey, arXiv preprint arXiv:2306.09193 (2023).

[39] A. K. Sinha, C. Possoz, A. Durand, J.-M. Desfontaines, F.-X. Barre, D. R. Leach, and B. Michel, Broken replication forks trigger heritable DNA breaks in the terminus

of a circular chromosome, PLoS genetics **14**, e1007256 (2018).

[40] M. R. Evans and S. N. Majumdar, Diffusion with stochastic resetting, Physical review letters **106**, 160601 (2011).

[41] M. R. Evans, S. N. Majumdar, and G. Schehr, Stochastic resetting and applications, Journal of Physics A: Mathematical and Theoretical **53**, 193001 (2020).

[42] M. M. Elshenawy, S. Jergic, Z.-Q. Xu, M. A. Sobhy, M. Takahashi, A. J. Oakley, N. E. Dixon, and S. M. Hamdan, Replisome speed determines the efficiency of the Tus- Ter replication termination barrier, Nature **525**, 394 (2015).

[43] D. Bertsimas and J. Tsitsiklis, Simulated Annealing, Statistical Science **8**, 10 (1993).