1 **Title (160 characters including spaces):** Lymphatic muscle cells are the innate pacemaker cells
2 regulating mouse lymphatic collecting vessel contractions
3

4 **Authors:** S.D. Zawieja[1], G.A. Pea[1], S.E. Broyhill[1], K.H. Bromert[1], C. E. Norton[1], H. J. Kim[1], M. Li[1],
5 J.A. Castorena-Gonzalez [2], B.T. Drumm[3], and M.J. Davis[1]
6
7 [1]Dept. of Medical Pharmacology & Physiology, University of Missouri, Columbia, Missouri;
8 [2]Dept. of Pharmacology, Tulane University, New Orleans, Louisiana
9 [3]Smooth Muscle Research Centre, Dundalk Institute of Technology, Dundalk, Co. Louth, A91
10 K584, Ireland.
11
12
13
Correspondence:
Michael J. Davis, PhD
Department of Medical Pharmacology and Physiology
One Hospital Drive
MA415 Medical Sciences Building
University of Missouri
Columbia, MO 65212
Tel: 573-884-5181
Fax: 573-884-4276
davismj@health.missouri.edu
24
25
Co-Correspondence:
Scott Zawieja PhD
Department of Medical Pharmacology and Physiology
One Hospital Drive
MA415 Medical Sciences Building
University of Missouri
Columbia, MO 65212
Tel: 573-882-8069
Fax: 573-884-4276
zawiejas@health.missouri.edu
36
37

38 **Running title: Lymphatic Muscle as Innate Pacemaker Cells**
39
Impact: The presence and functionality of proposed pacemaker cells in collecting lymphatic
vessels was tested with various mouse Cre models to drive expression of a recombination
reporter ROSA26mT/mG, a genetically encoded Ca^{2+} sensor GCaMP6f, or the optogenetic tool
channel-rhodopsin2. Lymphatic CD34[+] adventitial cells co-express PDGFRA[+] while cKit[+] cells are
mast cells; and neither cell type demonstrated pacemaking functionality. *Myh11CreER[T2]*

identified lymphatic muscle cells which exhibited pacemaker behaviors such as pressure-
dependent calcium events during diastole and propagated contraction induced by optical
stimulation of channel-rhodopsin2.
48
49 **Abbreviations**: Ano1, Anoctamin 1; ChR2, channel rhodopsin; cLV, collecting lymphatic vessel;
50 IALV, Inguinal Axillary Lymphatic Collecting Vessel; ICC, Interstitial Cell of Cajal; ICLC, Interstitial
51 Cell of Cajal Like Cell; LEC, Lymphatic Endothelial Cell; LMC, Lymphatic Muscle Cell; STDs,
52 Spontaneous transient depolarizations; STMs, Spatio-Temporal Maps
53
54 **Keywords:** Lymphatic collecting vessel, lymphatic muscle cell, pacemaking, interstitial cells of
55 Cajal like cells, mesenchymal stem cells
56
57
Abstract

Collecting lymphatic vessels (cLVs) exhibit spontaneous contractions with a pressure-dependent
frequency, but the identity of the lymphatic pacemaker cell is still debated. By analogy to
pacemakers in the GI and lower urinary tracts, proposed cLV pacemaker cells include interstitial
cells of Cajal like cells (ICLC), pericytes, as well as the lymphatic muscle (LMCs) cells themselves.
Here we tested the extent to which these cell types are invested into the mouse cLV wall and if
any cell type exhibited morphological and functional processes characteristic of pacemaker
cells: a contiguous network; spontaneous Ca^{2+} transients; and depolarization-induced
propagated contractions. We employed inducible Cre (iCre) mouse models routinely used to
target these specific cell populations including: c-kitCreER^{T2} to target ICLC; $PdgfrβCreER^{T2}$ to
target pericytes; $PdgfrαCreER^{TM}$ to target CD34$^+$ adventitial fibroblast-like cells or ICLC; and
$Myh11CreER^{T2}$ to target LMCs. These specific inducible Cre lines were crossed to the fluorescent
reporter ROSA26mT/mG, the genetically encoded Ca^{2+} sensor GCaMP6f, and the light-activated
cation channel rhodopsin2 (ChR2). c-KitCreER^{T2} labeled both a sparse population of LECs and
round adventitial cells that responded to the mast cell activator compound 48-80.
$PdgfrβCreER^{T2}$ drove recombination in both adventitial cells and LMCs, limiting its power to
discriminate a pericyte specific population. $PdgfrαCreER^{TM}$ labeled a large population of
interconnected, oak leaf-shaped cells primarily along the adventitial surface of the vessel.
Titrated induction of the smooth muscle-specific $Myh11CreER^{T2}$ revealed a LMC population with
heterogeneous morphology. Only LMCs consistently, but heterogeneously, displayed
spontaneous Ca^{2+} events during the diastolic period of the contraction cycle, and whose
frequency was modulated in a pressure-dependent manner. Optogenetic depolarization
through the expression of ChR2 by $Myh11CreER^{T2}$, but not $PdgfrαCreER^{TM}$ or c-KitCreER^{T2},
resulted in a propagated contraction. These findings support the conclusion that LMCs, or a
subset of LMCs, are responsible for mouse cLV pacemaking.

83
Introduction
The spontaneous contractions of collecting lymphatic vessels (cLV) are an integral
component to fluid and macromolecule homeostasis as they provide the force to transport fluid
from the interstitium back to the blood circulation (Scallan et al., 2016). In humans,
spontaneous contractile activity is estimated to account for 2/3 of lymph transport (Engeset et

al., 1977) and this function is significantly compromised in patients suffering from lymphedema,
whose cLVs typically display weak and irregular or entirely absent contractile activity
(Olszewski, 2002). *Ex vivo* studies, in which the intraluminal pressure can be precisely
controlled, have refined our understanding of the pressure-dependent regulation of
contraction frequency (Benoit et al., 1989; Gashev et al., 2004), with some mouse cLVs
displaying a 10-fold increase in contraction frequency over a 10 cmH$_2$O pressure gradient
(Scallan and Davis, 2013; Zawieja et al., 2018). The observation that cLVs, often cannulated at
various lengths for *ex vivo* preparations, retain a consistently tunable contraction frequency
points to the presence of a pacemaker cell(s) innate to the structure of the cLV wall and with a
seemingly ubiquitous presence along the length of the vessel (Zawieja et al., 1993; Castorena-
Gonzalez et al., 2018b).
Investigations into the cLV pacemaker identity have focused largely on cells termed
interstitial cells of Cajal like cells (ICLC; or telocytes) (McCloskey et al., 2002; Briggs Boedtkjer et
al., 2013), as they display some morphological and cell marker expression profiles similar to the
interstitial cells of Cajal (ICC), which are bona fide pacemakers in the gastrointestinal (GI) tract.
ICC are classically identified by either methylene blue staining and expression of cKit, and
coordinate GI smooth muscle contraction (Maeda et al., 1992; Ward et al., 1994; Ordog et al.,
1999). ICC also express the canonical Ca^{2+} activated chloride channel Anoctamin 1 (Ano1)
(Gomez-Pinilla et al., 2009), which is required for the electrical slow wave initiation and
formation that subsequently activates an action potential in GI smooth muscle (Hwang et al.,
2009; Zhu et al., 2009; Singh et al., 2014). Previous reports in sheep mesenteric lymphatic
vessels identified a population of cKit$^+$, vimentin$^+$, ICLC in the vessel wall between the
endothelial and LMC layer and running along the axis of the vessel (McCloskey et al., 2002).
Investigations in the human thoracic duct also identified a significant population of ICLC in close
proximity to the lymphatic muscle cells (LMCs) evident by methylene blue staining,
immunostaining for CD34, vimentin, and cKit, as well as the gold standard of electron
microscopy (Briggs Boedtkjer et al., 2013). However, neither study could determine if these
cells had functional electrical communication with the LMCs or demonstrate either a
membrane electrical clock or internal Ca^{2+} clock to drive the rhythmic lymphatic vessel
contractions observed *ex vivo*. LMCs share a functional similarity to ICC in that they also display
a Ca^{2+} activated chloride current (Van Helden, 1993; Toland et al., 2000; Mohanakumar et al.,
2018), mediated by Ano1 (Zawieja et al., 2019), that regulates pacemaking by modulating the
slope of diastolic depolarization. Spontaneous transient depolarizations, presumably Ano1
dependent, were recorded in mesenteric cLVs from guinea pigs (Van Helden, 1993; von der
Weid et al., 2008) providing a mechanism for membrane potential instability. Furthermore,
computational models have proposed LMC sarcoplasmic reticulum (SR) Ca^{2+} release as the
oscillator mechanism driving pacemaking (Imtiaz et al., 2007). SR Ca^{2+} release has also been
implicated in pericyte regulation of arterioles (Hashitani et al., 2015; van Helden and Imtiaz,
2019), in microvascular vasomotion (Boedtkjer et al., 2008; Aalkjaer et al., 2011; van Helden
and Imtiaz, 2019), and in the contraction waves of atypical muscle cells of the lower urinary
tract (Grainger et al., 2022).
130
Presently, no investigations have clearly identified the cellular identity of possible
pacemaker cells within the cLVs of the mouse. Murine cLVs exhibit contractile parameters and

conduction speed similar to those of human vessels (Castorena-Gonzalez et al., 2018b) and
their simplified architecture, compared to larger mammals, in combination with the genetic
tools developed for the mouse model, allowed us to test for a fundamental pacemaker cell in
the cLV. In this study we utilized multiple genetic mouse models, immunofluorescence imaging,
Ca^{2+} imaging, and optogenetic light-activated depolarization to both visualize and test the
functional aspects of putative pacemaker cells. We did not observe a significant or contiguous
cKit$^+$ cell population along the vessels. *PdgfraCreER^{TM}* was able to consistently delineate a
significant population of interconnected adventitial cells that were also CD34$^+$. However, an
absence of Ca^{2+} events in phase with the contractile activity and an inability to consistently
elicit a contraction upon photo-stimulated depolarization provides functional evidence against
a role for either PDGFRα$^+$ or cKit$^+$ adventitial cells as the pacemaker cell. In contrast, MYH11$^+$
LMCs exhibited diastolic Ca^{2+} events that, while asynchronous, were dynamically modulated by
pressure and could elicit a propagated contraction after optogenetic stimulation of LMCs
expressing ChR2.
147
Results
Methylene Blue Staining Reveals adventitial Cells in Murine cLVs
Methylene blue staining was used to identify an ICLC population in the human lymphatic
thoracic duct (Briggs Boedtkjer et al., 2013). In our isolated and cleaned IALVs ,methylene blue
stained a significant number of cells along the length of the vessel. The methylene blue$^+$ cells
had variable density along the length of the vessel and stained cells exhibited different patterns
of morphology and location (Figure 1A-C). A significant portion of the stained cells resembled
lymphatic vessel-associated macrophages with an elongated spindle shape, while other cells
were smaller and circular (Figure 1D-F). Methylene blue also appeared to stain mast cells as
there were large ovoid cells on the adventitia of the vessel with intracellular granules. In
addition, methylene blue stained a minor population of cells that exhibited long and thin axon-
like extensions which appeared to have a slight helical orientation, with a small central body
and nucleus (Figure 1C). None of these cell populations were aligned with the longitudinal axis
of the vessel that would permit efficient coupling or regulation across the circumferential LMCs
required for coordinated propagation along the length of the vessel.
163
Immunofluorescence Imaging of IALVs stained for ICLC, LEC, and LMC Markers
We first stained IALVs for the putative telocyte/ICLC markers cKit, CD34, and the intermediate
filament vimentin, which have been previously described in human and sheep lymphatic tissues
(McCloskey et al., 2002; Briggs Boedtkjer et al., 2013). Lymphatic muscle cells also express the
intermediate filament desmin (McCloskey et al., 2002). IALVs stained for cKit (Figure 2B) had
robust signal in large ovoid cells characteristic of mast cells, confirmed to contain a nucleus via
DAPI staining (Figure 2A), that were located in the outer part of the adventitia. Similarly, cKit
stained populations of elongated cells as well as circular cells with variable density throughout
the IALV wall, similar to methylene blue$^+$ cell populations (Figure 2B, J). Staining for CD34
revealed a striking population of cells that were seemingly contiguous along the length of the
vessel. These CD34$^+$ cells generally had multiple lobular processes and a "oak leaf" like
appearance, characteristic of fibroblasts, while some contained short, thin dendrite-like
extensions (Figure 2C, G, K). CD34$^+$ cells were negative for desmin (Figure 2H), which stained

the circumferential LMCs (Figure 2F; note that the largely non-circumferential cell organization
in this region is typical for a lymphatic endothelial valve site). Furthermore, the CD34[+] cells and
cKit[+] stained separate populations (Figure 2D, L). Vimentin stained lymphatic endothelial cells
(LECs) which exhibited a horizontal cobblestone morphology in parallel with the vessel axis
(Figure 2E, I), while also co-labeling the majority of the CD34[+] cells (Figure 2H) and cKit[+] cells
(Figure 2L). Videos of the half vessel z-stacks are provided (Supplemental Movies 1-3 for Figure
2D, H, and L respectively).
Of the cells stained in Figure 2, the CD34[+] population was intriguing due to its high
density and seemingly consistent presence through the length of the IALV, which potentially
would be conducive to effective regulation of LMC excitability. In addition to CD34 and
vimentin, PDGFRα expression is also commonly ascribed to both telocytes (Vannucchi et al.,
2013; Xiao et al., 2013; Zhou et al., 2015) as well as fibroblasts (Kimura et al., 2021; Clayton et
al., 2022). We performed immunofluorescence imaging for PGDFRα counterstained with CD34
and markers for LMCs, LECs, or myeloid-derived cells. As noted with desmin staining in Figure 2,
CD34[+] cells (Figure 3A) did not co-label LMCs (Figure 3D) that were smooth muscle actin[+]
(Figure 3B) and calponin[+] (Figure 3C). However, nearly all CD34[+] (Figure 3E) cells were also
PDGFRα[+] (Figure 3G, H) but not SMA+ (Figure 3G, H). Staining to outline LECs with CD31
(PECAM, Figure 3I) revealed the expected rectangular cobblestone morphology that was
distinct from the PDGFRα labeled cells (Figure 3J, L). Calponin specifically stained LMCs (Figure
3K) but not PDGFRα+ cells (Figure 3L). To determine if the CD34[+]PDGFRα[+] cells had
hematopoietic or mononuclear phagocytic origin, we stained IALVs from Macgreen mice (Csfr1-
EGFP) with CD45 and PDGFRα. Anti-GFP (Figure 3M) labeled the cLV myeloid cells (Macgreen),
which consists of predominantly macrophages and dendritic cells, that were co-stained by
CD45[+] (Figure 3O, P). CD45[+] but GFP- negative circular cells were also observed and are likely T-
Cells, B-cells, or mast cells. As expected, the PDGFRα[+] cell population (Figure 3N) was not CD45[+]
(Figure 3P). Lastly, we stained for PDGFRα, CD34, and PDGFRβ, a commonly used pericyte
marker (Figure 3 Q-T). As above, CD34 and PDGFRα were highly colocalized (Figure 3Q, R, T),
and many of the CD34[+] and PDGFRα[+] cells were also PDGFRβ[+] (Figure 3T). PDGFRβ also stained
some circumferential LMCs, albeit with a weaker signal than the adventitial cells (Figure 3U).
During the imaging of mouse IALVs for these markers, we also observed that the lymphatic
secondary endothelial valves were populated by elongated cells that stretched the length of the
valve leaflet and were positive for CD34, PDGFRα, and PDGFRβ, with varying intensities, and
found within both leaflets of the valve (Figure 3V,W) as determined from max projections of a
z-stack encompassing only the "z slices" within the lumen of the IALV. These cells had long, thin
extensions that were branched along with apparent dendrite extensions with a morphology
that closely resembled those described of pericytes or telocytes (Popescu and Faussone-
Pellegrini, 2010) (Figure 3V,W). PDGFRα[+] or CD34[+] cells with this morphology were only
observed in the valve leaflets and thus seemed insufficient to regulate pacemaking as normal
contractions are observed in cLVs without secondary valves (not shown). The z-stacks
demonstrating these valve-located "telocyte" shaped cells (Figure 3V,W) are provided as
Supplemental Movies 4 and 5.
The vast majority of the PDGFRα[+] cells were located in the adventitial layer Figure 4A-D,
which varied between 2-3 PDGFRα[+] cells thick (Figure 4E) for this particular mouse lymphatic
collecting vessel. Under this layer, we observed only a single layer of largely circumferential

LMCs stained by MYH11 (Figure 4B) sitting atop a single layer of CD31+ LECs (Figure 4A). We
also observed occasional PDGFRα+ cells or their extensions located in the sub-endothelial space
(Figure 4 E', E") positioned between the LECs and the LMCs.
We next determined the degree of colocalization between the CD34 and PDGFRα signal
given the significant overlap. Colocalization analysis of PDGFRα (Figure 5A) and CD34 (Figure
5B) and their colocalization (Figure 5C) was determined with the FIJI BIOP-JACoP tool. The
Pearson's coefficient was 0.83 (Figure 5 D) and Mander's coefficient of overlap 0.80 was for the
PDGFRα+ signal and 0.87 for the CD34 signal (Figure 5E). This high degree of colocalization CD34
and PDGFRα signal informed our use of the commercially available *PdgfrαCreTM* mouse model
to target these cells.
231
Use of iCre-Mediated Recombination of Rosa26mT/mG to delineate and characterize specific
IALV cell types
After confirming the presence of Vimentin+, cKit+, and CD34+ PDGFRα+ positive cells within the
mouse IALV, we sought to further investigate these cell populations by using constitutive and
inducible Cre recombinase expressing mouse lines. IALVs from the constitutively active
PdgfrαCre-ROSA26mTmG and *Ng2Cre*-ROSA26mTmG mice had GFP fluorescence in the
majority of LMCs as well as in the fibroblast-shaped cells found within the IALV wall (Figure 6
A,B). While informative of expression of the LMC progenitor cells, neither constitutive Cre
would be useful in delineating cell types. In contrast to the constitutively active *PdgfrαCre*, the
tamoxifen inducible *PdgfrαCreTM* line drove significant recombination in only the fibroblast-
shaped cells previously stained with CD34 and PDGFRα but not LMCs or LECs (Figure 6C).
PdgfrβCreT2, commonly used to label pericytes, drove recombination in both a minor
population of the LMCs and the fibroblast-shaped cells. *cKitCreT2*, which capably drives
recombination in the ICC of the GI (Baker et al., 2016), drove recombination only in a small
population of irregularly-spaced large ovoid cells on the surface of the IALV (Figure 6E), largely
corresponding with mast cell labeling, although recombination in 1 or 2 LECs could occasionally
be detected (not shown). Finally, *Myh11CreT2* drove recombination in nearly all LMCs which
were largely circumferentially oriented with dendrite-like, cell-cell contacts visible between
them and without significant GFP fluorescence in either LECs or the fibroblast-shaped CD34+
PDGFRα+ cell population (Figure 6F). Additionally, some LMCs maintained the bipolar shape but
had multiple extensions forming a "Y" shape in which an adjacent LMC typically filled the inner
void. A very minor population of recombined cells in the *Myh11CreT2*-ROSA26mTmG IALVs
were smaller and irregularly patterned with multiple fine axon-like projections or highly-
dendritic ruffled edges (Figure 6F).
To complement the morphological and cell density findings obtained with confocal
microscopy, we digested IALVs from the iCre-ROSA26mTmG lines, and the *Prox1-eGFP* line as a
control, into single cell suspensions and sorted the respective GFP+ populations (Figure 6G-J) for
RT-PCR profiling (Figure 6K). We first focused on determining the molecular fidelity of the
sorted cells based on the gene promoters used to drive each "iCre" model to discern cellular
overlap. In agreement with the confocal images, sorted GFP+ cells from *PdgfrβCreT2*-
ROSA26mT/mG IALVs expressed PDGFRβ but also MYH11 and PDGFRα. In contrast, GFP-sorted
cells from *PdgfrαCreTM* IALVs expressed PDGFRα and PDGFRβ, but with no detectable
expression of MYH11. Conversely, GFP+ cells from sorted *Myh11CreT2*-ROSA26mTmG IALVs

had high expression for MYH11, as well as PDGFRβ, but did not express PDGFRα. IALVs from
cKitCreER^{T2}-ROSA26mTmG mice were not pursued for FACS due to the exceptionally sparse
recombination along the IALV.
We further profiled each population of sorted cells with RT-PCR for other common
markers for endothelial cells, muscle cells, and pericytes. Similar to the Prox1 results (Figure
6K), endothelial nitric oxide synthase (eNOS) expression was observed only in the *Prox1-eGFP*
sorted cells, which also expressed vimentin and MCAM and had weak but detectable signal for
CD34 (Figure 7A). *Myh11CreER^{T2}* sorted cells showed expression of smooth muscle actin
(Acta2), the alpha subunit of the L-type voltage gated Ca^{2+} channel Cav1.2, desmin, MCAM, and
vimentin (Figure 7B). In addition to the genes expressed under *Myh11CreER^{T2}* recombination,
Cdh5, CD34, and Cspg4 (Ng2) were detected in cells sorted in *PdgfrβCreER^{T2}* IALVs (Figure 7C).
As expected, the GFP$^+$ cells sorted from *PdgfrαCreER^{TM}* IALVs expressed mRNA for CD34, Cspg4,
and vimentin, but not desmin, smooth muscle actin, nor the pericyte marker MCAM (Figure
7D). The alpha subunit of the voltage gated Ca^{2+} channel was positive in cells sorted from both
PDGFRα, PDGFRβ,and *Myh11CreER^{T2}* IALVs, but was also observed in the *PdgfrαCreER^{TM}* IALVs
without any evidence that MYH11 expressing muscle cells contaminated the latter. These
findings confirmed the separate cell populations achieved with *PdgfrαCre^{TM}* and *Myh11CreER^{T2}*
mediated recombination, at least as it pertains to ROSA26mTmG. We followed up the
identification of Cav1.2 expression in the *PdgfrαCreER^{TM}* sorted cell population by assessing the
expression of other genes involved in either pacemaking (Ano1) or electrical conduction (Cx45)
of IALVs (Figure 7E). Intriguingly, expression of both Ano1 and Cx45 was observed in the
PdgfrαCreER^{TM} sorted cells, which were further confirmed to lack either endothelial or
hematopoietic contamination as we did not detect expression of the endothelial marker CD31
or the hematopoietic marker CD45.
289
Inducible Deletion of Either Cav1.2, Ano1, and Cx45 with *PdgfrαCreER^{TM}* Did Not Affect cLV
Pacemaking
The expression of the genes critically involved in cLV function—Cav1.2, Ano1, and
Cx45—n the *PdgfrαCreER^{TM}*-ROSA26mTmG purified cells prompted us to generate
PdgfrαCreER^{TM}-Ano1fl/fl, *PdgfrαCreER^{TM}*-Cx45fl/fl, and *PdgfrαCreER^{TM}*-Cav1.2fl/fl mice for
functional testing. We isolated popliteal cLVs and tested their pacemaker and contractile
activity in response to physiological pressures over the range 0.5-10 cmH$_2$O, under normal
conditions. However, we did not detect any significant difference in pacemaking nor contractile
function of popliteal cLVs studied from *PdgfrαCreER^{TM}*-Ano1fl/fl (Figure 8A-F) or
PdgfrαCreER^{TM}-Cx45fl/fl (Figure9A-F), compared to their respective Ano1fl/f and Cx45fl/fl
controls. *PdgfrαCreER^{TM}*-Cav1.2fl/fl mice had no statistically significant differences in
normalized contraction amplitude (Figure 10A), contraction frequency (Figure 10C), fractional
pump flow (Figure 10D), end diastolic diameter (Figure 10E). However, we noted a mild but
statistically significant increase in ejection fraction at the lowest pressure, 0.5 cmH$_2$O. Vessels
isolated from *PdgfrαCreER^{TM}*-Cav1.2fl/fl mice also had a statistically significant increase in
vessel tone (Figure 10F) noted at the 2-way ANOVA level although we did not resolve
significance at any specific pressure with this sample size. Nonetheless, despite the presence of
transcript for these critical genes, *PdgfrαCreER^{TM}* mediated deletion failed to recapitulate

previous reports of contractile defects using the *Myh11CreER^T2* line (Castorena-Gonzalez et al.,
2018b; Zawieja et al., 2019; To et al., 2020; Davis et al., 2022).
310
Potential Role for PDGFRα⁺ Adventitial Fibroblasts As Pool of Progenitor Cells
Despite the lack of cLV pacemaking deficits in the *PdgfrαCreER^TM* genetic knockout lines, we
were curious to discern further insight into the role or function of the PDGFRα⁺ CD34⁺ cells
which comprise a significant portion of the lymphatic cLV wall. The *PdgfrαCreER^TM* recombined
cells exhibited expression of Krüppel-like factor 4 (Klf4), stem cell antigen 1 (Sca1, also referred
to as Ly6a), Gli1, CD29, CD105, and CD44 (Figure 11A, B). To again validate that this signature
was unique to the *PdgfrαCreER^TM* recombined cells, we performed RT-PCR on FACS from Prox1-
eGFP, *Myh11CreER^T2*-ROSA26mTmG, and *PdgfrαCreER^TM*-ROSA26mTmG IALVs. Recombined
(GFP⁺) cells from *Myh11CreER^T2*-ROSA26mTmG had weak expression of Klf4 and Gli1 and were
negative for Sca1, CD34, and PDGFRα (Figure 11C). Similarly, LECs sorted from *Prox1-eGFP* IALVs
were positive for Klf4, weak for Sca1, and positive for CD34 but negative for GLi1 and PDGFRα
Gli1. In contrast, *PdgfrαCreER^TM*- ROSA26mTmG recombined cells were positive for all markers
as were the unrecombined population (tdTomato⁺) cells in the *Myh11CreER^T2*- ROSA26mTmG
IALVs (Figure 11D). We performed immunofluorescence staining for one of these multipotent
markers, Sca1 (Figure 11E, I) while counter staining for LMCs, with SMA or MYH11 (Figure 11F,
J), and the adventitial cells with PDGFRα (Figure 11G, K). The morphology and staining pattern
of Sca1 overlapped significantly with PDGFRα staining and not the LMC staining (Figure 11H, L,
Supplemental Movie 6).
329
Optogenetic Stimulation of iCre-driven Channel Rhodopsin 2 to Induce Test Light-Stimulated
Depolarization Induced Lymphatic Contraction
We next used optogenetic methods to test whether the cell populations recombined by either
cKitCreER^T2, *PdgfrαCreER^TM*, or *Myh11CreER^T2* could elicit a coordinated contraction. The ChR2-
tdTomato construct appeared more sensitive to recombination than ROSA26mTmG, in some
cases resulting in LMC expression of ChR2-tdTomato in *PdgfrαCreER^TM* and *CKitCreER^T2* vessels.
Care was taken to image the vessel for tdTomato (Figure 12A,C,E) prior to stimulation at their
respective sites under brightfield conditions for diameter tracking (Figure 12B,D,F) to ensure
fidelity of the cell types and morphologies observed in Figure 3. As with ROSA26mTmG,
CKitCreER^T2 drove the ChR2-tdTomato expression primarily in large ovoid cells found on the
adventitia of the vessel. Photo-stimulation of these cells did not initiate coordinated
contractions (Figure 12G-J,S). Similarly, photo-stimulation of ChR2-tdTomato expressing cells
driven by *PdgfrαCreER^TM* failed to initiate a coordinated contraction (Figure 12K-N, T). In
contrast, photo-stimulation of LMCs, using *Myh11CreER*T2 to express Chr2-tdTomato, resulted
in a propagated contraction in the popliteal vessel (Figure 12O-R, U). In total, only 3.25% of
photo-stimulations of *cKitCreER^T2*-ChR2-TdTomato and 3.03% of photo-stimulations of
PdgfrαCreER^TM-ChR2-tdTomato were associated with a contraction, while 88.5% of photo-
stimulations of *Myh11CreER^T2*-ChR2-tdTomato photo-stimulations induced contractions (Figure
12V). The low percentages of optogenetic firing of contractions observed in *PdgfrαCreER^TM*-
ChR2-tdTomato and *cKitCreER^T2*-ChR2-TdTomato vessels are likely due to the happenstance of
spontaneous contractions occurring during the time and proximity of optogenetic stimulation.
As mast cells are not ascribed any tissue specific pacemaking behavior, the similar percentages

observed between these two groups is suggestive of coincidence. Brightfield videos of the
photo-stimulation and representative traces for *cKitCreER^{T2}*-ChR2-TdTomato, *PdgfαCreER^{TM}*-
ChR2-tdTomato, *Myh11CreER^{T2}*-ChR2-tdTomato are provided in Supplemental Movies 7-9.
355

Confocal Ca^{2+} Imaging of GCaMP6f Expression Driven by *cKitCreER^{T2}*, *PdgfαCreER^{TM}*, and
357 *Myh11CreER^{T2}* Over the Lymphatic Contraction Cycle
We imaged IALVs from *cKitCreER^{T2}*-GCaMp6f mice, which primarily resulted in expression of
GCaMp6f in the large ovoid cells in the adventitia (Figure 13A), although we occasionally
observed GCaMP6f expression in both LEC and LMCs (Figure 13A) as depicted in the maximum
projection of the acquisition period (Supplemental Movie 10) and the spatio-temporal maps
(STMS). The aberrant expressions of GCaMP6f in cells that demonstrated the typical
cobblestone morphology of LECs or the circumferential LMCs that exhibited Ca^{2+} flashes and
diastolic Ca^{2+} transients (Figure 13D,E green arrows) prior to contraction (Figure 13D,E) were
not included in the *cKitCreER^{T2}*-GCaMp6f analysis. Of the 39 *cKitCreER^{T2}*-GCaMp6f cells
analyzed, only 1 *cKitCreER^{T2}*-GCaMP6f cell exhibited a spontaneous Ca^{2+} transient during the
recording period (Figure 13B,C Cell 7). However, the Ca^{2+} transient in that cell did not align
temporally with the "Ca^{2+} flash" of the LMC with incidental GCaMP6f expression (Figure
13C,D). Despite the lack of Ca^{2+} transients under the baseline conditions throughout the IALV
contraction cycle, many *cKitCreER^{T2}*-GCaMP6f cells exhibited a robust and prolonged Ca^{2+} event
in response to stimulation with the mast cell activator compound 48-80 (Figure 13F, G, H).
Notably, the Ca^{2+} events in the ovoid cells elicited by administration of compound 48-80 did not
acutely alter the LMC Ca^{2+} activity (Figure 13I,J). Similar to *cKitCreER^{T2}*-GCaMp6f, the majority of
PDFRαCreER^{TM}-GCaMP6f expressing cells also largely lacked Ca^{2+} transients and also resulted in
incidental LMC GCaMP6f expression (Figure 14B, Supplemental Movie 11). Some cells exhibited
high basal Ca^{2+} (Figure 14A,D) sustained throughout the recording, but oscillations were not
observed (Figure 14B,C). In contrast, spurious GCaMP6f expression in a circumferentially
oriented LMC displayed Ca^{2+} flashes associated with contraction (Figure 14B,C). Of the 21
PDGFRα-GCaMP6f cells assessed, only 3 exhibited Ca^{2+} transients which were singular events
with limited spatial spread within the 20 sec imaging period (Figure 14E,F). The lack of either
global or consistent Ca^{2+} transients within either *cKitCreER^{T2}*-GCaMP6f or *PdgfαCreER^{TM}*-
GCaMP6f IALVs was in stark contrast to Ca^{2+} imaging of *Myh11CreER^{T2}*-GCaMP6f IALVs.
Myh11CreER^{T2} drove GCaMP6f expression in the circumferential LMCs (Figure 15A), which had
global and nearly synchronous Ca^{2+} flashes in 100% of the analyzed cells (Figure 15B, C).
Additionally, non-synchronous stochastic and localized Ca^{2+} transients during diastole were
commonly observed in the LMCs (Figure 15D, E, Supplemental Movie 12). Many LMCs exhibited
Ca^{2+} transients during each diastolic period while other LMCs displayed few Ca^{2+} transients or
lacked diastolic Ca^{2+} transients during the recording periods (Figure 15B). In aggregate, of the
39 *cKitCreER^{T2}*-GCAMP6f cells only 1 displayed a Ca^{2+} transient during recording, 3 of the 21
PdgfαCreER^{TM}-GCaMP6f cells, while 20 of 43 LMCs displayed at least one diastolic transient
apart from 43 of 43 LMCs with global flashes.
392

Pressure Dependency of Subcellular Ca^{2+} Transients in LMCs
We next sought to test whether diastolic Ca^{2+} transients were pressure-dependent, given that
cLVs exhibit pressure dependent chronotropy (Zawieja et al., 2019). GCaMP6f expressing LMCs

were studied in the presence of nifedipine, which blocks the calcium flashes but not local
calcium transients at intraluminal pressures of 0.5 -5 cmH$_2$O (Figure 16A). As intra-luminal
pressure was increased, there was a marked increase in the occurrence of Ca^{2+} transients
(Figure 16B, Supplemental Movies 13-15). By converting raw Ca^{2+} transients for particle
analysis (PTCLs), we generated activity maps of Ca^{2+} PTCL activity (Figure 16C) and determined
PTCL area (Figure 16D) and frequency at each pressure (Figure 16E). The maps show that as
pressure increased, the activity of PTCLs across the vessel also increased (as evident by the
increase in PTCL area activation). Across 11 experiments, the area of the field of view activated
by PTCLs/frame increased from 73.2 ± 17.7 mm^2/frame at 0.5 cmH$_2$0 to 108.6 ± 20.5
mm^2/frame at 2 cm H$_2$0 and further enhanced to 139.2 ± 26.9 mm^2/frame at 5 cm H$_2$O (Figure
16F). The number of PTCLs per frame also increased with pressure, from 2.9 ± 0.4 at 0.5 cmH$_2$0
to 4.1 ± 0.5 and 5.2 ± 0.6 PTCL/frame at 2 and 5 cmH$_2$0 respectively (Figure 16E).

Discussion
The identification of the cellular origin and signaling mechanisms underlying cLV pacemaking
will open up novel targets for pharmacological intervention in treating lymphedema and the
associated lymphatic contractile dysfunction. In this study we tested proposed pacemaker cell
types based on 3 parameters: 1) that the pacemaker cells are located along the entire length of
the cLV, to accommodate spontaneous contractions and coordinated electrical conduction
despite progressive shortening of cLVs; 2) that depolarization of the pacemaker cell can drive a
coordinated and propagated contraction of the vessel; and 3) that the presence of Ca^{2+}
transients precedes contraction, as commonly observed in other pacemaker cell types. We used
confocal microscopy and a combination of immunofluorescence and fluorescent reporters
under the control of various inducible Cres to identify various non-muscle cells that express the
markers CD34 and PDGFRα. From our initial fluorescence imaging studies, a role for intrinsic
pacemaking by LMCs (Van Helden, 1993; von der Weid et al., 2008), or by a novel population of
lymphatic ICLC (McCloskey et al., 2002; Briggs Boedtkjer et al., 2013), also referred to as
telocytes, were further examined. However, the PDGFRα cell population had minimal Ca^{2+}
activity and optogenetic depolarization of these cells failed to drive a spontaneous contraction.
In contrast, photo-stimulation of LMCs expressing ChR2 elicited robust propagating
contractions similar to spontaneous activity. Furthermore, Ca^{2+} imaging in LMCs revealed
diastolic Ca^{2+} transients in diastole that increased in frequency and spatial spread as pressure
was elevated. Our results, in addition to the recent findings using targeted deletion of Ano1
(Zawieja et al., 2019), Cx45 (Castorena-Gonzalez et al., 2018b), or Cav1.2 (To et al., 2020; Davis
et al., 2022) in lymphatic muscle support the model of LMCs as the intrinsic pacemaker as has
been previously proposed (Van Helden, 1993; Van Helden et al., 1996; Van Helden and Zhao,
2000).

Pacemaking in Smooth Muscle
In many smooth muscle organs, regulation of a coordinated contraction is a complex and
multicellular phenomenon. Multiple cell types integrate physical and biological information into
electrical activity to be transmitted to the force-producing smooth muscle cells, sometimes
across great distances relative to cell size, to regulate calcium influx by voltage dependent
calcium channels required for contraction. The intestine is one such documented tissue in

which cKit[+] ICCs and interstitial PDGFRα[+] cells form an electrical syncytium to regulate intestinal
motility (Sanders et al., 1999; Sanders et al., 2014). The pacemaking function of intestinal ICCs
relies heavily on Ano1, a calcium activated chloride channel, which is required for slow wave
activity in the ICCs. Both cKit and Ano1 can be used as a marker for ICCs in the intestine (Hwang
et al., 2009; Cobine et al., 2017; Malysz et al., 2017), cKit[+] and vimentin[+] ICLCs have been
observed in sheep lymphatic vessels (McCloskey et al., 2002), yet these cell populations do not
form gap junctions with the smooth muscle to form electrical connections (Briggs Boedtkjer et
al., 2013) as occurs in the intestinal ICCs. Our cKit staining of mouse IALVs revealed a sparse
population of large ovoid cells previously classified as mast cells (Chatterjee and Gashev, 2012;
Zawieja et al., 2019). Their identity as mast cells was further supported by the sustained global
Ca^{2+} event after stimulation with the mast cell degranulating agent compound 48-80. However,
both vimentin and CD34 showed robust staining throughout the mouse lymphatic vessel wall.
Vimentin stained LECs, as well as non-muscle stellate shaped cells, with many co-expressing
CD34, and other smaller circular cells some of which were cKit[+] as well and some whose
morphology was similar to that of the macrophage staining profile of the GFP[+] cells in IALVs
from MacGreen mice , as well as previous reports of macrophage staining in cLVs (Bridenbaugh
et al., 2013; Chakraborty et al., 2015; Zawieja et al., 2016). While vimentin[+] cells had a nuclear
and perinuclear staining profile, CD34 demarcated the cell membrane and was useful for
assessing the morphology of these cells. Of particular interest, the vimentin[+]CD34[+] cells densely
populated the length of the mouse IALV, with a majority displaying a flattened stellate
morphology characterized by the classic rounded oak leaf appearance, although some displayed
fine dendrite extensions. Contrasting with the previous findings in the human thoracic duct
(Briggs Boedtkjer et al., 2013), we did not observe a significant population of CD34[+] cells with a
bipolar morphology oriented axially along the vessel. However, z-stack reconstructions of
sections of the mouse IALV that included the secondary valves revealed interstitial
CD34[+]PDGFRα[+] cells that resembled the bipolar cells with multiple axon-like extensions
throughout the endothelial leaflets similar to the interstitial cells that were reported in the
lymphovenous valves (Geng et al., 2016). While these cells have not been frequently described
in the peripheral cLV valves, we observed these cells in each of the valve regions we imaged in
addition to labeling them with other Cre drivers, including NG2-Cre;ROSA26mTmG and
*Pdgfrβ CreER*T2 ROSA26mTmG (data not shown). Whether these cells play a role in the
extracellular matrix deposition or lymphatic valve integrity is unknown, but their role as a
critical pacemaker can be excluded as vessel segments without valves display normal
contractile behavior. Instead, the majority of the CD34[+]PDGFRα[+] cells were found in the
adventitia with 2-3 layers overtop the LMCs and were consistently observed in high density
along the IALV. Some CD34[+]PDGFRα[+] cells or their extensions were present between the
lymphatic endothelial and muscle layers as had been previously reported with electron
microscopy of human lymphatic vessels (Briggs Boedtkjer et al., 2013).
478
479   **PDGFRα[+]CD34[+] Cells are Not Involved in cLV Pacemaking Under Physiological Conditions**
480   CD34 and PDGFRα are described as telocyte markers, although PDGFRα routinely labels
481   fibroblasts and specific interstitial cells in the GI tract involved in purinergic neurotransmission
482   (Kurahashi et al., 2011; Kurahashi et al., 2013; Clayton et al., 2022), and CD34 expression has
483   been observed in multipotent cell populations of various origins (Sidney et al., 2014). Of course,

neither telocytes, hematopoietic or mesenchymal stem cells, nor fibroblasts, are monolithic in their expression patterns within and much less across tissues, and single cell RNA sequencing has provided immense detail about the sub clusters and spectrums within which these cells exist as well as their plasticity. Nonetheless, we attempted to gain further insight into the characteristics of CD34[+]PDGFRα[+] cells. CD34[+]PDGFRα[+] cells were consistently negative for the majority of smooth muscle markers such as desmin, calponin, smooth muscle actin, smooth muscle myosin heavy chain, and the pericyte marker MCAM. However, PDGFRβ expression was noted in sorted *PdgfrαCreER*[TM]-ROSA26mTmG cells, in addition to the staining of LMCs, and PDGFRβ protein expression was confirmed with variable immunofluorescence staining amongst the PDGFRα stained cells as well as LMCs. The *PdgfrβCreER*T2-ROSA26mTmG mice had only modest recombination in both the LMC and PDGFRα[+] cell population, but potentially highlighted a myofibroblast-like cell subpopulation, cells that might lie on the spectrum of differentiation from lymphatic muscle and PDGFRα[+] cells, or perhaps a cell with pacemaker activity as PDGFRβ is widely used as a pericyte marker and some pericytes display pacemaker activity (Hashitani et al., 2015). Adding to this intrigue, the *PdgfrαCreER*[TM] sorted cells expressed transcripts for Cav1.2, the voltage-gated L-type Ca[2+] channel critical for lymphatic contractions (Zawieja et al., 2018; To et al., 2020); Ano1, the ion channel underlying pressure dependent chronotropy (Mohanakumar et al., 2018; Zawieja et al., 2019); and Cx45, the primary connexin mediating electrical conduction in mouse lymphatic collecting vessels (Castorena-Gonzalez et al., 2018b; Hald et al., 2018). The presence of those gene transcripts does not appear to be due to muscle cell contamination or incidental recombination in LMCs as we did not detect LMC markers in the RT-PCR profiling of the sorted PDGFRα[+] cells nor were GFP-expressing cells with an LMC morphology observed in imaging of *PdgfrαCreER*[TM]-ROSA26mTmG vessels. Critically, however, deletion of Cav1.2, Cx45, or Ano1 through *PdgfrαCreER*[TM] mediated recombination neither recapitulated the previous phenotypes achieved with *Myh11CreER*[T2] (Castorena-Gonzalez et al., 2018b; Zawieja et al., 2019; To et al., 2020; Davis et al., 2022) nor significantly affected pacemaking in mouse popliteal cLVs. This is in stark contrast to the complete lack of contractions observed in *Myh11CreER*T2-Cav1.2 vessels (To et al., 2020) or the vessels from vascular muscle specific Itga8CreERT2-Cav1.2fl/fl mice (Davis et al., 2022; Warthi et al., 2022),and the significant loss in pressure-induced chronotropic modulation of pacemaker function in IALVs with *Myh11CreER*[T2]-mediated deletion of Ano1 that we have previously reported (Zawieja et al., 2019) . While a sub-population of CD34[+]PDGFRα[+] cells may share expression of critical pacemaker genes identified in the LMCs, they do not appear to be involved in cLV pacemaking or contractile function under physiological states. Instead, CD34[+]PDGFRα[+] cells co-stained significantly with Sca1[+], suggesting they may be primed to act as resident multipotent cells (Song et al., 2020; Kimura et al., 2021). To this point, the *PdgfrαCreER*[TM] FACS purified cells also expressed markers associated with "stemness" such as CD34, Klf4, Gli1, CD29, CD105, CD44, and vimentin, in addition to Sca1, and it is likely that the *PdgfrαCreER*[TM] population includes various distinct subpopulations (Jolly et al., 2022) expressing these markers. These cells may play a role in rebuilding the lymphatic collecting vessel vasculature following lymph node resection and further studies are required to assess their functional multipotency.

SR Ca[2+] Cycling in Pacemaking

The use of the mouse model, in addition to the simplicity of the vessel architecture, provided
the use of genetic tools that previously had been instrumental in identifying the cKit$^+$ ICC as the
pacemaker cells of the GI tract (Ward et al., 1994; Huizinga et al., 1995; Torihashi et al., 1995).
Through the use of the respective *PdgfraCreER^{TM}* and *Myh11CreER^{T2}* models, were able to
specifically image Ca^{2+} in each cell type in pressurized, contracting vessels. Pacemaking
initiating cells have an inherently unstable membrane potential, oftentimes utilizing the
oscillatory nature of Ca^{2+} release from the sarcoendoplasmic reticulum coupled to Ca^{2+} sensitive
electrogenic exchangers and ion channels to drive depolarization (Van Helden, 1993; Hashitani
et al., 2015; Baker et al., 2021b; Sanders et al., 2022). One such example is the pacemaker ICC
in the gastric corpus which exhibit abundant Ca^{2+} transients that couple to Ano1-mediated
chloride currents in both the intervening period between slow waves as well as the plateau
phase of the slow wave (Baker et al., 2021a), although such activity is not characteristic of all
pacemaker ICC types. The identification of a Ca^{2+} activated chloride current in LMCs (Van
Helden, 1993; Toland et al., 2000) and its correspondence with subcellular Ca^{2+} transients (Van
Helden, 1993; Ferrusi et al., 2004; von der Weid et al., 2008) led Van Helden to postulate that
LMCs had an intrinsic pacemaking capability (Van Helden, 1993; Van Helden et al., 1996). We
have previously reported that mouse LMCs in pressurized vessels routinely display subcellular
Ca^{2+} release events that reflect the kinetics and characteristics of Ca^{2+} puffs and waves in
addition to the coordinated global Ca^{2+} flash associated with influx during an action potential
(Castorena-Gonzalez et al., 2018b; Zawieja et al., 2018; Zawieja et al., 2019). Here we confirmed
the consistent presence of subcellular Ca^{2+} transients only in LMCs with GCaMP6f driven by
Myh11CreER^{T2} but not in the cells with GCaMP6f driven by *PdgfraCreER^{TM}*. Critically, we also
demonstrated that the Ca^{2+} transients increased in both frequency and spatial spread as
pressure was elevated in the vessel, as would be expected to account for the pressure
dependent chronotropy observed in lymphatic collecting vessels. This underscores the recent
finding that the genetic deletion of Ano1 in the LMCs dramatically reduced contraction
frequency and abolished pressure-dependent chronotropy in those vessels (Zawieja et al.,
2019). This phenotype was largely replicated with a similar reduction in frequency and loss of
pressure dependent chronotropy in our recent study utilizing *Myh11CreER^{T2}* to drive deletion of
IP3R1 from LMCs (In revision, JGP 2023) and fits with the central role of IP3R and subcellular
Ca^{2+} release as critical components of intrinsic LMC pacemaking (Van Helden et al., 1996; von
der Weid et al., 2008). The lack of Ca^{2+} transients in PDGFRα$^+$ cells across any stage of the
lymphatic contraction cycle diminishes any expected role for this cell type to perform as the
pacemaker for the mouse IALV. The contribution of specific LMC SR Ca^{2+} signals to lymphatic
pacemaking remains to be fully addressed.
563
A pacemaker cell would be expected to be electrically coupled to the LMC layer to permit the
nearly synchronous conduction velocity of the contraction wave (Zawieja et al., 1993;
Castorena-Gonzalez et al., 2018b; Hald et al., 2018) and to transmit their own depolarization
into their coupled LMCs to activate the voltage dependent Ca^{2+} channels that are responsible
for lymphatic muscle action potentials. Connexins are the molecular constituents of gap
junctions and, as stated above, we detected Cx45 expression in the *PdgfraCreER^{TM}* sorted cells.
However, we did not detect any impairment in pacemaking, nor were contraction conduction
speed deficits or multiple pacemakers noted in the *PdgfraCreER^{TM}* -Cx45fl/fl popliteal cLVs, in

contrast to the development of multiple pacemaker sites and the lack of entrainment that were
reported previously in *Myh11CreER^{T2}*-Cx45fl/fl cLVs (Castorena-Gonzalez et al., 2018b).
Admittedly, we did not perform an exhaustive assessment of the connexin expression profile of
the CD34$^+$PDGFRα$^+$ cells, and Cx45 may not be the dominant connexin expressed in the
CD34$^+$PDGFRα$^+$ cells, or heterotypic connexons could exist (Koval et al., 2014). However,
electron microscopy studies of the putative ICLC in the human thoracic duct did not detect any
gap junctions, although peg-and-socket connections were observed (Briggs Boedtkjer et al.,
2013). We utilized optogenetics to enforce channel rhodopsin expression in both the
PdgfrαCreER^{TM} and *Myh11CreER^{T2}* mouse models to directly depolarize the specific cell
populations in an attempt to drive a contraction. Local photo-stimulation of the PDGFRα cells
failed to initiate contraction while the stimulation of *Myh11CreER^{T2}* recombined cells resulted
in contractions that were indistinguishable from the spontaneously occurring contractions.
These results give functional credence to the lack of hetero-cellular coupling that was
previously reported . Just as critically, they also highlight the regenerative nature of the
lymphatic muscle action potential as local depolarization was sufficient to drive a coordinated
contraction along the vessel and that a single or few LMCs reaching threshold for AP initiation
are sufficient to drive the conducted activity observed at the tissue level.
589
Conclusions
Our present findings lend further support to the hypothesis that the LMCs are intrinsic
pacemakers (van Helden et al., 2006; Mitsui and Hashitani, 2020) and do not require an ICC-like
network to drive propagated contractions. These findings also underscore the significance of
lymphatic muscle Ca^{2+} handling as the driver of lymphatic pacemaking, which can be
compromised in disease states leading to impaired lymphatic contractile activity (Stolarz et al.,
2019; Lee et al., 2020; Van et al., 2021). Further studies delineating the specific SR Ca^{2+} release
and influx pathways, and the contributions of Ca^{2+} sensitive ion channels need to be identified
to develop sophisticated in silico models and identify potential therapeutic targets to rescue
lymphatic pacemaking in lymphedema patients (Olszewski, 2002, 2008).
600
Limitations
One assumption underlying our conclusions is that there is a conserved fundamental
pacemaking pathway in lymphatic muscle pacemaking across species, specifically pertaining to
the capability of lymphatic muscle to maintain pacemaking and coordination despite changes in
tissue complexity. Lymphatic collecting vessels in mice have similar pressure-dependent
chronotropy and contraction conduction velocity as recorded in human vessels (Castorena-
Gonzalez et al., 2018b). These similarities exist despite the fact that mouse lymphatic collecting
vessels are encircled by a single layer of lymphatic muscle while larger species have multiple
layers of LMCs in the wall. It is possible that vessels with multiple layers of LMCs need a
network of ICLC to coordinate their activity. The simplicity in the makeup of the mouse cLV and
the use of cell targeting Cre models provide great control over our experimental variables, but
other cell types may provide coordination of LMC pacemaking in other species where the
lymphatic cLV walls are larger with multiple muscle layers.
614

Our data demonstrates that limited staining of a few cell markers alone is insufficient to identify
discrete cell populations in the murine cLVs. Additionally, mRNA expression does not equal
protein translation nor guarantee specific function as we did not detect endothelial CD34 with
immunofluorescence despite detecting transcript; additionally, *PdgfrαCreER^{TM}* mediated
deletion of Ano1, Cx45, or Cav1.2 had no effect on cLV pacemaking. Hence, further
experimentation is also required to fully characterize expression of multipotent cell markers
and function of CD34$^+$PDGFRα$^+$Sca1$^+$ cells invested within the murine cLVs, although this was
beyond the scope of this study assessing pacemaker identity. Tangentially, another limitation of
our approach pertains to the specificity and recombination efficiency of inducible Cre
recombinase models, which can be a notable confounding variable (Chakraborty et al., 2019).
We observed that our inducible Cre models led to a degree of nonspecific recombination within
the murine cLV, with the GCaMP6f and ChR2 particularly susceptible to recombination
compared to the ROSA26mT/mG reporter. Recombination in multiple cell types was expected
with the constitutive Cre models we employed (*Ng2Cre* and *PdgfrαCre*), as vascular and
lymphatic muscle precursor cells can transiently express nestin, PDGFRα, and NG2 (Hill et al.,
2015; Castorena-Gonzalez et al., 2018b; Kenney et al., 2020). We also observed that
PdgfrβCreER^{T2} drove recombination in a sub population of LMCs and PDGFRα$^+$ cells. Whether
these are two distinct populations that only share expression for PDGFRβ or whether they
demonstrate a continuum of precursor LMCs and newly-formed LMCs are both plausible
explanations. PDGFB-PDGFRβ signaling is critical for normal mural cell recruitment to both the
blood and lymphatic vasculature (Gaengel et al., 2009; Wang et al., 2017) and proliferating
vascular smooth muscle cells and pericytes have both been documented to express PDGFRβ
(Andrae et al., 2008; Pitulescu and Adams, 2014). We have recently performed single cell
RNAseq on isolated IALVs and, while these this dataset is still being analyzed, sub-populations
of LMCs, LECs, and CD34$^+$ and PDGFRα$^+$ were readily identified (data not shown) highlighting
the difficulty in ascertaining functionality on the basis of expression of a few cell markers.
Ideally, novel Cre or combinatorial Cre models may be developed to further tease out the
functional role of these expected sub populations.
643
Materials and Methods
Mice Wild-type (WT) male mice (25-35 g) on the C57BL/6J background , ROSA26mT/mG reporter
(Muzumdar et al., 2007) (Strain#007676), transgenic *PdgfrαCre* (Strain#013148), CSFR1-EGFP
(MacGreen) (Sasmono et al., 2003) (Strain#018549), genetically encoded Ca^{2+} sensor GCaMP6f
(Chen et al., 2013) (Strain#028865), transgenic *PdgfrαCreER^{TM}* (Kang et al., 2010)
(Strain#018280), NG2-Cre (Strain #:008533) (Zhu et al., 2008), and ChR2 /tdTomato fusion mice
(Madisen et al., 2012) (Strain#012567) were purchased from The Jackson Laboratory (Bar Harbor,
MA, USA). *PdgfrβCreER^{T2}* (Gerl et al., 2015) mice were a gift from Ralf Adams (Mac Planck
Institute) and kindly provided by Lorin Olson (Oklahoma Medical Research Foundation), and are
currently available at Jax (Strain#029684). The *Myh11CreER^{T2}* mice (Wirth et al., 2008) were a gift
from Stefan Offermanns, Max-Planck-Intstitut fur Herz-und Lungendforschung, Bad Nauheim,
Germany, and are currently available at Jax (Strain #019079, Y-Linked). c-KitCreER^{T2} mice (Heger
et al., 2014) were a gift from Dieter Saur (Technical University of Munich). Prox1-eGFP mice (Choi
et al., 2011) were a gift from Young-Kwon Hong (University of Southern California. For
genotyping, we isolated genomic DNA from mouse tail clips using the HotSHOT method (Truett

et al., 2000). Specific mouse genotypes were confirmed via PCR using 2x PCR Super Master
Polymerase Mix (Catalog # B46019, Bimake, Houston, TX) performed as specified by the provider.
Mice were over 3-8 months of age during this study. All animal protocols were approved by the
University of Missouri Animal Care and Use Committee and conformed to the US Public Health
Service policy for the humane care and use of laboratory animals (PHS Policy, 1996).
664
iCre Tamoxifen Induction- Mice harboring *PdgfαCreER^{TM}*, *PdgfrβCreER^{T2}*, *Myh11CreER^{T2}*, and
cKitCreER^{T2} were crossed with ROSA26mT/mG mice to generate *PdgfαCreER^{TM}*-ROSA26mT/mG,
PdgfrβCreER^{T2}-ROSA26mT/mG, *Myh11CreER^{T2}*-ROSA26mT/mG, and *cKitCreER^{T2}*-ROSA26mT/mG
mice, respectively. The resulting iCre-ROSA26mT/mG mice were induced with tamoxifen 2-4
weeks after weaning. Tamoxifen induction was performed via consecutive 100 µL i.p. injections
of tamoxifen ranging from 1 to 5 days at concentration ranging from 0.2 -10 mg/mL in safflower
oil, using a titrated induction protocol to determine the extent of recombination in specific cell
populations. We used our maximal induction protocol, 100 µL of tamoxifen at 10 mg/mL over 5
consecutive days, for *cKitCreER^{T2}*-GCaMP6f, *Myh11CreER^{T2}*-GCaMP6f, and *PdgfαCreER^{TM}* -
GCaMP6f mice. Due to the paucity of recombined cells in the *cKitCreER^{T2}*-ROSA26mT/mG
reporter mice, we used our maximal tamoxifen induction protocol for *cKitCreER^{T2}*-ChR2 mice as
this still resulted in the ability to excite single recombined cells. *Myh11CreER^{T2}*-ChR2/tdTomato
mice were induced with one 100 µL i.p. injection of tamoxifen at 0.2 mg/mL while *PdgfαCreER^{TM}*-
ChR2/tdTomato were induced with 1 injection at 0.4 mg/mL tamoxifen to get mosaic induction
sufficient for single cell stimulation. All mice, regardless of duration, were given 2 weeks to
recover following tamoxifen injection.
681
Lymphatic Vessel Isolation- We utilized both popliteal and inguinal-axillary lymphatic collecting
vessels (IALVs) in this study, which were isolated as described previously (Zawieja et al., 2018). In
brief, mice were anaesthetized with a cocktail of 100/10 mg/mL) ketamine/xylazine mg/mL and
shaved along the flank or the legs for IALVs and popliteal cLVs respectively. The IALV (also
referred to as the flank cLV) is located adjacent to the thoracoepigastric vein and connects the
inguinal and axillary lymph node. A cut was made along the dorsal midline and the skin retracted
and pinned out to reveal the thoracoepigastric vascular bed. The thoracoepigastric vascular bed
and connected perivascular adipose containing the IALVs vessel was dissected out and pinned
onto a Sylgard coated dish in Krebs buffer. Popliteal lymphatic vessels were exposed through a
superficial incision in the leg, removed and transferred to the Krebs-albumin filled dissection
chamber. After removal, the vessel was carefully cleaned of adipocytes and excess matrix using
fine forceps and scissors through micro-dissection. For immunofluorescence, sections containing
2-3 valves were isolated, while smaller IALV sections consisting of 1-2 valves were isolated for
GCaMP6f Ca^{2+} imaging. Similarly, popliteal cLVs were isolated (Castorena-Gonzalez et al., 2018a)
following an incision along the skin overlying the saphenous vein for contractile function analysis
and for ChR2 optogenetic depolarization experiments.
Lymphatic Vessel Isobaric Function- *PdgfαCreER^{TM}* mice were crossed with Ano1fl/fl, Cx45fl/fl,
and Cav1.2fl/fl and to achieve *PdgfαCreER^{TM}*-Ano1fl/fl, *PdgfαCreER^{TM}*-Cx45fl/fl, and
PdgfαCreER^{TM}-Cav1.2fl/fl mice. Theses mice and their respective fl/fl controls were injected
with tamoxifen as described above for 5 days and given two weeks to recover. The popliteal

vessels were isolated, cleaned, and prepared for isobaric contractile tests as previously reported (Davis et al., 2023). Once equilibrated, inner diameter was tracked over a physiological pressure range (stepped from 3 to 2, 1, 0.5, 3, 5, 8, and 10 cmH$_2$O) with 2min of recording at each pressure. Following the pressure step protocol the vessels were equilibrated in with Ca^{2+}-free Krebs buffer (3mM EGTA) and diameter at each pressure recorded under passive conditions. The contractile parameters end diastolic diameter (EDD), end systolic diameter (ESD), and contraction frequency (FREQ) were recorded with a custom LabVIEW program and the following contractile parameters assessed:

1) Contraction Amplitude (AMP) = EDD−ESD

2) Normalized Contraction Amplitude = ((EDD−ESD)/DMAX) × 100

3) Ejection Fraction (EF) = (EDD^2−ESD^2)/EDD^2

4) Fractional Pump Flow (FPF) = EF × FREQ

5) Tone = ((DMAX−EDD)/DMAX) × 100

Methylene Blue- Isolated IALVs sections were transferred into a Krebs-BSA buffer filled 3-mL observation chamber, with a cover slip bottom, and cannulated onto two glass micropipettes (30-80 μm, outer diameter) held in place by pipette holders on a Burg-style V-track mounting system. The pipette holders were attached to a 3-way valve stop cock with polyethylene tubing filled with Krebs-BSA buffer. Vessels were pressurized to approximately 5 cmH$_2$O by raising the 3-way valve and the vessels were stretched to remove any slack. For methylene blue staining, IALVs from wild type C57Bl6 mice were stained with 50 μM methylene blue in Krebs-BSA buffer for two hours at room temperature and covered in foil to limit light induced phototoxicity. After the staining period, the vessel chambers were washed three times with Ca^{2+} free PSS to remove methylene blue. Brightfield images and manual Z-stack videos were collected on an inverted Leica DMi1 4X or 20X air objective, or a Leica DMi8 with a 25X water objective or an inverted DMi8 using a Leica Flexacam C1 color camera for image acquisition. Some Methylene blue images were also collected using a color Nikon DS-Fi3 camera. The collected z-stacks were analyzed using Image J and the "Stack Focuser" plugin (https://imagej.nih.gov/ij/plugins/stack-focuser.html). To accentuate the methylene blue stained cells, the color image stack was split into red, green, and blue channel stacks. The blue channel stack was then divided by the green channel stack using the "Image Calculator" function. The resulting 32-bit image was then converted into 16-bit image to permit the use of the Stack Focuser plugin with the 'n kernel value' set to 11.

Fluorescence Confocal Imaging- IALVs vessels from each respective iCre-ROSA26mT/mG mouse were prepared in a similar manner (excluding the addition of methylene blue). We performed confocal imaging to acquire z-stacks of 7-10 overlapping regions of interests to allow for manual stitching, with 1 μM z-steps (20X) or 0.5 μM steps at 40X. We imaged through to the midpoint of the vessel except when imaging the valve interstitial cells, in which case the entire vessel was

imaged. Max projections were made using FIJI. Following live imaging, the vessels were pressurized to 5 cmH$_2$O and fixed with 4% paraformaldehyde for 30 min at room temperature. IALVs were then washed with PBS containing 0.1% Triton X-100 (PBST) 3 times and blocked for a minimum of 2 hr with Blockaid® (B-10710, ThermoFisher Scientific). IALVs were then stained with the corresponding primary antibodies in BlockAid® Solution: anti-smooth muscle actin (SMA) 1:500 (Sigma, A2547), anti-GFP 1:200 (ThermoFisher, A11122), anti-cKit 1:100 (Cell Signaling, 3074), anti-Vimentin 1:100 (Thermofisher , OMA1-06001), anti-desmin 1:200 (Invitrogen, PA5-16705), anti-GFP 1:200 (Abcam, ab13970, anti-CD34 1:200 (Invitrogen, 14-0341-82), anti-PDGFRA 1:200 (R&DSystems, AF1062), anti-PDGFRβ 1:200 (eBiosciences, 14-1402-82), anti-calponin 1:500 (Abcam, AB46794), anti-MYH11 1:500 (Abcam, AB124679), anti-Sca1 1:200 (Biolegend, 108101). IALVs were then washed in PBS and incubated overnight with the corresponding donkey secondary antibodies (ThermoFisher®) at 1:200. After a final wash, IALVs were re-cannulated and pressurized for imaging using the aforementioned spinning disk confocal and Hamamatsu Orca Flash4 camera using a 20X air objective (Olympus UplanApo, 0.75) or 40X (Olympus UApo A340, 1.15) water objective. Images were taken as described above, and the resulting stacks were turned into a max projection using FIJI. Colocalization analysis of the max projections of CD34 and PDGFRα was performed using the BIOP JACoP colocalization plugin (Bolte and Cordelieres, 2006) with both Pearson's and Mander's coefficients reported.

LMC Dissociation and FACS Collection- IALVs vessels *PdgfrαCreER^{TM}*-ROSA26mT/mG, *PdgfrβCreER^{T2}*-ROSA26mT/mG, *Myh11CreER^{T2}*-ROSA26mT/mG, Macgreen, and *Prox1-eGFP* mice were dissected and cleaned of excess adventitia and adipose tissue in Krebs buffer. Isolated vessels were then transferred into a low Ca^{2+} PSS solution supplemented with 0.1 mg/mL bovine serum albumin (BSA, Amersham Life Science, Arlington Heights, IL). Primary LMCs were collected by enzymatic dissociation of IALVs. The dissected vessels were cleaned in room temperature Krebs-BSA buffer and then transferred into a 1-mL tube of low-Ca^{2+} PSS on ice, washed, and equilibrated for 10 min. Vessels were then digested in low-Ca^{2+} PSS with 26 U/mL papain (Sigma, St. Louis, MO) and 1 mg/mL dithioerythritol for 30 min at 37°C and were gently agitated every few minutes. This solution was then decanted and replaced with low-Ca^{2+} PSS with containing 1.95 collagenase H (U/mL, Sigma), 1.8 mg/mL collagenase F (Sigma), and 1mg/mL elastase (Worthington LS00635) and incubated for 3 - 5 min at 37° C. The mixture was then spun down at 1000 rpm for 4 min, the digestion buffer removed, and replaced with low-Ca^{2+} PS. This process was repeated twice to remove residual digestion buffer. The vessel was then triturated with a fire-polished Pasteur pipette to dissociate the cells into a single cell suspension, passed through a Falcon cap strainer (35 µm), and resuspended in ice-cold low-Ca^{2+} PSS for sorting. For iCre-ROSA26mT/mG mice, GFP$^+$RFP$^-$ cells or GFP$^+$ cells from Macgreen and *Prox1-eGFP* mice were then FACS purified straight into RNA isolation buffer for RT-PCR analysis. FACs was performed with a Beckamn-Coulter MoFlo XDP instrument using an excitation laser (488 m) and emission filter (530/40 m). Sorting was performed using 70-µm nozzle at a sheath pressure of 45 p.s.i. and sort rate of 100 events/s and with an efficiency of >90%. To maximize cell yield, we isolated both the left and right full-length IALVs vessels from 2 mice for digestions and subsequent FACS collection. For *Myh11CreER^{T2}*-ROSA26mT/mG and *PdgfraCreER^{TM}*-ROSA26mT/mG, the yield averaged 1000-2000 cells per mouse. For *Prox1-eGFP* mice, LEC yield was typically 1500-2000 cells per mouse.

RT-PCR Profiling of FACS Purified Cells- Total RNA was extracted from FACS purified GFP$^+$ cells from the isolated IALVs vessels using the Arcturus PicoPure RNA isolation kit (ThermoFisher Scientific, Waltham, MA) per the listed instructions. Prior to elution in 20 µl of water, on-column DNAse digestion (Qiagen, Valencia, CA) was performed to ensure removal of genomic DNA contaminants. RNA was converted into cDNA using SuperScript III First-Strand Synthesis System (Thermo Fisher Scientific, Waltham, MA) using oligo (dT) and random hexamer priming following the manufacturer's protocol. Each RT reaction used approximately 50-100 cells worth of RNA based on the sorted cells count number. Our PCR reaction mixture contained first-strand cDNA as the template, 2 mM $MgCl_2$, 0.25 µM primers, 0.2 mM deoxynucleotide triphosphates; and GoTaq® Flexi DNA polymerase (Promega, Madison, WI). The PCR program comprised an initial denaturation step at 95°C for four min; followed by 35 repetitions of the following cycle: denaturation (94° C, 30 s), annealing (58° C, 30 s), and extension (72° C, 30 s). This was followed by a final elongation step for 5 min at 72° C. PCR amplification products were separated on a 2% agarose gel by electrophoresis, stained with SYBR-Safe (Thermo Fisher Scientific, Waltham, MA), and visualized by UV trans-illumination. All primers were designed to amplify an intron-spanning region. Endpoint RT-PCR Primer sequences, amplicon size, accession numbers, and source are listed in Table 1.

***Ex vivo* Ca^{2+} imaging with the genetically encoded GCaMP6f Indicator**- *cKitCreER^{T2}*, *Myh11CreER^{T2}*, and *PdgfαCreER^{TM}* mice were crossed with GCaMP6f mice in a similar manner as described for ROSA26mT/mG. *cKitCreER^{T2}*-GCaMP6f, *PdgfαCreER^{TM}*-GCaMP6f, and *Myh11CreER^{T2}*-GCaMP6f were induced with tamoxifen (10 mg/mL) for 5 consecutive days by i.p. injection. IALVs isolated from *cKitCreER^{T2}*-GCaMP6f, *PdgfαCreER^{TM}*-GCaMP6f, and *Myh11CreER^{T2}*-GCaMP6f were cannulated as described above. The cannulated vessel, with micropipette holders, observation chamber and V-track mounting system, was transferred to the stage of the spinning disk confocal with a Prime95B scMOS camera (Photometrics), a Cascade II EMCCD (Photometrics), or an Ixon888 EMCCD camera (Andor) for Ca^{2+} imaging (Castorena-Gonzalez et al., 2018b). Pressures for the input and output cannula were connected to a T-junction which was set briefly to 8 cmH$_2$O and the vessel lengthened to remove axial slack. A peristaltic pump maintained constant perfusion of the observation chamber with Krebs buffer at a rate of 0.5 mL/min while the vessel equilibrated at 37°C for 30-60 min with pressures set to 3 cmH$_2$O. Spontaneous contractions were allowed to stabilize over a period of 30 min and then were blunted with 2 µM wortmannin to limit movement associated with contractions during Ca^{2+} imaging. A Windows-based computer was used to digitize the pressure transducer signals and video image of the vessel from a firewire camera at 30-40 Hz (Davis et al., 2012). A custom-written LabVIEW program (National Instruments; Austin, TX) detected the inner diameter of the vessel from the video (Davis et al., 2011). Once contractions were <5 µm in amplitude, Ca^{2+} recordings were made at 20FPS for 20-40 s.

Ca^{2+} Imaging and Analysis in IALVs Over the Contraction Cycle- Background noise was determined by using the histogram feature of FIJI in a rectangle in a region of the field of view without sample. This value was subtracted from the entire field of view. In some cases, the vessel movement due to contraction was offset with video stabilization with the FIJI plugin Image Stabilizer. A max projection was used to create non-overlapping ROIs of GCaMP6f$^+$ cells for each

iCre-GCaMp6f IALV. From these cell ROIs, the "reslice z" function was used to create a pseudo-
linescan STMs which were divided by their baseline values to obtain F/F_0 values for each
individual cell. At least 3 cells, except in the case of 1 *cKitCreER^{T2}*-GCaMp6f IALV, in which only
two cells were observed, were analyzed in this manner for each vessel segment. Max projections
of the image stack were then used to create non-overlapping cell masks of 3-5 muscle cells per
field of view of one vessel. Ca^{2+} traces for those cells contained 5-10 contraction cycles and Ca^{2+}
transients and were characterized for peak intensity (expressed as a baseline-referenced ratio,
F/F_0), frequency, and duration in seconds.
836
Analysis of Subcellular Ca^{2+} Transients in *Myh11CreER^{T2}*-GCaMP6f IALVs For *Myh11CreER^{T2}*-We
performed Ca^{2+} imaging as above in the presence of 1 µM nifedipine to stop the "Ca^{2+} flashes"
associated with action potentials (Zawieja et al., 2018) and focus on the subcellular activity at 3
different experimental pressures of 0.5, 2, and 5 cmH$_2$O. For this protocol, we used a particle
analysis approach to analyze all Ca^{2+} transients in the field of view. Ca^{2+} transients in intact vessels
were quantified by particle analysis as previously described (Drumm et al., 2017; Drumm et al.,
2019). Movies of Ca^{2+} transients in intact vessels were imported into custom built Volumetry
software (version G8d) and background subtracted. Movies were smoothed using a Gaussian
filter: 1.5 x 1.5 mM, StdDev 1.0). Raw Ca^{2+} transients were converted to Ca^{2+} particles (PTCLs)
using a flood-fill algorithm as previously described (Drumm et al., 2017; Drumm et al., 2019).
PTCLs <10 mm^2 were rejected to facilitate the removal of noise and then the total PTCL area and
PTCL count could be tabulated for each recording.
849
Light Activation of ChR2 to stimulate Popliteal Collecting Lymphatic Vessel Contractions.
As the IALV has a nearly continuous contractile cycle, we utilized the popliteal vessel for its much
slower contraction frequency in the experiments testing our ability to trigger a propagated
contraction upon stimulation of the enforced expression of ChR2. Popliteal vessels were isolated
from *cKitCreER^{T2}*-ChR2/tdTomato, *PdgfαCreER^{TM}* -ChR2/tdTomato, or *Myh11CreER^{T2}*-
ChR2/tdTomato mice as previously described (Scallan and Davis, 2013), although we intentionally
retained some connective tissue and adipose tissue to ensure we had a sufficient population of
recombined cells to test in the adventitia layer of the vessel. Contractions were allowed to
stabilize over a 30-min equilibration period with pressure set to 3 cmH$_2$O. If basal contraction
frequency was too high, we applied pinacidil to the bath in 100 nM increments, without
exceeding 600 nM, to further slow contraction frequency to around 6 contractions per minute.
Pinacidil at sub 1 µM doses can slow contraction frequency without causing overt
hyperpolarization of membrane potential (Davis et al., 2020). Supplemental 100 nM doses of
pinacidil were applied throughout the experiment to maintain a spontaneous contraction
frequency below 6 per minute to allow ample diastolic time for ChR2 stimulation. Throughout
this protocol the popliteal was allowed to contract spontaneously to ensure we had not overly
inhibited action potentials by the pacemaking cells with pinacidil. Occasionally spontaneous
contractions occurred just prior to contractions and could result in a potential false positive so
we performed multiple stimulations over a period of 5 - 10 min, typically waiting at least 3 s after
any spontaneous contraction before stimulating. Care was made to align the light fiber in such a
way that only part of the vessel would be directly illuminated and so target cells of interest would
be directly activated by 473 nm light using a Laser diode (Doric LD Fiber Light Source, Quebec,

Canada), through an optical probe with a 10-µm tip (Doric, OPT_200_0.22.010). To further limit
the excitation field, the optical probe was coated with black acrylic paint using an eyelash brush
so that the uncoated opening was ~2-3 µm. With the probe positioned within 5 µm of one side
of the vessel wall, the spread of light covered an area ~10-100 µm wide on the back side of the
vessel (depending on the diode amplitude setting). Light pulses, 200 ms in length, were triggered
by a Grass S9 stimulator (Harvard Apparatus, Holliston, MA) connected to the external TTL input
of the laser diode. Pulse amplitude was adjusted between 40-90 mA using the Laser Diode
Module Driver (Doric). A contraction was considered to be triggered if it occurred within 50ms of
stimulation. We performed photo-stimulation from 2-4 sites within each vessel, with 6-14
stimulations per site. If a photo-stimulation was triggered incidentally after the initiation of a
"spontaneous contraction" it was discarded from the analysis. For $Myh11CreER^{T2}$-ChR2-
tdTomato 6 vessels from 3 separate mice were tested. For $PdgfraCreER^{TM}$-ChR2-tdTomato 6
vessels from 4 separate mice were tested with a max of two vessels per mouse. For $cKitCreER^{T2}$-
ChR2-tdTomato 7 vessels from 3 separate mice were assessed. Diameter was recorded to align
photo-activation with the contraction cycle in a custom Labview program.
887
Solutions and Chemicals. Krebs buffer was composed of (in mM): 146.9 NaCl, 4.7 KCl, 2 $CaCl_2$,
1.2 $MgSO_4$, 1.2 $NaH_2PO_4 \cdot H_2O$, 3 $NaHCO_3$, 1.5 NaHEPES, and 5 d-glucose (pH = 7.4 at 37°C). Krebs-
BSA buffer was prepared with the addition of 0.5% (w/v) bovine serum albumin (BSA) while Krebs
Ca^{2+}-free replaced $CaCl_2$ with 3mM EGTA. Tamoxifen was dissolved to 10mg/ml in a Safflower Oil-
Ethanol (95%-5% v/v) solution with rocking agitation, separated into aliquots, and stored at -20
°C. Wortmannin was dissolved in DMSO to a stock solution of 1 mM. Pinacidil was dissolved in
DMSO to a stock concentration of 1 µM. Nifedipine was dissolved in DMSO to a stock
concentration of 1 mM. All chemicals were obtained from Sigma (St. Louis, MO), with the
exception of BSA (US Biochemicals; Cleveland, OH), $MgSO_4$ and NaHEPES (Fisher Scientific;
Pittsburgh, PA).
898
Statistical Tests Statistical differences in the isobaric contractile tests for popliteal cLVs isolated
from $PdgfraCreER^{TM}$-Ano1fl/fl, $PdgfraCreER^{TM}$-Cx45fl/fl, and $PdgfraCreER^{TM}$-Cav1.2fl/fl mice
over the various contractile parameters were assessed via 1) two-way ANOVAs with Tukey's
multiple comparison tests data performed using Prism9 (Graphpad). Data are plotted as mean ±
SEM and significance determined at $p < 0.05$. We used a categorical Chi-squared statistical test
for the experiments assessing our ability to trigger a contraction with activation of ChR2 cells.
Ca^{2+} PTCL area and frequency was compared using 1-way ANOVA with Tukey's post-hoc test.
Significance was determined at a p value of < 0.05.

907
908 **Acknowledgements**
909 We would like to thank Stefan Offermanns for donation of the $Myh11CreER^{T2}$ mice, Klaus
910 Willecke for his donation of $Cx45^{f/f}$ mice, Dieter Sauer for his donation of $cKitCreER^{T2}$ mice,
911 Ralph Adams for his donation of the $Pdgfr\beta CreER^{T2}$ mice and to Young Hong (University of
912 Southern California) for his donation of the $Prox1-eGFP$ mice (Choi et al., 2011).
913

Author contributions: S.D. Zawieja, J.A. Castorena-Gonzalez and M. J. Davis conceived and designed the experiments. S.D. Zawieja, S.E. Broyhill, J.A. Castorena-Gonzalez, H.J. Kim, B. Drumm, M. Li, C.E. Norton, and M.J. Davis performed the studies, collected, and analyzed and interpreted the data. S.D. Zawieja and M.J. Davis drafted the manuscript and all authors participated in critically revising the manuscript for intellectual concepts and content. All authors have approved the final draft of the manuscript and their contributions qualify them as authors.

The authors declare no competing financial interests. This work was supported by NIH HL-122608 and HL-122578 to MJD, HL-143198 to SDZ, and HL-141143 and HL-168568 to JAC-G, and AHA CDA-931652 to CEN.

Aalkjaer, C., D. Boedtkjer, and V. Matchkov. 2011. Vasomotion - what is currently thought? *Acta Physiol (Oxf)*. 202:253-269.

Andrae, J., R. Gallini, and C. Betsholtz. 2008. Role of platelet-derived growth factors in physiology and medicine. *Genes Dev*. 22:1276-1312.

Baker, S.A., B.T. Drumm, D. Saur, G.W. Hennig, S.M. Ward, and K.M. Sanders. 2016. Spontaneous Ca(2+) transients in interstitial cells of Cajal located within the deep muscular plexus of the murine small intestine. *J Physiol*. 594:3317-3338.

Baker, S.A., S.J. Hwang, P.J. Blair, C. Sireika, L. Wei, S. Ro, S.M. Ward, and K.M. Sanders. 2021a. Ca(2+) transients in ICC-MY define the basis for the dominance of the corpus in gastric pacemaking. *Cell Calcium*. 99:102472.

Baker, S.A., W.A. Leigh, G. Del Valle, I.F. De Yturriaga, S.M. Ward, C.A. Cobine, B.T. Drumm, and K.M. Sanders. 2021b. Ca(2+) signaling driving pacemaker activity in submucosal interstitial cells of Cajal in the murine colon. *Elife*. 10.

Basciani, S., S. Mariani, M. Arizzi, M. Brama, A. Ricci, C. Betsholtz, C. Bondjers, G. Ricci, A. Catizone, M. Galdieri, G. Spera, and L. Gnessi. 2004. Expression of platelet-derived growth factor (PDGF) in the epididymis and analysis of the epididymal development in PDGF-A, PDGF-B, and PDGF receptor beta deficient mice. *Biol Reprod*. 70:168-177.

Benoit, J.N., D.C. Zawieja, A.H. Goodman, and H.J. Granger. 1989. Characterization of intact mesenteric lymphatic pump and its responsiveness to acute edemagenic stress. *Am J Physiol*. 257:H2059-2069.

Boedtkjer, D.M., V.V. Matchkov, E. Boedtkjer, H. Nilsson, and C. Aalkjaer. 2008. Vasomotion has chloride-dependency in rat mesenteric small arteries. *Pflugers Arch*. 457:389-404.

Bolte, S., and F.P. Cordelieres. 2006. A guided tour into subcellular colocalization analysis in light microscopy. *J Microsc*. 224:213-232.

Bridenbaugh, E.A., W. Wang, M. Srimushnam, W.E. Cromer, S.D. Zawieja, S.E. Schmidt, D.C. Jupiter, H.C. Huang, V. Van Buren, and D.C. Zawieja. 2013. An immunological fingerprint differentiates muscular lymphatics from arteries and veins. *Lymphat Res Biol*. 11:155-171.

Briggs Boedtkjer, D., J. Rumessen, U. Baandrup, M. Skov Mikkelsen, N. Telinius, H. Pilegaard, C. Aalkjaer, and V. Hjortdal. 2013. Identification of interstitial Cajal-like cells in the human thoracic duct. *Cells Tissues Organs*. 197:145-158.

Castorena-Gonzalez, J.A., J.P. Scallan, and M.J. Davis. 2018a. Methods for Assessing the Contractile Function of Mouse Lymphatic Vessels Ex Vivo. *Methods in molecular biology*. 1846:229-248.

Castorena-Gonzalez, J.A., S.D. Zawieja, M. Li, R.S. Srinivasan, A.M. Simon, C. de Wit, R. de la
Torre, L.A. Martinez-Lemus, G.W. Hennig, and M.J. Davis. 2018b. Mechanisms of
Connexin-Related Lymphedema. *Circ Res*. 123:964-985.
Chakraborty, R., F.Z. Saddouk, A.C. Carrao, D.S. Krause, D.M. Greif, and K.A. Martin. 2019.
Promoters to Study Vascular Smooth Muscle. *Arterioscler Thromb Vasc Biol*. 39:603-
612.
Chakraborty, S., S.D. Zawieja, W. Wang, Y. Lee, Y.J. Wang, P.Y. von der Weid, D.C. Zawieja,
and M. Muthuchamy. 2015. Lipopolysaccharide modulates neutrophil recruitment and
macrophage polarization on lymphatic vessels and impairs lymphatic function in rat
mesentery. *Am J Physiol Heart Circ Physiol*.ajpheart 00467 02015.
Chatterjee, V., and A.A. Gashev. 2012. Aging-associated shifts in functional status of mast cells
located by adult and aged mesenteric lymphatic vessels. *Am J Physiol Heart Circ*
Physiol. 303:H693-702.
Chen, T.W., T.J. Wardill, Y. Sun, S.R. Pulver, S.L. Renninger, A. Baohan, E.R. Schreiter, R.A.
Kerr, M.B. Orger, V. Jayaraman, L.L. Looger, K. Svoboda, and D.S. Kim. 2013.
Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature*. 499:295-300.
Cheng, X., J. Liu, M. Asuncion-Chin, E. Blaskova, J.P. Bannister, A.M. Dopico, and J.H. Jaggar.
2007. A novel Ca(V)1.2 N terminus expressed in smooth muscle cells of resistance size
arteries modifies channel regulation by auxiliary subunits. *J Biol Chem*. 282:29211-
29221.
Choi, I., H.K. Chung, S. Ramu, H.N. Lee, K.E. Kim, S. Lee, J. Yoo, D. Choi, Y.S. Lee, B.
Aguilar, and Y.K. Hong. 2011. Visualization of lymphatic vessels by Prox1-promoter
directed GFP reporter in a bacterial artificial chromosome-based transgenic mouse.
Blood. 117:362-365.
Clayton, D.R., W.G. Ruiz, M.G. Dalghi, N. Montalbetti, M.D. Carattino, and G. Apodaca. 2022.
Studies of ultrastructure, gene expression, and marker analysis reveal that mouse bladder
PDGFRA(+) interstitial cells are fibroblasts. *Am J Physiol Renal Physiol*. 323:F299-
F321.
Cobine, C.A., E.E. Hannah, M.H. Zhu, H.E. Lyle, J.R. Rock, K.M. Sanders, S.M. Ward, and
K.D. Keef. 2017. ANO1 in intramuscular interstitial cells of Cajal plays a key role in the
generation of slow waves and tone in the internal anal sphincter. *J Physiol*. 595:2021-
2041.
Davis, M.J., J.A. Castorena-Gonzalez, H.J. Kim, M. Li, M. Remedi, and C.G. Nichols. 2023.
Lymphatic contractile dysfunction in mouse models of Cantu Syndrome with K(ATP)
channel gain-of-function. *Function (Oxf)*. 4:zqad017.
Davis, M.J., H.J. Kim, M. Li, and S.D. Zawieja. 2022. A vascular smooth muscle-specific
integrin-alpha8 Cre mouse for lymphatic contraction studies that allows male-female
comparisons and avoids visceral myopathy. *Front Physiol*. 13:1060146.
Davis, M.J., H.J. Kim, S.D. Zawieja, J.A. Castorena-Gonzalez, P. Gui, M. Li, B.T. Saunders,
B.H. Zinselmeyer, G.J. Randolph, M.S. Remedi, and C.G. Nichols. 2020. Kir6.1-
dependent KATP channels in lymphatic smooth muscle and vessel dysfunction in mice
with Kir6.1 gain-of-function. *J Physiol*.
Davis, M.J., E. Rahbar, A.A. Gashev, D.C. Zawieja, and J.E. Moore, Jr. 2011. Determinants of
valve gating in collecting lymphatic vessels from rat mesentery. *Am J Physiol Heart Circ*
Physiol. 301:H48-60.

Davis, M.J., J.P. Scallan, J.H. Wolpers, M. Muthuchamy, A.A. Gashev, and D.C. Zawieja. 2012.
Intrinsic increase in lymphangion muscle contractility in response to elevated afterload.
Am J Physiol Heart Circ Physiol. 303:H795-808.
Drumm, B.T., G.W. Hennig, S.A. Baker, and K.M. Sanders. 2019. Applications of spatio-
temporal mapping and particle analysis techniques to quantify intracellular Ca 2+
signaling in situ. *Journal of Visualized Experiments*. 2019:1-13.
Drumm, B.T., G.W. Hennig, M.J. Battersby, E.K. Cunningham, T.S. Sung, S.M. Ward, K.M.
Sanders, and S.A. Baker. 2017. Clustering of Ca2+ transients in interstitial cells of Cajal
defines slow wave duration. *The Journal of General Physiology*. 149:703-725.
Drumm, B.T., B.E. Rembetski, C.A. Cobine, S.A. Baker, G.P. Sergeant, M.A. Hollywood, K.D.
Thornbury, and K.M. Sanders. 2018. Ca(2+) signalling in mouse urethral smooth muscle
in situ: role of Ca(2+) stores and Ca(2+) influx mechanisms. *J Physiol*. 596:1433-1466.
Engeset, A., W. Olszewski, P.M. Jaeger, J. Sokolowski, and L. Theodorsen. 1977. Twenty-four
hour variation in flow and composition of leg lymph in normal men. *Acta physiologica
Scandinavica*. 99:140-148.
Ferrusi, I., J. Zhao, D. van Helden, and P.Y. von der Weid. 2004. Cyclopiazonic acid decreases
spontaneous transient depolarizations in guinea pig mesenteric lymphatic vessels in
endothelium-dependent and -independent manners. *Am J Physiol Heart Circ Physiol*.
286:H2287-2295.
Gaengel, K., G. Genove, A. Armulik, and C. Betsholtz. 2009. Endothelial-mural cell signaling in
vascular development and angiogenesis. *Arterioscler Thromb Vasc Biol*. 29:630-638.
Gashev, A.A., M.J. Davis, M.D. Delp, and D.C. Zawieja. 2004. Regional variations of
contractile activity in isolated rat lymphatics. *Microcirculation*. 11:477-492.
Geng, X., B. Cha, M.R. Mahamud, K.C. Lim, R. Silasi-Mansat, M.K.M. Uddin, N. Miura, L.
Xia, A.M. Simon, J.D. Engel, H. Chen, F. Lupu, and R.S. Srinivasan. 2016. Multiple
mouse models of primary lymphedema exhibit distinct defects in lymphovenous valve
development. *Dev Biol*. 409:218-233.
Gerl, K., L. Miquerol, V.T. Todorov, C.P. Hugo, R.H. Adams, A. Kurtz, and B. Kurt. 2015.
Inducible glomerular erythropoietin production in the adult kidney. *Kidney Int*. 88:1345-
1355.
Gomez-Pinilla, P.J., S.J. Gibbons, M.R. Bardsley, A. Lorincz, M.J. Pozo, P.J. Pasricha, M. Van
de Rijn, R.B. West, M.G. Sarr, M.L. Kendrick, R.R. Cima, E.J. Dozois, D.W. Larson, T.
Ordog, and G. Farrugia. 2009. Ano1 is a selective marker of interstitial cells of Cajal in
the human and mouse gastrointestinal tract. *Am J Physiol Gastrointest Liver Physiol*.
296:G1370-1381.
Grainger, N., C.C. Shonnard, S.K. Quiggle, E.B. Fox, H. Presley, R. Daugherty, M.C. Shonnard,
B.T. Drumm, and K.M. Sanders. 2022. Propagation of Pacemaker Activity and Peristaltic
Contractions in the Mouse Renal Pelvis Rely on Ca(2+)-activated Cl(-) Channels and T-
Type Ca(2+) Channels. *Function (Oxf)*. 3:zqac041.
Hald, B.O., J.A. Castorena-Gonzalez, S.D. Zawieja, P. Gui, and M.J. Davis. 2018. Electrical
Communication in Lymphangions. *Biophys J*. 115:936-949.
Hashitani, H., R. Mitsui, S. Masaki, and D.F. Van Helden. 2015. Pacemaker role of pericytes in
generating synchronized spontaneous Ca2+ transients in the myenteric microvasculature
of the guinea-pig gastric antrum. *Cell Calcium*. 58:442-456.
Heger, K., B. Seidler, J.C. Vahl, C. Schwartz, M. Kober, S. Klein, D. Voehringer, D. Saur, and
M. Schmidt-Supprian. 2014. CreER(T2) expression from within the c-Kit gene locus

allows efficient inducible gene targeting in and ablation of mast cells. *Eur J Immunol.* 44:296-306.

Hill, R.A., L. Tong, P. Yuan, S. Murikinati, S. Gupta, and J. Grutzendler. 2015. Regional Blood Flow in the Normal and Ischemic Brain Is Controlled by Arteriolar Smooth Muscle Cell Contractility and Not by Capillary Pericytes. *Neuron.* 87:95-110.

Huizinga, J.D., L. Thuneberg, M. Kluppel, J. Malysz, H.B. Mikkelsen, and A. Bernstein. 1995. W/kit gene required for interstitial cells of Cajal and for intestinal pacemaker activity. *Nature.* 373:347-349.

Hwang, S.J., P.J. Blair, F.C. Britton, K.E. O'Driscoll, G. Hennig, Y.R. Bayguinov, J.R. Rock, B.D. Harfe, K.M. Sanders, and S.M. Ward. 2009. Expression of anoctamin 1/TMEM16A by interstitial cells of Cajal is fundamental for slow wave activity in gastrointestinal muscles. *J Physiol.* 587:4887-4904.

Imtiaz, M.S., J. Zhao, K. Hosaka, P.Y. von der Weid, M. Crowe, and D.F. van Helden. 2007. Pacemaking through Ca2+ stores interacting as coupled oscillators via membrane depolarization. *Biophys J.* 92:3843-3861.

Jolly, A.J., S. Lu, K.A. Strand, A.M. Dubner, M.F. Mutryn, R.A. Nemenoff, M.W. Majesky, K.S. Moulton, and M.C.M. Weiser-Evans. 2022. Heterogeneous subpopulations of adventitial progenitor cells regulate vascular homeostasis and pathological vascular remodelling. *Cardiovasc Res.* 118:1452-1465.

Kang, S.H., M. Fukaya, J.K. Yang, J.D. Rothstein, and D.E. Bergles. 2010. NG2+ CNS glial progenitors remain committed to the oligodendrocyte lineage in postnatal life and following neurodegeneration. *Neuron.* 68:668-681.

Kenney, H.M., R.D. Bell, E.A. Masters, L. Xing, C.T. Ritchlin, and E.M. Schwarz. 2020. Lineage tracing reveals evidence of a popliteal lymphatic muscle progenitor cell that is distinct from skeletal and vascular muscle progenitors. *Sci Rep.* 10:18088.

Kimura, K., K. Ramirez, T.A.V. Nguyen, Y. Yamashiro, A. Sada, and H. Yanagisawa. 2021. Contribution of PDGFRalpha-positive cells in maintenance and injury responses in mouse large vessels. *Sci Rep.* 11:8683.

Koval, M., S.A. Molina, and J.M. Burt. 2014. Mix and match: investigating heteromeric and heterotypic gap junction channels in model systems and native tissues. *FEBS Lett.* 588:1193-1204.

Kramann, R., R.K. Schneider, D.P. DiRocco, F. Machado, S. Fleig, P.A. Bondzie, J.M. Henderson, B.L. Ebert, and B.D. Humphreys. 2015. Perivascular Gli1+ progenitors are key contributors to injury-induced organ fibrosis. *Cell Stem Cell.* 16:51-66.

Kurahashi, M., Y. Nakano, L.E. Peri, J.B. Townsend, S.M. Ward, and K.M. Sanders. 2013. A novel population of subepithelial platelet-derived growth factor receptor alpha-positive cells in the mouse and human colon. *Am J Physiol Gastrointest Liver Physiol.* 304:G823-834.

Kurahashi, M., H. Zheng, L. Dwyer, S.M. Ward, S.D. Koh, and K.M. Sanders. 2011. A functional role for the 'fibroblast-like cells' in gastrointestinal smooth muscles. *J Physiol.* 589:697-710.

Lee, Y., S. Chakraborty, and M. Muthuchamy. 2020. Roles of sarcoplasmic reticulum Ca(2+) ATPase pump in the impairments of lymphatic contractile activity in a metabolic syndrome rat model. *Sci Rep.* 10:12320.

Li, S.W., C.Y. Wang, Y.J. Jou, T.C. Yang, S.H. Huang, L. Wan, Y.J. Lin, and C.W. Lin. 2016.
SARS coronavirus papain-like protease induces Egr-1-dependent up-regulation of TGF-
beta1 via ROS/p38 MAPK/STAT3 pathway. *Sci Rep*. 6:25754.
Madisen, L., T. Mao, H. Koch, J.M. Zhuo, A. Berenyi, S. Fujisawa, Y.W. Hsu, A.J. Garcia, 3rd,
X. Gu, S. Zanella, J. Kidney, H. Gu, Y. Mao, B.M. Hooks, E.S. Boyden, G. Buzsaki, J.M.
Ramirez, A.R. Jones, K. Svoboda, X. Han, E.E. Turner, and H. Zeng. 2012. A toolbox of
Cre-dependent optogenetic transgenic mice for light-induced activation and silencing.
Nat Neurosci. 15:793-802.
Maeda, H., A. Yamagata, S. Nishikawa, K. Yoshinaga, S. Kobayashi, K. Nishi, and S.
Nishikawa. 1992. Requirement of c-kit for development of intestinal pacemaker system.
Development. 116:369-375.
Majesky, M.W., H. Horita, A. Ostriker, S. Lu, J.N. Regan, A. Bagchi, X.R. Dong, J. Poczobutt,
R.A. Nemenoff, and M.C. Weiser-Evans. 2017. Differentiated Smooth Muscle Cells
Generate a Subpopulation of Resident Vascular Progenitor Cells in the Adventitia
Regulated by Klf4. *Circ Res*. 120:296-311.
Malysz, J., S.J. Gibbons, S.A. Saravanaperumal, P. Du, S.T. Eisenman, C. Cao, U. Oh, D. Saur,
S. Klein, T. Ordog, and G. Farrugia. 2017. Conditional genetic deletion of Ano1 in
interstitial cells of Cajal impairs Ca(2+) transients and slow waves in adult mouse small
intestine. *Am J Physiol Gastrointest Liver Physiol*. 312:G228-G245.
McCloskey, K.D., M.A. Hollywood, K.D. Thornbury, S.M. Ward, and N.G. McHale. 2002. Kit-
like immunopositive cells in sheep mesenteric lymphatic vessels. *Cell Tissue Res*.
310:77-84.
Mitsui, R., and H. Hashitani. 2020. Synchrony of spontaneous Ca(2+) activity in microvascular
mural cells. *Journal of smooth muscle research = Nihon Heikatsukin Gakkai kikanshi*.
56:1-18.
Mohanakumar, S., J. Majgaard, N. Telinius, N. Katballe, E. Pahle, V.E. Hjortdal, and D.M.B.
Boedtkjer. 2018. Spontaneous and alpha-adrenoceptor-induced contractility in human
collecting lymphatic vessels require chloride. *Am J Physiol Heart Circ Physiol*.
Muzumdar, M.D., B. Tasic, K. Miyamichi, L. Li, and L. Luo. 2007. A global double-fluorescent
Cre reporter mouse. *Genesis*. 45:593-605.
Olszewski, W.L. 2002. Contractility patterns of normal and pathologically changed human
lymphatics. *Ann N Y Acad Sci*. 979:52-63; discussion 76-59.
Olszewski, W.L. 2008. Contractility patterns of human leg lymphatics in various stages of
obstructive lymphedema. *Ann N Y Acad Sci*. 1131:110-118.
Ordog, T., S.M. Ward, and K.M. Sanders. 1999. Interstitial cells of cajal generate electrical slow
waves in the murine stomach. *J Physiol*. 518:257-269.
Pitulescu, M.E., and R.H. Adams. 2014. Regulation of signaling interactions and receptor
endocytosis in growing blood vessels. *Cell Adh Migr*. 8:366-377.
Popescu, L.M., and M.S. Faussone-Pellegrini. 2010. TELOCYTES - a case of serendipity: the
winding way from Interstitial Cells of Cajal (ICC), via Interstitial Cajal-Like Cells
(ICLC) to TELOCYTES. *J Cell Mol Med*. 14:729-740.
Sanders, K.M., S.A. Baker, B.T. Drumm, and M. Kurahashi. 2022. Ca(2+) Signaling Is the Basis
for Pacemaker Activity and Neurotransduction in Interstitial Cells of the GI Tract. *Adv
Exp Med Biol*. 1383:229-241.
Sanders, K.M., T. Ordog, S.D. Koh, S. Torihashi, and S.M. Ward. 1999. Development and
plasticity of interstitial cells of Cajal. *Neurogastroenterol Motil*. 11:311-338.

Sanders, K.M., S.M. Ward, and S.D. Koh. 2014. Interstitial cells: regulators of smooth muscle
function. *Physiol Rev*. 94:859-907.
Sasmono, R.T., D. Oceandy, J.W. Pollard, W. Tong, P. Pavli, B.J. Wainwright, M.C. Ostrowski,
S.R. Himes, and D.A. Hume. 2003. A macrophage colony-stimulating factor receptor-
green fluorescent protein transgene is expressed throughout the mononuclear phagocyte
system of the mouse. *Blood*. 101:1155-1163.
Scallan, J.P., and M.J. Davis. 2013. Genetic removal of basal nitric oxide enhances contractile
activity in isolated murine collecting lymphatic vessels. *J Physiol*. 591:2139-2156.
Scallan, J.P., S.D. Zawieja, J.A. Castorena-Gonzalez, and M.J. Davis. 2016. Lymphatic
pumping: Mechanics, mechanisms and malfunction. *J Physiol*.
Sidney, L.E., M.J. Branch, S.E. Dunphy, H.S. Dua, and A. Hopkinson. 2014. Concise review:
evidence for CD34 as a common marker for diverse progenitors. *Stem Cells*. 32:1380-
1389.
Singh, R.D., S.J. Gibbons, S.A. Saravanaperumal, P. Du, G.W. Hennig, S.T. Eisenman, A.
Mazzone, Y. Hayashi, C. Cao, G.J. Stoltz, T. Ordog, J.R. Rock, B.D. Harfe, J.H.
Szurszewski, and G. Farrugia. 2014. Ano1, a Ca2+-activated Cl- channel, coordinates
contractility in mouse intestine by Ca2+ transient coordination between interstitial cells
of Cajal. *J Physiol*. 592:4051-4068.
Song, K., Y. Qing, Q. Guo, E.K. Peden, C. Chen, W.E. Mitch, L. Truong, and J. Cheng. 2020.
PDGFRA in vascular adventitial MSCs promotes neointima formation in arteriovenous
fistula in chronic kidney disease. *JCI Insight*. 5.
Spandidos, A., X. Wang, H. Wang, S. Dragnev, T. Thurber, and B. Seed. 2008. A
comprehensive collection of experimentally validated primers for Polymerase Chain
Reaction quantitation of murine transcript abundance. *BMC Genomics*. 9:633.
Spandidos, A., X. Wang, H. Wang, and B. Seed. 2010. PrimerBank: a resource of human and
mouse PCR primer pairs for gene expression detection and quantification. *Nucleic Acids
Res*. 38:D792-799.
Stolarz, A.J., M. Sarimollaoglu, J.C. Marecki, T.W. Fletcher, E.I. Galanzha, S.W. Rhee, V.P.
Zharov, V.S. Klimberg, and N.J. Rusch. 2019. Doxorubicin Activates Ryanodine
Receptors in Rat Lymphatic Muscle Cells to Attenuate Rhythmic Contractions and
Lymph Flow. *J Pharmacol Exp Ther*. 371:278-289.
To, K.H.T., P. Gui, M. Li, S.D. Zawieja, J.A. Castorena-Gonzalez, and M.J. Davis. 2020. T-type,
but not L-type, voltage-gated calcium channels are dispensable for lymphatic pacemaking
and spontaneous contractions. *Sci Rep*. 10:70.
Toland, H.M., K.D. McCloskey, K.D. Thornbury, N.G. McHale, and M.A. Hollywood. 2000.
Ca(2+)-activated Cl(-) current in sheep lymphatic smooth muscle. *Am J Physiol Cell
Physiol*. 279:C1327-1335.
Torihashi, S., S.M. Ward, S. Nishikawa, K. Nishi, S. Kobayashi, and K.M. Sanders. 1995. c-kit-
dependent development of interstitial cells and electrical activity in the murine
gastrointestinal tract. *Cell Tissue Res*. 280:97-111.
Truett, G.E., P. Heeger, R.L. Mynatt, A.A. Truett, J.A. Walker, and M.L. Warman. 2000.
Preparation of PCR-quality mouse genomic DNA with hot sodium hydroxide and tris
(HotSHOT). *Biotechniques*. 29:52, 54.
Van Helden, D.A., P.Y. Von Der Weid, and M. Crowe. 1996. Intracellular Ca Release: A basis
for electrical pacemaking in lymphatic smooth muscle. *In* Smooth muscle excitation. T.B.
Ed. Bolton and T. Tomita, editors. London: Academic Press. 355–373.

Van Helden, D.F. 1993. Pacemaker potentials in lymphatic smooth muscle of the guinea-pig mesentery. *J Physiol*. 471:465-479.

van Helden, D.F., K. Hosaka, and M.S. Imtiaz. 2006. Rhythmicity in the microcirculation. *Clin Hemorheol Microcirc*. 34:59-66.

van Helden, D.F., and M.S. Imtiaz. 2019. Venous Vasomotion. *Adv Exp Med Biol*. 1124:313-328.

Van Helden, D.F., and J. Zhao. 2000. Lymphatic vasomotion. *Clin Exp Pharmacol Physiol*. 27:1014-1018.

Van, S., S. Pal, B.R. Garner, K. Steed, V. Sridharan, S. Mu, N.J. Rusch, and A.J. Stolarz. 2021. Dantrolene Prevents the Lymphostasis Caused by Doxorubicin in the Rat Mesenteric Circulation. *Front Pharmacol*. 12:727526.

Vannucchi, M.G., C. Traini, M. Manetti, L. Ibba-Manneschi, and M.S. Faussone-Pellegrini. 2013. Telocytes express PDGFRalpha in the human gastrointestinal tract. *J Cell Mol Med*. 17:1099-1108.

von der Weid, P.Y., M. Rahman, M.S. Imtiaz, and D.F. van Helden. 2008. Spontaneous transient depolarizations in lymphatic vessels of the guinea pig mesentery: pharmacology and implication for spontaneous contractility. *Am J Physiol Heart Circ Physiol*. 295:H1989-2000.

Wang, X., and B. Seed. 2003. A PCR primer bank for quantitative gene expression analysis. *Nucleic Acids Res*. 31:e154.

Wang, Y., Y. Jin, M.A. Mae, Y. Zhang, H. Ortsater, C. Betsholtz, T. Makinen, and L. Jakobsson. 2017. Smooth muscle cell recruitment to lymphatic vessels requires PDGFB and impacts vessel size but not identity. *Development*. 144:3590-3601.

Ward, S.M., A.J. Burns, S. Torihashi, and K.M. Sanders. 1994. Mutation of the proto-oncogene c-kit blocks development of interstitial cells and electrical rhythmicity in murine intestine. *J Physiol*. 480 (Pt 1):91-97.

Warthi, G., J.L. Faulkner, J. Doja, A.R. Ghanam, P. Gao, A.C. Yang, O.J. Slivano, C.T. Barris, T.C. Kress, S.D. Zawieja, S.H. Griffin, X. Xie, A. Ashworth, C.K. Christie, W.B. Bryant, A. Kumar, M.J. Davis, X. Long, L. Gan, E.J. Belin de Chantemele, Q. Lyu, and J.M. Miano. 2022. Generation and Comparative Analysis of an Itga8-CreER (T2) Mouse with Preferential Activity in Vascular Smooth Muscle Cells. *Nat Cardiovasc Res*. 1:1084-1100.

Wirth, A., Z. Benyo, M. Lukasova, B. Leutgeb, N. Wettschureck, S. Gorbey, P. Orsy, B. Horvath, C. Maser-Gluth, E. Greiner, B. Lemmer, G. Schutz, J.S. Gutkind, and S. Offermanns. 2008. G12-G13-LARG-mediated signaling in vascular smooth muscle is required for salt-induced hypertension. *Nat Med*. 14:64-68.

Xiao, J., F. Wang, Z. Liu, and C. Yang. 2013. Telocytes in liver: electron microscopic and immunofluorescent evidence. *J Cell Mol Med*. 17:1537-1542.

Zawieja, D.C., K.L. Davis, R. Schuster, W.M. Hinds, and H.J. Granger. 1993. Distribution, propagation, and coordination of contractile activity in lymphatics. *Am J Physiol*. 264:H1283-1291.

Zawieja, S.D., J.A. Castorena, P. Gui, M. Li, S.A. Bulley, J.H. Jaggar, J.R. Rock, and M.J. Davis. 2019. Ano1 mediates pressure-sensitive contraction frequency changes in mouse lymphatic collecting vessels. *J Gen Physiol*. 151:532-554.

Zawieja, S.D., J.A. Castorena-Gonzalez, J. Scallan, and M.J. Davis. 2018. Differences in L-type
calcium channel activity partially underlie the regional dichotomy in pumping behavior
by murine peripheral and visceral lymphatic vessels. *Am J Physiol Heart Circ Physiol*.
Zawieja, S.D., W. Wang, S. Chakraborty, D.C. Zawieja, and M. Muthuchamy. 2016.
Macrophage alterations within the mesenteric lymphatic tissue are associated with
impairment of lymphatic pump in metabolic syndrome. *Microcirculation*. 23:558-570.
Zhou, Q., L. Wei, C. Zhong, S. Fu, Y. Bei, R.I. Huica, F. Wang, and J. Xiao. 2015. Cardiac
telocytes are double positive for CD34/PDGFR-alpha. *J Cell Mol Med*. 19:2036-2042.
Zhu, M.H., T.W. Kim, S. Ro, W. Yan, S.M. Ward, S.D. Koh, and K.M. Sanders. 2009. A
Ca(2+)-activated Cl(-) conductance in interstitial cells of Cajal linked to slow wave
currents and pacemaker activity. *J Physiol*. 587:4905-4918.
Zhu, X., D.E. Bergles, and A. Nishiyama. 2008. NG2 cells generate both oligodendrocytes and
gray matter astrocytes. *Development*. 135:145-157.
1244
1245

Table 1 Primer list for RT-PCR

Gene	Strand	Accession #	Sequence(5'-3')	Siza	Exon	Source
Prox1	s	NM_008937	GTA AGA CAT CAC CGC GTG C	218	1	NIH Primer Tool
	as		TCA TGG TCA GGC ATC ACT GG		2	
CD11b (Itgam)	s	NM_008401	ATG GAC GCT GAT GGC AAT ACC	203	13	MGH Primer Bank ID 668048a1
	as		TCC CCA TTC ACG TCT CCC A		14	
Pdgfrα	s	NM_011058	AGA GTT ACA CGT TTG AGC TGT C	252	8	MGH Primer Bank 26349287a1
	as		GTC CCT CCA CGG TAC TCC T		10	
						-
Myh11	s	NM_013607	AAG CTG CGG CTA GAG GTC A	238	33	MGH Primer Bank ID 7305295a1
	as		CCC TCC CTT TGA TGG CTG AG		34	
CD117 (cKit)	s	NM_021099	CGC CTG CCG AAA TGT ATG ACG	162	21	(Drumm et al., 2018)
	as		GGT TCT CTG GGT TGG GGT TGC		23	
Pdgfrβ	s	NM_008809	AGC TAC ATG GCC CCT TAT GA	367	16	(Basciani et al., 2004)
	as		GGA TCC AAA AAG ACC AGA CA		19	
CD144 (VE-cadherin)	s	NM_009868	CTT CCT TAC TGC CCT CAT TGT	313	3	IDT Primer Quest
	as		CTG TTT CTC TCG GTC CAA GTT		5	
Nos3 (eNOS)	s	NM_008713	ctg cca cct gat cct aac ttg	143	22	IDT Real time primer tool
	as		cag cca aac acc aaa gtc atg		23	
Acta2 (Smooth Muscle Actin)	s	NM_007392	GAG CTA CGA ACT GCC TGA C	129	7	IDT TaqMan Mm.PT.58.16320644
	as		CTG TTA TAG GTG GTT TCG TGG A		8	
CaV 1.2 exon1b	s	NM_001159533	ATG GTC AAT GAA AAC ACG AGG ATG		1	(Cheng et al., 2007)

	as		GGA ACT GAC GGT AGA GAT GGT TGC	234	2	
CD34	as	NM_001111059	GGT ACA GGA GAA TGC AGG TC	119	2	IDT Mm.PT.58.8626728
	s	NM_133654	CGT GGT AGC AGA AGT CAA GT		1	
Cspg4 (Ng2)	as	NM_139001	CTT CAC GAT CAC CAT CCT TCC	132	5	IDT Mm.PT.58.29461721
	s		CCC GAA TCA TTG TCT GTT CCC		6	
-	-	-	-	-	-	
Vimentin	s	NM_011701	CTG TAC GAG GAG GAG ATG CG	249	1	(Li et al., 2016)
	as		AAT TTC TTC CTG CAA GGA TT		3	
Desmin	s	NM_010043	GTG GAT GCA GCC ACT CTA GC	218	3	MGH Primer Bank ID 33563250a1
	as		TTA GCC GCG ATG GTC TCA TAC		4	
CD146 (Mcam)	s	NM_023061	CCC AAA CTG GTG TGC GTC TT	220	1	MGH Primer Bank 10566955a1
	as		GGA AAA TCA GTA TCT GCC TCT CC		3	
KLF4	s	NM_010637	ATT AAT GAG GCA GCC ACC TG	400	1	(Majesky et al., 2017)
	as		GGA AGA CGA GGA TGA AGC TG		3	
Ly6a (Sca1)	s	NM_001271416	CTC TGA GGA TGG ACA CTT CT	400	2	(Majesky et al., 2017)
	as		GGT CTG CAG GAG GAC TGA GC		4	
Gli1	s	NM_01029	ATC ACC TGT TGG GGA TGC TGG AT	316	8	(Kramann et al., 2015)
	as		CGT GAA TAG GAC TTC CGA CAG		10	
CD29 (Itgb1)	s	NM_010578	TCG ATC CTG TGA CCC ATT GC	170	14	NIH Primer Tool
	as		AAC AAT TCC AGC AAC CAC GC		15	

CD105 (Endoglin)	s	NM_007932	TGA GCG TGT CTC CAT TGA CC	416	11	NIH Primer Tool
	as		GGG GCC ACG TGT GTG AGA A		15	
CD44	as	NM_009851	CAC CAT TTC CTG AGA CTT GCT	148	19	IDT Mm.PT.58.12084136
	s		TCT GAT TCT TGC CGT CTG C		18	
CD31 (Pecam1)	s	NM_008816	CTG CCA GTC CGA AAA TGG AAC	218	7	MGH Primer Bank ID 6679273a1
	as		CTT CAT CCA CTG GGG CTA TC		8	
GJC1 (Connexin 45)	s	NM_008122	GGT AAC AGG AGT TCT GGT GAA	140	2	IDT Mm.PT.58.8383900
	as		TCG AAA GAC AAT CAG CAC AGT		3	
Anoctamin 1 (TMEM16A)	s	NM_178642	ggc att tgt cat tgt ctt cca g	141	25	IDT Real time primer tool
	as		tcc tca cgc ata aac agc tc		26	
CD45 (Ptprc)	s	NM_001111316	ATG CAT CCA TCC TCG TCC AC	225	29	NIH Primer Tool
	as		TGA CTT GTC CAT TCT GGG CG		31	
MGH Harvard Primer Bank (Wang and Seed, 2003; Spandidos et al., 2008; Spandidos et al., 2010)						

1246
1247
1248

Figure 1 Methylene blue staining of isolated mouse IALVs

(A) Representative image of an isolated and cleaned IALV after methylene blue staining which revealed cells of various morphology. (B) is the zoomed in image of the yellow dotted box in A which contained large ovoid cells with granular staining (B, yellow asterisks). Fine cellular extensions (red asterisks) stained by methylene blue in some cells were visualized with color channel separation and division (C). (D, E) Similar as B and C, but in a separate vessel which stained with a higher density of methylene blue stained cells some of which had limited cellular processes. F) Focal reconstruction from imaging a methylene blue stained vessel using an upright microscope and immersion objective.

1261
Figure 2 Staining Murine IALVs for ICLC Markers
Representative immunofluorescent max projections of half vessel confocal image stacks imaged
from mouse IALVs stained for ICLC markers. DAPI (A), cKit (B), and CD34 (C) and their merged

image (D). Representative max projections of the intermediate filament vimentin (E), the
intermediate filament desmin (F), CD34 (G) and their merged image (H). Representative max
projection of vimentin (I), cKit (J), CD34 (K) and their merged image (L). Scale bar = 100 μm for
all images.
1269
1270

1271
Figure 3 Immunofluorescent Labeling of Murine IALVs With Markers for ICLC, LMC, LEC, and
Immune Cell Populations

We stained isolated mouse IALVs with cellular markers used to differentiate various cell types
observed in cLVs. Half vessel image stacks were taken with confocal microscopy and the
resulting representative max projections are shown. (A) CD34 stained cells and LMC staining
with SMA (B) and calponin (C) and the corresponding merged (D) image. Significant overlap in
(E) CD34 staining along with the fibroblast marker PDGFRα compared to LMC staining with SMA
(G) and the merged (H) image. The endothelial marker CD31 (I) to delineate LECs with PDGFRα
staining (J), and the LMC marker calponin (K) with the merged image (L). Mononuclear
phagocyte cells were identified by anti-GFP (M) in IALVs from MacGreen mice and counter
stained with PDGFRα (N) and the hematopoietic marker CD45 (O) with (P) the merged image.
PDGFRβ (S) stained many cells that were CD34 (Q) and PDGFRα (R) positive in addition to signal
detected in the LMC layer (U). Max projections of only the luminal frames of a z-stack at
lymphatic valve locations revealed PDGFRβ, CD34, and PDGFRα labeling in bipolar shaped cells
with long extensions that traveled throughout the valve leaflets (V, W). Secondary only stained
control IALV (Y). Scale bar = 100 µm for all images.
1288
1289
1290

1291
Figure 4 PDGFRα⁺ Cells Reside Primarily in the Murine Lymphatic Collecting Vessel Adventitia and
Some in the Subendothelial Space
Max projection of confocal imaging of an IALV stained for LECs with CD31 (A), LMCs with
MYH11(B), and for PDGFRα (C) with the corresponding merge file (D). Orthogonal views of the
z-stack with (E) showing a single slice in the z stack and E' and E" the orthogonal views. White
dotted boxes outline locations where PDGFRα signal is observed between LMC and LEC layers.
Scale bar is 100 μm in (D) and 50 μm in (E).
1299
1300

1301
Figure 5 Colocalization of CD34 and PDGFRα
Representative max projections and their corresponding threshold adjusted images for
colocalization analysis for PDGFRα (A), CD34 (B), and their colocalized signal (C). Pearson's
coefficient and Mander's coefficients were calculated from 3 separate stained IALVS, each from a
separate mouse.

1309

Figure 6 iCre-ROSA26mTmG Labelling and Fidelity to Target Putative Pacemaker Cell Populations

Stitched montages of serial max projections of GFP and tdTomato signal from live IALVs isolated from *PdgfrαCre*-ROSA26mTmG (A), *Ng2Cre*-ROSA26mTmG (B), *PdgfrαCreER^{TM}*-ROSA26mTmG (C), *Pdgfrβ CreER^{T2}*-ROSA26mTmG (D), *cKitCreER^{T2}*-ROSA26mTmG (E), and *Myh11CreER^{T2}*-ROSA26mTmG (F). IALVs were digested into single cells and GFP$^+$ cells were purified via FACS from *Prox1-eGFP* (G), *Myh11CreER^{T2}*-ROSA26mTmG (H), *PdgfrαCreER^{TM}*-ROSA26mTmG (I), and *PdgfrβCreER^{T2}*-ROSA26mTmG (J) mice. Representative gels demonstrating RT-PCR products corresponding to the respective gene used in the promoter of the transgenes to drive GFP signal or Cre mediated recombination of ROSA26mTmG from each GFP$^+$ sorted population (K) to assess fidelity. Images are representative of IALVs from at least 3 separate mice. FACs and RT-PCR was repeated at least 3 times for each mouse.

1323
Figure 7 RT-PCR Profiling of FACs Sorted Cells from iCre-ROSA26mTmG
Expanded RT-PCR profiling of genes to discriminate LECs, LMCs, and other cell types in our GFP⁺
sorted cells from *Prox1-eGFP* (A), *Myh11CreER^T2^*-ROSA26mTmG (B), *PdgfrβCreER^T2^*-
ROSA26mTmG (C), and *PdgfrαCreER^TM^*-ROSA26mTmG (D). RT-PCR results for CD31, Cx45, Ano1,
and CD45 in GFP⁺ cells sorted from *PdgfrαCreER^TM^*-ROSA26mTmG IALVs. RT-PCR was repeated
at 2-4 times for each gene over each sorted cell population collected from different mice.
1330
1331

1332

Figure 8 Isobaric contractile Assessment of popliteal cLV from *PdgfraCre*TM-Ano1 fl/fl mice
Summary of the contractile parameters recorded from popliteal cLVs in *PdgfraCreER^TM*-Ano1fl/fl mice. Normalized contraction amplitude (A), ejection fraction (B), contraction frequency (C), fractional pump flow (D), end diastolic diameter (E), and vessel tone (F) were assessed. No statically significant differences observed. Mean and SEM shown, n=6 popliteal vessels from 3 mice *PdgfraCreER^TM*-Ano1fl/fl mice and n=10 popliteal vessels from 6 mice Ano1fl/fl mice.

1340

Figure 9 Isobaric contractile Assessment of popliteal cLV from *PdgfraCre*TM-Cx45 fl/fl mice

Summary of the contractile parameters recorded from popliteal cLVs in *PdgfraCreER^{TM}*-

CX45fl/fl mice. Normalized contraction amplitude (A), ejection fraction (B), contraction

frequency (C), fractional pump flow (D), end diastolic diameter (E), and vessel tone (F) were

assessed. No statically significant differences observed. Mean and SEM shown, n=5 popliteal

vessels from 3 mice *PdgfraCreER^{TM}*-CX45fl/fl mice and n=8 popliteal vessels from 11 mice

CX45fl/fl mice.

1348

1349
Figure 10 Isobaric contractile Assessment of popliteal cLV from *PdgfrαCre*TM-Cav1.2 fl/fl mice
Summary of the contractile parameters recorded from popliteal cLVs in *PdgfrαCreER*[TM]-
Cav1.2fl/fl mice. Normalized contraction amplitude (A), ejection fraction (B), contraction
frequency (C), fractional pump flow (D), end diastolic diameter (E), and vessel tone (F) were
assessed. Mean and SEM shown, n=6 popliteal vessels from 3 mice *PdgfrαCreER*[TM]-Cav1.2fl/fl
mice and n=9 popliteal vessels from 20 mice Cav1.2fl/fl mice. The contractile data from control
Cav1.2fl/fl vessels was previously published but was separated by sex (Davis et al., 2022) while
they are combined here. * denotes significance at p <0.05.

1358
Figure 11 PDGFRα Cells Express Multipotent Cell Makers
Representative RT-PCR results profiling purified GFP+ cells purified from IALVs isolated from
*PdgfraCreER*TM-ROSA26mTmG via FACS. PDGFRα cells expressed the multipotent markers Klf4,
Sca1, Gli1, CD29, CD105, and CD44 (A) with total brain cDNA serving as a positive control (B).
Representative RT-PCR results showing lack of expression of some of these markers in the GFP⁺ cells
purified from *Myh11CreER^T2*--ROSA26mTmG (C) or *Prox1-eGFP* mice, in contrast to the RFP⁺
population from *Myh11CreER^T2*--ROSA26mTmG mice (D). RT-PCRs were repeated at least 2 times
from separate purified cells populations from different mice. Representative max projections of
IALVs stained for Sca1 (E), PDGFRα (E), SMA (F) or Sca1 (I), PDGFRα (K), MYH11 (J), and their
corresponding merged file (H, L). Scale bar is 100 μm.
1369
1370
1371
1372

1373
Figure 12 ChR2-Mediated Depolarization in LMCs Only Triggers Contraction
Representative max projections of tdTomato-ChR2 signal in popliteal cLVs isolated from
cKitCreER^{T2}-ChR2-tdTomato (A), *PdgfαCreER^{TM}*-ChR2-tdTomato (C), and *Myh11CreER^{T2}*- ChR2-
tdTomato (E) with their corresponding brightfield image (B, D, F) respectively. Time-lapse

brightfield images every 0.5 s starting at stimulation t=0 for *cKitCreER^{T2}*-ChR2-tdTomato (G-J),
PdgfraCreER^{TM}-ChR2-tdTomato (K-N), and *Myh11CreER^{T2}*- ChR2-tdTomato (O-R). The I bar
denotes the inner diameter at t=0 over time and white asterisks denote the contraction.
Representative diameter trace for the popliteal cLV demonstrate spontaneous contractions
with the dotted boxes indicating the optical stimulation event in the respective brightfield
images of the time lapse images. Isolated cLVs from *cKitCreER^{T2}*-ChR2-tdTomato (S),
PdgfraCreER^{TM}-ChR2-tdTomato (T), and *Myh11CreER^{T2}*- ChR2-tdTomato (U) were stimulated
with light pulses (red dashed lines) and the summation of contraction triggering for each
genotype (V). Mean and SEM are shown, **** denotes $p < 0.0001$. Contraction recorded from at
least 6 popliteal cLVs from 3 mice per genotype.
1388
1389

A *cKitCreER^T2*-GCaMP6f Maximum Projection

B *cKitCreER^T2*-GCaMP6f STMs

Cell 1
Cell 2
Cell 3
Cell 4
Cell 5
Cell 6

Cell 7
Cell 8
Cell 9

Time (12.5s)

C *cKitCreER^T2*-GCaMP6f Cell Plots

Cell 1
Cell 2
Cell 3
Cell 4
Cell 5
Cell 6
Cell 9
Cell 7
Cell 8

D Incidental LEC and LMC

Cell 10 (LMC)
Cell 11 (LEC)

E

Cell 10
Cell 11

F Maximum Projection: 100 ug/ml C48-80

G *cKitCreER^T2*-GCaMP6f STMs
100 μg/ml Compound 48-80

Cell 1
Cell 2
Cell 3
Cell 4
Cell 5
Cell 6
Cell 7
Cell 8
Cell 9

Time (18.2s)

H *cKitCreER^T2*-GCaMP6f Cell Plots

100 μg/ml Compound 48-80

Cell 1
Cell 2
Cell 3
Cell 4
Cell 5
Cell 6
Cell 7
Cell 8
Cell 9

I Incidental LEC and LMC

100 μg/ml Compound 48-80

Cell 10 (LMC)
Cell 11 (LEC)

J

Cell 10
Cell 11

1390

Figure 13 *cKitCreER*T2 Drives GCaMP6f Expression in Primarily Mast Cells in Murine IALVs
Representative max projection of GCaMP6f signal over time in an IALV isolated from a
cKitCreER^{T2}-GCaMP6f mouse with ROI indicated around individual cells, primarily large ovoid
cells, but also including a circumferential LMC (Cell10) and a horizontal LEC (Cell 11). Of cells 1-
9, only cell 7 had any Ca^{2+} activity (red arrows) during the recording time as indicated by the
STMs from each ROI (B) and their normalized F/F_0 plots in (C). In contrast, the LMC in ROI 10
had both rhythmic global Ca^{2+} events (D) that spanned the cell axis (vertical axis) in the STM (E)
in addition to localized Ca^{2+} events intervening the time between global events (green arrows).
Representative max projection of GCaMP6f signal over time after stimulation with C48-80 (F)
with many large ovoid cells displaying long lasting global Ca^{2+} events (G,H) while not
immediately affecting the LMC Ca^{2+} dynamics (I,J).

1404
Figure 14 Lack of coordinated Ca²⁺ Activity Across Contraction Cycle in PDGFRα Cells
Representative max projections of GCaMP6f signal over time in an IALVs isolated from
PdgfrαCreER^TM-GCaMP6f mice (A, D). ROIs were made around cells and GCaMP6f recorded over
time to generate the corresponding STMs (B, E) for each cell and plots (C, F) respectively. Once
again, incidental recombination occurred in a LMC which displayed rhythmic Ca²⁺ flashes (C)
while the slight undulation in the other cells is due to movement artifact (B). Red arrows
indicate the limited local Ca²⁺ activity observed in two cells from a *PdgfrαCreER^TM*-GCaMP6f
IALV.
1413
1414

1415
Figure 15 Heterogeneous Diastolic Ca^{2+} Transient Activity in LMCs
Representative max projections of GCaMP6f signal over time in an IALVs isolated from
Myh11CreER^{T2}-GCaMP6f mice (A). LMCs were outlined with ROIs to assess GCaMp6F signal over
time. Rhythmic global flashes (B) were entrained across all the LMCs in the FOV (C) with many
cells exhibiting diastolic Ca^{2+} release events. Cells exhibiting at least one diastolic Ca^{2+} event,
within the context of our focal plane constraints, over the recorded time were denoted by the
red asterisks. The plot in (D) magnifies the first diastolic period, seconds 1-3, of C to assist in
visualizing the lack of coordination of the diastolic events. (E) Max projection of the pseudo-
linescan analysis across the axis of the vessel to highlight diastolic Ca^{2+} transients in all cells in
the field of view and their lack of coordination across the cells (x-axis). The white dotted box
shows the first diastolic period plotted in (D).
1427
1428

Figure 16 Pressure Dependency of Murine LMC Diastolic Ca^{2+} Transients
Representative max projection of GCaMP6f signal over 20 s in an IALVs isolated from
Myh11CreER^{T2}-GCaMP6f mice in the presence of the L-type blocker nifedipine (1μM) (A)
pressurized to 0.5 cmH$_2$O, 2 cmH$_2$O, 5 cmH$_2$O. The local diastolic Ca^{2+} transients persist in the
presence of nifedipine and increase with increasing pressure as demonstrated in the whole
vessel STMs (B). Particle occurrence maps highlight the Ca^{2+} activity in each LMC as pressure is
raised (C). Representative particle analysis plots for particle area (D) and particle counts/frame
at each pressure (E). Summary files for particle area (F) and count /frame (G0. * denotes
p<0.05, Mean and SEM shown with n=12 separate IALVs from 8 MYH11-CreER^{T2}-GCaMP6f mice.

A *cKitCreER^{T2}*-GCaMP6f Maximum Projection

B *cKitCreER^{T2}*-GCaMP6f STMs

Cell 1
Cell 2
Cell 3
Cell 4
Cell 5
Cell 6

Cell 7
Cell 8
Cell 9

Time (12.5s)

C *cKitCreER^{T2}*-GCaMP6f Cell Plots

F/F_0

Time (s)

Cell 1
Cell 2
Cell 3
Cell 4
Cell 5
Cell 6
Cell 9
Cell 7
Cell 8

D Incidental LEC and LMC

F/F_0

Time (s)

Cell 10 (LMC)
Cell 11 (LEC)

E
Cell 10
Cell 11

F Maximum Projection: 100 ug/ml C48-80

G *cKitCreER^{T2}*-GCaMP6f STMs
100 μg/ml Compound 48-80

Cell 1
Cell 2
Cell 3
Cell 4
Cell 5
Cell 6
Cell 7
Cell 8
Cell 9

Time (18.2s)

H *cKitCreER^{T2}*-GCaMP6f Cell Plots

↓ 100 μg/ml Compound 48-80

F/F_0

Time (s)

Cell 1
Cell 2
Cell 3
Cell 4
Cell 5
Cell 6
Cell 7
Cell 8
Cell 9

I Incidental LEC and LMC

↓ 100 μg/ml Compound 48-80

F/F_0

Time (s)

Cell 10 (LMC)
Cell 11 (LEC)

J
Cell 10
Cell 11

A *PdgfrαCreER^TM*-GCaMP6f Maximum Projection

100μm

D *PdgfrαCreER^TM*-GCaMP6f Maximum Projection

100μm

B *PdgfrαCreER^TM*-GCaMP6f STMs

Cell 1
Cell 2
Cell 3
Cell 4
LMC

Time (20s)

E *PdgfrαCreER^TM*-GCaMP6f STMs

Cell 1
Cell 2
Cell 3
Cell 4
Cell 5

Time (20s)

C *PdgfrαCreER^TM*-GCaMP6f Cell Plots

F/F_0

Time (s)

Cell 1
Cell 2
Cell 3
Cell 4
Cell 5
(LMC)

F *PdgfrαCreER^TM*-GCaMP6f Plots

F/F_0

time

Cell 1
Cell 2
Cell 3
Cell 4
Cell 5

A *Myh11CreER^{T2}*-GCaMP6f Maximum Projection

B *Myh11CreER^{T2}*-GCaMP6f Single Cell STMs

LMC 1
LMC 2
LMC 3 *
LMC 4 *
LMC 5 *
LMC 6 *
LMC 7

LMC 8 *
LMC 9 *
LMC 10 *
LMC 11 *
LMC 12 *
LMC 13*
LMC 14 *

Time (20s)

Time (20s)

C *Myh11CreER^{T2}*-GCaMP6f Cell Plots

F/F_0

Time (s)

— LMC1
— LMC2
— LMC3
— LMC4
— LMC5
— LMC6
— LMC7
— LMC8
— LMC9
— LMC10
— LMC11
— LMC12
— LMC13
— LMC14

D *Myh11CreER^{T2}*-GCaMP6f Cell Plots

F/F_0

Time (s)

E *Myh11CreER^{T2}*-GCaMP6f Whole Vessel STM

Vessel Axis

8 Sec

A P=0.5 cmH2O   P=2 cmH2O   P=5 cmH2O   *Myh11CreER^T2^-GCaMP6f*

B P=0.5 cmH2O   P=2 cmH2O   P=5 cmH2O

1μM Nifedipine   Time

5 sec

50μm

C P=0.5 cmH2O   P=2 cmH2O   P=5 cmH2O

D P=0.5 cmH$_2$O   P=2 cmH$_2$O   P=5 cmH$_2$O

PTCL Area μm^2/frame   Frame

E P=0.5 cmH$_2$O   P=2 cmH$_2$O   P=5 cmH$_2$O

PTCL Count per frame   Frame

F Particle Area

PTCL Area μm^2/frame

Pressure cmH$_2$O

**   *   **

G Particle Freq

PTCL Count per Frame

Pressure cmH$_2$O

**   *   **