A chromosome-level genome assembly of *Zasmidium syzygii* isolated from

banana leaves

A.C. van Westerhoven[1,2*], R. Mehrabi[1,3*], R. Talebi[3], M.B.F Steentjes[1], B. Corcolon[4],

Pablo A. Chong[5], G.H.J. Kema[1,#,*], M. F. Seidl[2,#,*]

[1]Laboratory of Phytopathology, Wageningen University & Research, the Netherlands

[2]Theoretical Biology & Bioinformatics, Department of Biology, Utrecht University, the

Netherlands

[3]Keygene N.V, P.O. Box 216, 6700 AE, Wageningen, Netherlands

[4]Research, Information, Compliance Department, Tadeco Inc., Panabo, Philippines

[5]ESPOL Polythecnic University, Escuela Superior Politécnica del Litoral, Centro de

Investigaciones Biotecnológicas del Ecuador, Laboratorio de Biología Molecular,

Guayaquil, Ecuador

*Authors contributed equally

[#]Corresponding authors: Gert H.J. Kema and Michael F. Seidl

Keywords: *Zasmidium syzygii,* genome assembly, banana leaf disease, foliar blight,

plant disease

Abstract

Accurate taxonomic classification of samples from infected host material is essential for disease diagnostics and genome analyses. Despite the importance, diagnosis of fungal pathogens that cause banana leaf diseases remains challenging. Foliar diseases of bananas are mainly caused by three *Pseudocercospora* species, of which the most predominant causal agent is *P. fijiensis*. Here, we sequenced and assembled four fungal isolates obtained from necrotic banana leaves in Bohol (Philippines) and obtained a high-quality chromosome-level genome assembly for one of these isolates. Using PCR diagnostics, the samples were initially identified as *P. fijiensis*, yet the assembly size was consistently 30 Mb smaller than expected. Based on the ITS sequences, we identified the samples as *Zasmidium syzygii*. The *Zasmidium syzygii* genome assembly is 42.5 Mb in size distributed over 16 chromosomes, eleven of which are complete. The genome contains 98.6% of the expected single-copy BUSCO genes and we predicted 14,789 protein-coding genes and 10.3% repeats. The three other genome assemblies based on short-read data are less continuous but have similar genome sizes (40.4 – 42.4 Mb) and contain between 96.5% and 98.4% BUSCO genes. All four isolates have identical ITS sequences and are distinct from *Zasmidium* isolates that have been previously sampled from banana leaves. We thus report the first chromosome-level genome assembly of a member of the *Zasmidium* genus, forming an essential resource for further analysis to enhance our understanding of the diversity of pathogenic fungal isolates as well as fungal diversity.

Introduction

Genome sequencing is important for the diagnostics and monitoring of diseases and forms an important step to help to understand the biology of pathogens and diseases. To enable sample identification and downstream genome analyses, accurate taxonomic classification and prevention of contamination in genome assemblies are essential (Francois et al., 2020; Lu & Salzberg, 2018; Rachtman et al., 2020). However, publicly available genome assemblies are occasionally reported to contain contaminants, which can lead to incorrect species classification (Cornet & Baurain, 2022; Kusch et al., 2023; Steinegger & Salzberg, 2020). Obtaining a clean genome assembly is especially challenging for pathogens that live in close association with other organisms; samples from pathogens are often contaminated with host material or other organisms that proliferate in or in proximity to host plants such as for example bacteria or endophytic fungi (Kusch et al., 2023; Zaccaron & Stergiopoulos, 2021).

Banana is an important food crop that provides food security in tropical and subtropical regions worldwide. Foliar blights are a major constraint to banana production and are mainly caused by a complex of three *Pseudocercospora* species, with *P. fijiensis*, the causal agent of black leaf streak disease or Black Sigatoka, being the dominant species. Control of this disease is responsible for up to 25% of the total costs of banana production (Drenth & Kema, 2021). The two other species also cause foliar blights, but are currently less prevalent; *P. musae* causes yellow Sigatoka and *P. eumusae* causes eumusae leaf spot (Chang et al., 2016). However, besides these three *Pseudocercospora* species, other fungal species can appear in association with symptomatic banana foliage. For example, a recent study that analysed fungal isolates associated with banana foliar diseases revealed the

presence of over 30 other fungal species, primarily belonging to the

Mycosphereallaceae family (Crous et al., 2021). Interestingly, before Crous and

colleagues revealed the identity of these isolates, most of these fungi were

considered to belong to the *Pseudocercospora* genus, highlighting that accurate

diagnosis of the causal agents of leaf symptoms remains challenging. However,

accurate identification and classification of the obtained isolates is important,

especially when developing disease diagnostics or enhancing the effectiveness of

disease management strategies (Kusch et al., 2023; Lu & Salzberg, 2018).

Here we sampled four fungal isolates from banana foliage showing blight

symptoms in Bohol, Philippines, and identified the fungal isolates by disease

symptoms, morphology, and diagnostic PCR assays (Arzanlou et al., 2008).

Interestingly, initial observations classified the isolates as *P. fijiensis*, but further

genome analyses revealed that these isolates rather belong to *Zasmidium syzygii*.

Until now, only three fragmented genome assemblies of *Zasmidium* species are

publicly available (Ametrano et al., 2022; Xu et al., 2017). Here, we assembled the

first chromosome-level genome assembly of a representative of the *Zasmidium*

genus. The availability of this genome assembly will help to improve molecular

diagnostics for pathogenic *Pseudocercospora* spp. on banana foliage. Moreover, the

genome adds to the diversity of available fungal genomes, providing the community

with a useful resource for future genomic studies.

Methods & Materials

Fungal isolation and sequencing

Banana leaves with symptoms of foliar disease were collected from a field in Bohol, Philippines. Leaf samples were cut into 1 cm^2 segments and used to discharge ascospores as described previously (Chong et al., 2019). Four single ascospore isolates (P121, P122, P123, and P124) were collected and grown on potato dextrose agar (PDA) plates supplemented with streptomycin (100 µg/ml) for three weeks at 25 °C. To obtain sufficient fungal biomass for DNA isolation, a piece of mycelium (2 cm^2) was blended for 20 sec. at 6,000 rpm in an Ultra Turrax Tube Drive homogenizer (IKA, Staufen, Germany) in 15 ml water using a sterile DT-20 tube (IKA, Staufen, Germany). The fragmented mycelium was transferred into a flask containing 100 ml of PDB amended with streptomycin (100 µg/ml) and kept at 25 °C on a rotary shaker with 150 rpm for about two weeks. The fungal mycelium was filtered through miracloth and subsequently washed with sterile water. Fungal mycelium was freeze-dried overnight and used for high-molecular-weight (HMW) DNA isolation based on the CTAB method (Murray & Thompson, 1980). After adding isopropanol, HMW DNA was collected from the extraction buffer using a sterile needle. AMPure XP purification kit (Becman Cluter Life Sciences, USA) was used to clean up the DNA. DNA quality and quantity were checked by gel electrophoresis, Nanodrop micro-volume spectrophotometers, and Qubit Fluorometric Quantitation (Thermo Fisher Scientific, USA). The HMW DNA of isolate P124 was sequenced using PromethION Oxford Nanopore (ONT) sequencing technologies. Additionally, all isolates were sequenced using the Illumina HiSeq platform, both sequencing platforms were located at Keygene B.V. (Wageningen, the Netherlands).

Fungal species diagnosis by PCR

117 To identify the isolated fungal species, the DNA from each isolate was extracted and

118 subjected to *P. fijiensis*-specific diagnostic PCR according to a previously

119 established protocol (Arzanlou et al., 2008). The list of primers used in PCRs is

120 shown in Table S1. Actin (820 bp) was used as a positive control to ensure

121 successful amplification and to assess the quality of DNA.

Pathogenicity assay

The pathogenicity of isolate P124 was tested on two banana genotypes; Cavendish

cv. Grand Naine (AAA) and a diploid (AA) genotype 'Pisang Berlin'. The inoculum

for isolate P124 as well as for the *P. fijiensis* reference isolate P78 were prepared

similarly. A piece of 1 cm^2 mycelium from three-week-old colonies grown on PDA

was collected in an Eppendorf tube containing 3-4 metal beads (3 mm diameter) and

was blended for 20 seconds at 3,000 rpm in a Bead Beater Homogenizer. About one

ml of sterile water was added to each tube and fragmented mycelium was spread on

PDA plates amended with 100 μg/ml streptomycin. The plates were kept at 25°C for

3-4 weeks and then a piece of mycelium (10 cm^2) was blended for 40 sec. at 6,000

rpm in an Ultra Turrax Tube Drive homogenizer (IKA, Staufen, Germany) in 15 ml of

distilled water using a sterile DT-20 tube (IKA, Staufen, Germany). The suspension

was passed through miracloth to remove non-fragmented mycelium. The collected

mycelial fragments were further diluted and adjusted to 5x10^5 fragments.ml^{-1} and

supplemented with 0.15% Tween 20. This suspension was used to inoculate two-

month-old banana plants on both sides of the leaves. Each treatment was repeated

tree times, as a control, water was used for mock inoculation. Inoculated plants were

kept for 48 h at 90% relative humidity at 25ºC in the dark in a growth cabinet and

subsequently for eight weeks in a greenhouse with >85% RH, and with a day length

of 12 hours.

De novo *genome assembly*

To construct a complete and continuous genome assembly, we assembled the ONT

long-read sequences of isolate P124. The raw Fast5 files were basecalled with

Guppy (v. 5.1.13) and filtered for a minimum quality score of 10. The filtered reads

were assembled *de novo* using Canu (v. 2.2, Koren et al., 2017) with the parameters

'-OvlMerThreshold=300' and '-corMaxEvidenceErate=0.15' to improve the assembly

of repetitive regions. The initial genome assembly was polished with Racon (v. 1.5.0)

using the raw ONT long reads. After polishing, missing telomeres were

reconstructed, when possible, using teloclip (v. 0.0.3,

https://github.com/Adamtaranto/teloclip) and a final polishing step was performed

based on shot-read Illumina data using Pilon (v. 1.24, Walker et al., 2014). The

presence of telomeres (TTAGGG) and coverage of the chromosomes in the final

polished assembly were determined using tapestry (v. 1.0.0, Davey et al., 2020). To

estimate the completeness of the genome assembly, the presence of conserved

single-copy genes was determined using BUSCO with the Capnodiales database

(Simão et al., 2015). The assembly continuity and potential contaminations were

accessed using BlobTools (v. 4.1.5, Challis et al., 2020). To estimate the size of the

assembled genome, all Illumina sequences were analyzed using GenomeScope (v.

2.0, Vurture et al., 2017) with the 21-mer profile determined by JellyFish (v 2.3.0,

Marçais & Kingsford, 2011). Isolate P121 showed high k-mer coverage outliers,

therefore the maximal k-mer coverage was set to 350x, similar to the maximum k-

mer coverage observed in the other genome. Repeats were predicted using

RepeatModeler (v. 2.0.3, Flynn et al., 2020) and annotated using RepeatMasker (v.

4.1.2, Smit et al., 2015). Finally, protein-coding genes were annotated using

Funannotate (v. 1.8.9, Palmer & Stajich, 2019) with the BUSCO lineage

Pezizomycotina and utilizing predicted *Zasmidium cellare* proteins as evidence

(GCF_010093935.1, Haridas et al., 2020). Additionally, we assembled the Illumina

sequenced isolates P121-P123 with Spades (v. 3.13, Bankevich et al., 2012). To

assess the quality and completeness of these assemblies, we ran QUAST (v. 5.2,

Gurevich et al., 2013) and BUSCO (v. 5.3.2, Simão et al., 2015) with the

Capnodiales odb10 database.

Identification of internal transcribed spaces sequence

The internal transcribed spacer (ITS) sequences from the four assemblies (P121-

P124) were retrieved using ITSx (v. 1.1.3, Bengtsson-Palme et al., 2013) and blastn

was used to detect similar ITS sequences in the NCBI blast database (accessed 19[th]

November 2022). Additionally, a maximum-likelihood tree was constructed to

compare the four ITS sequences to 227 publicly available ITS sequences obtained

from NCBI (accessed 13[th] January 2023). Sequences were aligned using mafft (v.7.

453) and a maximum-likelihood phylogenetic tree was constructed using RAxML,

with 500 bootstrap replicates (-m GTRCAT -p 1234 -b 100 -N 500) (v. 8.2.12). The

resulting phylogeny was visualized using iTOL (Letunic & Bork, 2021).

Genetic diversity

To assess the genetic diversity among isolates, short-read sequencing data of P121-

P123 were aligned to the P124 long-read genome assembly using BWA-mem (v.

0.7.17, Li, 2013). GATK4 was used to call variants (Van der Auwera et al., 2013),

and these variants were subsequently filtered using GATK Variant Filtration based

on the GATK best practices (Van der Auwera et al., 2013). The filtering process

involved excluding variants from reads with low mapping quality, variants

predominantly located at the edge of reads, and variants exhibiting a bias towards

reverse/forward strands.

Results and Discussion

Fungal isolates obtained from symptomatic banana foliage in Bohol

Foliar blights of banana is mainly caused by a complex of *Pseudocercospora* species (Chang et al., 2016; Drenth & Kema, 2021). However, additional fungal species predominantly belonging to the Mycosphaerellaceae family have also been associated with symptomatic banana leaves (Crous et al., 2021). To further analyze the fungal pathogens that cause disease on banana leaves, four samples were obtained from banana plants with necrotic lesions in Bohol, Philippines (Chong et al., 2021). For all four isolates (here called P121 – P124), we were able to amplify a PCR product using *P. fijiensis*-specific PCR primers (Arzanlou et al., 2008), indicating that the isolates can be identified as *P. fijiensis* (**Figure 1a**). To corroborate the identity of the isolates, we tested the pathogenicity of isolate P124 on banana cultivars Cavendish (cv. Grand Naine, AAA) and Pisang Berlin (AA) and compared the pathogenicity to the reference *P. fijiensis* isolate P78 originating from Tanzania. No leave spot symptoms were observed on Cavendish upon inoculation with P124, in contrast to the necrotic lesions caused by P78 (**Figure 1b**). However, both isolates caused necrotic lesions on Pisang Berlin four weeks after inoculation (**Figure 1c**). The necrotic lesions caused by isolate P124 were, however, less severe than the disease symptoms caused by *P. fijiensis* isolate P78. Even though we observe a difference in pathogenicity between the two isolates, both can cause necrosis on banana foliage.

Figure 1. PCR diagnostics and pathogenicity assay of isolate P124 in
comparison to the *Pseudocercospora fijiensis* reference strain P78 a) All
isolates show amplification with actin primers (left panel). *Pseudocercospora*-specific
primers amplified PCR products from *P. fijiensis* strain P78 (positive control) as well
as from isolates P121, P122, P123, and P124 (right panel), suggesting that the
isolates can be identified as *P. fijiensis*. b) Isolate P124 does not cause necrotic
lesions on Cavendish banana, in contrast to necrotic symptoms caused by P78. c)
Both P124 and P78 cause necrotic leaf symptoms on Pisang Berlin. Disease
symptoms were scored eight weeks after inoculation.
227

Complete genome assembly of P124

To generate a high-quality genome assembly for the isolates from Bohol, we

sequenced strain P124 using Oxford Nanopore Technology, which yielded 11.9 Gb

of reads with an average size of ~9.1 kb, corresponding to a ~150x genome

coverage based on the estimated genome size of 74 Mb (Arango et al., 2016). The

de novo genome assembly resulted in 16 contigs with an N50 of 3.4 Mb and a total

genome size of 42.5 Mb. Eleven of the 16 contigs had telomeric sequences

(TTAGGG) at both ends, and thus the assembly is highly contiguous and mostly

represents complete chromosomes (**Figure 2**). To assess genome completeness,

we queried the genome assembly for the presence of single-copy BUSCO genes

and identified 98.6% of the single-copy BUSCO genes that are expected in the

fungal order Capnodiales, indicating that the genome assembly of P124 covers the

conserved gene space (**Figure 2a**). We predicted a total of 14,789 protein-coding

genes in the genome and identified that 10.3% of the genome consists of repetitive

elements (**Figure 2b**). *De novo* genome assembly of isolates P121 – P123,

sequenced with short-read sequencing technology only, resulted in genome

assemblies with similar sizes (40.4 – 42.5 Mb). Although the assemblies of P121 -

P123 are more fragmented compared to the assembly of the nanopore-sequenced

isolate P124, they approximately contain an equally high number of expected

BUSCO genes (Table 1).

Table 1. The genome assembly statistics of four sequenced isolates obtained

from symptomatic banana foliage in Bohol, Philippines.

	P124*	P121**	P122**	P123**
Genome size (Mb) based on k-mer estimation (k=21)	40.6	39.8	39.9	41.9
Genome assembly size (Mb)	42.5	40.4	40.6	42.2
Number of contigs	16	1,225	1,563	796
Largest contig (Mb)	6.97	0.44	0.31	0.84
GC%	51.70	52.55	52.37	52.01
N50 (Mb)	3.40	0.98	0.60	1.9

N75 (Mb)	2.08	0.49	0.30	1.05
L50	2	134	201	67
BUSCO completeness (*n* = 3,578)	98.6%	97.8%	96.5%	98.4%
BUSCO fragmented genes	0.5%	0.9%	1.5%	0.5%
BUSCO missing genes	1.0%	1.3%	2.0%	1.1%

*P124 is assembled using long reads from Oxford Nanopore Technology (ONT).

**P121-P123 are assembled using Illumina short reads.

The *P. fijiensis* reference genome is 73.6 Mb (Arango et al., 2016), and remarkably,

the genome assemblies of the here sequenced fungal isolates are approximately 30

Mb smaller than expected. To verify the assembly size, we estimated genome sizes

using k-mer profiles, which resulted in an estimated genome size ranging from 38.9

to 41.9 Mb, similar to the size of the *de novo* assembled genomes (Table 1).

Furthermore, we mapped the sequencing reads of isolates P121 – P124 to the

chromosome-level genome assembly of P124, mapping on average 95% of the

reads with limited genetic diversity between isolates (on average 15 SNPs per kb;

Figure 2b). Moreover, mapping short reads of P124 to the P124 chromosome-level

genome assembly did not reveal any major coverage deviation (**Figure 2c**),

indicating that the P124 genome assembly does not contain extensively collapsed

repetitive regions that could account for the smaller genome size compared to *P.*

fijiensis. This validates that, although the genome assembly of strain P124 is smaller

than expected, it is highly complete and continuous, suggesting that the length

variation is likely not due to assembly errors. To capture possible contaminations in

the genome assemblies, we queried all contigs to the BLAST database to determine

their identity. All contigs showed high similarity to members of the Mycosphaerellaceae family; most contigs were similar to *Zasmidium* species (13 contigs, 35 Mb) and only one contig (3.3 Mb) showed similarity to *Pseudocercospora* (**Figure S1**). Whole-genome alignments of our assembly to the *P. fijiensis* reference genome assembly revealed that none of the contigs/chromosomes display significant similarity (**Figure S2**). Therefore, we considered that the difference in genome size between isolates P121 – P124 and the *P. fijiensis* reference isolate P78 is likely caused by the isolation of a different fungal species associated with necrotic symptoms on banana foliage (Crous et al., 2021), most likely by a member of genus *Zasmidium*.

Figure 2. Chromosome-level genome assembly of *Zasmidium syzygii* isolate P124. a) Genome assembly statistics for the *de novo* assembly of *Z. syzygii* isolate P124 based on Oxford Nanopore Technology (ONT). The genome assembly has 17 contigs (16 nuclear contigs of which at least 11 are complete chromosomes, and the mitochondrial genome) with a total genome size of 42.5 Mb and contains 98.7%

complete single-copy BUSCO genes. **b**) A circular representation of the chromosomes in P124. Dashed lines indicate missing telomeres for chromosome 3, 11, 13, 15, and 16. Genes (14,786) and repeats (10.3%) are distributed evenly over the chromosomes. Single nucleotide polymorphisms are found over all chromosomes, with two SNP dense regions on chromosome 8 and 9. **C**) Short-read coverage of P124 mapped to the assembled contigs of P124 shows minimal regions with exceptionally high or low coverage, suggesting that the assembly does not contain large repetitive regions that may have been collapsed during the assembly process.

The assembled genome sequence reveals that isolate P124 belongs to the fungal genus *Zasmidium*

To determine the identity of isolate P124, we retrieved the ITS sequence from the genome assembly and searched for related sequences in the non-redundant BLAST database on NCBI using the P124 ITS sequence as a query. We obtained a highly similar match (98.7% nucleotide identity) to *Zasmidium syzygii* (NR_111826.1), supporting our hypothesis that the assembled isolate belongs to the *Zasmidium* genus. In line with this finding, previous studies have found *Zasmidium* isolates on symptomatic banana leaves (Arzanlou et al., 2008; Crous et al., 2021), and other *Zasmidium* species are reported to cause leaf spot diseases on other plant species such as for example citrus (Aguilera-Cogley et al., 2017; An et al., 2021; Han et al., 2015). The *Zasmidium* genome assembly we report here is less fragmented and 4 Mb larger than the previous genome assembly of a *Zasmidium* species, yet shares extensive similarity to P124 (**Figure S2**). *Z. cellare* is known as the 'wine cellar fungus' because it thrives in walls and ceilings of wine cellars (Tribe et al., 2006), and was sequenced and its genome assembled into >267 scaffolds with a genome size of ~38 Mb (Tribe et al., 2006), illustrating that our genome offers a more continuous genome representation of a member of the *Zasmidium* genus.

To validate the identity and to determine the diversity of the *Zasmidium* isolates, we compared the ITS sequences of isolate P121-P124 with 43 other *Zasmidium* ITS sequences from NCBI (20 December 2022) as well as ITS sequences of 120 Mycosphaerellaceae strains obtained from banana leaves (Crous et al., 2021). A maximum-likelihood phylogeny based on the aligned ITS sequences confirmed that isolates P121 – P124 belong to the *Zasmidium* genus and shows that these four isolates encode identical ITS sequences (**Figure 3**). Notably, the set of

ITS sequences also contains sequences from five other *Zasmidium* isolates that had

been previously sampled from banana leaves from different geographic locations

(Martinique, Tonga, and Gabon) (Crous et al., 2021). Interestingly, these do not

cluster with isolates P121-P124, suggesting that pathogenicity towards banana is a

polyphyletic trait within the *Zasmidium* genus (**Figure 3**). Although *Zasmidium*

species have been linked to foliar blights in various hosts (Aguilera-Cogley et al.,

2017; Han et al., 2015; Osorio et al., 2021), the pathogenicity and global spread of

Zasmidium species has not been studied in depth. Based on our data, we conclude

that Z. *syzygii* occurs on banana leaves with necrotic symptoms and can be the

cause of mild necrotic lesions on banana foliage. However, the abundance and role

of Z. *syzygii* as a banana pathogen remains unknown and requires further research

to understand its prevalence, significance, and potential impact on banana

cultivation.

Figure 3. The four isolates sampled from Bohol (P121-P124) belong to the genus *Zasmidium*. A maximum-likelihood phylogenetic tree was constructed using the internally transcribed spacer (ITS) sequences of 43 *Zasmidium* species from NCBI and 120 Mycosphaerellaceae isolates from banana leaves (Crous et al., 2021). The ITS sequence extracted from the chromosome-level genome assembly (P124) as well as from the other isolates (P121-123) associated with an isolate classified as *Z. syzygii* and are not related to *Zasmidium* strains previously isolated from banana leaves. *Zasmidium* isolates associated with infected banana leaves (yellow labels) are genetically diverse and are distributed across various branches of the phylogenetic tree.

Accurate diagnostics of pathogens is essential to detect the emergence and trace the dispersal of diseases, which is pivotal for effective disease management. However, our data reveal that *P. fijiensis* and *Z. syzygii* are indistinguishable with the current PCR diagnostic (Arzanlou et al., 2008). To compare the similarity of the PCR

primers between *P. fijiensis* and *Z. syzygii*, we *in silico* detected the amplicon of the

supposedly *Pseudocercospora*-specific primers in the *Zasmidium syzygii* P124

genome assembly and in the *P. fijiensis* reference genome assembly (Cirad86;

Arango et al., 2016). Both isolates possess the primer sequence used to distinguish

P. fijiensis (**Figure S3**) from *P. musae* and *P. eumusae* and produce a similar sized

amplicon of 480 bp in *Z. syzygii* P124 and 478 bp in *P. fijiensis* Cirad86, which

explains the positive result for *Z. syzygii* in our PCR assay (**Figure 1**). The amplicons

share 88% sequence identity and *Zasmidium* or *Pseudocercospora* isolates can

therefore be distinguished only upon amplicon sequencing. Thus, novel primer pairs

need to be developed to enable easy and accurate diagnosis of fungal species

present in foliar blights of banana.

Conclusion

Here, we report the first chromosome-scale genome assembly of a *Zasmidium*

species, this adds a high-quality genome sequence to the thus far limited genetic

resources available for this genus. Our data show that *Z. syzygii* occurs on banana

foliage in Bohol and can cause leaf necrosis, comparable to the foliar blight

symptoms observed for *P. fijiensis*. The availability of the genome assembly will

facilitate further research into the association of *Zasmidium, Pseudocercospora*, and

possibly other fungal species related to foliar blights of banana. Moreover, it will

serve as a valuable resource for developing novel molecular diagnostics, enabling

the accurate identification and characterization of these fungal species.

Data Availability Statement

All sequencing data are deposited in the NCBI database under accession number

PRJNA931964.

Acknowledgments

We thank the Research Support Group (RSG) at KeyGene B.V. (Wageningen, the

Netherlands) for performing the DNA isolation, library preparation, and genome

sequencing. We thank Xiaoqian Shi-Kunne (Wageningen University & Research, the

Netherlands) for the data management and bioinformatics support.

Funder Information

ACW and GHJK were supported by the Bill and Melinda Gates Foundation, grant

number AG – 4425. Banana research at Wageningen University has been supported

by the Dutch Dioraphte Foundation, grant number 20 04 04 02.

Conflict of Interest

The authors report no conflict of interests.

References

Aguilera-Cogley, V. A., Berbegal, M., Català, S., Brentu, F. C., Armengol, J., & Vicent, A. (2017). Characterization of Mycosphaerellaceae species associated with citrus greasy spot in Panama and Spain. *PLOS ONE*, *12*(12), e0189585. https://doi.org/10.1371/journal.pone.0189585

Ametrano, C. G., Lumbsch, H. T., Di Stefano, I., Sangvichien, E., Muggia, L., & Grewe, F. (2022). Should we hail the Red King? Evolutionary consequences of a mutualistic lifestyle in genomes of lichenized ascomycetes. *Ecology and Evolution*, *12*(1), e8471. https://doi.org/10.1002/ece3.8471

An, Y.-Y., Zeng, X.-Y., Geng, K., Hyde, K. D., & Wang, Y. (2021). One new species and one new record of Zasmidium in China. *Biodiversity Data Journal*, *9*, e59001. https://doi.org/10.3897/BDJ.9.e59001

Arango, R. E. isaza, Diaz-Trujillo, C., Dhillon, B., Aerts, A., Carlier, J., Crane, C. F., Jong, T. V. de, Vries, I. de, Dietrich, R., Farmer, A. D., Fereira, C. F., Garcia, S., Guzman, M., Hamelin, R. C., Lindquist, E. A., Mehrabi, R., Quiros, O., Schmutz, J., Shapiro, H., … Kema, G. H. J. (2016). Combating a global threat to a clonal crop: Banana Black Sigatoka pathogen *Pseudocercospora fijiensis* (synonym *Mycosphaerella fijiensis*) genomes reveal clues for disease control. *PLOS Genetics*, *12*(8), e1005876. https://doi.org/10.1371/journal.pgen.1005876

Arzanlou, M., Groenewald, J. Z., Fullerton, R. A., Abeln, E. C. A., Carlier, J., Zapater, M.-F., Buddenhagen, I. W., Viljoen, A., & Crous, P. W. (2008). Multiple gene genealogies and phenotypic characters differentiate several novel species of *Mycosphaerella* and related anamorphs on banana. *Persoonia : Molecular*

Phylogeny and Evolution of Fungi, *20*, 19–37.

https://doi.org/10.3767/003158508X302212

Bankevich, A., Nurk, S., Antipov, D., Gurevich, A. A., Dvorkin, M., Kulikov, A. S.,

Lesin, V. M., Nikolenko, S. I., Pham, S., Prjibelski, A. D., Pyshkin, A. V.,

Sirotkin, A. V., Vyahhi, N., Tesler, G., Alekseyev, M. A., & Pevzner, P. A.

(2012). SPAdes: A new genome assembly algorithm and its applications to

single-cell sequencing. *Journal of Computational Biology*, *19*(5), 455–477.

https://doi.org/10.1089/cmb.2012.0021

Bengtsson-Palme, J., Ryberg, M., Hartmann, M., Branco, S., Wang, Z., Godhe, A.,

De Wit, P., Sánchez-García, M., Ebersberger, I., de Sousa, F., Amend, A.,

Jumpponen, A., Unterseher, M., Kristiansson, E., Abarenkov, K., Bertrand, Y.

J. K., Sanli, K., Eriksson, K. M., Vik, U., … Nilsson, R. H. (2013). Improved

software detection and extraction of ITS1 and ITS2 from ribosomal ITS

sequences of fungi and other eukaryotes for analysis of environmental

sequencing data. *Methods in Ecology and Evolution*, *4*(10), 914–919.

https://doi.org/10.1111/2041-210X.12073

Challis, R., Richards, E., Rajan, J., Cochrane, G., & Blaxter, M. (2020). BlobToolKit –

interactive quality assessment of genome assemblies. *G3*

Genes|Genomes|Genetics, *10*(4), 1361–1374.

https://doi.org/10.1534/g3.119.400908

Chang, T.-C., Salvucci, A., Crous, P. W., & Stergiopoulos, I. (2016). Comparative

genomics of the Sigatoka disease complex on banana suggests a link

between parallel evolutionary changes in *Pseudocercospora fijiensis* and

Pseudocercospora eumusae and increased virulence on the banana host.

PLOS Genetics, *12*(8), e1005904.

https://doi.org/10.1371/journal.pgen.1005904

Chong, P., Essoh, J. N., Arango Isaza, R. E., Keizer, P., Stergiopoulos, I., Seidl, M.

F., Guzman, M., Sandoval, J., Verweij, P. E., Scalliet, G., Sierotzski, H., de

Lapeyre de Bellaire, L., Crous, P. W., Carlier, J., Cros, S., Meijer, H. J. G.,

Peralta, E. L., & Kema, G. H. J. (2021). A world-wide analysis of reduced

sensitivity to DMI fungicides in the banana pathogen *Pseudocercospora*

fijiensis. Pest Management Science, *77*(7), 3273–3288.

https://doi.org/10.1002/ps.6372

Chong, P., Vichou, A.-E., Schouten, H. J., Meijer, H. J. G., Arango Isaza, R. E., &

Kema, G. H. J. (2019). Pfcyp51 exclusively determines reduced sensitivity to

14α-demethylase inhibitor fungicides in the banana black Sigatoka pathogen

Pseudocercospora fijiensis. PLOS ONE, *14*(10), e0223858.

https://doi.org/10.1371/journal.pone.0223858

Cornet, L., & Baurain, D. (2022). Contamination detection in genomic data: More is

not enough. *Genome Biology*, *23*(1), 60. https://doi.org/10.1186/s13059-022-

02619-9

Crous, P. W., Carlier, J., Roussel, V., & Groenewald, J. Z. (2021).

Pseudocercospora and allied genera associated with leaf spots of banana

(Musa spp.). *Fungal Systematics and Evolution*, *7*, 1–19.

https://doi.org/10.3114/fuse.2021.07.01

Davey, J. W., Davis, S. J., Mottram, J. C., & Ashton, P. D. (2020). *Tapestry: Validate*

and edit small eukaryotic genome assemblies with long reads (p.

2020.04.24.059402). bioRxiv. https://doi.org/10.1101/2020.04.24.059402

Drenth, A., & Kema, G. (2021). The vulnerability of bananas to globally emerging

disease threats. *Phytopathology®, 111*(12), 2146–2161.

https://doi.org/10.1094/phyto-07-20-0311-rvw

Flynn, J. M., Hubley, R., Goubert, C., Rosen, J., Clark, A. G., Feschotte, C., & Smit,

A. F. (2020). RepeatModeler2 for automated genomic discovery of

transposable element families. *Proceedings of the National Academy of*

Sciences, 117(17), 9451–9457. https://doi.org/10.1073/pnas.1921046117

Francois, C. M., Durand, F., Figuet, E., & Galtier, N. (2020). Prevalence and

implications of contamination in public genomic resources: A case study of 43

reference arthropod assemblies. *G3 Genes|Genomes|Genetics, 10*(2), 721–

730. https://doi.org/10.1534/g3.119.400758

Gurevich, A., Saveliev, V., Vyahhi, N., & Tesler, G. (2013). QUAST: Quality

assessment tool for genome assemblies. *Bioinformatics (Oxford, England)*,

29(8), 1072–1075. https://doi.org/10.1093/bioinformatics/btt086

Han, K. S., Park, M. J., Choi, I. Y., Park, J. H., & Shin, H. D. (2015). First report of

leaf spot caused by *zasmidium lythri* on *lythrum salicaria* in korea. *Plant*

Disease, 99(5), 728–728. https://doi.org/10.1094/PDIS-10-14-1039-PDN

Haridas, S., Albert, R., Binder, M., Bloem, J., LaButti, K., Salamov, A.,

Andreopoulos, B., Baker, S. E., Barry, K., Bills, G., Bluhm, B. H., Cannon, C.,

Castanera, R., Culley, D. E., Daum, C., Ezra, D., González, J. B., Henrissat,

B., Kuo, A., … Grigoriev, I. V. (2020). 101 Dothideomycetes genomes: A test

case for predicting lifestyles and emergence of pathogens. *Studies in*

Mycology, 96, 141–153. https://doi.org/10.1016/j.simyco.2020.01.003

Koren, S., Walenz, B. P., Berlin, K., Miller, J. R., Bergman, N. H., & Phillippy, A. M.

(2017). Canu: Scalable and accurate long-read assembly via adaptive k-mer

weighting and repeat separation. *Genome Research*, *27*(5), 722–736.

https://doi.org/10.1101/gr.215087.116

Kusch, S., Vaghefi, N., & Kiss, L. (2023). The good, the bad, and the misleading:

How to improve the quality of 'genome announcements'? *Molecular Plant-*

Microbe Interactions®. https://doi.org/10.1094/MPMI-01-23-0009-LE

Letunic, I., & Bork, P. (2021). Interactive Tree Of Life (iTOL) v5: An online tool for

phylogenetic tree display and annotation. *Nucleic Acids Research*, *49*(W1),

W293–W296. https://doi.org/10.1093/nar/gkab301

Li, H. (2013). *Aligning sequence reads, clone sequences and assembly contigs with*

BWA-mem. https://doi.org/10.6084/m9.figshare.963153.v1

Lu, J., & Salzberg, S. L. (2018). Removing contaminants from databases of draft

genomes. *PLOS Computational Biology*, *14*(6), e1006277.

https://doi.org/10.1371/journal.pcbi.1006277

Marçais, G., & Kingsford, C. (2011). A fast, lock-free approach for efficient parallel

counting of occurrences of k-mers. *Bioinformatics*, *27*(6), 764–770.

https://doi.org/10.1093/bioinformatics/btr011

Osorio, J. A., Linnakoski, R., Wingfield, M. J., de Beer, Z. W., & Roux, J. (2021). A

new species in the Mycosphaerellaceae *from Cecidomyiidae leaf galls* on

Avicennia marina in South Africa. *Antonie van Leeuwenhoek*, *114*(5), 515–

526. https://doi.org/10.1007/s10482-021-01537-3

Palmer, J., & Stajich, J. (2019). *nextgenusfs/funannotate: Funannotate v1.5.3*

[Computer software]. Zenodo. https://doi.org/10.5281/zenodo.2604804

Rachtman, E., Balaban, M., Bafna, V., & Mirarab, S. (2020). The impact of

contaminants on the accuracy of genome skimming and the effectiveness of

exclusion read filters. *Molecular Ecology Resources*, *20*(3), 649–661.

https://doi.org/10.1111/1755-0998.13135

Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V., & Zdobnov, E. M.

(2015). BUSCO: Assessing genome assembly and annotation completeness

with single-copy orthologs. *Bioinformatics*, *31*(19), 3210–3212.

https://doi.org/10.1093/bioinformatics/btv351

Smit, A. F. A., Hubley, R., & Green, P. (2015). *RepeatMasker Open-4.0.*

http://www.repeatmasker.org

Steinegger, M., & Salzberg, S. L. (2020). Terminating contamination: Large-scale

search identifies more than 2,000,000 contaminated entries in GenBank.

Genome Biology, *21*(1), 115. https://doi.org/10.1186/s13059-020-02023-1

Tribe, H. T., Thines, E., & Weber, R. W. S. (2006). Moulds that should be better

known: The wine cellar mould, *Racodium cellare*. *Mycologist*, *20*(4), 171–175.

https://doi.org/10.1016/j.mycol.2006.09.016

Van der Auwera, G. A., Carneiro, M. O., Hartl, C., Poplin, R., del Angel, G., Levy-

Moonshine, A., Jordan, T., Shakir, K., Roazen, D., Thibault, J., Banks, E.,

Garimella, K. V., Altshuler, D., Gabriel, S., & DePristo, M. A. (2013). From

fastQ data to high-confidence variant calls: The genome analysis toolkit best

practices pipeline. *Current Protocols in Bioinformatics*, *43*(1), 11.10.1-

11.10.33. https://doi.org/10.1002/0471250953.bi1110s43

Vurture, G. W., Sedlazeck, F. J., Nattestad, M., Underwood, C. J., Fang, H.,

Gurtowski, J., & Schatz, M. C. (2017). GenomeScope: Fast reference-free

genome profiling from short reads. *Bioinformatics*, *33*(14), 2202–2204.

https://doi.org/10.1093/bioinformatics/btx153

Walker, B. J., Abeel, T., Shea, T., Priest, M., Abouelliel, A., Sakthikumar, S., Cuomo,

C. A., Zeng, Q., Wortman, J., Young, S. K., & Earl, A. M. (2014). Pilon: An

integrated tool for comprehensive microbial variant detection and genome

assembly improvement. *PLOS ONE*, *9*(11), e112963.

https://doi.org/10.1371/journal.pone.0112963

Xu, C., Zhang, R., Sun, G., & Gleason, M. L. (2017). Comparative genome analysis

reveals adaptation to the ectophytic lifestyle of sooty blotch and flyspeck

fungi. *Genome Biology and Evolution*, *9*(11), 3137–3151.

https://doi.org/10.1093/gbe/evx229

Figures and tables

Figure 1. PCR diagnostics and pathogenicity assay of isolate P124 in
comparison to the *Pseudocercospora fijiensis* reference strain P78 a) All
isolates show amplification with actin primers (left panel). *Pseudocercospora*-specific
primers amplified PCR products from *P. fijiensis* strain P78 (positive control) as well
as from isolates P121, P122, P123, and P124 (right panel), suggesting that the
isolates can be identified as *P. fijiensis*. b) Isolate P124 does not cause necrotic
lesions on Cavendish banana, in contrast to necrotic symptoms caused by P78. c)
Both P124 and P78 cause necrotic leaf symptoms on Pisang Berlin. Disease
symptoms were scored eight weeks after inoculation.
558

Table 1. The genome assembly statistics of four sequenced isolates obtained from symptomatic banana foliage in Bohol, Philippines.

	P124*	P121**	P122**	P123**
Genome size (Mb) based on k-mer estimation (k=21)	40.6	39.8	39.9	41.9
Genome assembly size (Mb)	42.5	40.4	40.6	42.2
Number of contigs	16	1,225	1,563	796
Largest contig (Mb)	6.97	0.44	0.31	0.84
GC%	51.70	52.55	52.37	52.01
N50 (Mb)	3.40	0.98	0.60	1.9
N75 (Mb)	2.08	0.49	0.30	1.05
L50	2	134	201	67
BUSCO completeness (n = 3,578)	98.6%	97.8%	96.5%	98.4%
BUSCO fragmented genes	0.5%	0.9%	1.5%	0.5%
BUSCO missing genes	1.0%	1.3%	2.0%	1.1%

*P124 is assembled using long reads from Oxford Nanopore Technology (ONT).

**P121-P123 are assembled using Illumina short reads.

Figure 2. Chromosome-level genome assembly of *Zasmidium syzygii* isolate P124. a) Genome assembly statistics for the *de novo* assembly of *Z. syzygii* isolate P124 using Oxford Nanopore Technology (ONT). The genome assembly has 17 contigs (16 nuclear contigs of which at least 11 are complete chromosomes, and the mitochondrial genome) with a total genome size of 42.5 Mb and contains 98.7% complete single-copy BUSCO genes. **b**) A circular representation of the chromosomes in P124. Dashed lines indicate missing telomeres for chromosome 3, 11, 13, 15, and 16. Genes (14,786) and repeats (10.3%) are distributed evenly over the chromosomes. Single nucleotide polymorphisms are found over all chromosomes, with two SNP dense regions on chromosome 8 and 9. **C**) Short-read coverage of P124 mapped to the assembled contigs of P124 shows minimal regions with exceptionally high or low coverage, suggesting that the assembly does not contain repetitive regions that may have collapsed during the assembly process.

Figure 3. The four isolates sampled from Bohol (P121-P124) belong to the genus *Zasmidium*. Maximum-likelihood phylogenetic tree is constructed using the internally transcribed spacer (ITS) sequences of 43 *Zasmidium* species from NCBI and 120 Mycosphaerellaceae isolates from banana leaves (Crous et al., 2021). The ITS sequence extracted from the chromosome-level genome assembly (P124) as well as from the other isolates (P121-123) associated with an isolate classified as *Z. syzygii* and are not related to *Zasmidium* strains previously isolated from banana leaves. *Zasmidium* isolates associated with infected banana leaves (yellow labels) are genetically diverse and are distributed across various branches of the phylogenetic tree.

Scaffold statistics
- Log10 scaffold count (total 17)
- Scaffold length (total 42.5M)
- Longest scaffold (6.98M)
- N50 length (3.40M)
- N90 length (1.47M)

BUSCO capnodiales_odb10 (3578)
- Comp. (98.7%)
- Dupl. (0.6%)
- Frag. (0.4%)
- Missing (1.0%)

Scale
- 42.5M
- 6.98M

Composition
- GC (51.7%)
- AT (48.3%)

b

Chr1, Chr2, Chr3, Chr4, Chr5, Chr6, Chr7, Chr8, Chr9, Chr10, Chr11, Chr12, Chr13, Chr14, Chr15, Chr16

- Genes
- Repeats
- SNPs /10 kb
- 0   ≥500

c

Coverage bp (median)

240, 160, 80

Chromosome 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16

Scolecobasidium musicola
Scolecobasidium musicola
Scolecobasidium musicola
Scolecobasidium musicola
Mycosphaerella musae
Pantospora guazumae
Zymoseptoria tritici
Parapallidocercospora thailandica
Parapallidocercospora thailandica
Parapallidocercospora thailandica
P. fijiensis
P. musicola
P. longisporum
P. emusae
Zasmidium gahniicola
Zasmidium arcuatum
Zasmidium podocarpi
Zasmidium velutinum
Zasmidium hakeae
Zasmidium commune
Zasmidium pseudovespa
Zasmidium corymbiae
Zasmidium strelitziae
Zasmidium rothmanniae
Zasmidium schini
Zasmidium macluricola
Zasmidium iteae
Zasmidium elaeocarpi
Zasmidium queenslandicum
Zasmidium ducassei
Zasmidium thailandicum
Zasmidium musae
Zasmidium musae
Zasmidium musae
Zasmidium musae
Zasmidium musae
Zasmidium musae
Zasmidium dasypogonis
Zasmidium citrigriseum
Zasmidium anthuriicola
Zasmidium indonesianum
Zasmidium musicola
Zasmidium scaevolicola
Zasmidium musae-banksii
Zasmidium hakeicola
Zasmidium grevilleae
Zasmidium pseudoparkii
Zasmidium eucalyptorum
Zasmidium cellare
Zasmidium eucalypticola
Zasmidium syzygii
P122
P124
P123
P121
Zasmidium lonicericola
Zasmidium nocoxi
Zasmidium angulare
Zasmidium fructicola
Zasmidium fructigenum
Zasmidium cyatheae
Zasmidium cerophilum
Zasmidium xenoparkii

Tree scale: 0.25