Genotypic variation in resource exchange, use, and production traits in the legume-rhizobia mutualism.

McCall B. Calvert[1*], Maliha Hoque[1], and Corlett W. Wood[1*]

[1]Department of Biology, University of Pennsylvania, Philadelphia, PA, USA

*Correspondence
McCall B. Calvert, calvm@sas.upenn.edu
Corlett W. Wood, corlett@sas.upenn.edu

Abstract
Mutualisms, reciprocally beneficial interaction between two or more species, are ubiquitous in
nature. A common feature of mutualisms is extensive context-dependent variation in fitness
outcomes. This context-dependency is hypothesized to stem from the environment's mediation
of the relative costs and benefits associated with mutualisms. However, traits related to the
exchange of goods and services in mutualisms have received little attention in comparison to
net fitness outcomes. In this study, we quantified the contribution of host and symbiont
genotypes to variation in resource exchange, use, and production traits measured in the host
using the model mutualism between legumes and nitrogen-fixing rhizobia. We predicted that
plant genotype x rhizobia genotype (GxG) effects would be common to resource exchange traits
because resource exchange is hypothesized to be governed by both interacting partners
through bargaining. On the other hand, we predicted that plant genotype effects would dominate
host resource use and production traits because these traits are only indirectly related to the
exchange of resources. Consistent with our prediction for resource exchange traits, but not our
prediction for resource use and production traits, we found that rhizobia genotype and GxG
effects were the most common sources of variation in the traits that we measured. The results
of this study complement the commonly observed phenomenon of GxG effects for fitness by
showing that numerous mutualism traits also exhibit GxG variation. Furthermore, our results
highlight the possibility that the exchange of resources as well as how partners use and produce
traded resources can influence the evolution of mutualistic interactions. Our study lays the
groundwork for future work to explore the relationship between resource exchange, use and
production traits and fitness (i.e. selection) to test the competing hypotheses proposed to
explain the maintenance of fitness variation in mutualisms.
24
25 **Key words:** mutualism evolution, genotype by genotype interaction, resource exchange,
26 resource use, resource production, legume, rhizobia, *Medicago truncatula*, *Enisfer*
27

Introduction

Almost all multicellular life engages in some form of mutualism. Mutualisms are interspecific interactions in which partners increase their fitness by reciprocally exchanging goods and services. Genetic variation for fitness outcomes of mutualisms are notoriously context dependent, presumably because the environment influences the costs and benefits of the interaction for both partners. However, variation in the benefit and cost traits that underlie mutualisms has received comparatively little attention.

Fitness-related traits have been useful for inferring net interaction outcomes. However, the measurement of fitness-related traits also obscures the individual traits that underlie fitness variation. The fitness outcome of a mutualistic interaction is determined by the balance between its benefits and costs. Benefits can take a variety of forms, including transportation (e.g. the movement of gametes in plant-pollinator mutualisms; Albrecht et al. 2012), protection (e.g. defense against herbivory in ant-acacia mutualisms; Janzen 1966), and nutrition (e.g. the exchange of limiting resources between plants and microbes; Friesen et al. 2011). Costs are related to the energetic investment involved in producing benefits that are transferred to the interacting partner (Bronstein 2001, 2015). For example, in plant-microbe mutualisms that involve an exchange of resources, a principal cost for the host is the carbon that is transferred to microbial partners (Akçay and Roughgarden 2007; Clark et al. 2019).

Beyond the traits involved in the direct exchange of resources, traits related to the use or production of the traded resources can also influence the net fitness consequences of a mutualism by impacting its relative benefits and costs (Vidal and Segraves 2021). Traits like resource use efficiency dictate how efficiently the benefits received from an interacting partner are incorporated into physiological processes (e.g. growth, reproduction, defense) that are linked to fitness. Resource production traits, on the other hand, are involved in the creation of the goods and services that are traded to an interacting partner. These traits dictate the total resource pool available to invest in mutualism, and thereby affect the proportional cost of resource exchange. In plant-microbe interactions, a common resource production trait is photosynthetic rate, because it determines the size of a plant's carbon pool, a portion of which is traded to microbial partners (Grman et al. 2012; Clark et al. 2019).

Like other phenotypes, variation in resource exchange, use, and production traits is likely attributable to genetic variation in one or both partners. Indeed, partner genotype effects, where the fitness outcome of a mutualism depends on the genotype of one or both of the interacting partners, are a common feature of resource mutualisms (Hoeksema and Bruna 2015). An added layer of complication is that traits mediating species interactions can be jointly determined by both interacting partners (Moore et al. 1997; O'Brien et al. 2021b; De Lisle et al. 2022). Traits are jointly determined when variation in a trait is determined by contributions from the genomes of both interacting partners. Therefore, genetic variation in traits that mediate a host-microbe mutualism could arise from host genotype effects, microbe genotype effects, or from genotype-by-genotype (GxG) interactions (Heath 2010; Friesen et al. 2011).

Despite their ecological and evolutionary significance, however, few empirical studies have investigated whether there is heritable variation in resource exchange, use, and production traits in interacting populations. Quantifying genotypic differences in mutualism traits represents an important synthesis between the subfields of mutualism physiology, functional ecology, and evolution. The underlying metabolic and genetic pathways involved in mutualism function are

well defined in many systems, especially in nutritional mutualisms between hosts and symbionts
(Heath et al. 2012; Udvardi and Poole 2013; Levy et al. 2021). Furthermore, there is an
extensive body of theoretical work on the resource economics of mutualisms (Akçay and
Roughgarden 2007; Akçay and Simms 2011; Werner et al. 2014). However, little work has been
done that leverages measurements of traded resources to quantify variation in important
mutualism components like resource exchange, use, and production.
In this study, we quantified the contributions of host and symbiont genotype to resource
exchange and host resource use and production traits in the model mutualism between legumes
and nitrogen-fixing bacteria (rhizobia). The legume-rhizobia mutualism is centered on the
exchange of carbon for nitrogen, whereby the legume host trades a portion of its carbon budget
to rhizobia in exchange for fixed atmospheric nitrogen (Oldroyd et al. 2011; Udvardi and Poole
2013). Quantitative genetic experiments have established that fitness outcomes in the legume-
rhizobia mutualism are highly context-dependent, varying across plant and rhizobia genotypes
as well as environmental factors like resource availability (Heath and Tiffin 2007; Heath 2010;
Weese et al. 2015; Batstone et al. 2017; Burghardt et al. 2017). However, little is known about
how resource exchange, use, and production traits vary in natural populations of legumes and
rhizobia.
Here, we performed a factorial experiment in which we inoculated two plant genotypes
with two rhizobia genotypes and measured the contributions of plant genotype, rhizobia
genotype, and their interaction (GxG) to resource exchange, use, and production traits.
Exchange traits are those directly related to the transfer of nitrogen (benefits) to host plant
tissue and the transfer of carbon (costs) to rhizobia. Resource use and production traits are
those involved in how host plants allocate the nitrogen received from rhizobia and how
efficiently they fix (via photosynthesis) the carbon they transfer to rhizobia. We measured many
of these traits using elemental analysis, and as such, these traits represent the cumulative
endpoint of the dynamic processes. Given that the outcomes of resource exchange are
hypothesized to be governed by bargaining between both interacting partners (Sachs et al.
2004; Akçay and Roughgarden 2007; Clark et al. 2019), we predicted that GxG effects would be
common to exchange traits. On the other hand, we predicted that only plant genotype effects
would be found for resource use and production traits because those traits represent
components of plant physiology that should only be indirectly related to the exchange of
resources among partners. We found extensive differences between plant and rhizobia
genotypes, as well as GxG interactions, for all of the plant traits we measured. Our results
suggest that multiple resource-related traits likely contribute to variation in mutualism fitness
outcomes.
107
108 **Methods**
109
110 *Study system*
111
The mutualism between *Medicago truncatula* and *Ensifer* spp. is a deeply-studied model
system that has been used to uncover the molecular and genetic basis of symbiotic interactions
and to explore the evolution of mutualisms (Udvardi and Poole 2013). *M. truncatula* is an annual
plant native to the Mediterranean Basin (Urban 1873; Lesins and Lesins 1979). *M. truncatula*

forms an endosymbiotic interaction with rhizobia species belonging to the genus *Ensifer*
(Andrews and Andrews 2017). This interaction is localized to specialized structures on the host
plant's roots called nodules. As with other legume-rhizobia interactions, *M. truncatula* trades
photosynthetically-fixed carbon to rhizobia in nodules in exchange for fixed nitrogen. Previous
studies involving the *M. truncatula - Ensifer* mutualism have shown that fitness outcomes in this
interaction depend on the specific combination of plant and rhizobia genotypes (G x G) (Heath
2010). Past studies using *M. truncatula* have primarily used aboveground biomass as a proxy
for fitness.
For this study, we used two inbred lines of *M. truncatula*, HM001 and HM028, sourced
from the Medicago HapMap Project (Stanton-Geddes et al. 2013). These accessions are
derived from collections made in the natural range of *M. truncatula* in the Mediterranean Basin.
We chose these accessions because they, along with many other accessions, display genetic
variation for fitness proxies like aboveground biomass (Stanton-Geddes et al. 2013; Wood et al.
2018). We also used two strains of *Ensifer meliloti*, MAG 154 and MAG 282, sourced from a
panel of 191 strains isolated from *M. truncatula* sampled from the Mediterranean Basin (Katy
Heath, U. Illinois). We selected these strains because they had divergent effects on plant
performance when inoculated on *M. truncatula* genotype HM101. As with the inbred lines of *M.*
truncatula, we chose these strains because they exhibited variable effects on plant fitness
proxies like aboveground biomass (Batstone et al. 2022)**.**
135
136   *Trait measurement*
137
We measured a series of resource exchange, use, and production traits on host plants
(Table 1). We also measured two proxies for mutualism outcome: plant fitness and nodulation
(Table 1). Plant fitness was estimated by weighing the total dry aboveground biomass for an
individual plant. We measured three nodule traits: total nodule number, total nodule mass, and
average nodule size (total nodule number / total nodule mass). By harvesting plants and
measuring aboveground biomass at two different timepoints, we were also able to infer
genotype effects on growth rate. *M. truncatula* populations are thought to be subject to viability
selection in their dry, Mediterranean habitat, thus growth rate could be an important fitness
component (Urban 1873; Lesins and Lesins 1979).
Unless otherwise noted, all resource exchange, use, and production traits were
measured using carbon and nitrogen elemental composition data collected via combustion
analysis in the Department of Earth and Environmental Sciences at the University of
Pennsylvania. To perform combustion analysis, tissue was pulverized using either a mechanical
ball grinder or a mortar and pestle. Then, approximately 4 mg of pulverized tissue was weighed
using an ultra-micro balance (CP2P Microbalance, Sartorius GmbH, Goettingen, Germany),
packaged into tin capsules, and ran on a Costech ECS4010 Elemental Analyzer (Costech
Analytical Technologies, Inc. Valencia, CA, USA). The elemental analyzer returns
measurements of carbon and nitrogen composition (%) for the tissue sample analyzed. We
multiplied carbon and nitrogen composition by the total dry mass that the elemental analysis
sample came from to infer the amount of total biomass carbon and nitrogen in a given sample
(Table 1). We performed combustion analysis separately on one aboveground plant tissue
sample and one nodule tissue from between 6 and 10 individual plants from each genotype

Table 1: The mutualism outcome and resource exchange, use, and production traits measured in this study. Brief descriptions of how each trait was measured and calculated are also provided.

Mutualism component		Trait	Measurement / Calculation
Outcome	Plant fitness	Aboveground biomass	Dry weight of aboveground plant tissue
		Aboveground biomass growth rate	Dry weight of aboveground plant tissue measured at two timepoints (see methods)
	Nodulation	Total nodule mass	Dry weight of all nodules from an individual plant's root system
		Nodule number	Count of all nodules on individual plant's root system
		Average nodule mass	Total nodule mass / nodule number
Exchange traits	Benefit	Nitrogen composition	Proportion of aboveground plant tissue that is composed of nitrogen (measured via elemental analysis)
		Total nitrogen benefit	Nitrogen composition of aboveground tissue * aboveground biomass
		Total nitrogen benefit per symbiont	Slope of the relationship between total nitrogen benefit and total nodule mass (see "statistical analysis" in methods)
	Cost	Total carbon cost	Carbon composition of nodule tissue * total nodule mass
		Nodule carbon fraction	Total carbon cost / (total carbon cost + (aboveground tissue carbon composition * aboveground biomass)
	Resource exchange ratio	Carbon-for-nitrogen exchange ratio	Total carbon cost / total nitrogen benefit
Resource Use	Nitrogen use efficiency	Carbon:nitrogen ratio in aboveground tissue	Nitrogen composition in aboveground plant tissue / carbon composition in aboveground plant tissue
Resource Production		Leaf photosynthetic rate	See Methods and Supplementary Methods

combination. For elemental analysis on aboveground tissue, we pulverized the entirety of the
aboveground portion of the plant, while for elemental analysis on nodules, we pulverized all of
the nodules that were dissected from the plants root system.
We measured three benefit-associated exchange traits: nitrogen composition (%) in
aboveground tissue (hereafter referred to as nitrogen composition), total nitrogen in
aboveground tissue (hereafter referred to as total nitrogen benefit), and total nitrogen in
aboveground tissue as a function of total nodule mass (hereafter referred to as total nitrogen
benefit per symbiont; see statistical analysis section for more details; Table 1). We measured
two cost-associated exchange traits: total carbon in nodule tissue (hereafter referred to as total
carbon cost), and the fraction of total carbon in aboveground tissue and nodule tissue that is in
nodule tissue (hereafter referred to as nodule carbon fraction). These cost traits incorporate
both the carbon cost of constructing nodules as well as the carbon traded to the symbiont. This
approach is unable to quantify the portion of carbon transferred to rhizobia that was respired by
rhizobia, thus our analysis is making the simplifying assumption that respiration rates are
consistent across genotype combinations. We also calculated the ratio of total nitrogen in
aboveground tissue to total carbon in nodule tissue as a proxy of the carbon-for-nitrogen
exchange ratio.
We measured one resource use trait and one resource production trait. The resource
use trait was measured as the ratio of total carbon to total nitrogen in aboveground tissue as a
proxy for nitrogen use efficiency (Dovrat et al. 2020). The benefit production trait we measured
was leaf photosynthesis rate. Photosynthesis rate measurements were made using a LI-6400
Portable Photosynthesis System (LI-COR, Inc., Lincoln, NE, USA) with an 6400-02B LED Light
Source (LI-COR, Inc., Lincoln, NE, USA) attached to the leaf chamber to apply a consistent
level of light for all measurements. See the supplementary methods for details on how
photosynthesis rate was measured and calculated (Supplementary methods). Table 1 provides
short summaries of how each resource exchange, use, and production trait was quantified.
It should be emphasized that resource exchange, use, and production are dynamic and
interrelated processes. For example, how efficiently resources are produced can have knock on
effects on resource exchange and vice versa. In this study we have designated measured traits
as belonging to one of resource exchange, use, and production, but this does not necessarily
mean that variation in these traits is related to that process alone. Precise attribution of traits to
specific mutualism processes is a general challenge for the field of mutualism evolutionary
ecology as a whole and will emerge through more manipulative experiments.
193
194 *Experimental design*
195
To quantify the effect of plant genotype, rhizobia genotype, and GxG interactions on the
traits we measured, we grew our two *M. truncatula* accessions in fully factorial design with our
two *E. meliloti* strains for a total of four plant-rhizobia genotype combinations. Each of the four
genotype combinations was replicated 33 times totalling 132 experimental units.
Seeds were scarified, sterilized, imbibed, and stratified using the methods reported in
the "Seed storage and germination" chapter of the Medicago Handbook (Noble Institute;
Supplementary methods). Prior to planting, we filled 66 mL Cone-tainer pots (Stuewe & Sons,
Inc., Tangent, OR, USA) with a growing media consisting of a 1:4 mix of perlite and sand. We

sterilized the filled Cone-tainer by autoclaving them twice. One seedling was placed in each
Cone-tainer. One day after planting, seedlings were given 1 mL of 0.625 mM nitrogen Fahraeus
fertilizer (Batstone et al. 2017). Afterwards, plants were given 0.0 mM N Fahraeus fertilizer
every 2 to 3 days to ensure that rhizobia were the only source of nitrogen (Batstone et al. 2017).
Cone-tainer pots were placed in 98-well racks. To avoid the risk of cross contamination
of rhizobia strains, every plant in a rack received the same rhizobia strain. Each rack was
enclosed in a semi-permeable tent consisting of plastic wrap and PVC pipe to further reduce the
risk of cross contamination. Plants were grown in two growth chambers in a complete block
design. The height of light fixtures in both chambers was adjusted so that a photosynthetic
active radiation (PAR) reading of ~250 $\mu mol/m^{-2}/s^{-1}$ was consistently measured on the shelf of
the growth chambers. Chambers were set to 22°C and a 16:8 light:dark photoperiod. Seedings
were maintained in a high humidity environment for about one week, after which they were
exposed to ambient humidity, usually 40-50%.
Two weeks after planting, we inoculated plants with rhizobia. Plants received either
strain MAG 154 or MAG 282. We grew strains from glycerol stocks in sterile tryptone yeast
liquid media in a shaking incubator at 30 °C and 180 rpm for approximately two days. We then
streaked the media on sterile Petri plates containing tryptone yeast agar under a laminar flow
hood. We incubated the plates at 30°C until colonies started to form. Next, we selected an
individual colony from each strain to grow in tryptone yeast liquid media. Again, rhizobia cultures
were allowed to grow for approximately two days. We diluted the liquid media to an OD_{600} of 0.1
and inoculated each plant with 1 ml of the diluted culture.
Plants were harvested in three separate cohorts at 9, 10, and 11 weeks post planting.
Approximately 20 plants from each treatment group were harvested at 11 weeks. Of those
plants harvested at 11 weeks, approximately 10 from each treatment were used to measure the
resource exchange, use, and production traits described above via elemental analysis.
Approximately 3 plants from each treatment group were harvested at 10 weeks to measure
photosynthesis. Prior to harvest at the 10 week timepoint, photosynthesis rate measurements
were taken on 2-3 leaves of each plant. Approximately 10 plants from each treatment group
were harvested at 9 weeks. Plants in the 9-week cohort were compared to those in the 11-week
cohort to estimate growth rate (see statistical analysis section for more details).
At each harvest, we gently removed plants from their Cone-tainer pots and cut the
shoots from the roots. The shoots were placed in brown paper bags and dried at 60 °C for at
least a week before being weighed with an analytical scale to measure aboveground biomass.
Plants used for benefit and cost trait measurement via elemental analysis were processed as
described in the "Trait measurement" section. The roots were washed before being placed in
plastic bags and stored in a freezer at -20 °C.
To collect nodulation data, frozen roots were thawed and nodules were removed under a
dissecting microscope. Nodules were counted as they were removed to quantify nodule
number. To assess the consistency of nodule count estimates across researchers, we had each
researcher count nodules on the same subset of 10 plants. Nodule counts were highly
repeatable across researchers in a linear mixed effects model with nodule number as the
response, and plant and researcher as random effects (researcher explained only 0.4% of
variation in nodule counts). Removed nodules were placed in coin envelopes, dried at 60 °C for
at least a week, and weighed. Nodules from plants that were used for benefit and cost trait

measurement via elemental analysis were processed as described in the "Trait measurement"
section.
250
251   *Statistical analysis*
252
We used linear models to test for an effect of plant genotype, rhizobia genotype, and
plant genotype by rhizobia genotype (GxG) interactions on the traits we measured. All models
used a normal error distribution except for the model with nodule number as the response
variable, which used a negative binomial error distribution. Block was included as a fixed effect
in all models. Continuous predictors were mean-centered to avoid spurious inference on main
effects in the presence of interactions (Schielzeth 2010). For some models, response variables
were log-transformed to better conform with the assumption of normality. If a response variable
was log transformed, it will be indicated in the figure caption of the associated figure. For
analysis of elemental composition measurements (e.g. nitrogen composition), arcsine square
root transformations were performed to normalize the data prior to running the models. Models
with nodule traits as the response included total root mass as a covariate to account for
differences in nodule traits associated with differences in root system size.
To test for genotype effects on growth rate, we included the following interaction terms in
the model, plant genotype x timepoint, rhizobia genotype x timepoint, and plant genotype x
rhizobia genotype x timepoint.
To test for significant genotype effects on total nitrogen benefit per symbiont, we
included the interaction terms plant genotype x total nodule mass, rhizobia genotype x total
nodule mass, and plant genotype x rhizobia genotype x total nodule mass terms in our models.
Each of these interaction terms estimates differences in the slope of the relationship between
total nitrogen in aboveground tissue and nodule mass that are due to genotype effects.
Our model with photosynthesis rate as the response included each individual leaf
measurement as an independent replicate. To account for any variation due to measurements
made on leaves from the same plant, we included plant ID as a random effect.
The significance of model terms in each model was tested using type III sum of squares.
For linear models, significant testing was performed using *F*-statistics, and for generalized linear
models, significance testing was performed with Wald χ^2 tests.
Many of the traits that we measured are correlated with one another (Fig. S1) These
correlations could stem from a physiological relationship or a mathematical relationship resulting
from an overlap in the measurements used to calculate a given trait. We caution against
drawing robust interpretations from these correlations because of the widespread mathematical
non-independence, but we do suggest areas of future research that these correlations could
inform (see discussion).
All statistics and data visualization were performed in R v4.2.3 (R Core Team 2023). For
simple linear models, we used the base function lm; for linear mixed effects models we used the
lmer function in the lme4 package (Bates et al. 2014), and for generalized linear mixed effects
models we used the glmmtmb function in the glmmTMB package (Brooks et al. 2017). Model
assumptions were evaluated using the DHARMa package (Hartig and Lohse 2022). Significant
testing was performed with the Anova function in the car package (Fox and Weisberg 2018). All
plots include estimated marginal means that were calculated using the ggeffects package

(Lüdecke 2018). The ggplot2 (Wickham 2016), ghibli (Henderson et al. 2022), and ggpubr
(Kassambara 2023) packages were used to make all of the plots presented here. All data and
code used in this study can be found here [Dryad repository will be added if/when manuscript is
accepted for publication]
296
297 **Results**
298
299 *Genetic variation in fitness and mutualism outcomes*
300
We found a significant plant genotype x rhizobia genotype (GxG) interaction effect for
aboveground biomass, our proxy for plant fitness, but no main effects of either plant or rhizobia
genotype (Fig. 1, Table 2). We also found significant plant genotype, rhizobia genotype, and
GxG effects on aboveground biomass growth rate (Fig. 1, Table 2). During the analysis of
growth rate, we identified an outlier in the HM028 - MAG 282 - Timepoint 1 treatment group.
Removal of this outlier does not change the statistical significance of any model terms.
We observed a significant rhizobia genotype effect on total nodule mass, and significant
plant genotype effects and rhizobia genotype effects on total nodule number and average
nodule size (Fig. 1, Table 2).
310
311 *Genetic variation in exchange traits*
312
We measured three benefit-associated exchange traits, nitrogen composition, total
nitrogen benefit, and total nitrogen benefit per symbiont (Table 1). We observed a significant
rhizobia genotype effect and GxG effect for nitrogen composition in aboveground tissue (Fig. 2,
Table 3), but no significant genotype effects for total nitrogen biomass in aboveground tissue
(Fig. 2, Table 3). For total nitrogen per symbiont, we observed significant plant genotype x total
nodule mass, rhizobia genotype x total nodule mass, and G x G x total nodule mass effects,
indicating that our measure of benefit per symbiont varies in response to plant genotype,
rhizobia genotype, and the interaction of plant genotype and rhizobia genotype (Fig. 2, Table 3).
We measured two cost-associated exchange traits, total carbon cost, and the nodule
carbon fraction (Table 1). We observed a significant rhizobia genotype and GxG effect for total
carbon cost, but no significant genotype effects for nodule carbon fraction (Fig. 3, Table 3).
We also quantified the nitrogen-for-carbon exchange ratio by dividing total nitrogen
benefit by the total carbon cost (Table 1). We observed a significant rhizobia genotype effect for
the nitrogen-for-carbon exchange (Fig. 4, table 3).
327
328 *Genetic variation in resource use and production traits*
329
We measured variation in two traits associated with resource use and production, the
carbon:nitrogen ratio in aboveground tissue (a proxy for nitrogen use efficiency) and leaf
photosynthetic rate (Table 1). The carbon:nitrogen ratio in aboveground tissue was used to infer
how efficiently host plants are able to use nitrogen received through trade to incorporate carbon
in aboveground tissue (Dovrat et al. 2020). Leaf photosynthetic rate to infer the efficiency of the
production of carbon, some of which is ultimately traded away. We found a significant effect of

rhizobia genotype and the interaction between rhizobia genotype and plant genotype on
carbon:nitrogen ratio (Fig. 5, Table 3). We observed a significant plant genotype x rhizobia
genotype interaction on photosynthesis rate (Fig. 5, Table 3).
339
340 **Discussion**
341
342   In this study, we found that mutualism resource exchange, use, and production traits are
343 influenced by both plant and rhizobia genotype. Consistent with our expectations, we observed
344 rhizobia genotype and GxG effects on both benefit-associated and cost-associated resource

Figure 1: Genetic variation for (A) log transformed aboveground biomass, our proxy for plant fitness, (B) log transformed aboveground biomass growth rate, (C) total nodule mass, (D) nodule number, and (E) average nodule size. In all panels, transparent points are the raw data and opaque points are the estimated marginal means. Error bars are the 95% confidence interval. Significant genotype effects are listed on the figure: G_P = plant genotype effect, G_R = rhizobia genotype effect, $G_P x G_R$ = plant genotype x rhizobia genotype interaction, G_P x T = plant genotype x timepoint interaction, $G_R x T$ = rhizobia genotype x timepoint interaction, and $G_P x G_R x T$ = plant genotype x rhizobia genotype x timepoint interaction. Model term statistical significance did not change when the outlier in the HM028-MAG282-Timepoint 1 treatment group in panel B was removed.

Table 2: Genotype effects on mutualism outcome traits. Significant model terms are highlighted in gray.

	Outcomes									
	Plant Fitness				Nodulation					
	Above-ground biomass		Above-ground biomass growth rate		Total nodule mass		Nodule number		Average nodule mass	
	$F_{1,81}$	P	$F_{1,117}$	P	$F_{1,50}$	P	$\chi^2_{df=1}$	P	$F_{1,50}$	P
Rhizobia genotype (G_R)	1.09	0.3	**15.0**	**0.0002**	**7.17**	**0.004**	**16.74**	**<0.0001**	**5.41**	**0.024**
Plant genotype (G_P)	0.18	0.7	**20.76**	**<0.0001**	0.0004	0.98	**3.97**	**0.046**	**5.62**	**0.02**
G_R x G_P	**4.25**	**0.04**	**30.21**	**<0.0001**	3.96	0.05	3.25	0.071	0.002	0.96
Timepoint (T)	-	-	**64.9**	**<0.0001**	-	-	-	-	-	-
T x G_R	-	-	**13.28**	**0.0005**	-	-	-	-	-	-
T x G_P	-	-	**12.96**	**0.0004**	-	-	-	-	-	-
T x G_R x G_P	-	-	**29.81**	**0.0001**	-	-	-	-	-	-

exchange traits. However, we only observed a rhizobia genotype effect for the nitrogen-for-carbon exchange ratio. Contrary to our expectation, rhizobia genotype affected plant resource use and production traits as much or more than plant genotype. However, this result could also be a result of the small number of genotypes we used in our study. Our results suggest that rhizobia genetic variation shapes resource use and production traits in their hosts. Furthermore, our work indicates that the evolution of mutualisms may not be limited to resource exchange traits. Rather, how efficiently mutualistic partners can integrate the benefits of trade (i.e. resource use efficiency) and/or how efficiently they can produce the resource traded away may be just as important for determining the evolution of mutualisms. Together, these results suggest an important role for jointly determined traits in mediating mutualism evolution.

Genotype effects on resource exchange traits

We predicted that GxG effects, which occur when the genotype effect on trait expression in one partner depends on the genotype of the other partner, would be common to exchange traits because resource exchange involves cooperation between the plant and rhizobia (Sachs et al. 2004). Consistent with this prediction, we observed GxG effects for two benefit-associated traits: nitrogen composition and total nitrogen benefit per symbiont. For cost-associated traits we found GxG effects for total carbon cost. However, we did not observe a GxG effect for the

Table 3: Genotype effects on resource exchange, use, and production traits. Significant model terms are highlighted in gray.

	Exchange traits												Resource use		Resource production	
	Benefits						Costs				Benefit: cost					
	Nitrogen composition		Total nitrogen benefit		Total nitrogen benefit per symbiont		Total carbon cost		Nodule carbon fraction		Nitrogen-for-carbon exchange ratio		Nitrogen use efficiency		Leaf photosynthesis rate	
	$F_{1,36}$	P	$F_{1,36}$	P	$F_{1,29}$	P	$F_{1,26}$	P	$F_{1,35}$	P	$F_{1,30}$	P	$F_{1,36}$	P	$\chi^2_{df=1}$	P
Rhizobia genotype (G_R)	**11.48**	**0.002**	0.29	0.6	**7.54**	**0.01**	**5.04**	**0.03**	1.53	0.223	**5.6**	**0.02**	**6.72**	**0.01**	1.73	0.19
Plant genotype (G_P)	1.12	0.3	0.03	0.85	0.35	0.56	0.006	0.94	1.83	0.19	0.005	0.95	1.46	0.23	0.74	0.39
G_R x G_P	**6.3**	**0.02**	0.31	0.58	2.58	0.12	**4.19**	**0.049**	0.0003	0.99	1.88	0.18	**5.78**	**0.02**	**4.62**	**0.03**
Centered total nodule mass (tnm)	-	-	-	-	1.64	0.21	-	-	-	-	-	-	-	-	-	-
Tnm x G_R	-	-	-	-	**9.68**	**0.004**	-	-	-	-	-	-	-	-	-	-
Tnm x G_P	-	-	-	-	**4.63**	**0.04**	-	-	-	-	-	-	-	-	-	-
Tnm x G_P x G_R	-	-	-	-	**8.69**	**0.006**	-	-	-	-	-	-	-	-	-	-

Figure 2: Genetic variation for three benefit-associated exchange traits: (A) nitrogen composition (%), (B) total nitrogen benefit (g), and (C) total nitrogen benefit per symbiont. In panels A and B, transparent points are the raw data, opaque points are the estimated marginal means, and Error bars are 95% confidence intervals. For panel C, transparent points are the raw data, lines represent the estimated marginal slopes derived from the model, and ribbons are the 95% confidence interval for the slope. Significant genotype effects are listed on the figure: G_P = planet genotype effect, G_R = rhizobia genotype effect, $G_P \times G_R$ = plant genotype x rhizobia genotype interaction, $G_P \times TNM$, $G_R \times TNM$, $G_P \times G_R \times TNM$.

carbon-for-nitrogen exchange ratio. Our prediction that GxG effects should be common to
exchange traits is based on the hypothesis that the outcomes of trade in resource mutualisms
are governed by bargaining between partners (Sachs et al. 2004; Akçay and Roughgarden
2007). The GxG effects on exchange traits that we observed here could be mediated by the
numerous bargaining strategies that hosts and microbes employ. For example, legumes are
able to "sanction" poor quality partners resulting in beneficial trading outcomes (Kiers et al.
2003; Westhoek et al. 2021). Alternatively, GxG effects on exchange traits may also emerge via
a mismatch between partners (Barrett et al. 2012; Wendlandt et al. 2019). Partner mismatches
occur when some rhizobia are more effective partners for some plant genotypes compared to
others.
To our surprise, however, the carbon-for-nitrogen exchange rate was only affected by
rhizobia genotype, not by both partners. This result suggests that rhizobia may exert more
control over the overall outcome of trade than their host. This finding conflicts with theoretical
and empirical work that suggests that, in general, host plants exert more power in determining
the outcomes of trade (Daubech et al. 2017; Clark et al. 2019). Of course, a primary reason that
host plants are expected to exert more control over the trade is their ability to "bargain" with
multiple different microbial strains simultaneously (Akçay and Simms 2011; Westhoek et al.
2021); a scenario that was unable to occur in this study due to our single strain inoculation
design. It is also possible that the two plant genotypes we used in this experiment do not
measurably differ in their ability to manipulate the carbon-for-nitrogen exchange rate and that a
larger panel of plant genotypes would have revealed plant genotype variation for resource
exchange. Future research must include more plant and rhizobia genotypes to better test the
relative control that host plants rhizobia exert over the resource exchange.

Figure 3: Genetic variation for two cost-associated exchange traits: (A) total carbon cost and (B) the nodule carbon fraction. Transparent points are the raw data and opaque points are the estimated marginal means. Error bars are the 95% confidence interval. Significant genotype effects are listed on the figure: G_P = plant genotype effect, G_R = rhizobia genotype effect, and G_P X G_R = plant genotype X rhizobia genotype interaction.

Finally, there were a few exchange traits for which we observed no genotype effects.
Among benefit-associated traits, total nitrogen benefit, our metric of absolute benefit, did not
differ between plant or rhizobia genotypes. The same was true for nodule carbon fraction, our
metric of relative cost. The fact that we observed genotype differences in the relative but not
absolute benefit—and the absolute but not relative cost—suggests that absolute and relative
costs (and benefits) of mutualism may exhibit different evolutionary dynamics. Another
explanation is that there is genetic variation for these traits in nature, but was simply not
captured by our panel of two plant genotypes and two rhizobia genotypes.
396
397 *Genotype effects on resource use and production traits*
398
We predicted that plant genotype would be the sole contributor to differences in plant
resource use and production traits. Contrary to this prediction, we observed GxG effects for both
of the resource use and production traits that we measured: nitrogen use efficiency (our metric
of resource use) and leaf photosynthetic rate (our metric of resource production).
We predicted that host resource use and production traits would respond primarily to
plant genotype because these traits are indirectly related to resource exchange and represent
components of plant physiology. However, rhizobia effects on these host traits are perhaps not
surprising if hosts change how they allocate benefits or invest in the production of resources in
response to trade. Thus, rhizobia and GxG effects on resource use and production traits may
arise from the GxG effects for exchange traits that we observed. As mentioned above, it is also
possible that a lack of plant genotype effect is simply due to a lack of measurable differences in
resource use and production traits between the two plant genotypes we used here.

Figure 4: **Genetic variation for the nitrogen-for-carbon exchange ratio.**
Transparent points are the raw data and opaque points are the estimated marginal
means. Error bars are the 95% confidence interval. Significant genotype effects are
listed on the figure: G_P = planet genotype effect, G_R = rhizobia genotype effect, and
G_P X G_R = plant genotype X rhizobia genotype interaction.

Resource use traits, like nitrogen use efficiency, have previously been shown to be
microbially mediated (Dovrat et al. 2020; Zhang et al. 2020). For instance, legumes modify their
nitrogen use efficiency in response to both soil nitrogen availability and the presence of rhizobia
(Dovrat et al. 2020). Our study raises the important question of why nitrogen use efficiency
could vary across combinations of plant and rhizobia genotypes. A GxG effect on nitrogen use
efficiency could be due to differences in the efficiency of symbiotic nitrogen fixation across plant
and rhizobia combinations. In contrast to some legume species that down- or upregulate
nitrogen fixation in response to environmental conditions *M. truncatula* is an obligate fixer that
attempts to maintain a constant nitrogen fixation effort regardless of the activity of their rhizobia
partners or external supplies of nitrogen (Hedin et al. 2009). Variation in nitrogen use efficiency
in obligate fixers like *M. truncatula* could simply be a function of the variation in the efficiency of
nitrogen fixation across plant and rhizobia genotype combinations. Alternatively, it is also
plausible that legumes may have evolved strategies to adjust how they use and allocate the
benefits from trade in order to buffer against the consequences of interacting with poor quality
partners.
Resource production traits, like leaf photosynthetic rate, also have been shown to
respond to microbes (Adams et al. 2016). Legumes in particular exhibit significantly higher
photosynthetic rates in the presence versus the absence of rhizobia (Clark et al. 2019; Dovrat et
al. 2020). Photosynthesis is fundamental to the exchange of resources between legumes and
rhizobia because it provides the carbon necessary to support rhizobia and it also requires a

Figure 5: Genetic variation for (A) carbon : nitrogen ratio in aboveground tissue and (B) leaf photosynthetic rate. Transparent points are the raw data and opaque points are the estimated marginal means. Error bars are the 95% confidence interval. Significant genotype effects are listed on the figure: G_P = planet genotype effect, G_R = rhizobia genotype effect, and $G_P \times G_R$ = plant genotype X rhizobia genotype interaction.

large amount of nitrogen, which is provided by the rhizobia. A heightened photosynthetic rate may equate to a larger pool of carbon that can be devoted to the mutualism, and by extension, a large supply of nitrogen that can be returned to the host plant. Variation in photosynthetic rate may also simply be an indication of nitrogen limitation and the outcomes of trade. Photosynthesis involves nitrogen-rich machinery and therefore, genotype combinations with the highest photosynthetic rates may simply be the genotype combinations in which the largest amount of nitrogen is being transferred to the host plant. Regardless of the mechanisms responsible for GxG effects on photosynthetic rate, the fact that a portion of the genetic variation for leaf photosynthetic rate resides in rhizobia emphasizes the far-reaching effect that this mutualism has on host plant physiology.

The findings of this study underscore the possibility that the evolution of mutualisms may involve traits related to how interacting partners can integrate the benefits of trade (i.e. resource use efficiency) and how efficiently they can produce the resources traded away. This suggestion is corroborated by a recent study involving a synthetic mutualism between engineered strains of brewer's yeast (Vidal and Segraves 2021). The authors observed strong fluctuations in both resource use efficiency and resource production of both strains as they evolved in response to the availability of the limiting resources produced by their partners. Resource use and production traits may represent key components of mutualisms that allow interacting partners the ability to adjust their physiology in response to the outcomes of trade. Future studies should carefully consider selection on all mutualism traits, not just those related to the exchange of resources, as well as covariation among resource exchange, use and production traits.

Resource exchange, use, and production are all dynamic and interrelated processes. An
important challenge for the study of mutualisms is to better understand how each of these
processes influences the other. The widespread pairwise correlations among the traits we
quantified in the study suggests how tightly interwoven these processes are (Fig. S1). They also
raise the intriguing possibility that genotype effects on one trait can cause correlated changes in
the expression of genetic variation for other traits. However, as stated in the methods, many of
the traits we measured here are not mathematically independent owing to the fact that there is
some overlap in the variables used to calculate the traits we examined in the study. Future work
is needed to carefully isolate the different components of mutualisms to quantify their effects on
one another.
462
463   *The importance of jointly determined traits to predicting evolution in mutualism*
464
The ubiquity of rhizobia and GxG effects on mutualism traits has three important
implications for the evolution of mutualisms. First, this finding indicates that traits involved in
resource exchange, use, and production may be jointly determined by the genomes of both
interacting partners, just as is the case with fitness outcomes (Heath 2010). Jointly determined
traits, here defined as traits that exhibit variation in response to more than one genome as either
two independent main effects or a GxG effect, can have different evolutionary trajectories than
those determined by the genome of just one or the other interacting partner (O'Brien et al.
2021b). Thus, a trait-based framework, wherein quantitative genetic approaches can be
leveraged to identify the sources of variation in mutualism traits, is necessary to develop a
better understanding of mutualism evolution.
Second, these results add to the growing body of literature demonstrating that microbial
mutualists play an important role in the expression of host traits like resource use and
production. Microbes are known to mediate host nutrient acquisition (Hestrin et al. 2019; Harbort
et al. 2020), life history transitions (Wagner et al. 2014; O'Brien et al. 2021a), defense against
antagonists (Johnson et al. 1987; Berg and Koskella 2018), stress tolerance (Lau and Lennon
2012), and morphology (Ortíz-Castro et al. 2008; Batstone et al. 2017). Our work builds on this
knowledge by demonstrating that intraspecific variation in microbial populations, in addition to
interspecific variation, can also influence the expression of plant functional traits. This finding
complements two recent studies, O'Brien et al. (2021b) and De Lisle et al. (2022), which have
highlighted the potential evolutionary significance of the effect of intraspecific partner variation in
the expression of mutualism traits. Crucially, intraspecific microbial variation is invisible to the
16S sequencing that is commonly applied to characterize community composition in microbiome
studies because strains belonging to the same species have identical 16S sequences. Thus,
substantial cryptic variation in microbe-mediated plant traits measured in microbiome
manipulations may go unnoticed.
Third, variation in resource trade, use, and production traits across different
combinations of plant and rhizobia genotypes highlights the potential importance of the non-
random association of partner genotypes for mutualism evolution. If a mutualism trait responds
to a GxG effect and is a target of selection, then the plant genotypes favored by selection will be
dependent on which rhizobia genotype they are paired with. If associations between specific
plant and rhizobia genotypes are random, then the response to selection will be smaller than

expected because most genetic variation is non-additive and a function of which genotypes are paired together. On the other hand, if associations between specific plant and rhizobia genotypes are non-random, then the response to selection could be either larger (if associations between plant and rhizobia genotypes inflate genetic variation for the trait) or smaller (if associations between plant and rhizobia genotypes constrain genetic variation for the trait) than expected. While host plants have a clear influence on their rhizosphere microbes (Miranda-Sánchez et al. 2016; Greenlon et al. 2019) and can vary in their ability to select beneficial microbial genotypes (Burghardt et al. 2018, 2019), the frequency of non-random associations between specific plant and rhizobia genotypes in nature has been rarely measured. Given the apparent importance of rhizobia genotypes in influencing variation in mutualism traits, more studies are needed to address the degrees of non-random associations between interacting genotypes in order to develop a predictive framework for understanding mutualism evolution.

Ultimately, the relative importance of genotype and GxG variation for mutualism traits to mutualism evolution is dependent on the contribution of that variation to fitness. An important next step to expand our understanding of mutualisms is to quantify the relationship between resource exchange, use, and production traits and fitness, i.e. selection. While we did measure common fitness proxies for this system, such as plant aboveground biomass, these fitness proxies were also used in the calculation of most of the resource exchange, use, and production traits presented in this study. The mathematical non-independence between our fitness proxies and resource exchange, use, and production traits precluded us from being able to measure selection. Measurements of selection will reveal both the traits that underpin mutualism variation and will allow us to begin dissecting the selective agents that maintain fitness variation in mutualisms.

Acknowledgements
We thank Chang-Yu Chang, Addison Martin, and Eunnuri Yi for helpful comments on this manuscript. We also thank Chigozie Ibe, Anisa Robinson, Linda Wu, and Seokyoon Chang for help with executing the experiment and collecting data. *Medicago truncatula* genotypes were generously provided by Nevin Young and the Medicago HapMap project. *Ensifer meliloti* strains were generously provided by Liana Burghardt and Katy Heath. Growth chamber logistical support was provided by the University of Pennsylvania's Greenhouse support staff members Kathyrn Butler and Samara Gray. We are grateful to David Vann and the University of Pennsylvania's Earth and Environmental Science Department's Instrumentation Core for assistance with elemental analysis. This work was funded by a Society for the Study of Evolution R. C. Lewontin Award to McCall Calvert and NSF-DEB 2118397 to Corlett Wood

References

Adams, M. A., T. L. Turnbull, J. I. Sprent, and N. Buchmann. 2016. Legumes are different: Leaf nitrogen, photosynthesis, and water use efficiency. Proc. Natl. Acad. Sci. 113:4098–4103. Proceedings of the National Academy of Sciences.
Akçay, E., and J. Roughgarden. 2007. Negotiation of mutualism: rhizobia and legumes. Proc. R. Soc. B Biol. Sci. 274:25–32.
Akçay, E., and E. L. Simms. 2011. Negotiation, Sanctions, and Context Dependency in the

Legume-Rhizobium Mutualism. Am. Nat. 178:1–14.

Albrecht, M., B. Schmid, Y. Hautier, and C. B. Müller. 2012. Diverse pollinator communities
enhance plant reproductive success. Proc. R. Soc. B Biol. Sci. 279:4845–4852.

Andrews, M., and M. E. Andrews. 2017. Specificity in Legume-Rhizobia Symbioses. Int. J. Mol.
Sci. 18:705.

Barrett, L. G., L. M. Broadhurst, and P. H. Thrall. 2012. Geographic adaptation in plant–soil
mutualisms: tests using Acacia spp. and rhizobial bacteria. Funct. Ecol. 26:457–468.

Bates, D., M. Mächler, B. Bolker, and S. Walker. 2014. Fitting Linear Mixed-Effects Models
using lme4. arXiv.

Batstone, R. T., E. M. Dutton, D. Wang, M. Yang, and M. E. Frederickson. 2017. The evolution
of symbiont preference traits in the model legume *Medicago truncatula*. New Phytol.
213:1850–1861.

Batstone, R. T., H. Lindgren, C. M. Allsup, L. A. Goralka, A. B. Riley, M. A. Grillo, A. Marshall-
Colon, and K. D. Heath. 2022. Genome-wide association studies across environmental
and genetic contexts reveal complex genetic architecture of symbiotic extended
phenotypes. BioRxiv 2021–08. Cold Spring Harbor Laboratory.

Berg, M., and B. Koskella. 2018. Nutrient- and Dose-Dependent Microbiome-Mediated
Protection against a Plant Pathogen. Curr. Biol. 28:2487-2492.e3.

Bronstein, J. L. 2001. The Costs of Mutualism1. Am. Zool. 41:825–839.

Bronstein, J. L. 2015. The study of mutualism. P. 0 *in* J. L. Bronstein, ed. Mutualism. Oxford
University Press.

Brooks, M. E., K. Kristensen, K. J. van Benthem, A. Magnusson, C. W. Berg, A. Nielsen, H. J.
Skaug, M. Machler, and B. M. Bolker. 2017. glmmTMB balances speed and flexibility
among packages for zero-inflated generalized linear mixed modeling. R J. 9:378–400.
Technische Universitaet Wien.

Burghardt, L. T., B. Epstein, J. Guhlin, M. S. Nelson, M. R. Taylor, N. D. Young, M. J.
Sadowsky, and P. Tiffin. 2018. Select and resequence reveals relative fitness of bacteria
in symbiotic and free-living environments. Proc. Natl. Acad. Sci. 115:2425–2430.

Burghardt, L. T., B. Epstein, and P. Tiffin. 2019. Legacy of prior host and soil selection on
rhizobial fitness in planta. Evolution 73:2013–2023.

Burghardt, L. T., J. Guhlin, C. L. Chun, J. Liu, M. J. Sadowsky, R. M. Stupar, N. D. Young, and
P. Tiffin. 2017. Transcriptomic basis of genome by genome variation in a legume-
rhizobia mutualism. Mol. Ecol. 26:6122–6135.

Clark, T. J., C. A. Friel, E. Grman, M. L. Friesen, and Y. Shachar-Hill. 2019. Unfair trade
underground revealed by integrating data with Nash bargaining models. New Phytol.
222:1325–1337.

Daubech, B., P. Remigi, G. Doin de Moura, M. Marchetti, C. Pouzet, M.-C. Auriac, C. S.
Gokhale, C. Masson-Boivin, and D. Capela. 2017. Spatio-temporal control of mutualism
in legumes helps spread symbiotic nitrogen fixation. eLife 6:e28683. eLife Sciences
Publications, Ltd.

De Lisle, S. P., D. I. Bolnick, E. D. Brodie III, A. J. Moore, and J. W. McGlothlin. 2022.
Interacting phenotypes and the coevolutionary process: Interspecific indirect genetic
effects alter coevolutionary dynamics. Evolution 76:429–444.

Dovrat, G., H. Bakhshian, T. Masci, and E. Sheffer. 2020. The nitrogen economic spectrum of
legume stoichiometry and fixation strategy. New Phytol. 227:365–375.

Fox, J., and S. Weisberg. 2018. An R Companion to Applied Regression. 3rd ed. Sage
Publications, Inc., Washington DC.

Friesen, M. L., S. S. Porter, S. C. Stark, E. J. von Wettberg, J. L. Sachs, and E. Martinez-
Romero. 2011. Microbially Mediated Plant Functional Traits. Annu. Rev. Ecol. Evol. Syst.
42:23–46.

Greenlon, A., P. L. Chang, Z. M. Damtew, A. Muleta, N. Carrasquilla-Garcia, D. Kim, H. P.

Nguyen, V. Suryawanshi, C. P. Krieg, S. K. Yadav, J. S. Patel, A. Mukherjee, S. Udupa, I. Benjelloun, I. Thami-Alami, M. Yasin, B. Patil, S. Singh, B. K. Sarma, E. J. B. von Wettberg, A. Kahraman, B. Bukun, F. Assefa, K. Tesfaye, A. Fikre, and D. R. Cook. 2019. Global-level population genomics reveals differential effects of geography and phylogeny on horizontal gene transfer in soil bacteria. Proc. Natl. Acad. Sci. 116:15200–15209. Proceedings of the National Academy of Sciences.

Grman, E., T. M. P. Robinson, and C. A. Klausmeier. 2012. Ecological Specialization and Trade Affect the Outcome of Negotiations in Mutualism. Am. Nat. 179:567–581.

Harbort, C. J., M. Hashimoto, H. Inoue, Y. Niu, R. Guan, A. D. Rombolà, S. Kopriva, M. J. E. E. E. Voges, E. S. Sattely, R. Garrido-Oter, and P. Schulze-Lefert. 2020. Root-Secreted Coumarins and the Microbiota Interact to Improve Iron Nutrition in Arabidopsis. Cell Host Microbe 28:825-837.e6.

Hartig, F., and L. Lohse. 2022. DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression Models.

Heath, K. D. 2010. Intergenomic Epistasis and Coevolutionary Constraint in Plants and Rhizobia. Evolution 64:1446–1458.

Heath, K. D., P. V. Burke, and J. R. Stinchcombe. 2012. Coevolutionary genetic variation in the legume-rhizobium transcriptome. Mol. Ecol. 21:4735–4747.

Heath, K. D., and P. Tiffin. 2007. Context dependence in the coevolution of plant and rhizobial mutualists. Proc. R. Soc. Lond. B Biol. Sci. 274:1905–1912.

Hedin, L. O., E. N. J. Brookshire, D. N. L. Menge, and A. R. Barron. 2009. The Nitrogen Paradox in Tropical Forest Ecosystems. Annu. Rev. Ecol. Evol. Syst. 40:613–635.

Henderson, E., D. Desrosiers, and M. Chirico. 2022. ghibli: Studio Ghibli Colour Palettes.

Hestrin, R., E. C. Hammer, C. W. Mueller, and J. Lehmann. 2019. Synergies between mycorrhizal fungi and soil microbial communities increase plant nitrogen acquisition. Commun. Biol. 2:1–9. Nature Publishing Group.

Hoeksema, J. D., and E. M. Bruna. 2015. Context-dependent outcomes of mutualistic interactions. Pp. 181–202 in J. L. Bronstein, ed. Mutualism. Oxford University Press.

Janzen, D. H. 1966. Coevolution of Mutualism Between Ants and Acacias in Central America. Evolution 20:249–275.

Johnson, N. D., B. Liu, and B. L. Bentley. 1987. The effects of nitrogen fixation, soil nitrate, and defoliation on the growth, alkaloids, and nitrogen levels of Lupinus succulentus (Fabaceae). Oecologia 74:425–431.

Kassambara, A. 2023. ggpubr: "ggplot2" Based Publication Ready Plots.

Kiers, E. T., R. A. Rousseau, S. A. West, and R. F. Denison. 2003. Host sanctions and the legume–rhizobium mutualism. Nature 425:78–81.

Lau, J. A., and J. T. Lennon. 2012. Rapid responses of soil microorganisms improve plant fitness in novel environments. Proc. Natl. Acad. Sci. 109:14058–14062. Proceedings of the National Academy of Sciences.

Lesins, K. A., and I. Lesins. 1979. Genus Medicago (Leguminosae): A Taxogenetic Study. Dr. W. Junk Publishers, The Hague.

Levy, S., A. Elek, X. Grau-Bové, S. Menéndez-Bravo, M. Iglesias, A. Tanay, T. Mass, and A. Sebé-Pedrós. 2021. A stony coral cell atlas illuminates the molecular and cellular basis of coral symbiosis, calcification, and immunity. Cell 184:2973-2987.e18.

Lüdecke, D. 2018. ggeffects: Tidy Data Frames of Marginal Effects from Regression Models. J. Open Source Softw. 3:772.

Miranda-Sánchez, F., J. Rivera, and P. Vinuesa. 2016. Diversity patterns of Rhizobiaceae communities inhabiting soils, root surfaces and nodules reveal a strong selection of rhizobial partners by legumes. Environ. Microbiol. 18:2375–2391.

Moore, A. J., E. D. III Brodie, and J. B. Wolf. 1997. Interacting phenotypes and the evolutionary process: I. Direct and indirect genetic effects of social interactions. Evolution 51:1352–

643   1362.

644 O'Brien, A. M., N. A. Ginnan, M. Rebolleda-Gómez, and M. R. Wagner. 2021a. Microbial effects
645   on plant phenology and fitness. Am. J. Bot. 108:1824–1837.

646 O'Brien, A. M., C. N. Jack, M. L. Friesen, and M. E. Frederickson. 2021b. Whose trait is it
647   anyways? Coevolution of joint phenotypes and genetic architecture in mutualisms. Proc.
648   R. Soc. B Biol. Sci. 288:20202483.

649 Oldroyd, G. E. D., J. D. Murray, P. S. Poole, and J. A. Downie. 2011. The Rules of Engagement
650   in the Legume-Rhizobial Symbiosis. Annu. Rev. Genet. 45:119–144.

651 Ortíz-Castro, R., M. Martínez-Trujillo, and J. López-Bucio. 2008. N-acyl-L-homoserine lactones:
652   a class of bacterial quorum-sensing signals alter post-embryonic root development in
653   Arabidopsis thaliana. Plant Cell Environ. 31:1497–1509.

654 R Core Team. 2023. R: A language and environment for statistical computing. R Foundation fo
655   Statistical Computing, Vienna, Austria.

656 Sachs, J. L., U. G. Mueller, T. P. Wilcox, and J. J. Bull. 2004. The Evolution of Cooperation. Q.
657   Rev. Biol. 79:26.

658 Schielzeth, H. 2010. Simple means to improve the interpretability of regression coefficients.
659   Methods Ecol. Evol. 1:103–113.

660 Stanton-Geddes, J., T. Paape, B. Epstein, R. Briskine, J. Yoder, J. Mudge, A. K. Bharti, A. D.
661   Farmer, P. Zhou, R. Denny, G. D. May, S. Erlandson, M. Yakub, M. Sugawara, M. J.
662   Sadowsky, N. D. Young, and P. Tiffin. 2013. Candidate Genes and Genetic Architecture
663   of Symbiotic and Agronomic Traits Revealed by Whole-Genome, Sequence-Based
664   Association Genetics in Medicago truncatula. PLOS ONE 8:e65688. Public Library of
665   Science.

666 Udvardi, M., and P. S. Poole. 2013. Transport and Metabolism in Legume-Rhizobia Symbioses.
667   Annu. Rev. Plant Biol. 64:781–805.

668 Urban, I. 1873. Prodromus einer monographie der gattung Medicago L. Verh Bot Ver Brand.
669   15:1–85.

670 Vidal, M. C., and K. A. Segraves. 2021. Coevolved mutualists experience fluctuating costs and
671   benefits over time. Evolution 75:219–230.

672 Wagner, M. R., D. S. Lundberg, D. Coleman-Derr, S. G. Tringe, J. L. Dangl, and T. Mitchell-
673   Olds. 2014. Natural soil microbes alter flowering phenology and the intensity of selection
674   on flowering time in a wild Arabidopsis relative. Ecol. Lett. 17:717–726.

675 Weese, D. J., K. D. Heath, B. T. M. Dentinger, and J. A. Lau. 2015. Long-term nitrogen addition
676   causes the evolution of less-cooperative mutualists. Evolution 69:631–642.

677 Wendlandt, C. E., J. U. Regus, K. A. Gano-Cohen, A. C. Hollowell, K. W. Quides, J. Y. Lyu, E.
678   S. Adinata, and J. L. Sachs. 2019. Host investment into symbiosis varies among
679   genotypes of the legume Acmispon strigosus, but host sanctions are uniform. New
680   Phytol. 221:446–458.

681 Werner, G. D. A., J. E. Strassmann, A. B. F. Ivens, D. J. P. Engelmoer, E. Verbruggen, D. C.
682   Queller, R. Noë, N. C. Johnson, P. Hammerstein, and E. T. Kiers. 2014. Evolution of
683   microbial markets. Proc. Natl. Acad. Sci. 111:1237–1244. Proceedings of the National
684   Academy of Sciences.

685 Westhoek, A., L. J. Clark, M. Culbert, N. Dalchau, M. Griffiths, B. Jorrin, R. Karunakaran, R.
686   Ledermann, A. Tkacz, I. Webb, E. K. James, P. S. Poole, and L. A. Turnbull. 2021.
687   Conditional sanctioning in a legume–Rhizobium mutualism. Proc. Natl. Acad. Sci. 118.
688   National Academy of Sciences.

689 Wickham, H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag, New York.

690 Wood, C. W., B. L. Pilkington, P. Vaidya, C. Biel, and J. R. Stinchcombe. 2018. Genetic conflict
691   with a parasitic nematode disrupts the legume–rhizobia mutualism. Evol. Lett. 2:233–
692   245.

693 Zhang, J., N. He, C. Liu, L. Xu, Z. Chen, Y. Li, R. Wang, G. Yu, W. Sun, C. Xiao, H. Y. H. Chen,

694   and P. B. Reich. 2020. Variation and evolution of C:N ratio among different organs
695   enable plants to adapt to N-limited environments. Glob. Change Biol. 26:2534–2543.