Toward a trait-based framework for studying mutualism evolution: genotypic variation in benefit and cost traits in the legume-rhizobia mutualism.

McCall B. Calvert[1*], Maliha Hoque[1], and Corlett W. Wood[1*]

[1]Department of Biology, University of Pennsylvania, Philadelphia, PA, USA

*Correspondence
McCall B. Calvert, calvm@sas.upenn.edu
Corlett W. Wood, corlett@sas.upenn.edu

Abstract

Mutualisms exhibit extensive context-dependent genetic variation in fitness outcomes. However, the specific traits that underpin fitness variation in mutualisms remain understudied. Pinpointing fitness-associated traits in mutualisms is critical to understanding how mutualisms evolve, as well as identifying the mechanisms that maintain genetic variation for fitness in interacting populations. In this study, we quantified the contribution of host and symbiont genotypes to variation in resource exchange, use, and production traits measured in the host using the model mutualism between legumes and nitrogen fixing rhizobia as our study system. We predicted that plant genotype x rhizobia genotype (GxG) effects would be common to resource exchange traits because resource exchange is hypothesized to be governed by input from both interacting partners through bargaining. On the other hand, we predicted that plant genotype effects would be common to host resource use and production traits because these traits are indirectly related to the exchange of resources. Consistent with our prediction for resource exchange traits, but not our prediction for resource use and production traits, we found that rhizobia genotype and GxG effects were the most common sources of variation in the traits that we measured. The results of this study complement the commonly observed phenomenon of GxG effects for fitness by showing that numerous mutualism traits also exhibit GxG variation. Furthermore, they highlight the possibility that the exchange of resources as well as how partners use and produce traded resources can influence the evolution of mutualistic interactions. Future studies can now quantify the relationship between resource exchange, use and production traits and fitness (i.e. selection) to test the competing hypotheses proposed to explain the maintenance of fitness variation in mutualisms.

Introduction

27
Almost all multicellular life engages in some form of mutualism. Mutualisms are
interspecific interactions in which benefits are reciprocally exchanged. The benefits gained from
mutualistic interactions can have strong effects on the fitness of interacting partners, such as
size and fecundity (Bronstein 1994, 2015). Contrary to the expectation that natural selection
favoring the most beneficial partners would erode genetic variation for fitness in mutualism,
there is abundant variation in many mutualisms in the wild (Heath and Stinchcombe 2014). This
body of work has generated important insights into the potential evolutionary forces that drive
mutualism evolution and maintain genetic variation in populations of interacting species
(Thompson 2005; Heath and Stinchcombe 2014). The abundant variation in fitness outcomes in
mutualisms raises an important question: what underlying traits give rise to this variation?
Fitness-related traits, as commonly measured in many quantitative genetic studies on
mutualisms, have been useful for inferring net interaction outcomes and for identifying putative
selective agents. However, the measurement of fitness-related traits also obscures the
individual traits that underlie fitness variation. Pinpointing the traits underlying fitness variation in
mutualisms is key to understanding how mutualisms evolve, as well as the mechanisms that
maintain genetic variation for fitness in interacting populations. Trait-based approaches have
been employed to great effect in plant-pollinator mutualisms to test evolutionary hypotheses
about the forces acting on the underlying traits. For example, a combination of observation and
quantitative genetic experiments in the orchid family has revealed that increased pollen transfer
efficiency precipitated the emergence and maintenance of distinct pollination strategies,
ecological adaptation, and reproductive isolation (Peakall and Beattie 1996; Scopece et al.
2010, 2013, 2015). These studies along with others on plant-pollinator mutualisms, highlight the
power of applying trait-based approaches to studying evolution in mutualisms.
The net benefit of a mutualistic interaction, and by extension, fitness, is determined by
the balance between the benefits and costs associated with the interaction. Therefore, genetic
variation in net benefit could arise from variation in the benefits alone, the costs alone, or both.
Mutualism benefits can take a variety of forms, including transportation (e.g. the movement of
gametes in plant-pollinator mutualisms; Albrecht et al. 2012), protection (e.g. defense against
herbivory in ant-acacia mutualisms; Janzen 1966), and nutrition (e.g. the exchange of limiting
resources between plants and microbes; Friesen et al. 2011). Costs are related to the energetic
investment involved in producing benefits that are transferred to the interacting partner
(Bronstein 2001, 2015). For example, in plant-microbe mutualisms that involve an exchange of
resources, a principal cost for the host is the carbon that is transferred to microbial partners
(Akçay and Roughgarden 2007; Clark et al. 2019) .
Beyond the traits involved in the direct exchange of resources, traits related to the use or
production of the traded resources can also influence the net fitness consequences of a
mutualism (Vidal and Segraves 2021). Contrary to resource exchange traits, which capture the
total benefit and cost, resource use and production traits influence the *relative* benefits and
costs. Traits like resource use efficiency dictate how efficiently the benefits received from an
interacting partner are incorporated into physiological processes (e.g., growth, reproduction,
defense) that are linked to fitness. Resource production traits, on the other hand, are involved in
the creation of the goods and services that are traded to an interacting partner. These traits
dictate the total resource pool available to invest in mutualism, and thereby affect the

proportional cost of resource exchange. In plant-microbe interactions, a common resource
production trait is photosynthetic rate, because it determines the size of a plant's carbon pool, a
portion of which is traded to microbial partners (Grman et al. 2012; Clark et al. 2019).
An individual can achieve a large fitness benefit from mutualism via multiple trait
combinations: by maximizing the benefit it receives, minimizing the cost it pays, maximizing its
resource use efficiency, or modifying its rate of resource production (Vidal and Segraves 2021).
Genetic variation in fitness outcomes in mutualisms could therefore emerge from genotypic
differences in any trait involved in resource exchange, use, or production. An added layer of
complication is that traits mediating species interactions are often jointly determined by both
interacting partners (Moore et al. 1997; O'Brien et al. 2021b; De Lisle et al. 2022). Traits are
jointly determined when variation in a trait is determined by contributions from the genomes of
both interacting partners. Therefore, genetic variation in traits that mediate a host-microbe
mutualism could arise from host genotype effects, microbe genotype effects, or from genotype-
by-genotype (GxG) interactions (Heath 2010; Friesen et al. 2011).
Quantifying genotypic differences in mutualism traits represents an important synthesis
between the subfields of mutualism physiology, functional ecology, and evolution. The traits
involved in resource exchange, use, and production are well defined in many mutualisms,
especially in nutritional mutualisms between hosts and symbionts (Udvardi and Poole 2013;
Levy et al. 2021). For example, some legume species possess the ability to both "choose" the
high quality microbial partners and "sanction" low quality microbial partners (Denison 2000;
Kiers et al. 2003, 2011). Furthermore, there is an extensive body of theoretical work on the
resource economics of mutualism (Akçay and Roughgarden 2007; Akçay and Simms 2011;
Werner et al. 2014). Nevertheless, few empirical studies have investigated whether there is
heritable variation in these traits in interacting populations.
In this study, we quantified the contributions of host and symbiont genotype to host
resource exchange, use, and production traits in the model mutualism between legumes and
nitrogen-fixing bacteria (rhizobia). The legume-rhizobia mutualism is centered on the exchange
of carbon for nitrogen, whereby the legume host trades a portion of its carbon budget to rhizobia
in exchange for fixed atmospheric nitrogen (Oldroyd et al. 2011; Udvardi and Poole 2013). This
symbiosis is a powerful system to explore the traits underlying fitness variation. Quantitative
genetic experiments have established that fitness outcomes in the legume-rhizobia mutualism
are highly context-dependent, varying across plant and rhizobia genotypes as well as
environmental factors like resource availability (Heath and Tiffin 2007; Heath 2010; Weese et al.
2015; Batstone et al. 2017; Burghardt et al. 2017). However, little is known about how resource
exchange, use, and production traits vary in natural populations of legumes and rhizobia and
potentially underlie fitness variation.
Here, we performed a factorial experiment in which we inoculated two plant genotypes
with two rhizobia genotypes and measured the contributions of plant genotype, rhizobia
genotype, and their interaction (GxG) to resource exchange, use, and production traits.
Exchange traits are those directly related to the transfer of nitrogen (benefits) to host plant
tissue and the transfer of carbon (costs) to rhizobia. Resource use and production traits are
those involved in how host plants allocate the nitrogen received from rhizobia and how
efficiently they fix (via photosynthesis) the carbon they transfer to rhizobia. Given that the
outcomes of resource exchange are hypothesized to be governed by bargaining between both

interacting partners (Sachs et al. 2004; Akçay and Roughgarden 2007; Clark et al. 2019), we
predicted that GxG effects would be common to exchange traits. On the other hand, we
predicted that only plant genotype effects would be found for resource use and production traits
because those traits represent components of plant physiology that should only be indirectly
related to the exchange of resources among partners. We found extensive differences between
plant and rhizobia genotypes, as well as GxG interactions, for all of the plant traits we
measured. Our results suggest that multiple resource-related traits likely contribute to variation
in mutualism fitness outcomes, and suggest that mutualism evolution could be driven by
underlying change in resource exchange, use, and production traits.
124
125   **Methods**
126
127   *Study system*
128
The mutualism between *Medicago truncatula* and *Ensifer* spp. is a deeply-studied model
system that has been used to uncover the molecular and genetic basis of symbiotic interactions
and to explore the evolution of mutualisms (Udvardi and Poole 2013). *M. truncatula* is an annual
plant native to the Mediterranean Basin(Urban 1873; Lesins and Lesins 1979). *M. truncatula*
forms an endosymbiotic interaction with rhizobia species belonging to the genus *Ensifer*
(Andrews and Andrews 2017). This interaction is localized to specialized structures on the host
plant's roots called nodules. As with other legume-rhizobia interactions, *M. truncatula* trades
photosynthetically-fixed carbon to rhizobia in nodules in exchange for fixed nitrogen. Numerous
previous studies involving the *M. truncatula - Ensifer* mutualisms have observed genotype-
dependent variation in fitness in response to different genotypes of the interacting partner (G x
G) (Heath 2010). Past studies using *M. truncatula* have primarily used aboveground biomass as
a proxy for fitness.
For this study, we used two inbred lines of *M. truncatula*, HM001 and HM028, sourced
from the Medicago HapMap Project (Stanton-Geddes et al. 2013). These accessions are
derived from collections made in the natural range of *M. truncatula* in the Mediterranean Basin.
We chose these accessions because they, along with many other accessions, display genetic
variation for fitness proxies like aboveground biomass (Stanton-Geddes et al. 2013; Wood et al.
2018). We also used two strains of *Ensifer meliloti*, MAG 154 and MAG 282, sourced from a
panel of 191 strains isolated from *M. truncatula* sampled from the Mediterranean Basin (Katy
Heath, U. Illinois). As with the inbred lines of *M. truncatula,* we chose these strains because
they exhibited variable effects on plant fitness proxies like aboveground biomass (Batstone et
al. 2022)**.**
151
152   *Trait measurement*
153
154   We measured a series of resource exchange, use, and production traits on host plants
155   (Table 1). We also measured two sets of mutualism outcomes: plant fitness and nodulation
156   (Table 1). Plant fitness was estimated by weighing the total dry aboveground biomass for an
157   individual plant. We measured three nodule traits: total nodule number, total nodule mass, and
158   average nodule size (i.e., total nodule number / total nodule mass). By harvesting plants and

Table 1: The mutualism outcome and resource exchange, use, and production traits measured in this study. Brief descriptions of how each trait was measured and calculated are also provided.

Mutualism component		Trait	Measurement / Calculation
Outcome	Plant fitness	Aboveground biomass	Dry weight of aboveground plant tissue
		Aboveground biomass growth rate	Dry weight of aboveground plant tissue measured at two timepoints (see methods)
	Nodulation	Total nodule mass	Dry weight of all nodules from an individual plant's root system
		Nodule number	Count of all nodules on individual plant's root system
		Average nodule mass	Total nodule mass / nodule number
Exchange traits	Benefit	Nitrogen composition	Proportion of aboveground plant tissue that is composed of nitrogen (measured via elemental analysis)
		Total nitrogen benefit	Nitrogen composition of aboveground tissue * aboveground biomass
		Total nitrogen benefit per symbiont	Slope of the relationship between total nitrogen benefit and total nodule mass (see "statistical analysis" in methods)
	Cost	Total carbon cost	Carbon composition of nodule tissue * total nodule mass
		Nodule carbon fraction	Total carbon cost / (total carbon cost + (aboveground tissue carbon composition * aboveground biomass)
	Resource exchange ratio	Carbon-for-nitrogen exchange ratio	Total carbon cost / total nitrogen benefit
Resource Use	Nitrogen use efficiency	Carbon:nitrogen ratio in aboveground tissue	Nitrogen composition in aboveground plant tissue / carbon composition in aboveground plant tissue
Resource Production		Leaf photosynthetic rate	See Methods and Supplementary Methods

measuring aboveground biomass at two different timepoints, we were also able to infer
genotype effects on growth rate. *M. truncatula* populations are thought to be subject to viability
selection in their dry, Mediterranean habitat, thus growth rate could be an important fitness
component (Urban 1873; Lesins and Lesins 1979).
Unless otherwise noted, all resource exchange, use, and production traits were
measured using carbon and nitrogen elemental composition data collected via combustion
analysis. To perform combustion analysis, tissue was pulverized using either a mechanical ball
grinder or a mortar and pestle. Then, approximately 4 mg of pulverized tissue was weighed
using an ultra-micro balance (CP2P Microbalance, Sartorius GmbH, Goettingen, Germany),
packaged into tin capsules, and ran on a Costech ECS4010 Elemental Analyzer (Costech
Analytical Technologies, Inc. Valencia, CA, USA). The elemental analyzer returns
measurements of carbon and nitrogen composition (%) for the tissue sample analyzed. We
multiplied carbon and nitrogen composition by the total dry mass that the elemental analysis
sample came from to infer the amount of total biomass carbon and nitrogen in a given sample
(Table 1). We performed combustion analysis separately on one aboveground plant tissue
sample and one nodule tissue from between 6 and 10 individual plants from each genotype
combination.
We measured three benefit-associated exchange traits: nitrogen composition (%) in
aboveground tissue (hereafter referred to as nitrogen composition), total nitrogen in
aboveground tissue (hereafter referred to as total nitrogen benefit), and total nitrogen in
aboveground tissue as a function of total nodule mass (hereafter referred to as total nitrogen
benefit per symbiont; see statistical analysis section for more details; Table 1). We measured
two cost-associated exchange traits: total carbon in nodule tissue (hereafter referred to as total
carbon cost), and the fraction of total carbon in aboveground tissue and nodule tissue that is in
nodule tissue (hereafter referred to as nodule carbon fraction). These cost traits incorporate
both the carbon cost of constructing nodules as well as the carbon traded to the symbiont. This
approach is unable to quantify the portion of carbon transferred to rhizobia that was respired by
rhizobia, thus our analysis is making the simplifying assumption that respiration rates are
consistent across genotype combinations. We also calculated the ratio of total nitrogen in
aboveground tissue to total carbon in nodule tissue as a proxy of the carbon-for-nitrogen
exchange ratio.
We measured one resource use trait and one resource production trait. The resource
use trait was measured as the ratio of total carbon to total nitrogen in aboveground tissue as a
proxy for nitrogen use efficiency (Dovrat et al. 2020). The benefit production trait we measured
was leaf photosynthesis rate. Photosynthesis rate measurements were made using a LI-6400
Portable Photosynthesis System (LI-COR, Inc., Lincoln, NE, USA) with an 6400-02B LED Light
Source (LI-COR, Inc., Lincoln, NE, USA) attached to the leaf chamber to apply a consistent
level of light for all measurements. See the supplementary methods for details on how
photosynthesis rate was measured and calculated (Supplementary methods). Table 1 provides
short summaries of how each resource exchange, use, and production trait was quantified.
199
200   *Experimental design*
201

To quantify genotype-dependent variation (i.e., an effect of plant genotype, rhizobia
genotype, and GxG interactions) in the traits we measured we grew our two *M. truncatula*
accessions in fully factorial design with our two *E. meliloti* strains for a total of four plant-rhizobia
genotype combinations. Each of the four genotype combinations was replicated 33 times
totalling 132 experimental units.

Seeds from our two plant genotypes were scarified, sterilized, imbibed, and stratified
using the methods reported in the "Seed storage and germination" chapter of the Medicago
Handbook (Noble Institute; Supplementary methods). Prior to planting, we filled 66 mL Cone-
tainer pots (Stuewe & Sons, Inc., Tangent, OR, USA) with a growing media consisting of a 1:4
mix of perlite and sand. We sterilized the filled Cone-tainer by autoclaving them twice. One
seedling was placed in each Cone-tainer. One day after planting, seedlings were given 1 mL of
0.625 mM nitrogen Fahraeus fertilizer (Batstone et al. 2017). Afterwards, plants were given 0.0
mM N Fahraeus fertilizer every 2 to 3 days to ensure that rhizobia were the only source of
nitrogen (Batstone et al. 2017).

Cone-tainer pots were placed in 98-well racks. To avoid the risk of cross contamination
of rhizobia strains, every plant in a rack received the same rhizobia strain. Each rack was
enclosed in a semi-permeable tent consisting of plastic wrap and PVC pipe to further reduce the
risk of cross contamination. Plants were grown in two growth chambers in a complete block
design. The height of light fixtures in both chambers was adjusted so that a photosynthetic
active radiation (PAR) reading of ~250 $\mu mol/m^{-2}/s^{-1}$ was consistently measured on the floor of
the growth chambers. Chambers were set to 22 °C and a 16:8 light:dark photoperiod. Seedings
were maintained in a high humidity environment for about one week, after which they were
exposed to ambient humidity, usually 40-50%.

Two weeks after planting, we inoculated plants with rhizobia. Plants received either
strain MAG 154 or MAG 282. We grew strains from glycerol stocks in sterile tryptone yeast
liquid media in a shaking incubator at 30 °C and 180 rpm. We let the rhizobia cultures grow for
approximately two days. We then streaked the media on sterile Petri plates containing tryptone
yeast agar under a laminar flow hood. We incubated the plates at 30 °C until colonies started to
form. Next, we selected an individual colony from each strain to grow in tryptone yeast liquid
media. Again, rhizobia cultures were allowed to grow for approximately two days. We diluted the
liquid media to an OD_{600} of 0.1 and inoculated each plant with 1 ml of the diluted culture.

Plants were harvested in three separate cohorts at 9, 10, and 11 weeks post planting.
Approximately 20 plants from each treatment group were harvested at 11 weeks. Of those
plants harvested after 11 weeks, approximately 10 from each treatment were used to measure
the benefit and cost associated traits described above via elemental analysis. Approximately 3
plants from each treatment group were harvested at 10 weeks to measure photosynthesis. Prior
to harvest at the 10 week timepoint, photosynthesis rate measurements were taken on 2-3
leaves of each plant. Approximately 10 plants from each treatment group were harvested at 9
weeks. Plants in the 9-week cohort were compared to those in the 11-week cohort to estimate
growth rate (see statistical analysis section for more details).

At each harvest, we gently removed plants from their Cone-tainer pots and cut the
shoots from the roots. The shoots were placed in brown paper bags and dried at 60 °C for at
least a week before being weighed with an analytical scale to measure aboveground biomass.
Plants used for benefit and cost trait measurement via elemental analysis were processed as

described in the "Trait measurement" section. The roots were washed before being placed in
plastic bags and stored in a freezer at -20 °C.
To collect nodulation data, frozen roots were thawed and nodules were removed under a
dissecting microscope. Nodules were counted as they were removed to quantify nodule
number. To assess the consistency of nodule count estimates across researchers, we had each
research count nodules on the same subset of 10 plants. Nodule counts were highly repeatable
across researchers in a linear mixed effects model with nodule number as the response, and
plant and researcher as random effects (researcher explained only 0.4% of variation in nodule
counts). Removed nodules were placed in coin envelopes, dried at 60 °C for at least a week,
and weighed. Nodules from plants that were used for benefit and cost trait measurement via
elemental analysis were processed as described in the "Trait measurement" section.
257
258   *Statistical analysis*
259
We used linear models to test for an effect of plant genotype, rhizobia genotype, and
plant genotype by rhizobia genotype (GxG) interactions on the traits we measured. All models
used a normal error distribution except for the model with nodule number as the response
variable, which used a negative binomial error distribution. Block was included as a fixed effect
in all models. Continuous predictors were mean-centered to avoid spurious inference on main
effects in the presence of interactions (Schielzeth 2010). For some models, response variables
were log-transformed to better conform with the assumption of normality. If a response variable
was log transformed, it will be indicated in the figure caption of the associated figure. Models
with nodule traits as the response included total root mass as a covariate to account for
differences in nodule traits associated with differences in root system size.
To test for genotype effects on growth rate in aboveground biomass, we included the
following interaction terms in the model, plant genotype x timepoint, rhizobia genotype x
timepoint, and plant genotype x rhizobia genotype x timepoint.
For analysis of elemental composition measurements (e.g. nitrogen composition),
arcsine square root transformations were performed to normalize the data prior to running the
models.
To test for significant genotype effects on total nitrogen benefit per symbiont, we
included the interaction terms plant genotype x total nodule mass, rhizobia genotype x total
nodule mass, and plant genotype x rhizobia genotype x total nodule mass terms in our models.
Each of these interaction terms estimates differences in the slope of the relationship between
total nitrogen in aboveground tissue and nodule mass that are due to genotype effects.
Our model with photosynthesis rate as the response included each individual leaf
measurement as an independent replicate. To account for any variation due to measurements
made on leaves from the same plant, we included plant ID as a random effect.
The significance of model terms in each model was tested using type III sum of squares.
For linear models, significant testing was performed using *F*-statistics, and for generalized linear
models, significance testing was performed with Wald χ^2 tests.
All statistics and data visualization were performed in R v4.2.3 (R Core Team 2023). For
simple linear models, we used the base function lm(); for linear mixed effects models we used
the lmer() function in the lme4 package (Bates et al. 2014), and for generalized linear mixed

effects models we used the glmmtmb() function in the glmmTMB package (Brooks et al. 2017).
Model assumptions were evaluated using the DHARMa package (Hartig and Lohse 2022).
Significant testing was performed with the Anova() function in the car package (Fox and
Weisberg 2018). All plots include estimated marginal means that were calculated using the
ggeffects package (Lüdecke 2018). The ggplot2 (Wickham 2016), ghibli (Henderson et al.
2022), and ggpubr (Kassambara 2023) packages were used to make all of the plots presented
here. All data and code used in this study can be found here [Dryad repository will be added
if/when manuscript is accepted for publication]
298
299   **Results**
300
301   *Genetic variation in fitness and mutualism outcomes*
302
We found a significant plant genotype x rhizobia genotype (GxG) interaction effect on
aboveground biomass, our proxy for plant fitness, but no main effects of either plant or rhizobia
genotype (Fig. 1, Table 2). We also found significant plant genotype, rhizobia genotype, and
GxG effects on aboveground biomass growth rate (Fig. 1, Table 2). During the analysis of
growth rate, we identified an outlier in the HM028 - MAG 282 - Timepoint 1 treatment group.
Removal of this outlier does not change the statistical significance of any model terms.
We observed a significant rhizobia genotype effect on total nodule mass, and significant
plant genotype effects and rhizobia genotype effects on total nodule number and average
nodule size (Fig. 1, Table 2).
312
313   *Genetic variation in exchange traits*
314
We measured three benefit-associated exchange traits, nitrogen composition, total
nitrogen benefit, and total nitrogen benefit per symbiont (Table 1). We observed a significant
rhizobia genotype effect and GxG effect for nitrogen composition in aboveground tissue (Fig. 2,
Table 3), but no significant genotype effects were observed for total nitrogen biomass in
aboveground tissue (Fig. 2, Table 3). For total nitrogen per symbiont, we observed significant
plant genotype x total nodule mass, rhizobia genotype x total nodule mass, and G x G x total
nodule mass mass effects, indicating that our measure of benefit per symbiont varies in
response to plant genotype, rhizobia genotype, and the interaction of plant genotype and
rhizobia genotype (Fig. 2, Table 3).
We measured two cost-associated exchange traits, total carbon cost, and the nodule
carbon fraction (Table 1). We observed a significant rhizobia genotype and GxG effect for total
carbon cost, but no significant genotype effects for nodule carbon fraction (Fig. 3, Table 3).
We also quantified the nitrogen-for-carbon exchange ratio by dividing total nitrogen
benefit by the total carbon cost (Table 1). We observed a significant rhizobia genotype effect for
the nitrogen-for-carbon exchange (Fig. 4, table 3).
330

Figure 1: Genetic variation for (A) log transformed aboveground biomass, our proxy for plant fitness, (B) log transformed aboveground biomass growth rate, (C) total nodule mass, (D) nodule number, and (E) average nodule size. In all panels, transparent points are the raw data and opaque points are the estimated marginal means. Error bars are the 95% confidence interval. Significant genotype effects are listed on the figure: G_P = plant genotype effect, G_R = rhizobia genotype effect, $G_P \times G_R$ = plant genotype x rhizobia genotype interaction, $G_P \times T$ = plant genotype x timepoint interaction, $G_R \times T$ = rhizobia genotype x timepoint interaction, and $G_P \times G_R \times T$ = plant genotype x rhizobia genotype x timepoint interaction. Model term statistical significance did not change when the outlier in the HM028-MAG282-Timepoint 1 treatment group in panel B was removed.

Genetic variation in resource use and production traits

We measured variation in two traits associated with resource use and production, the carbon:nitrogen ratio in aboveground tissue (a proxy for nitrogen use efficiency) and leaf photosynthetic rate (Table 1). The carbon:nitrogen ratio in aboveground tissue was used to infer how efficiently host plants are able to use nitrogen received through trade to incorporate carbon in aboveground tissue (Dovrat et al. 2020). Leaf photosynthetic rate to infer the efficiency of the production of carbon, some of which is ultimately traded away. We found a significant effect of

Table 2: Genotype effects on mutualism outcome traits. Significant model terms are highlighted in gray.

	Outcomes									
	Plant Fitness				Nodulation					
	Above-ground biomass		Above-ground biomass growth rate		Total nodule mass		Nodule number		Average nodule mass	
	$F_{1,81}$	P	$F_{1,117}$	P	$F_{1,50}$	P	$\chi^2_{df=1}$	P	$F_{1,50}$	P
Rhizobia genotype (G_R)	1.09	0.3	**15.0**	**0.0002**	**7.17**	**0.004**	**16.74**	**<0.0001**	**5.41**	**0.024**
Plant genotype (G_P)	0.18	0.7	**20.76**	**<0.0001**	0.0004	0.98	**3.97**	**0.046**	**5.62**	**0.02**
G_R x G_P	**4.25**	**0.04**	**30.21**	**<0.0001**	3.96	0.05	3.25	0.071	0.002	0.96
Timepoint (T)	-	-	**64.9**	**<0.0001**	-	-	-	-	-	-
T x G_R	-	-	**13.28**	**0.0005**	-	-	-	-	-	-
T x G_P	-	-	**12.96**	**0.0004**	-	-	-	-	-	-
T x G_R x G_P	-	-	**29.81**	**0.0001**	-	-	-	-	-	-

339 rhizobia genotype and the interaction between rhizobia genotype and plant genotype on **Table 1:**
340 Genotype effects on mutualism outcome traits. Significant model terms are highlighted in gray.
341
342 carbon:nitrogen ratio (Fig. 5, Table 3). We observed a significant plant genotype x rhizobia
343 genotype interaction on photosynthesis rate (Fig. 5, Table 3).
344
345 **Discussion**
346
In this study, we found that variation in mutualism resource exchange, use, and
production traits are genotype-dependent. Consistent with our expectations, we observed
rhizobia genotype and GxG effects on both benefit-associated and cost-associated exchange
traits. However, we only observed differences between rhizobia genotypes for the nitrogen-for-
carbon exchange ratio. Contrary to our expectation, rhizobia genotype affected plant resource
use and production traits as much or more than plant genotype. Our results suggest that
rhizobia can contribute to the evolution of resource use and production traits in their hosts.
Furthermore, our work indicates that the evolution of mutualisms may not be limited to resource
exchange traits. Rather, how efficiently mutualistic partners can integrate the benefits of trade
(i.e. resource use efficiency) and/or how efficiently they can produce the resource traded away
may be just as important for determining the evolution of mutualisms. The ubiquity of concurrent

Table 3: Genotype effects on resource exchange, use, and production traits. Significant model terms are highlighted in gray.

	Exchange traits												Resource use		Resource production	
	Benefits						Costs				Benefit: cost		Nitrogen use efficiency		Leaf photosynthesis rate	
	Nitrogen composition		Total nitrogen benefit		Total nitrogen benefit per symbiont		Total carbon cost		Nodule carbon fraction		Nitrogen-for-carbon exchange ratio					
	$F_{1,36}$	P	$F_{1,36}$	P	$F_{1,29}$	P	$F_{1,26}$	P	$F_{1,35}$	P	$F_{1,30}$	P	$F_{1,36}$	P	$\chi^2_{df=1}$	P
Rhizobia genotype (G_R)	**11.48**	**0.002**	0.29	0.6	**7.54**	**0.01**	**5.04**	**0.03**	1.53	0.223	**5.6**	**0.02**	**6.72**	**0.01**	1.73	0.19
Plant genotype (G_P)	1.12	0.3	0.03	0.85	0.35	0.56	0.006	0.94	1.83	0.19	0.005	0.95	1.46	0.23	0.74	0.39
$G_R \times G_P$	**6.3**	**0.02**	0.31	0.58	2.58	0.12	**4.19**	**0.049**	0.0003	0.99	1.88	0.18	**5.78**	**0.02**	**4.62**	**0.03**
Centered total nodule mass (tnm)	-	-	-	-	1.64	0.21	-	-	-	-	-	-	-	-	-	-
Tnm x G_R	-	-	-	-	**9.68**	**0.004**	-	-	-	-	-	-	-	-	-	-
Tnm x G_P	-	-	-	-	**4.63**	**0.04**	-	-	-	-	-	-	-	-	-	-
Tnm x G_P x G_R	-	-	-	-	**8.69**	**0.006**	-	-	-	-	-	-	-	-	-	-

Figure 2: Genetic variation for three benefit-associated exchange traits: (A) nitrogen composition (%), (B) total nitrogen benefit (g), and (C) total nitrogen benefit per symbiont. In panels A and B, transparent points are the raw data, opaque points are the estimated marginal means, and Error bars are 95% confidence intervals. For panel C, transparent points are the raw data, lines represent the estimated marginal slopes derived from the model, and ribbons are the 95% confidence interval for the slope. Significant genotype effects are listed on the figure: G_P = planet genotype effect, G_R = rhizobia genotype effect, $G_P \times G_R$ = plant genotype x rhizobia genotype interaction, $G_P \times TNM$, $G_R \times TNM$, $G_P \times G_R \times TNM$.

358   plant genotype and rhizobia genotype effects as well as GxG effects that we observed on
359   resource exchange, use, and production traits complements previous research that has found
360   similar effects on plant fitness components and nodulation traits. Together, these results further
361   emphasize an important role for jointly determined traits in mediating mutualism evolution.
362
363   *Genotype effects on resource exchange traits*
364
We predicted that GxG effects, which occur when trait variation across genotypes in one
interacting partner is dependent on the genotype of the other partner, would be common to
exchange traits, because the exchange of resources involves cooperation between the plant
and rhizobia (Sachs et al. 2004). Consistent with this prediction, we observed GxG effects for
two benefit-associated traits: nitrogen composition and total nitrogen benefit per symbiont. For
cost-associated traits we found GxG effects for total carbon cost. However, we did not observe
a GxG effect for the carbon-for-nitrogen exchange ratio. Similarly, among our cost-associated
traits, we observed a GxG effect for total carbon cost. Total carbon cost is a metric of
cumulative cost because it is a representation of the gross amount of carbon that hosts provide
to their rhizobia (and to nodule formation).
Our prediction that GxG effects should be common to exchange traits is based on the
hypothesis that the outcomes of trade in resource mutualisms are governed by bargaining
between partners (Sachs et al. 2004; Akçay and Roughgarden 2007). There are numerous
proposed mechanisms by which interacting partners may strengthen their bargaining positions.
For example, legumes are able to "sanction" poor quality partners resulting in beneficial trading
outcomes (Kiers et al. 2003; Westhoek et al. 2021). Thus, the GxG effects on exchange traits

Figure 3: Genetic variation for two cost-associated exchange traits: (A) total carbon cost and (B) the nodule carbon fraction. Transparent points are the raw data and opaque points are the estimated marginal means. Error bars are the 95% confidence interval. Significant genotype effects are listed on the figure: G_P = plant genotype effect, G_R = rhizobia genotype effect, and G_P X G_R = plant genotype X rhizobia genotype interaction.

that we observed here could be mediated by the numerous bargaining strategies that hosts and
microbes employ. Alternatively, GxG effects on exchange traits may also emerge via a
mismatch between partners (Barrett et al. 2012; Wendlandt et al. 2019). Partner mismatches
occur when some rhizobia are more effective partners for some plant genotypes compared to
others. Partner mismatches can emerge from a variety of evolutionary mechanisms such as
rhizobia adaptation to different hosts (Batstone et al. 2020), coevolution between spatially
separated populations of legumes and rhizobia (Gomulkiewicz et al. 2003), and/or isolation by
distance.
To our surprise, however, the carbon-for-nitrogen exchange rate was only affected by
rhizobia genotype, not by both partners. This result suggests that rhizobia may exert more
control over the overall outcome of trade than their host. Rhizobia control conflicts with
theoretical and empirical work that suggests that, in general, host plants exert more power in
determining the outcomes of trade (Daubech et al. 2017; Clark et al. 2019). Of course, a primary
reason that host plants are expected to exert more control over the trade is their ability to
"bargain" with multiple different microbial strains simultaneously (Akçay and Simms 2011;
Westhoek et al. 2021); a scenario that was unable to occur in this study due to our single strain
inoculation design.
Finally, there were a few exchange traits for which we observed no genotype effects.
Among benefit-associated traits, total nitrogen benefit, our metric of absolute benefit, did not
differ between plant or rhizobia genotypes. The same was true for nodule carbon fraction, our
metric of relative cost. The fact that we observed genotype differences in the relative but not
absolute benefit—and the absolute but not relative cost—suggests that absolute and relative
costs (and benefits) of mutualism may exhibit different evolutionary dynamics.

Figure 4: Genetic variation for the nitrogen-for-carbon exchange ratio. **Transparent points are the raw data and** opaque points are the estimated marginal means. Error bars are the 95% confidence interval. Significant genotype effects are listed on the figure: G$_P$ = planet genotype effect, G$_R$ = rhizobia genotype effect, and G$_P$ X G$_R$ = plant genotype X rhizobia genotype interaction.

Genotype effects on resource use and production traits

We predicted that plant genotype would be the major contributor to differences in plant resource use and production traits. Contrary to this prediction, we observed GxG effects for both of the resource use and production traits that we measured: nitrogen use efficiency (our metric of resource use) and leaf photosynthetic rate (our metric of resource production).

We predicted that host resource use and production traits would respond primarily to plant genotype because these traits are indirectly related to resource exchange and represent components of plant physiology. However, rhizobia effects on these host traits are perhaps not surprising if hosts change how they allocate benefits or invest in the production of resources in response to trade. Thus, rhizobia and GxG effects on resource use and production traits may arise from the GxG effects for exchange traits that we observed.

Resource use traits, like nitrogen use efficiency, have previously been shown to be microbially mediated (Dovrat et al. 2020; Zhang et al. 2020). For instance, legumes modify their nitrogen use efficiency in response to both soil nitrogen availability and the presence of rhizobia (Dovrat et al. 2020). Our study raises the important question of why nitrogen use efficiency could vary across combinations of plant and rhizobia genotypes. A GxG effect on nitrogen use efficiency could be due to differences in the efficiency of symbiotic nitrogen fixation across plant

Figure 5: Genetic variation for (A) carbon : nitrogen ratio in aboveground tissue and (B) leaf photosynthetic rate. Transparent points are the raw data and opaque points are the estimated marginal means. Error bars are the 95% confidence interval. Significant genotype effects are listed on the figure: G_P = planet genotype effect, G_R = rhizobia genotype effect, and G_P X G_R = plant genotype X rhizobia genotype interaction.

and rhizobia combinations. Legume species vary in their strategies of regulation of nitrogen fixation. Some, like *M. truncatula,* are obligate fixers and will attempt to maintain a constant nitrogen fixation effort regardless of the activity of their rhizobia partners or external supplies of nitrogen, while others are facultative fixers and can down- or upregulate nitrogen fixation in response to environmental conditions (Hedin et al. 2009). Consequently, variation in nitrogen use efficiency in obligate fixers like *M. truncatula* could simply be a function of the variation in the efficiency of nitrogen fixation across plant and rhizobia genotype combinations. Alternatively, it is also plausible that legumes may have evolved strategies to adjust how they use and allocate the benefits from trade in order to buffer against the consequences of interacting with poor quality partners.

Resource production traits, like leaf photosynthetic rate, also have been shown to respond to microbes (Adams et al. 2016). Legumes in particular exhibit significantly higher photosynthetic rates in the presence versus the absence of rhizobia (Clark et al. 2019; Dovrat et al. 2020). Photosynthesis is fundamental to the exchange of resources between legumes and rhizobia because it provides the carbon necessary to support rhizobia and it also requires a large amount of nitrogen, which is provided by the rhizobia. A heightened photosynthetic rate may equate to a larger pool of carbon that can be devoted to the mutualism, and by extension, a large supply of nitrogen that can be returned to the host plant. Variation in photosynthetic rate may also simply be an indication of nitrogen limitation and the outcomes of trade. Photosynthesis involves nitrogen-rich machinery and therefore, genotype combinations with the

443   highest photosynthetic rates may simply be the genotype combinations in which the largest
444   amount of nitrogen is being transferred to the host plant. Regardless of the mechanisms
445   responsible for GxG effects on photosynthetic rate, the fact that a portion of the genetic variation
446   for leaf photosynthetic rate resides in rhizobia emphasizes the far-reaching effect that this
447   mutualism has on host plant physiology.
448   The findings of this study underscore the possibility that the evolution of mutualisms may
involve traits related to how interacting partners can integrate the benefits of trade (i.e. resource
use efficiency) and how efficiently they can produce the resources traded away. This suggestion
is corroborated by a recent study involving a synthetic mutualism between engineered strains of
brewer's yeast (Vidal and Segraves 2021). The authors observed strong fluctuations in both
resource use efficiency and resource production of both strains as they evolved in response to
the availability of the limiting resources produced by their partners. Resource use and
production traits may represent key components of mutualisms that allow interacting partners
the ability to adjust their physiology in response to the outcomes of trade. Future studies should
carefully consider selection on all mutualism traits, not just those related to the exchange of
resources, as well as covariation among resource exchange, use and production traits.
459
460   *The importance of jointly determined traits to predicting evolution in mutualism*
461
462   The ubiquity of rhizobia and GxG effects on mutualism traits has three important
implications for the evolution of mutualisms. First, this finding indicates that traits involved in
resource exchange, use, and production may be jointly determined by the genomes of both
interacting partners, just as is the case with fitness outcomes (Heath 2010). Jointly determined
traits, here defined as traits that exhibit variation in response to more than one genome as either
two independent main effects or a GxG effect, can have different evolutionary trajectories than
those determined by the genome of just one or the other interacting partner (O'Brien et al.
2021b). The evolution of jointly determined traits is a function of the optimal trait values for the
different interacting species, the strengths of selection, and fitness feedbacks (O'Brien et al.
2021b). Thus, a trait-based framework, wherein quantitative genetic approaches can be
leveraged to identify the sources of variation in mutualism traits, is necessary to develop a
better understanding of mutualism evolution.
474   Second, these results also show that microbial mutualists play an important role in
determining the expression of host traits like resource use and production. Microbes are known
to mediate host nutrient acquisition (Hestrin et al. 2019; Harbort et al. 2020), life history
transitions (Wagner et al. 2014; O'Brien et al. 2021a), defense against antagonists (Johnson et
al. 1987; Berg and Koskella 2018), stress tolerance (Lau and Lennon 2012), and morphology
(Ortíz-Castro et al. 2008; Batstone et al. 2017). The great majority of the research on microbial
mediation of plant traits involves observation and manipulative studies of either individual
species of microbes or whole microbiomes. Such a framework allows for questions about how
variation in community properties (e.g. relative abundance and composition of specific microbial
species) affects plant functional traits. Our work builds on this knowledge by demonstrating that
intraspecific variation, in addition to interspecific variation, can also influence the expression of
plant functional traits. This finding complements two recent studies, O'Brien et al. (2021b) and
De Lisle et al. (2022), which have highlighted the potential evolutionary significance of the effect

of intraspecific partner variation in the expression of mutualism traits. An important take away
from those studies and ours is that evolution of rhizobia populations can drive changes in host
plant traits. Furthermore, intraspecific microbial variation is invisible to the 16S sequencing that
is commonly applied to characterize community composition in microbiome studies because
strains belonging to the same species will have identical 16S sequences. Thus, substantial
cryptic variation in microbe mediated plant traits measured in microbiome manipulations may go
unnoticed.
Third, variation in resource trade, use, and production traits across different
combinations of plant and rhizobia genotypes highlights the potential importance of the non-
random association of partner genotypes for mutualism evolution. If a mutualism trait responds
to a GxG effect and is a target of selection, then the plant genotypes favored by selection will be
dependent on which rhizobia genotype they are paired with. If associations between specific
plant and rhizobia genotypes are random, then the response to selection will be smaller than
expected because most genetic variation is non-additive and a function of which genotypes are
paired together. On the other hand, if associations between specific plant and rhizobia
genotypes are non-random, then the response to selection could be either larger (if associations
between plant and rhizobia genotypes inflate genetic variation for the trait) or smaller (if
associations between plant and rhizobia genotypes constrain genetic variation for the trait) than
expected. While host plants have a clear influence on their rhizosphere microbes (Miranda-
Sánchez et al. 2016; Greenlon et al. 2019) and can vary in their ability to select beneficial
microbial genotypes (Burghardt et al. 2018, 2019), the frequency of non-random associations
between specific plant and rhizobia genotypes in nature has been rarely measured. Given the
apparent importance of rhizobia genotypes in influencing variation in mutualism traits, more
studies are needed to address the degrees of non-random associations between interacting
genotypes in order to develop a predictive framework for understanding mutualism evolution.
512
513   *Conclusions and future directions*
514
Our study has identified a number of mutualism traits that could underlie variation in
fitness outcomes in mutualisms and are jointly determined by the genotypes of both interacting
partners. More studies are now needed to quantify (1) the relationship between mutualism traits
measured here and the fitness of both interacting partners, (2) the variation and covariation
among these traits, and (3) the non-random association among the genotypes of interacting
species. Each of these objectives offers different, but important, extensions of the work
presented here to advance our understanding of mutualism evolution. Measuring the
relationship between mutualism traits and fitness (i.e. selection) is necessary to both identify the
traits that could underlie fitness variation and to identify the evolutionary forces that maintain
fitness variation. Quantifying variation and covariation among mutualism traits is necessary to
evaluate the physiological interdependence of mutualism traits. For example, if traits like
resource use help host plants buffer the outcomes of trade, then we might expect to see high
covariation among exchange and resource use traits. Finally, the broad prevalence of rhizobia
and GxG effects highlights the potential importance of non-random associations among
genotypes in mutualism evolution.
530

Acknowledgements

We thank Chang-Yu Chang, Addison Martin, and Eunnuri Yi for helpful comments on this manuscript. We also thank Chigozie Ibe, Anisa Robinson, Linda Wu, and Seokyoon Chang for help with executing the experiment and collecting data. *Medicago truncatula* genotypes were generously provided by Nevin Young and the Medicago HapMap project. *Ensifer meliloti* strains were generously provided by Liana Burghardt and Katy Heath. Growth chamber logistical support was provided by the University of Pennsylvania's Greenhouse support staff members Kathyrn Butler and Samara Gray. We are grateful to David Vann and the University of Pennsylvania's Earth and Environmental Science Department's Instrumentation Core for assistance with elemental analysis. This work was funded by a Society for the Study of Evolution R. C. Lewontin Award to McCall Calvert and NSF-DEB 2118397 to Corlett Wood

References

Adams, M. A., T. L. Turnbull, J. I. Sprent, and N. Buchmann. 2016. Legumes are different: Leaf nitrogen, photosynthesis, and water use efficiency. Proc. Natl. Acad. Sci. 113:4098–4103. Proceedings of the National Academy of Sciences.

Akçay, E., and J. Roughgarden. 2007. Negotiation of mutualism: rhizobia and legumes. Proc. R. Soc. B Biol. Sci. 274:25–32.

Akçay, E., and E. L. Simms. 2011. Negotiation, sanctions, and context dependency in the legume-rhizobium mutualism. Am. Nat. 178:1–14.

Albrecht, M., B. Schmid, Y. Hautier, and C. B. Müller. 2012. Diverse pollinator communities enhance plant reproductive success. Proc. R. Soc. B Biol. Sci. 279:4845–4852.

Andrews, M., and M. E. Andrews. 2017. Specificity in legume-rhizobia symbioses. Int. J. Mol. Sci. 18:705.

Barrett, L. G., L. M. Broadhurst, and P. H. Thrall. 2012. Geographic adaptation in plant–soil mutualisms: tests using Acacia spp. and rhizobial bacteria. Funct. Ecol. 26:457–468.

Bates, D., M. Mächler, B. Bolker, and S. Walker. 2014. Fitting linear mixed-effects models using lme4. arXiv.

Batstone, R. T., E. M. Dutton, D. Wang, M. Yang, and M. E. Frederickson. 2017. The evolution of symbiont preference traits in the model legume *Medicago truncatula*. New Phytol. 213:1850–1861.

Batstone, R. T., H. Lindgren, C. M. Allsup, L. A. Goralka, A. B. Riley, M. A. Grillo, A. Marshall-Colon, and K. D. Heath. 2022. Genome-wide association studies across environmental and genetic contexts reveal complex genetic architecture of symbiotic extended phenotypes. BioRxiv 2021–08. Cold Spring Harbor Laboratory.

Batstone, R. T., A. M. O'Brien, T. L. Harrison, and M. E. Frederickson. 2020. Experimental evolution makes microbes more cooperative with their local host genotype. Science 370:476–478. American Association for the Advancement of Science.

Berg, M., and B. Koskella. 2018. Nutrient- and dose-dependent microbiome-mediated protection against a plant pathogen. Curr. Biol. 28:2487-2492.e3.

Bronstein, J. L. 1994. Conditional outcomes in mutualistic interactions. Trends Ecol. Evol. 9:214–217.

Bronstein, J. L. 2001. The Costs of mutualism. Am. Zool. 41:825–839.

Bronstein, J. L. 2015. The study of mutualism. P. 0 *in* J. L. Bronstein, ed. Mutualism. Oxford University Press.

Brooks, M. E., K. Kristensen, K. J. van Benthem, A. Magnusson, C. W. Berg, A. Nielsen, H. J.
Skaug, M. Machler, and B. M. Bolker. 2017. glmmTMB balances speed and flexibility
among packages for zero-inflated generalized linear mixed modeling. R J. 9:378–400.
Technische Universitaet Wien.
Burghardt, L. T., B. Epstein, J. Guhlin, M. S. Nelson, M. R. Taylor, N. D. Young, M. J.
Sadowsky, and P. Tiffin. 2018. Select and resequence reveals relative fitness of bacteria
in symbiotic and free-living environments. Proc. Natl. Acad. Sci. 115:2425–2430.
Burghardt, L. T., B. Epstein, and P. Tiffin. 2019. Legacy of prior host and soil selection on
rhizobial fitness in planta. Evolution 73:2013–2023.
Burghardt, L. T., J. Guhlin, C. L. Chun, J. Liu, M. J. Sadowsky, R. M. Stupar, N. D. Young, and
P. Tiffin. 2017. Transcriptomic basis of genome by genome variation in a legume-
rhizobia mutualism. Mol. Ecol. 26:6122–6135.
Clark, T. J., C. A. Friel, E. Grman, M. L. Friesen, and Y. Shachar-Hill. 2019. Unfair trade
underground revealed by integrating data with Nash bargaining models. New Phytol.
222:1325–1337.
Daubech, B., P. Remigi, G. Doin de Moura, M. Marchetti, C. Pouzet, M.-C. Auriac, C. S.
Gokhale, C. Masson-Boivin, and D. Capela. 2017. Spatio-temporal control of mutualism
in legumes helps spread symbiotic nitrogen fixation. eLife 6:e28683. eLife Sciences
Publications, Ltd.
De Lisle, S. P., D. I. Bolnick, E. D. Brodie III, A. J. Moore, and J. W. McGlothlin. 2022.
Interacting phenotypes and the coevolutionary process: Interspecific indirect genetic
effects alter coevolutionary dynamics. Evolution 76:429–444.
Denison, R. F. 2000. Legume Sanctions and the Evolution of Symbiotic Cooperation
by Rhizobia. Am. Nat. 156:567–576. The University of Chicago Press.
Dovrat, G., H. Bakhshian, T. Masci, and E. Sheffer. 2020. The nitrogen economic spectrum of
legume stoichiometry and fixation strategy. New Phytol. 227:365–375.
Fox, J., and S. Weisberg. 2018. An R companion to applied regression. 3rd ed. Sage
Publications, Inc., Washington DC.
Friesen, M. L., S. S. Porter, S. C. Stark, E. J. von Wettberg, J. L. Sachs, and E. Martinez-
Romero. 2011. Microbially mediated plant functional traits. Annu. Rev. Ecol. Evol. Syst.
42:23–46.
Gomulkiewicz, R., S. L. Nuismer, and J. N. Thompson. 2003. Coevolution in variable
mutualisms. Am. Nat. 162:S80–S93. The University of Chicago Press.
Greenlon, A., P. L. Chang, Z. M. Damtew, A. Muleta, N. Carrasquilla-Garcia, D. Kim, H. P.
Nguyen, V. Suryawanshi, C. P. Krieg, S. K. Yadav, J. S. Patel, A. Mukherjee, S. Udupa,
I. Benjelloun, I. Thami-Alami, M. Yasin, B. Patil, S. Singh, B. K. Sarma, E. J. B. von
Wettberg, A. Kahraman, B. Bukun, F. Assefa, K. Tesfaye, A. Fikre, and D. R. Cook.
2019. Global-level population genomics reveals differential effects of geography and
phylogeny on horizontal gene transfer in soil bacteria. Proc. Natl. Acad. Sci. 116:15200–
15209. Proceedings of the National Academy of Sciences.
Grman, E., T. M. P. Robinson, and C. A. Klausmeier. 2012. Ecological specialization and trade
affect the outcome of negotiations in mutualism. Am. Nat. 179:567–581.
Harbort, C. J., M. Hashimoto, H. Inoue, Y. Niu, R. Guan, A. D. Rombolà, S. Kopriva, M. J. E. E.
E. Voges, E. S. Sattely, R. Garrido-Oter, and P. Schulze-Lefert. 2020. Root-secreted
coumarins and the microbiota interact to improve iron nutrition in *Arabidopsis*. Cell Host
Microbe 28:825-837.e6.
Hartig, F., and L. Lohse. 2022. DHARMa: Residual Diagnostics for Hierarchical (Multi-Level /
Mixed) Regression Models.
Heath, K. D. 2010. Intergenomic epistasis and coevolutionary constraint in plants and rhizobia.
Evolution 64:1446–1458.
Heath, K. D., and J. R. Stinchcombe. 2014. Explaining mutualism variation: A new evolutionary

paradox? Evolution 68:309–317.

Heath, K. D., and P. Tiffin. 2007. Context dependence in the coevolution of plant and rhizobial
mutualists. Proc. R. Soc. Lond. B Biol. Sci. 274:1905–1912.

Hedin, L. O., E. N. J. Brookshire, D. N. L. Menge, and A. R. Barron. 2009. The nitrogen paradox
in tropical forest ecosystems. Annu. Rev. Ecol. Evol. Syst. 40:613–635.

Henderson, E., D. Desrosiers, and M. Chirico. 2022. ghibli: Studio Ghibli colour palettes.

Hestrin, R., E. C. Hammer, C. W. Mueller, and J. Lehmann. 2019. Synergies between
mycorrhizal fungi and soil microbial communities increase plant nitrogen acquisition.
Commun. Biol. 2:1–9. Nature Publishing Group.

Janzen, D. H. 1966. Coevolution of mutualism between ants and acacias in central america.
Evolution 20:249–275.

Johnson, N. D., B. Liu, and B. L. Bentley. 1987. The effects of nitrogen fixation, soil nitrate, and
defoliation on the growth, alkaloids, and nitrogen levels of *Lupinus succulentus*
(Fabaceae). Oecologia 74:425–431.

Kassambara, A. 2023. ggpubr: "ggplot2" Based Publication Ready Plots.

Kiers, E. T., M. Duhamel, Y. Beesetty, J. A. Mensah, O. Franken, E. Verbruggen, C. R.
Fellbaum, G. A. Kowalchuk, M. M. Hart, A. Bago, T. M. Palmer, S. A. West, P.
Vandenkoornhuyse, J. Jansa, and H. Bücking. 2011. Reciprocal rewards stabilize
cooperation in the mycorrhizal symbiosis. Science 333:880–882. American Association
for the Advancement of Science.

Kiers, E. T., R. A. Rousseau, S. A. West, and R. F. Denison. 2003. Host sanctions and the
legume–rhizobium mutualism. Nature 425:78–81.

Lau, J. A., and J. T. Lennon. 2012. Rapid responses of soil microorganisms improve plant
fitness in novel environments. Proc. Natl. Acad. Sci. 109:14058–14062. Proceedings of
the National Academy of Sciences.

Lesins, K. A., and I. Lesins. 1979. Genus Medicago (Leguminosae): a taxogenetic study. Dr. W.
Junk Publishers, The Hague.

Levy, S., A. Elek, X. Grau-Bové, S. Menéndez-Bravo, M. Iglesias, A. Tanay, T. Mass, and A.
Sebé-Pedrós. 2021. A stony coral cell atlas illuminates the molecular and cellular basis
of coral symbiosis, calcification, and immunity. Cell 184:2973-2987.e18.

Lüdecke, D. 2018. ggeffects: Tidy data frames of marginal effects from regression models. J.
Open Source Softw. 3:772.

Miranda-Sánchez, F., J. Rivera, and P. Vinuesa. 2016. Diversity patterns of Rhizobiaceae
communities inhabiting soils, root surfaces and nodules reveal a strong selection of
rhizobial partners by legumes. Environ. Microbiol. 18:2375–2391.

Moore, A. J., E. D. III Brodie, and J. B. Wolf. 1997. Interacting phenotypes and the evolutionary
process: I. Direct and indirect genetic effects of social interactions. Evolution 51:1352–
1362.

O'Brien, A. M., N. A. Ginnan, M. Rebolleda-Gómez, and M. R. Wagner. 2021a. Microbial effects
on plant phenology and fitness. Am. J. Bot. 108:1824–1837.

O'Brien, A. M., C. N. Jack, M. L. Friesen, and M. E. Frederickson. 2021b. Whose trait is it
anyways? Coevolution of joint phenotypes and genetic architecture in mutualisms. Proc.
R. Soc. B Biol. Sci. 288:20202483.

Oldroyd, G. E. D., J. D. Murray, P. S. Poole, and J. A. Downie. 2011. The rules of engagement
in the legume-rhizobial symbiosis. Annu. Rev. Genet. 45:119–144.

Ortíz-Castro, R., M. Martínez-Trujillo, and J. López-Bucio. 2008. N-acyl-L-homoserine lactones:
a class of bacterial quorum-sensing signals alter post-embryonic root development in
Arabidopsis thaliana. Plant Cell Environ. 31:1497–1509.

Peakall, R., and A. J. Beattie. 1996. Ecological and genetic consequences of pollination by
sexual deception in the orchid *Caladenia tentaculata*. Evolution 50:2207–2220.

R Core Team. 2023. R: A language and environment for statistical computing. R Foundation for

681   Statistical Computing, Vienna, Austria.
682   Sachs, J. L., U. G. Mueller, T. P. Wilcox, and J. J. Bull. 2004. The evolution of cooperation. Q.
683   Rev. Biol. 79:26.
684   Schielzeth, H. 2010. Simple means to improve the interpretability of regression coefficients.
685   Methods Ecol. Evol. 1:103–113.
686   Scopece, G., S. Cozzolino, S. D. Johnson, and F. P. Schiestl. 2010. Pollination efficiency and
687   the evolution of specialized deceptive pollination systems. Am. Nat. 175:98–105. The
688   University of Chicago Press.
689   Scopece, G., A. Croce, C. Lexer, and S. Cozzolino. 2013. Components of reproductive isolation
690   between *Orchis mascula* and *Orchis pauciflora*. Evolution 67:2083–2093.
691   Scopece, G., F. P. Schiestl, and S. Cozzolino. 2015. Pollen transfer efficiency and its effect on
692   inflorescence size in deceptive pollination strategies. Plant Biol. 17:545–550.
693   Stanton-Geddes, J., T. Paape, B. Epstein, R. Briskine, J. Yoder, J. Mudge, A. K. Bharti, A. D.
694   Farmer, P. Zhou, R. Denny, G. D. May, S. Erlandson, M. Yakub, M. Sugawara, M. J.
695   Sadowsky, N. D. Young, and P. Tiffin. 2013. Candidate genes and genetic architecture
696   of symbiotic and agronomic traits revealed by whole-genome, sequence-based
697   association genetics in *Medicago truncatula*. PLOS ONE 8:e65688. Public Library of
698   Science.
699   Thompson, J. N. 2005. The geographic mosaic of coevolution. University of Chicago Press.
700   Udvardi, M., and P. S. Poole. 2013. Transport and metabolism in legume-rhizobia symbioses.
701   Annu. Rev. Plant Biol. 64:781–805.
702   Urban, I. 1873. Prodromus einer monographie der gattung Medicago L. Verh Bot Ver Brand.
703   15:1–85.
704   Vidal, M. C., and K. A. Segraves. 2021. Coevolved mutualists experience fluctuating costs and
705   benefits over time. Evolution 75:219–230.
706   Wagner, M. R., D. S. Lundberg, D. Coleman-Derr, S. G. Tringe, J. L. Dangl, and T. Mitchell-
707   Olds. 2014. Natural soil microbes alter flowering phenology and the intensity of selection
708   on flowering time in a wild Arabidopsis relative. Ecol. Lett. 17:717–726.
709   Weese, D. J., K. D. Heath, B. T. M. Dentinger, and J. A. Lau. 2015. Long-term nitrogen addition
710   causes the evolution of less-cooperative mutualists. Evolution 69:631–642.
711   Wendlandt, C. E., J. U. Regus, K. A. Gano-Cohen, A. C. Hollowell, K. W. Quides, J. Y. Lyu, E.
712   S. Adinata, and J. L. Sachs. 2019. Host investment into symbiosis varies among
713   genotypes of the legume Acmispon strigosus, but host sanctions are uniform. New
714   Phytol. 221:446–458.
715   Werner, G. D. A., J. E. Strassmann, A. B. F. Ivens, D. J. P. Engelmoer, E. Verbruggen, D. C.
716   Queller, R. Noë, N. C. Johnson, P. Hammerstein, and E. T. Kiers. 2014. Evolution of
717   microbial markets. Proc. Natl. Acad. Sci. 111:1237–1244. Proceedings of the National
718   Academy of Sciences.
719   Westhoek, A., L. J. Clark, M. Culbert, N. Dalchau, M. Griffiths, B. Jorrin, R. Karunakaran, R.
720   Ledermann, A. Tkacz, I. Webb, E. K. James, P. S. Poole, and L. A. Turnbull. 2021.
721   Conditional sanctioning in a legume–Rhizobium mutualism. Proc. Natl. Acad. Sci. 118.
722   National Academy of Sciences.
723   Wickham, H. 2016. ggplot2: elegant graphics for data analysis. Springer-Verlag, New York.
724   Wood, C. W., B. L. Pilkington, P. Vaidya, C. Biel, and J. R. Stinchcombe. 2018. Genetic conflict
725   with a parasitic nematode disrupts the legume–rhizobia mutualism. Evol. Lett. 2:233–
726   245.
727   Zhang, J., N. He, C. Liu, L. Xu, Z. Chen, Y. Li, R. Wang, G. Yu, W. Sun, C. Xiao, H. Y. H. Chen,
728   and P. B. Reich. 2020. Variation and evolution of C:N ratio among different organs
729   enable plants to adapt to N-limited environments. Glob. Change Biol. 26:2534–2543.