Sex chromosome dosage compensation in a sex reversing skink is not influenced by sexual phenotype

Authors: Benjamin J. Hanrahan[1], J King Chang[1], Ashley M. Milton[1], Nicholas C. Lister[1], Duminda S.B. Dissanayake[2], Jillian M. Hammond[3,4], Andre L.M. Reis[3,4,5], Ira W. Deveson[3,4,5], Aurora Ruiz-Herrera[6,7], Hardip R. Patel[8], Jennifer A. Marshall Graves[2,9], Arthur Georges[2], Paul D. Waters[1]*

Affiliations:

[1] Evolution & Ecology Research Centre, School of Biotechnology and Biomolecular Sciences, Faculty of Science, University of New South Wales, Sydney, NSW 2052, Australia.

[2] Institute for Applied Ecology, University of Canberra, Canberra, ACT 2601 Australia.

[3] Genomics Pillar, Garvan Institute of Medical Research, Sydney, NSW, Australia.

[4] Centre for Population Genomics, Garvan Institute of Medical Research and Murdoch Children's Research Institute, Australia.

[5] Faculty of Medicine, University of New South Wales, Sydney, NSW, Australia.

[6] Departament de Biologia Cel·lular, Fisiologia i Immunologia, Universitat Autònoma de Barcelona, Cerdanyola del Vallès, 08193, Spain.

[7] Genome Integrity and Instability Group, Institut de Biotecnologia i Biomedicina, Universitat Autònoma de Barcelona, Cerdanyola del Vallès, 08193, Spain.

[8] National Centre for Indigenous Genomics, Australian National University, ACT 2601, Australia.

[9] Department of Environment and Genetics, La Trobe University, Melbourne, Victoria 3068, Australia.

26 *corresponding author. Email: p.waters@unsw.edu.au

ABSTRACT

Background

Lizards have sex determination systems that can differ between even closely related species. These include XY and ZW systems, and thermolabile systems where genes and temperature interact to determine sex. The eastern three-lined skink (*Bassiana duperreyi*) has a differentiated XY sex determination system, in which low temperature incubation during development can cause female to male sex reversal, producing XX males. This provides a unique opportunity to investigate how genotype and sexual phenotype affect dosage compensation.

Results

Here, we present a draft genome assembly of the Eastern three-lined skink generated from nanopore sequencing. We also generated transcriptomes from brain and heart tissue of normal adult males and females, along with brain tissue of sex-reversed XX males. We observed partial dosage compensation between XX females and XY males in both brain and heart, with median gene expression from the X in normal males being 0.7 times that of normal females. In brain of sex reversed XX males the median X chromosome output matched that of the normal XX female level, and not that of normal XY males.

Conclusions

Partial dosage compensation in the Eastern three-lined skink is similar to several other species of lizard. However, here for the first time we describe dosage compensation

in a lizard with natural sex reversal, and show that in sex reversed individuals dosage

compensation of the X chromosome follows genotypic sex and not phenotypic sex.

Key words: dosage compensation, sex chromosomes, *Bassiana duperreyi*, sex

reversal

INTRODUCTION

Sex chromosome dosage compensation in mammals was proposed to have evolved

in response to loss of gene function from the mammalian Y chromosome. Gene

expression from the single X in males should need to be upregulated to restore

ancestral autosomal levels. This X upregulation carried through to females resulting in

disproportionately high X expression in XX females (1), and this was countered by X

chromosome inactivation (XCI) to silence one X chromosome in the somatic cells of

females (2). This classic model of dosage compensation appears to hold true for

marsupial mammals, but transcriptional upregulation of the single X appears

incomplete in eutherian mammals (3, 4). In monotremes, median gene expression

from the X chromosome is increased compared to the autosomes in XY males, as it

is for the bird Z in ZW females. However, global X (or Z) transcriptional output is not

evenly balanced between the sexes (5).

Mammals and birds have ancient and relatively stable sex chromosomes and sex

determination systems. Most mammals have a conserved XY system and birds have

a conserved ZW system. In contrast, reptiles display a wide array of sex determination

and sex chromosome systems, even in closely related species (6, 7). These organisms exhibit not only XY and ZW genetic sex determination (GSD) systems, but also temperature-dependent sex determination (TSD), whereby the sex of offspring is determined by the temperature at which the egg is incubated during a thermosensitive window. GSD and TSD systems are often observed in closely related lizard species, and some species have a GSD system that can be overridden by temperature to cause sex reversal (8-11).

Studies of reptile dosage compensation have revealed a variety of non-canonical dosage compensation systems. In snakes, there are reports of partial dosage compensation of the Z chromosome by upregulation of Z-borne genes in females (12, 13). The green anole (*Anolis carolinensis*), a lizard with an XY sex chromosome system, has complete dosage compensation via 2-fold upregulation of the single X in males to match expression with the ancestral autosomal level and balancing expression with the two Xs in females (14), which is mediated by a lncRNA (15). The blacktail brush lizard also has two-fold upregulation of the single X in males (16), but with unknown mechanism. In contrast, the Komodo dragon (*Varanus komodoensis*), which has one of the oldest ZW sex chromosome systems, shows no evidence of dosage compensation of Z-specific genes (17). Partial dosage compensation has been observed in the brown basilisk (*Basiliscus vittatus*) such that the XY males had expression intermediate between full dosage compensation of the two X's in females, and half the expression (18). Incomplete dosage compensation was also observed in Gila monster (19), which has a ZW sex chromosome system.

The scincid lizard *Bassiana duperreyi* (Australian eastern three-lined skink) has a

differentiated XY sex chromosome system (20) that is thought to predate skink

radiation (21). However, the genetic sex determining switch can be overridden by

temperature (8, 9, 11). Reduced incubation temperature of eggs during the

thermosensitive period results in sex reversed XX males in the adult population and

the natural nests of both captive and wild populations (22, 23). This gives rise to three

different genotypic and phenotypic sexes: XX females, XY males and sex reversed

XX males. Therefore, *B. duperreyi* provides a unique opportunity not only to examine

dosage compensation in a reptile with differentiated sex chromosomes, but to also test

if dosage compensation in sex reversed XX males follows genotype or sexual

phenotype.

Here we present a chromosome level *B. duperreyi* draft genome assembly from a male

individual, in which we have identified both the X and Y chromosomes. We generated

mRNA sequence data from brain and heart tissue of genotypically normal XX females

and XY males, and brain tissue of XX male individuals to examine the effect of

genotype and phenotype on dosage compensation. In brain and heart of normal XY

male and XX female individuals, there was partial dosage compensation, such that the

single X of XY males was overexpressed. Sex reversed XX males were observed to

have the same median X gene expression as XX females, showing for the first time

that dosage of the X follows genotype in a sex reversed lizard.

RESULTS

Genome assembly summary

Genome length was 1.485 Gbp, assembled into 5,369 scaffolds. The six largest

scaffolds (288.7 - 75.5 Mbp) corresponded to autosome macrochromosomes (from 1

to 6), with the next six largest scaffolds (54.7 – 18.1 Mbp) representing the

microchromosomes (from 7 to 14) (Figure 1A, Table 1). Based on HiC contacts, contig

235 was found to contain merged microchromosomes, likely chromosomes 12, 13 and

14 (Figure 1A, inset). Including the sex chromosome scaffolds this comprised all the

chromosomes in *B. duperreyi* (2n = 30). The chromosome-scale scaffolds comprised

98.5% of the total assembled sequence. BUSCO analysis showed 96.0% complete

conserved orthologs in the tetrapod database and 96.6% complete conserved

orthologs in the vertebrate database (Table 2).

Figure 1. Genome summary of the *Bassiana duperreyi* assembly

A) Hi-C contact map generated with HiC explorer (53) showing scaffold (chromosome)

boundaries. Inset: magnified view of the HiC contact map for microchromosomes with

size 18.1 – 55.4 Mb, and the sex chromosomes. Number of contacts are between 100

kbp bins. Below: mean inter-chromosomal interactions for 100 kbp windows between

each chromosome. **B)** Number of reads mapped in 20kb windows of male Illumina

genome sequencing on a representative autosome (contig 31), the putative Y (contig

93) and X contigs (contig 1241). Vertical red line is the PAR boundary on the X contig.

C) Synteny across chicken, bassiana and blue tailed skink (*Cryptoblepharus egeriae*)

genomes, generated with GENESPACE (54). The bassiana Y scaffold (contig 93, not

shown) shares no synteny with the other two species.

Table 1. Chromosome identity based on scaffold size

Scaffold Name	Size (bp)	Chromosome Identity
BASDU_flyeontv1_Contig31	288676622	1
BASDU_flyeontv1_Contig51	258390987	2
BASDU_flyeontv1_Contig32	215829809	3
BASDU_flyeontv1_Contig88	205722031	4
BASDU_flyeontv1_Contig66	107145552	5
BASDU_flyeontv1_Contig53	75523483	6
BASDU_flyeontv1_Contig1241	74033703	X
BASDU_flyeontv1_Contig298	54678650	7
BASDU_flyeontv1_Contig235	50055479	12,13,14
BASDU_flyeontv1_Contig328	39280427	8
BASDU_flyeontv1_Contig38	25948193	9
BASDU_flyeontv1_Contig604	22020833	10
BASDU_flyeontv1_Contig93	21659042	Y

| BASDU_flyeontv1_Contig1096 | 18146980 | 11 |

150 **Table 2. BUSCO assessment of the *B. duperreyi* genome**

BUSCO Assessment	Total	Complete and single-copy	Complete and duplicated	Fragmented	Missing	Groups searched
tetrapod_odb10	5156	5098 (96.0%)	58 (1.1%)	44 (0.8%)	110 (2.1%)	38
vertebrate_odb 10	3290	3240 (96.6%)	50 (1.5%)	32 (1.0%)	32 (0.9%)	67

Sex chromosome summary

Contig 93 was identified as the Y chromosome by BLASTn with male specific *B.*

duperreyi k-mer contigs (Supplemental Figure 1). Read depth analysis of short read

Illumina whole genome sequencing from a male individual confirmed contig 1241 as

the X (Figure 1C), and revealed the pseudoautosomal region (PAR) boundary location

at which read coverage increased compared to the X and Y specific regions. The X

and Y chromosomes were 71.2 Mbp and 21.7 Mbp, respectively, and shared a small

PAR of approximately 2.8 Mbp that was assembled onto the X scaffold (Figure 1C).

The Y chromosome had more interactions with the macrochromosomes, than the X

chromosome (Figure 1B). Fewer interactions of the X with autosomes is similar to

eutherian XY systems that also have fewer interactions between the X chromosome

and autosomes (24). As expected, microchromosomes interacted with each other

more than with the macrochromosomes (25).

Annotation identified a total of 13,979 genes in the genome. Of these, 667 were X-

specific genes, sharing homology with chicken chromosome 1 (Figure 1D). A total of

62 genes were annotated on the male specific region of the Y: 47 of these were X/Y

shared, with the remaining 15 having no detected X-borne partner. A total of 39 genes

were annotated in the PAR.

There were no XY shared or Y specific genes with a known role in the sex determining

pathway of other vertebrates (Supplemental Table 1). The only gene conserved on the

Y chromosome of the Christmas Island blue-tailed skink was *Ifit5*, which encodes for

an interferon that has not been implicated in sex determining pathways. Gene Onology

(GO) analysis of the 15 Y specific genes showed no enrichment of genes involved in

male related functions. In fact, these genes were not enriched for any biological

function. GO analysis of XY shared genes, as well as all genes on the Y (XY shared

and Y specific genes), revealed an enrichment of genes involved in histone H4

acetyltransferase activity (Supplemental Tables 2 and 3). In summary, there was no

candidate sex determining gene identified.

The karyotypic difference between the X and Y (20), and the disparity between their

gene contents, confirmed their differentiation. This suggested an unbalanced output

of X-specific genes in XX females and XY males that might be balanced by a sex

chromosome dosage compensation mechanism.

Dosage compensation of the X chromosome

In brain and heart, median gene expression from the autosome scaffolds and PAR

was equivalent between XY males and XX females (Figure 2A). However, median X

chromosome transcriptional output in XY males was reduced compared to XX females

(to 70% in brain and 65% in heart: Mood's median test, $p < 0.0001$).

We next tested if phenotypic sex influenced total transcriptional output from the X

versus the autosomes in brain. We measured the ratios of median X specific to

autosome gene expression levels within the three genotypic and phenotypic sexes. In

XX females and XX males there was no significant difference between X and

autosomal expression levels. In XY males, median transcriptional output from the X

was 72% that of the autosomal median (permutation test, $p < 0.0001$) (Figure 2B).

We then compared transcriptional output of the X specific region between the three

phenotypic and genotypic sexes. Median gene expression from autosome scaffolds of

XX females, XY males and sex reversed XX males were equivalent (Figure 2C),

individual autosome scaffolds were also equivalent (Supplemental Figure 2). PAR

region showed slightly higher expression in XY males, however this was only

significant in the XYm/XXf comparison. As observed for heart and brain above, median

gene expression from the X chromosome was significantly lower in normal XY males

compared to XX females (down to 69%: Mood's median test, $p < 0.0001$) (Figure 2C).

Median gene expression from the X chromosome was also lower in XY males than XX

males (down to 71%: Mood's median test, $p < 0.0001$). These ratios are both greater

than the 50% expected from a non-dosage compensated X chromosome, implying

significant upregulation of the single X in XY males relative to both XX females and

XX males (Figure 2C). This upregulation of genes on the X in XY males compared to

both XX females and XX males was not region specific on the X (Supplemental Figure

3), nor was it restricted to a subset of genes (Supplemental Figure 4).

We also compared expression from the X between sex reversed XX males and normal

XX females. Both have two X chromosomes, so we could directly address if sex

chromosome dosage compensation strictly follows genotype, or was also influenced

by phenotype. We observed that XX males and XX females had equivalent

transcriptional output from the X, with the same expression ratio as the autosomes

(Mood's median test, $p = 0.886$) (Figure 2C), and that the median output from the X

was not different compared to autosomes in either XX males (permutation test, $p =$

0.333) or XX females (permutation test, $p = 0.433$) (Figure 2B).

Having established that sex chromosome dosage compensation followed genotype,

we investigated expression of X genes with a Y gametologue. These X genes (n=42)

had significantly lower total transcriptional output in XY males than did XX females

(Mood's median test, $p < 0.0001$) and XX males (Mood's median test, $p < 0.0001$)

(Figure 2C). However, when counts for X and Y gametologues in XY males were

summed their expression was equal to that of X gametologues in XX males and XX

females (Figure 2D). Median expression of the XY shared genes was found to be

higher than total X gene expression in all sexes (Mood's median test: XXf ,$p < 0.001$;

XXm, $p < 0.001$; XYm, $p < 0.05$) (Figure 2D).

Figure 2. Dosage of gene expression between sexes

A) Median expression male (XY) to female (XX) ratios were calculated for adult brain

and heart tissue (n=1 for both sexes and tissues). A ratio above zero indicates higher

expression in male and a ratio below zero indicates higher expression in female.

Median is plotted in the box with exact value above each median. Boxes represent the

middle 50% of the data, and whiskers represent 1.5 times the interquartile range.

Outliers not plotted. **B)** Brain transcriptomes were used to calculate mean RPKM for

each sex condition (XYm, n=3; XXm, n=2; and XXf, n=3) and plotted for autosomal, X

specific and PAR genes as a ratio to autosomal median. Ratio to the autosomal

median is plotted in the box with exact value above each median. Permutation tests

were used to calculate whether median PAR and X specific RPKMs were statistically

different from the median autosome RPKM, as well as if the median PAR RPKM was

different from the median X specific RPKM (**** $p<0.0001$, *** $p<0.001$, * $p<0.05$). **C)**

Median expression ratios were calculated for pairwise comparisons of the three sexes

(XYm, XXm and XXf). Genomic regions were separated and plotted as follows;

autosomes (green), PAR (blue), total X output (red) and XY shared genes (pink).

Mood's median tests were used to calculate if the ratios for the autosomes and X, as

well as Xs between sexes, were statistically different (**** $p<0.0001$). **D)** As above

mean RPKM values were calculated for each sex conditions and represented as a

boxplot. Genomic regions on the X chromosome were separated as follows; X (red),

XY shared Y (light pink), XY shared X (pink) and sum of X and Y ametologues (dark

pink).

DISCUSSION

Here we present a draft assembly for *B. duperreyi* with chromosome level scaffolds

that included both X and Y chromosomes. Gene expression of the X chromosome

between XY males and XX females revealed partial dosage compensation of the X in

XY males. Dosage of the genes on the X chromosome in sex reversed XX males was

equal to the X in XX females, therefore sex dosage compensation followed genotype.

In normal XY male *B. duperreyi* there was partial dosage compensation of the X

chromosome (Figure 2C), with median X chromosome gene expression 69% that of

XX females. Therefore, the single X chromosome is being over-expressed by a factor

of approximately 1.45. This is similar to observations in non-therian mammal

vertebrates such as platypus (5). It is also similar to the dosage system seen in the

brown basilisk (18) and Gila monster (19). However, it is different from dosage

compensation in other lizard species; the absence of sex chromosome dosage

compensation in the ZW system of the Komodo dragon (17), and complete dosage

compensation in the green anole (14) and the blacktail brush lizard (16).

Thus, we demonstrate that *B. duperreyi* represents a partial dosage compensation

system in a lizard that is chromosome-wide with no region-specific effect. Thus

different dosage compensation systems have evolved repeatedly and independently

in different lizard lineages, perhaps depending on the dosage sensitivity of the

progenitor autosomes from which the sex chromosome systems evolved. Study of

more lizards will elucidate the breadth of reptile sex chromosome dosage

compensation and highlight different evolutionary strategies.

As well as comparing normal XY males and XX females, for the first time, we describe

dosage compensation in a model with naturally occurring sex reversal. We observed

that gene expression in the sex reversed XX males was equivalent to that in normal

XX females, with median X expression 99% that of XX females and 140% that of XY

males. Therefore, it is genotype, as opposed to sexual phenotype, that dictates sex

chromosome dosage compensation in *B. duperreyi*. This is the first example showing

dosage compensation in a sex reversed animal matches that of the sex with the same

genotype. It remains to be seen if the offspring of normal XX females and sex reversed

XX males also follow this trend of dosage compensation matching normal females, as

it is unknown if sex reversed XX males are reproductively viable (22, 26), though it is

likely this is the case.

Temperature-induced sex reversal in lizards is not unique to *B. duperreyi,* having been

observed in several species (8, 10). It is possible that sex reversal in lizards is more

widespread than previously believed, as it has been systematically investigated in only

a few species (27), and there may be numerous other species whose sex reversal

status is yet to be confirmed (see 28, 29, 30). Examining dosage compensation in

other sex reversing models will confirm whether our finding that phenotype does not

influence sex chromosome dosage is ubiquitous.

Recent evidence suggests that in systems with partial or incomplete dosage

compensation in the transcriptome, the abundance of proteins can remain balanced

(31). This should be investigated further in organisms such as *B. duperreyi,* brown

basilisk, Gila monster and komodo dragon that show partial or lack of dosage

compensation in the transcriptome, to observe if the sex chromosome output is

balanced in the proteome.

CONCLUSIONS

Here we analysed transcriptomes from brain and heart of adult *B. duperreyi* to show

that in canonical XY males there is partial dosage compensation of the X chromosome.

The brain transcriptome of hatchling brains also revealed that sex reversed XX males

have the same transcriptomic output from the X chromosome as canonical XX

females. This is the first description of X chromosome dosage in a naturally sex-

reversing system and demonstrates that, in *B. duperreyi*, dosage follows the

underlying genotype, regardless of an individual's sex.

MATERIALS AND METHODS

Sample collection and sexing

In December 2020, samples (eggs) of alpine *B. duperreyi* were collected from nests

in the field location within the Brindabella Range (Piccadilly Circus – 1240 m a.s.l.,

35°21'42.0"S 148°48'12.5"E) after ensuring that approximately 90% of the

development period had passed in natural conditions (9). The eggs were then

collected and transported to the University of Canberra, where they were buried in

moist vermiculite with a ratio of 4 parts water to 5 parts vermiculite by weight, and

placed in incubators (LabWit, ZXSDR1090) that maintained 23°C, which produces a

balanced sex ratio (9). Details egg collection methods (23) and a description of the

alpine study site can be found in (32).

To determine phenotypic sex of the hatchlings, tail bases of 7-day old hatchlings were

squeezed to evert the hemipenes (33) and sex was checked again by hemipene

transillumination after 5 weeks (23). Tail snips were collected to determine the

genotype of the lizards. Genotypic sex was determined for *B. duperreyi* using

polymerase chain reaction (PCR)-based molecular sex tests from extracted DNA

collected from tissue samples. DNA purity was determined using a NanoDrop 1000

spectrophotometer (NanoDrop Technologies Inc., Wilmington, DE, USA) and

quantified using the Qubit 2.0 Fluorometric Quantitation (Invitrogen, Life technologies,

Sydney, N.S.W., Australia). The sex-reversal status was determined for *B. duperreyi*

using PCR as described by Dissanayake et al. (20), where the genotypic sex was

identified based on Y-specific markers allowing identification of XY males (XYm) and

XX males (XXm). The lizards were euthanised via intraperitoneal injection of sodium

pentobarbitone (100–150 µg/g body weight) as approved by the University of

Canberra Animal Ethics Committee (AEC 17–26).

RNA extraction and sequencing

Total RNA was extracted from the brain tissue of three XY males (XYm), three XX

females (XXfm), and two XX males (XXm). Tissue extracts were homogenized using

T10 Basic ULTRA-TURRAX® Homogenizer (IKA, Staufen im Breisgau, Germany) and

extracted using TRIzol reagent following the manufacturer's instructions, purifying with

an isopropanol precipitation. Seventy-five bp single-ended reads were generated on

the Illumina NextSeq 500 platform at the Ramaciotti Centre for Genomics (UNSW,

Sydney, Australlia).

DNA extraction and sequencing for assembly

Genomic DNA was extracted from 13 mg of ethanol-preserved muscle tissue from a

male (XY) *B. duperreyi*, using the Circulomics Nanobind tissue kit as per the

manufacturer's protocols, including the specified pre-treatment for ethanol removal.

Library preparation was performed with 3 µg of DNA as input, using the SQK-LSK109

kit from Oxford Nanopore Technologies and sequenced across two promethION (FLO-

PRO002) flow cells, with washes (EXP-WSH004) performed every 24 hours.

Genome assembly

The genome assembly pipeline relied on using a combination of whole-genome

sequencing ONT long reads, Illumina short reads and Hi-C reads. Firstly, adapters

were removed from Illumina and Hi-C reads using TrimGalore v0.6.6 with default

parameters. A primary assembly was generated using Flye v2.8.3 (34) and the ONT

long reads, with the following parameters "--trestle --iterations 2". Illumina short reads

were aligned to the primary assembly with bwa-mem v0.7.17-r1188 and ONT long

reads were aligned with minimap2 v2.24-r1122 (35), and both alignments were used

to polish the assembly with hypo v0.5.1. Homologous contigs were identified and

removed using Purge Haplotigs v1.1.1 (36). Chromosome-length scaffolding was

performed with Hi-C data using Juicer v1.6 (37) to generate a contact matrix of the

connections between contigs and 3d-dna v180922 to organise the contigs into larger

scaffolds. Gaps in the assembly were filled using PBjelly v15.8.24 (38) with the ONT

reads. The final assembly was assessed for completeness using BUSCO and

accuracy using Merqury v1.3 (39), by comparing the assembly k-mer spectrum to

those found in the Illumina reads.

Annotation

The annotation used in this study was generated using AUGUSTUS v3.4.0 (40) with

RNA sequencing data from brain, heart and gonad from male and female individuals.

Prior to running AUGUSTUS, a transcriptome assembly and a soft-masked genome

assembly were generated with Trinity v2.12.0 (41) and RepeatMasker v4.1.2-p1 (42),

respectively. AUGUSTUS parameters were optimised with a training set of 500 genes,

including 5 rounds of optimisation with exon and UTR parameters, before gene

prediction and stitching. Peptides were inferred from the final annotation using gffread

v0.12.7 (43) and blasted against the uniprot database with blast v2.11.0 (44) for gene

identification. A github repository of this pipeline can be found at

https://github.com/kango2/Annotation.

Y specific sequences (20) in *B. duperreyi* were blasted against the generated genome

assembly with blast v2.11.0 (44).

Read depth analysis

DNA was extracted from muscle samples of the individual animals using the Gentra

Puregene Tissue Kit (QIAGEN, Australia), following the manufacturer's protocols with

the modifications described below. The volume and reagent amounts were adjusted

according to the size of the tissue sample, using three times more reagent than

specified in the manufacturer's protocols. Additionally, we made modifications to the

DNA precipitation steps outlined in the manufacturer's protocol. The DNA thread was

spooled out using a galls rod and submerged in 300ul of 70% ethanol, then air dried

for one minute. Subsequently, we used TE buffer for DNA hydration and allowed it to

dissolve overnight at room temperature.

Raw read quality of the DNA sequencing was assessed using FastQC v0.11.9 (45)

and trimmed accordingly with trimmomatic v0.38 (46). Trimmed reads were aligned to

the newly generated *B. duperreyi* genome assembly with subread-align command in

subread v2.0.1 (47). Read depth in 20000 bp windows was calculated using the

bamCoverage command, from deeptools v3.5.1 (48), for the X and Y scaffolds as well

as a representative autosomal scaffold (contig 328). The generated BED files were

then used to plot read depth across these scaffolds using the ggplot2 package in R

v4.2.1.

Bioinformatics analysis of RNA-seq

419
Raw read quality of RNA sequencing was assessed using FastQC v0.11.9 (45) and

trimmed accordingly with trimmomatic v0.38 (46). Trimmed reads were aligned to the

newly generated *B. duperreyi* genome assembly with subread-align function in

subread v2.0.1 (47). The subread-featurecount function from subread v2.0.1 (49) was

used to count reads that overlapped genomic features using the settings "-0 -2 -t

CDS,five_prime_utr,three_prime_utr", the remaining settings were left as default.

Library size and read counts were normalised by counts per million and RPKM

respectively using the edgeR package in R v4.2.1 (50). Gene expression ratios were

calculated and plotted with the ggplot2 package. Gene expression values were

calculated from the RNA-seq data (hatchling brain tissue) of XX females (n=3), XY

males (n=3) and sex reversed XX males (n=2). Normalised expression values were

used to create male to female median expression ratios between the three genotypic

and phenotypic sexes for each gene. Genes were binned according to their location

on an autosome, the PAR, X as total output and X genes with Y gametologues (XY

shared).

Gene ontology and gene function

437 Gene ontology analysis was performed using g:Profiler (51) with gene lists identified

438 using the *B. duperreyi* annotation. Results were restricted to biological process and

439 molecular function and used the human database.

441 Protein functions of Y genes were obtained using the panther database v19.0 (52).

DECLARATIONS

Ethics approval and consent to participate

The animals involved in this study were collected from nests in the field under permits issued by the Australian state government: ACT permit numbers (LT201826, LT2017956) and NSW government (SL102002). The collection was carried out in accordance with their respective guidance and protocols, as well as the University of Canberra Animal Ethics Committee (Approval Number AEC 17–26). All husbandry practices complied with the Australian Code for the Care and Use of Animals for Scientific Purposes, 8th edition (2013), particularly sections 3.2.13–3.2.2.

Consent for publication

Not applicable

Availability of data and materials

All raw sequencing data generated in this study have been submitted to the NCBI BioProject database (https://www.ncbi.nlm.nih.gov/bioproject/) under accession number PRJNA980841. The genome assembly presented in this study is available from the NCBI Genome database under GenBank accession GCA_041722995.1.

Competing interests

The authors declare that they have no competing interests.

Funding

B.J.H. is supported by an Australian Government Research Training Program (RTP)

Scholarship. P.D.W. and J.A.M. are supported by Australian Research Council

Discovery Projects (DP210103512 and DP220101429). P.D.W. and H.R.P. are

supported by an NHMRC Ideas Grant (2021172). A.R-H. acknowledges the Spanish

Ministry of Science and Innovation (PID2020-112557GB-I00 founded by

AEI/10.13039/501100011033), the Agència de Gestió d'Ajuts Universitaris i de

Recerca, AGAUR (2021SGR00122) and the Catalan Institution for Research and

Advanced Studies (ICREA). Fieldwork and initial laboratory work were supported by

Australian Research Council Grants DP110104377 and DP170101147, awarded to

A.G.

Authors' contributions

B.J.H wrote the original draft. P.D.W and A.G designed the study. D.S.B.D collected

and harvested samples. B.J.H and J.K.C conducted extraction experiments to

generate data. A.L.M.R, J.M.H and J.K.C generated the genome assembly and

annotation. B.J.H and J.K.C conducted data analysis with input from A.M.M and N.C.L.

D.S.B.D, I.W.D, A.R-H, H.R.P, J.A.M.G and A.G provided insight into data

interpretation and writing. All authors reviewed and approved the submitted

manuscript.

Acknowledgements

Sequencing costs for this project were met by grants from Bioplatforms Australia

(AusARG initiative) and the Australian Research Council (DP170101147). We

acknowledge the Garvan Sequencing Platform.

REFERENCES

1.   Ohno S. Sex chromosomes and sex-linked genes. Berlin, New York etc.:
Springer-Verlag; 1967. x, 192 p. p.

2.   Lyon MF. X-chromosome inactivation and developmental patterns in
mammals. Biol Rev Camb Philos Soc. 1972;47(1):1-35.

3.   Xiong Y, Chen X, Chen Z, Wang X, Shi S, Wang X, et al. RNA sequencing
shows no dosage compensation of the active X-chromosome. Nat Genet.
2010;42(12):1043-7.

4.   Lin F, Xing K, Zhang J, He X. Expression reduction in mammalian X
chromosome evolution refutes Ohno's hypothesis of dosage compensation. Proc
Natl Acad Sci U S A. 2012;109(29):11752-7.

5.   Julien P, Brawand D, Soumillon M, Necsulea A, Liechti A, Schutz F, et al.
Mechanisms and evolutionary patterns of mammalian and avian dosage
compensation. PLoS Biol. 2012;10(5):e1001328.

6.   Sarre SD, Georges A, Quinn A. The ends of a continuum: genetic and
temperature-dependent sex determination in reptiles. Bioessays. 2004;26(6):639-45.

7.   Ezaz T, Sarre SD, O'Meally D, Graves JAM, Georges A. Sex chromosome
evolution in lizards: independent origins and rapid transitions. Cytogenet Genome
Res. 2009;127(2-4):249-60.

8.   Quinn AE, Georges A, Sarre SD, Guarino F, Ezaz T, Graves JA. Temperature
sex reversal implies sex gene dosage in a reptile. Science. 2007;316(5823):411.

9.   Shine R, Elphick MJ, Donnellan S. Co-occurrence of multiple, supposedly
incompatible modes of sex determination in a lizard population. Ecol Lett.
2002;5(4):486-9.

10.   Holleley CE, O'Meally D, Sarre SD, Graves JAM, Ezaz T, Matsubara K, et al.
Sex reversal triggers the rapid transition from genetic to temperature-dependent sex.
Nature. 2015;523(7558):79-82.

11.   Radder RS, Quinn AE, Georges A, Sarre SD, Shine R. Genetic evidence for
co-occurrence of chromosomal and thermal sex-determining systems in a lizard. Biol
Lett. 2008;4(2):176-8.

12.   Vicoso B, Emerson JJ, Zektser Y, Mahajan S, Bachtrog D. Comparative sex

chromosome genomics in snakes: differentiation, evolutionary strata, and lack of

global dosage compensation. PLoS Biol. 2013;11(8):e1001643.

13.   Schield DR, Card DC, Hales NR, Perry BW, Pasquesi GM, Blackmon H, et al.

The origins and evolution of chromosomes, dosage compensation, and mechanisms

underlying venom regulation in snakes. Genome Res. 2019;29(4):590-601.

14.   Marin R, Cortez D, Lamanna F, Pradeepa MM, Leushkin E, Julien P, et al.

Convergent origination of a Drosophila-like dosage compensation mechanism in a

reptile lineage. Genome Res. 2017;27(12):1974-87.

15.   Tenorio M, Cruz-Ruiz S, Encarnacion-Guevara S, Hernandez M, Corona-

Gomez JA, Sheccid-Santiago F, et al. MAYEX is an old long noncoding RNA

recruited for X chromosome dosage compensation in a reptile. Science.

2024;385(6715):1347-54.

16.   Davalos-Dehullu E, Baty SM, Fisher RN, Scott PA, Dolby GA, Munguia-Vega

A, et al. Chromosome-Level Genome Assembly of the Blacktail Brush Lizard,

Urosaurus nigricaudus, Reveals Dosage Compensation in an Endemic Lizard.

Genome Biol Evol. 2023;15(12).

17.   Rovatsos M, Rehak I, Velenský P, Kratochvil L. Shared Ancient Sex

Chromosomes in Varanids, Beaded Lizards, and Alligator Lizards. Mol Biol Evol.

2019;36(6):1113-20.

18.   Acosta A, Suarez-Varon G, Rodriguez-Miranda LA, Lira-Noriega A, Aguilar-

Gomez D, Gutierrez-Mariscal M, et al. Corytophanids Replaced the Pleurodont XY

System with a New Pair of XY Chromosomes. Genome Biol Evol. 2019;11(9):2666-

77.

19.   Webster TH, Vannan A, Pinto BJ, Denbrock G, Morales M, Dolby GA, et al.

Lack of Dosage Balance and Incomplete Dosage Compensation in the ZZ/ZW Gila

Monster (Heloderma suspectum) Revealed by De Novo Genome Assembly.

Genome Biol Evol. 2024;16(3).

20.   Dissanayake DSB, Holleley CE, Hill LK, O'Meally D, Deakin JE, Georges A.

Identification of Y chromosome markers in the eastern three-lined skink (Bassiana

duperreyi) using in silico whole genome subtraction. BMC Genomics.

2020;21(1):667.

21.   Kostmann A, Kratochvil L, Rovatsos M. Poorly differentiated XX/XY sex
chromosomes are widely shared across skink radiation. Proc Biol Sci.
2021;288(1943):20202139.
22.   Dissanayake DSB, Holleley CE, Deakin JE, Georges A. High elevation
increases the risk of Y chromosome loss in Alpine skink populations with sex
reversal. Heredity (Edinb). 2021;126(5):805-16.
23.   Dissanayake DSB, Holleley CE, Georges A. Effects of natural nest
temperatures on sex reversal and sex ratios in an Australian alpine skink. Sci Rep.
2021;11(1):20093.
24.   Álvarez-González L, Arias-Sardá C, Montes-Espuña L, Marín-Gual L, Vara C,
Lister NC, et al. Principles of 3D chromosome folding and evolutionary genome
reshuffling in mammals. Cell Rep. 2022;41(12):111839.
25.   Waters PD, Patel HR, Ruiz-Herrera A, Álvarez-González L, Lister NC,
Simakov O, et al. Microchromosomes are building blocks of bird, reptile, and
mammal chromosomes. Proc Natl Acad Sci U S A. 2021;118(45).
26.   Dissanayake DSB. Sex reversal in the alpine skink Bassiana duperreyi –
response to natural environment [Doctoral Thesis]: University of Canberra; 2022.
27.   Holleley CE, Sarre SD, O'Meally D, Georges A. Sex Reversal in Reptiles:
Reproductive Oddity or Powerful Driver of Evolutionary Change? Sex Dev.
2016;10(5-6):279-87.
28.   Whiteley SL, Castelli MA, Dissanayake DSB, Holleley CE, Georges A.
Temperature-Induced Sex Reversal in Reptiles: Prevalence, Discovery, and
Evolutionary Implications. Sex Dev. 2021;15(1-3):148-56.
29.   Cornejo-Páramo P, Dissanayake DSB, Lira-Noriega A, Martínez-Pacheco ML,
Acosta A, Ramírez-Suástegui C, et al. Viviparous Reptile Regarded to Have
Temperature-Dependent Sex Determination Has Old XY Chromosomes. Genome
Biol Evol. 2020;12(6):924-30.
30.   Van Dyke JU, Thompson MB, Burridge CP, Castelli MA, Clulow S,
Dissanayake DSB, et al. Australian lizards are outstanding models for reproductive
biology research. Australian Journal of Zoology. 2020;68(4):168-99.
31.   Lister NC, Milton AM, Patel HR, Waters SA, Hanrahan BJ, McIntyre KL, et al.
Incomplete transcriptional dosage compensation of chicken and platypus sex
chromosomes is balanced by post-transcriptional compensation. Proc Natl Acad Sci
U S A. 2024;121(32):e2322360121.

32.   Dissanayake DSB, Holleley CE, Sumner J, Melville J, Georges A. Lineage

diversity within a widespread endemic Australian skink to better inform conservation

in response to regional-scale disturbance. Ecol Evol. 2022;12(3):e8627.

33.   Harlow PS. A Harmless Technique for Sexing Hatchling Lizards. Herpetol

Rev. 1996;27(2):71-.

34.   Kolmogorov M, Yuan J, Lin Y, Pevzner PA. Assembly of long, error-prone

reads using repeat graphs. Nat Biotechnol. 2019;37(5):540-6.

35.   Li H. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics.

2018;34(18):3094-100.

36.   Roach MJ, Schmidt SA, Borneman AR. Purge Haplotigs: allelic contig

reassignment for third-gen diploid genome assemblies. BMC Bioinformatics.

2018;19(1):460.

37.   Durand NC, Shamim MS, Machol I, Rao SS, Huntley MH, Lander ES, et al.

Juicer Provides a One-Click System for Analyzing Loop-Resolution Hi-C

Experiments. Cell Syst. 2016;3(1):95-8.

38.   English AC, Richards S, Han Y, Wang M, Vee V, Qu J, et al. Mind the gap:

upgrading genomes with Pacific Biosciences RS long-read sequencing technology.

PLoS One. 2012;7(11):e47768.

39.   Rhie A, Walenz BP, Koren S, Phillippy AM. Merqury: reference-free quality,

completeness, and phasing assessment for genome assemblies. Genome Biol.

2020;21(1):245.

40.   Stanke M, Diekhans M, Baertsch R, Haussler D. Using native and syntenically

mapped cDNA alignments to improve de novo gene finding. Bioinformatics.

2008;24(5):637-44.

41.   Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, et al.

Full-length transcriptome assembly from RNA-Seq data without a reference genome.

Nat Biotechnol. 2011;29(7):644-52.

42.   Smit AFA, Hubley R, Green P. RepeatMasker Open-4.0. 2013-2015.

43.   Pertea G, Pertea M. GFF Utilities: GffRead and GffCompare. F1000Res.

2020;9.

44.   Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al.

BLAST+: architecture and applications. BMC Bioinformatics. 2009;10:421.

45.   Andrews S. FastQC: a quality control tool for high throughput sequence data.

2010.

46.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinform. 2014;30(15):2114-20.

47.   Liao Y, Smyth GK, Shi W. The Subread aligner: fast, accurate and scalable

read mapping by seed-and-vote. Nucleic Acids Res. 2013;41(10):e108.

48.   Ramirez F, Ryan DP, Gruning B, Bhardwaj V, Kilpert F, Richter AS, et al.

deepTools2: a next generation web server for deep-sequencing data analysis.

Nucleic Acids Res. 2016;44(W1):W160-5.

49.   Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose

program for assigning sequence reads to genomic features. Bioinformatics.

2014;30(7):923-30.

50.   Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for

differential expression analysis of digital gene expression data. Bioinformatics.

2010;26(1):139-40.

51.   Kolberg L, Raudvere U, Kuzmin I, Adler P, Vilo J, Peterson H. g:Profiler-

interoperable web service for functional enrichment analysis and gene identifier

mapping (2023 update). Nucleic Acids Res. 2023;51(W1):W207-W12.

52.   Thomas PD, Ebert D, Muruganujan A, Mushayahama T, Albou LP, Mi H.

PANTHER: Making genome-scale phylogenetics accessible to all. Protein Sci.

2022;31(1):8-22.

53.   Ramirez F, Bhardwaj V, Arrigoni L, Lam KC, Gruning BA, Villaveces J, et al.

High-resolution TADs reveal DNA sequences underlying genome organization in

flies. Nat Commun. 2018;9(1):189.

54.   Lovell JT, Sreedasyam A, Schranz ME, Wilson M, Carlson JW, Harkess A, et

al. GENESPACE tracks regions of interest and gene copy number variation across

multiple genomes. Elife. 2022;11.

SUPPLEMENTAL FIGURE LEGENDS

Supplemental Figure 1 Blastn hits to the the largest 15 scaffolds representing the *B.*

duperreyi chromosomes with males specific kmer contigs.

Supplemental Figure 2. Expression ratios for each autosomal scaffold for the three

pairwise sex/ genotype comparisons presented as boxplots. Number of genes (n) is

show in the key for each contig.

Supplemental Figure 3. Expression ratios were calculated for each gene on **A**) a

representative autosome (contig 31) and **B)** the X chromosome (contig 1241), in a

pairwise fashion for each sex (XYm, XXm and XXf). Values are plotted on a log2 scale

and median values for each comparison are plotted as a red line. For panel B median

for the X specific region is a red line and PAR is a blue line.

Supplemental Figure 4 Scatterplot of counts per million (CPM) for each gene for each

pairwise comparison of XXf to XYm, XXm to XYm and XXf to XXm. X genes are plotted

in black with slope plotted as a red line. Autosomal genes are plotted in blue with slope

as a green line. Slope and R-squared of each trendline is shown above each plot.

Supplemental Table 1 List of genes on the Y chromosome (contig 93) including

protein function

Supplemental Table 2 Gene ontology results for XY shared genes

Supplemental Table 3 Gene ontology results for all Y genes