Non-Invasive Perfusion MR Imaging of the Human Brain via Breath-Holding

J.B. Schulman[1,2*], S. Kashyap[2], S.G. Kim[3,4], K. Uludağ[1,2,3,4*]

[1] Department of Medical Biophysics, University of Toronto, Toronto, ON, Canada

[2] Krembil Brain Institute, University Health Network, Toronto, ON, Canada

[3] Center for Neuroscience Imaging Research, Institute for Basic Science, Suwon, Republic of Korea

[4] Department of Biomedical Engineering, Sungkyunkwan University, Suwon, Republic of Korea

Correspondence:

Jacob Schulman

jacob.schulman@mail.utoronto.ca

Dr. Kamil Uludağ

Kamil.uludag@uhn.ca

Work Address:

Krembil Brain Institute

Toronto Western Hospital, University Health Network

399 Bathurst Street, Room 12MCL405

Toronto, Ontario, Canada M5T 2S8

Abstract

Dynamic susceptibility contrast (DSC) MRI plays a pivotal role in the accurate diagnosis and prognosis of several neurovascular diseases, but the technique's reliance on gadolinium, an intravascularly injected metal, presents a major limitation. Here, we determined the feasibility of using breath-holds to non-invasively induce DSC perfusion contrast. To achieve this, we acquired T_2^*-weighted MRI data at both 3T and 7T from ten healthy participants who engaged in eight consecutive breath-holds (16 s each separated by 44 s of rest). By pairing a novel arterial input function with a standard DSC MRI analysis, we successfully quantified essential perfusion parameters, including the cerebral blood volume, flow, and transit delay, and found these perfusion values to be congruent with those documented in the literature using gadolinium. Breath-holding resulted in significantly higher contrast-to-noise (6.2 at 3T *vs* 8.5 at 7T) and gray matter-to-white matter contrast at higher field strength. For the first time, we have demonstrated the feasibility of DSC using breath-holds—we are confident that this technique holds major

36 promise, both clinically and in research, by being a safe alternative to invasive methods, particularly at
37 ultra-high field strength.
38

39 **Keywords: MRI, Perfusion, DSC, Contrast, Deoxyhemoglobin**

40
41

Introduction

Cerebral perfusion imaging allows clinicians to investigate diseases characterized by vascular
abnormality, including stroke, cancer, and even neurodegenerative disease (Copen *et al.,* 2011; Harris *et*
al., 1998; Law *et al.,* 2008; Qiao *et al.,* 2016). Dynamic susceptibility contrast (DSC) MRI is a standard
perfusion imaging technique wherein a bolus of paramagnetic contrast agent is tracked as it passes
through the cerebral vasculature—the associated time course properties reflect underlying tissue
perfusion in accordance with the principles of indicator dilution theory (Jahng *et al.,* 2014; Rosen *et al.,*
1990; Welker *et al.,* 2015; Zierler, 1962). DSC yields perfusion metrics possessing significant research and
clinical utility, including the cerebral blood volume (*CBV*), cerebral blood flow (*CBF*), and mean transit
time (*MTT*) (Copen *et al.,* 2011; Harris *et al.,* 1998; Law *et al.,* 2008; Qiao *et al.,* 2016; Tanaka *et al.,* 2011). Of
note, all DSC perfusion metrics are technically 'relative' in that they are dependent on numerous
acquisition and analysis parameters, as has been previously documented (Kiselev, 2001; Schulman *et al.,*
2023).
56
While gadolinium (Gd), an exogenous paramagnetic metal, is the standard contrast agent used clinically
for DSC, researchers have recently exploited the paramagnetic properties of deoxyhemoglobin (dOHb) as
an endogenous contrast alternative (Lee *et al.,* 2022; Poublanc *et al.,* 2021; Sayin *et al.,* 2022; Schulman *et al.,*
2023; Vu *et al.,* 2021; Vu *et al.,* 2023)—in essence, both Gd and dOHb induce T_2^* signal changes that can be
exploited in a DSC analysis. However, Gd is limited in that it is invasive, expensive, and exhibits toxicity
in certain patient populations (Kanda, *et al.,* 2015; Rogowska *et al.,* 2018; Schlaudecker and Bernheisel,
2009). On the other hand, generating dOHb contrast has thus far relied on gas control systems that
modify dOHb concentration in the blood through the induction of hypoxia (Lee *et al.,* 2022; Poublanc *et*
al., 2021; Sayin *et al.,* 2022; Schulman *et al.,* 2023; Vu *et al.,* 2021), hyperoxia (MacDonald *et al.,* 2018), or
hyper/hypocapnia (Vu *et al.,* 2023). Given the associated cost, set-up time, and expertise required for gas
control system utilization, the widespread clinical implementation of dOHb contrast as an alternative to
Gd is currently limited.
69
Unlike hypoxia and hyperoxia, hypercapnia (i.e., an increase in arterial CO_2) does not directly result in a
change to the concentration of dOHb—rather, the CO_2 bolus induces vasodilation through a process
known as cerebrovascular reactivity (CVR), resulting in a simultaneous rise of *CBF* and reduction of
dOHb in the tissue (Liu *et al.,* 2019). Given these inherent mechanistic differences, the use of hypercapnia
for DSC relies on certain assumptions, including the constancy of oxygen metabolism, and most
importantly, the ability to measure an arterial input function (AIF), which represents the input time
course of contrast agent, or in this case, vasodilatory agent (see Discussion).
77
Physiologically, the effect of breath-holding is roughly equivalent to that of hypercapnia induced by a gas
control system (Bright *et al.,* 2011; Bright and Murphy, 2013; Liu *et al.,* 2019). However, there is no study
that has developed and/or applied a DSC analysis pipeline to obtain perfusion information from signal
changes induced by breath-holding. Thus, a breath-hold DSC (bhDSC) approach may provide, for the
first time, DSC perfusion metrics without the use of Gd or a gas control system. To develop the bhDSC

approach and test its feasibility, we acquired T_2*-weighted MRI data at 3T and 7T from ten healthy
subjects who performed eight breath-holds (16 s each separated by 44 s of rest). In summary, bhDSC
holds major promise as a perfusion imaging alternative that may circumvent the need for invasive
contrast agents and expensive gas control system equipment.
87
88

89 ## Results
90
91 Gradient-echo 2D-EPI (GRE-EPI) imaging data were acquired on Siemens 3T Prisma and 7T Terra
92 scanners from ten healthy subjects—each subject was scanned at both field strengths. The subjects were
93 instructed to fixate on a countdown timer projected onto the back of the scanner, which indicated when
94 to breathe regularly and when to perform a breath-hold. The first eight breath-hold "boluses" were
95 averaged, with each bolus allotted a 72 s window centered at the bolus maximum (Figure 1).
96

97
Figure 1. Summary of the Breath-hold Experiment. A. Description of the cerebrovascular reactivity phenomenon. At
baseline, tissue blood oxygenation is roughly between 60-80%. Increased CO_2 from breath-holding causes
vasodilation in the upstream arterial/arteriolar vasculature, leading to increased blood flow, and subsequently, a
decrease in the downstream tissue dOHb concentration (hyperoxia), resulting in an MRI signal increase. **B.** Signal
time course (blue) resulting from nine consecutive breath-holds (green) in a venous voxel. **C.** The signal time course
resulting from the temporal average of the first eight boluses was then used for perfusion quantification.
104

105 *Time Course Properties in bhDSC*
106
107 The temporally averaged signal time courses were then converted to relaxation rate time courses (ΔR_2*(t))
108 by normalizing to the pre- and post- breath-hold baselines (see Methods).
109

Figure 2. Breath-Hold-Induced Relaxation Rate Time Course Dynamics. Time courses were averaged across subjects for gray matter (GM), white matter (WM), vein (VOF), and artery (AIF) at 3T and 7T.

Figure 2 shows the breath-hold-induced ΔR_2^* time courses averaged across all subjects, for arterial (AIF), venous (VOF), gray matter (GM), and white matter (WM) voxels at 3T and 7T. The $\Delta R_2^*(t)$ bolus roughly follows the shape of a gamma variate function, with dispersion and delay increasing from input (AIF), to tissue (GM then WM), and ultimately to output (VOF), as is expected physiologically (Figure 2). Unlike the other time courses, the AIF has a positive ΔR_2^* (to be discussed in Figure 4). This contrasts with Gd-DSC experiments where the $\Delta R_2^*(t)$ has the same sign in the AIF as it does in the GM, WM and VOF. The bolus magnitude (defined here as the $\Delta R_2^*(t)$ integral (AUC)) was found to be 1.5-2 times higher in GM relative to WM. Importantly, breath-holding at 7T resulted in larger bolus magnitudes in tissue (by ~2 times), artery (by ~2 times), and vein (by ~3 times) relative to 3T—this is also illustrated in the subject-wise data (Figure S1) and simulation results (Figure S2).

Breath-Hold CNR Properties

The contrast-to-noise ratio (CNR) and associated GM-to-WM contrast were subsequently quantified (see Methods for details). Although eight breath-holds were averaged in the main results of our work, we also studied the effect of averaging fewer breath-holds to determine whether shorter scan time (fewer breath-holds) would yield sufficient CNR.

Figure 3. Breath-hold *CNR* Properties. A. *CNR* maps were calculated (Eq. 2), transformed to MNI152 2 mm anatomical space, and averaged across subjects for both 3T and 7T (axial view is slightly dorsal to the lateral ventricles). For illustration purposes, *CNR* maps are shown when there is no bolus averaging or 2, 4, 6, or 8 boluses averaged at 7T. **B.** Plot of subject-averaged *CNR*$_{GM}$ at 3T (dashed) and 7T (solid) as a function of the number of breath-holds averaged (error bars represent the standard error). **C.** Linear regression of subject-wise GM (red) and WM (blue) *CNR* values at 3T *vs* 7T. *m* represents the regression slope.

Figure 3 shows the subject-averaged *CNR* at 3T and 7T when modifying the number of breath-holds averaged (Figure 3A and 3B), along with a subject-wise regression for *CNR* values between 3T and 7T (Figure 3C). As more breath-holds were averaged, *CNR* increased significantly for both 3T ($p = 0.00006$) and 7T ($p = 0.00007$)—however, at both field strengths, we found the smallest gain in *CNR* when comparing 6 *vs* 8 breath-holds averaged (Figure 3B). *CNR* and GM-to-WM contrast were significantly higher at 7T relative to 3T ($p_{CNR} = 0.0012$; $p_{GM\text{-}to\text{-}WM} = 0.00075$) (Table 1). In fact, 1 min of scan time at 7T resulted in the same *CNR* as 2 min 50 s of scan time at 3T and the same GM-to-WM contrast as 4 min 50 s of scan time at 3T. Of note, the subject-wise regression of *CNR* values between field strengths (Figure 3C), which provides some insight into subject repeatability, was very high ($R^2 = 0.7$)—according to the regression slope, *CNR* values were 1.45 times higher at 7T relative to 3T.

Novel Determination of an AIF for bhDSC

A prerequisite for DSC MRI is the presence of signal change upstream of tissue (typically at the arterial level), which is then used as an input function (i.e., AIF) for perfusion quantification in the tissue and veins. In Gd- or hypoxia-based methods, arterial signal change results from an increase in paramagnetic contrast agent in the artery, which subsequently passes through to the tissue (Rosen *et al.*, 1990; Zierler, 1962). However, given that blood oxygenation in healthy subjects is typically fully saturated in the major cerebral arteries, an increase in blood flow induced by CO_2 following breath-holding is not expected to significantly change the amount of dOHb in the arteries (Jain *et al.*, 2011; Severinghaus, 1979; Vestergaard and Larsson, 2019). Consequently, arterial magnetic susceptibility is not expected to change much as a result of breath-holding. However, hypercapnia uniquely results in **vasodilation** (Grubb Jr *et al.*, 1974)—

162 therefore, while MRI signal is not expected to change in the arteries from paramagnetic contrast agent, we
163 expect that arterial vasodilation may result in a signal decrease, particularly at higher magnetic field
164 strength (see Discussion). Thus, we developed a novel, alternative framework for selecting the AIF when
165 using breath-holds (or hypercapnia in general).
166

Figure 4. The AIF. A. ΔR_2* *AUC* maps were calculated subject-wise, transformed to MNI152 2 mm anatomical space,
and averaged at both 3T and 7T (sagittal and axial views are shown at the level of the middle cerebral artery). Pink
arrows indicate an example of the AIF's location. *AUC* maps are overlaid onto the MNI152 2 mm anatomical
template. **B.** The associated AIF ΔR_2* time courses are shown at 3T and 7T, with subject-wise time courses in gray and
subject-averaged time courses in pink. **C.** Illustration of vasodilation mechanism which results in arterial signal loss
during breath-holding.

Figure 4 shows voxels exhibiting a negative signal change (i.e., a positive relaxation change), and
following averaging of these voxels in the middle cerebral artery (MCA), displays the subject-averaged
AIF time courses at 3T and 7T in response to breath-holding. Voxels with a positive relaxation rate
change are located adjacent to the ventricles and in regions containing/adjacent to veins or arteries—there
are approximately two times more voxels fitting this criterion at 7T (Figure 4A), and the magnitude of the
associated relaxation rate changes are significantly higher ($p < 0.0007$) in comparison to 3T. The presence
of voxels displaying a positive relaxation rate change is, at first glance, counterintuitive given that the
predominant effect of breath-holding is a decrease in dOHb (i.e., decrease in the relaxation rate).
However, when we simulated (Figure S2) a voxel containing a large vessel, with little-to-no change in
dOHb in combination with vasodilation, a substantial relaxation rate increase was observed, in
agreement with the experimental results. Thus, the simulations and experimental findings support the
hypothesis that arterial vasodilation is responsible for the substantial relaxation rate increase observed in
the arteries, particularly at 7T, as a result of breath-holding.
188

189 *bhDSC Perfusion Quantification*
190

The AIF was first scaled by the integral of the VOF, composed of voxels in the superior sagittal sinus (SSS)—this step was conducted as, like in tissue, the VOF integral is representative of the amount of contrast agent (dOHb), whereas unlike in tissue, the AIF integral is representative of the degree of vasodilation. Following this scaling step, *CBV*, *CBF*, and *MTT* maps were calculated using a standard, truncated singular value decomposition (SVD) analysis, with an SVD noise threshold of 20% and hematocrit correction factor of 1.45 (Bjornerud and Emblem, 2010; Meier and Zierler, 1954; Østergaard *et al.,* 1996; Wirestam *et al.,* 2000; Schulman *et al.,* 2023). Note that although these measurements have been converted to physiological units, all reported perfusion values are still considered **relative**, in that they depend on numerous acquisition and analysis parameters (refer to Methods).

		ΔS (%)		CNR		CBV (mL/100g)		CBF (mL/100g/min)		MTT (s)
		GM	WM	GM	WM	GM	WM	GM	WM	GM
3T	Mean	**1.78**	**1.28**	**6.2**	**4.01**	**5.99**	**4.3**	**59.34**	**43.14**	**6.96**
	St.dev	*0.31*	*0.12*	*1.25*	*0.52*	*1.16*	*0.97*	*20.69*	*16.13*	*2.11*
	CV	0.17	0.09	0.20	0.13	0.19	0.23	0.35	0.37	0.30
7T	Mean	**2.2**	**1.27**	**8.52**	**4.8**	**5.2**	**3.3**	**44.29**	**25.48**	**7.36**
	St.dev	*0.41*	*0.23*	*1.87*	*0.94*	*2.15*	*1.39*	*10.05*	*6.05*	*2.52*
	CV	0.19	0.18	0.22	0.20	0.41	0.42	0.23	0.24	0.34

Table 1. Summary of bhDSC Perfusion Quantification Statistics. Average (mean), standard deviation (st.dev), and coefficient of variation (CV) values for 3T and 7T are shown.

Figure 5. bhDSC Perfusion Quantification Results. A. *CBV* (mL/100g) and *CBF* (mL/100g/min) maps were calculated subject-wise, transformed to MNI152 2 mm anatomical space, and averaged at both 3T and 7T (sagittal (x = 30) and axial views (z = 38, 46) are shown). **B.** Box and whisker plots show the calculated *CBV* and *CBF* (for GM and WM) at 3T and 7T. **C.** Voxel-wise regressions for *CBV* and *CBF* at 3T *vs* 7T. The regressions are based on voxels from the subject-averaged perfusion maps, and only voxels from the GM and WM are displayed (masks defined in Methods). Further masking was conducted to remove voxels at the periphery of the brain, where values differed substantially between 3T and 7T due to registration differences (i.e., in regions where the ratio of ΔS_{7T} to ΔS_{3T} was

above the 95th percentile). Regressions are constrained such that the line of best fit intersects with the origin. *m*
represents the regression slope.

Figure 5 shows the subject-averaged *CBV* and *CBF* perfusion maps (Figure 5A), subject-wise GM and WM
perfusion values (Figure 5B), and voxel-wise regressions for *CBV* and *CBF* between 3T and 7T (Figure
5C). At 3T, *CBV* values are slightly higher ($p = 0.108$) and *CBF* values are significantly higher ($p = 0.0027$)
than at 7T (Figure 5B). The attained GM-to-WM perfusion contrast (i.e., ratio of GM-to-WM perfusion
values) was significantly higher at 7T relative to 3T for both *CBF* ($p = 0.018$) and *CBV* ($p = 0.016$) values. In
addition, as shown in the subject-wise *CBF* maps (Figure S3), clear GM-to-WM contrast is observed in all
subjects at 7T, but not at 3T. A high coefficient of determination (voxel-wise $R^2 = 0.53$) is associated with
the voxel-wise linear regression of 3T and 7T *CBV* values—this is similarly the case for *CBF* (voxel-wise R^2
$= 0.58$) (Figure 5C). Of note, *MTT* values in GM (Figure S4) are slightly higher, although not significantly
($p = 0.74$), at 7T as opposed to 3T ($MTT_{7T,GM} = 7.36 \pm 2.52$ *vs* $MTT_{3T,GM} = 6.96 \pm 2.11$)—due to low *CNR*, *MTT*
values are not reported from the WM.

Delay Quantification

Finally, average bolus delay maps, relative to the AIF, were generated from the bhDSC data using a
model-independent approach. First, the subject-wise relaxation rate time courses were transformed to
MNI152 2 mm anatomical space and averaged across subjects. The average time courses were then
interpolated to a temporal resolution of 0.5 s. The AIF time courses from each subject were time-shifted
such that AIF bolus maximums were temporally aligned between subjects—these time courses were then
averaged. Finally, this subject-averaged AIF was shifted forward in 0.5 s steps, up to a maximum of 15 s,
and the correlation between the AIF and tissue time course at each temporal shift was recorded. The
temporal shift resulting in the highest correlation was determined to be the delay.

238
239 **Figure 6. bhDSC Delay Map at 7T.** Tissue delay map relative to the AIF is shown in MNI152 2 mm anatomical space
240 (sagittal and axial views are shown).

	GM	WM	VOF	Putamen	Occipital Lobe
Delay (s)	5.3	5.8	7.0	4.4	6.6

Table 2. bhDSC Delay at 7T. Calculated average delay values from various cerebral regions (note that the masks for
the occipital GM and putamen are from the MNI structural atlas (Mazziotta *et al.,* 2001)).

The averaged bhDSC bolus delay map at 7T and associated delay values across various cortical regions
are shown in Figure 6 and Table 2—results at 3T are also shown (Figure S5), although due to a lower
CNR, 3T results may not accurately capture the delay properties of the tissue. Within the GM, subcortical
regions such as the putamen yield a shorter delay (4.4 s) in comparison to average cortical GM (5.3 s),
while the occipital lobe (perfused by the posterior cerebral artery) resulted in a longer delay (6.6 s). The
perfusion output (VOF) of the brain was characterized by the longest observed delay (7 s).

Discussion

255 We have shown, for the first time, that DSC-based perfusion quantification is feasible using a breath-hold
256 task, without exogenous contrast or additional equipment. To that end, we acquired T_2*-weighted data at
257 3T and 7T from ten healthy subjects who performed short-duration breath-holds (16 s each separated by
258 44 s of rest) and then analyzed the data using a DSC MRI framework to quantify perfusion. We have
259 found the following:

- The AIF during a breath-hold is characterized by a negative signal change, likely representing vasodilation, as supported by both simulations and the literature
- bhDSC-calculated *CBV* and *CBF* were generally within the physiological range of values reported in the literature using established perfusion MRI approaches and demonstrated strong voxel-wise linear agreement between 3T and 7T
- While all subjects at 7T had observable GM-to-WM perfusion contrast, this was not the case at 3T
- At 7T, the breath-hold task yielded significantly higher *CNR* ($p = 0.0012$) and GM-to-WM contrast ($p = 0.00075$) relative to 3T
- *CNR* gains roughly plateau following approximately 5-to-6 min of scan time (averaging 5-6 breath-holds) at both 3T and 7T
- Bolus delay maps showed a clear delineation between various cortical regions, and the distribution of delay values across these territories agreed with those reported in the literature

From Breath-Hold to Signal Change

It is well established that hypercapnia induced by breath-holding produces a quantifiable T_2* signal
change in the tissue, which has been particularly useful in CVR and calibrated BOLD studies (Bright *et al.,*
2011; Bright and Murphy, 2013; Bulte *et al.,* 2009). Of note, mild hypoxia is also known to accompany
hypercapnia during breath-holding—but, given a roughly 0.65% reduction in arterial oxygen saturation
during a 16 s breath-hold (Sasse *et al.,* 1996; Severinghaus, 1979), the impact of breath-hold-induced
hypoxia on T_2* signal change is expected to be negligible in comparison to the effect attributed to
hypercapnia.

It has long been known that increased P_aCO_2 (arterial partial pressure of CO_2) leads to increased blood
flow in the arteries, and ultimately, in tissue capillaries and veins (Kety and Schmidt, 1948). This occurs
either directly through CO_2 acting on the smooth muscle cells surrounding the arteries/arterioles, or
indirectly through CO_2 acting on the vascular endothelium, in both cases resulting in vasodilation and
increased blood flow (Liu *et al.,* 2019). Therefore, a breath-hold which yields a transient rise and fall (i.e.,
bolus) of P_aCO_2 is a relatively simple way to induce a transient bolus of *CBF*. Of note, the breath-hold
duration has a roughly sigmoidal relationship with the resulting change in P_aCO_2 (Sasse et al., 1996),
which then also has a roughly sigmoidal relationship with the resulting change in *CBF* (Bhogal *et al.,*
2014)—however, the short duration breath-hold in our study occupies the linear range within this

sigmoidal relationship. Given that oxygen metabolism is assumed to be constant during mild
hypercapnia (Blockley *et al.,* 2013; Chen and Pike, 2010; Jain *et al.,* 2011; Vestergaard and Laarson, 2019,
Zappe *et al.,* 2008), the *CBF* bolus resulting from arterial/arteriolar/capillary vasodilation now permits the
cerebral tissue to be supplied with more oxygen than it requires, leading to a transient surplus in oxygen,
and thus, an increase in blood oxygenation and a reduction in dOHb, roughly in accordance with the
bolus shape and magnitude of P_aCO_2 in tissue.

We refer to what has just been described as the CVR hypothesis, wherein the observed arterial, tissue,
and venous signal changes are a result of CO_2-induced vasodilation throughout the vascular tree. That is,
the increased CO_2 following breath-holding leads to an increase in arterial blood volume, in turn
resulting in increased flow. In this hypothesis, it is predominantly brain-originating CO_2-induced
vasodilation/flow increases prior to the tissue which result in an oxygenation bolus at the tissue level.
However, a second mechanism—the cardiac hypothesis—may alternatively describe the observed
oxygenation bolus. Here, the previously described CVR effect may occur in addition to a cardiac output
increase which propagates throughout the cerebral vasculature as a bolus of increased blood flow (in
accordance with the breath-hold duration). According to the literature, the likelihood of a cardiac
hypothesis is very low given that cardiac output can increase, decrease, or stay constant depending on
the breath-hold maneuver (Paulev and Wetterqvist, 1968). In fact, Sakuma *et al.* found that large lung
volume breath-holding (end-inspiration) results in a reduction of cardiac output—this would
theoretically yield the reverse effect of that observed in our data (Sakuma *et al.,* 2001).

Regardless, both the CVR and cardiac hypotheses predict a change in paramagnetic dOHb concentration,
which is the basis for T_2* signal change in fMRI (Bandettini *et al.,* 1992; Kwong *et al.,* 1992; Ogawa *et al.,*
1990; Ogawa *et al.,* 1992). The magnitude of these signal changes reflect the underlying tissue *CBV,*
allowing for quantification within a DSC framework. In the current study, we observed a roughly 10-15 s
delay between the initiation of hypercapnia and the resulting T_2* signal change bolus onset (Figure 1),
which is similarly observed in previous hypercapnia studies and is attributed to the time for CO_2 build-
up in the lungs and the lung-to-brain travel time of blood (Blockley *et al.,* 2010; Thomas *et* al., 2014), or in
the case of the cardiac hypothesis, the delay between cardiac output initiation and propagation to the
cerebral tissue. Ultimately, the decrease in dOHb results in a transient increase in MRI signal, and the
magnitude of the signal change is dependent on the magnitude of ΔP_aCO_2, field strength, pulse sequence,
and tissue composition (Uludag *et al.,* 2009).

Determination of an AIF for bhDSC

To quantify *CBV, CBF,* and *MTT* using DSC MRI, the tissue bolus must be deconvolved with an input
bolus, ideally from the artery (AIF), that acquires dispersion and delay as it travels to and through the
tissue, in accordance with indicator dilution theory (Rosen *et al.,* 1990; Zierler, 1962). While an AIF is
expected and routinely determined when using Gd or hypoxia contrast, the presence of an AIF and its
associated properties are unknown in bhDSC—to address this, we examined the physiological processes
underlying arterial signal change as a result of breath-holding.

As argued above, dOHb levels in healthy subjects do not change much in the arteries during breath-
holding as there is effectively no oxygen exchange at this level of the vasculature and arterial blood
oxygenation is already near-saturation (Jain *et al.,* 2011; Vestergaard and Larsson, 2019). Thus, given that
most voxels in a typical T_2*-weighted MRI acquisition (2-4 mm voxel dimensions) likely consist of more
than one tissue/vessel type, any observed signal increase in and around the MCA likely represents signal

changes from surrounding, early-perfused cortical tissue and nearby veins—using this as an AIF would
violate basic tenets of indicator dilution theory.
341
Experimentally, we noticed that the MRI signal, which increased throughout most of the brain in
response to a breath-hold, decreased in many voxels containing large arteries and/or veins, particularly
when adjacent to CSF, with a larger observed effect at 7T (Figure 4). To better understand this
phenomenon, we modeled (Figure S2) an arterial voxel with vasodilation (i.e., *CBV* change) but little to
no dOHb change—in doing so, we found that vasodilation resulted in a sizeable signal decrease (i.e.,
relaxation rate increase) in accordance with our experimental findings, particularly at 7T. The reason for
this phenomenon relies on the fact that the intravascular transverse relaxation rate is substantially higher
than the extravascular tissue and CSF relaxation rates at 7T, and to a much lesser extent, at 3T where the
intravascular transverse relaxation rate is higher than CSF relaxation (Uludag, 2009). Thus, an increase in
CBV with little-to-no change in blood oxygenation leads to an MRI signal decrease (Figure 4C). This
signal origin has also been hypothesized as an explanation for the initial signal dip observed in fMRI
studies (Uludag, 2010) and has been posited in other work (Jin and Kim, 2010). Although its magnitude
depends on vessel orientation and voxel composition, measurable signal change is expected from our
simulations to occur in any voxel where a relatively large vessel increases in volume with negligible
change in blood oxygenation, displacing extravascular tissue and/or CSF, particularly at 7T. Of note, this
means that tissue voxels co-localized with larger vessels and CSF may display **both** an initial signal
decrease from vasodilation and a subsequent signal increase from the resulting dOHb decrease.
359
Our findings are also supported by the literature, where negative signal change during hypercapnia was
found in and adjacent to the ventricles (Blockley *et al.*, 2011; Thomas *et al.*, 2013). At 3T, this effect results
almost entirely from the difference between CSF and intravascular signal, as previously described
(Thomas *et al.*, 2013). Voxels characterized by a signal decrease were observed in higher abundance at the
arterial and venous level in a previous fMRI study at 7T (Bianciardi *et al.*, 2011), as is similarly observed
in our study.
366
In sum, there is very strong experimental and theoretical evidence that the positive relaxation rate change
at the arterial level represents a time course of arterial vasodilation in direct response to changes in
P_aCO_2—this phenomenon is not expected in Gd or hypoxia studies, where relaxation rate is known to
increase in the artery due to increased paramagnetic contrast agent, and not vasodilation. Although it is
not the traditional input of contrast agent-induced signal change, arterial vasodilation does in fact
represent the input induced by hypercapnia, essentially reflecting the input P_aCO_2 time course in the
brain, which then passes through and acts on downstream vasculature to yield tissue signal change in
accordance with the shape and magnitude of the P_aCO_2 bolus. The careful selection and averaging of
voxels in and around the MCA exhibiting this positive relaxation rate change provided us with a novel
solution for deconvolving the tissue data with an input function. The fact that arterial signal change is
opposite in sign to signal change observed in the majority of the brain will likely make it easier to identify
an AIF using bhDSC in comparison to traditional DSC techniques.
379
380 *Bolstering CNR in bhDSC*
381
There is a significant increase in *AUC, CNR,* and GM-to-WM contrast as field strength increases (Figures
2 and 3; Table 1). These findings are supported by our simulations (Figure S2) and studies that have
demonstrated a higher transverse relaxivity of dOHb as a function of field strength (Blockley *et al.*, 2008;
Kjolby *et al.*, 2009; Uludag *et al.*, 2009)—thus, for the same breath-hold duration and performance, higher

signal change will be observed at 7T relative to 3T. This supports the notion that a bhDSC analysis will
yield more robust and, hence, clinically valuable results at higher field strength.
388
Although there is also a significant increase in *CNR* as the number of breath-holds averaged increases, we
found the smallest *CNR* and GM-to-WM contrast gained when comparing 6 *vs* 8 breath-holds averaged
(Figures 3A and 3B). This suggests a saturation/plateau in *CNR* after approximately 5-to-6 min of scan
time. Thus, in roughly the same amount of time that it takes to acquire a 2 mm isotropic arterial spin
labeling (ASL) scan at 7T, the *CNR* is roughly 5-6 times higher when using the bhDSC technique (Ivanov
et al., 2017). Even a single breath-hold at 7T—approximately 1 minute of scan time—provides roughly 2
times the *CNR* relative to previous studies that used gas control system-induced hypoxia at 3T (Poublanc
et al., 2021; Sayin *et al.,* 2023), again demonstrating the supremacy of ultra-high field strength. Although
CNR using bhDSC is lower than when using Gd (Schulman *et al.,* 2023), the increased *CNR* of our
technique relative to current non-invasive approaches, including ASL, is noteworthy.
399
bhDSC Relative Perfusion Quantification
401
The subject-averaged *CBV* and *CBF* values and maps (Figures 4 and S3; Table 1) show sufficient GM-to-
WM contrast and are generally within the documented range of previously reported values (CBV_{GM} = 3-11
mL/100g; CBF_{GM} = 52-137 mL/100g/min) (Bjornerud and Emblem, 2010; Carroll *et al.,* 2008; Knutsson *et al.,*
2010). While high GM-to-WM contrast is observed in all individual subject maps at 7T, this is not always
the case at 3T (Figure S3), suggesting that bhDSC will consistently yield perfusion maps that are more
clinically valuable at higher field strength.
408
The discrepancy between 3T and 7T values supports the notion that DSC quantification is **relative**, with a
dependance on field strength, amongst many other parameters. The high-level explanation for the *CBV*
and *CBF* quantification discrepancy is that tissue *AUC* increases less than VOF *AUC* as field strength
increases (Figures 2 and S2). Specifically, we found that AUC_{GM} roughly doubles while the average
AUC_{VOF} nearly triples as field strength increases from 3T to 7T, leading to lower calculated values of *CBV*
and *CBF* at higher field strength—this is also supported by our simulations. This scaling difference likely
results from the non-linear intravascular contribution, which is a larger signal component at lower field
strength (approximately 70-90% of the signal at 3T in comparison to roughly 0-10% of the signal at 7T in
accordance with Uludag *et al.,* 2009). To reiterate, as there is no commonly accepted pipeline for absolute
DSC MRI quantification due to the many quantification dependencies that exist, all calculated perfusion
values are considered relative. Regardless, there is a strong voxel-wise linear agreement/regression
between 3T and 7T tissue perfusion values (Figure 5C), indicating that there are no significant regional
scaling differences between field strengths.
422
As previously mentioned, we do not consider *MTT* values in the WM to be reliable in bhDSC due to the
low *CNR* in combination with the fact that *MTT* is calculated using already-noisy blood volume and flow
information—thus the following discussion will focus on the GM. The MTT_{GM} values (Figure S4; Table 1)
are twice as high as values obtained in studies using Gd contrast—however, the values in this study are
similar to, if not lower than, those reported in studies using hypoxia contrast (Bjornerud and Emblem,
2010; Knutsson *et al.,* 2010; Schulman *et al.,* 2023; Vu *et al.,* 2021). Once again, this discrepancy likely
results from another known relative DSC quantification dependency—namely, in previous work, we
found longer bolus durations to result in higher calculated *MTT* values (Schulman *et al.,* 2023). A breath-
hold of 16 s induces a roughly 35 s signal bolus, which is substantially longer than a Gd bolus (but similar
in duration to a typical hypoxia bolus), and consequently yields a higher calculated *MTT* due to the

inherent limitations of SVD. Although slightly higher at 7T, *MTT* quantification is not statistically
different between field strengths. Of note, in accordance with the central volume principle (Meier and
Zierler, 1954), the higher *MTT* values explain the slight underestimation of *CBF,* particularly at 7T, in
comparison to average values reported in the literature (Bjornerud and Emblem, 2010; Carroll *et al.,* 2008;
Knutsson *et al.,* 2010).

Finally, the delay maps are useful for addressing whether the breath-hold induces a bolus that traverses
the brain in a manner dependent on the vascular topography, acquiring delay/dispersion as it progresses.
According to our delay maps (Figures 6 and S5; Table 2), the bhDSC method yields delay values across
the brain in accordance with the ordering of bolus arrival times reported in the literature when using Gd
as a contrast agent (Ibaraki *et al.,* 2007) and breath-holding for CVR analysis (Zvolnek *et al.,* 2023). This
provides strong evidence that bhDSC involves a bolus whose delay and dispersion properties are
dependent on the location within the cerebral vascular network where tissue is perfused. However, our
7T delay values are globally higher than those reported in Ibaraki *et al.* by approximately 4.5 s. This
discrepancy is not entirely clear, but may be partially owed to the fact that our input function, reflecting
arterial vasodilation, is a 'pure' input function, whereas that used in Ibaraki *et al.* likely suffers
substantially from the partial volume effect given their reported 5 mm slice resolution, leading to a
shorter measured delay as a result.

Clinical and Physiological Considerations for bhDSC

There are many benefits to the clinical implementation of bhDSC, including the lack of exogenous
contrast, additional medical device equipment, and the limitations therein (i.e., patient discomfort,
contrast agent extraversion, medical device maintenance, and additional costs) (Calamante, 2013;
Poublanc *et al.,* 2021). There are also notable benefits in comparison to ASL, such as bolstered *CNR,* and
the ability to acquire multiple perfusion parameters without the necessity of scanner-restrictive multi-
delay acquisitions. However, it is important to consider whether there are physiological and/or physical
processes, unique to breath-holding, which limit bhDSC's clinical utility in comparison to DSC acquired
with Gd or hypoxia.

In a standard DSC analysis (**Østergaard *et al.,* 1996**), it is assumed that *CBV* does not change, a condition
which is fulfilled during mild hypoxia or Gd. In bhDSC, signal change induced by a dynamic
vasodilation time course in tissue may confound the calculated perfusion metrics. To investigate, we
simulated (see Supplementary Materials) a 16 s breath-hold in addition to tissue vessel vasodilation and
found moderate, global *CBV* and *CBF* underestimation (Figure S6). We found the underestimation to be
almost identical in magnitude for voxels with differing vascular compositions and *CBV* values—thus,
relative values are expected to be preserved across the brain. On the other hand, given the presence of
other global quantification errors in DSC, global scaling underestimation from vasodilation is not
expected to hinder the utility of this method, particularly if a global correction factor is introduced. Thus,
from our simulations, vasodilation in the tissue is not expected to confound bhDSC perfusion
calculations. However, vasodilation from arterial and venous pial vessels co-localized with CSF and
tissue would likely result in more substantial inaccuracies to *CBV* and *CBF* estimation.

Another consideration is the potential variability in the oxygen metabolism rate (CMRO$_2$) during breath-
holding, as this would additionally violate an important DSC analysis assumption. There is very
compelling evidence that shows a lack of significant CMRO$_2$ variability during mild hypercapnia or

breath-holding, especially for the short, 16 s breath-hold duration employed in our study (Blockley *et al.,*
2013; Chen and Pike, 2010; Jain *et al.,* 2011; Vestergaard and Laarson, 2019; Zappe *et al.,* 2008).

Causality of flow in the vascular network is another important consideration. In healthy subjects, the
cerebral vasculature is assumed to respond homogeneously to a vasodilatory stimulus (Bhogal *et al.,* 2014;
Fisher and Mikulis, 2021). However, in certain patients (i.e., those with steno-occlusive disease and/or
vascular steal physiology), the affected arteries may have reduced vasodilatory capacity, which would
result in tissue downstream from the affected vessels yielding reduced calculated *CBV* values in
comparison to tissue supplied by other vessels (and potentially negative values in the case of vascular
steal (Poublanc *et al.,* 2013)), even though the ground truth blood volume values may not be reduced. In
these patients, apparent hypoperfusion relative to the rest of the brain may be a result of combined
hypoperfusion and reduced vasodilatory capacity in the supplying vasculature, constraining bhDSC's
validity in this cohort.

Finally, unlike Gd- and hypoxia-based methods, the signal change in bhDSC reflects an oxygenation
bolus that only begins at the tissue level—thus, signal change in bhDSC is capillary/venous dominated.
When comparing bhDSC with Gd-DSC perfusion results, there will likely be some incongruency in brain
regions which have substantially more arterial blood—however, this difference also creates an
opportunity to combine information from multiple methods to isolate different blood pools.

Limitations and Future Directions

As previously mentioned, bhDSC has reduced *CNR* in comparison to when using Gd (by approximately
5 times at 3T). This stems from the fact that dOHb has a far lower molar susceptibility than Gd (Schulman
et al., 2023) and ΔdOHb is relatively small during breath-holding (Jain *et al.,* 2011; Sasse *et al.,* 1996).
However, as illustrated in this work, both the accuracy and precision of perfusion quantification with
bhDSC is still sufficient on an individual subject level at 7T and on a group level at 3T. Also,
physiological and physical noise correction methods, such as those based on multi-echo T_2^* acquisition
(Caballero-Gaudes and Reynolds, 2017), may enable bhDSC to yield robust perfusion metrics for
individual subjects at lower field strengths.

The breath-hold itself confers some limitations to the proposed method. Patient compliance during the
repeated breath-holds of 16 s may be a somewhat difficult task for certain subjects, particularly the
elderly and those with neurological disease (Jahanian *et al.,* 2017; Spano *et al.,* 2013). However, breath-
holds are often applied in patients for body imaging, and as previously described, sufficient contrast can
be obtained at 7T with fewer breath-holds, limiting the duration required for patient compliance. Another
potential limitation is that motor activity associated with breath-holding can result in detectable motion
during the scan—we found this to be minimal and easily corrected for using motion correction software
in our study. Finally, there is inter-subject variability in the breath-holds, which is reflected in the
variability of bolus shape and magnitude between subjects (Figure S1). The variability (i.e., coefficient of
variation (CV)) of our calculated perfusion metrics (Table 1) is on par with that observed in a previous
study (Vu *et al.,* 2021) which used a gas control system for DSC quantification at 3T (*CBV*$_{CV}$ = 0.33; *CBF*$_{CV}$ =
0.42), limiting the concern that the contrast bolus needs to be standardized across subjects to ensure
precision of the calculated perfusion values. Regardless, precision should be enhanced in future work by
optimizing the breath-hold approach—providing more detailed instructions to the subjects, including
whether an end-inspiration or end-expiration is employed (Scouten and Schwarzbauer, 2008).

While we have compared literature values to our results, a direct subject-wise bhDSC *vs* Gd-DSC
comparison will allow for a stronger validation of the technique's utility in perfusion imaging. Although,
another group that induced hypercapnia with a gas control system to obtain DSC metrics found relatively
high *CBF* regional correlation with Gd-DSC (Vu *et al.*, 2023)—this is very promising given that our
technique relies on a similar physiological mechanism. As well, the clinical utility of the bhDSC technique
should be validated in different patient cohorts, to determine whether all or a subset of those with
vascular brain pathology will benefit from bhDSC alone or in combination with additional perfusion
imaging techniques (i.e., ASL). Finally, to provide more clinical centers with access to this method, it will
be prudent to assess the viability of and optimize the bhDSC technique at lower field strengths.

535
536

Conclusion

538
For the first time, we used breath-hold-induced T_2*-MRI signal changes to quantify perfusion using a
DSC analysis, at both 3T and 7T. This was made possible by developing and applying a novel arterial
input function strategy, to be used specifically in breath-hold and hyper/hypocapnic paradigms. In doing
so, we found that perfusion values (i.e., *CBF* and *CBV*) showed sufficient GM-to-WM contrast and were
within the range of physiological and literature-reported values. We hope that the methods and results
presented in this study will help pave the way toward contrast-free perfusion imaging, both clinically
and in research.

546
547

Methods

549
550 *Subjects*

551
552 This study was approved by the Research Ethics Board of Sungkyunkwan University and all procedures
553 followed the principles expressed in the Declaration of Helsinki. Informed consent was obtained in all 10
554 healthy volunteers (age: 30.4 ± 9.4 years, 3 female). Please note that the same subjects were scanned at
555 both 3T and 7T.

556
557 *MRI Sequences and Experimental Protocols*

558
559 MRI data were acquired at the Center for Neuroscience Imaging Research at Sungkyunkwan University
560 on the Siemens 3T Prisma (Siemens Healthineers, Erlangen, Germany) and Siemens 7T Terra (Siemens
561 Healthineers, Erlangen, Germany) using the commercially available 64 channel head/neck and 32
562 channel head coils (Nova Medical, Wilmington, USA).

563
564 *Structural MRI*
565 The 3T parameters were as follows: 3D-MP2RAGE (Marques *et al.*, 2010) with 1.0 mm isotropic spatial
566 resolution (176 sagittal slices; GRAPPA = 3; Ref lines PE = 32; FoVread = 250 mm; phase-encoding = A-P;
567 TI1/TI2 = 700/2500 ms; □1/□2 = 4°/5°; TE/TR = 2.98/5000 ms; bandwidth = 240 Hz/px; echo-spacing = 7.1
568 ms; TA = 8:22 min).

569
570 The 7T parameters were as follows: 3D-MP2RAGE (Marques *et al.* 2010) with 0.7 mm isotropic spatial
571 resolution (240 sagittal slices; GRAPPA = 3; Ref lines PE = 36; FoVread = 224 mm; phase-encoding = A-P;

572 TI1/TI2 = 1000/3200 ms; $\alpha1/\alpha2$ = 4°/4°; TE/TR = 2.29/4500 ms; partial-Fourier$_{slice}$ = 6/8; bandwidth = 200
573 Hz/px; echo-spacing = 7.3 ms; TA = 9:15 min).

574
Breath-hold MRI
The 3T parameters were as follows: Gradient-echo 2D-EPI (GRE-EPI) with 2.0 mm isotropic spatial
resolution (64 interleaved axial slices; GRAPPA = 2; Ref lines PE = 46; SMS = 2; Ref. scan = EPI; FatSat =
True; FoVread = 192 mm; phase-encoding = P-A; TE = 30 ms; α = 70°; TR = 2000 ms; bandwidth = 2312
Hz/px; echo-spacing = 0.53 ms; EPI factor = 94) was acquired for the breath-hold experiment (260 TRs,
TA = 8:53 min). 6 measurements of opposite phase-encoded (A-P) data were also acquired for
susceptibility distortion correction.

582
The 7T parameters were as follows: GRE-EPI with 2.0 mm isotropic spatial resolution (74 interleaved
axial slices; GRAPPA = 3; Ref lines PE = 54; SMS = 2; Ref. scan = EPI; partial-Fourier$_{phase}$ = 6/8; FatSat =
True; FoVread = 192 mm; phase-encoding = P-A; TE = 20 ms; α = 50°; TR = 2000 ms; bandwidth = 2368
Hz/px; echo-spacing = 0.53 ms; EPI factor = 96) was acquired for the breath-hold experiment (250 TRs,
TA = 8:46 min). 6 measurements of opposite phase-encoded (A-P) data were also acquired for
susceptibility distortion correction. Please note that the spatial resolution has **not** been optimized for 7T
in this study—we have chosen the same spatial resolution as at 3T to allow for easier comparison.

590
For the breath-hold scans, subjects were instructed to fixate on a black cross at the center of an iso-
luminant gray screen and a countdown timer which indicated when to breathe regularly and when to
perform a breath-hold. Specifically, each breath-hold block was composed of three sections: 10 s of
preparation with regular breathing, 16 s of breath-holding, and 34 s of regular breathing (Figure 1). This
was repeated nine times to obtain multiple boluses for averaging, with an added final baseline of 80 s.
Only the first eight boluses were used for each subject as the response to the final breath-hold bolus was
not fully included during the acquisition for some subjects—we highly recommend that quality control is
conducted here to select only a subset of the boluses for averaging, if necessary (i.e., if one or more
boluses (breath-holds) were not performed properly by the subject).

600
601 *Preprocessing*
602
603 FSL (version 6.0.4), AFNI (version 23.0.07), and in-house Python scripts (see Code Availability) were used
604 for image pre-processing (Cox, 1996; Jenkinson *et al.,* 2012). Anatomical and breath-hold T$_2$* data from
605 both scanners were corrected for gradient nonlinearities using the Human Connectome Project's version
606 of the gradunwarp tool (https://github.com/Washington-University/gradunwarp).

607
The MP2RAGE structural data were pre-processed using presurfer (https://github.com/srikash/presurfer)
and skull-stripped using SynthStrip (Hoopes *et al.,* 2022). The background denoised T1-weighted UNI
image was segmented using FSL FAST (Zhang *et al.,* 2001) to obtain three tissue classes corresponding to
gray matter (GM), white matter (WM), and cerebrospinal fluid (CSF). The breath-hold data then
underwent slice-timing (3dTshift, AFNI), motion (mcflirt, FSL), and distortion correction (topup, FSL)
(Andersson *et al.,* 2003; Jenkinson *et al.,* 2002). The pre-processed data were linearly detrended and
temporally filtered by averaging each signal time point with a 1x5 Gaussian kernel. The first eight breath-
hold blocks were then averaged, wherein each of the eight "boluses" was allotted a 72 s window centered
at the bolus maximum and averaged—an example of an original and averaged time course is shown in
Figure 1. The percentage maximal signal change (ΔS) and ΔS-to-noise ratio, hereafter termed contrast-to-
noise ratio (*CNR*), were then calculated from the bolus-averaged signal time course:

$$\Delta S = \frac{|S_{Max} - S_0|}{S_0} * 100 \quad (1)$$

$$CNR = \frac{|S_{Max} - S_0|}{\varepsilon_t} \quad (2)$$

622 In Eq. 1, S_0 is the average signal of ten temporal volumes before and after (roughly defined as the time
623 that the signal returns to the average pre-bolus baseline) the bolus and S_{Max} is the maximum signal
624 increase. The CNR is computed as shown in Eq. 2, where ε_t is the standard deviation of signal for ten total
625 temporal volumes before and after the selected bolus divided by S_0. GM-to-WM contrast ($GW_{contrast}$) was
626 also calculated as follows:

$$GW\,contrast = \frac{(CNR_{GM} - CNR_{WM})}{sCNR_{GM}} \quad (3)$$

629 Where CNR_{GM} and CNR_{WM} represent the mean GM and WM CNR, respectively, and $sCNR_{GM}$ represents
630 the standard deviation of CNR in GM. $S(t)$ was then converted to the change in transverse relaxation rate
631 time course ($\Delta R_2^*(t)$):

$$\Delta R_2^*(t) = -\left(\frac{1}{TE}\right) * ln\left(\frac{S(t)}{S_0}\right) \quad (4)$$

The above equations were applied to T_2*-weighted breath-hold data from all subjects at 3T and 7T to
generate ΔS, CNR, and $\Delta R_2^*(t)$ maps.

To facilitate group analyses, these maps, along with the later described perfusion maps, were
transformed to FSL's MNI152 2 mm space. The processed bhDSC data were first registered to the
subject's anatomical space (6 dof flirt, FSL) and the anatomical data were then non-linearly registered
(fnirt, FSL) to the FSL MNI152 2 mm template (Jenkinson and Smith, 2001; Jenkinson et al., 2012). The two
transformation matrices were then combined into a subject-specific native-to-MNI space warp matrix and
applied to all the subject-specific maps. The MNI space-transformed maps were then averaged across all
subjects to generate mean bhDSC$_{3T}$ and bhDSC$_{7T}$ perfusion maps. Data were also subjected to visual
quality control following each processing step.

Gray and White Matter Segmentation

GM and WM masks were generated, at both 3T and 7T, using whole-brain anatomical data (FSL FAST).
The partial volume estimate maps were then thresholded at 0.9, binarized, and transformed from
anatomical to GRE-EPI space for each subject (nearest-neighbor interpolation). Arterial and venous voxels
were removed from the masks by thresholding the standard deviation map (only voxels with values in
the lowest 10% of the ε_t range were kept in the mask) and using the output to mask the binarized FAST
segmentations—the assumption being that voxels with high CBV have higher physiological noise due to
pulsatility, leading to higher values of ε_t which can be removed from further analysis (Kashyap et al.,
2018; Schulman et al., 2023). The whole-brain GM and WM masks were then applied to each subject's
perfusion, ΔS, and CNR maps to obtain average GM and WM values (Table 1). GM and WM masks were
also generated in standard space from the MNI152 T_1 2 mm template using the procedure described
above.

Defining the AIF and VOF

In DSC MRI, it is critical to define an input function that is then used to normalize and deconvolve the
tissue data—this input is defined at the level of the artery and is consequently termed the arterial input
function (AIF) (Calamante *et al.*, 2013). Typically, to represent the AIF, 10-15 arterial voxels within or
adjacent to the M1 segment of the middle cerebral artery (MCA) are identified, with ΔS values above the
95th percentile, and zero delay (see delay quantification below). However, given the spatial resolution of
GRE-EPI (typically 2-4 mm in each dimension), signal change derived at the arterial level is likely
contaminated by signal changes from cortical tissue and nearby veins. With this in mind, we developed
an alternative, novel method to obtain an AIF from the breath-hold data, which can be used in any study
that uses hypercapnia to calculate perfusion at high or ultra-high field strength. In essence, the selection
criteria are the same as those described above, but voxels yielding **negative** Δ values, with magnitude
above the 95th percentile (S_{Max} is replaced with S_{Min} in Eq. 1), were used instead. We hypothesize that the
resulting signal time course reflects vasodilation at the arterial level in response to the CO_2 stimulus (see
Results and Discussion).

In DSC MRI, it is also useful to define an output function from the cerebral draining veins, whose
magnitude can be used to scale the measured AIF (Calamante *et al.*, 2013). Using both GRE-EPI and GRE-
EPI-registered anatomical data, 10-15 venous voxels within or adjacent to the superior sagittal sinus (SSS),
with ΔS values above the 99th percentile, long delay (~ 2-3 s), and low S_0 (as vein contains more dOHb
than artery, reducing the baseline GRE signal) were defined and averaged. This averaged venous time
course is hereafter termed the venous output function (VOF).

Perfusion Quantification

All voxel time courses were truncated to commence at the start of the AIF bolus and finish at the end of
the VOF bolus. The AIF was then scaled by the integral of the VOF using the following equation:

$$AIF_{Scaled}(t) = \frac{\int_0^\infty VOF(t)dt}{\int_0^\infty AIF(t)dt} * AIF(t) \quad (5)$$

This step was conducted as, like in tissue, the VOF integral is representative of the amount of contrast
agent (dOHb), whereas unlike in tissue, the AIF integral is representative of the degree of vasodilation.
Normalizing to venous signal also reduces the effect of various perfusion quantification dependencies
(Schulman *et al.*, 2023).

Cerebral blood volume (*CBV*), cerebral blood flow (*CBF*), and mean transit time (*MTT*) maps were then
calculated using a standard, truncated singular value decomposition (SVD) analysis, with an SVD noise
threshold of 20%, brain density factor of 1.05 mL/100g (introduces physiological units), and hematocrit
correction factor of 1.45 (Bjornerud and Emblem, 2010; Meier and Zierler, 1954; Østergaard *et al.*, 1996;
Wirestam *et al.*, 2000; Schulman *et al.*, 2023). Note that although reported in physiological units, these
perfusion values are **relative**, not absolute; research has shown perfusion imaging values to depend on
various contrast agent and acquisition parameters (Kiselev *et al.*, 2001; Schulman *et al.*, 2023). Regardless,
we still report values in physiological units to adhere to the standard DSC analysis pipeline and allow for
comparison with the literature. Also, note that all voxels yielding a negative *CBV* or *CBF* were removed
from subsequent analysis, and are not included in the summary results (Table 1).

Delay Quantification

Average bolus delay maps were also generated from the bhDSC data at 3T and 7T. Unlike the other perfusion metrics where DSC quantification was conducted in native space, the delay maps were calculated using the group average data.

First, the relaxation time courses were transformed to MNI152 2 mm anatomical space (using the previously described warp matrices) and averaged across subjects. The average time courses were then linearly interpolated to a temporal resolution of 0.5 s. The AIF time course from each subject was time-shifted to align with the timing of the median AIF peak onset across—these time courses were then averaged. The subject-averaged AIF was then shifted forward in 0.5 s steps, up to a maximum of 15 s, and the correlation between the AIF and tissue time courses for each voxel at each temporal shift was recorded. The temporal shift resulting in the highest correlation was determined to be the delay in a model-independent manner. A similar delay quantification method has been employed in previous studies using resting-state fMRI data (Khalil *et al.,* 2017; Lv *et al.,* 2013).

Note that the calculated delay maps reflect a combination of artery-to-tissue delay, tissue transit time, and dispersion, given that the AIF time courses were simply shifted to find the maximal correlation with each voxel time course (i.e., there was no separately modeled dispersion parameter in this analysis). The masks used for delay quantification in the occipital GM and putamen come from the MNI structural atlas (Mazziotta *et al.,* 2001).

Statistics

Parameters measured in native space have a calculated mean, standard deviation, and coefficient of variation (CV) for GM and WM, at both field strengths (Table 1). Note that CV is calculated as the inter-subject standard deviation divided by the parameter mean. Linear regressions were conducted for voxel-wise comparisons of *CBV, CBF,* and *MTT,* and subject-wise comparisons of *CNR*.

We found the data to be normally distributed for all measured parameters in GM and WM using the Shapiro-Wilk test. Thereafter, any statistical testing that was conducted for the experimental data (i.e., comparing mean GM and WM values at different field strengths) was done by applying a two-tailed paired student's t-test, with an alpha of 0.05.

Data and Code Availability

Anonymized subject data will be shared with any investigator for the purpose of ensuring reproducibility of the results presented in this work.

Analysis and simulation code will be **made available on GitHub and appended here** upon acceptance of the manuscript.

Conflict of Interest

The listed authors have no conflicts of interest to declare.

Author Contributions

J.B.S: Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Software, Visualization, and Writing

S.K: Data Curation, Software, and Writing

S.G.K: Funding Acquisition, Resources, Supervision, and Writing

K.U: Conceptualization, Formal Analysis, Funding Acquisition, Investigation, Methodology, Project Administration, Resources, Supervision, and Writing

Funding

The research conducted in this paper was supported by funding from the Canadian Institutes of Health Research (CIHR) to K.U. The study was also supported by the Institute for Basic Science, Suwon, Republic of Korea (IBS-R015-D1) to S.G.K.

Acknowledgments

We would like to thank Boohee Choi and Suji Jeong at the Institute for Basic Science for helping us acquire the data. Figures 1A and 4C were created with BioRender.com.

References

Andersson, J.L.R., Skare, S., Ashburner, J., 2003. How to correct susceptibility distortions in spin-echo echo-planar images: application to diffusion tensor imaging. Neuroimage 20, 870–888. https://doi.org/10.1016/S1053-8119(03)00336-7

Aslan, S., Xu, F., Wang, P.L., Uh, J., Yezhuvath, U.S., van Osch, M., Lu, H., 2010. Estimation of labeling efficiency in pseudocontinuous arterial spin labeling. Magn Reson Med 63, 765–771. https://doi.org/10.1002/mrm.22245

Bandettini, P.A., Wong, E.C., Hinks, R.S., Tikofsky, R.S., Hyde, J.S., 1992. Time course EPI of human brain function during task activation. Magnetic Resonance in Medicine 25, 390–397. https://doi.org/10.1002/mrm.1910250220

Bhogal, A.A., Siero, J.C.W., Fisher, J.A., Froeling, M., Luijten, P., Philippens, M., Hoogduin, H., 2014. Investigating the non-linearity of the BOLD cerebrovascular reactivity response to targeted hypo/hypercapnia at 7T. NeuroImage 98, 296–305. https://doi.org/10.1016/j.neuroimage.2014.05.006

Bianciardi, M., Fukunaga, M., van Gelderen, P., de Zwart, J.A., Duyn, J.H., 2011. Negative BOLD-fMRI signals in large cerebral veins. J Cereb Blood Flow Metab 31, 401–412. https://doi.org/10.1038/jcbfm.2010.164

Bjørnerud, A., Emblem, K.E., 2010. A fully automated method for quantitative cerebral hemodynamic

analysis using DSC–MRI. J Cereb Blood Flow Metab 30, 1066–1078. https://doi.org/10.1038/jcbfm.2010.4

Blockley, N.P., Jiang, L., Gardener, A.G., Ludman, C.N., Francis, S.T., Gowland, P.A., 2008. Field strength dependence of R1 and R relaxivities of human whole blood to prohance, vasovist, and deoxyhemoglobin. Magnetic Resonance in Medicine 60, 1313–1320. https://doi.org/10.1002/mrm.21792

Blockley, N.P., Driver, I.D., Francis, S.T., Fisher, J.A., Gowland, P.A., 2011. An improved method for acquiring cerebrovascular reactivity maps. Magn Reson Med 65, 1278–1286. https://doi.org/10.1002/mrm.22719

Blockley, N.P., Griffeth, V.E.M., Simon, A.B., Buxton, R.B., 2013. A review of calibrated blood oxygenation level-dependent (BOLD) methods for the measurement of task-induced changes in brain oxygen metabolism. NMR Biomed 26, 987–1003. https://doi.org/10.1002/nbm.2847

Bouten, J., Bourgois, J.G., Boone, J., 2020. Hold your breath: peripheral and cerebral oxygenation during dry static apnea. Eur J Appl Physiol 120, 2213–2222. https://doi.org/10.1007/s00421-020-04445-y

Bright, M.G., Donahue, M.J., Duyn, J.H., Jezzard, P., Bulte, D.P., 2011. The effect of basal vasodilation on hypercapnic and hypocapnic reactivity measured using magnetic resonance imaging. J Cereb Blood Flow Metab 31, 426–438. https://doi.org/10.1038/jcbfm.2010.187

Bright, M.G., Murphy, K., 2013. Reliable quantification of BOLD fMRI cerebrovascular reactivity despite poor breath-hold performance. Neuroimage 83, 559–568. https://doi.org/10.1016/j.neuroimage.2013.07.007

Bulte, D.P., Drescher, K., Jezzard, P., 2009. Comparison of hypercapnia-based calibration techniques for measurement of cerebral oxygen metabolism with MRI. Magn Reson Med 61, 391–398. https://doi.org/10.1002/mrm.21862

Bulte, D.P., Kelly, M., Germuska, M., Xie, J., Chappell, M.A., Okell, T.W., Bright, M.G., Jezzard, P., 2012. Quantitative measurement of cerebral physiology using respiratory-calibrated MRI. Neuroimage 60, 582–591. https://doi.org/10.1016/j.neuroimage.2011.12.017

Caballero-Gaudes, C., Reynolds, R.C., 2017. Methods for cleaning the BOLD fMRI signal. Neuroimage 154, 128–149. https://doi.org/10.1016/j.neuroimage.2016.12.018

Calamante, F., 2013. Arterial input function in perfusion MRI: a comprehensive review. Prog Nucl Magn Reson Spectrosc 74, 1–32. https://doi.org/10.1016/j.pnmrs.2013.04.002

Calamante F., Gadian DG., Connelly A., 2000. Delay and dispersion effects in dynamic susceptibility contrast MRI: simulations using singular value decomposition. Magn Reson Med 44(3), 466-73. https://doi.org/10.1002/1522-2594(200009)44:3<466::aid-mrm18>3.0.co;2-m.

Carroll, T.J., Horowitz, S., Shin, W., Mouannes, J., Sawlani, R., Ali, S., Raizer, J., Futterer, S., 2008. Quantification of Cerebral Perfusion Using the "Bookend Technique": an Evaluation in CNS Tumors. Magn Reson Imaging 26, 1352–1359. https://doi.org/10.1016/j.mri.2008.04.010

Cavuşoğlu, M., Pfeuffer, J., Uğurbil, K., Uludağ, K., 2009. Comparison of pulsed arterial spin labeling encoding schemes and absolute perfusion quantification. Magn Reson Imaging 27, 1039–1045. https://doi.org/10.1016/j.mri.2009.04.002

Chen, J.J., Pike, G.B., 2010. Global cerebral oxidative metabolism during hypercapnia and hypocapnia in humans: implications for BOLD fMRI. J Cereb Blood Flow Metab 30, 1094–1099. https://doi.org/10.1038/jcbfm.2010.42

Chen, J.J., Pike, G.B., 2010. MRI measurement of the BOLD-specific flow-volume relationship during hypercapnia and hypocapnia in humans. Neuroimage 53, 383–391. https://doi.org/10.1016/j.neuroimage.2010.07.003

Copen, W.A., Schaefer, P.W., Wu, O., 2011. MR Perfusion Imaging in Acute Ischemic Stroke. Neuroimaging Clin N Am 21, 259–283. https://doi.org/10.1016/j.nic.2011.02.007

Cox, R.W., 1996. AFNI: software for analysis and visualization of functional magnetic resonance

neuroimages. Comput Biomed Res 29, 162–173. https://doi.org/10.1006/cbmr.1996.0014

De Vis, J.B., Hendrikse, J., Bhogal, A., Adams, A., Kappelle, L.J., Petersen, E.T., 2015. Age-related changes
in brain hemodynamics; A calibrated MRI study. Hum Brain Mapp 36, 3973–3987.
https://doi.org/10.1002/hbm.22891

Fisher, J.A., Mikulis, D.J., 2021. Cerebrovascular Reactivity: Purpose, Optimizing Methods, and
Limitations to Interpretation – A Personal 20-Year Odyssey of (Re)searching. Frontiers in
Physiology 12.

Gauthier, C.J., Madjar, C., Desjardins-Crépeau, L., Bellec, P., Bherer, L., Hoge, R.D., 2013. Age
dependence of hemodynamic response characteristics in human functional magnetic resonance
imaging. Neurobiol Aging 34, 1469–1485. https://doi.org/10.1016/j.neurobiolaging.2012.11.002

Grubb, R.L., Raichle, M.E., Eichling, J.O., Ter-Pogossian, M.M., 1974. The effects of changes in PaCO2 on
cerebral blood volume, blood flow, and vascular mean transit time. Stroke 5, 630–639.
https://doi.org/10.1161/01.str.5.5.630

Harris, G.J., Lewis, R.F., Satlin, A., English, C.D., Scott, T.M., Yurgelun-Todd, D.A., Renshaw, P.F., 1998.
Dynamic Susceptibility Contrast MR Imaging of Regional Cerebral Blood Volume in Alzheimer
Disease: A Promising Alternative to Nuclear Medicine 6.

Hoopes, A., Mora, J.S., Dalca, A.V., Fischl, B., Hoffmann, M., 2022. SynthStrip: skull-stripping for any
brain image. NeuroImage 260, 119474. https://doi.org/10.1016/j.neuroimage.2022.119474

Hornak, J., 2010. The basics of MRI.

Ibaraki, M., Ito, H., Shimosegawa, E., Toyoshima, H., Ishigame, K., Takahashi, K., Kanno, I., Miura, S.,
2007. Cerebral Vascular Mean Transit Time in Healthy Humans: A Comparative Study with PET
and Dynamic Susceptibility Contrast-Enhanced MRI. J Cereb Blood Flow Metab 27, 404–413.
https://doi.org/10.1038/sj.jcbfm.9600337

Ito, H., Kanno, I., Ibaraki, M., Hatazawa, J., Miura, S., 2003. Changes in Human Cerebral Blood Flow and
Cerebral Blood Volume during Hypercapnia and Hypocapnia Measured by Positron Emission
Tomography. J Cereb Blood Flow Metab 23, 665–670.
https://doi.org/10.1097/01.WCB.0000067721.64998.F5

Ivanov, D., Gardumi, A., Haast, R.A.M., Pfeuffer, J., Poser, B.A., Uludağ, K., 2017. Comparison of 3T and
7T ASL techniques for concurrent functional perfusion and BOLD studies. Neuroimage 156, 363–376.
https://doi.org/10.1016/j.neuroimage.2017.05.038

Jahanian, H., Christen, T., Moseley, M.E., Pajewski, N.M., Wright, C.B., Tamura, M.K., Zaharchuk, G.,
2017. Measuring vascular reactivity with resting-state blood oxygenation level-dependent
(BOLD) signal fluctuations: A potential alternative to the breath-holding challenge? J Cereb Blood
Flow Metab 37, 2526–2538. https://doi.org/10.1177/0271678X16670921

Jahng, G.-H., Li, K.-L., Ostergaard, L., Calamante, F., 2014. Perfusion Magnetic Resonance Imaging: A
Comprehensive Update on Principles and Techniques. Korean J Radiol 15, 554–577.
https://doi.org/10.3348/kjr.2014.15.5.554

Jain, V., Langham, M.C., Floyd, T.F., Jain, G., Magland, J.F., Wehrli, F.W., 2011. Rapid magnetic resonance
measurement of global cerebral metabolic rate of oxygen consumption in humans during rest and
hypercapnia. J Cereb Blood Flow Metab 31, 1504–1512. https://doi.org/10.1038/jcbfm.2011.34

Jenkinson, M., Bannister, P., Brady, M., Smith, S., 2002. Improved optimization for the robust and
accurate linear registration and motion correction of brain images. Neuroimage 17, 825–841.
https://doi.org/10.1016/s1053-8119(02)91132-8

Jenkinson, M., Beckmann, C.F., Behrens, T.E.J., Woolrich, M.W., Smith, S.M., 2012. FSL. Neuroimage 62,
782–790. https://doi.org/10.1016/j.neuroimage.2011.09.015

Jenkinson, M., Smith, S., 2001. A global optimisation method for robust affine registration of brain
images. Med Image Anal 5, 143–156.

Jin, T., Kim, S.-G., 2010. Change of the cerebrospinal fluid volume during brain activation investigated by

T(1rho)-weighted fMRI. Neuroimage 51, 1378–1383.
https://doi.org/10.1016/j.neuroimage.2010.03.047

Kanda, T., Fukusato, T., Matsuda, M., Toyoda, K., Oba, H., Kotoku, J., Haruyama, T., Kitajima, K., Furui,
S., 2015. Gadolinium-based Contrast Agent Accumulates in the Brain Even in Subjects
without Severe Renal Dysfunction: Evaluation of Autopsy Brain Specimens with
Inductively Coupled Plasma Mass Spectroscopy. Radiology 276, 228–232.
https://doi.org/10.1148/radiol.2015142690

Kashyap, S., Ivanov, D., Havlicek, M., Poser, B.A., Uludağ, K., 2018. Impact of acquisition and analysis
strategies on cortical depth-dependent fMRI. NeuroImage, Neuroimaging with Ultra-high Field
MRI: Present and Future 168, 332–344. https://doi.org/10.1016/j.neuroimage.2017.05.022

Kety, S.S., Schmidt, C.F., 1948. THE EFFECTS OF ALTERED ARTERIAL TENSIONS OF CARBON
DIOXIDE AND OXYGEN ON CEREBRAL BLOOD FLOW AND CEREBRAL OXYGEN
CONSUMPTION OF NORMAL YOUNG MEN 1. J Clin Invest 27, 484–492.

Khalil, A.A., Ostwaldt, A.-C., Nierhaus, T., Ganeshan, R., Audebert, H.J., Villringer, K., Villringer, A.,
Fiebach, J.B., 2017. Relationship Between Changes in the Temporal Dynamics of the Blood-
Oxygen-Level-Dependent Signal and Hypoperfusion in Acute Ischemic Stroke. Stroke 48, 925–
931. https://doi.org/10.1161/STROKEAHA.116.015566

Kiselev, V. g., 2001. On the theoretical basis of perfusion measurements by dynamic susceptibility
contrast MRI. Magnetic Resonance in Medicine 46, 1113–1122. https://doi.org/10.1002/mrm.1307

Kjølby, B.F., Mikkelsen, I.K., Pedersen, M., Østergaard, L., Kiselev, V.G., 2009. Analysis of partial volume
effects on arterial input functions using gradient echo: a simulation study. Magn Reson Med 61,
1300–1309. https://doi.org/10.1002/mrm.21849

Knutsson, L., Ståhlberg, F., Wirestam, R., 2010. Absolute quantification of perfusion using dynamic
susceptibility contrast MRI: pitfalls and possibilities. Magn Reson Mater Phy 23, 1–21.
https://doi.org/10.1007/s10334-009-0190-2

Knutsson, L., Xu, X., Ståhlberg, F., Barker, P.B., Lind, E., Sundgren, P.C., van Zijl, P.C.M., Wirestam, R.,
2017. Dynamic Susceptibility Contrast MRI at 7 T: Tail-Scaling Analysis and Inferences About
Field Strength Dependence. Tomography 3, 74–78. https://doi.org/10.18383/j.tom.2017.00001

Kwong, K.K., Belliveau, J.W., Chesler, D.A., Goldberg, I.E., Weisskoff, R.M., Poncelet, B.P., Kennedy,
D.N., Hoppel, B.E., Cohen, M.S., Turner, R., 1992. Dynamic magnetic resonance imaging of
human brain activity during primary sensory stimulation. Proc Natl Acad Sci U S A 89, 5675–
5679.

Law, M., Young, R.J., Babb, J.S., Peccerelli, N., Chheang, S., Gruber, M.L., Miller, D.C., Golfinos, J.G.,
Zagzag, D., Johnson, G., 2008. Gliomas: predicting time to progression or survival with cerebral
blood volume measurements at dynamic susceptibility-weighted contrast-enhanced perfusion
MR imaging. Radiology 247, 490–498. https://doi.org/10.1148/radiol.2472070898

Lee, D., Le, T.T., Im, G.H., Kim, S.-G., 2022. Whole-brain perfusion mapping in mice by dynamic BOLD
MRI with transient hypoxia. J Cereb Blood Flow Metab 0271678X221117008.
https://doi.org/10.1177/0271678X221117008

Liu, P., De Vis, J.B., Lu, H., 2019. Cerebrovascular reactivity (CVR) MRI with CO2 challenge: A technical
review. Neuroimage 187, 104–115. https://doi.org/10.1016/j.neuroimage.2018.03.047

Lv, Y., Margulies, D.S., Cameron Craddock, R., Long, X., Winter, B., Gierhake, D., Endres, M., Villringer,
K., Fiebach, J., Villringer, A., 2013. Identifying the perfusion deficit in acute stroke with resting-
state functional magnetic resonance imaging. Ann Neurol 73, 136–140.
https://doi.org/10.1002/ana.23763

MacDonald ME, Berman AJL, Mazerolle EL, Williams RJ, Pike GB., 2018. Modeling hyperoxia-induced
BOLD signal dynamics to estimate cerebral blood flow, volume and mean transit time.
Neuroimage 178, 461-474. doi: 10.1016/j.neuroimage.2018.05.066.

Marques, J.P., Kober, T., Krueger, G., van der Zwaag, W., Van de Moortele, P.-F., Gruetter, R., 2010.
MP2RAGE, a self bias-field corrected sequence for improved segmentation and T1-mapping at
high field. Neuroimage 49, 1271–1281. https://doi.org/10.1016/j.neuroimage.2009.10.002
Mazziotta, J., Toga, A., Evans, A., Fox, P., Lancaster, J., Zilles, K., Woods, R., Paus, T., Simpson, G., Pike,
B., Holmes, C., Collins, L., Thompson, P., MacDonald, D., Iacoboni, M., Schormann, T., Amunts,
K., Palomero-Gallagher, N., Geyer, S., Parsons, L., Narr, K., Kabani, N., Le Goualher, G.,
Boomsma, D., Cannon, T., Kawashima, R., Mazoyer, B., 2001. A probabilistic atlas and reference
system for the human brain: International Consortium for Brain Mapping (ICBM). Philos Trans R
Soc Lond B Biol Sci 356, 1293–1322. https://doi.org/10.1098/rstb.2001.0915
Meier, P., Zierler, K.L., 1954. On the theory of the indicator-dilution method for measurement of blood
flow and volume. J Appl Physiol 6, 731–744. https://doi.org/10.1152/jappl.1954.6.12.731
Ogawa, S., Lee, T.M., Kay, A.R., Tank, D.W., 1990. Brain magnetic resonance imaging with contrast
dependent on blood oxygenation. Proc Natl Acad Sci U S A 87, 9868–9872.
Ogawa, S., Tank, D.W., Menon, R., Ellermann, J.M., Kim, S.G., Merkle, H., Ugurbil, K., 1992. Intrinsic
signal changes accompanying sensory stimulation: functional brain mapping with magnetic
resonance imaging. Proc Natl Acad Sci U S A 89, 5951–5955.
https://doi.org/10.1073/pnas.89.13.5951
Ostergaard, L., Weisskoff, R.M., Chesler, D.A., Gyldensted, C., Rosen, B.R., 1996. High resolution
measurement of cerebral blood flow using intravascular tracer bolus passages. Part I:
Mathematical approach and statistical analysis. Magn Reson Med 36, 715–725.
https://doi.org/10.1002/mrm.1910360510
Paulev, P., Wetterqvist, H., 1968. Cardiac Output during Breath-holding in Man. Scandinavian Journal of
Clinical and Laboratory Investigation 22, 115–123. https://doi.org/10.3109/00365516809160955
Poublanc, J., Han, J.S., Mandell, D.M., Conklin, J., Stainsby, J.A., Fisher, J.A., Mikulis, D.J., Crawley, A.P.,
2013. Vascular Steal Explains Early Paradoxical Blood Oxygen Level-Dependent Cerebrovascular
Response in Brain Regions with Delayed Arterial Transit Times. Cerebrovasc Dis Extra 3, 55–64.
https://doi.org/10.1159/000348841
Poublanc, J., Sobczyk, O., Shafi, R., Sayin, E.S., Schulman, J., Duffin, J., Uludag, K., Wood, J.C., Vu, C.,
Dharmakumar, R., Fisher, J.A., Mikulis, D.J., 2021. Perfusion MRI using endogenous
deoxyhemoglobin as a contrast agent: Preliminary data. Magnetic Resonance in Medicine 86(6),
3012-3021. https://doi.org/10.1002/mrm.28974
Qiao, P.-G., Han, C., Zuo, Z.-W., Wang, Y.-T., Pfeuffer, J., Duan, L., Qian, T., Li, G.-J., 2017. Clinical
assessment of cerebral hemodynamics in Moyamoya disease via multiple inversion time arterial
spin labeling and dynamic susceptibility contrast-magnetic resonance imaging: A comparative
study. Journal of Neuroradiology 44, 273–280. https://doi.org/10.1016/j.neurad.2016.12.006
Rogowska, J., Olkowska, E., Ratajczyk, W., Wolska, L., 2018. Gadolinium as a new emerging contaminant
of aquatic environments. Environ Toxicol Chem 37, 1523–1534. https://doi.org/10.1002/etc.4116
Rosen, B.R., Belliveau, J.W., Vevea, J.M., Brady, T.J., 1990. Perfusion imaging with NMR contrast agents.
Magn Reson Med 14, 249–265. https://doi.org/10.1002/mrm.1910140211
Sakuma, H., Kawada, N., Kubo, H., Nishide, Y., Takano, K., Kato, N., Takeda, K., 2001. Effect of breath
holding on blood flow measurement using fast velocity encoded cine MRI. Magnetic Resonance
in Medicine 45, 346–348. https://doi.org/10.1002/1522-2594(200102)45:2<346::AID-
MRM1044>3.0.CO;2-I
Sasse, S.A., Berry, R.B., Nguyen, T.K., Light, R.W., Mahutte, C.K., 1996. Arterial blood gas changes during
breath-holding from functional residual capacity. Chest 110, 958–964.
https://doi.org/10.1378/chest.110.4.958
Sayin, E.S., Schulman, J., Poublanc, J., Levine, H.T., Raghavan, L.V., Uludag, K., Duffin, J., Fisher, J.A.,
Mikulis, D.J., Sobczyk, O., 2023. Investigations of hypoxia-induced deoxyhemoglobin as a

contrast agent for cerebral perfusion imaging. Hum Brain Mapp 44, 1019–1029.
https://doi.org/10.1002/hbm.26131

Schlaudecker, J.D., Bernheisel, C.R., 2009. Gadolinium-associated nephrogenic systemic fibrosis. Am Fam
Physician 80, 711–714.

Schulman, J.B., Sayin, E.S., Poublanc, J., Manalac, A., Fisher, J.A., Sobczyk, O., Duffin, J., Levine, H.,
Mikulis, D.J., Uludag, K., 2022. Deoxyhemoglobin versus gadolinium as contrast in dynamic
susceptibility perfusion imaging: simulations and scan validations. Proc Intl Soc Mag Reson Med
30, 337.

Schulman, J.B., Sayin, E.S., Manalac, A., Poublanc, J., Sobczyk, O., Duffin, J., Fisher, J.A., Mikulis, D.,
Uludağ, K., 2023. DSC MRI in the human brain using deoxyhemoglobin and gadolinium—
Simulations and validations at 3T. Frontiers in Neuroimaging 2.

Scouten, A., Schwarzbauer, C., 2008. Paced respiration with end-expiration technique offers superior
BOLD signal repeatability for breath-hold studies. Neuroimage 43, 250–257.
https://doi.org/10.1016/j.neuroimage.2008.03.052

Severinghaus, J.W., 1979. Simple, accurate equations for human blood O2 dissociation computations. J
Appl Physiol Respir Environ Exerc Physiol 46, 599–602. https://doi.org/10.1152/jappl.1979.46.3.599

Spano, V.R., Mandell, D.M., Poublanc, J., Sam, K., Battisti-Charbonney, A., Pucci, O., Han, J.S., Crawley,
A.P., Fisher, J.A., Mikulis, D.J., 2013. CO2 blood oxygen level-dependent MR mapping of
cerebrovascular reserve in a clinical population: safety, tolerability, and technical feasibility.
Radiology 266, 592–598. https://doi.org/10.1148/radiol.12112795

Tanaka, Y., Nagaoka, T., Nair, G., Ohno, K., Duong, T.Q., 2011. Arterial spin labeling and dynamic
susceptibility contrast CBF MRI in postischemic hyperperfusion, hypercapnia, and after mannitol
injection. J Cereb Blood Flow Metab 31, 1403–1411. https://doi.org/10.1038/jcbfm.2010.228

Thomas, B.P., Liu, P., Aslan, S., King, K.S., van Osch, M.J.P., Lu, H., 2013. Physiologic underpinnings of
negative BOLD cerebrovascular reactivity in brain ventricles. Neuroimage 83, 505–512.
https://doi.org/10.1016/j.neuroimage.2013.07.005

Thomas, B.P., Liu, P., Park, D.C., van Osch, M.J., Lu, H., 2014. Cerebrovascular reactivity in the brain
white matter: magnitude, temporal characteristics, and age effects. J Cereb Blood Flow Metab 34,
242–247. https://doi.org/10.1038/jcbfm.2013.194

Uludağ, K., Müller-Bierl, B., Uğurbil, K., 2009. An integrative model for neuronal activity-induced signal
changes for gradient and spin echo functional imaging. NeuroImage 48, 150–165.
https://doi.org/10.1016/j.neuroimage.2009.05.051

Uludağ, K., 2010. To dip or not to dip: Reconciling optical imaging and fMRI data. Proceedings of the
National Academy of Sciences of the United States of America 107, E23.
https://doi.org/10.1073/pnas.0914194107

Vestergaard, M.B., Larsson, H.B., 2019. Cerebral metabolism and vascular reactivity during breath-hold
and hypoxic challenge in freedivers and healthy controls. J Cereb Blood Flow Metab 39, 834–848.
https://doi.org/10.1177/0271678X17737909

Vu, C., Chai, Y., Coloigner, J., Nederveen, A.J., Borzage, M., Bush, A., Wood, J.C., 2021. Quantitative
perfusion mapping with induced transient hypoxia using BOLD MRI. Magn Reson Med 85, 168–
181. https://doi.org/10.1002/mrm.28422

Vu, C., Xu, B., González-Zacarías, C., Shen, J., Baas, K., Choi, S., Nederveen, A., Wood, J., 2023. Sinusoidal
CO2 respiratory challenge for concurrent perfusion and cerebrovascular reactivity MRI. Frontiers
in Physiology 14. https://doi.org/10.3389/fphys.2023.1102983

Welker, K., Boxerman, J., Kalnin, A., Kaufmann, T., Shiroishi, M., Wintermark, M., American Society of
Functional Neuroradiology MR Perfusion Standards and Practice Subcommittee of the ASFNR
Clinical Practice Committee, 2015. ASFNR recommendations for clinical performance of MR
dynamic susceptibility contrast perfusion imaging of the brain. AJNR Am J Neuroradiol 36, E41-

51. https://doi.org/10.3174/ajnr.A4341

Wirestam, R., Andersson, L., Ostergaard, L., Bolling, M., Aunola, J.P., Lindgren, A., Geijer, B., Holtås, S.,
Ståhlberg, F., 2000. Assessment of regional cerebral blood flow by dynamic susceptibility contrast
MRI using different deconvolution techniques. Magn Reson Med 43, 691–700.
https://doi.org/10.1002/(sici)1522-2594(200005)43:5<691::aid-mrm11>3.0.co;2-b

Zappe, A.C., Uludağ, K., Oeltermann, A., Uğurbil, K., Logothetis, N.K., 2008. The influence of moderate
hypercapnia on neural activity in the anesthetized nonhuman primate. Cereb Cortex 18, 2666–
2673. https://doi.org/10.1093/cercor/bhn023

Zhang, Y., Brady, M., Smith, S., 2001. Segmentation of brain MR images through a hidden Markov
random field model and the expectation-maximization algorithm. IEEE Trans Med Imaging 20,
45–57. https://doi.org/10.1109/42.906424

Zierler, K.L., 1962. Theoretical Basis of Indicator-Dilution Methods For Measuring Flow and Volume.
Circulation Research 10, 393–407. https://doi.org/10.1161/01.RES.10.3.393

Zvolanek, K.M., Moia, S., Dean, J.N., Stickland, R.C., Caballero-Gaudes, C., Bright, M.G., 2023.
Comparing end-tidal CO2, respiration volume per time (RVT), and average gray matter signal for
mapping cerebrovascular reactivity amplitude and delay with breath-hold task BOLD fMRI.
Neuroimage 272, 120038. https://doi.org/10.1016/j.neuroimage.2023.120038

1052
1053

Figure Captions

1055

Figure 1. Summary of the Breath-hold Experiment. A. Description of the cerebrovascular reactivity phenomenon. At
baseline, tissue blood oxygenation is roughly between 60-80%. Increased CO_2 from breath-holding causes
vasodilation in the upstream arterial/arteriolar vasculature, leading to increased blood flow, and subsequently, a
decrease in the downstream tissue dOHb concentration (hyperoxia), resulting in an MRI signal increase. **B.** Signal
time course (blue) resulting from nine consecutive breath-holds (green) in a venous voxel. **C.** The signal time course
resulting from the temporal average of the first eight boluses was then used for perfusion quantification.

1062

1063 **Figure 2. Breath-Hold-Induced Relaxation Rate Time Course Dynamics.** Time courses were averaged across
1064 subjects for gray matter (GM), white matter (WM), vein (VOF), and artery (AIF) at 3T and 7T.

1065

Figure 3. Breath-hold *CNR* Properties. A. *CNR* maps were calculated (Eq. 2), transformed to MNI152 2 mm
anatomical space, and averaged across subjects for both 3T and 7T (axial view is slightly dorsal to the lateral
ventricles). For illustration purposes, *CNR* maps are shown when there is no bolus averaging or 2, 4, 6, or 8 boluses
averaged at 7T. **B.** Plot of subject-averaged *CNR*GM at 3T (dashed) and 7T (solid) as a function of the number of
breath-holds averaged (error bars represent the standard error). **C.** Linear regression of subject-wise GM (red) and
WM (blue) *CNR* values at 3T *vs* 7T. *m* represents the regression slope.

1072

Figure 4. The AIF. A. $\Delta R_2{}^*$ *AUC* maps were calculated subject-wise, transformed to MNI152 2 mm anatomical space,
and averaged at both 3T and 7T (sagittal and axial views are shown at the level of the middle cerebral artery). Pink
arrows indicate an example of the AIF's location. *AUC* maps are overlaid onto the MNI152 2 mm anatomical
template. **B.** The associated AIF $\Delta R_2{}^*$ time courses are shown at 3T and 7T, with subject-wise time courses in gray and
subject-averaged time courses in pink. **C.** Illustration of vasodilation mechanism which results in arterial signal loss
during breath-holding.

1079

Figure 5. bhDSC Perfusion Quantification Results. A. *CBV* (mL/100g) and *CBF* (mL/100g/min) maps were
calculated subject-wise, transformed to MNI152 2 mm anatomical space, and averaged at both 3T and 7T (sagittal (x =
30) and axial views (z = 38, 46) are shown). **B.** Box and whisker plots show the calculated *CBV* and *CBF* (for GM and
WM) at 3T and 7T. **C.** Voxel-wise regressions for *CBV* and *CBF* at 3T *vs* 7T. The regressions are based on voxels from
the subject-averaged perfusion maps, and only voxels from the GM and WM are displayed (masks defined in

Methods). Further masking was conducted to remove voxels at the periphery of the brain, where ΔS values differed
substantially between 3T and 7T due to registration differences (i.e., in regions where the ratio of ΔS_{7T} to ΔS_{3T} was
above the 95[th] percentile). Regressions are constrained such that the line of best fit intersects with the origin. m
represents the regression slope.
1089
1090 **Figure 6. bhDSC Delay Map at 7T.** Tissue delay map relative to the AIF is shown in MNI152 2 mm anatomical space
1091 (sagittal and axial views are shown).
1092
1093
1094 ## Tables
1095
1096 **Table 1. Summary of bhDSC Perfusion Quantification Statistics.** Average (mean), standard deviation (st.dev), and
1097 coefficient of variation (CV) values for 3T and 7T are shown.

		ΔS (%)		CNR		CBV (mL/100g)		CBF (mL/100g/min)		MTT (s)
		GM	WM	GM	WM	GM	WM	GM	WM	GM
3T	**Mean**	**1.78**	**1.28**	**6.2**	**4.01**	**5.99**	**4.3**	**59.34**	**43.14**	**6.96**
	St.dev	*0.31*	*0.12*	*1.25*	*0.52*	*1.16*	*0.97*	*20.69*	*16.13*	*2.11*
	CV	0.17	0.09	0.20	0.13	0.19	0.23	0.35	0.37	0.30
7T	**Mean**	**2.2**	**1.27**	**8.52**	**4.8**	**5.2**	**3.3**	**44.29**	**25.48**	**7.36**
	St.dev	*0.41*	*0.23*	*1.87*	*0.94*	*2.15*	*1.39*	*10.05*	*6.05*	*2.52*
	CV	0.19	0.18	0.22	0.20	0.41	0.42	0.23	0.24	0.34

1098
1099 **Table 2. bhDSC Delay at 7T.** Calculated average delay values from various cerebral regions (note that the masks for
1100 the occipital GM and putamen are from the MNI structural atlas (Mazziotta *et al.,* 2001)).

	GM	WM	VOF	Putamen	Occipital Lobe
Delay (s)	5.3	5.8	7.0	4.4	6.6

1101
1102