Identification of clade-defining single nucleotide polymorphisms for improved rabies virus surveillance

Ankeet Kumar[1], Sheetal Tushir[1], Yashas Devasurmutt[1], Sujith S Nath[1] & Utpal Tatu[1]*

Affiliation:

[1] Department of Biochemistry, Division of Biological Sciences, Indian Institute of Science, Bangalore, India

Corresponding Author:

*Utpal Tatu, Department of Biochemistry, Division of Biological Sciences, Indian Institute of Science, Bangalore, India

Email: tatu@iisc.ac.in

Abstract

Background: Rabies is an ancient disease that remains endemic in many countries. It causes many human deaths annually, predominantly in resource-poor countries. Over evolutionary timelines, several rabies virus (RABV) genotypes have stabilised, forming distinct clades. Extensive studies have been conducted on the origin, occurrence and spread of RABV clades. Single nucleotide polymorphisms (SNPs) distribution across the RABV genome and its clades remains largely unknown, highlighting the need for comprehensive whole-genome analyses.

Methods: We accessed whole genome sequences for RABV from public databases and identified SNPs across the whole genome sequences. Then, we annotated these SNPs using an R script, and these SNPs were categorised into different categories; universal, clade-specific, and clade-defining, based on the frequency of occurrence.

Results: In this study, we present the SNPs occurring in the RABV based on whole genome sequences belonging to 8 clades isolated from 7 different host species likely to harbour dog-related rabies. We classified mutations into several classes based on their location within the genome and assessed the effect of SNP mutations on the viral glycoprotein.

Conclusions: The clade-defining mutations have implications for targeted surveillance and classification of clades. Additionally, we investigated the effects of these mutations on the Glycoprotein of the virus. Our findings contribute to expanding knowledge about RABV clade diversity and evolution, which has significant implications for effectively tracking and combatting RABV transmission.

36   **Keywords:** rabies virus (RABV), single nucleotide polymorphisms (SNP) analysis, whole
37   genome SNP analysis, canine rabies, mutations in RABV clades.

Introduction

Rabies virus (RABV), along with some other species of the genus *Lyssavirus*, causes a disease called rabies. It is a deadly zoonotic viral encephalitis that is fatal in almost 100% of cases and significantly threatens human and animal health [1]. The *Lyssavirus* genus comprises 17 species, including the RABV, type species of the genus and other rabies-related viruses [2]. These species are classified into phylogroups based on genetic composition and antigenic differences [3]. Most Lyssavirus species have a limited impact on human health due to their narrow host range. RABV displays broad host tropism and is responsible for approximately 59,000 human deaths annually across the globe [4]. Almost 99% of human deaths caused by RABV globally are attributed to dog bites [1]. In India, dogs are the only known reservoir of RABV, and India bears the highest burden of rabies-related human deaths, accounting for around 20,000 fatalities annually [5], followed by China [6].

The RABV genome is approximately 12 kilobases long. The genome is non-segmented, single-stranded negative-sense RNA enclosed within an enveloped, bullet-shaped virion [7]. The genome codes for 5 proteins are arranged in the following order: Nucleoprotein (N), Phosphoprotein (P), Matrix (M), Glycoprotein (G), and Large (L) protein genes. The L gene is the largest, comprising approximately 53% of the total nucleotide content. It codes for a crucial, multifunctional protein called RNA-dependent RNA polymerase (RdRp) [8]. Like most RNA viruses, RABV RdRp lacks proofreading activity [9], resulting in variations, ultimately leading to the emergence of new variants and the formation of quasi-species during infection [10]. P protein codes for various isoforms; its full-length P protein interacts with the RdRp protein via its N-terminal region, while its shorter isoforms (P2-P5) with truncated N-termini likely serve alternative functions [11]. The N protein forms the core of the nucleocapsid, encapsulating the viral RNA in a helical structure [12]. The M protein interacts with the N protein and G protein facilitating viral envelopment [13]. The G protein is a trimeric and is essential for viral entry [14]. The G protein is known to interact with multiple host receptors [15]. It is also a target for neutralising antibodies and a therapeutic target in the virus [16].

Previously, studies primarily relied on sub-genomic sequences[17,18] , which has constrained our ability to understand whole-genome-level attributes comprehensively. Additionally, to our knowledge, there is no systematic study tabulating the global occurrence of single nucleotide polymorphisms (SNPs) in the RABV and comparing SNPs across the clades; a notable gap exists in the characterisation of clades. In this study, we present a comprehensive analysis of SNPs based on whole genome sequences of 8 clades of RABV isolated from 7 host species (Dog, Cat, Cattle, Horse, Sheep, Goat and Human). The SNPs result in various types of mutation at the protein level. The percentage of these mutations differs across the clades. We report some mutations occurring with high frequency (clade-specific, universal and clade-defining) and further analyse their impact on the G protein.

Methods

Collection and Curation of the WGS Data

To identify SNPs in the WGSs of the virus, we accessed data from 2 databases- Bacterial and Viral Bioinformatics Resource Center (BV-BRC) [19] and RABV-GLUE (RABV-GLUE) [20] (databases accessed on 1/11/2023). The databases contained varying numbers of WGSs

for the representative hosts. Comprehensive datasets were generated through manual curation, which involved merging non-overlapping sequences from two databases and removing redundancies. Annotations were further verified against NCBI databases (Nucleotide and Bio-sample). The pipeline of manual curation is presented in S1 Fig.

This study focuses on the dog-variant of RABV; therefore, we downloaded sequences isolated from the Dog host and 6 other hosts likely to harbour the dog variant of RABV. The other hosts were cat with 43 sequences, cattle with 106 sequences, horse with 8 sequences, sheep with 12 sequences, goat with 23 sequences and human with 47 sequences.

Sequence metadata files were downloaded from both databases. Metadata lacking clade information was supplemented using RABV-GLUE, a web-based tool from the University of Glasgow [21]. Only relevant information was retained in the refined metadata. Clade distribution across different hosts and the corresponding genome count is presented in Table 1.

The final metadata file for 778 genome sequences utilised in the study is present in the S1 File.

Identification of Single Nucleotide Polymorphisms

The identification of SNPs involved several steps. The multi-fasta sequence files for hosts were stored in different directories. These fasta files were subjected to SNP analysis using the NUCMER v3.1 [22]. This tool generates pairwise alignment of each sequence against the reference sequence of RABV (NC_001542.1) and produces a table of SNPs. The process was repeated for each host. This SNP table contained information regarding the variant position and nucleotide change compared to the reference sequence. The table generated from Nucmer was used for further analysis using R (v4.3.1).

Classification of Single Nucleotide Polymorphisms using R programming

The original R script for SNP analysis [23] was modified (S2 File) for improved accuracy by changing the "round" function to "ceiling". This was necessary because the "round" function could lead to misinterpretation of changes occurring at the first nucleotide position of a codon. This script utilises the SNP table generated using Nucmer in the previous step. The script utilises a General Feature Format 3 (GFF3) file containing information about the start and end positions of the genes. The GFF3 file for the RABV reference genome is available as S3 File. This GFF3 was manually curated to keep the information only for coding regions. The script utilises Seqinr (v4.2.30) [24] and Biostrings (v2.68.1) [25] tools to load a reference fasta file and translate the nucleotide changes into amino acids using the reference genome backbone. The translated amino acid changes are labelled as SNP (non-synonymous change) mutation, SNP-silent (synonymous change) mutation, deletion-frameshift, insertion frameshift and SNP-stop. The region falling outside the genes is labelled as extragenic. The generated data frame containing amino acid changes includes duplicate rows for an amino acid where more than one nucleotide change is responsible. To eliminate these duplications, an R script (S4 File) was written to merge rows with multiple nucleotide changes corresponding to the same amino acid residue.

Categorisation of High-Frequency Mutations

Most observed mutations were occurring with low frequencies. However, our analysis
focused on high-frequency mutations because of their potential significance in the biology of
the virus. Initially, we pinpointed mutations prevalent within individual hosts. The mutations
appearing in over 90% of sequences within a given host were denoted as "universal
mutations". Subsequently, given the presence of multiple clades within each host, our
attention turned to "clade-specific mutations". These mutations represented unique and non-
overlapping mutations localised within distinct clades coexisting within the same host. To
ensure the specificity of clade-specific mutations, universal mutations were excluded from
the dataset during the process. Finally, as similar clades were found in more than one host, we
discerned "clade-defining mutations" by identifying common clade-specific mutations across
different hosts for a clade. These mutations, characterised by their high frequency and
consistent presence, served as distinctive markers reliably delineating specific clades.

Due to the reliance on percentage values for classification, a minimum threshold of four
sequences in a host and the presence of at least two clades for comparison were established as
criteria for inclusion in the high-frequency mutation analysis. Consequently, horse and goat
data were excluded from this analysis as they failed to meet the criteria (S2 Fig.).

Assessing the Impact of Mutations on the Glycoprotein

To elucidate the mutational effects on the structure of RABV glycoprotein, homology
modelling and protein threading techniques were implemented. The RABV glycoprotein
backbone was modelled using the post-fusion Mokola virus structure (PDB ID: 6TMR),
while the stalk region utilised the H5N1 influenza virus hemagglutinin (PDB ID: 4UJM) as a
template, both in SWISS-MODEL [26]. DynaMut [27], which provides computational
predictions on the impact of mutations on proteins, was employed to predict the impact of
some of these mutations on protein.

Data Visualisation and Bioinformatics Tools

We utilised R (v4.3.1) for data analysis and visualisation. A map was created to understand
the global distribution of the RABV clades found in dog hosts (Fig. 1A) using the Maps
v3.4.1 tool from the R library. Genetic diversity at the nucleotide level was calculated using
DnaSP6 program [28]. The calculation of average mutations for the 5 proteins of RABV
clades is done using base R functions and dplyr package v1.1.2. The count of mutation is
normalised to protein length and averaged by sample number for that clade. The library
drawProteins v1.20.0 [29] was used to draw domains for different proteins in Fig. 5A. The
structure was visualised in the PyMol and ggplot2 v3.4.2 [30] was utilised to make figures.

Results

Global Distribution and Host Association of Rabies Virus Clades

Rabies virus shows global presence and, over time, has diversified into distinct genotypes
forming monophyletic clades [31]. Fig. 1A displays the spatial distribution of the 6 RABV
clades isolated from the dog hosts. The cosmopolitan clade demonstrates the broadest spatial
spread. Most countries predominantly harbour a single clade; China exhibits 3 clades, and
Russia, Iran, and Brazil show 2 clades each. The Asian clade ranks second in numbers for
dog hosts in our dataset; the clade was reported from China and neighbouring Southeast
Asian countries. The Arctic clade spans three continents (Europe, Asia, and the USA).

However, dog-mediated rabies is not endemic to America and Europe [32]. The Africa-2
clade is restricted to western Africa. The Indian-Sub clade is reported from the UK, and the
Bats clade in dogs is found in South American countries.

Fig. 1B shows the occurrence of hosts in 78 countries. None of the countries in the dataset
showed the sequencing data for all 7 hosts simultaneously. The maximum number of hosts
observed in any single country was 5- China, Hungary, Kenya, Tanzania, and the USA
demonstrated RABV sequences from 5 hosts, followed by Mexico and Moldova, which
represented 4 hosts each. Dog hosts were found to be the most common host (observed in 67
countries), followed by cattle (in 22 countries), humans (in 21 countries), and cats (in 18
countries). Conversely, goats, horses, and sheep were the least common, each represented in
only 5 countries. Forty-three countries had data only from 1 host; out of these 43 countries,
around 72% showed sequences only from dogs.

To understand the co-occurrence of hosts and clades in the dataset, we looked for countries
with either more than 1 host or more than 1 clade; most of the countries passed the criteria,
while Indonesia and Bangladesh had only one host harbouring a single clade, so they were
excluded. As shown in Fig. 1C, the Cosmopolitan clade is the most common clade across all
7 hosts, followed by the Arctic. Bats clade is reported from the USA, Mexico, and Brazil, and
RAC-SK is only reported from the USA.

Characterisation of Mutation Classes and Nucleotide Position Biasness in RABV Genome Across Diverse Hosts

RNA viruses show a high mutation rate. To understand the differences in RABV clades at the
molecular level, we identified nucleotide variations and categorised them into six classes:
SNP-silent, SNP, SNP-stop, extragenic, insertion-frameshift and deletion-frameshift mutation
using an R script. A detailed breakdown of mutation counts categorised by functional class
for each RABV clade in the five proteins and the extragenic region, with respect to different
host species, can be found in the S1 Table. Fig. 2A presents the percentage of various classes
of mutations based on 539 dog RABV sequences belonging to 6 clades (Africa-2, Arctic,
Asian, Bats, Cosmopolitan, and Indian-Sub). The synonymous mutation or SNP-silent is the
most prevalent (70-82%) class of mutations across the clades. The subsequent mutation
classes are SNP mutation and extragenic mutation class, which account for most of the
remaining variations (approximately 30%). Less than 1% of mutations fall into insertion-
frameshift and deletion-frameshift classes. The percentage of non-synonymous mutation in
different clades ranges between 11-15%. The extragenic class ranged from 6-15% across the
clades. The cosmopolitan clade displayed the highest percentage of extragenic and SNP
mutations in all the hosts. The reason for displaying higher SNP mutations in cosmopolitan
could be because multiple sub-clades have become associated with geographical locations.

Additionally, we analysed the contribution of the nucleotide positions within a codon on SNP
and SNP silent mutations. Fig. 2B shows that SNP silent mutations predominantly result from
III nucleotide change within a codon. Only a minor fraction of SNP silent mutations occur
due to changes in I nucleotide and even less due to II. Conversely, all three nucleotide
changes contribute to SNP mutations significantly. Both I and II nucleotide changes
contribute almost equally to SNP mutations in RABV proteins except for G and P, which
exhibited lower contributions from II nucleotide change. While SNPs primarily arise from
individual nucleotide changes at the I and II positions of codons, a considerable portion also

results from simultaneous substitutions within a single codon. The co-occurrence of SNPs
within a codon depicted contribution disparity within clades and proteins, as shown in Fig.
2C.

Fig. 2D represents the occurrence of various classes of mutations in clades of other hosts
present in the dataset (Cat, Cattle, Sheep, Goat, Horse and Human). Like dogs, the variations
observed in other host species also showed SNP silent as the predominant mutation, followed
by SNP and extragenic mutation. The Cosmopolitan clade demonstrated the lowest
percentage of SNP-silent mutations throughout all hosts compared to other clades. These
results highlight differences in the selection pressures across the RABV clades and hosts.
Data on the mutations found for different hosts is presented in the S5 File.

219 Phosphoprotein Exhibits Highest Mutation and C>T is Major Transition Across Diverse
220 Clades of RABV

Previous studies have reported that the Phosphoprotein shows the highest mutation rate [33].
We investigated the mutation patterns across diverse RABV clades present in 7 hosts. Fig. 3A
depicts mutation trends across 5 proteins of RABV clades isolated from dogs. In our study,
most clades showed Phosphoprotein as the highest mutating protein among the 5 RABV
proteins in the sequences isolated from dogs. The Glycoprotein and RdRp exhibited similar
average mutations despite the huge difference in protein length. Notably, the Bats clade
exhibited a distinct pattern of variation, with the highest number of mutations observed in the
Nucleoprotein, followed by fewer mutations in the Phosphoprotein, and no mutations
detected in the Matrix, Glycoprotein, and RdRp proteins.

The nucleotide transitions and transversion across different clades found in the dog host are
presented in Fig. 3B. The top 4 nucleotide transitions (C>T, A>G, T>C, G>A) were
consistent throughout the clades. Cytosine to Thymine (C>T) was found to be the
predominant transition in 5 clades, while A> G was found to occur with the highest frequency
in the Cosmopolitan clade. Transversions were present, but the frequency was lower than the
transitions. The trend of these changes varied across the clades in dogs.

Fig. 3C delineates mutations across 5 proteins within diverse RABV clades found in 6 hosts.
While the overall mutation trend (P>G~L>M>N) is the same as that found for the dog host,
distinct patterns emerged in the hosts harbouring Bats and RAC-SK clades. For instance, the
Bats clade showcased heightened N protein mutations in the cattle and human hosts, whereas
P protein displayed elevated mutations in horse and cat hosts. In the cattle host, the mutations
were significantly less in Phosphoprotein and RdRp for the Bats clade. The RAC-SK clade
consistently exhibited more mutations in the N protein across all the hosts. Interestingly, no
mutations were observed in the Glycoprotein and Matrix proteins across any host for the
RAC-SK clade, while mutations were seen for RdRp in the cattle host.

The 4 most common nucleotide transitions found in dog RABV were checked in other hosts
as well (Fig. 3D). C>T transition was dominant in sheep, goat, cattle, and horse hosts.
However, the T>C transition prevailed in human and cat hosts. In the sheep, host A>G was
the second most common nucleotide transition, while in all other hosts, it was third.
Collectively, these results highlight the different evolutionary pressures across RABV clades
in different hosts.

Mapping the Distribution of Rabies Virus Mutations Across Clades and Identifying Universal Mutations

Similar to other RNA viruses, RABV lacks a proofreading mechanism which drives its ability to accumulate mutations quickly [34]. Fig. 4A sheds light on the distribution of mutations identified across the four RABV clades circulating in dog hosts. The analysis revealed that the Cosmopolitan clade harboured the most mutations, followed by Asian, Arctic, and Africa-2 clades in dogs. This trend also remains consistent for unique and non-overlapping mutations within each clade. These findings suggest that the Cosmopolitan clade may have undergone greater evolutionary pressure or higher adaptability, facilitating its survival in diverse environments. Moreover, the Cosmopolitan clade exhibits the highest degree of overlap with the Asian clade, sharing 489 mutually, followed by the Arctic clade. Notably, 1539 mutations are common across all four clades of dogs and most of these mutations primarily occur at low frequencies (data not shown). Fig. 4B depicts the distribution of universal mutations found across 5 hosts. The sheep host displays the highest unique universal mutations, followed by dog, cat, human and cattle hosts. Sheep possessed 144 unique universal mutations, while 78 were common in dogs and sheep. Human hosts shared 2 universal mutations with dogs, and 1 mutation was exclusive to human hosts. Cattle and cats shared no mutation exclusively with the dogs. A total of 9 mutations were found to be conserved across all the hosts, potentially essential for viral functions that remain unaltered across diverse hosts.

The distribution of these universal mutations across proteins is showcased in Fig. 4C. RdRp protein displayed the highest universal mutations, but the mutations were only seen for dog and sheep hosts. This was followed by G and M proteins, which showed mutations for dog, human and sheep hosts. The N and M showed universal mutations for all the hosts. Fig. 4D highlights 9 conserved mutations common to RABV and found across all hosts in high frequency. Only 3 belong to the SNP mutation class, and the rest are SNP silent mutations. This subset of non-synonymous mutations may contribute to subtle functional shifts that enhance viral fitness. A list of unique mutations in hosts with the universal mutations for the respective hosts is provided in the S6 File. This mutation landscape sheds light on the evolutionary dynamics of RABV and suggests that mutation accumulation, particularly in receptor-binding and replication-related proteins, may contribute to the virus's ability to persist and adapt across different hosts.

Assessing the Impact of High-Frequency Mutations on RABV Glycoprotein

The RABV glycoprotein is crucial in facilitating virus entry and serves as the primary target for neutralising antibodies, making it a key protein for host adaptation and immune evasion [14]. Fig. 5A highlights the clade-specific SNP mutations present in the 4 proteins. M protein did not display any clade-specific SNP mutation. N protein displayed 1 mutation in Africa and the Arctic each. P displayed 1 and 2 mutations in Arctic and Africa-2, respectively. RdRp protein, essential for viral replication, displayed 5 clade-specific SNP mutations in Africa-2 and 4 in Asian clades. G protein displayed 5 clade-specific mutations in Africa and 2 mutations in the Arctic clade. The distribution of these mutations may clade-specific adaptations and immune evasion strategies. The stability values for mutations in the G protein are available in the S2 Table. A complete list of all the clade-specific mutations observed in dog RABV clades is presented in Table 2. The RdRp protein showed the highest number of

clade-specific mutations in dogs, and the mutations in RdRp were found for the Africa-2 and
Asian clades. Africa-2 showed clade-specific mutations in all 5 proteins. All the clade-
specific mutations seen in Glycoprotein belonged to the SNP class.

Fig. 5B shows the post-fusion structure of a monomer of Glycoprotein of RABV, with
mutations in the stalk and ectodomain region. The surface structure of trimeric Glycoprotein
from the top and front view is also presented. Fig. 5C shows the trimeric structure of the
Glycoprotein; the zoomed-in portion highlights the 6 mutations occurring in 2 regions. The
mutations in blue (ectodomain) occur in the snake-toxin-like region of the Glycoprotein
(V181I, A182T, and N177K are universal mutations, while V183I is the only clade-specific
mutation occurring in the Arctic clade in this region). 2 universal mutations E444K and
R445Q are located in the stalk region of the Glycoprotein, and this region is essential for the
interaction of 3 monomers of the Glycoprotein [35]. Fig. 5D highlights the altered
interactions between residue 181 and the neighbouring residues, likely impacting protein
function. The free energy values suggest it to be one of the highest stabilising mutations, and
the interactions are seen to have increased in the mutated version of the residue. This
increased stability may enhance viral entry. The clade-specific mutations for other hosts and
clade-defining mutations are presented in the S7 File. The accumulation of clade-specific
mutations in different proteins across RABV clades may reflect the evolutionary pressures.

Discussion

Rabies is one of the oldest zoonotic diseases known to mankind. Regardless of the
availability of ample WGSs, the characterisation and comparison of SNPs across the clades
were lacking. We report SNPs in RABV clades leveraging a dataset comprising 778 whole
genome sequences isolated from 7 hosts. Despite the global presence of RABV, the
sequencing data available for it is highly biased, severely affected regions lack surveillance
regarding the numbers and diversity of the hosts. This leads to the underestimation of
infected cases and the types of hosts infected. The re-emergence cases of rabies [36,37]
demand robust surveillance mechanisms in dogs as well as other hosts to reach the "Zero by
2030" goal [38].

Most of the mutations identified in this study occurred with a low frequency, yet some low-
frequency mutations could still play a crucial role in viral evolution and adaptation. Though
these mutations are not extensively described here, their longitudinal tracking is essential to
understanding emerging variants. The P protein displayed the highest mutation across most of
the clades, likely due to host immune pressure and clade-adaptive strategies. While in some
cases for the Bats clade and all the cases for the RAC-SK clade, the mutations were higher in
the N protein, emphasising the divergence in mutation pattern due to host-specific factors,
although more studies are required to understand these dynamics. SNP silent was found to be
the major class of RABV mutations, highlighting the negative selection in RABV and
limiting the changes in the protein sequence. Even though the SNP-silent mutations do not
change amino acids, the difference at the nucleotide level contributes to changes in codon
usage [39,40] and is also known to change protein structure [40], suggesting that these
mutations are not entirely inconsequential. The percentage distribution of mutation types
varies across the clades. This analysis presents that there are inherent differences in the
clades, and their association with the host must be explored to underscore the adaptive
mutations of RABV across different clades.

The presence of universal mutations, along with distinct clade-specific mutations, indicates
that each clade may require a unique set of mutations for adaptation and survival. These
mutations appear to have stabilised within their respective clades over the course of evolution
despite the rapid mutation rate of RABV, suggesting their importance in viral biology.
Bonnaud et al. (2019) identified adaptive mutations in the N and G proteins of RABV that
facilitated the adaptation of dog variant in foxes [41]. They reported that transmission was
unidirectional, and several mutations observed in their experiment were seen in the sequences
isolated from natural hosts. However, the study only used cosmopolitan strain, and we found
that residues described for the adaptation in their study belonged to the universal class in dog
hosts (residue 61 in N protein and residue 480 in G protein). This suggests that studying
SNPs will help unravel the adaptive mechanisms of the virus and the evolutionary forces
responsible for the changes and cross-species transmission.

The clade-specific mutations represent key mutations required for viral adaptation. We
compared the clade-specific mutations across the same clade in different hosts and identified
clade-defining mutations (S7 File). These clade-defining mutations offer a promising avenue
for detecting and categorising clades based on SNPs. We have developed a PCR-based rapid
typing tool based on SNPs (data not shown), which will improve the epidemiological tracking
of RABV clades by providing an alternative way to determine clades where genome
sequencing is inaccessible. A summary table for clade-specific and clade-defining mutations
for dog hosts is presented as S4 Table. Similar strategies have been proven effective in
tracking viral lineages, notably in SARS-CoV-2 [42]. Our analysis unveils the absence of
clade-specific mutations within the cosmopolitan clade. Also, the presence of high-frequency
mutations for subclades (S8 File) hints that the diversity at the sub-clade level is higher in the
Cosmopolitan clade. The Cosmopolitan showed high extragenic mutations, although
extragenic mutations do not directly contribute to the protein sequence but are crucial for
protein synthesis and regulation [43].

The genetic diversity calculated for RABV clades found in dogs showed no discernible
correlation with prevalence or the temporal origins in the host. Notably, the Asian clade in
dogs exhibited the highest nucleotide diversity, as indicated by the Pi (π) values (S3 Table).
This particular clade, present in dog hosts, stands out as one of the oldest identified clades
after the Indian-Sub clade [44]. The occurrence of similar clades in multiple hosts in a
country highlights the events of cross-species transmission. However, currently, only
phylogeny-based methods are available to identify clades; we believe the clade-defining
SNPs reported in this study will serve as an alternative approach for clade identification,
which will aid in the tracking of RABV clades.

Rabies is known to persist in two distinct forms: furious form and paralytic form. Hueffer et
al. (2017) reported that the glycoprotein can modify host behaviour through a region that
binds to the AchR receptor [45]. We find several mutations in an area of G protein (snake-
toxin-like region, 175-203 residues in Glycoprotein) that binds to AchR. Most of the
mutations in this region are universal (V181I, A182T, N177K). However, a stabilising
mutation (V183I) is only present in Arctic clade sequences, further supporting the idea that
clade-specific mutations may influence viral fitness and adaptation. The study of mutations
also becomes important for understanding the differences in the fitness of the virus strains
[46].

While this study encompassed sequences from 7 distinct hosts, dog hosts constituted more than half of the dataset. Countries significantly affected by RABV reported less WGS, impeding an in-depth understanding of RABV in these regions. The low-frequency mutations are not described, but these mutations are important for a comprehensive understanding of the virus evolution and circulation pattern. Furthermore, the inability to validate the functional implications of these pivotal mutations through wet lab experimentation remains a notable limitation. The diversity and distinct mutational repertoire of clades make evaluating vaccine efficacy against RABV clades crucial. Addressing these points would enhance our understanding of RABV evolution and facilitate more targeted and effective control measures against this global health concern.

Acknowledgements

The authors would like the whole Utpal Tatu lab for their discussions.

Author contributions

Conceptualisation: Ankeet Kumar and Utpal Tatu.

Data curation: Ankeet Kumar.

Formal analysis: Ankeet Kumar and Yashas Devasurmutt.

Investigation: Ankeet Kumar and Utpal Tatu.

Methodology: Ankeet Kumar and Utpal Tatu.

Supervision: Utpal Tatu.

Validation: Ankeet Kumar, Sheetal Tushir and Utpal Tatu.

Writing – original draft: Ankeet Kumar and Sujith S Nath.

Writing – review & editing: Ankeet Kumar, Sheetal Tushir and Utpal Tatu.

Funding

UT acknowledges the DBT-IISc partnership.

AK acknowledges CSIR for financial support.

YD and ST acknowledge fellowship from the institute.

No external funding was obtained to support the project.

Data Availability

The data supporting the findings in the study is available in the manuscript and as supporting data.

Conflict of interest

The authors declare no conflict of interest.

Ethics approval

Not applicable

References

[1] Brunker K, Mollentze N. Rabies Virus. Trends Microbiol 2018;26:886–7. https://doi.org/10.1016/j.tim.2018.07.001.

[2] Shipley R, Wright E, Smith SP, Selden D, Fooks AR, Banyard AC. Taiwan Bat Lyssavirus: In Vitro and In Vivo Assessment of the Ability of Rabies Vaccine-Derived Antibodies to Neutralise a Novel Lyssavirus. Viruses 2022;14. https://doi.org/10.3390/V14122750.

[3] Černe D, Hostnik P, Toplak I, Presetnik P, Maurer-Wernig J, Kuhar U. Discovery of a novel bat lyssavirus in a Long-fingered bat (Myotis capaccinii) from Slovenia. PLoS Negl Trop Dis 2023;17:e0011420. https://doi.org/10.1371/journal.pntd.0011420.

[4] BANYARD AC, TORDO N. Rabies pathogenesis and immunology. Rev Sci Tech l'OIE 2018;37:323–30. https://doi.org/10.20506/rst.37.2.2805.

[5] Yale G, Sudarshan S, Taj S, Patchimuthu GI, Mangalanathan BV, Belludi AY, et al. Investigation of protective level of rabies antibodies in vaccinated dogs in Chennai, India. Vet Rec Open 2021;8. https://doi.org/10.1002/vro2.8.

[6] TU C, FENG Y, WANG Y. Animal rabies in the People's Republic of China. Rev Sci Tech l'OIE 2018;37:519–28. https://doi.org/10.20506/rst.37.2.2820.

[7] Itakura Y, Tabata K, Saito T, Intaruck K, Kawaguchi N, Kishimoto M, et al. Morphogenesis of Bullet-Shaped Rabies Virus Particles Regulated by TSG101. J Virol 2023;97. https://doi.org/10.1128/jvi.00438-23.

[8] Morin B, Liang B, Gardner E, Ross RA, Whelan SPJ. An In Vitro RNA Synthesis Assay for Rabies Virus Defines Ribonucleoprotein Interactions Critical for Polymerase Activity. J Virol 2017;91. https://doi.org/10.1128/JVI.01508-16.

[9] Tao YJ, Ye Q. RNA Virus Replication Complexes. PLOS Pathog 2010;6:e1000943. https://doi.org/10.1371/JOURNAL.PPAT.1000943.

[10] Conselheiro JA, Barone GT, Miyagi SAT, de Souza Silva SO, Agostinho WC, Aguiar J, et al. Evolution of Rabies Virus Isolates: Virulence Signatures and Effects of Modulation by Neutralizing Antibodies. Pathogens 2022;11:1556. https://doi.org/10.3390/pathogens11121556.

[11] Okada K, Ito N, Yamaoka S, Masatani T, Ebihara H, Goto H, et al. Roles of the Rabies Virus Phosphoprotein Isoforms in Pathogenesis. J Virol 2016;90:8226–37. https://doi.org/10.1128/JVI.00809-16.

[12] Masatani T, Ito N, Shimizu K, Ito Y, Nakagawa K, Sawaki Y, et al. Rabies Virus Nucleoprotein Functions To Evade Activation of the RIG-I-Mediated Antiviral Response. J Virol 2010;84:4002–12. https://doi.org/10.1128/JVI.02220-09.

[13] Liu X, Li F, Zhang J, Wang L, Wang J, Wen Z, et al. The ATPase ATP6V1A facilitates rabies virus replication by promoting virion uncoating and interacting with the viral matrix protein. J Biol Chem 2021;296:100096. https://doi.org/10.1074/jbc.RA120.014190.

[14] Yang F, Lin S, Ye F, Yang J, Qi J, Chen Z, et al. Structural Analysis of Rabies Virus Glycoprotein Reveals pH-Dependent Conformational Changes and Interactions with a Neutralizing Antibody. Cell Host Microbe 2020;27:441-453.e7.

https://doi.org/10.1016/J.CHOM.2019.12.012.

[15]   Lian M, Hueffer K, Weltzin MM. Interactions between the rabies virus and nicotinic
acetylcholine receptors: A potential role in rabies virus induced behavior
modifications. Heliyon 2022;8:e10434.
https://doi.org/10.1016/J.HELIYON.2022.E10434.

[16]   Shi C, Sun P, Yang P, Liu L, Tian L, Liu W, et al. Research progress on neutralizing
epitopes and antibodies for the Rabies virus. Infect Med 2022;1:262–71.
https://doi.org/10.1016/J.IMJ.2022.09.003.

[17]   Tsai KJ, Hsu WC, Chuang WC, Chang JC, Tu YC, Tsai HJ, et al. Emergence of a
sylvatic enzootic formosan ferret badger-associated rabies in Taiwan and the
geographical separation of two phylogenetic groups of rabies viruses. Vet Microbiol
2016;182:28–34. https://doi.org/10.1016/j.vetmic.2015.10.030.

[18]   Davis R, Nadin-Davis SA, Moore M, Hanlon C. Genetic characterization and
phylogenetic analysis of skunk-associated rabies viruses in North America with special
emphasis on the central plains. Virus Res 2013;174:27–36.
https://doi.org/10.1016/j.virusres.2013.02.008.

[19]   Olson RD, Assaf R, Brettin T, Conrad N, Cucinell C, Davis JJ, et al. Introducing the
Bacterial and Viral Bioinformatics Resource Center (BV-BRC): a resource combining
PATRIC, IRD and ViPR. Nucleic Acids Res 2023;51:D678.
https://doi.org/10.1093/NAR/GKAC1003.

[20]   Singer JB, Thomson EC, McLauchlan J, Hughes J, Gifford RJ. GLUE: a flexible
software system for virus sequence data. BMC Bioinformatics 2018;19.
https://doi.org/10.1186/S12859-018-2459-9.

[21]   Campbell K, Gifford RJ, Hill V, Toole O', Hampson K, Brunker &. Making Genomic
Surveillance Deliver: A Lineage Classification and Nomenclature System to Inform
Rabies Elimination n.d. https://doi.org/10.1101/2021.10.13.464180.

[22]   Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, et al.
Versatile and open software for comparing large genomes. Genome Biol 2004;5:R12.
https://doi.org/10.1186/gb-2004-5-2-r12.

[23]   Mercatelli D, Giorgi FM. Geographic and Genomic Distribution of SARS-CoV-2
Mutations. Front Microbiol 2020;11:1800.
https://doi.org/10.3389/FMICB.2020.01800/BIBTEX.

[24]   Charif D, Lobry JR. SeqinR 1.0-2: A Contributed Package to the R Project for
Statistical Computing Devoted to Biological Sequences Retrieval and Analysis
2007:207–32. https://doi.org/10.1007/978-3-540-35306-5_10.

[25]   H Pagès, P Aboyoun, R. Gentleman and SD. Package "Biostrings" Title Efficient
manipulation of biological strings 2013.

[26]   Schwede T, Kopp J, Guex N, Peitsch MC. SWISS-MODEL: an automated protein
homology-modeling server. Nucleic Acids Res 2003;31:3381.
https://doi.org/10.1093/NAR/GKG520.

[27]   Rodrigues CHM, Pires DEV, Ascher DB. DynaMut: predicting the impact of
mutations on protein conformation, flexibility and stability. Nucleic Acids Res
2018;46:W350. https://doi.org/10.1093/NAR/GKY300.

[28]   Rozas J, Ferrer-Mata A, Sanchez-DelBarrio JC, Guirao-Rico S, Librado P, Ramos-
504   Onsins SE, et al. DnaSP 6: DNA Sequence Polymorphism Analysis of Large Data
505   Sets. Mol Biol Evol 2017;34:3299–302. https://doi.org/10.1093/MOLBEV/MSX248.

506   [29]   Brennan P. drawProteins: a Bioconductor/R package for reproducible and
507   programmatic generation of protein schematics. F1000Research 2018;7.
508   https://doi.org/10.12688/F1000RESEARCH.14541.1.

509   [30]   Wickham H. ggpolt2 Elegant Graphics for Data Analysis. Use R! Ser 2016:211.

510   [31]   Dellicour S, Troupin C, Jahanbakhsh F, Salama A, Massoudi S, Moghaddam MK, et
511   al. Using phylogeographic approaches to analyse the dispersal history, velocity and
512   direction of viral lineages — Application to rabies virus spread in Iran. Mol Ecol
513   2019;28:4335–50. https://doi.org/10.1111/mec.15222.

514   [32]   Tiwari HK, Gogoi-Tiwari J, Robertson ID. Eliminating dog-mediated rabies:
515   challenges and strategies. Anim Dis 2021;1:1–13. https://doi.org/10.1186/S44149-021-
516   00023-7/FIGURES/1.

517   [33]   Troupin C, Dacheux L, Tanguy M, Sabeta C, Blanc H, Bouchier C, et al. Large-Scale
518   Phylogenomic Analysis Reveals the Complex Evolutionary History of Rabies Virus in
519   Multiple Carnivore Hosts. PLOS Pathog 2016;12:e1006041.
520   https://doi.org/10.1371/JOURNAL.PPAT.1006041.

521   [34]   Domingo E. RNA Genetics: Volume III: Variability of RNA Genomes. 2018.

522   [35]   Fernando BG, Yersin CT, José CB, Paola ZS. Predicted 3D Model of the Rabies Virus
523   Glycoprotein Trimer. Biomed Res Int 2016;2016.
524   https://doi.org/10.1155/2016/1674580.

525   [36]   Botto Nuñez G, Becker DJ, Plowright RK. The emergence of vampire bat rabies in
526   Uruguay within a historical context. Epidemiol Infect 2019;147:e180.
527   https://doi.org/10.1017/S0950268819000682.

528   [37]   Sharma B, Dhand NK, Timsina N, Ward MP. Reemergence of Rabies in Chhukha
529   District, Bhutan, 2008. Emerg Infect Dis 2010;16:1925–30.
530   https://doi.org/10.3201/eid1612.100958.

531   [38]   Fooks AR, Cliquet F, Finke S, Freuling C, Hemachudha T, Mani RS, et al. Rabies. Nat
532   Rev Dis Prim 2017 31 2017;3:1–19. https://doi.org/10.1038/nrdp.2017.91.

533   [39]   Cuevas JM, Domingo-Calap P, Sanjuán R. The Fitness Effects of Synonymous
534   Mutations in DNA and RNA Viruses. Mol Biol Evol 2012;29:17–20.
535   https://doi.org/10.1093/MOLBEV/MSR179.

536   [40]   Rosenberg AA, Marx A, Bronstein AM. Codon-specific Ramachandran plots show
537   amino acid backbone conformation depends on identity of the translated codon. Nat
538   Commun 2022;13:1–11. https://doi.org/10.1038/s41467-022-30390-9.

539   [41]   Bonnaud EM, Troupin C, Dacheux L, Holmes EC, Monchatre-Leroy E, Tanguy M, et
540   al. Comparison of intra- and inter-host genetic diversity in rabies virus during
541   experimental cross-species transmission. PLOS Pathog 2019;15:e1007799.
542   https://doi.org/10.1371/JOURNAL.PPAT.1007799.

543   [42]   Nasir A, Aamir UB, Kanji A, Bukhari AR, Ansar Z, Ghanchi NK, et al. Tracking
544   SARS-CoV-2 variants through pandemic waves using RT-PCR testing in low-resource

settings. PLOS Glob Public Heal 2023;3:e0001896.
https://doi.org/10.1371/journal.pgph.0001896.

[43]   Palusa S, Ndaluka C, Bowen RA, Wilusz CJ, Wilusz J. The 3′ Untranslated Region of
the Rabies Virus Glycoprotein mRNA Specifically Interacts with Cellular PCBP2
Protein and Promotes Transcript Stability. PLoS One 2012;7:e33561.
https://doi.org/10.1371/journal.pone.0033561.

[44]   Holtz A, Baele G, Bourhy H, Zhukova A. Integrating full and partial genome
sequences to decipher the global spread of canine rabies virus. Nat Commun 2023 141
2023;14:1–13. https://doi.org/10.1038/s41467-023-39847-x.

[45]   Hueffer K, Khatri S, Rideout S, Harris MB, Papke RL, Stokes C, et al. Rabies virus
modifies host behaviour through a snake-toxin like region of its glycoprotein that
inhibits neurotransmitter receptors in the CNS. Sci Rep 2017;7:12818.
https://doi.org/10.1038/s41598-017-12726-4.

[46]   Bloom JD, Neher RA. Fitness effects of mutations to SARS-CoV-2 proteins. Virus
Evol 2023;9. https://doi.org/10.1093/VE/VEAD055.

Tables.

Host Number	Host Name	Clade Number	Major Clade	Number of genome sequences
1	Dog (*Canis lupus familiaris*)	1	Cosmopolitan	383
		2	Asian	65
		3	Arctic	50
		4	Africa-2	37
		5	Bats	3
		6	Indian subcontinent (Indian-Sub)	1
2	Cat (*Felis catus*)	1	RAC-SK (Raccoon-Skunk)	18
		2	Cosmopolitan	13
		3	Africa-3	3
		4	Arctic	3
		5	Asian	3
		6	Africa-2	2
		7	Bats	1
3	Cattle (*Bos taurus*)	1	Cosmopolitan	70
		2	Arctic	12
		3	RAC-SK	11
		4	Bats	7
		5	Asian	5
		6	Indian-Sub	1
4	Goat (*Capra hircus*)	1	Cosmopolitan	22
		2	Arctic	1
5	Horse (*Equus caballus*)	1	RAC-SK	3
		2	Cosmopolitan	2
		3	Bats	1
		4	Asian	1
		5	Arctic	1
6	Human (*Homo sapiens*)	1	Cosmopolitan	14
		2	Asian	13
		3	Arctic	8
		4	Africa-2	6
		5	Indian-Sub	4
		6	Bats	2
7	Sheep (*Ovis aries*)	1	Asian	7
		2	Cosmopolitan	4
		3	Arctic	1

563 **Table 1.** Dataset of hosts and their clades with the associated count.

Clade	Protein	Mutations
	Nucleoprotein	**F213Y**
	Phosphoprotein	K139K, **P159L**

Africa-2	Matrix Protein	S185S
	Glycoprotein	**F8V**, **Q263R**, **N355D**, **N375H**, **N484K**
	RdRp	**M93I**, **T436A**, **N683D**, Q745Q, **K1624R**, **T1961A**
Arctic	Nucleoprotein	**M60L**, P135P
	Phosphoprotein	**E165D**, **A267G**
	Glycoprotein	**L16M**, **V183I**
Asian	RdRp	**L2I**, **L424I**, **A1535S**, **N1839E**

Table 2. Clade-specific mutations were observed in 3 clades of RABV in dog hosts (bold
values highlight SNP mutations)**.**

Fig. 1. Geographic distribution of different clades of RABV based on whole genome
sequences (N=778). **A.** The world map shows the prevalence of RABV clades found in the
dog host; colours highlight different clades. **B.** Distribution of hosts across 78 countries. The
coloured bars represent different hosts. **C.** Represents countries with more than one host or a
single host with more than one clade. Major clades are represented in different colours

Fig. 2. Analysis of mutation classes and underlying nucleotide positions involved in
RABV. A. The percentage of different mutation classes occurring in the RABV clades of dog
host. **B.** The contribution of nucleotide position within a codon for the SNP and SNP silent
mutation class in RABV genes. The y-axis contains the fraction contributed by three
nucleotides of a codon. **C.** The fraction of different (individual and co-occurring) nucleotide
positions to SNP mutations in the RABV genes. **D.** The percentage of mutation classes
observed in different clades of the other six hosts.

Fig. 3. Mutation counts across five proteins of RABV and underlying nucleotide
transitions and transversions. A. Mutation count normalised over protein length in the
RABV proteins. The x-axis presents the proteins of RABV, and the y-axis represents the
mutation count normalised to the protein length. **B.** The top 12 most common nucleotide
transition and transversion in different RABV clades (dog host). The coloured bars represent
nucleotide change. **C.** Mutations observed for RABV proteins in other hosts. **D.**
Representation of the four most common nucleotide transitions in the clades of other hosts.

Fig. 4. The distribution of mutations in dog RABV and universal mutations in other
hosts. A. Venn diagram shows the distribution of SNP and SNP silent mutations in RABV
clades isolated from dog hosts. The "N" indicates the total number of unique mutations for
the clades. **B.** Venn diagram depicts the distribution of universal mutations across five hosts.
C. Highlights the distribution of universal mutations in the five proteins across different
hosts; the colours represent hosts. **D.** Representation of nine common mutations in RABV
sequences found across all hosts.

Fig. 5. Clade-specific mutations in the glycoprotein. A. G protein with different
ectodomains, showing clade-specific mutations. **B.** RABV Glycoprotein structure predicted
using the SWISS-MODEL, showing the post-fusion form of protein. The transmembrane
region is highlighted in pink, the stalk is coloured red, and the ectodomain is represented with
blue colour (left). The surface structure of trimeric-RABV Glycoprotein top and front view
(right). **C.** The trimeric ribbon representation showing the interaction of three monomers of
Glycoprotein (left). Zoomed-in box represents mutations occurring in the AchR receptor
binding region. The green and orange represent universal and clade-specific mutations,
respectively. Two universal mutations, E444K and R445Q occur in the stalk region of the
glycoprotein. **D.** Represents the alteration of the interaction upon V to I change at 181
residue.

Supplementary Data:

S1 Fig. The pipeline used to curate the RABV dataset.

The number of sequences is shown with the corresponding hosts and databases. The
manual curation was done to identify the wrong annotations and add clade-related
information.

S2 Fig. The hosts and clades used for high-frequency mutation analysis.

The clades that were used for the analysis (> 4 sequences) are represented in black. The
clades and hosts that are shown in red were not used for the analysis.

S1 File. The list of genome accession utilised in the study with clade information.

S2 File. The modified script used to translate nucleotide variations (SNPs) to amino acids.

S3 File. A gff3 file with the information of coding regions for five proteins of RABV.

S4 File. An R script employed to merge the rows with nucleotide variation corresponding to
the   same amino acid.

S5 File. Host-wise characterisation of total mutations.

S6 File. A list of total and universal mutations represented in different clades of RABV.

S7 File. A list of clade-specific and clade-defining mutations for different hosts.

S8 File. Clade-specific mutations for the Cosmopolitan clade at the subclade level.

S1 Table.   Count of mutations in different clades across hosts.

S2 Table. Energy change values for mutations in the G protein.

S3 Table. A summary table for clades found in Dogs.

S4 Table. A table representing diversity for the clades found in dogs.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Supplementary Figures

S1 Fig

Database	Filters (Sequence Length > 11,000)	Manual Curation	Curated Dataset		Remarks
1.RABV-GLUE	**2,387**	**Intersect Accessions** Merge data from two databases	Dog	539	Some sequences were uncommon between two databases
Total 26,319			Cat	43	A list of misannotated cat accessions is provided below
Dog 6,014	328	**Manual Inspection** Incorrect wrong annotation	Cattle	106	
Cat 1,010	49				
Cattle 2,491	64	Add missing information on the metadata file like collection year, etc	Goat	23	
Goat 165	21		Horse	8	
Horse 285	7				
Human 840	41	**Classification into clades**	Human	47	One sequence (LT839613) belonging to EBLV was labelled as RABV
Sheep 92	12	RABV-GLUE was used to classify the sequences into clades			
2.BV-BRC	**(Complete Genome)**		Sheep	12	
Total 29,102	**2,484**				
Dog 8,547	503				
Cat 437	36				
Cattle 22	96				
Goat 171	21				
Horse 201	7				
Human 718	35				
Sheep 119	9				

Cattle sequences labeled as Cat in RABV-GLUE

S.No	Accession Number
1	ON366706
2	ON366707
3	ON366708
4	ON366709
5	ON366710
6	MK596362
7	MK596364
8	MN175989
9	MK540669
10	KX149163
11	KX149164
12	KR230089
13	KR230090
14	LC029869
15	KC171643
16	KC169985

The data were downloaded on November 1st 2023

S2 Fig

Criteria:
More than two clades with sequences above the cutoff of 4 sequences

(N =778) Sequences

(N = 727) Sequences

RAC-SK (N= 18)
Cosmopolitan (N=13)
Africa-3 (N=3)
Arctic (N=3)
Asian (N=3)
Africa-2 (N=2)
Bats (N=1)
Cat (N=43)

Cosmopolitan (N=22)
Arctic (N=1)
Goat(N=23)

Cosmopolitan (N=14)
Asian (N=13)
Arctic (N=8)
Africa-2 (N=6)
Indian-Sub (N=4)
Bats (N=2)
Human (N=47)

Dog (N= 539)
Cosmopolitan (N=383)
Asian (N=65)
Arctic (N=50)
Africa-2 (N=37)
Bats (N=3)
Indian-Sub (N=1)

Cattle (N=106)
Cosmopolitan (N=70)
Arctic (N=12)
RAC-SK (N=11)
Bats (N=7)
Asian (N=5)
Indian-Sub (N=1)

Horse(N=8)
RAC-SK (N=3)
Cosmopolitan (N=2)
Bats (N=1)
Asian (N=1)
Arctic (N=1)

Sheep (N=12)
Asian (N=7)
Cosmopolitan (N=4)
Arctic (N=1)