Mapping threatened Thai bovids provides opportunities for improved conservation
outcomes in Asia

Authors:
Wantida Horpiencharoen[1*], Renata L. Muylaert[1], Reju Sam John[1,2], Jonathan C. Marshall[1],
Antony J. Lynam[3], Alex Riggio[4], Alexander Godfrey[5], Dusit Ngoprasert[6], George A. Gale[6], Eric
Ash[7], Francesco Bisi[8], Giacomo Cremonesi[9], Gopalasamy Reuben Clements[10], Marnoch
Yindee[11], Nay Myo Shwe[12], Chanratana Pin[13], Thomas N. E. Gray[14], Saw Soe Aung[15], Seree
Nakbun[16], Stephanie Schuttler[17], Robert Steinmetz[18], Rungnapa Phoonjampa[18], Naret
Seuaturien[18], Worrapan Phumanee[18], David T. S. Hayman[1]

Affiliations:
[1] Molecular Epidemiology and Public Health Laboratory, Hopkirk Research Institute, Massey University,
Palmerston North 4472 New Zealand
[2] Department of Physics, Faculty of Science, University of Auckland, Auckland, New Zealand
[3] Centre for Global Conservation, Wildlife Conservation Society, New York, USA
[4] Mahidol University, Salaya, Phutthamonthon, Nakhon Pathom, Thailand
[5] Panthera, 8 West 40th Street 18th Floor New York, NY 10018 United States
[6] Conservation Ecology Program, King Mongkut's University of Technology Thonburi, Bangkok, Thailand
[7] Wildlife Conservation Research Unit, University of Oxford, UK
[8] Environmental Analysis and Management Unit, Guido Tosi Research Group, Department of Theoretical
and Applied Sciences, University of Insubria, Italy
[9] Istituto Oikos E.T.S. Via Crescenzago 1, 20134 Milan, Italy
[10] Department of Biological Sciences, School of Medical and Life Sciences, Sunway University, Selangor,
Malaysia
[11] Akkhraratchakumari Veterinary College, Walailak University, Nakhon Si Thammarat 80160, Thailand
[12] Friends of Wildlife, Yan-Aung Street (1), Building 296, Room 15 Quarter No.(2), Yankin Township,
Yangon, Myanmar
[13] Ministry of Environment, 48 Samdach Preah Sihanouk Blvd., Phnom Penh 12301, Cambodia
[14] WWF Tigers Alive Initiative, Phnom Penh, Cambodia
[15] Fauna & Flora International, Myanmar Programme, Bahan Township, Yangon, Myanmar
[16] Khaonampu Nature and Wildlife Education Center, Department of National Park, Wildlife and Plant
Conservation, Kanchanaburi, Thailand
[17] North Carolina Department of Natural and Cultural Resources, NC, USA
[18] WWF-Thailand, 9 Pisit Building, Pradiphat Road Soi 10, Phayathai, Bangkok 10400, Thailand

* Corresponding author Email: wantidah@gmail.com
35

Abstract

Wild bovids provide important ecosystem services throughout their native range. In Asia, most are threatened with extinction. Five wild bovids remain in Thailand: gaur (*Bos gaurus*), banteng (*Bos javanicus*), wild water buffalo (*Bubalus arnee*), mainland serow (*Capricornis sumatraensis*) and Chinese goral (*Naemorhedus griseus*). However, their populations and habitats have declined substantially and become fragmented. Here, we identified potentially suitable habitat for these five threatened bovids using ecological niche models, first throughout the species entire distribution and second within Thailand, and quantified how much suitable area remains within protected areas. We combined species occurrence data with 28 environmental variables for modelling and used a spatially-restricted Biotic-Abiotic-Mobility framework for two accessible areas: 1) species-specific accessible areas and 2) a single large accessible area. We applied spatially restricted and weighted average ensembles from eight algorithms when generating maps. For *B. gaurus* and *B. javanicus,* the best models predicted suitable habitat was mostly within Southeast Asia*,* with *B. gaurus* having predicted large areas in Thailand and India. *B. arnee* suitable habitat was mostly in India. *C. sumatraensis* suitable habitat was mostly in Thailand and Myanmar. *N. griseus* was mainly restricted to China. In Thailand, the highest bovid potential richnesses were in the Northern Forest, Western Forest, Eastern Forest and Dong Phayayen-Khao Yai Forest Complexes. We identified unprotected hotspots with >50% of overall suitable habitat located outside protected areas. *B. arnee* had the smallest proportion of protected habitat (9%). Suitable areas identified in and out protected areas may guide habitat management and conflict mitigation strategies.

Keywords: species distribution, herbivores, large mammals, wildlife conservation, habitat suitability, protected areas

1. Introduction

An important task of wildlife research and conservation is to define the distributional ecology of species, and to understand how they relate to the environment, climate and other organisms (Franklin, 2009). Ecological niche models (ENM) are applied to predict the geographic distribution that is suitable for a species by using ecological niche dimensions combined with species' presence data (Soberon & Peterson, 2005). ENM can be approached using the 'Biotic-Abiotic-Mobility' (BAM) framework which considers the relationship between the species' distribution, geographical and climatic factors and explains the influence of factors on predicted habitat suitability (Peterson & Soberón, 2012). Abiotic (A) factors generally determine the potential distribution (or fundamental niche) of a species, and the intersection of abiotic and biotic (B) factors form the realised niche, or the part of this potential distribution where species actually live (Soberón & Nakamura, 2009). Mobility (M) is the area that is accessible by species related to their distribution over periods of time (the 'accessible area'; Barve et al., 2011). Selecting the extent of species' accessible areas, including buffer zones, impacts model prediction results (Anderson & Raza, 2010; Barve et al., 2011).

Wild Bovidae (Mammalia: Artiodactyla) play significant ecological roles in tropical forests and grasslands (Hassanin, 2014). Bovids are grazers and browsers, modifying plant diversity and abundance within ecosystems (Ripple et al., 2015; Romero et al., 2015). Large wild bovids are also the prey of predators such as tigers (*Panthera tigris*) and leopards (*Panthera pardus*) (Simcharoen et al., 2018). Throughout Asia, wild bovid populations are threatened by poaching (Gray et al., 2018) and habitat loss (Nguyen, 2009), especially in South to Southeast Asia (Giam & Wilcove, 2012). Natural habitats have been disturbed by free-grazing livestock, which can lead to interbreeding (e.g. between domestic and wild water buffalo, Kaul et al., 2019), increased competition for food and natural resources (Bhandari et al., 2022), and increased risk of disease transmission between wildlife and livestock (Hassell et al., 2017).

In South and Southeast Asia, there are 27 recognised bovid species (IUCN, 2021), of which seven species are listed as vulnerable, five as endangered and three as critically endangered with extinction. Thailand has five bovids species (Gaur *Bos gaurus*, banteng *Bos javanicus*, wild water buffalo *Bubalus arnee*, mainland serow *Capricornis sumatraensis* and Chinese goral *Naemorhedus griseus*) are remaining in their natural habitat. These species are distributed in the other countries from South to Southeast Asia (Fig. 1) and also have different suitable habitats. For example, gaur can be found in evergreen forest or grassland and range from India, Nepal, across Southeast Asia to Peninsula Malaysia (Duckworth et al., 2016). *C. sumatraensis* also has a wide distribution from Nepal to Indonesia through hill forests to shrubland habitats (Phan et al., 2020). Nevertheless, evaluations of their habitat suitability in Thailand and the other

countries have been conducted only in some protected areas (Chaiyarat et al., 2019;
Pintana & Lakamavichian, 2013), but not at the regional or national level.

Understanding historical and present distributions of threatened species is fundamental
for planning and needs to be constructed at suitable scales, yet are typically lacking in
Asia. Regional scale modelling provides an overview of potential habitat and benefits
conservation planning (Catullo et al., 2008), such as by informing threatened species
reintroduction programmes (Bellis et al., 2021) and management through identifying
high-quality habitat connectivity and fragmentation (Crooks et al., 2011), along with
global biodiversity trend predictions (Araújo et al., 2019). Several studies have predicted
habitat suitability for some of these five wild bovids in local areas, including in Malaysia
(Ariffin et al., 2021), China (Ding et al., 2018), Indonesia (Rahman et al., 2019),
Thailand (Prayoon et al., 2021) and Nepal (Thapa et al., 2020), but habitat suitability
studies for larger extents across their distribution are lacking.

Here, we built ENM for five the Thai bovid species: *B. gaurus*, *B. javanicus*, *B. arnee*,
C.sumatraensis and *N. griseus* at two scales: first, at the regional scale throughout the
entire distribution and, second, at the country scale in Thailand. We aim to 1) identify
the potential distribution for these five species in South to Southeast Asia, and 2)
identify conservation areas in their geographical distribution, with a particular focus on
Thailand.

2. Materials and methods
Our workflow consisted of two main processes of data preparation and model building
(summarised in Supplementary Fig. S1) that generated habitat suitability maps for all
species and accessible areas used. Data preparation consisted of gathering the species
occurrence data and environmental data and selecting the accessible areas. Then, the
model building consisted of pre-processing, processing and post-processing steps.

2.1. Study area
The study area consists of 13 Asian countries: Bhutan, Bangladesh, Cambodia, China,
India, Indonesia, Laos, Malaysia, Myanmar, Nepal, Sri Lanka, Thailand and Vietnam
(Fig. 1), that cover the distribution of *Bos gaurus*, *Bos javanicus, Bubalus arnee*,
Capricornis sumatraensis, and *Naemorhedus griseus*, based on the literature (Table
S1).

Fig. 1. Species occurrence data (yellow circles), IUCN polygons (blue areas) and study areas (grey areas) used in model building for five wild bovid species. First, a common large 'accessible area' (A) was used for all species for model building, and then species-specific accessible areas (B-F) for individual species.

2.2. Species Occurrence data

We compiled species occurrence data from January 2000 to June 2021 from researchers, government, NGOs (World Wildlife Fund, Wildlife Conservation Society, Freeland, Panthera, Fauna & Flora International, Friends of Wildlife and RIMBA) and open data sources, including GBIF (https://www.gbif.org/) and eMammal (https://emammal.si.edu/). The data coverage by country can be found in Table S2. The occurrence data were collected through observation of animal signs (e.g. footprint and dung) during forest patrols, direct observation during wildlife surveys, camera trapping, and radio-collar signals (Table S3). We excluded occurrence records that fell outside the species-specific accessible area, duplicated records and museum collections.

2.3. Environmental variables

Hypothesised environmental variables were selected based on species' habitat and distribution related literature (Table S1). We used 28 variables (Table S4) for model construction, including 19 bioclimatic variables (average for 1970-2000) from WorldClim

v2 (Fick & Hijmans, 2017), elevation (Shuttle Radar Topography Mission-SRTM) from WorldClim (Fick & Hijmans, 2017), slope (Amatulli et al., 2020), five land cover fractions (grass, tree, urban, water and crop) (Buchhorn et al., 2019), human population density (Stevens et al., 2015) and greenness through the normalized difference vegetation index (NDVI) (Didan, 2015). All layers were processed using the geographic coordinates system (Datum WGS84) and ~1 km^2 spatial resolution. We transformed the human population density using logarithm base 10 to adjust for skewness. We rescaled the NDVI layer by multiplying all values with a scale factor (0.0001), based on the Moderate Resolution Imaging Spectroradiometer (MODIS) User's guidelines (Didan et al., 2015).

2.4. Accessible areas

The accessible area refers to the parts of the world that have been accessible to species via dispersal over time (Barve et al., 2011). The extent of the accessible area and the inclusion of a buffer zone has an important effect on ENM performance (Anderson & Raza, 2010; Barve et al., 2011). Therefore, we used two accessible area sizes to delimitate our modelling extent (Fig. 1), and compare ENMs with and without spatial restriction (MSDM). To reduce overprediction and make our predictions closer to realised niche estimates, we used an occurrences-based threshold (OBR) method with ensemble models from Mendes et al. (2020) for creating the spatially restricted ENM (hereon MSDM). OBR is an *a posteriori* method that restricts the suitable areas of our final ensemble models based on presence and the largest nearest neighbour distance among pairs of occurrences. The first larger accessible area (hereon LA) includes most of the Asian continent and its ecoregions, and all species distributions are included as a common extent. We downloaded current IUCN range maps for each species, then overlapped those on ecoregions (Olson et al., 2001), then combined the results with selected ecoregions based on biogeographic knowledge of the species distributions and habitat preference from the literature reviews. For example, gaur (*Bos gaurus*) habitat typically contains moist evergreen, semi-evergreen, and dry evergreen forests (Steinmetz et al., 2008; Tanasarnpaiboon, 2016), so we included these regions in our accessible areas. The second accessible area was more restricted and cropped based on individual species-specific distributions (hereon SSA) from literature reviews (Table S1), IUCN polygons or 'ranges' (IUCN, 2020) and the terrestrial ecoregions where they occur. Further details on ecoregions included in accessible areas are in Supplementary Table S5. Overall, we built four combinations between two accessible areas with and without MSDM methods for each species, including 1) No MSDM-SSA; 2) No MSDM-LA; 3) MSDM-SSA and 4) MSDM-LA.

2.5. Model building

We processed the species occurrence files and environmental datasets in R 4.0.1 (R Core Team, 2020). We developed reproducible ecological niche models with optimized

processing times using the ENMTML package (Andrade et al., 2020), following three
main steps: 1) pre-processing, 2) processing and 3) post-processing.

In pre-processing, we performed occurrence thinning using 2 times the cell-size (1 km^2)
(Velazco et al., 2019) to reduce clustering of species records and sampling bias. We
used principal component (PC) analysis (PCA) to reduce the collinearity of the
predictors. We assigned species' accessible areas to determine the species'
distributions using a mask function. We used random sampling to create pseudo-
absence background points in a 1:1 ratio with presence points (Barbet-Massin et al.,
2012). The occurrence and pseudo-absence data was divided into two sets for fitting
the model (75%) and evaluating the fitted models (25%), using the bootstrapping
partition method with 10 replications for each algorithm.

In the processing step, eight algorithms were used to build the ENMs, namely: Bioclims
(Booth et al., 2014), Generalized Linear Models (McCullagh & Nelder, 1989),
Generalized Additive Models (Hastie, 2017), Random Forest (Liaw & Wiener, 2002),
Support Vector Machine (Karatzoglou et al., 2004), Maximum Entropy default (Phillips
et al., 2006), Maximum Likelihood (Royle et al., 2012) and Bayesian Gaussian Process
(Golding, 2014). All models used the default settings from the ENMTML package, which
included the functions from different packages (e.g. dismo, maxnet) based on the
algorithms that use to fit the models. The data type used for each algorithm is in
Supplementary Table S6.

In the post-processing step, we created ensemble models using the weighted average
(WMEAN) method based on the True Skill Statistic (TSS) values for building final
habitat suitability and binary maps. The benefits of ensemble models are 1) robust
decision making (Ahmad et al., 2020); 2) reducing uncertainty (Marmion et al., 2009);
and 3) a combination of several models into one model prediction (Kindt, 2018). We
used TSS to calculate threshold values to convert habitat suitability maps into binary
suitability maps (0 = unsuitable and 1 = suitable). We used TSS and area under the
curve (AUC) for evaluating model performance. The TSS threshold is calculated using
the maximum summed specificity and sensitivity and is not based on prevalence, where
an equal TSS score for given models means similar performance (Allouche et al.,
2006). Therefore, we selected the final models from the best TSS of weighted average
ensemble models.

For calculating total suitable areas, we summed the suitable area of the best TSS binary
map models using the zonal function in the raster R package (Hijmans, 2021). Then, we
created a species richness map ranging from 0 (no suitable areas) to 5 (suitable areas
for all five species) by summing the best TSS binary maps of each species.

2.6. Protected area analyses

We used the binary (suitable and unsuitable) predictions from the best performing
model SSA and LA ensembles for calculating the suitable areas inside and outside
protected areas. The source for our protected areas map was the World Database of
Protected Area (WDPA) (UNEP-WCMC and IUCN, 2021). We classified protected
areas based on IUCN protected areas from WPDA into 8 categories including:
categories 1 to 6 as IUCN management categories I to VI; category 7 as 'not applicable'
which is included 'not reported', 'not applicable' and 'not assigned' protected areas; and
category 8 as non-protected areas, which are the remaining areas that have not been
classified as IUCN categories 1 to 7 (UNEP-WCMC and IUCN, 2021). Then, we used
the zonal function again to calculate overlapping areas between binary map predictions
and protected areas for each species. We used 1 km resolution and WGS 1984
projection for all the calculations.

We calculated the percentage of suitable habitat in WDPA polygons using the threshold
values from the best TSS binary suitability maps to separate suitable and unsuitable
areas. The areas that had suitability values higher than each species' threshold were
classified as suitable and those lower as unsuitable. We used the exact_extract function
in exactextractr package (Baston et al., 2021) for extracting the suitable areas (values
=1) from binary map rasters in each WDPA polygon. Then, we classified each PA into 5
different suitability categories, based on the percentage of suitable habitat in the PA:
low suitability (0 - 20%); low - medium suitability (>20 - 40%); medium suitability (>40 -
60%); high suitability (>60 - 80%) and very high suitability (≥80%), and selected only the
PAs that have the proportion of suitable area larger than species home range in the
result. We have provided the code for creating the models in a GitHub repository.

3. Results

We compiled 33,664 occurrence records (Table S3, TableS12). After filtering and
spatial thinning, we used 3,708 points for modelling: 2,732 for *B. gaurus*, 361 for *B.
javanicus*, 92 for *B. arnee*, 388 for *C. sumatraensis*, and 135 for *N. griseus.*

We found that the PCA reduced the 28 environmental variables into 12 PCs that
explained 95% of the environmental variance in the variables for the LA models for all
species. The PCs for SSA models explained more than 96% of the total variance and
the PC number varied by species, comprising 13 PCs (*B. arnee*), 11 PCs (*B. gaurus, C.
sumatraensis*), and 10 PCs (*B. javanicus, N. griseus*). The bioclimatic variables were
the important variables in all species models. For LA models, the first two axes (PC1
and PC2) have high contributions from the annual mean temperature (bio01), mean
temperature of the coldest month (bio06), mean temperature of the driest quarter
(bio09) and mean temperature of the warmest quarter (bio10). The first two axes of SSA
models showed high positive contributions from mean temperature of the coldest month
(*B. gaurus*), minimum temperature of coldest month (*B. javanicus, C. sumatraensis*),

annual mean temperature (*B. arnee, C. sumatraensis)*, and precipitation of the wettest
quarter (*N. griseus*). We also found that NDVI, elevation, slope and human population
density have less effect on explaining the variability for the first two PCs for all species.
The correlations between PCs and individual environmental variables, PC biplots and
percentage of explained variance are summarised in Table S7 and Fig. S2.

3.1 Ecological niche models

Overall, all ensemble models showed high performance both for TSS and the area
under the curve (AUC) with the highest performing models over 0.8 for all species
(Table S8). Models with species-specific accessible areas were not always the best
performing models, but most ensemble models performed above 0.7 TSS. The habitat
suitability prediction maps using the best model ensembles are in Supplementary Fig.
S3 (SSA) and Fig. S4 (LA), and the binary maps which were used for calculating the
suitable area in Supplementary Fig. S5. The performance of spatially restricted
ensembles was higher in comparison with the No MSDM models, as the TSS was
improved for *B. javanicus*, *N. griseus* and *B. arnee.* The lowest performing model for *B.*
arnee was the No MSDM-SSA (TSS = 0.57). The best model for *B. gaurus* was no
MSDM-LA, *B. javanicus* and *N. griseus* is MSDM-LA, *B. arnee* is MSDM-SSA, and *C.*
sumatraensis is No MSDM-SSA. We found that all species have small predicted
suitable habitats. Moreover, all species models predicted less than 50% of the suitable
areas inside PAs.

For *B. gaurus*, the best model predicted the total suitable area was 612,548 km^2, of
which Thailand has the largest area (29%; 177,216 km^2). *B. javanicus* has a similar
predicted distribution to *B. gaurus*, but is more restricted to Southeast Asia. The total
suitable area for *B. javanicus* predicted was 446,075 km^2 (MSDM-LA) and most of the
suitable areas were located in mainland Southeast Asia (96%, 429,759 km^2). *B. arnee*
has a patchy distribution of suitable habitat in the southern tip of India, Sri Lanka,
through the south of Thailand. The best model (MSDM-SSA) predicted the greatest
suitable area in India (35%, 213,247 km^2), followed by Thailand (135,552 km^2) and
Myanmar (99,844 km^2). *C. sumatraensis* suitable habitat is predicted throughout the
region modelled, down through the south of Thailand - Myanmar border forest patches
and in the southwestern parts of Cambodia in the species accessible area. The total
predicted suitable area for *C. sumatraensis* for the best performing model (no MSDM-
SSA) was around 290,000 km^2. Myanmar had the greatest suitable area of around 23%
(66,892 km^2). *N. griseus* suitable habitat predictions ranged through northeastern India,
Myanmar and northern Thailand to parts of China, with a total suitable area of 169,882
km^2 with the MSDM-LA model. China has the greatest suitable area, with 96% (162,929
km^2) of the total suitable area. Thailand has no suitable areas predicted by the best LA
models but around 4% (14,847 km^2 from 331,704 km^2) of the total suitable area was

predicted in Thailand by the best SSA models. Total of the suitable area in km^2 for each species and countries is shown in Fig. 2, and the prediction maps is in Fig. 3.

Fig. 2. Total of the suitable area in km^2 for each species and countries. Blue is the species-specific accessible area (SSA) and grey is the large accessible area models (LA) (see details in Supplementary Table S9).

Fig. 3. Habitat suitability prediction maps for five species (A-E) using the best model from either accessible areas and the weighted average ensemble. The value ranges from 0-1: yellow represents low suitability and dark brown represents high suitability. (F) presents the map of species numbers for five species using the binary suitability results, ranging from 0 (not suitable area for the five species) to 4 (suitable area for four species), which is the highest value range for the best model predictions. Interactive maps are provided in the supplementary material (link).

3.2. Identifying priority areas for conservation

3.2.1 Asia

Most suitable habitats in protected areas are located in IUCN category Ia (Strict nature reserve), Ib (Wilderness area) and II (National park) areas for the best TSS models for all species, while IUCN category V (Protected landscape or seascape) has the least. Overall, more than half of the species' suitable habitat is not under any form of protection defined by the WDPA (Supplementary Table S10, Fig. S6, Fig. S7). The proportion of the suitable area in each WDPA of the best models is shown in Fig. 4; and the proportion of the best models from SSA and LA for each species are presented in Supplementary Fig. S8 and Fig. S9.

Our results predicted *B. gaurus* has 27% (166,092 km^2) of the total suitable area inside PAs with the best model (No MSDM-LA). We identified 179 PAs (from 2132 PAs; Fig. 4) that had a high proportion of suitable area (≥80%). We identified several suitable PAs in the large connected landscape, both in Dawna-Tanintharyi Landscape (DTL) with approximately 21,000 km^2 (18 PAs) on the Thailand and Myanmar border, and in Cardamom Rainforest Landscape (CRL) in Cambodia, covering around 20,680 km^2 (12 PAs).

Our best models (MSDM-LA) predicted *B. javanicus'* suitable area to include 33% (145,790 km^2) of the total predicted suitable area inside PAs. From 2132 PAs evaluated (Fig. 4), 151 PAs had ≥80% of their area classified as suitable. We identified connected PAs that had greater than 80% of the total suitable areas in Cambodia's PAs (CRL, Eastern Plains Landscape), Thailand's PAs (Western, Dong Phayayen-Khao Yai and Eastern forest complexes) and areas along the border of Thailand (Panom Dong Rak WS), Cambodia (North West Corridor, Preah Vihear, Chhaeb WS) and Laos (non-PA).

Bubalus arnee had around 9% (55,378 km^2) of the total predicted suitable area inside PAs predicted with the best model (MSDM-SSA). We identified high habitat suitability in 146 PAs from 1304 PAs where suitable areas cover over 80% of the total PA and most suitable PAs were identified in Sri Lanka (128 PAs from 146 PAs, Fig. 4). India had the highest proportion of suitable area in PAs (≥80%), identified in Kaziranga NP and Manas WS.

For *C. sumatraensis*, our findings showed 41% (110,535 km^2) of the predicted suitable
area inside PAs by the best model (No MSDM-SSA). A high percentage of suitable area
(≥80%) was identified within 64 PAs (from 835 PAs, Fig. 4). Most suitable PAs are
found in Southeast Asia, including Thailand, Cambodia, Myanmar, Laos and Malaysia.
The connected suitable PAs with a high percentage of suitable area (≥80%) are DTL
and CRL. Other connected suitable areas were identified in Gunung Aais, Sungai Ketiar
and Taman Negara (Terengganu) NP in Peninsula Malaysia, and the part of Greater
Annamites complexes located in Laos and Cambodia.

Only 2% (3,904 km^2) of the suitable area predicted for *N. griseus* is currently protected
in a PA, and 12% were in 'not applicable' PAs, according to the best model (MSDM-LA).
Our best model had highly suitable areas in only 10 of 2132 PAs (Fig. 4). The highest
suitable areas were found in Jin Zhai Goa scenic area (75%) in China and Emae bum
NP (75%) in Myanmar, followed by Sichuan Giant Panda Sanctuaries (70%), Three
Parallel Rivers of Yunnan (60%) and Hkakaborazi National Park in Myanmar (40%).

Fig. 4. The proportion of predicted suitable areas within protected areas. Proportions
range from 0 (all unsuitable) to 1 (all suitable), with suitability determined by thresholds
from species best performing models. (A) Gaur (*Bos gaurus*), (B) Banteng (*Bos
javanicus*), (C) wild water buffalo (*Bubalus arnee*), (D) mainland serow (*Capricornis
sumatraensis*), and (E) Chinese goral (*Naemorhedus griseus*).

3.2.2. Thailand

We identified a high percentage (≥80%) of suitable area of Thailand for *B. gaurus* in 122 PAs (74,268 km^2; 15% of Thailand), *B. javanicus* in 102 PAs (59,528 km^2; 12% of Thailand), *C. sumatraensis* in 46 PAs (31,076 km^2; 6% of Thailand), *N. griseus* in 2 PAs (969 km^2; 0.19% of Thailand) and *B. arnee* in 3 PAs (559 km^2; 0.1 % of Thailand). A high proportion of suitable area for *B. gaurus* and *B. javanicus* is in Thungyai Naresuan, Kaengkrachan and Huai Kha Khaeng, for *C. sumatraensis* Ton Pariwat, Khlong Yan and Khlong Naka, for *B. arnee* Phu Wua and Dong Yai, and for *N. griseus* Mae Wa, Doi Wiang La and Mae Ping (Fig. 5 and 6). The hotspots for each species can be found in the supplementary material (Fig. S10). We found that the highest percentage of suitable area was mixed deciduous forest for all species, followed by evergreen forest for *B. gaurus, B. javanicus* and *C. sumatraensis,* and dry dipterocarp forest for *B. arnee* and montane forest for *N. griseus*. We found a percentage of non-forest areas identified from the total suitable for all species: *B. arnee* (71%), *B. javanicus* (33%), *B. gaurus* (24%), *C. sumatraensis* (10%) and Chinese goral (6%). For more details of forest types by suitable areas see Table S11 and Fig. S11.

Fig. 5 Habitat suitability for five species in Thailand. The value ranges from 0-1: yellow represents low suitability and dark brown represents high suitability. (F) presents the map of species numbers for five species using the binary suitability results.

Fig. 6. Estimated species richness of wild bovid in Thailand. Frames A-E focus on (A) Lum Nam Pai-Salawim, (B1) Mae Ping-Om koi and (B2) Srilanna-Khun Tan, (C) Western Forest Complex (WEFCOM), (D) Dong Phayayen-Khao Yai Forest Complex (DPKY) and (E) Eastern Forest Complex where the highest species numbers (n=4) and connected conservation areas are present.

Western, Dong Payayen-Khao Yai and Eastern forests have suitable areas for 3 – 4 species inside PAs and also in the surrounding areas. Overall, most protected areas had two to three species in the south, such as Kaeng Krachan and Khlong Saeng-Khao Sok forest complex. *N. griseus* suitable areas are restricted only to the northern forests (Lum Nam Pai-Salawim, Mae Ping-Om koi and Srilanna-Khun Tan), which make these PAs suitable for all species (Fig. 6 B1 and B2), but only in a small area compared to total PAs areas. The best model by TSS for *N. griseus* (MSDM-LA) showed no suitable areas in Thailand, though the SSA model did.

4. Discussion

We modelled the potential distribution for the five threatened wild bovid species, distributed in South and Southeast Asia, providing high resolution habitat suitability. Our aim was to build predictive models to identify conservation areas, and potential species richness maps in their entire geographical distribution and then with a particular focus on Thailand. Therefore, we used ensembles of models that make better predictions and achieve better performance than any single contributing model, reducing dispersion of

predictions. We identified that suitable areas were fragmented and often (50%) located
outside PAs. Those suitable areas outside PAs could possibly be managed as corridors
or buffer zones to connect currently fragmented bovid populations, thereby enhancing
long-term wild bovid conservation success (Karanth, 2016; Penjor et al., 2021), which
requires further investigation.

Our study found most suitable areas for gaur were similar to IUCN range assessments
(Duckworth et al., 2016) and consistent with studies that have confirmed species
presences, such as in Thailand's PAs (Prayoon et al., 2021), Myanmar (Hein et al.,
2020) and Western Ghats in southwestern India and Manas WS in the Himalayan
foothills (Choudhury, 2002). Our study predicted larger gaur suitable habitats in
Thailand inside (~82,400 km^2) and outside (95,000 km^2) PAs than Prayoon et al (2021),
who predicted 39,508 km^2 of total suitable habitat. Our predictions used NDVI and land
coverage fractions (Table S4) for predicting greenness, which may be useful for
predicting the vegetation quality and availability for ungulates (Borowik et al., 2013).
However, NDVI is difficult to differentiate vegetation variations (Didan et al., 2015;
Martinez & Labib, 2023), such as between specific agricultural areas, grassland, and
dense forest canopy. This may include other vegetation types other than the species'
habitat in suitable areas and estimated larger suitable areas predicted in non-forest
areas and non-PAs to be identified in our study, compared to Prayoon et al's study.
Other studies suggest *B. gaurus* does use crop plantations or man-made grasslands,
which may increase the suitable areas in our prediction, even if these are not their
natural habitats and lead to conflict between human and gaur (Chaiyarat et al., 2021).

Our best model predicted larger suitable areas (446,075 km^2) for *B. javanicus* than the
IUCN-SSC report released in 2010 (~ 209,000 km^2) (IUCN-SSC AWCS Group, 2010).
We found a high percentage of predicted suitable areas in Eastern Plains Landscape
(ELP) and Chhaeb WS in Cambodia; the former supports the likely largest *B. javanicus*
population globally (Gray et al., 2012). However, our results showed low habitat
suitability in Sundaic Southeast Asia, with just 2% of the total suitable area in Indonesia
(mainly in Alas Purwo NP, Java) and 2% of the total suitable area in Malaysia. *B.
javanicus* populations and habitats in Southeast Asian islands (Borneo, Kalimantan,
Java, and Bali) are threatened due to hunting for horn and meat consumption and
habitat loss (Dewi et al. 2020). In Thailand, we found high suitability similar to previous
studies in Eastern (Menkham et al., 2019) and Western forest complexes (Jornburom et
al., 2020), including reintroduction areas in Salak Pra WS (Chaiyarat et al., 2019) and
where recent recolonization by natural population movement has occurred in Mae Wong
NP (Phoonjampa et al., 2021).

Bubalus arnee has been domesticated and bred as livestock, making it hard to
distinguish between the free-grazing domestic buffalo and *B. arnee* as domesticated

animals may replace wild animals in suitable habitats and cause the high suitable area prediction outside PAs, especially in overlapping habitats (Zhang et al., 2020). We estimate the highest percentages of suitable areas at Kaziranga NP in India, currently with the largest population of wild water buffalo (Kaul et al., 2019). Grasslands and flood plain areas of Manas NP (500 km^2) and Kaziranga NP (>850 km^2) in India contain the most suitable habitat and are the main population strongholds for *B. arnee* (Choudhury, 2014). Moreover, we identified suitable areas for *B. arnee* in Huai Kha Kheang WS in Thailand, located close to streams and plains, but it is around 43% of Huai Kha Kheang WS. Thus, we recommend protecting these habitats to ensure the protection of wild cattle, such as active patrolling to reduce illegal intrusions, snare removal and habitat management based on their diet diversity (McShea et al., 2019).

We estimated suitable *C. sumatraensis* habitats in 12 countries, all consistent with IUCN polygons. However, our models predicted 293,883 km^2 of suitable area, substantially smaller than the IUCN polygons (around 3,500,000 km^2) (IUCN, 2021). We identified suitable *C. sumatraensis* and *B. gaurus* habitats in Dawna-Tanintharyi Landscape (DTL), Western Forest Complex, Kaeng Krachan and Kuiburi, which are transboundary parks connecting Myanmar and Thailand. These areas were identified as important conservation areas to build habitat connectivity for maintaining large felids, their prey, and other animal populations (Greenspan et al., 2021). However, our models predicted low suitability areas where *C. sumatraensis* has been recorded in Myanmar, such as Hkakaborazi NP and Hponkanrazi (Lwin et al., 2021; Rao et al., 2010). Forty-seven percent (185/388) of the data occurrences were collected from Thailand (Table S2), and only 13% were collected in Myanmar, so we suggest collecting more occurrence data from these PAs in other regions for validating and improving the model predictions.

We predicted *N. griseus* to have the lowest total suitable habitat area (around 170,000 km^2) among the five species, using its best model (MSDM-LA), with suitable areas restricted to China and Myanmar. This area is smaller than the IUCN polygon (Duckworth et al., 2008). We identified suitable habitat along the border of northern Myanmar (Hkakaborazi NP), China (Three Parallel Rivers of Yunnan PA), and part of the far-eastern Himalayan Landscapes (FHL), from which *N. griseus* as been observed in field surveys (Li et al., 2014). In Thailand, we predicted *N. griseus* may only be distributed in the northern part, with 13 PAs having medium to very high (40 - 90%) suitable areas for the MSDM-SSA models (Supplementary Fig. S8) at higher elevations, close to a recent study that reported the species in 11 PAs (Jangtarwan et al., 2020). Pintana & Lakamavichian (2013) report potential locations outside Qinling PAs, supporting our findings that more than 50% of suitable areas are found outside PAs.

We acknowledge sampling deficiencies across the regions. We had fewer occurrences
in Vietnam, Laos, Myanmar and Indonesia compared to Thailand, from which a large
number of our data points came (30,512 points in Thailand, 3,152 points outside
Thailand, Supplementary Table S2). Occurrence data based on data accessibility can
have sampling bias, particularly clustered points for *B. gaurus*, *B. javanicus*, and *C.*
sumatraensis. We minimised these biases through spatial thinning. Since we found
large amounts of the suitable area outside of Thailand, the imbalance has not likely
impacted our results, however, we suggest that future studies should focus on ensuring
monitoring of bovid populations in other countries, especially in India and Myanmar.
However, missing data has impacted some results. The model TSS values for
endangered *B. javanicus* and *N. griseus* are over 0.8, yet our models predict unsuitable
areas in part of Indonesia (east and central Kalimantan; Dewi et al. 2020) for *B.*
javanicus and China (e.g. Beijing and northeast Inner Mongolia; Yang et al. 2019) for *N.*
griseus from which these species have been reported. This would likely be improved if
more spatial data were available for these species.

In this study, we included all subspecies data points in our model ensembles as we aim
to extrapolate and predict the entire range of species' habitat suitability, but this may
increase uncertainty (Dormann, 2007). These five bovids have multiple subspecies,
including 3 subspecies of *B. gaurus* (Duckworth et al., 2016)*, B. javanicus* (Gardner et
al., 2016) *B. arnee* (Kaul et al., 2019) and *C. sumatraensis* (Mori et al., 2019), and 2
subspecies of *N. griseus* (Duckworth et al., 2008). Subspecies may vary in niche,
climate and biological interactions that could affect the model predictions. For example,
C. s. thar live in Himalayan areas, and our models predicted the only suitable area was
in Khaptad NP, whereas the Annapurna Conservation Areas, where the species has
been detected (Aryal, 2009), were unsuitable. The low habitat suitability of our study in
Borneo for *B. javanicus* could be because climatic and geographic conditions differ for
B.j. lowi and those in mainland Asia, affecting model transferability across different
regions (Zhu et al., 2021). Equally, Mori et al. (2019) suggest that *N. griseus* should be
reclassified as a single species with Himalayan (*N. bedfordi*) and Burmese goral (*N.*
evansi). Future analyses must consider these taxonomic reclassifications. However, we
modeled species level habitat suitability, rather than the subspecies, as we assume that
there is less likely to be habitat and environmental condition variation at the subspecies
level for these bovids (Smith et al., 2019).

We found that using the MSDM OBR technique made the predicted ecological niche
closer to a realised niche for species with more restricted ranges like *B. javanicus*, *B.*
arnee and *N. griseus*, with higher performance TSS values compared to No MSDM
models. We recommend restricting the accessible area for predicting *B. arnee* potential
habitat to reduce overprediction caused by overlapping areas with domestic water
buffalo. However, OBR can be sensitive to the distribution of occurrence data, because

it keeps predicted suitable areas close to the occurrence locations. This may lead to the
exclusion of potential suitable areas driven by a lack of occurrence data in those areas.
For example, the *B. arnee* No MSDM predicted potentially suitable habitat around the
Sre Pok Wildlife Sanctuary in Cambodia where the species is distributed (Gray et al.,
2012; IUCN, 2008), but after the MSDM, this potential habitat was excluded as we lack
occurrence data in Cambodia. Although our study showed slightly different TSS values
between two different accessible area extents, we encourage testing the different
accessible areas as it affects the model results (Anderson & Raza, 2010). Moreover,
model performance varied with accessible area sizes and spatial restrictions,
emphasising the need for careful accessible area definition in ecological modelling
(Barve et al., 2011). Still, we suggest further studies assess habitat quality and fine-
scale validation to assess high suitability PAs and their capacity to maintain populations.

5. Conclusion

Our study provided an overview of the suitable remaining habitat for threatened bovid
species at a regional scale using high-resolution environmental variables and species
occurrence data from multiple observation methods. Our predictions showed that the
suitable areas are small and fragmented for all five species, and more than 50% of
suitable areas are outside of protected areas. Those suitable areas outside PAs could
possibly become efficient conservation areas, such as forest corridors or buffer zones to
connect fragmented bovid populations and enhance long-term habitat conservation. Our
predictions may inform conservation actions to avoid further defaunation of wild bovidae
such as the management of human-wildlife conflicts and habitat quality for long-term
species survival.

Acknowledgements

We thank all the data contributions and collaborations from these institutions:

Dr Supagit Vinitpornsawan (Director of Wildlife Conservation Area Management and
Education Center, Wildlife Conservation Office, Department of National Parks, Wildlife
and Plant Conservation, Thailand) for animal track and sign data in Thailand. For
camera trap data in Thailand: Naret Seuaturien (WWF Thailand), Manoon
Pliosungnoen, Department of Environmental and Forest Biology, State University of
New York College of Environmental Science), Wanlop Chutipong (Conservation
Ecology Program, Pilot Plant Development and Training Institute, King Mongkut's
University of Technology Thonburi), Nucharin Songsasen (Smithsonian's National Zoo
and Conservation Biology Institute), Lon I. Grassman, Jr. (Feline Research Program,

Caesar Kleberg Wildlife Research Institute, Texas A&M University, USA), Freeland
Foundation, WCS Thailand, Smithsonian Institution, PANTHERA USA in Thailand.
WWF Thailand would like to thank: WWF Germany, WWF Sweden, WWF US,
B.Grimm, WWF Japan, and WWF Switzerland for wonderful support of field projects,
and Department of National Parks, Wildlife, and Plant Conservation for kind permission
and collaboration. WCS Cambodia, Friends of Wildlife, Wildlife Alliance and Ministry of
Environment, Royal Government of Cambodia for the camera trap data in Cambodia.
CarBi Project of WWF Lao and WWF Greater Mekong for the camera trap data in Laos.
Camera trap in Myanmar: Friends of Wildlife and Data collecting in Tanintharyi,
Myanmar was partially funded by the European Union, Helmsley Charitable Trust and
mainly funded by Integrated Tiger Habitat Conservation Project, through the Fauna &
Flora International (FFI). Open source databases: https://www.gbif.org, camera trap
data from eMammals (https://emammal.si.edu/) by William J. McShea (Conservation
Ecology Center, Smithsonian Conservation Biology Institute) and Megan Baker-Whatton
(Smithsonian Conservation Biology Institute, and George Mason University) under the
project: Qionglai Mountains Project, Liangshan Mountains Project, Habitat Connectivity
in Minshan Mountains, Qinling Project, HKK ForestGEO Project, and Carnivore
Intraguild Interactions in Select Thailand Reserves. Also, the other researchers who
share the species occurrence data with us. We thank the IUCN specialists for
commenting on the early version of the habitat suitability maps. The authors wish to
acknowledge the use of New Zealand eScience Infrastructure (NeSI:
https://www.nesi.org.nz) high-performance computing facilities, consulting support
and/or training services as part of this research.

577 References

Ahmad, S., Yang, L., Khan, T. U., Wanghe, K., Li, M., & Luan, X. (2020). Using an
ensemble modelling approach to predict the potential distribution of Himalayan gray
goral (*Naemorhedus goral bedfordi*) in Pakistan. *Global Ecology and Conservation,*
21, e00845. https://doi.org/10.1016/j.gecco.2019.e00845
Alves-Pinto, H., Geldmann, J., Jonas, H., Maioli, V., Balmford, A., Latawiec, A.E.,
Crouzeilles, R. and Strassburg, B., 2021. Opportunities and challenges of other
effective area-based conservation measures (OECMs) for biodiversity conservation.

Perspectives in Ecology and Conservation, 19(2), pp.115-120.
https://doi.org/10.1016/j.pecon.2021.01.004

Allouche, O., Tsoar, A., & Kadmon, R. (2006). Assessing the accuracy of species distribution models: prevalence, kappa and the true skill statistic (TSS). *Journal of Applied Ecology, 43*(6), 1223-1232. https://doi.org/10.1111/j.1365-2664.2006.01214.x

Amatulli, G., McInerney, D., Sethi, T., Strobl, P., & Domisch, S. (2020). Geomorpho90m, empirical evaluation and accuracy assessment of global high-resolution geomorphometric layers. *Scientific Data, 7*(1), 162. https://doi.org/10.1038/s41597-020-0479-6

Anderson, R. P., & Raza, A. (2010). The effect of the extent of the study region on GIS models of species geographic distributions and estimates of niche evolution: preliminary tests with montane rodents (genus Nephelomys) in Venezuela. *Journal of Biogeography, 37*(7), 1378-1393. https://doi.org/10.1111/j.1365-2699.2010.02290.x

Andrade, A. F. A. d., Velazco, S. J. E., & De Marco Júnior, P. (2020). ENMTML: An R package for a straightforward construction of complex ecological niche models. *Environmental Modelling & Software, 125*, 104615. https://doi.org/10.1016/j.envsoft.2019.104615

Ariffin, N., Mustapha, M., Taher, T., Khodri, N., Abdullah, N., & Nor, S. (2021). Potential habitat connectivity for Malayan gaur (*Bos gaurus*) in a fragmented forest area in Peninsular Malaysia. *Journal of Environmental Biology, 42*, 798-805.

Araújo, M. B., Anderson, R. P., Márcia Barbosa, A., Beale, C. M., Dormann, C. F., Early, R., Garcia, R. A., Guisan, A., Maiorano, L., & Naimi, B. (2019). Standards for distribution models in biodiversity assessments. *Science Advances, 5*(1), eaat4858.

Aryal, A. (2009). Habitat ecology of Himalayan serow (Capricornis sumatraensis ssp. *thar*) in Annapurna Conservation Area of Nepal. *Tiger paper, 34*(4), 12-20.

Barbet-Massin, M., Jiguet, F., Albert, C. H., & Thuiller, W. (2012). Selecting pseudo-absences for species distribution models: how, where and how many? *Methods in Ecology and Evolution, 3*(2), 327-338. https://doi.org/10.1111/j.2041-210X.2011.00172.x

Barve, N., Barve, V., Jiménez-Valverde, A., Lira-Noriega, A., Maher, S. P., Peterson, A. T., Soberón, J., & Villalobos, F. (2011). The crucial role of the accessible area in ecological niche modeling and species distribution modeling. *Ecological Modelling, 222*(11), 1810-1819.

Baston, D. (2021). exactextractr: Fast Extraction from Raster Datasets using Polygons. R package version 0.7.2. https://CRAN.R-project.org/package=exactextractr

Bellis, J., Longden, M., Styles, J., & Dalrymple, S. (2021). Using macroecological species distribution models to estimate changes in the suitability of sites for threatened species reintroduction. *Ecological Solutions and Evidence, 2*(1), e12050. https://doi.org/10.1002/2688-8319.12050

Bhandari, S., Crego, R. D., & Stabach, J. A. (2022). Spatial segregation between wild
ungulates and livestock outside protected areas in the lowlands of Nepal. *PLoS One*,
17(1), e0263122. https://doi.org/10.1371/journal.pone.0263122
Booth, T. H., Nix, H. A., Busby, J. R., & Hutchinson, M. F. (2014). bioclim: the first
species distribution modelling package, its early applications and relevance to most
current MaxEnt studies. *Diversity and distributions, 20*(1), 1-9.
https://doi.org/10.1111/ddi.12144
Borowik, T., Pettorelli, N., Sönnichsen, L., & Jędrzejewska, B. (2013). Normalized
difference vegetation index (NDVI) as a predictor of forage availability for ungulates
in forest and field habitats. *European Journal of Wildlife Research, 59*(5), 675-682.
https://doi.org/10.1007/s10344-013-0720-0
Buchhorn, M., Smets, B., Bertels, L., Lesiv, M., Tsendbazar, N.-E., Herold, M., & Fritz,
S. (2019). Copernicus Global Land Service: Land Cover 100m: collection 2: epoch
2015: Globe (Version V2.0.2). ESA Living Planet Symposium 2019 (LPS2019), Milan,
Italy: Zenodo.
Catullo, G., Masi, M., Falcucci, A., Maiorano, L., Rondinini, C., & Boitani, L. (2008). A
gap analysis of Southeast Asian mammals based on habitat suitability models.
Biological Conservation, 141(11), 2730-2744.
https://doi.org/10.1016/j.biocon.2008.08.019
Chaiyarat, R., Lauhachinda, V., Kutintara, U., Bhumpakphan, N., & Prayurasiddhi, T.
(2004). Population of wild water buffalo (*Bubalus bubalis*) in Huai Kha Khaeng
Wildlife Sanctuary, Thailand. *Natural History Bulletin of the Siam Society, 52*(2), 151-
162.
Chaiyarat, R., Prasopsin, S., & Bhumpakphan, N. (2021). Food and nutrition of Gaur
(*Bos gaurus* C.H. Smith, 1827) at the edge of Khao Yai National Park, Thailand.
Scientific Reports, 11(1), 3281. https://doi.org/10.1038/s41598-021-82858-1
Chaiyarat, R., Youngpoy, N., Kongsurakan, P., & Nakbun, S. (2019). Habitat
preferences of reintroduced banteng (*Bos javanicus*) into the Salakphra Wildlife
Sanctuary, Thailand. *Wildlife Research, 46*(7), 573-586.
https://doi.org/10.1071/WR18184
Choudhury, A. (2002). Distribution and conservation of the Gaur Bos gaurus in the
Indian Subcontinent. *Mammal Review, 32*(3), 199-226. https://doi.org/10.1046/j.1365-
2907.2002.00107.x
Choudhury, A. (2014). Wild water buffalo *Bubalus arnee* (Kerr, 1792). In J. Burton & M.
Melletti (Eds.), *Ecology, Evolution and Behaviour of Wild Cattle: Implications for
Conservation* (pp. 255-301). Cambridge University Press.
https://doi.org/10.1017/CBO9781139568098.018
Choudhury, A. (2022). Upward trend in numbers of the wild water buffalo (Bubalus
arnee). *Journal of the IUCN SSC Asian Wild Cattle Specialist Group* (7).

Conry, P. J. (1989). Gaur *Bos gaurus* and development in Malaysia. *Biological*
Conservation, 49(1), 47-65. https://doi.org/https://doi.org/10.1016/0006-
3207(89)90112-2

Crooks, K. R., Burdett, C. L., Theobald, D. M., Rondinini, C., & Boitani, L. (2011).
Global patterns of fragmentation and connectivity of mammalian carnivore habitat.
Philosophical transactions of the Royal Society of London. Series B, Biological
sciences, 366(1578), 2642-2651. https://doi.org/10.1098/rstb.2011.0120

De Alban, J. D. T., Leong, B. P. I., Venegas-Li, R., Connette, G. M., Jamaludin, J., Latt,
K. T., Oswald, P., Reeder, C., & Webb, E. L. (2021). Conservation beyond the
existing protected area network is required to improve species and habitat
representation in a global biodiversity hotspot. *Biological Conservation, 257*, 109105.
https://doi.org/10.1016/j.biocon.2021.109105

Dewi, M. C., Sapari, I., & Santoso, E. (2020). Threats And Conservation Efforts of the
Last Remaining Banteng (*Bos javanicus lowi*) in unprotected areas at Belantikan
HuLu, Central Kalimantan. *Media Konservasi, 25*(1), 64-72.

Didan, K. (2015). MOD13A2 MODIS/Terra Vegetation Indices 16-Day L3 Global 1km
SIN Grid V006. https://doi.org/10.5067/MODIS/MOD13A2.006

Didan, K., Munoz, A. B., Solano, R., & Huete, A. (2015). MODIS vegetation index
user's guide (MOD13 series). *University of Arizona: Vegetation Index and Phenology*
Lab.

Ding, C., Hu, Y., Li, C., & Jiang, Z. (2018). Distribution and habitat suitability
assessment of the gaur *Bos gaurus* in China. *Biodiversity Science, 26*(9), 951-961.

Dormann, C. F. (2007). Promising the future? Global change projections of species
distributions. *Basic and applied ecology, 8*(5), 387-397.

Duckworth, J. W., Steinmetz, R., & Chaiyarat, R. (2008). *Naemorhedus griseus*. In
IUCN 2012. IUCN red list of threatened species. Version 2012.2.

Duckworth, J. W., Sankar, K., Williams, A. C., Samba Kumar, N., & Timmins, R. J.
(2016). *Bos gaurus. The IUCN Red List of Threatened Species 2016:*
e.T2891A46363646. https://dx.doi.org/10.2305/IUCN.UK.2016-
2.RLTS.T2891A46363646.en. Accessed on 11 February 2021

eMammal. (2021). Available from: https://emammal.si.edu/. Accessed on 21 June 2021

Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: new 1-km spatial resolution climate
surfaces for global land areas. *International Journal of Climatology, 37*(12), 4302-
4315. https://doi.org/10.1002/joc.5086

Franklin, J. (2009). *Mapping species distributions: spatial inference and prediction*.
Cambridge University Press.

Gardner, P., Hedges, S., Pudyatmoko, S., Gray, T. N. E., & Timmins, R. J. (2016). *Bos*
javanicus. The IUCN Red List of Threatened Species 2016: e.T2888A46362970.
https://dx.doi.org/10.2305/IUCN.UK.2016-2.RLTS.T2888A46362970.en.

GBIF.org (25 September 2020) GBIF Occurrence Download (*Capricornis
milneedwardsii*)https://doi.org/10.15468/dl.gnsbut
GBIF.org (26 September 2020) GBIF Occurrence (*Bos javanicus*) Download
https://doi.org/10.15468/dl.4znm7y
GBIF.org (14 October 2020) GBIF Occurrence (*Bubalus arnee*)
Download https://doi.org/10.15468/dl.5dek9t
GBIF.org (16 October 2020) GBIF Occurrence (*Naemorhedus griseus*) Download
https://doi.org/10.15468/dl.vb9cn7
GBIF.org (02 November 2020) GBIF Occurrence (C*apricornis thar*) Download
https://doi.org/10.15468/dl.bw5gdb
GBIF.org (1 December 2020) GBIF Occurrence (*Bos gaurus*) Download
https://doi.org/10.15468/dl.kwkw6q
GBIF.org (02 December 2020) GBIF Occurrence (*Capricornis sumatraensis*) Download
https://doi.org/10.15468/dl.ra6d3f
Giam, X., & Wilcove, D. S. (2012). The geography of conservation ecology research in
Southeast Asia: current biases and future opportunities. *Raffles Bull. Zool, 25*, 29-36.
Giri, S., Aryal, A., Koirala, R., Adhikari, B., & Raubenheimer, D. (2011). Feeding
ecology and distribution of Himalayan serow (*Capricornis thar*) in Annapurna
Conservation Area, Nepal. *World Journal of Zoology, 6*(1), 80-85.
Golding, N. (2014). GRaF: Species distribution modelling using latent Gaussian
random fields. *R package version 0.1-12*.
Gray, T. N. E., Prum, S., Pin, C., & Phan, C. (2012). Distance sampling reveals
Cambodia's Eastern Plains Landscape supports the largest global population of the
Endangered banteng *Bos javanicus. Oryx, 46*(4), 563-566.
https://doi.org/10.1017/S0030605312000567
Gray, T. N. E., Hughes, A., Laurance, W., Long, B., Lynam, A., O'Kelly, H., Ripple, W.,
Seng, T., Scotson, L., & Wilkinson, N. (2018). The wildlife snaring crisis: an insidious
and pervasive threat to biodiversity in Southeast Asia. *Biodiversity and Conservation,
27*. https://doi.org/10.1007/s10531-017-1450-5
Greenspan, E., Montgomery, C., Stokes, D., Wantai, S., & Moo, S. S. B. (2021). Large
felid habitat connectivity in the transboundary Dawna-Tanintharyi landscape of
Myanmar and Thailand. *Landscape Ecology, 36*(11), 3187-3205.
https://doi.org/10.1007/s10980-021-01316-5
Hassanin, A. (2014). Systematic and evolution of Bovini. *Ecology, Evolution and
Behavior of Wild Cattle: Implications for Conservation. Cambridge University Press,
Cambridge*, 7-20. https://doi.org/https://doi.org/10.1017/CBO9781139568098.003
Hassell, J. M., Begon, M., Ward, M. J., & Fèvre, E. M. (2017). Urbanization and
Disease Emergence: Dynamics at the Wildlife–Livestock–Human Interface. *Trends in
ecology & evolution, 32*(1), 55-67. https:/doi.org/10.1016/j.tree.2016.09.012

Hastie, T. J. (2017). Generalized additive models. In *Statistical models in S* (pp. 249-
307): Routledge.
Herrera-Sánchez, F. J., Gil-Sánchez, J. M., Álvarez, B., Cancio, I., de Lucas, J.,
Arredondo, Á., Díaz-Portero, M. Á., Rodríguez-Siles, J., Sáez, J. M., Pérez, J.,
McCain, E., Qninba, A., & Abáigar, T. (2020). Identifying priority conservation areas
in a Saharan environment by highlighting the endangered Cuvier's Gazelle as a
flagship species. *Scientific Reports, 10*(1), 8241. https://doi.org/10.1038/s41598-020-
65188-6
Hein, Z. M., Williams, A. C., Soe, P., Cox, N. J., Htun, N. Z., Thaung, N. O., Aye, Y. Y.,
Htun, Y. L., & Yoganand, K. (2020). Status of two species of threatened wild cattle
(Bos gaurus and Bos javanicus birmanicus) in North Zamari Wildlife Sanctuary, Bago
Region, Myanmar. *BULLetin*.
Hijmans, R. J. (2021). raster: Geographic Data Analysis and Modeling. R package
version 3.5-11. https://CRAN.R-project.org/package=raster
IUCN-SSC Asian Wild Cattle Specialist Group 2010. (2010). Regional Conservation
Strategy for Wild Cattle and Buffaloes in South-east Asia
IUCN. 2020. The IUCN Red List of Threatened Species Version 2020-3.
https://www.iucnredlist.org. Accessed on March 2020.
IUCN. 2021. The IUCN Red List of Threatened Species Version 2021-1.
https://www.iucnredlist.org. Accessed on April 2021.
Jangtarwan, K., Kamsongkram, P., Subpayakom, N., Sillapaprayoon, S., Muangmai,
N., Kongphoemph, A., Wongsodchuen, A., Intapan, S., Chamchumroon, W.,
Safoowong, M., Peyachoknagul, S., Duengkae, P., & Srikulnath, K. (2020). Predictive
genetic plan for a captive population of the Chinese goral (*Naemorhedus griseus*)
and prescriptive action for ex situ and in situ conservation management in Thailand.
PLoS One, 15(6), e0234064. https://doi.org/10.1371/journal.pone.0234064
Janžekovič, F., & Novak, T. (2012). PCA–a powerful method for analyze ecological
niches. *Principal component analysis-multidisciplinary applications, 29*, 127-142.
Jornburom, P., Duangchantrasiri, S., Jinamoy, S., Pattanavibool, A., Hines, J. E.,
Arnold, T. W., Fieberg, J., & Smith, J. L. D. (2020). Habitat use by tiger prey in
Thailand's Western Forest Complex: What will it take to fill a half-full tiger landscape?
Journal for Nature Conservation, 58, 125896.
https://doi.org/https://doi.org/10.1016/j.jnc.2020.125896
Karanth, K. K. (2016). Wildlife in the matrix: spatio-temporal patterns of herbivore
occurrence in Karnataka, India. *Environmental management, 57*(1), 189-206.
Karatzoglou, A., Smola, A., Hornik, K., & Zeileis, A. (2004). kernlab-an S4 package for
kernel methods in R. *Journal of statistical software, 11*(9), 1-20.
https://doi.org/10.18637/jss.v011.i09
Kaul, R., Williams, A. C., rithe, k., Steinmetz, R., & Mishra, R. (2019). *Bubalus arnee.*
The IUCN Red List of Threatened Species 2019: e.T3129A46364616.

https://dx.doi.org/10.2305/IUCN.UK.2019-1.RLTS.T3129A46364616.en. Accessed on 28 April 2020.

Kindt, R. (2018). Ensemble species distribution modelling with transformed suitability values. *Environmental Modelling & Software, 100*, 136-145. https://doi.org/10.1016/j.envsoft.2017.11.009.

Li, X., Buzzard, P., & Jiang, X. (2014). Habitat associations of four ungulates in mountain forests of southwest China, based on camera trapping and dung counts data. *Population Ecology, 56*(1), 251-256. https://doi.org/10.1007/s10144-013-0405-2

Liaw, A., & Wiener, M. (2002). Classification and regression by randomForest. *R news, 2*(3), 18-22.

Lim, H. Y., Gardner, P. C., Abram, N. K., Yusah, K. M., & Goossens, B. (2021). Identifying habitat and understanding movement resistance for the Endangered Bornean banteng Bos javanicus lowi in Sabah, Malaysia. *Oryx*, *55*(1), 122-130.

Lwin, Y. H., Wang, L., Li, G., Maung, K. W., Swa, K., & Quan, R.-C. (2021). Diversity, distribution and conservation of large mammals in northern Myanmar. *Global Ecology and Conservation, 29*, e01736. https://doi.org/10.1016/j.gecco.2021.e01736

Marmion, M., Parviainen, M., Luoto, M., Heikkinen, R. K., & Thuiller, W. (2009). Evaluation of consensus methods in predictive species distribution modelling. *Diversity and distributions, 15*(1), 59-69. https://doi.org/10.1111/j.1472-4642.2008.00491.x

Martinez, A. d. l. I., & Labib, S. (2023). Demystifying normalized difference vegetation index (NDVI) for greenness exposure assessments and policy interventions in urban greening. *Environmental Research, 220*, 115155. https://doi.org/10.1016/j.envres.2022.115155

McCullagh, P., & Nelder, J. A. (1989). *Generalized linear models.* (2nd ed.). chapman and hall. London, UK.

McShea, W. J., Sukmasuang, R., Erickson, D. L., Herrmann, V., Ngoprasert, D., Bhumpakphan, N., & Davies, S. J. (2019). Metabarcoding reveals diet diversity in an ungulate community in Thailand. *Biotropica*, *51*(6), 923-937. https://doi.org/10.1111/btp.12720

Mendes, P., Velazco, S. J. E., Andrade, A. F. A. d., & De Marco, P. (2020). Dealing with overprediction in species distribution models: How adding distance constraints can improve model accuracy. *Ecological Modelling, 431*, 109180. https://doi.org/10.1016/j.ecolmodel.2020.109180

Menkham, K., Sukmasuang, R., Pla-ard, M., Charaspet, K., Panganta, T., Trisurat, Y., & Bhumpakphan, N. (2019). Population and habitat use of Asian elephants (*Elephas maximus*) and five ungulate species in Khao Ang Rue Nai Wildlife Sanctuary, Chachoengsao Province, Thailand. *Biodiversitas Journal of Biological Diversity*, *20*(8).

Mori, E., Nerva, L., & Lovari, S. (2019). Reclassification of the serows and gorals: the end of a neverending story?. *Mammal Review*, *49*(3), 256-262. https://doi.org/10.1111/mam.12154

Nguyen, M. H. (2009). The status of Vulnerable gaur Bos gaurus and Endangered banteng *Bos javanicus* in Ea So Nature Reserve and Yok Don and Cat Tien National Parks, Vietnam. *Oryx, 43*(1), 129-135. https://doi.org/10.1017/S0030605307000440

Olson, D. M., Dinerstein, E., Wikramanayake, E. D., Burgess, N. D., Powell, G. V. N., Underwood, E. C., D'amico, J. A., Itoua, I., Strand, H. E., Morrison, J. C., Loucks, C. J., Allnutt, T. F., Ricketts, T. H., Kura, Y., Lamoreux, J. F., Wettengel, W. W., Hedao, P., & Kassem, K. R. (2001). Terrestrial Ecoregions of the World: A New Map of Life on Earth: A new global map of terrestrial ecoregions provides an innovative tool for conserving biodiversity. *BioScience, 51*(11), 933-938. https://doi.org/10.1641/0006-3568(2001)051[0933:Teotwa]2.0.Co;2

Penjor, U., Kaszta, Ż., Macdonald, D. W., & Cushman, S. A. (2021). Prioritizing areas for conservation outside the existing protected area network in Bhutan: the use of multi-species, multi-scale habitat suitability models. *Landscape Ecology, 36*(5), 1281-1309. https://doi.org/10.1007/s10980-021-01225-7

Peterson, A. T., & Holt, R. D. (2003). Niche differentiation in Mexican birds: using point occurrences to detect ecological innovation. *Ecology Letters, 6*(8), 774-782. https://doi.org/10.1046/j.1461-0248.2003.00502.x

Peterson, A. T., & Soberón, J. (2012). Species distribution modeling and ecological niche modeling: getting the concepts right. *Natureza & Conservação, 10*(2), 102-107. https://doi.org/10.4322/natcon.2012.019

Phan, T.D., Nijhawan, S., Li, S. & Xiao, L. 2020. *Capricornis sumatraensis*. The IUCN Red List of Threatened Species 2020: e.T162916735A162916910. https://dx.doi.org/10.2305/IUCN.UK.2020-2.RLTS.T162916735A162916910.en. Accessed on 26 October 2020.

Phillips, S. J., Anderson, R. P., & Schapire, R. E. (2006). Maximum entropy modeling of species geographic distributions. *Ecological Modelling, 190*(3-4), 231-259.

Phoonjampa, R., Steinmetz, R., Phumanee, W., Bunchornratana, K., Kaewsrisod, T., Srirattanaporn, S., Taraphibarl, K., Bejraburnin, T., & Bhumpakphan, N. (2021). Recolonization of Former Range by Endangered Banteng Bos j avanicus in Mae Wong National Park, Thailand. *Tropical Conservation Science, 14*, 19400829211065359.

Pintana, D., & Lakamavichian, S. (2013). Ecology and conservation of Goral outside protected areas by community-based approach in Thailand. *Tigerpaper, 40*(3), 12-21.

Prayoon, U., Suksavate, W., Chaiyes, A., Winitpornsawan, S., Tunhikorn, S., Faengbubpha, K., Angkaew, C., Pattanakiat, S., & Duengkae, P. (2021). Past, present and future habitat suitable for gaur (*Bos gaurus*) in Thailand. *Agriculture and Natural Resources, 55*(5), 743–756-743–756.

R Core Team. (2020). R: A language and environment for statistical computing. In R
Foundation for Statistical Computing.

Rahman, D. A., Herliansyah, R., Rianti, P., Rahmat, U. M., Firdaus, A. Y., &
Syamsudin, M. (2019). Ecology and conservation of the Endangered banteng (*Bos
javanicus*) in Indonesia tropical lowland forest. *HAYATI Journal of Biosciences, 26*(2),
68-68.

Rahman, H. A., McCarthy, K. P., McCarthy, J. L., & Faisal, M. M. (2021). Application of
Multi-Species Occupancy Modeling to assess mammal diversity in northeast
Bangladesh. *Global Ecology and Conservation, 25*, e01385.
https://doi.org/10.1016/j.gecco.2020.e01385

Rao, M., Htun, S., Zaw, T., & Myint, T. (2010). Hunting, Livelihoods and Declining
Wildlife in the Hponkanrazi Wildlife Sanctuary, North Myanmar. *Environmental
management, 46*(2), 143-153. https://doi.org/10.1007/s00267-010-9519-x

Ripple, W. J., Newsome, T. M., Wolf, C., Dirzo, R., Everatt, K. T., Galetti, M., Hayward,
M. W., Kerley, G. I. H., Levi, T., Lindsey, P. A., Macdonald, D. W., Malhi, Y., Painter,
L. E., Sandom, C. J., Terborgh, J., & Van Valkenburgh, B. (2015). Collapse of the
world's largest herbivores. *Science Advances, 1*(4), e1400103-e1400103.
https://doi.org/10.1126/sciadv.1400103

Romero, G. Q., Gonçalves-Souza, T., Vieira, C., & Koricheva, J. (2015). Ecosystem
engineering effects on species diversity across ecosystems: a meta-analysis.
Biological Reviews, 90(3), 877-890. https://doi.org/10.1111/brv.12138

Royle, J. A., Chandler, R. B., Yackulic, C., & Nichols, J. D. (2012). Likelihood analysis
of species occurrence probability from presence-only data for modelling species
distributions. *Methods in Ecology and Evolution, 3*(3), 545-554.
https://doi.org/10.1111/j.2041-210X.2011.00182.x

Shakya, B., Uddin, K., Yi, S., Bhatta, L. D., Lodhi, M. S., Htun, N. Z., & Yang, Y. (2021).
Mapping of the ecosystem services flow from three protected areas in the far-eastern
Himalayan Landscape: An impetus to regional cooperation. *Ecosystem Services, 47*,
101222. https://doi.org/10.1016/j.ecoser.2020.101222

Simcharoen, A., Simcharoen, S., Duangchantrasiri, S., Bump, J., & Smith, J. L. D.
(2018). Tiger and leopard diets in western Thailand: Evidence for overlap and
potential consequences. *Food Webs, 15*, e00085.
https://doi.org/10.1016/j.fooweb.2018.e00085

Smith, A. B., Godsoe, W., Rodríguez-Sánchez, F., Wang, H. H., & Warren, D. (2019).
Niche estimation above and below the species level. *Trends in Ecology & Evolution*,
34(3), 260-273.

Soberón, J., & Nakamura, M. (2009). Niches and distributional areas: Concepts,
methods, and assumptions. *Proceedings of the National Academy of Sciences,
106*(Supplement 2), 19644. https://doi.org/10.1073/pnas.0901637106

Soberon, J., & Peterson, A. T. (2005). Interpretation of models of fundamental
ecological niches and species' distributional areas. *Biodiversity Informatics*, 2.
https://doi.org/10.17161/bi.v2i0.4
Steinmetz, R., Chutipong, W., Seuaturien, N., & Chirngsaard, E. (2008). Community
Structure of Large Mammals in Tropical Montane and Lowland Forest in the
Tenasserim-Dawna Mountains, Thailand. *Biotropica, 40*(3), 344-353.
www.jstor.org/stable/30043637
Stevens, F. R., Gaughan, A. E., Linard, C., & Tatem, A. J. (2015). Disaggregating
Census Data for Population Mapping Using Random Forests with Remotely-Sensed
and Ancillary Data. *PLoS One, 10*(2), e0107042.
https://doi.org/10.1371/journal.pone.0107042
Tanasarnpaiboon, S. (2016). *Gaur (Bos gaurus)* abundance, distribution, and habitat
use patterns in Kuiburi National Park, Southwestern Thailand [Doctoral dissertation,
University of Nebraska-Lincoln].
https://digitalcommons.unl.edu/dissertations/AAI10102227
Thapa, R., Neupane, B., Ranabhat, S., Poudel, M., & Panthi, S. (2020). Habitat
suitability of wild water buffalo (*Bubalus arnee*) in Babai flood plain of Bardia National
Park, Nepal. *Global Ecology and Conservation, 23*, e01172.
https://doi.org/10.1016/j.gecco.2020.e01172
UNEP-WCMC and IUCN. (2021). *Protected Planet: The World Database on Protected
Areas (WDPA)*. www.protectedplanet.net. Accessed on 01 May 2021
Velazco, S. J. E., Villalobos, F., Galvão, F., & De Marco Júnior, P. (2019). A dark
scenario for Cerrado plant species: Effects of future climate, land use and protected
areas ineffectiveness. *Diversity and distributions, 25*(4), 660-673.
https://doi.org/10.1111/ddi.12886
Yang, J., Zhu, G. F., Jiang, J., Xiang, C. L., Gao, F. L., & Bao, W. D. (2019). Non-
invasive genetic analysis indicates low population connectivity in vulnerable Chinese
gorals: concerns for segregated population management. *Zoological Research*,
40(5), 439.
Zhang, Y., Colli, L., & Barker, J. (2020). Asian water buffalo: domestication, history and
genetics. *Animal Genetics, 51*, 177-191. https://doi.org/10.1111/age.12911
Zhu, G., Fan, J., & Peterson, A. T. (2021). Cautions in weighting individual ecological
niche models in ensemble forecasting. *Ecological Modelling, 448*, 109502.
https://doi.org/10.1016/j.ecolmodel.2021.109502