**Mapping threatened Thai bovids provides opportunities for improved**
**conservation outcomes in Asia**

Authors:
Wantida Horpiencharoen[1*], Renata L. Muylaert[1] Jonathan C. Marshall[1], Reju Sam John[1,2], Antony J.
Lynam[3], Alex Riggio[4], Alexander Godfrey[5], Dusit Ngoprasert[6], George A. Gale[6], Eric Ash[7], Francesco
Bisi[8], Giacomo Cremonesi[8], Gopalasamy Reuben Clements[9], Marnoch Yindee[10], Nay Myo Shwe[11],
Chanratana Pin[12], Thomas N. E. Gray[13], Saw Soe Aung[14], Seree Nakbun[15], Stephanie G. Manka[16], Robert
Steinmetz[17], Rungnapa Phoonjampa[17], Naret Seuaturien[17], Worrapan Phumanee[17], David T. S.
Hayman[1]

Affiliations:
[1] Molecular Epidemiology and Public Health Laboratory, Hopkirk Research Institute, Massey University,
Palmerston North 4472 New Zealand
[2] Department of Physics, Faculty of Science, University of Auckland, Auckland, New Zealand
[3] Centre for Global Conservation, Wildlife Conservation Society, New York, USA
[4] Mahidol University, Salaya, Phutthamonthon, Nakhon Pathom, Thailand
[5] Panthera, 8 West 40th Street 18th Floor New York, NY 10018 United States
[6] Conservation Ecology Program, King Mongkut's University of Technology Thonburi, Bangkok, Thailand
[7] Wildlife Conservation Research Unit, University of Oxford, UK
[8] Istituto Oikos E.T.S., Milan, Italy
[9] Zoological Society of London, Regents Park, London, UK
[10] Akkhraratchakumari Veterinary College, Walailak University, Nakhon Si Thammarat 80160, Thailand
[11] Friends of Wildlife, Yan-Aung Street (1), Building 296, Room 15 Quarter No.(2), Yankin Township, Yangon,
Myanmar
[12] Ministry of Environment, 48 Samdach Preah Sihanouk Blvd., Phnom Penh 12301, Cambodia
[13] WWF Tigers Alive Initiative, Phnom Penh, Cambodia
[14] Fauna & Flora International, Myanmar Programme, Bahan Township, Yangon, Myanmar
[15] Khaonampu Nature and Wildlife Education Center, Department of National Park, Wildlife and Plant
Conservation, Kanchanaburi, Thailand
[16] Fancy Scientist LLC
[17] WWF-Thailand, 9 Pisit Building, Pradiphat Road Soi 10, Phayathai, Bangkok 10400, Thailand

* Corresponding author Email: wantidah@gmail.com

Keywords: species distribution, herbivores, large mammals, wildlife conservation, habitat
suitability, protected areas

**Abstract**

Wild bovids provide important ecosystem functions throughout their ranges. Five wild bovids remain in Thailand: gaur (*Bos gaurus*), banteng (*Bos javanicus*), wild water buffalo (Bubalus arnee), mainland serow (*Capricornis sumatraensis*) and Chinese goral (*Naemorhedus griseus*). However, their populations and habitats have declined substantially and become fragmented. Here, we identify potentially suitable habitat for these threatened bovids using ecological niche models and quantify how much suitable area remains within protected areas. We combined species occurrence data with environmental variables and used spatially-restricted Biotic-Abiotic-Mobility frameworks with species-specific and single large accessible areas. We used ensembles from eight algorithms for generating maps and out-of-sample predictions to validate model performance against new data. Gaur, banteng, and buffalo models performed well throughout the entire distribution (≥62%) and in Thailand (≥80%). Mainland serow and Chinese goral performed poorly for the entire distribution and in Thailand, though a 5 km movement buffer markedly improved model performance for serow. Particularly large suitable areas were in Thailand and India for gaur, Cambodia and Thailand for banteng, and India for buffalo. Over 50% of overall suitable habitat is located outside protected areas, with just 9% for buffalo in Thai protected areas, highlighting area for potential habitat management and conflict mitigation.

**Introduction**

An important task of wildlife research and conservation is to define the distributional ecology of species and to understand how they relate to the environment, climate and other organisms (Franklin, 2009). Ecological niche models (ENM) are applied to predict the geographic distribution suitable for a species by using ecological niche dimensions combined with species' presence data (Soberon & Peterson, 2005). ENM can be approached using the 'Biotic-Abiotic-Mobility' (BAM) framework, which considers the relationship between the species' distribution, geographical and climatic factors and explains the influence of factors on predicted habitat suitability (Peterson & Soberón, 2012). Abiotic (A) factors generally determine the potential distribution (or fundamental niche) of a species, and the intersection of abiotic and biotic (B) factors form the realised niche, or the part of this potential distribution where species actually live (Soberón & Nakamura, 2009). Mobility (M) is the area accessible by species related to their distribution over periods of time (the 'accessible area'; (Barve et al., 2011)). Selecting the extent of species' accessible areas, including buffer zones, impacts model prediction results (Anderson & Raza, 2010; Barve et al., 2011).

Wild Bovidae (Mammalia: Artiodactyla) play significant ecological roles in tropical forests and grasslands (Hassanin, 2014). Bovids are grazers and browsers, modifying plant diversity and abundance within ecosystems (Ripple et al., 2015; Romero et al., 2015). Large wild bovids are also the prey of predators such as tigers (*Panthera tigris*) and leopards (*Panthera pardus*) (Simcharoen et al., 2018). Throughout Asia, wild bovid populations are threatened by poaching (Gray et al., 2018) and habitat loss (Nguyen, 2009), especially in South

to Southeast Asia (Giam & Wilcove, 2012). Natural habitats have been disturbed by free-
grazing livestock, which can lead to interbreeding (e.g. between domestic and wild water
buffalo, (Kaul et al., 2019), increased competition for food and natural resources (Bhandari et
al., 2022), and increased risk of disease transmission between wildlife and livestock (Hassell
et al., 2017). Moreover, habitat destruction is likely to influence the species' distribution and
behaviour adaptation, which could lead to shared natural resources and conflict between
humans and wild bovids.

In South and Southeast Asia, there are 27 recognised bovid species (IUCN, 2021), of
which seven species are listed as vulnerable, five as endangered and three as critically
endangered with extinction. Thailand has five bovid species (gaur; *Bos gaurus*, banteng; *Bos
javanicus*, wild water buffalo; *Bubalus arnee*, mainland serow; *Capricornis sumatraensis* and
Chinese goral; *Naemorhedus griseus*) remaining in their natural habitat. These species are
distributed in other countries from South to Southeast Asia (Figure 1) and also have different
suitable habitats. For example, gaur can be found in evergreen forest or grassland and range
from India, Nepal, across Southeast Asia to Peninsula Malaysia (Duckworth et al., 2016).
Mainland serow also has a wide distribution from Nepal to Sumatra in Indonesia through hill
forests to shrubland habitats (Phan et al., 2020). Nevertheless, the prediction of the remaining
habitat quality and suitability in Thailand and other countries have been conducted only in
some protected areas (Chaiyarat et al., 2019; Pintana & Lakamavichian, 2013), but not at the
regional or national level.

Species distribution modelling provides an overview of potential habitats for
threatened species and aids in conservation planning (Catullo et al., 2008). For instance,
previous studies have focused on identifying potentially high-quality habitat connectivity and
fragmentation (Crooks et al., 2011) as well as predicting global biodiversity trends (Araújo et
al., 2019). In Thailand, there are several studies that have predicted habitat suitability for
some of these five wild bovids in local areas (Prayoon et al., 2021), but habitat suitability
studies for larger extents across their distribution are lacking.

Here, we built ENM for the five Thai wild bovid species: gaur, banteng, wild water
buffalo, mainland serow and Chinese goral at two scales: first, at the regional scale
throughout the entire distribution and, second, at the country scale in Thailand. We aim to 1)
identify the potential distribution for these five species in South to Southeast Asia, and 2)
identify conservation areas in their geographical distribution, with a particular focus on
Thailand.

**Materials and methods**

108      Our workflow consisted of two main processes of data preparation and model building
109  (summarised in Figure S1) that generated habitat suitability maps for all species and
110  accessible areas used. Data preparation consisted of gathering the species occurrence data
111  and environmental data and selecting the accessible areas. Then, the model building
112  consisted of pre-processing, processing and post-processing steps.

**Study area**

The study area consists of 13 Asian countries: Bhutan, Bangladesh, Cambodia, China,
India, Indonesia, Laos, Malaysia, Myanmar, Nepal, Sri Lanka, Thailand and Vietnam (Figure 1),
that cover the distribution of gaur, banteng, wild water buffalo, mainland serow and Chinese
goral based on the literature (Table S1).

Figure 1 Species occurrence data before thinning (yellow circles), IUCN polygons (blue areas)
and study areas (grey areas) used in model building for five wild bovid species. First, a
common large 'accessible area' (A) was used for all species for model building, and then
species-specific accessible areas (B-F) for individual species.**Species Occurrence data**

We compiled species occurrence data collected from GPS records collected between
January 2000 and June 2021 from researchers, government, NGOs (World Wildlife Fund,
Wildlife Conservation Society, Freeland [Ash et al. (2021)], Panthera, Fauna & Flora
International, Friends of Wildlife and RIMBA) and open data sources, including GBIF
(https://www.gbif.org/) and eMammal (https://emammal.si.edu/). The data coverage by
country can be found in Table 1. The occurrence data were collected through observation of
animal signs (e.g. footprint and dung) during forest patrols, direct observation during wildlife
surveys, camera trapping, and radio-collar signals (Table S2). We used only the research grade
observation for GBIF data, which included the photo for species identification. We filtered all
the occurrences and excluded occurrence records outside the species-specific accessible area,
duplicated records from the same species and museum collections.

Table 1 The number of raw and after spatial thinning occurrence points is shown by species
and country.

| | Raw data | | | | | |
|---|---|---|---|---|---|---|
| Country | *Bos gaurus* | *Bos javanicus* | *Bubalus arnee* | *Capricornis sumatraensis* | *Naemorhedus griseus* | Total |
| Bangladesh | X | X | X | ? | X | 0 |
| Bhutan | 1 | X | ? | ? | X | 1 |
| Cambodia | 44 | 355 | ? | 38 | X | 437 |
| China | ? | ? | X | 109 | 301 | 410 |
| India | 286 | X | 78 | 2 | ? | 366 |
| Indonesia | X | 6 | X | ? | X | 6 |
| Laos | 2 | 1 | X | 11 | X | 14 |
| Malaysia | 1,067 | ? | X | 603 | X | 1,671 |
| Myanmar | 114 | 5 | ? | 99 | ? | 218 |
| Nepal | 4 | X | 1 | 2 | X | 7 |
| Sri Lanka | X | X | 21 | X | X | 21 |
| Thailand | 24,258 | 5,383 | 50 | 805 | 16 | 30,512 |
| Vietnam | 1 | ? | ? | ? | ? | 1 |
| **Grand Total** | **25,777** | **5,751** | **150** | **1,669** | **317** | **33,664** |

| | Thinning data | | | | | |
|---|---|---|---|---|---|---|
| Country | *Bos gaurus* | *Bos javanicus* | *Bubalus arnee* | *Capricornis sumatraensis* | *Naemorhedus griseus* | Total |
| Bangladesh | X | X | X | ? | X | |
| Bhutan | 1 | X | ? | ? | X | 1 |
| Cambodia | 14 | 48 | ? | 28 | X | 90 |
| China | ? | ? | X | 64 | 130 | 194 |
| India | 244 | X | 64 | 2 | ? | 310 |
| Indonesia | X | 6 | X | ? | X | 6 |
| Laos | 2 | 1 | X | 8 | X | 11 |
| Malaysia | 26 | 1 | X | 49 | X | 76 |
| Myanmar | 53 | 2 | ? | 51 | ? | 106 |
| Nepal | 4 | X | 1 | 1 | X | 6 |
| Sri Lanka | X | X | 20 | X | X | 20 |
| Thailand | 2,387 | 303 | 7 | 185 | 5 | 2,887 |
| Vietnam | 1 | ? | ? | ? | ? | 1 |
| **Grand Total** | **2,732** | **361** | **92** | **388** | **135** | **3,708** |

| Colour definitions | |
|---|---|
| Number | species presence with occurrence data |
| ? | species presence without occurrence data |
| X | no species presence |

**Environmental variables**

Hypothesized environmental variables were selected based on species' habitat and distribution related literature Table S1. We used 28 variables (supplementary material, Table S3) for model construction, including 19 bioclimatic variables (Booth et al., 2014) (average for 1970-2000) from WorldClim v2 (Fick & Hijmans, 2017), elevation (Shuttle Radar Topography Mission-SRTM) from WorldClim (Fick & Hijmans, 2017), slope (Amatulli et al., 2020), five land cover fractions (grass, tree, urban, water and crop) (Buchhorn et al., 2019), human population density (Stevens et al., 2015) and greenness through the normalized difference vegetation index (NDVI) (Didan, 2015). All layers were processed using the geographic coordinates system (Datum WGS84) and ~1 km$^2$ spatial resolution. We transformed the human population density using logarithm base 10 to adjust for skewness. We rescaled the NDVI layer by multiplying all values with a scale factor (0.0001), based on the Moderate Resolution Imaging Spectroradiometer (MODIS) User's guidelines (Didan et al., 2015).

**Accessible areas**

The accessible area refers to the parts of the world accessible to species via dispersal over time (Barve et al., 2011). The extent of the accessible area and the inclusion of a buffer zone have an important effect on ENM performance (Anderson & Raza, 2010; Barve et al., 2011). We used two accessible area sizes to delimitate our modelling extent (Figure 1). The first larger accessible area (hereon LA) includes most of the Asian continent and its ecoregions, and all species distributions are included as a common extent. The second accessible area was more restricted and cropped based on individual species-specific distributions (hereon SSA) from literature reviews (Table S1), IUCN polygons or 'ranges' (IUCN, 2020) and the terrestrial ecoregions where they occur. For creating the extent, we downloaded the current IUCN range maps for each species, then intersected those on ecoregions (Olson et al., 2001), then combined the results with selected ecoregions based on biogeographic knowledge of the species distributions and habitat preference from the literature reviews. For example, gaur habitat typically contains moist evergreen, semi-evergreen, and dry evergreen forests (Steinmetz et al., 2008; Tanasarnpaiboon, 2016), so we included these regions in our accessible areas. Further details on ecoregions included in accessible areas are in supplementary material, Table S4. To reduce overprediction and make our predictions closer to realised niche estimates, we used an occurrences-based threshold (OBR) method with ensemble models from (Mendes et al., 2020) for creating the spatially restricted ENM (hereon MSDM). OBR is an *a posteriori* method that restricts the suitable areas of our final ensemble models based on presence and the largest nearest neighbour distance among pairs of occurrences. Overall, we built four combinations between two accessible areas with and without MSDM methods for each species, including 1) No MSDM-SSA; 2) No MSDM-LA; 3) MSDM-SSA and 4) MSDM-LA.

**Model building**

We processed the species occurrence files and environmental datasets in R 4.0.1 (R
Core Team, 2020). We developed reproducible ecological niche models with optimized
processing times using the ENMTML package (Andrade et al., 2020), following three main
steps: 1) pre-processing, 2) processing and 3) post-processing.

In pre-processing, we performed occurrence thinning using 2 times the cell-size (1
km$^2$) (Velazco et al., 2019) to reduce clustering of species records and sampling bias. We used
principal component (PC) analysis (PCA) to reduce the collinearity of the predictors. We
assigned species' accessible areas to determine the species' distributions using a mask
function. We used random sampling to create pseudo-absence background points in a 1:1
ratio with presence points (Barbet-Massin et al., 2012). The occurrence and pseudo-absence
data was divided into two sets for fitting the model (75%) and evaluating the fitted models
(25%), using the bootstrapping partition method with 10 replications for each algorithm.

In the processing step, eight algorithms were used to build the ENMs, namely:
BIOCLIM (Booth et al., 2014), Generalized Linear Models (McCullagh & Nelder, 1989),
Generalized Additive Models (Hastie, 2018), Random Forest (Liaw & Wiener, 2002), Support
Vector Machine  (Karatzoglou et al., 2004), Maximum Entropy default (Phillips et al., 2006),
Maximum Likelihood (Royle et al., 2012) and Bayesian Gaussian Process (Golding, 2014). All
models used the default settings from the ENMTML package, which included the functions
from different packages (e.g. dismo, maxnet) based on the algorithms that use to fit the
models. The data type used for each algorithm is in supplementary materials, Table S5.

In the post-processing step, we created ensemble models using the weighted average
(WMEAN) method based on the True Skill Statistic (TSS) values for building final habitat
suitability and binary maps. The benefits of ensemble models are 1) robust decision-making
(Ahmad et al., 2020); 2) reducing uncertainty (Marmion et al., 2009); and 3) a combination of
several models into one model prediction (Kindt, 2018). We used TSS to calculate threshold
values to convert habitat suitability maps into binary suitability maps (0 = unsuitable and 1 =
suitable). We used TSS and area under the curve (AUC) for evaluating model performance.
The TSS threshold is calculated using the maximum summed specificity and sensitivity and is
not based on prevalence, where an equal TSS score for given models means similar
performance (Allouche et al., 2006). Therefore, we selected the final models from the best
TSS of weighted average ensemble models. We assessed the model's accuracy by plotting a
new dataset of species occurrences obtained from camera traps and human observations
(https://www.gbif.org/) on the binary maps. Because, for example, gaur have been recorded
to walk up to 6.3 km a day (mean 1.6 km (Rizal et al., 2020)), we created a 5 km buffer zone
measured from the edges of the suitable pixels to include occurrences within the travel
distance of wild bovids' movement (Ahrestani & Karanth, 2014; Gardner et al., 2014). The
percentage of points inside and outside the suitable areas and the buffer zone was calculated
for each species. We present all the results, then only models with high prediction accuracy
(greater than 80%,(Zhang et al., 2015)) are selected for further analyses. The total suitable
areas of the best TSS binary map models were calculated using the zonal function in the raster

R package (Hijmans, 2023). Then, we summed the pixels of the best TSS binary maps to
generate the map of species number.

**Protected area analyses**

The source for our protected areas map was the World Database of Protected Area
(WDPA) (UNEP-WCMC & IUCN, 2021). We classified protected areas based on IUCN protected
areas from WPDA into 8 categories, including categories 1 to 6 as IUCN management
categories I to VI; category 7 as 'not applicable', which includes 'not reported', 'not applicable'
and 'not assigned' protected areas; and category 8 as non-protected areas, which are the
remaining areas that have not been classified as IUCN categories 1 to 7 (UNEP-WCMC and
IUCN, 2021). Then, we used the zonal function in the Raster package to calculate overlapping
areas between the suitable areas and protected areas for each species.

We calculated the percentage of suitable areas in WDPA polygons using the
exact_extract function in exactextractr package (Baston et al., 2021) for extracting the
suitable areas (values = 1) from binary map rasters in each WDPA polygon. Then, we classified
each PA into 5 different suitability categories based on the percentage of suitable habitat in
the PA: low suitability (0 - 20%); low - medium suitability (>20 - 40%); medium suitability (>40
- 60%); high suitability (>60 - 80%) and very high suitability (≥80%), and selected only the PAs
that have the proportion of suitable area larger than species home range in the result. We
have provided the code for creating the models in a GitHub repository.

**Results**

We compiled 33,664 occurrence records. After filtering and spatial thinning, we used
3,708 points for modelling: 2,732 for gaur, 361 for banteng, 92 for wild water buffalo, 388 for
mainland serow, and 135 for Chinese goral. The majority of the thinning occurrences (77%)
were collected in Thailand, India and other countries in mainland SEA; see Table 1**Error!**
**Reference source not found.** for details on the data coverage by country and supplementary
materials Table S10 for details on the study sites.

We found that the PCA reduced the 28 environmental variables into 12 PCs that
explained 95% of the environmental variance in the variables for the LA models for all species.
The PCs for SSA models explained more than 96% of the total variance and the PC number
varied by species, comprising 13 PCs (wild water buffalo), 11 PCs (gaur, mainland serow), and
10 PCs (banteng, Chinese goral). The bioclimatic variables were important variables in all
species models. For LA models, the first two axes (PC1 and PC2) have high contributions from
the annual mean temperature (bio01), mean temperature of the coldest month (bio06), mean
temperature of the driest quarter (bio09) and mean temperature of the warmest quarter
(bio10). The first two axes of SSA models showed high positive contributions from mean
temperature of the coldest month (gaur), minimum temperature of coldest month (banteng,
mainland serow), annual mean temperature (wild water buffalo, mainland serow), and
precipitation of the wettest quarter (Chinese goral). We also found that NDVI, elevation, slope

and human population density have less effect on explaining the variability for the first two
PCs for all species. The correlations between PCs and individual environmental variables, PC
biplots and percentage of explained variance are summarised in supplementary materials,
Table S6 and Figure S2.

**Ecological niche models**

Overall, all ensemble models showed high performance both for TSS and the area
under the curve (AUC) with the highest performing models over 0.8 for all species (Table 2).
Models with species-specific accessible areas were not always the best performing models,
but most ensemble models performed above 0.7 TSS. The habitat suitability prediction maps
using the best model ensembles are in supplementary materials, Figure S3 (SSA) Figure S4
(LA), Figure S5 (selected the best model of SSA and LA) and the binary maps which were used
for calculating the suitable area in Figure S6. The performance of spatially restricted
ensembles was higher in comparison with the No MSDM models, as the TSS was improved
for banteng, Chinese goral and wild water buffalo. The lowest performing model for wild
water buffalo was the No MSDM-SSA (TSS = 0.57). The best model for gaur was No MSDM-
LA, banteng and Chinese goral is MSDM-LA, wild water buffalo is MSDM-SSA, and mainland
serow is No MSDM-SSA. We found that all species have small predicted suitable habitats.
Moreover, all species models predicted less than 50% of the suitable areas inside PAs.

Table 2 True Skill Statistics (TSS) and Area Under the Curve (AUC) values of the weighted
average ensemble, and the threshold values for binary maps for five species classified by
accessible area type and MSDM method.Best performing models for each accessible area by
TSS are shown in **Boldface**.

| | Large accessible area | | | | | | Species specific accessible area | | | | | |
| | No MSDM[a] | | AUC | MSDM (OBR)[b] | | AUC | No MSDM[a] | | AUC | MSDM (OBR)[b] | | AUC |
| | TSS | | | TSS | | | TSS | | | TSS | | |
| Species | Score | Threshold | Score | Score | Threshold | Score | Score | Threshold | Score | Score | Threshold | Score |
| Gaur *Bos gaurus* | **0.92** | 0.49 | 0.99 | 0.92 | 0.44 | 0.99 | **0.88** | 0.39 | 0.98 | 0.88 | 0.41 | 0.98 |
| Banteng *Bos javanicus* | 0.93 | 0.55 | 0.99 | **0.94** | 0.41 | 1 | 0.85 | 0.33 | 0.96 | **0.83** | 0.42 | 0.97 |
| Wild water buffalo *Bubalus arnee* | 0.67 | 0.47 | 0.88 | **0.72** | 0.6 | 0.9 | 0.57 | 0.58 | 0.83 | **0.85** | 0.44 | 0.95 |
| Mainland serow *Capricornis sumatraensis* | **0.87** | 0.55 | 0.97 | 0.76 | 0.47 | 0.94 | **0.93** | 0.57 | 0.98 | 0.93 | 0.52 | 0.98 |
| Chinese goral *Naemorhedus griseus* | 0.91 | 0.29 | 0.98 | **0.91** | 0.59 | 0.98 | 0.87 | 0.47 | 0.96 | **0.9** | 0.39 | 0.97 |

276 [a] spatially restricted ENM
277 [b] occurrences-based threshold

Figure 2 Total of the suitable area in km² for each species and countries. Blue is the species-specific accessible area (SSA) and grey is the large accessible area models (LA) (see details in the supplementary Table S7).

The total of the suitable areas in km² for each species and country are shown in Figure 2 and Suitable areas calculated from the best model are in supplementary materials Table S8 and the IUCN protected areas for all types of models are in Figure S7.

Our model's out-of-sample predictions with new species occurrences demonstrated a higher prediction accuracy within Thailand than the entire distribution, and this was further improved by including 5 km buffer zones, with the exception of Chinese goral, which exhibited poor accuracy across all scales (Table 3 and Figure 3). Implementing a buffer zone improves the accuracy of all four remaining species. For large herbivore species gaur, banteng and wild water buffalo, the model cropped to Thailand showed a higher accuracy (>80%) compared to the entire distribution (~60-80%). We selected only model predictions with a high accuracy percentage, greater than 80%, for further analyses. As a result, three species, including gaur, banteng, and wild water buffalo, were retained, while two species, mainland serow and Chinese goral, were excluded from the rest of the study. Furthermore, we cropped the entire distribution to focus only on the result within Thailand as the number of data collection and model predictions is higher compared to the entire species distribution. The result of the entire distribution for all species can be found in the supplementary material, Figure S3 and Figure S4.

299 Table 3 Comparison of the accuracy of the selected best models[a] in predicting out-of-sample
300 data for the entire accessible areas range and Thailand.

| | | No buffer | | | Buffer | | | |
|---|---|---|---|---|---|---|---|---|
| Entire accessible areas | Total | Unsuitable | Suitable | Accuracy % | Unsuitable | Suitable | Buffer 5 km | Accuracy % |
| *B. gaurus* (Gaur) | 221 | 85 | 136 | 62 | 23 | 136 | 62 | 90 |
| *B. javanicus* (Banteng) | 12 | 4 | 8 | 67 | 2 | 8 | 2 | 83 |
| *B. arnee* (Buffalo) | 35 | 4 | 31 | 89 | 0 | 31 | 4 | 100 |
| *C. sumatraensis* (Mainland serow) | 21 | 17 | 4 | 19 | 7 | 4 | 10 | 67 |
| *N. griseus* (Chinese goral) | 10 | 9 | 1 | 10 | 7 | 1 | 2 | 30 |

| | | No buffer | | | Buffer | | | |
|---|---|---|---|---|---|---|---|---|
| Thailand | Total | Unsuitable | Suitable | Accuracy % | Unsuitable | Suitable | Buffer 5 km | Accuracy % |
| *B. gaurus* (Gaur) | 52 | 8 | 44 | 85 | 2 | 44 | 6 | 96 |
| *B. javanicus* (Banteng) | 10 | 2 | 8 | 80 | 0 | 8 | 2 | 100 |
| B. arnee (Wild water buffalo) | 1 | 0 | 1 | 100 | 0 | 1 | 1 | 100 |
| *C. sumatraensis* (Mainland serow) | 14 | 9 | 5 | 36 | 2 | 4 | 8 | 86 |
| *N. griseus* (Chinese goral) | 2 | 2 | 0 | 0 | 2 | 0 | 0 | 0 |

301 [a] The best models for gaur is No MSDM-LA, banteng is MSDM-LA, wild water buffalo and Chinese goral is
302 MSDM-SSA and mainland serow is No MSDM-SSA.

303 Nearest distance from out of sample points to suitable area

| | | Distance (km) | | |
|---|---|---|---|---|
| Species | point | Min | Mean | Max |
| Gaur | 52 | 0.0047 | 1.54 | 22.4 |
| Banteng | 10 | 0.0323 | 4.72 | 39.9 |
| Wild water buffalo | 1 | 0.811 | 0.811 | 0.811 |
| Mainland Serow | 14 | 0.00668 | 6.07 | 38.1 |
| Chinese goral | 2 | 0.147 | 1.54 | 2.93 |

Figure 3 Model prediction testing for five bovid species (A-E) by calculating the percentage of the out of sample points that fall inside the model predicted suitable areas (blue). The model fitting datasets (red) were mainly within the suitable areas compared to the new occurrence dataset (green). IUCN ranges show greater areas than the predictions for mainland serow and Chinese goral. Some of the occurrence data were distributed outside both the model predicted suitable area and IUCN range.

Figure 4 Habitat suitability prediction maps of three wild bovids species in Thailand: gaur (*B.*
*gaurus*), banteng (*B. javanicus*) and wild water buffalo (*B. arnee*) species (A-C) using the best
model from the weighted average ensemble. The value ranges from 0-1: yellow represents
low suitability and dark brown represents high suitability. Interactive maps are provided in
the online supplementary material (link).

**Identifying priority areas for conservation**

Most suitable habitats in protected areas are located in IUCN category Ia (Strict nature
reserve), Ib (Wilderness area) and II (National Park) areas for the best TSS models for all
species, while IUCN category V (Protected landscape or seascape) has the least. Overall, more
than half of the species' suitable habitat is not under any form of protection defined by the
WDPA (supplementary materials, Table S8, Figure S8). The proportion of the suitable area in
each WDPA of the best models from SSA and LA for each species are presented in
supplementary materials, Figure S9 and Figure 10.

In Thailand, we identified a high percentage (≥80%) of suitable area of Thailand for
gaur in 122 PAs (74,268 km$^2$; 15% of Thailand), banteng in 102 PAs (59,528 km$^2$; 12% of
Thailand), and wild water buffalo in 3 PAs (559 km$^2$; 0.1 % of Thailand). A high proportion of
the suitable area for gaur and banteng is in Thungyai Naresuan, Kaengkrachan and Huai Kha
Khaeng, and for wild water buffalo in Phu Wua WS and Dong Yai WS in eastern DPKY-FC
(Figure 4 and Figure 5). The hotspots for five species can be found in supplementary materials,
Figure S11.

Proportions range from 0 (all unsuitable) to 1 (all suitable), with suitability determined by
thresholds from species best performing models. (A) gaur (*Bos gaurus*), (B) banteng (*Bos*
*javanicus*), (C) wild water buffalo (*Bubalus arnee*).

Figure 5 Estimated species richness of three wild bovids in Thailand. The species are gaur,
banteng, and wild water buffalo. Frames A-C focus on (A) Western Forest Complex
(WEFCOM), (B) Dong Phayayen-Khao Yai Forest Complex (DPKY-FC) and (C) Eastern Forest
Complex, where the overlapping suitable areas of all species (n=3). Western, Dong Phayayen-
Khao Yai and Eastern forests have suitable areas for gaur, banteng and wild water buffalo for
both inside PAs and also in the surrounding areas.

We found that the highest percentage of suitable area was comprised of mixed
deciduous forest for all species, followed by evergreen forest for gaur and banteng, and dry
dipterocarp forest for wild water buffalo. We found a percentage of non-forest areas
identified from the total suitable for all species: wild water buffalo (71%), banteng (33%), and
gaur (24%). For more details of forest types by suitable areas, see Table 4 and Figure S12.
346

347    Table 4 The suitable areas of five bovid species classified by forest types in Thailand.

| Forest types | Gaur (*Bos gaurus*) | | Banteng (*Bos javanicus*) | | Wild water buffalo (*Bubalus arnee*) | | Mainland serow (*Capricornis sumatraensis*) | | Chinese goral (*Naemorhedus griseus*) | |
|---|---|---|---|---|---|---|---|---|---|---|
| | km² | % | km² | % | km² | % | km² | % | km² | % |
| Bamboo Forest | 390 | 0.22 | 348 | 0.19 | 178 | 0.13 | 250 | 0.37 | 2 | 0.02 |
| Beach Forest | 3 | - | 8 | - | 26 | 0.02 | 1 | - | - | - |
| Dry Dipterocarp Forest | 11,119 | 6.26 | 12,876 | 7.13 | 7,365 | 5.43 | 1,415 | 2.07 | 3,546 | 23.91 |
| Dry Evergreen Forest | 20,730 | 11.68 | 19,209 | 10.63 | 5,944 | 4.38 | 13,893 | 20.3 | 1,027 | 6.93 |
| Freshwater Swamp Forest | 66 | 0.04 | 134 | 0.07 | 24 | 0.02 | - | - | - | - |
| Mangrove Forest | 609 | 0.34 | 1,072 | 0.59 | 1,028 | 0.76 | 121 | 0.18 | - | - |
| Mixed Deciduous Forest | 66,132 | 37.25 | 59,211 | 32.77 | 18,837 | 13.88 | 25,243 | 36.88 | 7,347 | 49.54 |
| Moist Evergreen Forest | 14,802 | 8.34 | 15,729 | 8.7 | 1,975 | 1.46 | 12,213 | 17.84 | - | - |
| Montane Forest | 16,693 | 9.4 | 8,497 | 4.7 | 812 | 0.6 | 7,532 | 11 | 1,878 | 12.66 |
| Peat Swamp Forest | 49 | 0.03 | 2 | - | 201 | 0.15 | - | - | - | - |
| Pine Forest | 634 | 0.36 | 185 | 0.1 | 87 | 0.06 | 78 | 0.11 | 15 | 0.1 |
| Savanna | 548 | 0.31 | 348 | 0.19 | 108 | 0.08 | 312 | 0.46 | 7 | 0.05 |
| Secondary Forest | 2,017 | 1.14 | 1,856 | 1.03 | 1,189 | 0.88 | 602 | 0.88 | 153 | 1.03 |
| Teak Plantation | 846 | 0.48 | 1,045 | 0.58 | 919 | 0.68 | 60 | 0.09 | 12 | 0.08 |
| Vegetation on Pen Rock Platform | 201 | 0.11 | 208 | 0.11 | 118 | 0.09 | 90 | 0.13 | 2 | 0.01 |
| Other Plantations | 37 | 0.02 | 42 | 0.02 | 29 | 0.02 | 9 | 0.01 | - | - |
| Non-forest Area | 42,649 | 24.02 | 59,923 | 33.16 | 96,883 | 71.38 | 6,631 | 9.69 | 842 | 5.68 |
| Total | 177,526 | 100 | 180,693 | 100 | 135,725 | 100 | 68,452 | 100 | 14,831 | 100 |

**Discussion**

We modelled the potential distribution for the five threatened wild bovid species,
distributed in East, South and Southeast Asia. Our aim was to build predictive models to
identify conservation areas, and potential species richness maps in their entire geographical
distribution. However, the model predictions were better for Thailand, where most data were
from for all but Chinese goal (Table 3), therefore we were focus on Thailand. We found our
models were able to predict the presence of out of sample observations well for three species,
gaur, banteng, and wild water buffalo throughout the entire distribution (≥62%), but not
mainland serow or Chinese goral (≤19%). We identified that suitable areas were fragmented
and often (50%) located outside PAs. Those suitable areas outside PAs could possibly be
managed as corridors or buffer zones to connect currently fragmented bovid populations,
thereby enhancing long-term wild bovid conservation success (Karanth, 2016; Penjor et al.,
2021) which requires further investigations. For example, a corridor was built within DPKY-FC

and showed the possibilities of connecting the western forest complex and Kaengkrachan NP
to conserve the endangered tiger population (Sukmasuang et al., 2020; Suttidate et al., 2021).

Our study found that most suitable areas for gaur were similar to IUCN range
assessments (Duckworth et al., 2016) and consistent with studies that have confirmed species
presences, such as in Thailand's PAs (Prayoon et al., 2021), Myanmar (Hein et al., 2020) and
Western Ghats in southwestern India and Manas WS in the Himalayan foothills (Choudhury,
2002). However, there are differences. Our study predicted larger gaur suitable habitats in
Thailand inside (~82,400 km$^2$) and outside (95,000 km$^2$) PAs than Prayoon et al. (2021), who
predicted 39,508 km$^2$ of total suitable habitat. Choudhury (2002) predicted their distributions
in Western Ghats, Central and North-eastern India which included larger than our predictions.
Our predictions used NDVI and land coverage fractions (Table S3) for predicting greenness,
which may be useful for predicting the vegetation quality and availability for ungulates
(Borowik et al., 2013). However, NDVI is difficult to differentiate vegetation variations (Didan
et al., 2015; Martinez & Labib, 2023), such as between specific agricultural areas, grassland,
and dense forest canopy. This may include other vegetation types other than the species'
habitat in suitable areas and estimated larger suitable areas predicted in non-forest areas and
non-PAs to be identified in our study, compared to Prayoon et al's study. Other studies
suggest that gaur does use crop plantations or man-made grasslands, which may increase the
suitable areas in our prediction, even if these are not their natural habitats and lead to conflict
between humans and gaur (Chaiyarat et al., 2021).

Our best model predicted larger suitable areas (446,075 km$^2$) for banteng than the
IUCN-SSC report released in 2010 (~ 209,000 km$^2$) (IUCN-SSC AWCS Group, 2010). We found
a high percentage of predicted suitable areas in Eastern Plains Landscape (ELP) and Chhaeb
WS in Cambodia; the former supports the likely largest banteng population globally (Gray et
al., 2012). However, our results showed low habitat suitability in Sundaic Southeast Asia, with
just 2% of the total suitable area in Indonesia (mainly in Alas Purwo NP, Java) and 2% of the
total suitable area in Malaysia. Banteng populations and habitats in Southeast Asian islands
(Borneo, Java, and Bali) are threatened due to hunting for horn and meat consumption and
habitat loss (Dewi et al., 2020). In Thailand, we found high suitability similar to previous
studies in Eastern (Menkham et al., 2019) and Western forest complexes (Jornburom et al.,
2020), including reintroduction areas in Salak Pra WS (Chaiyarat et al., 2019) and where recent
recolonisation by natural population movement has occurred in Mae Wong NP (Phoonjampa
et al., 2021).

Wild water buffalo has been domesticated and bred as livestock, making it hard to
distinguish between the free-grazing domestic buffalo and wild water buffalo as
domesticated animals may replace wild animals in suitable habitats and cause the high
suitable area prediction outside PAs, especially in overlapping habitats (Zhang et al., 2020).
We estimate the highest percentages of suitable areas at Kaziranga NP in India, currently with
the largest population of wild water buffalo (Kaul et al., 2019). Grasslands and flood plain
areas of Manas NP (500 km$^2$) and Kaziranga NP (>850 km$^2$) in India contain the most suitable

habitat and are the main population strongholds for wild water buffalo (Choudhury, 2014). In
Thailand, this type of habitat can be found in many places, but it is not often represented in
protected areas. Wild water buffalo are only found in Huai Kha Kheang WS parts of the
Western Forest Complex. Our model predicts that only 43% of Huai Kha Kheang Wildlife
Sanctuary is suitable for this species, primarily because the floodplains are mainly situated
close to the mainstream in the middle of the PA. Additionally, the population has remained
constant for decades, which could be attributed to a single population group or constraints
within suitable habitats.

The three selected species showed overlapping suitable areas in the Western Forest
Complex, Eastern Forest Complex, and Dong Phayayen-Khao Yai Forest Complexes (DPKY-FC).
These forest complexes encompass extensive areas of high wildlife biodiversity and diverse
forest types, including several contiguous protected areas (PAs) situated at the borders of
Cambodia and Myanmar. The Western Forest Complex is the largest conservation area in
Thailand where these wild bovids still exist, while the DPKY-FC maintains a higher population
of gaur as they are mainly covered by evergreen forest. The Eastern Forest Complex sustains
a large population of banteng because most of the main vegetation consists of deciduous and
dipterocarp forest. Gaur uses a diversity of types of habitats and prefers denser canopy at
higher elevation than banteng, which tends to inhabit in dry and open habitats such as dry
dipterocarp and deciduous forests (Gray & Phan, 2011; Steinmetz, 2004). Wild water buffalo
also shares overlapping areas with these two species, despite its distribution being found
exclusively in Huai Kha Khaeng Wildlife Sanctuary. We recommend protecting these
important suitable habitats to ensure the protection of wild bovids. This may involve
implementing active patrolling to reduce illegal intrusions, snare removal and habitat
management based on their diet diversity (McShea et al., 2019). Additionally, one option to
maintain wild water buffalo populations is to reintroduce them into their historical range,
from which they have been extirpated. This method could be evaluated by combining
predicted suitable areas with several important factors such as vegetation types, forage
biomass, carrying capacity and hunting pressure (Bora et al., 2024).

In this study, we included all subspecies data points in our model ensembles as we aim
to extrapolate and predict the entire range of species' habitat suitability, but this may increase
uncertainty (Dormann, 2007). These five bovids have multiple subspecies, including 3
subspecies of gaur (Duckworth et al., 2016), banteng (Gardner et al., 2016) wild water buffalo
(Kaul et al., 2019) and mainland serow (Mori et al., 2019), and 2 subspecies of Chinese goral
(Duckworth et al., 2008). Subspecies may vary in niche, climate and biological interactions
that could affect the model predictions. The low habitat suitability of our study in Borneo for
banteng could be because climatic and geographic conditions differ for *B.j. lowi* compared to
those in mainland Asia, affecting model transferability across different regions (Zhu et al.,
2021). Equally, Mori et al. (2019) suggest that Chinese goral (*N. griseus*) should be reclassified
within Brown goral (*N. goral*) together and Burmese goral (*N. evansi*) that together with *N.*
*griseus* should be split to become an individual species. Future analyses must consider these

taxonomic reclassifications. However, we modelled species level habitat suitability, rather than the subspecies, as we assume that there is less likely to be habitat and environmental condition variation at the subspecies level for these bovids (Smith et al., 2019).

We found that using the MSDM OBR technique showed a better predicted suitable area of the ecological niche, closer to the real distribution for species with more restricted ranges like banteng, wild water buffalo and Chinese goral, with higher performance TSS values compared to No MSDM models. We recommend restricting the accessible area for predicting wild water buffalo potential habitat to reduce overprediction caused by overlapping areas with domestic water buffalo.

We also used ensemble approaches, to obtain better predictive performance than from any single model type, but further analyses could also look at individual model results using different parameters, such as differing pseudo-absence background point ratios. The equal ratio of presence to pseudo-absence (1:1 ratio) has been used in several types of model like general linear models, artificial neural networks, and Maxent models, and it is also recommended for use in ensemble models when dealing with small sample sizes (Liu et al., 2019).

Limitations

We acknowledge sampling deficiencies across the regions. We had fewer occurrences in Vietnam, Laos, Myanmar and Indonesia compared to Thailand, from which a large number of our data points came (30,512 points in Thailand, 3,152 points outside Thailand, **Error! Reference source not found.**). Occurrence data based on data accessibility may have sampling bias, particularly with clustered points for gaur, banteng, and mainland serow. We minimised these biases through spatial thinning (Aiello-Lammens et al., 2015). Since we found large amounts of suitable areas outside of Thailand, we suggest that future studies should focus on monitoring bovid populations in other countries, especially in India and Myanmar. However, because of this and the model performance, we focused on Thailand.

Missing data data has impacted some results. The model TSS values for endangered banteng and Chinese goral are over 0.8, yet our models predict unsuitable areas in part of Indonesia (east and central Kalimantan; Dewi et al. 2020) for banteng and China (e.g. Beijing and northeast Inner Mongolia; (Yang et al., 2019) for Chinese goral from which these species have been reported. This would likely be improved if more spatial data were available for these species.

We used a new dataset of species occurrences to assess our model's performance with a 5 km buffer zone, aiming to enhance modelling accuracy. Given these species have quite large home ranges and daily movements, adding a buffer to represent this movement unsurprisingly lead to better model predictions for all species, but most notably for mainland serow, changing the out of sample prediction from 19% to 67% for the entire region and 36% to 86% for Thailand. The buffer zone may indicate the utilisation of unsuitable areas of the

species near forested regions, such as secondary forests, agricultural areas, or water resources, which possibly extend these buffer areas from the protected aeras to enhance the wildlife protection.

The spatial restriction method, OBR, can be sensitive to the distribution of occurrence data, because it keeps predicted suitable areas close to the occurrence locations. This may lead to the exclusion of potential suitable areas driven by a lack of occurrence data in those areas. For example, the wild water buffalo No MSDM predicted potentially suitable habitat around the Sre Pok Wildlife Sanctuary in Cambodia where the species is distributed (Gray et al., 2012), but after the MSDM, this potential habitat was excluded as we lack occurrence data in Cambodia. Although our study showed slightly different TSS values between two different accessible area extents, we encourage testing the different accessible areas as it affects the model results (Anderson & Raza, 2010). Moreover, model performance varied with accessible area sizes and spatial restrictions, emphasising the need for careful accessible area definition in ecological modelling (Barve et al., 2011). Further, future analyses may try to better account for the current presence of species by accounting for factors such as hunting using other proxies, such as other human-disturbance metrics like distance from roads (Lim et al., 2021).

**Conclusion**

Our study provided an overview of the suitable remaining habitat for threatened bovid species at a regional scale using high-resolution environmental variables and species occurrence data from multiple observation methods. Our predictions showed that the suitable areas are small and fragmented for all species, and more than 50% of suitable areas are outside of protected areas. Those suitable areas outside PAs could possibly become efficient conservation areas, such as forest corridors or buffer zones to connect fragmented bovid populations and enhance long-term habitat conservation. Our predictions may inform conservation actions to avoid further defaunation of wild bovidae such as the management of human-wildlife conflicts and habitat quality for long-term species survival.

**Acknowledgements**

We thank all the data contributions and collaborations from these institutions:

Dr Supagit Vinitpornsawan (Director of Wildlife Conservation Area Management and Education Center, Wildlife Conservation Office, Department of National Parks, Wildlife and Plant Conservation, Thailand) for animal track and sign data in Thailand. For camera trap data in Thailand: Naret Seuaturien (WWF Thailand), Manoon Pliosungnoen, Department of Environmental and Forest Biology, State University of New York College of Environmental Science), Wanlop Chutipong (Conservation Ecology Program, Pilot Plant Development and Training Institute, King Mongkut's University of Technology Thonburi), Nucharin Songsasen (Smithsonian's National Zoo and Conservation Biology Institute), Lon I. Grassman, Jr. (Feline Research Program, Caesar Kleberg Wildlife Research Institute, Texas A&M University, USA), Freeland Foundation, WCS Thailand, Smithsonian Institution, PANTHERA USA in Thailand.

WWF Thailand would like to thank: WWF Germany, WWF Sweden, WWF US, B.Grimm, WWF Japan, and WWF Switzerland for wonderful support of field projects, and Department of National Parks, Wildlife, and Plant Conservation for kind permission and collaboration. WCS Cambodia, Friends of Wildlife, Wildlife Alliance

and Ministry of Environment, Royal Government of Cambodia for the camera trap data in Cambodia. CarBi
Project of WWF Lao and WWF Greater Mekong for the camera trap data in Laos. Camera trap in Myanmar:
Friends of Wildlife and Data collecting in Tanintharyi, Myanmar was partially funded by the European Union,
Helmsley Charitable Trust and mainly funded by Integrated Tiger Habitat Conservation Project, through the
Fauna & Flora International (FFI). Open source databases: https://www.gbif.org, camera trap data from
eMammals (https://emammal.si.edu/) by William J. McShea (Conservation Ecology Center, Smithsonian
Conservation Biology Institute) and Megan Baker-Whatton (Smithsonian Conservation Biology Institute, and
George Mason University) under the project: Qionglai Mountains Project, Liangshan Mountains Project, Habitat
Connectivity in Minshan Mountains, Qinling Project, HKK ForestGEO Project, and Carnivore Intraguild
Interactions in Select Thailand Reserves. Also, the other researchers who share the species occurrence data with
us. We thank the IUCN specialists for commenting on the early version of the habitat suitability maps. The
authors wish to acknowledge the use of New Zealand eScience Infrastructure (NeSI) high-performance
computing facilities, consulting support and/or training services as part of this research.

**Funding Statement**

WH was supported by Manaaki New Zealand Scholarships. DTSH, RLM, RSJ were supported by Bryce Carmine
and Anne Carmine (née Percival), through the Massey University Foundation. DTSH was supported by the
Percival Carmine Chair in Epidemiology and Public Health and Royal Society Te Apūrangi (grant no. RDF-
MAU1701). EA supported by U.S. Fish & Wildlife Service Rhinoceros and Tiger Conservation Fund, David
Shepherd Wildlife Foundation, Care for the Wild International/Born Free Foundation, Point Defiance Zoo &
Aquarium and 21st Century Tiger. DN was supported by King Mongkut's University of Technology Thonburi
(grant no. WOR1-2557–2558); International Association for Bear Research and Management (Research &
Conservation grant), 2012 & 2014 and The Asahi Glass Foundation (Research Grant), 2003. Antony Lynam was
supported by TRF/BIOTEC Special Program for Biodiversity Research and Training. George A. Gale was
supported by TRF/BIOTEC Special Program for Biodiversity Research and Training (BRT R 346001) and National
Science and Technology Development Agency (grant no. NSTDA P-11-00390). Data collecting in Tanintharyi,
Myanmar was partially funded by the European Union, Helmsley Charitable Trust and mainly funded by
Integrated Tiger Habitat Conservation Project - grant no. ITHCP1338, through the Fauna & Flora International.

**Data Accessibility**

Data available upon request

**Competing Interests**

We have no competing interests.

**Authors' Contributions**

**Alex Riggio**: Investigation, Resources, Writing - Review and Editing.

**Alexander Godfrey**: Investigation, Resources, Writing - Review and Editing.

**Anony Lynam**: Investigation, Resources, Writing - Review and Editing

**David T. S. Hayman:** Conceptualization, Methodology, Validation, Supervision, Writing - Original Draft, Writing
- review and Editing, Project administration, Funding acquisition.

**Dusit Ngoprasert**: Investigation, Resources, Writing - Review and Editing.

**Eric Ash**: Investigation, Resources, Writing - Review and Editing.

**Francesco Bisi**: Investigation, Resources, Writing - Review and Editing.

**George A. Gale**: Investigation, Resources.

**Giacomo Cremonesi**: Investigation, Resources, Writing - Review and Editing.

**Gopalasamy Reuben Clements**: Investigation, Resources, Writing - Review and Editing.

**Jonathan C Marshall**: Methodology, Software, Validation, Formal analysis
Supervision, Writing - Original Draft, Review and Editing.

**Marnoch Yindee**: Investigation, Resources, Writing - Review and Editing.

**Nay Myo Shwe:** Investigation, Resources.
**Pin Chanratana**: Investigation, Resources.
**Reju Sam John**: Methodology, Software, Validation, Formal analysis, Supervision, Writing - Original Draft,
Review and Editing.
**Renata L. Muylaert:** Methodology, Software, Validation, Formal analysis, Supervision, Writing - Original Draft,
Review and Editing.
**Robert Steinmetz**, Investigation, Resources, Writing - Review and Editing.
**Rungnapa Phoonjampa**: Investigation, Resources.
**Saw Soe Aung**: Investigation, Resources.
**Seri Nakbun**: Investigation, Resources.
**Stephanie Schuttler**: Investigation, Resources, Writing - Review and Editing.
**Thomas N. E. Gray**: Investigation, Resources, Writing - Review and Editing.
**Wantida Horpiencharoen**: Conceptualization, Methodology, Software, Formal analysis, Data Curation, Writing
- Original Draft, Writing - review and Editing, Project administration, Funding acquisition.
**Worrapan Phumanee**: Investigation, Resources.
All authors read the text, provided comments, suggestions and corrections, and approved the final version.

**References**

Ahmad, S., Yang, L., Khan, T. U., Wanghe, K., Li, M., & Luan, X. (2020). Using an ensemble modelling approach
to predict the potential distribution of Himalayan gray goral (Naemorhedus goral bedfordi) in Pakistan.
*Global Ecology and Conservation, 21*, e00845. https://doi.org/https://doi.org/10.1016/j.gecco.2019.e00845
Ahrestani, F. S., & Karanth, K. U. (2014). Gaur Bos gaurus C.H. Smith, 1827. In M. Melletti & J. Burton (Eds.),
*Ecology, Evolution and Behaviour of Wild Cattle* (pp. 174-193). Cambridge University Press.
Aiello-Lammens, M. E., Boria, R. A., Radosavljevic, A., Vilela, B., & Anderson, R. P. (2015). spThin: an R package
for spatial thinning of species occurrence records for use in ecological niche models
[https://doi.org/10.1111/ecog.01132]. *Ecography, 38*(5), 541-545.
https://doi.org/https://doi.org/10.1111/ecog.01132
Allouche, O., Tsoar, A., & Kadmon, R. (2006). Assessing the accuracy of species distribution models:
prevalence, kappa and the true skill statistic (TSS) [https://doi.org/10.1111/j.1365-2664.2006.01214.x].
*Journal of Applied Ecology, 43*(6), 1223-1232. https://doi.org/https://doi.org/10.1111/j.1365-
2664.2006.01214.x
Amatulli, G., McInerney, D., Sethi, T., Strobl, P., & Domisch, S. (2020). Geomorpho90m, empirical evaluation
and accuracy assessment of global high-resolution geomorphometric layers. *Scientific Data, 7*(1), 162.
https://doi.org/10.1038/s41597-020-0479-6
Anderson, R. P., & Raza, A. (2010). The effect of the extent of the study region on GIS models of species
geographic distributions and estimates of niche evolution: preliminary tests with montane rodents (genus
Nephelomys) in Venezuela [https://doi.org/10.1111/j.1365-2699.2010.02290.x]. *Journal of Biogeography,*
*37*(7), 1378-1393. https://doi.org/https://doi.org/10.1111/j.1365-2699.2010.02290.x
Andrade, A. F. A. d., Velazco, S. J. E., & De Marco Júnior, P. (2020). ENMTML: An R package for a
straightforward construction of complex ecological niche models. *Environmental Modelling & Software,*
*125*, 104615. https://doi.org/https://doi.org/10.1016/j.envsoft.2019.104615
Araújo, M. B., Anderson, R. P., Márcia Barbosa, A., Beale, C. M., Dormann, C. F., Early, R., Garcia, R. A., Guisan,
A., Maiorano, L., & Naimi, B. (2019). Standards for distribution models in biodiversity assessments. *Science*
*Advances, 5*(1), eaat4858.
Ash, E., Kaszta, Ż., Noochdumrong, A., Redford, T., Chanteap, P., Hallam, C., Jaroensuk, B., Raksat, S.,
Srinoppawan, K., & Macdonald, D. W. (2021). Opportunity for Thailand's forgotten tigers: assessment of
the Indochinese tiger Panthera tigris corbetti and its prey with camera-trap surveys. *Oryx, 55*(2), 204-211.
https://doi.org/10.1017/S0030605319000589
Barbet-Massin, M., Jiguet, F., Albert, C. H., & Thuiller, W. (2012). Selecting pseudo-absences for species
distribution models: how, where and how many? *Methods in Ecology and Evolution, 3*(2), 327-338.
https://doi.org/https://doi.org/10.1111/j.2041-210X.2011.00172.x
Barve, N., Barve, V., Jiménez-Valverde, A., Lira-Noriega, A., Maher, S. P., Peterson, A. T., Soberón, J., &
Villalobos, F. (2011). The crucial role of the accessible area in ecological niche modeling and species
distribution modeling. *Ecological Modelling, 222*(11), 1810-1819.

Baston, D., ISciences, L., & Baston, M. D. (2021). Package 'exactextractr'. *R Foundation for Statistical*
*Computing. URL: https://cran. r-project. org*.
Bhandari, S., Crego, R. D., & Stabach, J. A. (2022). Spatial segregation between wild ungulates and livestock
outside protected areas in the lowlands of Nepal. *PLoS One, 17*(1), e0263122.
https://doi.org/10.1371/journal.pone.0263122
Booth, T. H., Nix, H. A., Busby, J. R., & Hutchinson, M. F. (2014). bioclim: the first species distribution modelling
package, its early applications and relevance to most current MaxEnt studies
[https://doi.org/10.1111/ddi.12144]. *Diversity and distributions, 20*(1), 1-9.
https://doi.org/https://doi.org/10.1111/ddi.12144
Bora, J. K., Vardhan, V., Vijh, R. K., Deshmukh, A. V., Srinivas, Y., Mungi, N. A., Goswami, S., Jhala, H., Chauhan,
J. S., Kumar, U., & Jhala, Y. (2024). Evaluating the potential for reintroducing the endangered wild water
buffalo (Bubalus arnee) in Kanha National Park, central India. *Restoration Ecology, 32*(3), e14079.
https://doi.org/https://doi.org/10.1111/rec.14079
Borowik, T., Pettorelli, N., Sönnichsen, L., & Jędrzejewska, B. (2013). Normalized difference vegetation index
(NDVI) as a predictor of forage availability for ungulates in forest and field habitats. *European Journal of*
*Wildlife Research, 59*(5), 675-682. https://doi.org/10.1007/s10344-013-0720-0
Buchhorn, M., Smets, B., Bertels, L., Lesiv, M., Tsendbazar, N.-E., Herold, M., & Fritz, S. (2019). Copernicus
Global Land Service: Land Cover 100m: collection 2: epoch 2015: Globe (Version V2.0.2) [Data set]. ESA
Living Planet Symposium 2019 (LPS2019), Milan, Italy: Zenodo.
Catullo, G., Masi, M., Falcucci, A., Maiorano, L., Rondinini, C., & Boitani, L. (2008). A gap analysis of Southeast
Asian mammals based on habitat suitability models. *Biological Conservation, 141*(11), 2730-2744.
https://doi.org/https://doi.org/10.1016/j.biocon.2008.08.019
Chaiyarat, R., Prasopsin, S., & Bhumpakphan, N. (2021). Food and nutrition of Gaur (Bos gaurus C.H. Smith,
1827) at the edge of Khao Yai National Park, Thailand. *Scientific Reports, 11*(1), 3281.
https://doi.org/10.1038/s41598-021-82858-1
Chaiyarat, R., Youngpoy, N., Kongsurakan, P., & Nakbun, S. (2019). Habitat preferences of reintroduced
banteng (*Bos javanicus*) into the Salakphra Wildlife Sanctuary, Thailand. *Wildlife Research, 46*(7), 573-586.
https://doi.org/https://doi.org/10.1071/WR18184
Choudhury, A. (2002). Distribution and conservation of the Gaur *Bos gaurus* in the Indian Subcontinent.
*Mammal Review, 32*(3), 199-226. https://doi.org/https://doi.org/10.1046/j.1365-2907.2002.00107.x
Crooks, K. R., Burdett, C. L., Theobald, D. M., Rondinini, C., & Boitani, L. (2011). Global patterns of
fragmentation and connectivity of mammalian carnivore habitat. *Philosophical transactions of the Royal*
*Society of London. Series B, Biological sciences, 366*(1578), 2642-2651.
https://doi.org/10.1098/rstb.2011.0120
Dewi, M. C., Sapari, I., & Santoso, E. (2020). THREATS AND CONSERVATION EFFORTS OF THE LAST REMAINING
BANTENG (Bos javanicus lowi) IN UNPROTECTED AREAS AT BELANTIKAN HULU, CENTRAL KALIMANTAN.
*Media Konservasi, 25*(1), 64-72.
Didan, K. (2015). *MOD13A2 MODIS/Terra Vegetation Indices 16-Day L3 Global 1km SIN Grid V006*
https://doi.org/https://doi.org/10.5067/MODIS/MOD13A2.006
Didan, K., Munoz, A. B., Solano, R., & Huete, A. (2015). MODIS vegetation index user's guide (MOD13 series).
*University of Arizona: Vegetation Index and Phenology Lab*.
Dormann, C. F. (2007). Promising the future? Global change projections of species distributions. *Basic and*
*applied ecology, 8*(5), 387-397.
Duckworth, J., Steinmetz, R., & Chaiyarat, R. (2008). Naemorhedus griseus. In *IUCN 2012. IUCN red list of*
*threatened species. Version 2012.2*.
Duckworth, J. W., Sankar, K., Williams, A. C., Samba Kumar, N., & Timmins, R. J. (2016). *Bos gaurus. The IUCN*
*Red List of Threatened Species 2016: e.T2891A46363646*. Retrieved 11 February 2021 from
https://dx.doi.org/10.2305/IUCN.UK.2016-2.RLTS.T2891A46363646.en.
Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: new 1-km spatial resolution climate surfaces for global land
areas. *International Journal of Climatology, 37*(12), 4302-4315.
https://doi.org/https://doi.org/10.1002/joc.5086
Franklin, J. (2009). *Mapping species distributions: spatial inference and prediction*. Cambridge University Press.
Gardner, P., Hedges, S., Pudyatmoko, S., Gray, T. N. E., & Timmins, R. J. (2016). *Bos javanicus. The IUCN Red List*
*of Threatened Species 2016: e.T2888A46362970*. Retrieved 11 February 2021 from
https://dx.doi.org/10.2305/IUCN.UK.2016-2.RLTS.T2888A46362970.en.
[Record #143 is using a reference type undefined in this output style.]

Giam, X., & Wilcove, D. S. (2012). The geography of conservation ecology research in Southeast Asia: current
biases and future opportunities. *Raffles Bull. Zool, 25*, 29-36.
Golding, N. (2014). GRaF: Species distribution modelling using latent Gaussian random fields. *R package*
*version 0.1-12*.
Gray, T., Hughes, A., Laurance, W., Long, B., Lynam, A., O'Kelly, H., Ripple, W., Seng, T., Scotson, L., &
Wilkinson, N. (2018). The wildlife snaring crisis: an insidious and pervasive threat to biodiversity in
Southeast Asia. *Biodiversity and Conservation, 27*. https://doi.org/10.1007/s10531-017-1450-5
Gray, T. N., & Phan, C. (2011). Habitat preferences and activity patterns of the larger mammal community in
Phnom Prich Wildlife Sanctuary, Cambodia. *The Raffles Bulletin of Zoology, 59*(2), 311-318.
Gray, T. N. E., Prum, S., Pin, C., & Phan, C. (2012). Distance sampling reveals Cambodia's Eastern Plains
Landscape supports the largest global population of the Endangered banteng Bos javanicus. *Oryx, 46*(4),
563-566. https://doi.org/10.1017/S0030605312000567
Hassanin, A. (2014). Systematic and evolution of Bovini. *Ecology, Evolution and Behavior of Wild Cattle:*
*Implications for Conservation. Cambridge University Press, Cambridge*, 7-20.
https://doi.org/https://doi.org/10.1017/CBO9781139568098.003
Hassell, J. M., Begon, M., Ward, M. J., & Fèvre, E. M. (2017). Urbanization and disease emergence: dynamics at
the wildlife–livestock–human interface. *Trends in ecology & evolution, 32*(1), 55-67.
Hastie, T. (2018). Gam: Generalized Additive Models.
Hein, Z. M., Williams, A. C., Soe, P., Cox, N. J., Htun, N. Z., Thaung, N. O., Aye, Y. Y., Htun, Y. L., & Yoganand, K.
(2020). Status of two species of threatened wild cattle (Bos gaurus and Bos javanicus birmanicus) in North
Zamari Wildlife Sanctuary, Bago Region, Myanmar. *BULLetin*.
Hijmans, R. J. (2023). *raster: Geographic Data Analysis and Modeling.* In https://rspatial.org/raster
IUCN. (2020). *The IUCN Red List of Threatened Species Version 2020-3*. Retrieved 14 February 2021 from
https://www.iucnredlist.org
IUCN. (2021). *The IUCN Red List of Threatened Species*. Retrieved 16 October from https://www.iucnredlist.org
IUCN-SSC AWCS Group. (2010). *IUCN Regional Conservation Strategy for Wild Cattle and Buffaloes in South-*
*east Asia, 2011 to 2020*. Oxford, UK, IUCN SSC AWCSG https://portals.iucn.org/library/node/9870
Jornburom, P., Duangchantrasiri, S., Jinamoy, S., Pattanavibool, A., Hines, J. E., Arnold, T. W., Fieberg, J., &
Smith, J. L. D. (2020). Habitat use by tiger prey in Thailand's Western Forest Complex: What will it take to
fill a half-full tiger landscape? *Journal for Nature Conservation, 58*, 125896.
https://doi.org/https://doi.org/10.1016/j.jnc.2020.125896
Karanth, K. K. (2016). Wildlife in the matrix: spatio-temporal patterns of herbivore occurrence in Karnataka,
India. *Environmental management, 57*(1), 189-206.
Karatzoglou, A., Smola, A., Hornik, K., & Zeileis, A. (2004). kernlab-an S4 package for kernel methods in R.
*Journal of statistical software, 11*(9), 1-20. https://doi.org/10.18637/jss.v011.i09
[Record #597 is using a reference type undefined in this output style.]
Kindt, R. (2018). Ensemble species distribution modelling with transformed suitability values. *Environmental*
*Modelling & Software, 100*, 136-145. https://doi.org/https://doi.org/10.1016/j.envsoft.2017.11.009
Liaw, A., & Wiener, M. (2002). Classification and regression by randomForest. *R news, 2*(3), 18-22.
Lim, H. Y., Gardner, P. C., Abram, N. K., Yusah, K. M., & Goossens, B. (2021). Identifying habitat and
understanding movement resistance for the Endangered Bornean banteng Bos javanicus lowi in Sabah,
Malaysia. *Oryx, 55*(1), 122-130. https://doi.org/10.1017/S0030605318001126
Liu, C., Newell, G., & White, M. (2019). The effect of sample size on the accuracy of species distribution
models: considering both presences and pseudo-absences or background sites. *Ecography, 42*(3), 535-548.
https://doi.org/https://doi.org/10.1111/ecog.03188
Marmion, M., Parviainen, M., Luoto, M., Heikkinen, R. K., & Thuiller, W. (2009). Evaluation of consensus
methods in predictive species distribution modelling [https://doi.org/10.1111/j.1472-4642.2008.00491.x].
*Diversity and distributions, 15*(1), 59-69. https://doi.org/https://doi.org/10.1111/j.1472-4642.2008.00491.x
Martinez, A. d. l. I., & Labib, S. (2023). Demystifying normalized difference vegetation index (NDVI) for
greenness exposure assessments and policy interventions in urban greening. *Environmental Research, 220*,
115155. https://doi.org/https://doi.org/10.1016/j.envres.2022.115155
McCullagh, P., & Nelder, J. A. (1989). Generalized linear models 2nd edition chapman and hall. *London, UK*.
McShea, W. J., Sukmasuang, R., Erickson, D. L., Herrmann, V., Ngoprasert, D., Bhumpakphan, N., & Davies, S. J.
(2019). Metabarcoding reveals diet diversity in an ungulate community in Thailand
[https://doi.org/10.1111/btp.12720]. *Biotropica, 51*(6), 923-937.
https://doi.org/https://doi.org/10.1111/btp.12720

Mendes, P., Velazco, S. J. E., Andrade, A. F. A. d., & De Marco, P. (2020). Dealing with overprediction in species
distribution models: How adding distance constraints can improve model accuracy. *Ecological Modelling,*
*431*, 109180. https://doi.org/https://doi.org/10.1016/j.ecolmodel.2020.109180

Menkham, K., Sukmasuang, R., Pla-ard, M., Charaspet, K., Panganta, T., Trisurat, Y., & Bhumpakphan, N.
(2019). Population and habitat use of Asian elephants (Elephas maximus) and five ungulate species in Khao
Ang Rue Nai Wildlife Sanctuary, Chachoengsao Province, Thailand. *Biodiversitas Journal of Biological*
*Diversity, 20*(8).

Mori, E., Nerva, L., & Lovari, S. (2019). Reclassification of the serows and gorals: the end of a neverending
story? [https://doi.org/10.1111/mam.12154]. *Mammal Review, 49*(3), 256-262.
https://doi.org/https://doi.org/10.1111/mam.12154

Nguyen, M. H. (2009). The status of Vulnerable gaur Bos gaurus and Endangered banteng Bos javanicus in Ea
So Nature Reserve and Yok Don and Cat Tien National Parks, Vietnam [Article]. *Oryx, 43*(1), 129-135.
https://doi.org/10.1017/s0030605307000440

Olson, D. M., Dinerstein, E., Wikramanayake, E. D., Burgess, N. D., Powell, G. V. N., Underwood, E. C., D'amico,
J. A., Itoua, I., Strand, H. E., Morrison, J. C., Loucks, C. J., Allnutt, T. F., Ricketts, T. H., Kura, Y., Lamoreux, J.
F., Wettengel, W. W., Hedao, P., & Kassem, K. R. (2001). Terrestrial Ecoregions of the World: A New Map of
Life on Earth: A new global map of terrestrial ecoregions provides an innovative tool for conserving
biodiversity. *BioScience, 51*(11), 933-938. https://doi.org/10.1641/0006-
3568(2001)051[0933:Teotwa]2.0.Co;2

Penjor, U., Kaszta, Ż., Macdonald, D. W., & Cushman, S. A. (2021). Prioritizing areas for conservation outside
the existing protected area network in Bhutan: the use of multi-species, multi-scale habitat suitability
models. *Landscape Ecology, 36*(5), 1281-1309. https://doi.org/10.1007/s10980-021-01225-7

Peterson, A. T., & Soberón, J. (2012). Species distribution modeling and ecological niche modeling: getting the
concepts right. *Natureza & Conservação, 10*(2), 102-107. https://doi.org/10.4322/natcon.2012.019

Phillips, S. J., Anderson, R. P., & Schapire, R. E. (2006). Maximum entropy modeling of species geographic
distributions. *Ecological Modelling, 190*(3-4), 231-259.

Phoonjampa, R., Steinmetz, R., Phumanee, W., Bunchornratana, K., Kaewsrisod, T., Srirattanaporn, S.,
Taraphibarl, K., Bejraburnin, T., & Bhumpakphan, N. (2021). Recolonization of Former Range by Endangered
Banteng Bos j avanicus in Mae Wong National Park, Thailand. *Tropical Conservation Science, 14*,
19400829211065359.

Pintana, D., & Lakamavichian, S. (2013). Ecology and conservation of Goral outside protected areas by
community-based approach in Thailand. *Tigerpaper, 40*(3), 12-21.

Prayoon, U., Suksavate, W., Chaiyes, A., Winitpornsawan, S., Tunhikorn, S., Faengbubpha, K., Angkaew, C.,
Pattanakiat, S., & Duengkae, P. (2021). Past, present and future habitat suitable for gaur (Bos gaurus) in
Thailand. *Agriculture and Natural Resources, 55*(5), 743–756-743–756.

R Core Team. (2020). *R: A language and environment for statistical computing.* In R Foundation for Statistical
Computing.

Ripple, W. J., Newsome, T. M., Wolf, C., Dirzo, R., Everatt, K. T., Galetti, M., Hayward, M. W., Kerley, G. I. H.,
Levi, T., Lindsey, P. A., Macdonald, D. W., Malhi, Y., Painter, L. E., Sandom, C. J., Terborgh, J., & Van
Valkenburgh, B. (2015). Collapse of the world's largest herbivores. *Science Advances, 1*(4), e1400103-
e1400103. https://doi.org/10.1126/sciadv.1400103

Rizal, A. R. M., Magintan, D., Rahman, Z. A., Nibong, O., Rahim, M. A. B., Hassan, Z. M., & Patah, P. A. (2020).
Home range size estimation of a satellite collared female gaur at Ulu Jelai Forest Reserve, Pahang by using
minimum convex polygon and fixed kernel analysis: a preliminary study. *Journal of Wildlife and Parks, 35*,
49-59.

Romero, G. Q., Gonçalves-Souza, T., Vieira, C., & Koricheva, J. (2015). Ecosystem engineering effects on species
diversity across ecosystems: a meta-analysis [https://doi.org/10.1111/brv.12138]. *Biological Reviews,*
*90*(3), 877-890. https://doi.org/https://doi.org/10.1111/brv.12138

Royle, J. A., Chandler, R. B., Yackulic, C., & Nichols, J. D. (2012). Likelihood analysis of species occurrence
probability from presence-only data for modelling species distributions [Article]. *Methods in Ecology and*
*Evolution, 3*(3), 545-554. https://doi.org/10.1111/j.2041-210X.2011.00182.x

Simcharoen, A., Simcharoen, S., Duangchantrasiri, S., Bump, J., & Smith, J. L. D. (2018). Tiger and leopard diets
in western Thailand: Evidence for overlap and potential consequences. *Food Webs, 15*, e00085.
https://doi.org/https://doi.org/10.1016/j.fooweb.2018.e00085

Smith, A. B., Godsoe, W., Rodríguez-Sánchez, F., Wang, H.-H., & Warren, D. (2019). Niche Estimation Above
and Below the Species Level. *Trends in ecology & evolution, 34*(3), 260-273.
https://doi.org/https://doi.org/10.1016/j.tree.2018.10.012

Soberón, J., & Nakamura, M. (2009). Niches and distributional areas: Concepts, methods, and assumptions.
*Proceedings of the National Academy of Sciences, 106*(Supplement 2), 19644.
https://doi.org/10.1073/pnas.0901637106
Soberon, J., & Peterson, A. T. (2005). Interpretation of Models of Fundamental Ecological Niches and Species'
Distributional Areas. *Biodiversity Informatics, 2*. https://doi.org/10.17161/bi.v2i0.4
Steinmetz, R. (2004). Gaur (Bos gaurus) and Banteng (B. javanicus) in the lowland forest mosaic of Xe Pian
Protected Area, Lao PDR: abundance, habitat use, and conservation. *Mammalia, 68*(2-3), 141-157.
https://doi.org/https://doi.org/10.1515/mamm.2004.015
Steinmetz, R., Chutipong, W., Seuaturien, N., & Chirngsaard, E. (2008). Community Structure of Large
Mammals in Tropical Montane and Lowland Forest in the Tenasserim-Dawna Mountains, Thailand.
*Biotropica, 40*(3), 344-353. www.jstor.org/stable/30043637
Stevens, F. R., Gaughan, A. E., Linard, C., & Tatem, A. J. (2015). Disaggregating Census Data for Population
Mapping Using Random Forests with Remotely-Sensed and Ancillary Data. *PLoS One, 10*(2), e0107042.
https://doi.org/10.1371/journal.pone.0107042
Sukmasuang, R., Charaspet, K., Panganta, T., Pla-ard, M., Khioesree, N., & Thongbanthum, J. (2020). Diversity,
abundance, activity period, and factors affecting the appearance of wildlife around the corridors between
Khao Yai- Thap Lan National Parks, Thailand by camera trapping. *Biodiversitas Journal of Biological*
*Diversity, 21*(5). https://doi.org/10.13057/biodiv/d210563
Suttidate, N., Steinmetz, R., Lynam, A. J., Sukmasuang, R., Ngoprasert, D., Chutipong, W., Bateman, B. L., Jenks,
K. E., Baker-Whatton, M., Kitamura, S., Ziółkowska, E., & Radeloff, V. C. (2021). Habitat connectivity for
endangered Indochinese tigers in Thailand. *Global Ecology and Conservation, 29*, e01718.
https://doi.org/https://doi.org/10.1016/j.gecco.2021.e01718
Tanasarnpaiboon, S. (2016). *Gaur (Bos gaurus) abundance, distribution, and habitat use patterns in Kuiburi*
*National Park, Southwestern Thailand*. The University of Nebraska-Lincoln.
https://digitalcommons.unl.edu/dissertations/AAI10102227
UNEP-WCMC, & IUCN. (2021). *Protected Planet: The World Database on Protected Areas (WDPA) [Online]*
www.protectedplanet.net
Velazco, S. J. E., Villalobos, F., Galvão, F., & De Marco Júnior, P. (2019). A dark scenario for Cerrado plant
species: Effects of future climate, land use and protected areas ineffectiveness
[https://doi.org/10.1111/ddi.12886]. *Diversity and distributions, 25*(4), 660-673.
https://doi.org/https://doi.org/10.1111/ddi.12886
Yang, J., Zhu, G.-F., Jiang, J., Xiang, C.-L., Gao, F.-L., & Bao, W.-D. (2019). Non-invasive genetic analysis indicates
low population connectivity in vulnerable Chinese gorals: concerns for segregated population
management. *Zoological Research, 40*(5), 439.
Zhang, L., Liu, S., Sun, P., Wang, T., Wang, G., Zhang, X., & Wang, L. (2015). Consensus Forecasting of Species
Distributions: The Effects of Niche Model Performance and Niche Properties. *PLoS One, 10*(3), e0120056.
https://doi.org/10.1371/journal.pone.0120056
Zhang, Y., Colli, L., & Barker, J. (2020). Asian water buffalo: domestication, history and genetics. *Animal*
*Genetics, 51*, 177-191. https://doi.org/10.1111/age.12911
Zhu, G., Fan, J., & Peterson, A. T. (2021). Cautions in weighting individual ecological niche models in ensemble
forecasting. *Ecological Modelling, 448*, 109502.
https://doi.org/https://doi.org/10.1016/j.ecolmodel.2021.109502

831