Systematic creation and phenotyping of Mendelian disease models in *C. elegans*: towards large-scale drug repurposing

Thomas J. O'Brien[1,2*], Ida L. Barlow[1,2*], Luigi Feriani[1,2] and André E.X. Brown[1,2]

[1]Institute of Clinical Sciences, Imperial College London, London, UK
[2]MRC London Institute of Medical Sciences, London, UK
*These authors contributed equally
Correspondence: andre.brown@imperial.ac.uk

Abstract

There are thousands of Mendelian diseases with more being discovered weekly and the majority have no approved treatments. To address this need, we require scalable approaches that are relatively inexpensive compared to traditional drug development. In the absence of a validated drug target, phenotypic screening in model organisms provides a route for identifying candidate treatments. Success requires a screenable phenotype. However, the right phenotype and assay may not be obvious for pleiotropic neuromuscular disorders. Here we show that high-throughput imaging and quantitative phenotyping can be conducted systematically on a panel of *C. elegans* disease model strains. We used CRISPR genome-editing to create 25 worm models of human Mendelian diseases and phenotyped them using a single standardised assay. All but two strains were significantly different from wild-type controls in at least one feature. The observed phenotypes were diverse, but mutations of genes predicted to have related functions led to similar behavioural differences in worms. As a proof-of-concept, we performed a drug repurposing screen of an FDA approved compound library, and identified two compounds that rescued the behavioural phenotype of a model of UNC80 deficiency. Our results show that a single assay to measure multiple phenotypes can be applied systematically to diverse Mendelian disease models. The relatively short time and low cost associated with creating and phenotyping multiple strains suggests that high-throughput worm tracking could provide a scalable approach to drug repurposing commensurate with the number of Mendelian diseases.

Introduction

By definition, rare genetic diseases each affect a small number of patients. However, because there are many kinds of rare diseases, they are collectively common and affect more than 300 million people worldwide (Nguengang Wakap et al., 2020). Over 80% of rare genetic diseases affect children and many have severe symptoms (Bavisetty et al., 2013). Furthermore, 95% of rare diseases have no approved treatment (Nguengang Wakap et al., 2020). Given the severity and prevalence of rare diseases, the lack of treatments represents an important unmet need. Genome and exome sequencing has become faster and cheaper, enabling its widespread use in the diagnosis of genetic disease. In turn, this has led to the rapid identification of genetic lesions associated with rare genetic diseases (Gonzaga-Jauregui et al., 2012). Genetic diagnosis is an important first step, but often presents patients and physicians with a research problem rather than a treatment.

Genetic diagnoses rarely lead directly to a drug target hypothesis, especially for loss-of-function mutations. In the absence of a validated drug target, phenotypic screens in a model organism provide an alternative route to candidate treatments. Three things are required for a phenotypic screen to be successful: (1) a conserved gene; (2) a genetically tractable organism or culture system in which the disease mutation can be introduced; and (3) a measurable phenotype compatible with high-

throughput screening. For the model nematode *C. elegans*, more than half of the
human genes associated with rare genetic diseases in the OMIM database are
conserved (Kropp et al., 2021) and CRISPR-based editing has made the generation
of *in vivo* models containing disease-relevant mutations faster and easier. However,
the third condition, a screenable phenotype, remains a bottleneck for many
mutations and models.
If all three conditions are met, a phenotypic screen for candidate treatments
can be performed. Because there is limited investment in drug development for rare
diseases, lead optimisation and safety testing may not be possible. Hence, there is a
particular focus on screening approved drugs that have already been shown to be
safe and bioavailable in humans to reduce the cost and time it takes to translate
screening hits to human trials (Roessler et al., 2021). Two recent examples of
promising repurposing screens using multiple model organisms illustrate this
potential.
Firstly, modelling amyotrophic lateral sclerosis (ALS) with disease-associated
genetic variants of conserved genes in *C. elegans*, zebrafish, and mice led to clinical
trials for repurposing of the antipsychotic drug pimozide. By exploiting high
throughput drug screening in *C. elegans* carrying point mutations in *TDP-43*, a panel
of drugs were identified that rescued ALS movement and neurodevelopment
phenotypes in all animal models. Early clinical trials found pimozide improved clinical
and physiological outcomes in ALS patients (Patten et al., 2017). Secondly, the
glycosylation disorder, PMM2-CDG, is caused by mutations in *PMM2* and presents
clinically with developmental delay, psychomotor retardation, and axial hypotonia,
among other symptoms. *C. elegans* carrying the same disease-associated mutation
in their endogenous *pmm-2* gene exhibited larval arrest upon pharmacological ER
stress. *In vivo* screening identified two compounds that rescued development in *C.
elegans* and enzymatic activity in PMM2-CDG patient fibroblasts (Iyer et al., 2019).
One of the compounds, the aldose reductase inhibitor epalrestat, is currently being
tested in clinical trials for PMM2-CDG patients (National Clinical Trial number:
NCT04925960).
A screenable phenotype was critical for the success of these studies. In the
ALS model, worms became paralysed after 2 hours in liquid and in the PMM2-CDG
model, brood size was severely affected. The majority (~74%) of rare genetic
diseases affect nervous system function (Lee et al., 2020), and many of these
mutations will not cause strong effects on development or gross motility when
mutated in worms. High-content phenotyping that measures multiple features
simultaneously could be useful in these cases to identify differences between strains
that are not captured with univariate measures. In this study, we use a combination
of high-throughput imaging (Barlow et al., 2022) and quantitative phenotyping (Javer,
Currie, et al., 2018; Javer, Ripoll-Sánchez, et al., 2018) to phenotype a diverse panel
of 25 *C. elegans* strains with mutations in orthologs of human disease genes. All but
two of the mutant strains had detectable phenotypes compared to controls in at least
one aspect of their morphology, posture, or motion. No single feature was different in
all strains highlighting the need for a multidimensional phenotype. Mutation of genes
predicted to have similar functions (e.g., *bbs-1*, *bbs-2,* and *tub-1*) led to similar
phenotypes. As a proof-of-concept, we then performed a repurposing screen using a
library of 743 FDA approved compounds to identify drugs that improved the
behavioural phenotype of *unc-80* loss-of-function mutants. Liranaftate and
atorvastatin rescued the core behavioural features associated with *unc-80* loss of
function and did not cause a large number of detectable side effects.

The ability to detect phenotypic difference in diverse strains using a
standardised 16-minute assay will make it possible to perform repurposing screens
for existing and newly described rare diseases efficiently.

Results

Selection and molecular characterisation of a diverse panel of C. elegans disease
models

The problem of large-scale disease modelling can be thought of as a problem of
sampling and connecting two related genotype-phenotype (GP) maps (Fig. 1A). The
first is the human disease map. The genomic space is sampled by mutation and
mating in humans and the map is elucidated by sequencing and clinical phenotyping
of patients with genetic diseases. The second is the model organism map. Here, the
genomic space is sampled using mutagenesis and genome editing and the map is
elucidated using laboratory experiments. Given the accelerated characterisation of
the human GP map, we sought to accelerate the characterisation of the
corresponding *C. elegans* GP map. We follow a phenolog-inspired approach to
connecting the two maps. Provided there is a causal connection between a genetic
variant and a disease in humans and that the causal gene is conserved in worms,
the observed worm phenotype may be a useful disease model even if the connection
to the human phenotype is non-obvious (McGary et al., 2010).

To identify a set of *C. elegans* genes that samples a broad genotypic space
with connections to the human disease GP map, we first filtered the full database of
human-*C. elegans* gene orthologs, Ortholist 2 (Kim et al., 2018), using three criteria:
(1) at least two orthology prediction algorithms agree the human and worm genes
are orthologs; (2) the WormBase (version WS270) (Harris et al., 2020) gene
description includes either 'neuro' or 'musc'; and (3) mutations in the human gene
are associated with a Mendelian disease. A total of 767 *C. elegans* genes with 2558
orthologous human protein-coding genes met these criteria (Figure 1B). We used
keyword searches to further reduce the list to 543 genes. Genes were kept if their
WormBase gene description included any of the following: ion, ion channel,
dopamine, serotonin, 5-HT, glutamate, acetylcholine, behavior, behaviour, disease,
epilepsy, autism, Parkinson, schizophrenia, bipolar, ADHD, seizure, GPCR, coupled
receptor, antidepressant, antipsychotic). We prioritised genes associated with
autosomal recessive disease in humans to increase the chance that a gene deletion
would yield an appropriate model, and we focussed on those where loss-of-function
mutants were known or likely to be viable. To get the list of 25 genes presented in
this study, the final selection criterion was subjective interestingness based on our
prior knowledge, WormBase gene descriptions, and/or brief literature searches.

In the final panel of 25 worm genes selected, 22 genes have >60% sequence
similarity to their human ortholog and 11 of the genes share >90% sequence
similarity to their human counterpart (Figure 1C). Furthermore, 24/25 genes are
predicted to be orthologous to human genes across >3 orthology prediction
algorithms (Figure 1C), and all selected genes have a BLAST E-value (McGinnis &
Madden, 2004) smaller than 3×10^{-12} (Supplementary Table 1).

Large CRISPR-Cas9 deletions (mean 4.4 kb) were made in each of the target
genes to generate 25 strains containing a >55% deletion (average 76%) of the
chosen gene. We direct the reader to the strain-specific gene cards (Supplementary
Information) for specific details on the size and position of genomic deletions for
each of the individual mutants as well as a brief summary of the associated disease
and worm phenotypes. Although we have focused on genes expected to have direct

Figure 1: Overview of genotype-phenotype mapping and disease model panel.
Disease modelling as genotype-phenotype mapping in humans and a model organism. Arrows show progression from symptom identification (a phenotype outside the healthy range) to genotyping (placing a patient in genotype space by identifying a genetic variant), disease model creation (making a corresponding mutation in a model organism), and model organism phenotyping. Phenotypic drug screens identify compounds that move a disease model towards the wild-type phenotype (green arrow). These candidates can then be tested in humans (dashed green arrow). Thin lines show other symptom-gene-ortholog-phenotype connections. (B) Venn diagram showing number of conserved genes (Ortholist 2), those involved in neuron or muscle function (Wormbase), and those associated with human genetic disorders according to the Online Mendelian Inheritance in Man (OMIM) database. (C) Sequence similarity between human and C. elegans genes and the total number of orthology programs predicting that the gene is an ortholog.

effects on neurons or muscles, even this small subset of disease-associated genes
is predicted to affect diverse cellular processes including
neurotransmission, excitability, development, and cellular structure (Table 1).
According to OMIM, the mutated genes are associated with 31 rare genetic disorders
in humans including intellectual disability, developmental delay, and disorders
affecting the muscular and/or nervous systems, and are associated with >70 clinical
presentations of disease according to the Human Disease Ontology database
(Schriml et al., 2022) (Table 1).
161
162 *Disease-associated mutations result in diverse phenotypes that are captured by*
163 *multidimensional behavioural fingerprints*
164 Given that there are thousands of rare diseases, methods to characterise the
165 corresponding disease models should be as scalable as possible. Scalability is
166 improved if a single standardised assay can be used to detect model phenotypes

Table 1. Summary of the associated human ortholog(s), predicted functional class and key associated human disease phenotypes (Human Phenotype Ontology database) for each C. elegans disease model mutant.

regardless of the mutation. High-resolution worm tracking and behavioural
fingerprinting is a promising candidate for a widely applicable assay because it is
both fast and multidimensional: a single minutes-long recording can capture
differences in morphology, posture, and behaviour that can classify the effect of
mutations and drug treatments (Baek et al., 2002; Barlow et al., 2022; Geng et al.,
2004; Javer, Ripoll-Sánchez, et al., 2018; McDermott-Rouse et al., 2021; Perni et al.,
2018; Ramot et al., 2008; Restif et al., 2014; Swierczek et al., 2011; Tsibidis &
Tavernarakis, 2007; S. J. Wang & Wang, 2013; Yemini et al., 2013).
175   We tracked disease model worms on 96-well plates (Figure 2A) in three
periods: (1) a 5-minute baseline recording (prestim); (2) a 6-minute video with three
ten-second blue light stimuli separated by 90 seconds (blue light), and (3) a 5-minute
post-stimulus recording (poststim). For each recording we extract 2763 features
covering morphology, posture, and motion and concatenate the feature vectors
together to represent the phenotype of each strain (8289 features in total). Many of
these features are highly correlated and so for clustering we use a pre-selected
subset of 256 features (Javer, Ripoll-Sánchez, et al., 2018) (768 across the three
recordings). Using these reduced feature vectors, we performed hierarchical
clustering across all strains which demonstrates phenotypic diversity (many of the
strains are both different from the N2 wild-type strain and from each other) as well as
some similarities. For example, *bbs-1, bbs-2* and *tub-1* mutants affect cilia formation
and result in similar phenotypes that cluster together. The same is true for *kcc-2, snf-*
11 and *unc-25* mutants, which all affect GABAergic function. An interactive heatmap
containing all behavioural feature vectors extracted for every disease model strain is
also available for download (https://doi.org/10.5281/zenodo.12684130).
191   Principal component analysis (PCA) of the same data, but now separated into
prestimulus, blue light, and poststimlus recordings again shows the separation of
many of the mutant strains from the wild-type N2 (Figure 2D). The effect of the blue
light stimulus on behaviour is clear from the motion of the points through this
phenotype space and also reveals new phenotypes. While most strains show a
partial recovery in the post-stimulus recording, some strains including *bbs-1*, *bbs-2*,
glr-1, and *unc-43* display a sustained photophobic response (Figure 2D).
From the panel of 25 disease model strains, 23 are significantly different from wild-
type in at least one feature (Figure 2E). The strains that are most different from wild-
type show significant differences across a large number of features. These strains
are particularly useful for high-throughput drug screens because the phenotypes can
be reliably detected with a small number of replicates. Strain-specific summaries
(gene cards) of each disease model mutant, its associated phenotype, and individual
molecular characteristics are available in the supplementary information. Here we
provide more detailed characterisation of two classes of mutants modelling
ciliopathies and channelopathies in humans.
207
Ciliopathies
Heritable mutations that affect cilia function are associated with a group of rare
genetic disorders (ciliopathies) that share diverse symptoms including retinal
dystrophy, developmental polydactyly, obesity, cognitive impairment, and renal
dysfunction (Reiter & Leroux, 2017). The pleotropic nature of ciliopathies is
highlighted by significant interfamilial and intrafamilial phenotypic variability
(Shaheen et al., 2016), and the resulting poor gene-phenotype correlation
significantly complicates the discovery of effective therapeutics. As a result, current
treatment regimens are based on treating individual symptoms.

Figure 2: Diverse multidimensional behavioural phenotypes are obtained across the diverse panel of disease model mutants.
(A) Representative field of view for a single camera channel with individual skeletonised worm images of *bbs-1(syb1588)* (top) and N2 (bottom) imaged on the same plate viewed using Tierpsy Tracker. **(B)** Representative behavioural phenotypes extracted by Tierpsy, representing changes in morphology, posture and locomotion of different disease mutant strains. *P*-values are for comparisons to wild-type N2 worms, block permutation t-test, corrected for multiple comparisons. **(C)** Hierarchical clustering of behavioural fingerprints. Features are *Z*-normalised. The top barcode shows the period of image acquisition where the behavioural feature was extracted by Tierpsy: pre-stimulation (pink), blue light (blue) and post-stimulation (green). **(D)** Principal component analysis of the disease model mutants and N2 reference (blue). Strains move in phenospace between pre-stimulus (circular points), blue light (crosses) and post-stimulation (squares) recordings. Strains with an aberrant blue light response are shown in red. Error bars represent standard error of the mean. **(E)** The total number of statistically significant behavioural features for each strain compared to N2 (from the total set of 8289 features extracted by Tierpsy). *P*-values for each feature were calculated using block permutation t-tests, using n = 100,000 permutations and *P* < 0.05 considered statistically significant after correcting for multiple comparisons using the Benjamini-Ye-kutieli method.

In the longlist of 543 disease genes, there were 35 genes (6.4% of the total list) predicted to be involved in cilia function. We selected four genes (*bbs-1*, *bbs-2*, *tmem-231* and *tub-1*) involved in regulating nonmotile (primary) cilia development and function that have confirmed expression in the cilia of amphid sensory neurons (Taylor et al., 2021; Ward et al., 2008; Zhang et al., 2022).

Of the genes affecting cilia function, two are orthologs of the Bardet-Biedl syndrome (BBS) family of genes (*bbs-1* and *bbs-2*). BBS is a pleiotropic syndrome often diagnosed in late childhood and is characterised by retinitis pigmentosa, cognitive impairment, obesity, renal dysfunction, hypogonadism, and polydactyly (Forsythe & Beales, 2013). The most prevalent BBS mutations are in *BBS1*, accounting for 23.4% of cases (Forsyth & Gunay-Aygun, 2023), and with 50% embryonic lethality and highly variable phenotypes, *Bbs1-/-* mouse models demonstrate model validity but have not yet yielded insights into potential therapies (Forsyth & Gunay-Aygun, 2023). Mutations in *BBS2* account for 8% of BBS cases (Forsythe & Beales, 2013) and are also implicated in other ciliopathies, such as Meckel syndrome (Karmous-Benailly et al., 2005).

Mutations in *bbs-1* and *bbs-2* led to strong phenotypes including changes in morphology, posture, and locomotion (Figure 3A-D). *bbs-1(syb1588)* and *bbs-2(syb1547)* mutants were shorter, wider, and had decreased curvature compared to wild-type worms (Figures 3A, 3B and 3C). They are also hyperactive, with a greater fraction of worms moving during pre-stimulus (baseline) recordings (see individual gene cards), and had faster body bends (derivative of curvature) compared to N2 (Figure 3D and 3F). BBS proteins also regulate the *C. elegans* photoreceptor protein LITE1 in ASH sensory neurons through a DLK-MAPK signalling pathway independent of their function in cilia (Zhang et al., 2022). Consistent with this, we find that *bbs-1(syb1588)* and *bbs-2(syb1547)* mutants have attenuated blue light sensitivity, with both strains exhibiting a delayed forward, but enhance backward photophobic escape response upon 10-second stimulation with blue light (Figures 3F-H).

The vertebrate family of tubby-like proteins (TUB, TULP) are required for correct GPCR localisation and intra-flagellar transport in primary cilia (Mukhopadhyay & Jackson, 2011). In humans, mutations in this family are associated with rod-cone dystrophy, obesity, and retinitis pigmentosa (North et al., 1997). Furthermore, *Tub1-/-* mice reflect the associated human disease pleiotropy, with insulin resistance, obesity, and cochlea and visual degeneration (Sun et al., 2012). *C. elegans* have a single ortholog of tubby-like proteins, *tub-1*, and mutants have defects in chemotaxis and insulin signalling as well as increased lipid accumulation (Mak et al., 2006).

tub-1(syb1562) mutants are shorter, wider, and hyperactive compared to wild-type worms, similar to *bbs-1* and *bbs-2* mutants (Figures 3A, 3B and 3E). They also have a defect in their response to blue light (Figures 3F-H). Unlike BBS mutants, cilia integrity is maintained in *tub-1* loss-of-function mutants (Mak et al., 2006), which could explain the similar but less severe phenotype of *tub-1* mutants compared to *bbs-1* and *bbs-2* mutants.

Compartmentalisation of cilia-specific signalling components is regulated by a complex of transmembrane proteins including TMEM231 and mutations in this compartmentalisation complex are associated with neurodevelopmental limb defects and pathologies of the brain and kidneys found in Joubert and Meckel syndrome (Chih et al., 2012). Mouse *Tmem231-/-* mutations are embryonic lethal and the *C. elegans* ortholog *tmem-231* has been shown to have conserved molecular and cellular function (Roberson et al., 2015). However, unlike the other cilia-related

Figure 3: Ciliopathies disease model phenologs.
(A-D) Key behavioural features altered in loss-of-function mutant strains associated with ciliopathies: *bbs-1(syb1588)*, *bbs-2(syb1547)*, *tub-1(syb1562)* and *tmem-231(syb1575)*, under baseline (pre-stimulus) imaging conditions. Individual points marked on the box plots are well averaged values (three worms per well) for each feature across the independent days of tracking. *P*-values are for comparisons to wild-type N2 worms using block permutation t-tests (*n* = 100,000 permutations, correcting for multiple comparisons using the Benjamini-Yekutieli method). **(E-H)** Changes in selected features in response to stimulation with a single 10-second blue light pulse (blue shaded region). Feature values were calculated using 10 second windows centred around 5 seconds before, 10 seconds after, and 20 seconds after the beginning of each blue light pulse.
(I) Heatmap of the entire set of 8289 behavioural features extracted by Tierpsy for the disease model strains associated with ciliopathies and N2. The 'stim type' barcode denotes when during image acquisition the feature was extracted: pre-stimulation (pink), blue light stimulation (blue) and post-stimulation (green). Asterisks show the location of selected features present in A-E.

268   mutants, *tmem-231* mutants do not have significant differences in body morphology
269   or posture compared to N2 (Figure 3). We do note that *tmem-231(syb1575)* is
270   slightly more active during baseline tracking (see strain-specific gene card).
271
272   *Channelopathies*
273   The Na+ cation leak channel (NALCN) is expressed throughout the central
nervous system, and in parts of the endocrine (pancreas, adrenal, thyroid gland),
respiratory and cardiac systems (Cochet-Bissuel et al., 2014). NALCN is a voltage-
independent, nonselective cation channel that regulates resting potential (Senatore
et al., 2013), and plays a role in neuromodulation by neurotransmitters (Cochet-
Bissuel et al., 2014). Mutations in NALCN are associated with neuromuscular
disorders including severe hypotonia, infantile neuroaxonal dystrophy (INAD),
congenital contractures, cognitive delay, autism, epilepsy, bipolar disorder, and
cardiac/respiratory problems (Aoyagi et al., 2015; Cochet-Bissuel et al., 2014; Gal et
al., 2016). NALCN-knock-out mice die 12 hours after birth (Lu et al., 2007), so
development of appropriate non-mammalian animal models to understand the
associated human diseases is essential.
285   *C. elegans* encodes two functionally redundant, but differentially expressed,
NALCN homologs, NCA-2 and UNC-77, whose proper expression and axonal
localization in cholinergic neurons is regulated by UNC-80 (Zhou et al., 2019). Unlike
murine models, *nca-2*, *unc-77*, and *unc-80* loss-of-function mutants are viable.
Similar to *Drosophila melanogaster* or mouse neonates lacking the cation leak
channel (Cochet-Bissuel et al., 2014), loss-of-function mutations in *nca-2* and *unc-80*
result in morphological changes in *C. elegans*: both mutants are significantly shorter
than wild-type N2 (Figure 4A).
293   Consistent with their differential expression (Jospin et al., 2007; Yeh et al.,
2008), we observe phenotypic differences between *nca-2(syb1612)* and *unc-*
77(syb1688) deletion mutants (Figure 4). Gain-of-function mutations in *unc-77* have
previously been reported to cause deeper body bends (Topalidou et al., 2017)
whereas *unc-77(syb1688)* deletion mutants have decreased curvature (Figure 4B).
In contrast, *nca-2(syb1612)* has no significant change in curvature, but is slower than
the wild-type strain (Figures 4B-C). We find that *unc-80(syb1531)* mutants, which
affect the localisation of NALCN channel subunits, have the most severe phenotype
exhibiting a decrease in both curvature and speed (Figures 4B-4C).
302   Mutations in both NALCN channel subunits in *C. elegans* cause a 'fainter'
phenotype when stimulated mechanically or immersed in liquid (Pierce-Shimomura
et al., 2008). Similarly, we found that a ten second blue light pulse resulted in
increased post-exposure pausing and a decreased forward escape response in *nca-*
2(syb1612) and *unc-80(syb1531)* single mutants (Figures 4E-H). *unc-77(syb1688)*
mutants did not 'faint' after blue light exposure, but consistent with findings that the
NALCN channelsome regulates reversal behaviour (Zhou et al., 2019) both *unc-*
77(syb1688) and *unc-80(syb1531)* fail to initiate a backwards escape response.
Pausing and 'fainting' after blue light exposure represents a novel screenable
phenotype for NALCN-channelopathies.
312
Drug repurposing screen in unc-80 mutants
Because *unc-80* mutants have a clear behavioural phenotype without a strong
developmental phenotype, we reasoned that they were a useful test case for a drug
repurposing screen using an acute 4-hour treatment which might rescue behavioural
'symptoms' in fully developed animals. When screening compound libraries, it is not
practical to perform a large number of replicates which would be necessary to detect

Figure 4: NALCN disease model phenologs.

(A-D) Key behavioural and postural features altered in loss-of-function mutant strains associated with NALCN mutants: *nca-2(syb1612)*, *unc-77(syb1688)* and *unc-80(syb1531)*, under baseline (pre-stimulus) imaging conditions. Individual points marked on the box plots are well averaged values (three worms per well) for each feature across the independent days of tracking. *P*-values are for comparisons to wild-type N2 worms using block permutation t-tests (n = 100,000 permutations correcting for multiple comparisons using the Benjamini-Yekutieli method). **(E-H)** Changes in selected features in response to stimulation with a single 10-second blue light pulse (blue shaded region). Feature values were calculated using 10 second windows centred around 5 seconds before, 10 seconds after, and 20 seconds after the beginning of each blue light pulse. **(E)** A representative 'fainting phenotype' for *unc-80(syb1531)* and *nca-2(syb1612)*, characterised by an increase in pausing following the cessation of stimulation with blue light. **(I)** Heatmap of the entire set of 8289 behavioural features extracted by Tierpsy for the disease model strains associated with NALCN disease and N2. The 'stim type' barcode denotes when during image acquisition the feature was extracted: pre-stimulation (pink), blue light stimulation (blue) and post-stimulation (green). Asterisks show the location of selected features present in A-D.

subtle phonotypes and overcome the reduced statistical power that comes from
correcting for multiple comparisons in a large feature set (Supplementary Figure 1).
We therefore defined a reduced set of core features to capture the *unc-80*
behavioural phenotype consisting of curvature, speed, and fraction of paused worms
all during blue light stimulation (Figure 4B-D).
For the repurposing screen, we used a library of 743 FDA approved drugs.
We prepared tracking plates where each well contained a drug at a final
concentration of 100 μM with 1% DMSO. Three adult worms were added to each
well, incubated for 4 hours, and then tracked. The screen was repeated across three
independent tracking days with three independent wells per day (9 total well
replicates). Hits were defined as compounds that significantly improved all three of
the core *unc-80* features, meaning that treated worms were significantly different
from *unc-80* DMSO controls and that the direction of the effect was towards wild-type
controls. These hit compounds shift *unc-80* towards N2 in a three-dimensional
phenotype space (Figure 5A). To test the utility of reducing the feature set in a
screen compared to testing a larger feature set, we repeated this with larger feature
sets. Testing larger feature sets reduced our power to detect differences. No hits
were detected using a predefined set of 256 features (Javer, Ripoll-Sánchez, et al.,
2018) (Supplementary Figure 1).
We then performed a confirmation screen on 30 of the hit compounds from
the initial screen with a larger number of replicates (24 wells per day over three
independent days). 13 of the 30 hits were reproduced in the sense of showing
significant differences from *unc-80* DMSO controls in the direction of wild-type,
however several showed a reduced effect size and were clustered around the *unc-*
80 DMSO controls (Figure 5B). Having a larger number of replicates, we also tested
for 'side effects' in treated worms which we define as features where there was no
significant difference between *unc-80* mutants and wild-type N2 animals but where
there is a significant difference between *unc-80* mutants treated with a drug
compared to wild-type N2 animals. About half of the confirmed hit compounds lead
to more than 1000 side effects (Figure 5C) and are coloured yellow in the core *unc-*
80 phenotype space (Figure 5B). The same trend is observed when using 256
predefined features (Javer, Ripoll-Sánchez, et al., 2018) (Supplementary Figure 2).
Of the confirmed hit compounds with fewer than 1000 side effects, liranaftate
and atorvastatin (Lipitor) lead to good rescue of the core *unc-80* behavioural
phenotypes. Interestingly, both compounds act upon the melavonate biosynthesis
pathway. The main trunk of this pathway is conserved across species and converts
acetyl-CoA to farnesyl diphosphate (Rauthan & Pilon, 2011). Atorvastatin is an
inhibitor of HMG-CoA reductase that regulates lipid homeostasis (Goldstein et al.,
2006) and catalyses the rate limiting step of the melavonate pathway. Liranaftate, an
antifungal inhibitor of squalene epoxidase, shows the greatest behavioural
improvement, with the fewest number of side effects. However, *C. elegans* lack the
branch of the mevalonate pathway responsible for converting squalene to
cholesterol, instead relying on dietary cholesterol. In this case the effect may be due
to an off-target interaction on one of the other enzymes in the pathway.
Fenofibrate, a PPAR-alpha activator that increases lipolysis, is also a hit but
has a large number of side effects. Given that it overshoots perfect rescue by shifting
unc-80 animals past wild-type controls in core phenospace, it could be that at a
lower dose it would provide good rescue with fewer side effects. Since UNC-80 is
involved in localisation of the NACLN+ channel complex, we hypothesise that
changes in membrane composition might be indirectly affecting the localisation or
function of the channel complex.

Figure 5: Drug repurposing screening
(A) Phenotypes of unc-80(syb1531) mutant (red star) and N2 (blue star) worms treated with 1% DMSO, and unc-80(syb1531) mutants treated with a library of 743 FDA approved drugs at a concentration of 100 μM for 4 hours (circles). Each point represents an average of 3 well replicates, across 3 independent days of tracking (n = 9 total). Blue points are the top 30 compounds that significantly improved all three of the core behavioural features, pushing the unc-80 mutant strain towards the control in phenospace. (B) Confirmation screen of the top 30 compounds identified in the initial library screen. Again unc-80(syb1531) and N2 DMSO treated controls are represented by red and blue stars, respectively, and each circular point represents unc-80(syb1531) treated with 100 μM compound for 4 hours. The 13 compounds coloured in yellow lead to the worsening of >1000 behavioural features (see below). Liranaftate and atorvastatin both lead to a rescue of the core mutant phenotype with a low number of side effects. (C) Total number of behavioural side effects following treatment of unc-80(syb1531) with the 30 compounds in the confirmation screen. Side effects are defined as features that are not significant between unc-80 mutants and wild-type N2 worms treated with 1% DMSO but where there is a significant difference between unc-80 mutants treated with a drug compared to N2. Red dashed line separates drug treatments that lead to a worsening of >1000 behavioural features that correspond to the points coloured in yellow in the 3D scatterplot.

Discussion

We framed the problem of disease modelling as one of sampling and elucidating a model organism genotype-phenotype map that is connected to a human disease genotype-phenotype map through genetic conservation. However, both the genomic and phenomic spaces are large and high-dimensional and so it was not clear that a systematic approach would identify useful phenologs for disease modelling. Using an

initial set of genes that are diverse in function (but biased towards roles in neurons
and muscles) and that lead to diverse symptoms when mutated in humans, we found
that it is possible to systematically create and phenotype worm models of rare
diseases using a uniform assay and protocol. Without adding additional
perturbations or experimental conditions, we found that 23 out of the 25 disease
models we made had detectable differences compared to the wild-type.
Of the disease models with detectable phenotypes, approximately half had
strong phenotypes that would support high throughput phenotypic screens for
candidate treatments. To test the feasibility of performing drug repurposing screens
using the same uniform assay, we chose a worm model of UNC80 deficiency and,
focussing on a reduced core phenotype, identified an initial list of 30 FDA-approved
drugs that rescued the mutant phenotype. A confirmation screen with more
replicates identified several compounds that still rescued the core disease model
phenotype and did not have a large number of side effects. For UNC80 deficiency
we chose a short treatment time of four hours to model treatment in patients where
some developmental effects will have already occurred. Our approach is also
compatible with longer treatments where animals at the first larval stage are added
to tracking plates and allowed to develop in the presence of drugs. This might reveal
other hit compounds and would be particularly interesting for mutants with
developmental delays including some in this study (e.g. *cat-4*, *dys-1*, *figo-1*, *gpb-2*
and *kcc-2*).
For this proof-of-principle study, we focused on the generation of large
deletion alleles to help ensure the target gene is functionally disrupted. However,
patient-specific missense mutations can also be introduced at conserved loci within
genes and these mutations can lead to detectable behavioural phenotypes (see for
example (Barlow et al., 2022; D. Wang et al., 2021)). There is also value in studying
multiple alleles even for loss-of-function mutations. The large deletions we use here
can delete regulatory elements or non-coding RNAs that could lead to phenotypes
not related to the modelled disease. Alternative methods of creating loss-of-function
alleles such as introducing premature stop codons can also be done at the scale of
dozens of strains (Pu et al., 2023). Furthermore, to study loss of function in essential
genes, balanced heterozygous strains can be created where this makes sense for a
particular disease model.
With a current discovery rate of >100 new genetic diseases each year, the
number of diseases with little aetiological understanding is going to grow at a faster
rate than biomedical and pharmaceutical research will find therapies (Boycott et al.,
2017). Therefore, novel approaches that can systematically screen the effects of
large numbers of genetic variants associated with different diseases are required.
CRISPR genome editing in model organisms including *C. elegans* allows the
creation of new disease models at the required rate and cost. We have shown that
high-throughput worm tracking can identify screenable phenotypes and be used for
repurposing screens. However, we do not yet know whether the phenotypes and
repurposing hits will be useful for translation. It is possible that the phenotypes we
detect are not direct results of key disease mechanisms but are instead secondary
effects on connected systems. We expect that secondary effects are less likely to be
driven by conserved mechanisms but this can only be assessed on a case-by-base
basis. Similarly, the 'side effects' we detect in worms may not be predictive in
humans. Still, we think quantifying side effects will be useful in some cases. For
example, if a disease model is hyperactive and gross motility were the only measure
used in a screen, compounds that are broadly toxic would be detected as hits. This
kind of broad toxicity would result in a large number of side effects and could be

avoided. Ultimately, the translatability of hits in invertebrate screens can only be
determined in clinical trials and choosing hits from phenotypic screens to pursue
clinically will require judgement that takes into account the full picture available from
multiple data sources including the strength of rescue, the putative drug mechanism
of action, knowledge of disease biology, and an assessment of the risk of testing a
given drug in patients. We are optimistic that we will see more human trials based on
screens in diverse small animal models.
435
436

Materials and Methods
Mutant generation
CRISPR guide RNAs were designed to target large deletions (>1000bp) that start
close to the start codon and excise several exons from the gene in order to give high
confidence of loss of function. For exceptionally large genes (e.g. *dys-1* or *mpz-1*)
the entire protein-coding region was excised. Mutants were designed and made by
SunyBiotech in an N2 background.

Worm preparation
All strains were cultured on Nematode Growth Medium at 20°C and fed with *E. coli*
(OP50) following standard procedure (Stiernagle, 2006). Synchronised populations
of young adult worms for imaging were cultured by bleaching unsynchronised gravid
adults, and allowing L1 diapause progeny to develop for 2.5 days at 20°C (detailed
protocol: https://dx.doi.org/10.17504/protocols.io.2bzgap6). Several strains were
developmentally delayed and were allowed to grow for longer before imaging. *cat-4*(*syb1591*), *gpb-2*(*syb1577*), *kcc-2*(*syb2673*), and *unc-25*(*syb1651*) were allowed to
develop for 3.5 and *dys-1*(*syb1688*), *figo-1*(*syb1562*), and *pink-1*(*syb1546*) were
allowed to develop for 5.5 days prior to imaging. On the day of imaging, young adults
were washed in M9 (detailed protocol:
https://dx.doi.org/10.17504/protocols.io.bfqbjmsn), transferred to the imaging plates
(3 worms per well) using a COPAS 500 Flow Pilot (detailed protocol:
https://dx.doi.org/10.17504/protocols.io.bfc9jiz6), and returned to a 20°C incubator
for 3.5 hours. Plates were then transferred onto the multi-camera tracker for another
30 minutes to habituate prior to imaging (detailed protocol:
https://dx.doi.org/10.17504/protocols.io.bsicncaw).
        For drug repurposing experiments, the MRCT FDA-approved compound
library (Catalog No.L1300) was supplied pre-dissolved in DMSO by LifeArc
(Stevenage, UK). The day prior to tracking, imaging plates were dosed with the
compound library to achieve a final well concentration of 100 µM prior to seeding
with bacteria (see below for details). Plates were left to dry (~30 mins), before being
stored in the dark at room temperature overnight. Following the methods described
above, age synchronized young adult *unc-80*(*syb1531*) worms were dispensed into
the imaging plate wells and incubated at 20°C for 4 hours before tracking. The
behaviour of mutant worms dosed with the drugs was then compared to wild-type N2
and *unc-80*(*syb1531*) worms (also age-synchronized young adults) dispensed into
the wells of the same tracking plates dosed with an identical volume (1% w/v) of
DMSO only (detailed protocol:
https://dx.doi.org/10.17504/protocols.io.5jyl8p5yrg2w/v1).

Plate preparation

Low peptone (0.013%) nematode growth medium (detailed protocol:
https://dx.doi.org/10.17504/protocols.io.2rcgd2w) was prepared as follows: 20 g agar
(Difco), 0.13 g Bactopeptone, and 3 g NaCl were dissolved in 975 mL of milliQ water.
After autoclaving, 1 mL of 10 mg/mL cholesterol was added along with 1 mL CaCl2
(1 M), 1 mL MgSO4 (1 M) and 25 mL KPO4 buffer (1M, pH 6.0). Molten agar was
cooled to 50-60°C and 200 µL was dispensed into each well of 96-square well plates
(Whatman UNIPLATE: WHAT-77011651) using an Integra VIAFILL (detailed
protocol: https://dx.doi.org/10.17504/protocols.io.bmxbk7in). Poured plates were
stored agar-side up at 4°C until required.
One day prior to imaging, plates were placed without lids in a drying cabinet to
lose 3-5% of weight by volume. Wells were then seeded with 5 µL OP50 (OD600
1.0) using an Integra VIAFILL dispenser, and plates were stored with lids (WHAT-
77041001) on at room temperature overnight.
490
Image acquisition
All videos were acquired and processed following methods previously described
(Barlow et al., 2022). In brief, videos were acquired at 25 frames per second in a
room with a nominal temperature of 20°C using a shutter time of 25 ms and a
resolution of 12.4 µm/px. Three videos were taken sequentially: a 5-minute pre-
stimulus video, a 6-minute blue light recording with three 10-second blue light pulses
starting at 60, 160 and 260 seconds, and a 5-minute post-stimulus recording. The
script for controlling recording timings and photostimulation was made using
LoopBio's API for their Motif software (https://github.com/loopbio/python-motifapi).
500
Image processing and feature extraction
Videos were segmented and tracked using Tierpsy Tracker (Javer, Currie, et al.,
2018). After segmentation of worm skeletons, our previously described convolutional
neural network classifier was used to exclude non-worm objects from being
classified during feature extraction (Barlow et al., 2022). Skeletons that did not meet
the following criteria were removed from the analysis: 700 – 1300 µM length, 20 –
200 µM width. As an additional quality control measure, we used Tierpsy Tracker's
viewer to mark wells with visible contamination, agar damage, compound
precipitation, or excess liquid as "bad", and exclude these wells from downstream
analysis.
Following tracking, we extracted a previously-defined set of 3076 behavioural
features for each well in each of the three videos (pre-stimulus, blue light, and post-
stimulus) (Javer, Ripoll-Sánchez, et al., 2018). Feature values are averaged over
tracks to produce a single feature vector for each well.
515
Statistical analysis
Statistically significant differences in the pre-stimulus, post-stimulus and blue-light
behavioural feature sets extracted from each disease model strain compared to our
N2 reference were calculated using block permutation t-tests
(https://github.com/Tierpsy/tierpsy-tools-
python/blob/master/tierpsytools/analysis/statistical_tests.py). Python (version 3.8.5)
was used to perform the analysis, using n = 10000 permutations that were randomly
shuffled within, but not between, the independent days of image acquisition for each
strain to control for day-to-day variation in the experiments. The *p*-values were then
corrected for multiple comparisons using the Benjamini-Yekutieli procedure to control
the false discovery rate at 5% (Benjamini & Yekutieli, 2001).

Heatmaps, cluster maps and principal component analysis of the extracted feature sets for each disease model strain compared to our N2 reference were calculated using in-built Seaborn (version 0.11.2) packages (Waskom, 2021). All scripts used for statistical analysis and the generation of figures are available at: https://github.com/Tom-OBrien/Disease-Modelling.

Data Availability

The datasets (Tierpsy features, tracking data, and metadata) produced in this study are available on Zenodo: https://doi.org/10.5281/zenodo.12684118. Any remaining information can be obtained from the corresponding author upon reasonable request.

Acknowledgements

This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (Grant agreement No. 714853) and was supported by the Medical Research Council through grant MC-A658-5TY30.

References

Aoyagi, K., Rossignol, E., Hamdan, F. F., Mulcahy, B., Xie, L., Nagamatsu, S., Rouleau, G. A.,

Zhen, M., & Michaud, J. L. (2015). A Gain-of-Function Mutation in NALCN in a Child

with Intellectual Disability, Ataxia, and Arthrogryposis. *Human Mutation*, *36*(8), 753–

757. https://doi.org/10.1002/humu.22797

Baek, J.-H., Cosman, P., Feng, Z., Silver, J., & Schafer, W. R. (2002). Using machine vision to

analyze and classify Caenorhabditis elegans behavioral phenotypes quantitatively.

Journal of Neuroscience Methods, *118*, 9–21. https://doi.org/10.1016/S0165-

0270(02)00117-6

Barlow, I. L., Feriani, L., Minga, E., McDermott-Rouse, A., O'Brien, T. J., Liu, Z., Hofbauer, M.,

Stowers, J. R., Andersen, E. C., Ding, S. S., & Brown, A. E. X. (2022). Megapixel camera

arrays enable high-resolution animal tracking in multiwell plates. *Communications*

Biology, *5*(1), Article 1. https://doi.org/10.1038/s42003-022-03206-1

Bavisetty, S., Grody, W. W., & Yazdani, S. (2013). Emergence of pediatric rare diseases:

Review of present policies and opportunities for improvement. *Rare Diseases*, *1*.

https://doi.org/10.4161/rdis.23579

Benjamini, Y., & Yekutieli, D. (2001). The control of the false discovery rate in multiple

testing under dependency. *The Annals of Statistics*, *29*(4), 1165–1188.

https://doi.org/10.1214/aos/1013699998

Boycott, K. M., Rath, A., Chong, J. X., Hartley, T., Alkuraya, F. S., Baynam, G., Brookes, A. J.,

Brudno, M., Carracedo, A., den Dunnen, J. T., Dyke, S. O. M., Estivill, X., Goldblatt, J.,

Gonthier, C., Groft, S. C., Gut, I., Hamosh, A., Hieter, P., Höhn, S., … Lochmüller, H.

(2017). International Cooperation to Enable the Diagnosis of All Rare Genetic

Diseases. *The American Journal of Human Genetics*, *100*(5), 695–705.

https://doi.org/10.1016/j.ajhg.2017.04.003

Chih, B., Liu, P., Chinn, Y., Chalouni, C., Komuves, L. G., Hass, P. E., Sandoval, W., & Peterson,

A. S. (2012). A ciliopathy complex at the transition zone protects the cilia as a

privileged membrane domain. *Nature Cell Biology*, *14*(1), Article 1.

https://doi.org/10.1038/ncb2410

Cochet-Bissuel, M., Lory, P., & Monteil, A. (2014). The sodium leak channel, NALCN, in

health and disease. *Frontiers in Cellular Neuroscience*, *8*.

https://www.frontiersin.org/article/10.3389/fncel.2014.00132

Forsyth, R., & Gunay-Aygun, M. (2023). Bardet-Biedl Syndrome Overview. In M. P. Adam, G.

M. Mirzaa, R. A. Pagon, S. E. Wallace, L. J. Bean, K. W. Gripp, & A. Amemiya (Eds.),

GeneReviews®. University of Washington, Seattle.

http://www.ncbi.nlm.nih.gov/books/NBK1363/

Forsythe, E., & Beales, P. L. (2013). Bardet–Biedl syndrome. *European Journal of Human*

Genetics, *21*(1), Article 1. https://doi.org/10.1038/ejhg.2012.115

Gal, M., Magen, D., Zahran, Y., Ravid, S., Eran, A., Khayat, M., Gafni, C., Levanon, E. Y., &

Mandel, H. (2016). A novel homozygous splice site mutation in NALCN identified in

siblings with cachexia, strabismus, severe intellectual disability, epilepsy and

abnormal respiratory rhythm. *European Journal of Medical Genetics*, *59*(4), 204–209.

https://doi.org/10.1016/j.ejmg.2016.02.007

Geng, W., Cosman, P., Berry, C. C., Feng, Z., & Schafer, W. R. (2004). Automatic tracking,

feature extraction and classification of C elegans phenotypes. *IEEE Transactions on*

Bio-Medical Engineering, *51*(10), 1811–1820.

https://doi.org/10.1109/TBME.2004.831532

Goldstein, J. L., DeBose-Boyd, R. A., & Brown, M. S. (2006). Protein sensors for membrane

sterols. *Cell*, *124*(1), 35–46. https://doi.org/10.1016/j.cell.2005.12.022

Gonzaga-Jauregui, C., Lupski, J. R., & Gibbs, R. A. (2012). Human genome sequencing in

health and disease. *Annual Review of Medicine*, *63*, 35–61.

https://doi.org/10.1146/annurev-med-051010-162644

Harris, T. W., Arnaboldi, V., Cain, S., Chan, J., Chen, W. J., Cho, J., Davis, P., Gao, S., Grove, C.

A., Kishore, R., Lee, R. Y. N., Muller, H.-M., Nakamura, C., Nuin, P., Paulini, M., Raciti,

D., Rodgers, F. H., Russell, M., Schindelman, G., … Sternberg, P. W. (2020).

WormBase: A modern Model Organism Information Resource. *Nucleic Acids*

Research, *48*(D1), D762–D767. https://doi.org/10.1093/nar/gkz920

Iyer, S., Sam, F. S., DiPrimio, N., Preston, G., Verheijen, J., Murthy, K., Parton, Z., Tsang, H.,

Lao, J., Morava, E., & Perlstein, E. O. (2019). Repurposing the aldose reductase

inhibitor and diabetic neuropathy drug epalrestat for the congenital disorder of

glycosylation PMM2-CDG. *Disease Models & Mechanisms*, *12*(11), dmm040584.

https://doi.org/10.1242/dmm.040584

Javer, A., Currie, M., Lee, C. W., Hokanson, J., Li, K., Martineau, C. N., Yemini, E., Grundy, L.

J., Li, C., Ch'ng, Q., Schafer, W. R., Nollen, E. A., Kerr, R., & Brown, A. E. (2018). An

open source platform for analyzing and sharing worm behavior data. *Nature*

Methods, *15*(9), 645–646. https://doi.org/10.1038/s41592-018-0112-1

Javer, A., Ripoll-Sánchez, L., & Brown, A. E. X. (2018). Powerful and interpretable

behavioural features for quantitative phenotyping of Caenorhabditis elegans.

Philosophical Transactions of the Royal Society B: Biological Sciences, *373*(1758),

20170375. https://doi.org/10.1098/rstb.2017.0375

Jospin, M., Watanabe, S., Joshi, D., Young, S., Hamming, K., Thacker, C., Snutch, T. P.,

Jorgensen, E. M., & Schuske, K. (2007). UNC-80 and the NCA ion channels contribute

to endocytosis defects in synaptojanin mutants. *Current Biology: CB*, *17*(18), 1595–

1600. https://doi.org/10.1016/j.cub.2007.08.036

Karmous-Benailly, H., Martinovic, J., Gubler, M.-C., Sirot, Y., Clech, L., Ozilou, C., Augé, J.,

Brahimi, N., Etchevers, H., Detrait, E., Esculpavit, C., Audollent, S., Goudefroye, G.,

Gonzales, M., Tantau, J., Loget, P., Joubert, M., Gaillard, D., Jeanne-Pasquier, C., …

Attié-Bitach, T. (2005). Antenatal Presentation of Bardet-Biedl Syndrome May Mimic

Meckel Syndrome. *American Journal of Human Genetics*, *76*(3), 493–504.

Kim, W., Underwood, R. S., Greenwald, I., & Shaye, D. D. (2018). OrthoList 2: A New

Comparative Genomic Analysis of Human and Caenorhabditis elegans Genes.

Genetics, *210*(2), 445–461. https://doi.org/10.1534/genetics.118.301307

Kropp, P. A., Bauer, R., Zafra, I., Graham, C., & Golden, A. (2021). Caenorhabditis elegans for

rare disease modeling and drug discovery: Strategies and strengths. *Disease Models*

& Mechanisms, *14*(8), dmm049010. https://doi.org/10.1242/dmm.049010

Lee, C. E., Singleton, K. S., Wallin, M., & Faundez, V. (2020). Rare Genetic Diseases: Nature's

Experiments on Human Development. *iScience*, *23*(5), 101123.

https://doi.org/10.1016/j.isci.2020.101123

Lu, B., Su, Y., Das, S., Liu, J., Xia, J., & Ren, D. (2007). The neuronal channel NALCN

contributes resting sodium permeability and is required for normal respiratory

rhythm. *Cell*, *129*(2), 371–383. https://doi.org/10.1016/j.cell.2007.02.041

Mak, H. Y., Nelson, L. S., Basson, M., Johnson, C. D., & Ruvkun, G. (2006). Polygenic control

of Caenorhabditis elegans fat storage. *Nature Genetics*, *38*(3), Article 3.

https://doi.org/10.1038/ng1739

McDermott-Rouse, A., Minga, E., Barlow, I., Feriani, L., Harlow, P. H., Flemming, A. J., &

Brown, A. E. X. (2021). Behavioral fingerprints predict insecticide and anthelmintic

mode of action. *Molecular Systems Biology*, *17*(5), e10267.

https://doi.org/10.15252/msb.202110267

McGary, K. L., Park, T. J., Woods, J. O., Cha, H. J., Wallingford, J. B., & Marcotte, E. M. (2010).

Systematic discovery of nonobvious human disease models through orthologous

phenotypes. *Proceedings of the National Academy of Sciences*, *107*(14), 6544–6549.

https://doi.org/10.1073/pnas.0910200107

McGinnis, S., & Madden, T. L. (2004). BLAST: At the core of a powerful and diverse set of

sequence analysis tools. *Nucleic Acids Research*, *32*(Web Server issue), W20–W25.

https://doi.org/10.1093/nar/gkh435

Mukhopadhyay, S., & Jackson, P. K. (2011). The tubby family proteins. *Genome Biology*,

12(6), 225. https://doi.org/10.1186/gb-2011-12-6-225

Nguengang Wakap, S., Lambert, D. M., Olry, A., Rodwell, C., Gueydan, C., Lanneau, V.,

Murphy, D., Le Cam, Y., & Rath, A. (2020). Estimating cumulative point prevalence of

rare diseases: Analysis of the Orphanet database. *European Journal of Human*

Genetics, *28*(2), Article 2. https://doi.org/10.1038/s41431-019-0508-0

North, M. A., Naggert, J. K., Yan, Y., Noben-Trauth, K., & Nishina, P. M. (1997). Molecular

characterization of TUB, TULP1, and TULP2, members of the novel tubby gene family

and their possible relation to ocular diseases. *Proceedings of the National Academy*

of Sciences of the United States of America, *94*(7), 3128–3133.

Patten, S. A., Aggad, D., Martinez, J., Tremblay, E., Petrillo, J., Armstrong, G. A. B., Fontaine,

A. L., Maios, C., Liao, M., Ciura, S., Wen, X.-Y., Rafuse, V., Ichida, J., Zinman, L., Julien,

J.-P., Kabashi, E., Robitaille, R., Korngut, L., Parker, J. A., & Drapeau, P. (2017).

Neuroleptics as therapeutic compounds stabilizing neuromuscular transmission in

amyotrophic lateral sclerosis. *JCI Insight*, *2*(22).

https://doi.org/10.1172/jci.insight.97152

Perni, M., Challa, P. K., Kirkegaard, J. B., Limbocker, R., Koopman, M., Hardenberg, M. C.,

Sormanni, P., Müller, T., Saar, K. L., Roode, L. W. Y., Habchi, J., Vecchi, G., Fernando,

N. W., Casford, S., Nollen, E. A. A., Vendruscolo, M., Dobson, C. M., & Knowles, T. P.

J. (2018). Massively parallel C. elegans tracking provides multi-dimensional

fingerprints for phenotypic discovery. *Journal of Neuroscience Methods*.

https://doi.org/10.1016/j.jneumeth.2018.02.005

Pierce-Shimomura, J. T., Chen, B. L., Mun, J. J., Ho, R., Sarkis, R., & McIntire, S. L. (2008).

Genetic analysis of crawling and swimming locomotory patterns in C. elegans.

Proceedings of the National Academy of Sciences of the United States of America,

105(52), 20982–20987. https://doi.org/10.1073/pnas.0810359105

Pu, L., Wang, J., Lu, Q., Nilsson, L., Philbrook, A., Pandey, A., Zhao, L., Schendel, R. van, Koh,

A., Peres, T. V., Hashi, W. H., Myint, S. L., Williams, C., Gilthorpe, J. D., Wai, S. N.,

Brown, A., Tijsterman, M., Sengupta, P., Henriksson, J., & Chen, C. (2023). Dissecting

the genetic landscape of GPCR signaling through phenotypic profiling in C. elegans.

Nature Communications, *14*(1), 8410. https://doi.org/10.1038/s41467-023-44177-z

Ramot, D., Johnson, B. E., Berry, T. L., Carnell, L., & Goodman, M. B. (2008). The Parallel

Worm Tracker: A Platform for Measuring Average Speed and Drug-Induced Paralysis

in Nematodes. *PLoS ONE*, *3*, e2208. https://doi.org/10.1371/journal.pone.0002208

Rauthan, M., & Pilon, M. (2011). The mevalonate pathway in C. Elegans. *Lipids in Health and*

Disease, *10*, 243. https://doi.org/10.1186/1476-511X-10-243

Reiter, J. F., & Leroux, M. R. (2017). Genes and molecular pathways underpinning

ciliopathies. *Nature Reviews Molecular Cell Biology*, *18*(9), Article 9.

https://doi.org/10.1038/nrm.2017.60

Restif, C., Ibáñez-Ventoso, C., Vora, M. M., Guo, S., Metaxas, D., & Driscoll, M. (2014).

CeleST: Computer vision software for quantitative analysis of C. elegans swim

behavior reveals novel features of locomotion. *PLoS Computational Biology*, *10*(7),

e1003702. https://doi.org/10.1371/journal.pcbi.1003702

Roberson, E. C., Dowdle, W. E., Ozanturk, A., Garcia-Gonzalo, F. R., Li, C., Halbritter, J.,

Elkhartoufi, N., Porath, J. D., Cope, H., Ashley-Koch, A., Gregory, S., Thomas, S.,

Sayer, J. A., Saunier, S., Otto, E. A., Katsanis, N., Davis, E. E., Attié-Bitach, T.,

Hildebrandt, F., … Reiter, J. F. (2015). TMEM231, mutated in orofaciodigital and

Meckel syndromes, organizes the ciliary transition zone. *The Journal of Cell Biology*,

209(1), 129–142. https://doi.org/10.1083/jcb.201411087

Roessler, H. I., Knoers, N. V. A. M., van Haelst, M. M., & van Haaften, G. (2021). Drug

Repurposing for Rare Diseases. *Trends in Pharmacological Sciences*, *42*(4), 255–267.

https://doi.org/10.1016/j.tips.2021.01.003

Schriml, L. M., Munro, J. B., Schor, M., Olley, D., McCracken, C., Felix, V., Baron, J. A.,

Jackson, R., Bello, S. M., Bearer, C., Lichenstein, R., Bisordi, K., Dialo, N. C., Giglio, M.,

& Greene, C. (2022). The Human Disease Ontology 2022 update. *Nucleic Acids*

Research, *50*(D1), D1255–D1261. https://doi.org/10.1093/nar/gkab1063

Senatore, A., Monteil, A., Minnen, J. van, Smit, A. B., & Spafford, J. D. (2013). NALCN Ion

Channels Have Alternative Selectivity Filters Resembling Calcium Channels or Sodium

Channels. *PLOS ONE*, *8*(1), e55088. https://doi.org/10.1371/journal.pone.0055088

Shaheen, R., Szymanska, K., Basu, B., Patel, N., Ewida, N., Faqeih, E., Al Hashem, A., Derar,

N., Alsharif, H., Aldahmesh, M. A., Alazami, A. M., Hashem, M., Ibrahim, N.,

Abdulwahab, F. M., Sonbul, R., Alkuraya, H., Alnemer, M., Al Tala, S., Al-Husain, M., …

Alkuraya, F. S. (2016). Characterizing the morbid genome of ciliopathies. *Genome*

Biology, *17*, 242. https://doi.org/10.1186/s13059-016-1099-5

Stiernagle, T. (2006). Maintenance of C. elegans. *WormBook: The Online Review of C.*

Elegans Biology, 1–11. https://doi.org/10.1895/wormbook.1.101.1

Sun, X., Haley, J., Bulgakov, O. V., Cai, X., McGinnis, J., & Li, T. (2012). Tubby is required for

trafficking G protein-coupled receptors to neuronal cilia. *Cilia*, *1*(1), 21.

https://doi.org/10.1186/2046-2530-1-21

Swierczek, N. A., Giles, A. C., Rankin, C. H., & Kerr, R. A. (2011). High-throughput behavioral

analysis in C. elegans. *Nature Methods*, *8*(7), 592–598.

https://doi.org/10.1038/nmeth.1625

Taylor, S. R., Santpere, G., Weinreb, A., Barrett, A., Reilly, M. B., Xu, C., Varol, E.,

Oikonomou, P., Glenwinkel, L., McWhirter, R., Poff, A., Basavaraju, M., Rafi, I.,

Yemini, E., Cook, S. J., Abrams, A., Vidal, B., Cros, C., Tavazoie, S., … Miller, D. M.

(2021). Molecular topography of an entire nervous system. *Cell*, *184*(16), 4329-

4347.e23. https://doi.org/10.1016/j.cell.2021.06.023

Topalidou, I., Cooper, K., Pereira, L., & Ailion, M. (2017). Dopamine negatively modulates

the NCA ion channels in C. elegans. *PLoS Genetics*, *13*(10), e1007032.

https://doi.org/10.1371/journal.pgen.1007032

Tsibidis, G. D., & Tavernarakis, N. (2007). Nemo: A computational tool for analyzing

nematode locomotion. *BMC Neuroscience*, *8*, 86. https://doi.org/10.1186/1471-

2202-8-86

Wang, D., Dao, M., Muntean, B. S., Giles, A. C., Martemyanov, K. A., & Grill, B. (2021).

Genetic modeling of GNAO1 disorder delineates mechanisms of Gαo dysfunction.

Human Molecular Genetics, *31*(4), 510–522. https://doi.org/10.1093/hmg/ddab235

Wang, S. J., & Wang, Z.-W. (2013). Track-A-Worm, An Open-Source System for Quantitative

Assessment of C. elegans Locomotory and Bending Behavior. *PLoS ONE*, *8*(7),

e69653. https://doi.org/10.1371/journal.pone.0069653

Ward, A., Liu, J., Feng, Z., & Xu, X. Z. S. (2008). Light-sensitive neurons and channels mediate

phototaxis in C. elegans. *Nature Neuroscience*, *11*(8), Article 8.

https://doi.org/10.1038/nn.2155

Waskom, M. L. (2021). seaborn: Statistical data visualization. *Journal of Open Source*

Software, *6*(60), 3021. https://doi.org/10.21105/joss.03021

Yeh, E., Ng, S., Zhang, M., Bouhours, M., Wang, Y., Wang, M., Hung, W., Aoyagi, K., Melnik-

Martinez, K., Li, M., Liu, F., Schafer, W. R., & Zhen, M. (2008). A putative cation

channel, NCA-1, and a novel protein, UNC-80, transmit neuronal activity in C.

elegans. *PLoS Biology*, *6*(3), e55. https://doi.org/10.1371/journal.pbio.0060055

Yemini, E., Jucikas, T., Grundy, L. J., Brown, A. E. X., & Schafer, W. R. (2013). A database of

Caenorhabditis elegans behavioral phenotypes. *Nature Methods*, *10*(9), 877–879.

https://doi.org/10.1038/nmeth.2560

Zhang, X., Liu, J., Pan, T., Ward, A., Liu, J., & Xu, X. Z. S. (2022). A cilia-independent function

of BBSome mediated by DLK-MAPK signaling in C. elegans photosensation.

Developmental Cell, *57*(12), 1545-1557.e4.

https://doi.org/10.1016/j.devcel.2022.05.005

Zhou, C., Luo, J., He, X., Zhou, Q., He, Y., Wang, X., & Ma, L. (2019). The NALCN Channel

Regulator UNC-80 Functions in a Subset of Interneurons To Regulate Caenorhabditis

elegans Reversal Behavior. *G3: Genes|Genomes|Genetics*, *10*(1), 199–210.

https://doi.org/10.1534/g3.119.400692

760