Estimating the number of breeders from helminth larvae with genomic data.

Tristan. P. W. Dennis[1,2]✉, William Sands[2], Millicent Opoku[2,8,9], Alex Debrah3,4, Linda Batsa,

Kenneth Pfarr[6,7], Ute Klarmann-Schulz[6,7], Achim Hoerauf [6,7], Sabine Specht[10], Ivan

Scandale[10], Lisa C. Ranford-Cartwright[2]▯, Poppy H. L. Lamberton[2]▯✉

▯Joint sr author
▯ Corresponding authors
1. Department of Vector Biology, Liverpool School of Tropical Medicine, Liverpool, UK
2. School of Biodiversity, One Health and Veterinary Medicine, University of Glasgow, Glasgow, UK
3. Kumasi Centre for Collaborative Research in Tropical Medicine, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana
4. Department of Medical Diagnostics, Faculty of Allied Health Sciences, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana
5. Department of Clinical Microbiology, School of Medicine and Dentistry, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana.
6. Institute for Medical Microbiology, Immunology and Parasitology, University Hospital Bonn, Bonn, Germany
7. German Center for Infection Research (DZIF), Partner Site Bonn-Cologne, Bonn, Germany
8. Department of Parasitology, Noguchi Memorial Institute for Medical Research, University of Ghana, Accra, Ghana
9. Department of Environment and Genetics, La Trobe University, Bundoora, Melbourne, Australia
10. Drugs For Neglected Diseases Initiative, Geneva, Switzerland

Abstract

Effective control of helminth infections requires the application of mathematical models to inform control efforts and policy, the development of product profiles for new drugs, and the monitoring of existing drugs for resistance. Key to the success of these approaches is accurately estimating the number of worms within a host, as well as distinguishing, in drug efficacy trials and monitoring, between adults surviving treatment and adults who have reinfected a host following drug treatment. In practice, observing adult worms is often extremely challenging in a live patient, as many adult helminths are embedded deep in host tissues. Genetic approaches to infer kinship between larvae or eggs offer a solution to establish adult worm burdens, and to distinguish between treatment failure or treatment success followed by reinfection. Here, we use low-coverage whole-genome, and mitochondrial sequencing, of *Onchocerca volvulus* larvae to estimate the number of adults contributing to pools of offspring of known and unknown parentage. lcWGS reconstructs full-sibling relationships, resolving the number of unique adult worms contributing to a pool of offspring. Mitochondrial genotyping reconstructs maternal sib-ship, thus estimating the minimum number of adult females within a patient. Further development will improve these techniques for evaluating adult worm burden and trial outcome.

Intro

Human helminthiases exert a devastating worldwide public-health burden. Many are neglected tropical diseases (NTDs), causing poor health, morbidity, and mortality, disproportionately for the world's poorest people (1). Mass drug administration (MDA) with anthelmintic drugs remains the cornerstone of treatment for many NTDs. Monitoring and evaluation of these programmes is vital to support continued drug donation, to detect potential drug resistance, and to identify areas for improved interventions. The complex life cycle of many indirectly transmitted macroparasites makes elucidation of adult worm burden, the evaluation of control and drug trial success, and establishment of baselines of infection levels in pre-clinical trials difficult, as adult worms of many species (e.g. *Schistosoma mansoni* (2), *Wuchereria bancrofti* (3), *Onchocerca volvulus* (4), *Opisthorchis viverrini* (5) or *Trichuris trichiura* (6)) are either very difficult or impossible to retrieve

from a patient. As such, inferring whether a trial macrofilaricide has successfully killed all adult
worms, with any subsequent infection being due to reinfection, or whether treatment failure has
been followed by repopulation of microfilariae (mf), or renewed reproduction and excretion of
larvae from surviving adults, is impossible using current diagnostics. Moreover, estimating adult
worm burden, a key parameter for assessing infection severity and individual contribution to
community transmission, based on sampled offspring is problematic, as helminth larval abundance
does not necessarily linearly predict adult worm burden, and instead often follows a density-
dependent pattern (5,7,8), with adults only sporadically contributing to the larval pool at any given
time (9). This gap introduces a considerable level of uncertainty for mathematical models – key
tools for informing policy and control success.
59
Onchocerca volvulus is a filarial nematode parasite of humans, and is the agent of
onchocerciasis, also known as river blindness, a disease characterised by blindness and
debilitating, disfiguring skin disease. Onchocerciasis afflicts over 20 million people, with ~1 million
affected by blindness (10). The severe disease burden exerted by *O. volvulus* in Africa, South
America and Yemen have led to concerted efforts to eliminate onchocerciasis. Since 1997,
Community Directed Treatment with the microfilaricidal drug Ivermectin (Mectizan®) (CDTI) has
been the mainstay of *O. volvulus* control and is the current World Health Organization (WHO)
recommended strategy, administered as yearly MDA. Despite decades of CDTI, transmission and
disease burden have remained stubbornly high in many regions (11–14). This difficulty
necessitates both improved monitoring of existing CDTI, and the development and/or repurposing
of anthelmintic drugs such as moxidectin (4), emodepside (15,16), oxfendazole (17) or
Corallopyronin A (18). Evaluation of anthelmintics for *O. volvulus* is challenging. The adult worm is
embedded in fibrous nodules, and whilst some of these nodules can be detected if near the skin's
surface, the adult worms themselves are accessible only through surgical nodulectomy - an
invasive procedure. Diagnostics therefore rely on the counting of larval worms - microfilaria (mf)
from skin biopsies, also an invasive procedure (albeit less so than a nodulectomy).
76
Genetic approaches for evaluating sib-ship and relatedness enable the estimation of the
number of adults contributing to a sample of larvae (number of breeders, (*Nb)*), as well as
establishing whether the adults contributing to serially sampled larvae have changed, e.g. if
treatment has cleared the first set of parents, and a new set of parents have infected the patient. If
full-siblings (FS) or maternal half-siblings (HS) between pre- and post-treatment samples are
identified, treatment failure can be asserted due to the presence of surviving fecund females.
Estimating the relationships and parentage of helminth larvae in the absence of adult information
has been performed with multilocus genotyping methods such as microsatellite genotyping, with
sib-ship and *Nb* reconstruction being performed most commonly with likelihood-based methods
such as COLONY (19) for *Schistosoma mansoni* (20,21) and MASTERBAYES (22) for
Trichostrongylus tenuis, methods which rely on well-validated microsatellite, SNP or multilocus
sequence typing (MLST) panels.
89
Estimation of pairwise relatedness with genome data makes it possible to infer close familial
relationships between individuals in the absence of validated multilocus data, and this is chiefly
performed by estimating the proportion of alleles between two individuals that share 0, 1 or 2

copies identical-by-descent (IBD) (23). Whole-genome-sequencing is becoming increasingly
affordable, especially with the advent of low-coverage WGS (lcWGS) approaches (24). lcWGS is
particularly valuable if achieving high sequencing depth is problematic due to low-input and
degraded samples - common for many helminth larvae (25). WGS is now a widely used tool for
parasite molecular epidemiology (26,27), is possible from single helminth larvae (25), and can be
used to determine relatedness (28,29). Another approach toward assigning parentage is
mitochondrial genome sequencing. Mitochondria are inherited maternally, meaning that two
individuals sharing identical, or near-identical mitochondrial genomes, can be assigned as
probable maternal siblings. This is particularly useful in non-monogamous systems, such as many
helminths including *O. volvulus (*which is thought to be polyandrous) (30), where a combination of
relatedness estimation and mitochondrial genotyping can be used to identify maternal half-sibs.
Due to higher copy number of the mitochondrial genome in the cell opposed to the nuclear
genome, it is often easier to selectively amplify mitochondrial genomes than whole nuclear
genomes for sequencing and analysis (24).
107
In this study, we perform a proof-of-concept with lcWGS, relatedness estimation, and
mitochondrial genotyping in a complex indirectly transmitted macroparasite, *O. volvulus.* Our aims
were to: 1) establish parentage and the *Nb* contributing to a sample of mf from which parentage is
known, dissected from the uterus of gravid adult females, and 2) identify parentage of a sample of
mf taken from a skin biopsy, of which parentage is unknown. Finally, we discuss the extent to
which the approaches taken here may be more widely used toward estimating larval relatedness,
adult worm burden, and treatment failure for any parasite where adults are inaccessible, but
offspring, (eggs or larvae) can be obtained.
116

Results

1.   Parentage of intrauterine mf
We sequenced the genomes of 10 mf dissected from three adult females, from two separate
human patients (**Figure 1**). The mean number of reads ('X') covering each position was 10.1 over
the nuclear genome, and 47.6 over the mitochondrial genome. The exact number of per-sample
reads is given in **Supplementary Table 1**. To calculate relatedness between each mf, we used
genotype-likelihoods inferred from aligned reads as input for *NGSRelate,* which implements the
framework for estimating relatedness and relationship described by (28). The inferred R0, R1, and
KING kinship statistics from the intrauterine mf showed that mf collected from the same female
worms/broods all formed a discrete cluster within the kinship coefficient thresholds for full- or half-
sibling (28) , (**Supplementary Figures 1A & 1B**). One pair from the same female fell outside of
this (**Supplementary Figure 1A, 1B**). Based on these plots, we defined thresholds for full-sibling
relationships at R0 ≤ 0.1, R1 ≥ 0.5, and KING ≥ 0.25, and thresholds for half-sibling relationships

as R0 ≥ 0.1, R1 ≤ 0.5, and KING ≤ 0.25. These thresholds were then applied to our intrauterine mf
data (**Supplementary Figure 2**).

To further evaluate whether this method reconstructed sib-ship according to our expectations,
we drew networks from the relatedness data and plotted them (**Figure 2**). We found that

individuals dissected from the same female were at least full or half siblings. However, we found
that the assignments also predicted half-sibling relationships to mf dissected from different
females. As it is difficult to resolve relationships accurately below FS without nonsensical results
(28,31), we discarded our HS estimations and retained only FS relationships (**Figure 2**). The
network showed four FS groups, implying three female worms, and four male worms, contributed
to the mf sample.
142
We extracted mitochondrial genome *data in silico* from the intrauterine mf WGS data and used
them to build a minimum-spanning network (**Figure 3**). The network showed a high-level of within-
brood sequence similarity, with members of the same brood (having been dissected out of the
same female uterus) being a maximum of three SNPs different to one another. This method
reconstructed mitochondrial group membership with all offspring corresponding to the mother that
they were dissected from (**Figure 3**). When the number of mitochondrial groups was considered
alongside the number of full-sibling groups inferred from the intrauterine mf sample, we estimated
that seven unique parents (three females and four males) had contributed to this sample of 10 mf.
151
152　*2.　Maternal　assignment　of　skin-snip　microfilariae　from　the　same　patient*
153
To infer the number of breeders contributing to a sample of mf *for which the parentage was*
unknown, we attempted to sequence the genomes of 39 mf from two skin snips taken from the left
and right trochanter of a single patient. (**Figure 4**). Due to uneven and low nuclear genome
coverage (**Supplementary Table 2**) (mean 4.1), we were unable to obtain reliable relatedness
data for these samples. However, mitochondrial genome coverage across all samples was
sufficient to genotype whole mitochondria genomes (mean depth-of-coverage 216.1X)
(**Supplementary Table 2**). The medium-spanning network disclosed five major mitochondrial
groups (**Figure 5**) indicating at least five adult females contributing to the 39 microfilariae for which
we successfully obtained sufficient mitochondrial genome data from this patient. There was no
structuring in mf haplotype by location (left or right-hand trochanter) **(Figure 5).**
164

Discussion
For many helminth parasites, adult worms are not readily available for diagnosis, either at all
from living patients (e.g. *Schistosoma spp.),* or only through invasive surgery (e.g. *O. volvulus),* or
inefficient adult worm excretion methods (e.g. *Ascaris*, *Opisthorchis*). Despite many transmission
models and drug efficacy measurements being based on estimated adult numbers (32,33),
diagnostics often only focus on the offspring. Here, we use intrauterine- collected mf from known
O. volvulus adult females, as well as mf of unknown parentage collected from two skin-snips from
one patient, to test methods to predict sib-ship relationships, maternal parentage and number of
breeders (*Nb)*. We report the outcome of estimating helminth parentage using low-coverage WGS
and relatedness estimation, as well as mitochondrial genotyping. For the intrauterine mf, we were
successfully able to infer full-sibling relationships (including possible contribution of multiple fathers
to the mf pool (30)), and, in the absence of successful half-sibling inference, we combined
relatedness data with mitochondrial haplotypes to infer the *Nb* contributing to the offspring sample.
For the skin-snip mf, we were hampered by degraded, low-input DNA leading to uneven genome
coverage and unsuccessful relatedness estimation. However, good coverage over the

mitochondrial genome enabled us to infer a minimum of five females contributing to the offspring
samples.

Identification of shared parentage and *Nb* in parasitic helminths is a vital tool for discerning adult
worm burdens, as well as for distinguishing between treatment failure followed by repopulation and
treatment success followed by reinfection. This has the potential to inform the outcome of
macrofilaricide clinical trials (e.g. (16,34)), as well as the establishment of key parasite infection
dynamic parameters, such as the minimum adult worm burden and density-dependence, with the
assistance of statistical models (e.g. (35,36)). Both relatedness estimation and mitochondrial
genotyping have the potential to help address these questions in *O. volvulus* and other indirectly
transmitted parasites. For *O. volvulus*, a chromosome level genome assembly has been generated
(37), but no microsatellite, SNP or MLST panels exist for parentage analysis. Generating these
resources typically require laboratory systems (20), as well as time, money, and access to
samples. For pathogens for which laboratory life cycles are problematic to achieve, such as *O.*
volvulus, and for which samples are in limited supply – for a variety of different reasons, including
lack of funding, or difficult-to-obtain samples - generating panels of markers is difficult. Thus,
methods such as ours, that rely on low-coverage WGS to infer relatedness, may be particularly
suitable for studying infection dynamics and drug trial outcomes. Furthermore, once siblings have
been excluded, unrelated samples can be used for parasite population genomic analysis without
introducing biases from siblings. This method can easily be generalised to several parasite
systems, if a reference genome assembly has been developed. Reference genomes are becoming
increasingly common for even the most neglected of human helminthiases. However, methods for
SNP discovery such as RAD-Seq also offer a potential stopgap for relatedness estimation in the
absence of a reference genome, using relatedness methods such as SEQOUIA (38).

The sequencing effort reported here aimed for 5X coverage per-sample, which cost £30 per-
sample (around $36 US at the time of writing). Options for improving the cost-effectiveness include
performing downsampling experiments to establish the minimum depth-of-coverage required to
identify kinships. Simulations suggest that *NGSRelate* may be robust to 1X coverage (39).
Moreover, library preparation costs can be greatly reduced by adapting commercial library
preparation kits (e.g. dilution) (24,40).

Another possibility toward reducing cost and increasing ease of access is to restrict parentage
analysis in this form to the resequencing of mitochondrial genomes alone, which could be
performed by designing long-range PCR primers to conserved mitochondrial regions (26).
However, in the absence of population mitochondrial haplotype frequencies, it is impossible to
determine whether two individuals with the same mitochondrial haplotype are truly maternal
siblings, or whether their mothers are siblings, or if they have unrelated mothers with the same
mitochondrial lineage. Here, our data indicate either there is homogeneous distribution of mf
across an infected person despite adult worm location, or that there are multiple females with the
same mitochondrial haplotype on either side of the body. Given extensive *O. volvulus*
mitochondrial diversity and large population sizes (26), mitochondrial genotyping can at least
establish the minimum number of females contributing to a pool of larvae, a valuable piece of

information and a key parameter in models of helminth transmission (8,32,36), but it cannot
determine the maximum number of females contributing to the offspring being assessed.
225
Despite technical improvements in single-larva DNA extraction and low-input library preparation
(25), single-mf WGS is still technically challenging, with often unreliable results. The skin-snip mf,
for example, sequenced very poorly, with highly heterogeneous coverage across the nuclear
genome, perhaps indicative of degraded genomic DNA. Further work is needed to improve single-
mf WGS sample storage and preparation both for the employment of this approach for use in
epidemiological investigations and clinical trials, but also for improving *O. volvulus* population
genomic analyses which hitherto has only been possible with adult worms (41,42). Population
genomics based on mf would make evolutionary analysis of *O. volvulus* much easier and more
ethical, as it removes the need for surgical nodulectomies to access adult worms, although the
invasiveness of skin-snips remains a problem.
236
Ultimately, development of this method to the point where it can be reliably used in clinical trials
will require optimisation of sample storage, DNA extraction and library preparation techniques, as
well as more extensive testing against samples of mf for which parentage is known, so that the
variation in the range of kinship coefficients corresponding to a given relationship and the error
rates can be quantified. This is possible by performing further sequencing studies where adult
worms are available.
243
244 **Conclusion**
245
The decreasing cost of WGS, combined with technical and theoretical advances that
make it possible to infer relationships between helminth larvae from low-coverage WGS data, are
facilitating the development of techniques that enable interrogation of fine scale helminth infection
dynamics, as well as identifying treatment failure, versus repopulation from surviving adult worms.
This is important because many adult helminths, such as *O. volvulus* are, often inaccessible, with
their offspring being comparatively easy to sample. Here, we employed lcWGS, relatedness
estimation, and mitochondrial genotyping to identify parentage, maternal parentage and *Nb*,
showing that, when sample data quality is sufficient, we are able to identify the number of males
and females contributing to a sample of offspring and, even when nuclear genome data quality is
poor, the minimum number of females may still be estimated. This method, when developed
further has potential use in investigating the outcome of macrofilaricide trials for *O. volvulus*, as
well as for further probing multiple parasite infection dynamics, in systems for which validated
marker panels have not yet been developed.
259
260 **Materials and Methods**
261
262   Data (sequencing metadata, relatedness data, mitochondrial group data) were parsed
263 and analysed in *R* version *4.0.2*, (43), particularly using the *dplyr, gpplot2,* (44), *RColorbrewer*
264 (45)and *cowplot* (46) packages**.**
265
266 *Sample Collection*

267 **Part 1:** Onchocercomata containing adult female worms were surgically removed from patients as
268 part of the field study "Comparison of doxycycline alone vs doxycycline plus rifampicin in their
269 efficacy against onchocerciasis' in Ghana in 2009.
270
Part 2: Skin-snips were collected in Oct 2019. All infected patients presented at least one palpable
nodule. For Mf analysis, two skin biopsies of 1±3 mg were taken from the buttocks using a
corneoscleral (Holth) punch. Each biopsy was immersed in 100 ml of 0.9% NaCl solution in a well
of a microtiter plate. The skin biopsies were incubated overnight at room temperature to enable mf
to emerge. The solution was then transferred onto a slide for microscopic examination. The
biopsies were weighed using an electronic balance and mf load was calculated per mg skin. The
skin biopsies and the solution containing the free mf were then stored at -20 degrees C for
subsequent processing and analyses.
279
Sample Processing, DNA Extraction and Sequencing Library Preparation
Part 1: Intrauterine mf: 10 mf were dissected from three gravid *O. volvulus* adult females,
obtained by surgical excision from two nodules. (**Figure 1**) Adult females were washed once in
70% ethanol and three times in PBS. Insulin needles (29G) were used to dissect mf from the adult
female uterus. Individual mf were pipetted into a 96 well-plate, and DNA extracted using the
Beckman Coulter Agencourt DNAdvance gDNA purification kit following the manufacturer's
instructions. DNA was eluted into 35μl. The presence of *O. volvulus* was determined using O-150
primers (47). 26 μl of DNA were fragmented and made into libraries using the New England
Biolabs NEBNext Ultra FS II kit with unique dual barcodes. Libraries were pooled, and each pool
size selected using a Sage Science Blue Pippin with a range of 150-500bp. Further QC was
performed on an Agilent Bioanalyzer 2100 High-Sensitivity Chip.
291
292 **Part 2: Skin-snip mf:** 39 mf were dissected into PBS with insulin needles from two skin-snips, one
293 of the left and one of the right trochanter of a single patient. (**Figure 5**). The remaining processing
294 followed the methods used for the intrauterine mf.
295
296 The intrauterine mf and skin-snip mf were then sequenced in two separate runs on an Illumina
297 NextSeq Nano and an Illumina NovaSeq 6000, aiming for 13^e6 paired-end reads (10X mean
298 nuclear genome coverage of 2x75 paired-end reads) and 3.3^e6 (5X coverage of 2x150 reads)
299 respectively.
300
Mapping and Genotype-Likelihood Calling
Demultiplexed reads were trimmed using *Trimmomatic* v0.39 (48)**,** and aligned to the *O. volvulus*
reference genome vWBPS15 (37) using bwa-mem (49). PCR duplicates were identified and
removed with *GATK* (50). Samtools was used for SAM/BAM file manipulation, and calculation of
mapping statistics (51). BAM files were used as input for ANGSD (52), which was used to calculate
genotype likelihoods with the following filters: depth > 3, mapQ > 20, max-missingness <20% of
individuals, Hardy Weinberg Equilibrium (HWE) p-value 1e-6. See **Supplementary Tables 1 and**
2 for per-sample information, and [https://github.com/tristanpwdennis/onchogenome] for code.
309
Relatedness Estimation

The final genotype likelihood file was used as input for the *NGSRelate* software (39), which infers,
amongst others, the measures of relatedness: *R0, R1* and *KING-Robust-kinship*, as previously
described (28,29,39). The relationship between pairs of full siblings was based on previously
defined thresholds *.* All analyses downstream of *NGSRelate* were performed in R *4.0.2* (28,29)
(see https://github.com/tristanpwdennis/onchogenome). R0/R1 and R1/KING-Robust-kinships
were plotted with *ggplot2*. Graphs of related individuals were plotted in *iGraph* (53).
317
Mitochondrial Genotyping
Reads mapping to the *O. volvulus* mitochondrial genome were extracted from the BAM alignments
with samtools. SNPs were called using *Freebayes* (54) with a ploidy setting of 1, and filtered in
bcftools (51), based on depth > 3 reads, maximum missingness < 0.25, QUAL > 30. The resulting
filtered, multisample VCF file was converted into a FASTA alignment with bcftools *bcf-consensus*.
The alignment was used as an input for a minimum-spanning network in PEGAS (55), in R.
Clusters were defined based on the maximum level of divergence within-brood < 3 SNP
differences.
326

Ethics Statements

Funding

DNDi is grateful to the Bill & Melinda Gates Foundation (#INV008203) for funding this project.
PHLL is funded by the European Research Council (starting grant SCHISTO_PERSIST_680088,
Engineering and Physical Sciences Research Council (gEP/T003618/1) and the Medical Research
Council (MR/P025447/1). The Institute for Medical Microbiology, Immunology and Parasitology,
University Hospital Bonn received funding through a grant from the Liverpool School of Tropical
Medicine as part of the A-WOL Consortium funded by the Bill and Melinda Gates Foundation
(39284). Achim Hoerauf is funded by the Deutsche Forschungsgemeinschaft (DFG, German
Research Foundation) under Germany's Excellence Strategy – EXC2151 – 390873048"

Conflict of Interest Declaration

351
352 The authors declare no competing interests.
353

Data accessibility

355
356 All raw read data used in this study have been deposited in ENA under study accession
357 PRJEB63468. All the code (analysis code as an RMarkdown file, sequence alignment/processing
358 wrapping as shell wrappers), sequencing metadata, relatedness data, ANGSD output and
359 mitochondrial genome alignments are available at
360 https://github.com/tristanpwdennis/onchogenome.
361

362 **Acknowledgements**

364 Many thanks to James Cotton and Thomas Crellen for helpful discussions and feedback on the
365 manuscript.
366
367

 Figures

370
371 **Figure 1:** Schematic of *Onchocerca volvulus* microfilariae (mf) dissections from known parents.
372 (Results Section 1) 10 mf were dissected from the uteri of 3 adult female worms, derived from 3
373 nodulectomies from 2 patients. mf were whole-genome-sequenced.

375
Figure 2: Network of full-sibling relationships between intrauterine *Onchocerca volvulus* microfilariae. Nodes indicate intrauterine *O. volvulus* microfilariae and edges indicate full-sibling relationships. Nodes are coloured by known maternal parentage.

379
380
381   **Figure 4:** Medium-spanning network of intrauterine *Onchocerca volvulus* microfilariae
382   mitochondrial haplotypes. Colour indicates shared known maternal parentage. Dots on
383   connecting lines indicate the number of single nucleotide polymorphisms (SNPs) between
384   haplotype groups.

385
386
Figure 5: Schema of identifying parents of mf for which parentage is unknown. (Results Section 2)
39 mf were eluted from two skin-snips, from the left and right trochanter of a single patient.

389
390 **Figure 5:** Medium-spanning network of mitochondrial haplotypes of skin-snip collected
391 *Onchocerca volvulus* microfilariae from two locations on the body of a single patient. Pie colours
392 indicate the proportion of samples taken from the left or right trochanter of the patient. Dots on
393 connecting lines indicate the number of single nucleotide polymorphisms (SNPs)   between
394 mitochondrial haplotype groups.

Supplementary Figures

Supplementary Figure 1: Inferred kinship statistics for R0, R1 and KING-robust-kinship for pairs
of intrauterine *Onchocerca volvulus* microfilariae, with points coloured by whether individuals in a
pair were sampled from the same (within brood) or different (between brood) *O. volvulus* adult
female worms.

Supplementary Figure 2: Kinship statistics R0 and R1 for pairs of intrauterine *Onchocerca
volvulus* microfilariae, coloured by the inferred relationship using thresholds for full-sibling (Full-
Sib) relationships at R0 < 0.1, R1 > 0.5, and KING > 0.25 and thresholds for half-sibling (Half-Sib)
relationships as R0 > 0.1, R1 < 0.5, and KING < 0.25, informed from Figure 1.

Bibliography

1. Hotez PJ, Alvarado M, Basáñez MG, Bolliger I, Bourne R, Boussinesq M, et al. The Global Burden of Disease Study 2010: Interpretation and Implications for the Neglected Tropical Diseases. PLoS Negl Trop Dis. 2014 Jul 24;8(7):e2865.

2. Utzinger J, Booth M, N'goran EK, Müller I, Tanner M, Lengeler C. Relative contribution of day-to-day and intra-specimen variation in faecal egg counts of Schistosoma mansoni before and after treatment with praziquantel. Parasitology. 2001 May;122(5):537–44.

3. Mand S, Debrah A, Batsa L, Adjei O, Hoerauf A. Reliable and frequent detection of adult Wuchereria bancrofti in Ghanaian women by ultrasonography. Trop Med Int Health. 2004;9(10):1111–4.

4. Opoku NO, Bakajika DK, Kanza EM, Howard H, Mambandu GL, Nyathirombo A, et al. Single dose moxidectin versus ivermectin for Onchocerca volvulus infection in Ghana, Liberia, and the Democratic Republic of the Congo: a randomised, controlled, double-blind phase 3 trial. The Lancet. 2018 Oct;392(10154):1207–16.

5. Crellen T, Haswell M, Sithithaworn P, Sayasone S, Odermatt P, Lamberton PHL, et al. Diagnosis of helminths depends on worm fecundity and the distribution of parasites within hosts. Proc R Soc B Biol Sci. 2023 Jan 18;290(1991):20222204.

6. Hall A. Quantitative variability of nematode egg counts in faeces: a study among rural Kenyans. Trans R Soc Trop Med Hyg. 1981 Jan;75(5):682–7.

7. Duerr HP, Dietz K, Schulz-Key H, Büttner DW, Eichner M. Density-dependent parasite establishment suggests infection-associated immunosuppression as an important mechanism for parasite density regulation in onchocerciasis. Trans R Soc Trop Med Hyg. 2003 Mar;97(2):242–50.

8. Churcher TS, Filipe JAN, Basáñez MG. Density dependence and the control of helminth parasites. J Anim Ecol. 2006 Nov;75(6):1313–20.

9. Picq JJ, Jardel JP. Une methode d'evaluation des densites microfilariennes d'Onchocerca volvulus Leuckart, 1 893 chez des onchocerquiens.

10. Vos T, Flaxman AD, Naghavi M, Lozano R, Michaud C, Ezzati M, et al. Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. The Lancet. 2012 Dec;380(9859):2163–96.

11. Bakajika D, Senyonjo L, Enyong P, Oye J, Biholong B, Elhassan E, et al. On-going transmission of human onchocerciasis in the Massangam health district in the West Region of Cameroon: Better understanding transmission dynamics to inform changes in programmatic interventions. PLoS Negl Trop Dis. 2018 Nov;12(11):e0006904.

12. Koala L, Nikiema A, Post RJ, Paré AB, Kafando CM, Drabo F, et al. Recrudescence of onchocerciasis in the Comoé valley in Southwest Burkina Faso. Acta Trop. 2017 Feb;166:96–105.

13. Koala L, Nikièma AS, Paré AB, Drabo F, Toé LD, Belem AMG, et al. Entomological assessment of the transmission following recrudescence of onchocerciasis in the Comoé Valley, Burkina Faso. Parasit Vectors. 2019 Jan;12(1):34.

14. Eisenbarth A, Achukwi MD, Renz A. Ongoing Transmission of Onchocerca volvulus after 25 Years of Annual Ivermectin Mass Treatments in the Vina du Nord River Valley, in North Cameroon. PLoS Negl Trop Dis. 2016 Feb;10(2):e0004392.

15. Bah GS, Schneckener S, Hahnel SR, Bayang NH, Fieseler H, Schmuck GM, et al. Emodepside targets SLO-1 channels of Onchocerca ochengi and induces broad anthelmintic effects in a bovine model of onchocerciasis. PLOS Pathog. 2021 Jun;17(6):e1009601.

16. Krücken J, Holden-Dye L, Keiser J, Prichard RK, Townson S, Makepeace BL, et al. Development of emodepside as a possible adulticidal treatment for human onchocerciasis—

The fruit of a successful industrial–academic collaboration. PLOS Pathog. 2021 Jul;17(7):e1009682.

17. Hübner MP, Martin C, Specht S, Koschel M, Dubben B, Frohberger SJ, et al. Oxfendazole mediates macrofilaricidal efficacy against the filarial nematode Litomosoides sigmodontis in vivo and inhibits Onchocerca spec. motility in vitro. PLoS Negl Trop Dis. 2020 Jul 6;14(7):e0008427.

18. Schiefer A, Hübner MP, Krome A, Lämmer C, Ehrens A, Aden T, et al. Corallopyronin A for short-course anti-wolbachial, macrofilaricidal treatment of filarial infections. PLoS Negl Trop Dis. 2020 Dec 7;14(12):e0008930.

19. Jones OR, Wang J. COLONY: a program for parentage and sibship inference from multilocus genotype data. Mol Ecol Resour. 2010;10(3):551–5.

20. Steinauer ML, Christie MR, Blouin MS, Agola LE, Mwangi IN, Maina GM, et al. Non-Invasive Sampling of Schistosomes from Humans Requires Correcting for Family Structure. PLoS Negl Trop Dis. 2013 Sep;7(9):e2456.

21. Gower CM, Gehre F, Marques SR, Lamberton PHL, Lwambo NJ, Webster JP. Phenotypic and genotypic monitoring of Schistosoma mansoni in Tanzanian schoolchildren five years into a preventative chemotherapy national control programme. Parasit Vectors. 2017 Dec;10(1):593.

22. Johnson PCD, Hadfield JD, Webster LMI, Adam A, Mable BK, Keller LF. Bayesian paternity analysis and mating patterns in a parasitic nematode, Trichostrongylus tenuis. Heredity. 2010 Jun;104(6):573–82.

23. Sticca EL, Belbin GM, Gignoux CR. Current Developments in Detection of Identity-by-Descent Methods and Applications. Front Genet. 2021;12:722602.

24. Therkildsen NO, Palumbi SR. Practical low-coverage genomewide sequencing of hundreds of individually barcoded samples for population and evolutionary genomics in nonmodel species. Mol Ecol Resour. 2017;17(2):194–208.

25. Doyle SR, Sankaranarayanan G, Allan F, Berger D, Jimenez Castro PD, Collins JB, et al. Evaluation of DNA Extraction Methods on Individual Helminth Egg and Larval Stages for Whole-Genome Sequencing. Front Genet [Internet]. 2019 [cited 2022 Dec 18];10. Available from: https://www.frontiersin.org/articles/10.3389/fgene.2019.00826

26. Hedtke SM, Kuesel AC, Crawford KE, Graves PM, Boussinesq M, Lau CL, et al. Genomic epidemiology in filarial nematodes: transforming the basis for elimination program decisions. Front Genet. 2020;10:1282.

27. Durrant C, Thiele EA, Holroyd N, Doyle SR, Sallé G, Tracey A, et al. Population genomic evidence that human and animal infections in Africa come from the same populations of Dracunculus medinensis. Basáñez MG, editor. PLoS Negl Trop Dis. 2020 Nov;14(11):e0008623.

28. Waples RK, Albrechtsen A, Moltke I. Allele frequency-free inference of close familial relationships from genotypes or low-depth sequencing data. Mol Ecol. 2019;28(1):35–48.

29. Manichaikul A, Mychaleckyj JC, Rich SS, Daly K, Sale M, Chen WM. Robust relationship inference in genome-wide association studies. Bioinformatics. 2010 Nov;26(22):2867–73.

30. Hildebrandt JC, Eisenbarth A, Renz A, Streit A. Reproductive biology of Onchocerca ochengi, a nodule forming filarial nematode in zebu cattle. Vet Parasitol. 2014 Sep;205(1):318–29.

31. Ackerman MW, Hand BK, Waples RK, Luikart G, Waples RS, Steele CA, et al. Effective number of breeders from sibship reconstruction: empirical evaluations using hatchery steelhead. Evol Appl. 2017 Feb;10(2):146–60.

32. Anderson RM, Turner HC, Farrell SH, Truscott JE. Chapter Four - Studies of the Transmission Dynamics, Mathematical Model Development and the Control of Schistosome Parasites by Mass Drug Administration in Human Communities. In: Basáñez MG, Anderson RM, editors. Advances in Parasitology [Internet]. Academic Press; 2016 [cited 2023 Jun 27].

p. 199–246. (Mathematical Models for Neglected Tropical Diseases; vol. 94). Available from:
https://www.sciencedirect.com/science/article/pii/S0065308X16300720

33.   Minter A, Pellis L, Medley GF, Hollingsworth TD. What Can Modeling Tell Us About
Sustainable End Points for Neglected Tropical Diseases? Clin Infect Dis. 2021 Jun
15;72(Supplement_3):S129–33.

34.   Geary TG, Mackenzie CD, Silber SA. Flubendazole as a macrofilaricide: History and
background. Lustigman S, editor. PLoS Negl Trop Dis. 2019 Jan;13(1):e0006436.

35.   Neves MI, Gower CM, Webster JP, Walker M. Revisiting density-dependent fecundity in
schistosomes using sibship reconstruction. PLoS Negl Trop Dis. 2021 May;15(5):e0009396.

36.   Neves MI, Webster JP, Walker M. Estimating helminth burdens using sibship
reconstruction. Parasit Vectors. 2019 Dec;12(1):441.

37.   Cotton JA, Bennuru S, Grote A, Harsha B, Tracey A, Beech R, et al. The genome of
Onchocerca volvulus, agent of river blindness. Nat Microbiol. 2016 Nov;2(2):1–12.

38.   Huisman J. Pedigree reconstruction from SNP data: parentage assignment, sibship
clustering and beyond. Mol Ecol Resour. 2017;17(5):1009–24.

39.   Hanghøj K, Moltke I, Andersen PA, Manica A, Korneliussen TS. Fast and accurate
relatedness estimation from high-throughput sequencing data in the presence of inbreeding.
GigaScience [Internet]. 2019 May [cited 2023 Jan 1];8(5). Available from:
https://academic.oup.com/gigascience/article/doi/10.1093/gigascience/giz034/5481763

40.   Baym M, Kryazhimskiy S, Lieberman TD, Chung H, Desai MM, Kishony R. Inexpensive
Multiplexed Library Preparation for Megabase-Sized Genomes. PLOS ONE. 2015
May;10(5):e0128036.

41.   Choi YJ, Tyagi R, McNulty SN, Rosa BA, Ozersky P, Martin J, et al. Genomic diversity in
Onchocerca volvulus and its Wolbachia endosymbiont. Nat Microbiol. 2016 Nov;2(2):16207.

42.   Doyle SR, Bourguinat C, Nana-Djeunga HC, Kengne-Ouafo JA, Pion SDS, Bopda J, et
al. Genome-wide analysis of ivermectin response by Onchocerca volvulus reveals that
genetic drift and soft selective sweeps contribute to loss of drug sensitivity. Unnasch TR,
editor. PLoS Negl Trop Dis. 2017 Jul;11(7):e0005816.

43.   R Core Team. R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria. [Internet]. [cited 2023 May 3]. Available from:
https://www.eea.europa.eu/data-and-maps/indicators/oxygen-consuming-substances-in-
rivers/r-development-core-team-2006

44.   Wickham H, Averick M, Bryan J, Chang W, McGowan LD, François R, et al. Welcome to
the Tidyverse. J Open Source Softw. 2019 Nov;4(43):1686.

45.   Neuwirth E. RColorBrewer: ColorBrewer Palettes [Internet]. 2022 [cited 2023 Jan 1].
Available from: https://CRAN.R-project.org/package=RColorBrewer

46.   Wilke CO. cowplot: Streamlined Plot Theme and Plot Annotations for 'ggplot2' [Internet].
2020 [cited 2023 Jan 1]. Available from: https://CRAN.R-project.org/package=cowplot

47.   Detection of Onchocerca volvulus Infection by O-150 Polymerase Chain Reaction
Analysis of Skin Scratches on JSTOR [Internet]. [cited 2023 Jun 27]. Available from:
https://www.jstor.org/stable/30114150

48.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence
data. Bioinformatics. 2014 Aug;30(15):2114–20.

49.   Li H, Durbin R. Fast and accurate short read alignment with Burrows–Wheeler
transform. Bioinformatics. 2009 Jul 15;25(14):1754–60.

50.   McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, et al. The
Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA
sequencing data. Genome Res. 2010 Sep;20(9):1297–303.

51.   Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, et al. Twelve years of
SAMtools and BCFtools. GigaScience. 2021 Jan 29;10(2):giab008.

52.   Korneliussen TS, Albrechtsen A, Nielsen R. ANGSD: Analysis of Next Generation

Sequencing Data. BMC Bioinformatics. 2014 Nov 25;15(1):356.
53. Csardi G, Nepusz T. The igraph software package for complex network research.
54. Garrison, Erik, Marth, Gabor. Haplotype-based variant detection from short-read
sequencing. arRXIV. 2012 Jul 4;1207.3907.
55. Paradis E. pegas: an R package for population genetics with an integrated-modular
approach. Bioinformatics. 2010 Feb;26(3):419–20.