Tick-borne zoonotic flaviviruses and *Borrelia* infections in wildlife hosts: what have field studies contributed?

Armelle Poisson[1,2], Thierry Boulinier[2], Laure Bournez[3], Gaëlle Gonzalez[4], Camille V. Migné[4], Sara Moutailler[5], Bruno Faivre[6], Raphaëlle Métras[1]

[1]Sorbonne Université, INSERM, Institut Pierre Louis d'Epidémiologie et de Santé Publique (IPLESP, UMRS 1136), Paris, France.

[2]Centre d'Ecologie Fonctionnelle et Evolutive, UMR 5175, CNRS, Université de Montpellier, IRD, EPHE, Montpellier, France.

[3]Laboratoire de la Rage et de la Faune Sauvage, ANSES, Malzéville, France.

[4]ANSES, INRAE, Ecole Nationale Vétérinaire d'Alfort, UMR Virologie, Laboratoire de Santé Animale, Maisons-Alfort, France.

[5]ANSES, INRAE, Ecole Nationale Vétérinaire d'Alfort, UMR BIPAR, Laboratoire de Santé Animale, Maisons-Alfort, France.

[6]Biogéosciences, UMR 6282, CNRS, Université de Bourgogne, EPHE, Dijon, France.

Abstract

Tick-borne flaviviruses and *Borrelia* spp. are globally spread pathogens of zoonotic potential that are maintained by a transmission cycle at the interface between ticks and vertebrate hosts, mainly wild animals. Aside data on pathogen burden in ticks, information on the status of various hosts relative to infection could be critical. We reviewed how those infections have been studied in wildlife species in the field to discuss how collected data provided relevant epidemiological information and identify needs for further studies. The literature was screened for populational studies on direct or antibody detection for tick-borne *Borrelia* spp. and flaviviruses in animals, in the wild. Overall, *Borrelia* spp. were more studied (73% of case studies, representing 297 host species) than flaviviruses (27% of case studies, representing 114 host species). Studies on both *Borrelia* spp. and flaviviruses focused mainly on the same species, namely bank vole and yellow-necked mouse. Most studies were order-specific and cross-sectional, reporting prevalence at various locations, but with little insight into the underlying epidemiological dynamics. Species with potential to act as reservoir hosts were overlooked, notably passerine birds. We highlight the necessity of collecting both demographics and infection data in wildlife studies, and to consider communities of species, to better estimate zoonotic risk potential in the One Health context.

Keywords
Tick-borne diseases, wildlife, *Borrelia*, flavivirus, reservoir, sentinel, host.

I-Introduction
Vector-borne and zoonotic infections are emerging threats to public health[1]. Tick-borne pathogens (TBP) are maintained by multi-hosts complex epidemiological cycles[2]. Transmission to humans occur on

specific occasions, and understanding and quantifying it is necessary to control diseases. Persistence of
tick-borne and zoonotic pathogens is allowed at the interface between tick vectors and a large variety of
vertebrate hosts permitting ticks to feed and complete their life cycles. In those systems, early tick stages
(larvae and nymph) usually feed on small mammals and birds, while mature stages mainly feed on larger
vertebrates (such as deer, boar, cattle)[3].

Beyond their feeding role for ticks, vertebrate species can act as pathogen reservoir or sentinel hosts.
Infection and exposure data from both give precious information on dynamics of the studied TBP[4]. Whilst
reservoir hosts allow the pathogen to multiply and be transmitted further, sentinel species are
characterized by their ability to reflect underlying epidemiological phenomena and by being of easy
access[5]. For instance, wild red and Arctic foxes have been suggested as sentinels for human and animal
toxoplasma risk in Canada, based on serology and direct detection on carcasses[6]. Also, animal ecology
can drive variations in risk of transmission to human[7]. Therefore, because understanding pathogens'
transmission and distribution in wildlife is prerequisite to addressing risk of transmission, quantitatively
tracking infection and exposure to pathogens in those hosts is necessary.

Tick-borne *Borreliaceae* and flaviviruses are TBP of zoonotic potential which are the most prevalent in the
temperate regions of the world[8], and are thus of interest for human and veterinary public health. Tick-
borne encephalitis virus (TBEV), first described in Austria in 1931, is the most common tick-borne flavivirus
(TBFV) in Europe with several endemic foci in Europe and Asia and growing number of human cases[9]. Its
ecology and epidemiology have early been studied in Europe, especially in Czech Republic[10]. Other TBFV
have later been isolated from wild animals and their ticks, such as Meaban virus (MEAV) in Brittany,
France[11], or from human cases, such as Powassan virus (POWV) in Powassan town, Canada[12].
Prominent vectors of TBFV are hard ticks *Ixodes ricinus* in Western and Central Europe and *I. persulcatus*
in Eastern Europe, as well as local tick species in other parts of the world. Similarly transmitted bacterial
genospecies of the *Borrelia burgdorferi* sensu lato (Bbsl) complex are the causative agents for Lyme
disease, one of the most common TBD in the Northern Hemisphere. Other tick-borne *Borrelia* spp. cause
relapsing fever in humans[13].

Recent reviews on tick-borne *Borrelia* spp. and TBFV have focused on getting information on the biology
of infection in vertebrate hosts[14,15], reporting prevalence and clinical cases in human[16,17] and
discussing diagnostic methodologies[18,19], the role of non-vector transmissions for TBFV[20,21], or on
the importance of modelling TBP regarding climate change[22,23]. Some reviews have focused on wildlife
hosts but only at a national or continental scale[24,25], or on reservoirs associated with flaviviruses in
general, with little focus on TBFV[26]. Finally, a lot of research effort has focused on studying the infectious
agents in the ticks[27,28].

Despite their major importance in zoonotic transmission, no study has reviewed the body of evidence in
wildlife hosts from field data at a global scale. Yet, data on the level of infection and exposure in free-
ranging populations are critical to model the dynamics of infectious diseases[29]. Strong inference about
the transmission processes underlying those dynamics often requires a combination of approaches,
notably experiments to ascertain the role of reservoir[30], but data on infection and exposure of hosts are
prerequisite in most cases. When it comes to wildlife species, a strong heterogeneity in the type of field
data is nevertheless expected, from local cross-sectional sampling of particular species, to broad spatial
surveys over series of years. Estimating epidemiological parameters also requires accounting for potential
biases in detection probabilities and host population parameters[31], which demands specific designs.

The aim of this review was thus to investigate how tick-borne *Borrelia* spp. and flaviviruses infection
burdens have been studied in free-ranging wildlife (that is, excluding studies of TBP in ticks) to discuss how
collected data provided relevant epidemiological information, and identify needs for further studies. We
predicted that strong heterogeneity would appear in the types of hosts investigated, with a high number
of studies conducted on common species sharing landscape use with humans in the Northern Hemisphere.
We expected few studies including whole communities of potential hosts and few studies estimating TBP
burden while accounting for temporal and spatial variations and uncertainties. We anticipated that
monitoring of wildlife diseases would have been difficult and that knowledge of what is precisely measured
would have been uncertain.

II-Studied hosts and their potential role in epidemiological cycles

95 We reconstructed 947 "case studies" (unique combinations of species-pathogen-area-study period)
96 studying overall 347 host species, from the 314 articles matching the selection criteria (Figure 1). Detailed
97 methodology on the literature search and metrics on the results are available in SuppInfo1. Figure 2 links
98 the order of the tested species to the investigated pathogen.

99
100 *Figure 1:* **Diagram of inclusion and exclusion in database screening.** See SupInfo1 for details on literature search and additional
101 searches.

a-Historical papers shaped research on rodents as reservoirs

Rodentia was the most investigated order, represented by more than half (363/689) of the *Borrelia* study
cases and more than a third (91/258) of the TBFV study cases. White-footed mouse (*Peromyscus leucopus*)
was investigated in the first published studies from Northern America, and has been regarded since the
eighties as the most common *Borrelia* spp. reservoir host in wildlife in Northern America[32]. Regarding
TBFV, a series of studies were conducted initially in Northern America too. The first published studies
looked at Powassan virus (POWV) in *Peromyscus* genus, only present in Northern America, and secondly
in *Microtus* and *Mus* genera, present in the continent as well. In addition, in Canada, a series of studies

from the same authors, presented rodents as primary reservoirs for POWV. For this, species such as
Tamias, *Urocitellus* and *Callospermophilus* chipmunks and *Tamiasciurus* and *Sciurus* squirrels were
investigated, with average seroprevalence of 15%[33–41]. Groundhogs (*Marmota monax*) sampled in the
same studies reported the highest seroprevalence (around 43%)[37–39]. In parallel, several studies on the
eco-epidemiology of TBFV were conducted in the former Union of Soviet Socialist Republics but were not
published in English, although some articles are available with translation systems such as those of
NAMRU-3 (see limitations in SuppInfo1). Rodents were later sampled on other continents and TBFV
detections were several times successful in Europe in *Apodemus*, *Microtus* and *Myodes* genera[42], and in
Micromys, *Mus* and *Sciurus* for *Borrelia*[42,43]. In Europe, the bank vole (*Myodes glareolus*) and yellow-
necked mouse (*Apodemus flavicollis*) were recognized as reservoirs for TBEV[44]. Bbsl was detected in
Tamias, *Tamiasciurus* and *Neotoma* in North America[45]. Furthermore, *Gerbillus*, *Mastomys*, *Praomys*
and *Apodemus*, *Cricetulus*, *Eothenomys*, *Meriones*, *Mus*, *Niviventer* and *Rattus* bore relapsing fever
associated *Borrelia* respectively in Africa and in Asia. Notably, Eastern grey squirrels (*Sciurus carolinensis*)
similarly carried *Borrelia* DNA, both in their native countries and where they are invasive in Europe.

b-Deer, broadly studied as sentinels

The second most investigated order by the number of tested individuals was Artiodactyla. Artiodactyls
were mainly represented by the *Cervidae* family, in terms of numbers of tested animals (82%,
26,090/31,862) and investigated species (56%, 9/16) (Figure 3B and 3D). Deer are considered incompetent
reservoirs for *Borrelia* spp. by some authors because of borreliacidal properties of their sera[46]. Yet,
studies showed that the complement borreliacidal activity depends on strain and also exists on other
Borrelia strains for other vertebrates, including rodents[47]. Despite their poor competence as reservoirs,
deer are frequently met across the world and can be infected by feeding adult ticks they commonly
host[48]. In *Cervidae*, species locally present in America, Europe and Asia showed similar level of TBP
contact, with average seroprevalence of 18%. Restraint home ranges, of no more than 10km² among the
different deer species[49,50], allowed investigating infection processes at small spatial scales. Moreover,
surveillance activities and pathogen detection can be facilitated through deer hunting, allowing
widespread sampling programmes on a regular basis for reasonable costs and working efforts[51]. As such,
deer could help identifying endemic or emerging risk areas and give information on underlying processes,
and were repeatedly considered as sentinels. Other game artiodactyls identified in this review, such as
wild boar (*Sus scrofa*), offer the same kind of possibilities for sentinel surveillance.

c-Birds, reservoirs, have been overlooked

Birds were also reported as reservoirs in the literature[52]. Contrary to rodents, birds may have been
overlooked. Indeed, although birds represented about 19% (129/689) of the *Borrelia* study cases (Figure
2) and 33% (85/258) of TBFV cases, they originated from only 12% (28/228) and 18% (16/89) of the
publications. In those, pathogen presence was often screened in birds caught in a banding configuration.
For instance, in one study, Newman *et al.* caught birds on the University of California's campus in 2003-
2004[53]. Their wide-range study allowed detecting Bbsl in 23 species out of 53 caught species. Mistnetting
is a common field method efficient for capturing birds, hence for detecting potential reservoirs. However,
this approach is mostly favourable for Passeriformes, and passerines were the avian order the most often
studied as potential reservoirs for TBP. In addition, for migratory species or those exploiting large areas
during their life cycle, the infection status might not always reflect the local infection processes.
Nevertheless, combined with movement information, they can unravel critical information on pathogen
dissemination over long distances and between various ecosystems[54]. We hypothesized that limited

availability of data on birds might be due to logistical difficulties related to blood sampling in the wild
(training and legal requirements, ethical concerns for endangered species, practical implementation).
However, antibodies against Gadgets Gully and Saumarez Reef viruses were extensively studied in a
collection of Australian seabirds' sera[55]. This constitutes a unique occurrence of exposure information
on rare TBFV, but sadly, no clarification was made on which species retrieved seropositivity.

d-Least studied species with growing interest
Other vertebrate species of interest have been less studied. Recently, however, new species of tick-borne
Borrelia spp. have been sought and encountered in vertebrates from other orders. For example,
squamates were investigated for *Borrelia* in America, Africa and Europe, and 18 species associated with
Bbsl. Anti-Bbsl antibodies were detected in raccoon (*Procyon lotor*) in the USA and raccoon dog
(*Nyctereutes procyonoides*) in Asia, as well as where they are invasive in Europe or Asia. In India, primates
were found seropositive to KFDV, a TBFV close to POWV and DTV[56]. In the UK, an isolated clinical case
of lethal borreliosis was reported in a *Pipistrellus* spp. bat[57]. Neotropical bats have been identified to
carry particular strains of relapsing fever *Borrelia* and Bbsl in Central and Southern America. Studies in
Central America have been conducted as follow-back of a paper published in 1968 by Marinkelle &
Grose[58]. Some areas are currently investigated more in depth or monitored. For instance, Tribeč
mountain region in Czech Republic is continuously investigated for TBEV since the seventies.

III-Diagnostic tests: current *versus* past infections
Direct and indirect detection methods were at different use depending on the pathogen. Whilst most
Borrelia studies (71%) looked for pathogen's genome with molecular techniques to identify ongoing
infections, 89% of flaviviruses studies used serological methods. Long-lasting in the organisms, *Borrelia*
spp. were often sought by PCR (60% of case studies overall). Skin and blood were used for spirochetes PCR
detection, allowing positive samples to be sequenced and species identified, although the use of blood
compared with other organs to directly detect *Borrelia* is discussed[59]. Determination of *Borrelia*
subspecies was useful in understanding the role of hosts in disease cycles[60]. Different *Borrelia* have been
preferentially detected in different hosts, such as *B. lusitaniae* which was the only Bbsl genospecies found
in lizards in Europe.

In contrast, the short viremia noted after flaviviruses infection as demonstrated by experimental
infections[61], makes direct viral detection challenging especially in wildlife. PCR was therefore rarely used
for flaviviruses (in only 11% of TBFV studies), but allowed sequencing of TBEV subtypes that clustered with
TBEV-Eur (European) in Croatia, Slovakia and the Netherlands[62–64], TBEV-Eur and TBEV-Sib (Siberian) in
Finland[65], TBEV-Eur, TBEV-Sib and TBEV-FE (Far Eastern) in Russia[66] and TBEV-Him (Himalayan) in
China[67], and evidenced co-infection in Russia[66]. Identifying flaviviruses will help understanding further
the ecology and dynamics of pathogens and will assist disease surveillance with phylogenetics providing
information on circulation and evolution of TBP[68]. For example, TBEV could be mistaken for Looping Ill
virus (LIV) causing severe conditions mainly in sheep[51].

Flaviviruses infections induce immune response with neutralizing antibodies that could be detected
lifelong in blood in human[69] and up to 168 days after experimental infection in bank voles[70]. ELISA
were the most used serological techniques (89% of flaviviruses cases, and 38% of overall cases). Serological
information in combination with metadata can provide precious information on pathogen
transmission[71]. Yet, cross-reactivity in wide-range immunoassays may hamper data interpretation[72].
For instance, anti-MEAV antibodies were surprisingly detected in *Cervidae*[73], whilst MEAV belongs to

the seabird group in TBFV phylogeny[74]. Antibodies against LIV have been unusually identified in seabirds
Fratercula arctica and *Hydrobates leucorhous*[75]. In addition, whilst ELISA cut-off is generally reproduced
from manufacturer, some authors proposed its determination via mathematical fit on observed data to
optimize sensitivity and specificity[76]. Others even chose to present raw ELISA results as a signal of
contact with TBFV, without indicating cut-off nor prevalence[77].

Publications about TBFV outside of TBEV and POWV were marginal and specific detection for these TBFV
antibodies was not always commercially available. Arnal *et al.* found MEAV-seropositivity in *Larus*
michaellis michaellis egg yolks in France using flavivirus-ELISA cross-reactivity properties, with the ID-Vet
ID Screen® kit designed for the mosquito-borne flavivirus West Nile virus (WNV)[78]. Similarly, through
both COMPAC® WNV-ELISA and exclusion via serum neutralization tests (SNT), Jurado-Tarifa *et al.*
investigated flaviviruses in Spanish birds of prey and detected evidence of exposure to flaviviruses
different from the mosquito-borne WNV and Usutu (USUV) viruses and from MEAV in *Circus pygargus*,
Falco tinnunculus, *Hieraaetus pennatus* and *Tyto alba*[79]. This relies on lack of specificity of commercially
available anti-WNV antibodies detection kits. Some commercial TBEV ELISA have also been compared for
their reliability according to the TBEV-strain used and for the at-risk cross-reactivity[80]. In the UK, Holding
et al. recorded similar LIV and TBEV seroprevalences in cervids although TBEV was though not to be
present on the British Isles[51]. Holding et *al.* used the FSME/TBE All-Species Progen® ELISA and a
hemagglutination assay specific to LIV. For confirmation, Holding *et al.* sequenced TBEV in a tick itself.
Ytrehus *et al.* also showed presence of both LIV and TBEV antibodies in Southern Norway in *Alces alces*
and *Capreolus capreolus* using ELISA kits for TBEV and hemagglutination for LIV as well as SNT for
verification[81]. However, their results appeared differentiable for the two pathogens. On the opposite,
Bournez *et al.* chose to deliver their final TBEV micro-neutralization test serology results under the
TBEV/LIV label, after a first antibodies screening in *Capreolus capreolus* and *Sus scrofa* with the ID-Vet ID
Screen® WNV-ELISA[82]. TBEV and LIV as well as the Turkish Sheep Encephalomyelitis virus (TSEV) and the
Spanish Sheep Encephalomyelitis virus (SSEV) belong to the TBEV serocomplex[56], hence can particularly
easily be mistaken in serology.

IV-Infection and exposure in hosts: summary measures

A total of 297 wild animal species (mammals, birds and lizards) were investigated for *Borrelia* (Figure 3A
and 3B). When considering all *Borrelia* species, all studies confounded, overall detection and
seroprevalence tended to uniformize between 15% and 30% as the number of tested individuals increased
(Figure 4A). In studies with over 50 tested individuals, the proportion of active infections ranged from 0.1%
for *B. lusitaniae* in Eastern grey squirrel (*Sciurus carolinensis*) to 76% for Bbsl in white-footed mouse[83],
and from 0.2% for relapsing fever associated species in house mouse (*Mus musculus*)[84] to 50% for *B.*
miyamotoi in wild turkey (*Meleagris gallopavo*)[85]. In mammals, including rodents, artiodactyls and
carnivores, the level of direct detection and seroprevalence of *Borrelia* spp. was between 20 and 25% for
orders with over 100 specimens tested (Figure 3A and 3B).
In rodent studies testing more than 50 individuals, the average proportion of *Borrelia* direct detection was
11.2% (median 5.3%, range 0.2%-76%) and average seroprevalence was twice higher (mean 21.2%, median
15%, range 0.7%-89%) (Table 1, Figure S2). Variations were reported along study sites and periods
regardless of taxonomy. Beyond that such TBP might be widespread, this could be explained by the ecology
broadly shared by rodents, living close to the ground where foraging infected ticks can be
encountered[86], and therefore leading to similar exposure. Variations in prevalence across studies were
observed notably for a same species, such as for Bbsl direct detection in yellow-necked mouse tissues

varying from 5.8% in Hungary in 2014-2015[87] to 23% in Germany in 2012-2014[88]. In artiodactyls studies with more than 50 sampled animals, the average prevalence of *Borrelia* DNA detection was 14.4% and average seroprevalence was 23% (Table 1, Figure S2). They were of 12.6% and 24.2% in birds (Table 1, Figure S2). Among birds, Columbiformes, Passeriformes, Charadriiforms and Galliformes were the most studied in terms of number of tested individuals, with Charadriiforms and Passeriformes reaching 29% of seroprevalence (Figure 3B). Among the least studied species, Chiropters presented both seropositivity (36%) and pathogen prevalence (25%) of rather high level (Figure 3A).

TBFV or antibodies were searched and detected in 114 mammal and bird species (Figure 3C and 3D). The overall average seroprevalence ranged from 8% for MEAV to 51% for TYUV (Figure 4B), but with heterogeneous sample sizes. Among studies with more than 50 tested individuals, seroprevalences ranged from 0.07% for MEAV in red deer (*Cervus elaphus*)[73] to 68% for TYUV/MEAV/SREV-like in European herring gull (*Larus argentatus*)[89]. For TBFV of the mammalian group, seroprevalence ranged from 0,2% for TBEV/LIV in roe deer (*Capreolus capreolus*)[82] to 63% for TBEV in European bison (*Bison bonasus*)[90]. In rodent studies with at least 50 samples, the average TBFV-seropositivity was 8.9% (median 4.95%, range 0.4-44%) (Table 1, Figure S2), lower than for *Borrelia*. No studies looking at rodent host exposure to Kyanasur Forest Disease virus (KFDV) and Yamaguchi virus (YV) were recorded, although they have terrestrial hosts[91]. The average seroprevalence in artiodactyls was 10% and more than twice in birds reaching 21.9% (Table 1, Figure S2). Most studied avian orders were Passeriformes, Charadriiforms, Galliformes and Procellariforms, with highest seroprevalence for Passeriformes (about 40%), followed by Charadriiforms (27%) and Galliformes (21%) when Procellariforms tended to have less seropositive individuals (Figure 3C). Whilst 80% of Pelecaniformes had antibodies to TYUV, for this order only 8 European shags (*Phalacrocorax aristotelis*) were tested on a single location (in Brittany, France) in the eighties[89], highlighting the weight small scale studies could have. Exceptionally, mammalian Eulipotyphla were well-studied by PCR techniques and showed high rates of viremia (of about 50%)[92].

Table 1: **Prevalence of direct detection and seroprevalence by Order, extracted from studies with more than 50 individuals.** n = number of study cases, median values, mean, range and inter-quartile range (IQ). See plots of distributions in Figure S2.

Studies with >=50 tested individuals in:		Rodents (n=230)	Artiodactyls (n=83)	Birds (n=55)
Borrelia	**Direct detection**	n = 135; median 5.3%; mean 11.2%; range 0.2-76%; IQR 2-15%;	n = 15; median 10.6%; mean 14.4%; range 0.8%-49.7%; IQR 6.4-17.4%;	n = 22; median 8.8%; mean 12.6%; range 1-51%; IQR 4.4-14%;
	Seroprevalence	n = 43; median 15%; mean 21.2%; range 0.7-89%; IQR 9.2-31.5%;	n = 25; median 19%; mean 23%; range 0.8-68.8%; IQR 7.2-30%;	n = 14; median 17%; mean 24.2%; range 3-78.6%; IQR 9.5-26.6%;
TBFV	**Seroprevalence**	n = 44; median 5.0%; mean 8.9%; range 0.4-44%; IQR 3.5-10.5%;	n = 42; median 5.4%; mean 10%; range 0.07-62.7%; IQR 2.2-10%;	n = 18; median 19.7%; mean 21.9%; range 0.5-68.1%; IQR 4.1-33.8%;

Order

Pathogen

 Figure 2: **Sankey diagram pairing study cases by pathogen and order.** Width of the ribbon is proportional to the number of cases
 included in this order-pathogen couple. Data are available in Supplementary Dataset 1.

Figure 3A-D: **Investigation efforts for TBP by Orders.** Both serological and "active infection" data are presented on the same plot. Top row shows for *Borrelia* spp. **(A)** the average percentage of animals positive to *Borrelia* by order, **(B)** the number of tested animals in each order. Bottom row shows for TBFV **(C)** the percentage of animals positive to TBFV by order, **(D)** the number of tested animals in each order. Most publications regarding TBFV used an indirect methodology (antibody detection), direct detection was used in only 11% of TBFV cases. **(B)** and **(D)** show the number of tested individuals on a log10 scale, and the number of different host species tested in each order is shown at the bottom of each bar. Orders are grouped by class, Mammalia are shown to the left, Aves to the right, and for *Borrelia* spp., Lepidosauria at the bottom. Data are available in Supplementary Dataset 1.

Figure 4A-B: **Average prevalence of pathogen detection (dark blue labels, direct tests) and antibodies (light grey labels, indirect tests) in the tested individuals, rounded to 0 digit.** The x-axis represents the log10 of the sum of individuals that have been tested for **(A)** *Borrelia* **spp.** and **(B) Flaviviruses** in the selected publications, the y-axis represents the % of positive. DTV (Deer Tick Virus) in oldest publications might have been detected under the name of POWV, because DTV formerly corresponded to POWV lineage II. Data are available in Supplementary Dataset 1.

V-Study designs and eco-epidemiological parameters

Implemented study designs varied in spatial and temporal dimensions, resulting in varying quality in data acquisition. Most publications conducted cross-sectional studies (57%, 180/314), and 20 did not inform a study period. Some studies conducted repeated sampling without taking into consideration the sampling period in data presentation. Spatial resolution varied from locality to country scale, with often provinces chosen as spatial scale. Inconstancy came from different focuses, either on locally targeted area or over broader-scale exposure. In each study, sample sizes were usually limited, with median sample size at 40 (IQR 14-119). Overall, the way data were collected did not allow taking geography as well as other confounding factors into account. Forty-one (41) publications marked captured individuals but marking techniques were not always specific and trustable, some authors simply relying on scars made by biopsy or on feather clipping. The aim of marking was sometimes only to avoid resampling, with immediate release in case of recapture. However, 24 publications used individual marks and repeated capture occasions. Nevertheless, capture-mark-recapture data was not systematically analysed using relevant techniques. It is however an important specific sampling approach that permits accounting for detection uncertainty. For instance, seroprevalence and capture-mark-recapture data were recently used in a *Borrelia* host system[93]. Similar approaches were also used in other wildlife and infection systems, such as brucellosis in Alpine ibex (*Capra ibex*)[94].

Moreover, metadata informing basic demographics were lacking in almost all studies whereas data on age structure can be key in understanding susceptibility to pathogen at the individual level[95], but also transmission processes at the population level. Mammals and birds or squamates were not studied together in the same areas. Designs tended to investigate a single order by setting field material only fit for this order, whilst theoretical definitions of reservoir and sentinel suggest that reservoirs are communities rather than species[30]. The underlying community of hosts was seldomly considered. Furthermore, although sometimes convenient, getting material from hunted, road-killed or rescued

animals implies biases in detection that should be accounted for. Such sampling strategies often lead to small sample sizes and difficult interpretation. Moreover, as recommended by Yoccoz *et al.* on biodiversity monitoring, sampling designs and data analyses need to account for uncertainty on host detectability and spatial variability, for better estimating the state and rates of change of relevant ecological communities[96]. Thereby, acquiring infection and exposure data in wildlife should target communities of species in longitudinal studies with capture-marking-recapture or patch-occupancy modelling.

VI-Complementarity of laboratory and tick data

In our review, some infectious agents received less attention, such as TYUV, MEAV and SREV seabird flaviviruses. Whilst such agents may be less pathogenic, they can still be useful to understand transmission processes. In addition, their virulence could vary between host species and they could emerge as possible pathogens of concern. For example, TUYV was unexpectedly found pathogenic and deadly in seabirds in a study conducted by Berezina *et al.* in 1974[97]. Therefore, combining data on pathogenicity from the field and the laboratory is still needed.

In the literature, models and experimental infections have attempted to characterize reservoir competence[97]. Especially, xenodiagnoses are used to detect host reservoir capacities and tick vector competences in experimental settings[98]. Ticks from the field have been extensively examined for pathogens. From the 827 articles matching the literature search, 224 publications were directly looking into ticks for pathogens with methods ranging from PCR to metagenomics. Studies on ticks can be complementary to data from hosts insofar as they analyse tick-host interactions. Their deployment is less invasive on hosts than blood or tissue sampling. They need less expertise in the field and are subject to less strict ethical protocols. Such studies can serve as early detection methods when the pathogen is poorly known. For instance, whilst data on TBP hosts in Australia were limited, studying pathogens in ticks demonstrated circulation of *Borrelia* in squamates and monotremes[99]. In addition, identification of ticks' life stages allows to infer ticks' contamination capacity, although other ways of transmission such as vertical transmission in vertebrate hosts have been reported for some agents[100]. However, accounting for the whole local host community and their infection status remains as important.

VII-Conclusion

TBP are present in a wide range of wild animal species, and rodents and artiodactyls were the most investigated. Some infectious agents, rare TBFV or particular *Borrelia* strains, were little studied and information on their infecting capacities is still needed. Diagnostic detection methods varied with the targeted pathogens. Selected publications showed that study designs were heterogenous, both in data collection and analysis. Key information on host age and precise location was often unavailable; 58% (200/342) of the studies conducted fieldwork at one time point, and very few studies used quantitative methods to analyse collected data. Quantifying transmission necessitates gathering empirical data from populational studies, including hosts demographics and identifying which species or community act as reservoirs and sentinels. Whilst we acknowledge that gathering this information is difficult, this review highlights the necessity to estimate eco-epidemiological parameters in wildlife infection studies. The subject discussed in this review concerns a multidisciplinary scientific community (ecology, epidemiology, microbiology, parasitology, modelling, public health), emphasizing that getting and interpreting zoonotic infection data from wildlife in a rigorous way is important to build workable One Health programmes.

Data availability

The data that support this study are available as Supplementary Dataset 1. Source data are provided with this paper.

Reference

[1]   M.E.J. Woolhouse, Population biology of emerging and re-emerging pathogens, Trends Microbiol. 10 (2002) s3–s7. https://doi.org/10.1016/S0966-842X(02)02428-9.

[2]   A. Estrada-Peña, J. De La Fuente, The ecology of ticks and epidemiology of tick-borne viral diseases, Antiviral Res. 108 (2014) 104–128. https://doi.org/10.1016/j.antiviral.2014.05.016.

[3]   K.D. McCoy, N. Boulanger, Tiques et maladies à tiques: biologie, écologie évolutive, épidémiologie, IRD, Marseille, 2015.

[4]   D.J. Becker, S.N. Seifert, C.J. Carlson, Beyond Infection: Integrating Competence into Reservoir Host Prediction, Trends Ecol. Evol. 35 (2020) 1062–1065. https://doi.org/10.1016/j.tree.2020.08.014.

[5]   J.E.B. Halliday, A.L. Meredith, D.L. Knobel, D.J. Shaw, B.M.D.C. Bronsvoort, S. Cleaveland, A framework for evaluating animals as sentinels for infectious disease surveillance, J. R. Soc. Interface 4 (2007) 973–984. https://doi.org/10.1098/rsif.2007.0237.

[6]   É. Bouchard, R. Sharma, A. Hernández-Ortiz, K. Buhler, B. Al-Adhami, C. Su, H. Fenton, G. G.-Gouin, J.D. Roth, C.W. Rodrigues, C. Pamak, A. Simon, N. Bachand, P. Leighton, E. Jenkins, Are foxes (Vulpes spp.) good sentinel species for Toxoplasma gondii in northern Canada?, Parasit. Vectors 15 (2022) 115. https://doi.org/10.1186/s13071-022-05229-3.

[7]   P. Eby, A.J. Peel, A. Hoegh, W. Madden, J.R. Giles, P.J. Hudson, R.K. Plowright, Pathogen spillover driven by rapid changes in bat ecology, Nature 613 (2023) 340–344. https://doi.org/10.1038/s41586-022-05506-2.

[8]   N. Riccardi, R.M. Antonello, R. Luzzati, J. Zajkowska, S. Di Bella, D.R. Giacobbe, Tick-borne encephalitis in Europe: a brief update on epidemiology, diagnosis, prevention, and treatment, Eur. J. Intern. Med. 62 (2019) 1–6. https://doi.org/10.1016/j.ejim.2019.01.004.

[9]   D. Amicizia, A. Domnich, D. Panatto, P.L. Lai, M.L. Cristina, U. Avio, R. Gasparini, Epidemiology of tick-borne encephalitis (TBE) in Europe and its prevention by available vaccines, Hum. Vaccines Immunother. 9 (2013) 1163–1171. https://doi.org/10.4161/hv.23802.

[10]   Z. Hubálek, History of Arbovirus Research in the Czech Republic, Viruses 13 (2021) 2334. https://doi.org/10.3390/v13112334.

[11]   C. Chastel, A.J. Main, C. Guiguen, G. Le Lay, M.C. Quillien, J.Y. Monnat, J.C. Beaucournu, The isolation of Meaban virus, a new Flavivirus from the seabird tick Ornithodoros (Alectorobius) maritimus in France, Arch. Virol. 83 (1985) 129–140. https://doi.org/10.1007/BF01309911.

[12]   D.M. McLean, W.L. Donohue, Powassan virus: isolation of virus from a fatal case of encephalitis, Can. Med. Assoc. J. 80 (1959) 708–711.

[13]   S. Rebaudet, P. Parola, Epidemiology of relapsing fever borreliosis in Europe, FEMS Immunol. Med. Microbiol. 48 (2006) 11–15. https://doi.org/10.1111/j.1574-695X.2006.00104.x.

[14]   C. Anderson, C.A. Brissette, The Brilliance of Borrelia: Mechanisms of Host Immune Evasion by Lyme Disease-Causing Spirochetes, Pathogens 10 (2021) 281. https://doi.org/10.3390/pathogens10030281.

[15]   E.T. Stone, A.K. Pinto, T Cells in Tick-Borne Flavivirus Encephalitis: A Review of Current Paradigms in Protection and Disease Pathology, Viruses 15 (2023) 958. https://doi.org/10.3390/v15040958.

[16]   Y. Dong, G. Zhou, W. Cao, X. Xu, Y. Zhang, Z. Ji, J. Yang, J. Chen, M. Liu, Y. Fan, J. Kong, S. Wen, B. Li, P. Yue, A. Liu, F. Bao, Global seroprevalence and sociodemographic characteristics of *Borrelia burgdorferi sensu lato* in human populations: a systematic review and meta-analysis, BMJ Glob. Health 7 (2022) e007744. https://doi.org/10.1136/bmjgh-2021-007744.

[17] M.A. Wondim, P. Czupryna, S. Pancewicz, E. Kruszewska, M. Groth, A. Moniuszko-Malinowska,
Epidemiological Trends of Trans-Boundary Tick-Borne Encephalitis in Europe, 2000–2019,
Pathogens 11 (2022) 704. https://doi.org/10.3390/pathogens11060704.
[18] E. Chou, A. Minor, N.C. Cady, Quantitative multiplexed strategies for human Lyme disease
serological testing, Exp. Biol. Med. 246 (2021) 1388–1399.
https://doi.org/10.1177/15353702211003496.
[19] M. Gudowska-Sawczuk, B. Mroczko, Selected Biomarkers of Tick-Borne Encephalitis: A Review,
Int. J. Mol. Sci. 22 (2021) 10615. https://doi.org/10.3390/ijms221910615.
[20] M. Elbaz, A. Gadoth, D. Shepshelovich, D. Shasha, N. Rudoler, Y. Paran, Systematic Review and
Meta-analysis of Foodborne Tickborne Encephalitis, Europe, 1980–2021, Emerg. Infect. Dis. 28
(2022). https://doi.org/10.3201/eid2810.220498.
[21] E. Martello, E.L. Gillingham, R. Phalkey, C. Vardavas, K. Nikitara, T. Bakonyi, C.M. Gossner, J.
Leonardi-Bee, Systematic review on the non-vectorial transmission of Tick-borne encephalitis
virus (TBEv), Ticks Tick-Borne Dis. 13 (2022) 102028.
https://doi.org/10.1016/j.ttbdis.2022.102028.
[22] O.M. Zannou, A.S. Ouedraogo, A.S. Biguezoton, E. Abatih, M. Coral-Almeida, S. Farougou, K.P. Yao,
L. Lempereur, C. Saegerman, Models for Studying the Distribution of Ticks and Tick-Borne
Diseases in Animals: A Systematic Review and a Meta-Analysis with a Focus on Africa, Pathogens
10 (2021) 893. https://doi.org/10.3390/pathogens10070893.
[23] C. Voyiatzaki, S.I. Papailia, M.S. Venetikou, J. Pouris, M.E. Tsoumani, E.G. Papageorgiou, Climate
Changes Exacerbate the Spread of Ixodes ricinus and the Occurrence of Lyme Borreliosis and Tick-
Borne Encephalitis in Europe—How Climate Models Are Used as a Risk Assessment Approach for
Tick-Borne Diseases, Int. J. Environ. Res. Public. Health 19 (2022) 6516.
https://doi.org/10.3390/ijerph19116516.
[24] B. Defaye, S. Moutailler, V. Pasqualini, Y. Quilichini, A Systematic Review of the Distribution of
Tick-Borne Pathogens in Wild Animals and Their Ticks in the Mediterranean Rim between 2000
and 2021, Microorganisms 10 (2022) 1858. https://doi.org/10.3390/microorganisms10091858.
[25] B. Mubemba, M.M. Mburu, K. Changula, W. Muleya, L.C. Moonga, H.M. Chambaro, M. Kajihara, Y.
Qiu, Y. Orba, K. Hayashida, C.G. Sutcliffe, D.E. Norris, P.E. Thuma, P. Ndubani, S. Chitanga, H.
Sawa, A. Takada, E. Simulundu, Current knowledge of vector-borne zoonotic pathogens in Zambia:
A clarion call to scaling-up "One Health" research in the wake of emerging and re-emerging
infectious diseases, PLoS Negl. Trop. Dis. 16 (2022) e0010193.
https://doi.org/10.1371/journal.pntd.0010193.
[26] P.S. Pandit, M.M. Doyle, K.M. Smart, C.C.W. Young, G.W. Drape, C.K. Johnson, Predicting wildlife
reservoirs and global vulnerability to zoonotic Flaviviruses, Nat. Commun. 9 (2018) 5425.
https://doi.org/10.1038/s41467-018-07896-2.
[27] S. Hussain, A. Hussain, U. Aziz, B. Song, J. Zeb, D. George, J. Li, O. Sparagano, The Role of Ticks in
the Emergence of Borrelia burgdorferi as a Zoonotic Pathogen and Its Vector Control: A Global
Systemic Review, Microorganisms 9 (2021) 2412.
https://doi.org/10.3390/microorganisms9122412.
[28] K.M. Hansford, B.W. Wheeler, B. Tschirren, J.M. Medlock, Questing *Ixodes ricinus* ticks and
Borrelia spp. in urban green space across Europe: A review, Zoonoses Public Health 69 (2022)
153–166. https://doi.org/10.1111/zph.12913.
[29] M.J. Keeling, P. Rohani, Modeling Infectious Diseases in Humans and Animals, Princeton
University Press, 2011. https://doi.org/10.2307/j.ctvcm4gk0.
[30] M. Viana, R. Mancy, R. Biek, S. Cleaveland, P.C. Cross, J.O. Lloyd-Smith, D.T. Haydon, Assembling
evidence for identifying reservoirs of infection, Trends Ecol. Evol. 29 (2014) 270–279.
https://doi.org/10.1016/j.tree.2014.03.002.

[31] B.T. McClintock, J.D. Nichols, L.L. Bailey, D.I. MacKenzie, William.L. Kendall, A.B. Franklin, Seeking
a second opinion: uncertainty in disease ecology: Uncertainty in disease ecology, Ecol. Lett. 13
(2010) 659–674. https://doi.org/10.1111/j.1461-0248.2010.01472.x.
[32] K. LoGiudice, R.S. Ostfeld, K.A. Schmidt, F. Keesing, The ecology of infectious disease: Effects of
host diversity and community composition on Lyme disease risk, Proc. Natl. Acad. Sci. 100 (2003)
567–571. https://doi.org/10.1073/pnas.0233733100.
[33] D.M. McLean, L.W. MacPherson, S.J. Walker, G. Funk, Powassan virus: surveys of human and
animal sera, Am. J. Public Health Nation Health 50 (1960) 1539–1544.
[34] D.M. McLean, S.J. Walker, L.W. MacPherson, T.H. Scholten, K. Ronald, J.C. Wyllie, E.J. McQueen,
Powassan Virus: Investigations of Possible Natural Cycles of Infection, J. Infect. Dis. 109 (1961) 19–
23.
[35] D.M. McLean, E.J. McQueen, H.E. Petite, L.W. Macpherson, T.H. Scholten, K. Ronald, Powassan
Virus: Field Investigations in Northern Ontario, 1959 to 1961, Can. Med. Assoc. J. 86 (1962) 971–
974.
[36] D.M. Mclean, R.P. Larke, Powassan and Silverwater viruses: ecology of two Ontario arboviruses,
Can. Med. Assoc. J. 88 (1963) 182–185.
[37] D.M. Mclean, J.M. Best, S. Mahalingam, M.A. Chernesky, W.E. Wilson, Powassan virus: Summer
infection cycle, 1964, Can. Med. Assoc. J. 91 (1964) 1360–1362.
[38] D.M. McLean, P.A. Smith, S.E. Livingstone, W.E. Wilson, A.G. Wilson, Powassan virus: vernal
spread during 1965, Can. Med. Assoc. J. 94 (1966) 532–536.
[39] D.M. McLean, C. Cobb, S.E. Gooderham, C.A. Smart, A.G. Wilson, W.E. Wilson, Powassan virus:
persistence of virus activity during 1966, Can. Med. Assoc. J. 96 (1967) 660–664.
[40] D.M. McLean, S.R. Ladyman, K.W. Purvin-Good, Westward extension of Powassan virus
prevalence, Can. Med. Assoc. J. 98 (1968) 946–949.
[41] D.M. McLean, M.A. Chernesky, S.J. Chernesky, E.J. Goddard, S.R. Ladyman, R.R. Peers, K.W.
Purvin-Good, Arbovirus prevalence in the East Kootenay Region, 1968, Can. Med. Assoc. J. 100
(1969) 320–326.
[42] F. Frandsen, J. Bresciani, H.G. Hansen, Prevalence of antibodies to Borrelia burgdorferi in Danish
rodents, APMIS Acta Pathol. Microbiol. Immunol. Scand. 103 (1995) 247–253.
[43] B. Pisanu, J.-L. Chapuis, A. Dozières, F. Basset, V. Poux, G. Vourc'h, High prevalence of Borrelia
burgdorferi s.l. in the European red squirrel Sciurus vulgaris in France, Ticks Tick-Borne Dis. 5
(2014) 1–6. https://doi.org/10.1016/j.ttbdis.2013.07.007.
[44] A. Michelitsch, K. Wernike, C. Klaus, G. Dobler, M. Beer, Exploring the Reservoir Hosts of Tick-
Borne Encephalitis Virus, Viruses 11 (2019) 669. https://doi.org/10.3390/v11070669.
[45] G.M. Hacker, R.N. Brown, N. Fedorova, Y.A. Girard, M. Higley, B. Clueit, R.S. Lane, Spatial
clustering of Borrelia burgdorferi sensu lato within populations of Allen's chipmunks and dusky-
footed woodrats in northwestern California, PLOS ONE 13 (2018) e0195586.
https://doi.org/10.1371/journal.pone.0195586.
[46] S.I. Moore, M.L. Wilson, A. Spielman, S.R. Telford, T.N. Mather, Incompetence of Deer as
Reservoirs of the Lyme Disease Spirochete, Am. J. Trop. Med. Hyg. 39 (1988) 105–109.
https://doi.org/10.4269/ajtmh.1988.39.105.
[47] K. Kurtenbach, H.-S. Sewell, N.H. Ogden, S.E. Randolph, P.A. Nuttall, Serum Complement
Sensitivity as a Key Factor in Lyme Disease Ecology, Infect. Immun. 66 (1998) 1248–1251.
https://doi.org/10.1128/IAI.66.3.1248-1251.1998.
[48] J.D. Radolf, M.J. Caimano, B. Stevenson, L.T. Hu, Of ticks, mice and men: understanding the dual-
host lifestyle of Lyme disease spirochaetes, Nat. Rev. Microbiol. 10 (2012) 87–99.
https://doi.org/10.1038/nrmicro2714.

[49]   P. Kjellander, A.J.M. Hewison, O. Liberg, J.-M. Angibault, E. Bideau, B. Cargnelutti, Experimental
489   evidence for density-dependence of home-range size in roe deer (Capreolus capreolus L.): a
490   comparison of two long-term studies, Oecologia 139 (2004) 478–485.
491   https://doi.org/10.1007/s00442-004-1529-z.

492   [50]   W.D. Walter, T.S. Evans, D. Stainbrook, B.D. Wallingford, C.S. Rosenberry, D.R. Diefenbach,
493   Heterogeneity of a landscape influences size of home range in a North American cervid, Sci. Rep. 8
494   (2018) 14667. https://doi.org/10.1038/s41598-018-32937-7.

495   [51]   M. Holding, S.D. Dowall, J.M. Medlock, D.P. Carter, S.T. Pullan, J. Lewis, R. Vipond, M.S. Rocchi, M.
496   Baylis, R. Hewson, Tick-Borne Encephalitis Virus, United Kingdom, Emerg. Infect. Dis. 26 (2020)
497   90–96. https://doi.org/10.3201/eid2601.191085.

498   [52]   L. Gern, A. Estrada-Peña, F. Frandsen, J.S. Gray, T.G.T. Jaenson, F. Jongejan, O. Kahl, E. Korenberg,
499   R. Mehl, P.A. Nuttall, European Reservoir Hosts of Borrelia burgdorferi sensu lato, Zentralblatt Für
500   Bakteriol. 287 (1998) 196–204. https://doi.org/10.1016/S0934-8840(98)80121-7.

501   [53]   E.A. Newman, L. Eisen, R.J. Eisen, N. Fedorova, J.M. Hasty, C. Vaughn, R.S. Lane, Borrelia
502   burgdorferi Sensu Lato Spirochetes in Wild Birds in Northwestern California: Associations with
503   Ecological Factors, Bird Behavior and Tick Infestation, PLOS ONE 10 (2015) e0118146.
504   https://doi.org/10.1371/journal.pone.0118146.

505   [54]   T. Boulinier, S. Kada, A. Ponchon, M. Dupraz, M. Dietrich, A. Gamble, V. Bourret, O. Duriez, R.
506   Bazire, J. Tornos, T. Tveraa, T. Chambert, R. Garnier, K.D. McCoy, Migration, Prospecting,
507   Dispersal? What Host Movement Matters for Infectious Agent Circulation?, Integr. Comp. Biol. 56
508   (2016) 330–342. https://doi.org/10.1093/icb/icw015.

509   [55]   I. Humphery-Smith, D.H. Cybinski, K.A. Byrnes, T.D. St George, Seroepidemiology of arboviruses
510   among seabirds and island residents of the Great Barrier Reef and Coral Sea, Epidemiol. Infect.
511   107 (1991) 435–440. https://doi.org/10.1017/S0950268800049086.

512   [56]   K.L. Carpio, J.K. Thompson, S.G. Widen, J.K. Smith, T.L. Juelich, D.E. Clements, A.N. Freiberg, A.D.T.
513   Barrett, Differences in Genetic Diversity of Mammalian Tick-Borne Flaviviruses, Viruses 15 (2023)
514   281. https://doi.org/10.3390/v15020281.

515   [57]   N.J. Evans, K. Bown, D. Timofte, V.R. Simpson, R.J. Birtles, Fatal Borreliosis in Bat Caused by
516   Relapsing Fever Spirochete, United Kingdom, Emerg. Infect. Dis. 15 (2009) 1331–1333.
517   https://doi.org/10.3201/eid1508.090475.

518   [58]   C.J. Marinkelle, E.S. Grose, Species of Borrelia from a Colombian Bat (Natalus tumidirostris),
519   Nature 218 (1968) 487–487. https://doi.org/10.1038/218487a0.

520   [59]   A.C. Norte, I. Lopes De Carvalho, M.S. Núncio, P.M. Araújo, E. Matthysen, J. Albino Ramos, H.
521   Sprong, D. Heylen, Getting under the birds' skin: tissue tropism of Borrelia burgdorferi s.l. in
522   naturally and experimentally infected avian hosts, Microb. Ecol. 79 (2020) 756–769.
523   https://doi.org/10.1007/s00248-019-01442-3.

524   [60]   K.R. O'Keeffe, Z.J. Oppler, D. Brisson, Evolutionary ecology of Lyme Borrelia, Infect. Genet. Evol. 85
525   (2020) 104570. https://doi.org/10.1016/j.meegid.2020.104570.

526   [61]   J. Salat, M. Hunady, P. Schanilec, P. Strakova, M. Stefanik, P. Svoboda, L. Strelcova, J. Bojcukova,
527   M. Palus, D. Růžek, Experimental and Natural Infections of Tick-Borne Encephalitis Virus in Dogs,
528   Viruses 13 (2021) 2039. https://doi.org/10.3390/v13102039.

529   [62]   L. Jemeršić, D. Dežđek, D. Brnić, J. Prpić, Z. Janicki, T. Keros, B. Roić, A. Slavica, S. Terzić, D.
530   Konjević, R. Beck, Detection and genetic characterization of tick-borne encephalitis virus (TBEV)
531   derived from ticks removed from red foxes (Vulpes vulpes) and isolated from spleen samples of
532   red deer (Cervus elaphus) in Croatia, Ticks Tick-Borne Dis. 5 (2014) 7–13.
533   https://doi.org/10.1016/j.ttbdis.2012.11.016.

534   [63]   T. Csank, K. Bhide, E. Bencúrová, S. Dolinská, P. Drzewnioková, P. Major, Ľ. Korytár, E. Bocková, M.
535   Bhide, J. Pistl, Detection of West Nile virus and tick-borne encephalitis virus in birds in Slovakia,

using a universal primer set, Arch. Virol. 161 (2016) 1679–1683. https://doi.org/10.1007/s00705-
016-2828-5.
[64] H.J. Esser, S.M. Lim, A. De Vries, H. Sprong, D.J. Dekker, E.L. Pascoe, J.W. Bakker, V. Suin, E. Franz,
B.E.E. Martina, C.J.M. Koenraadt, Continued Circulation of Tick-Borne Encephalitis Virus Variants
and Detection of Novel Transmission Foci, the Netherlands, Emerg. Infect. Dis. 28 (2022) 2416–
2424. https://doi.org/10.3201/eid2812.220552.
[65] E. Tonteri, A.E. Jääskeläinen, T. Tikkakoski, L. Voutilainen, J. Niemimaa, H. Henttonen, A. Vaheri, O.
Vapalahti, Tick-borne Encephalitis Virus in Wild Rodents in Winter, Finland, 2008–2009, Emerg.
Infect. Dis. 17 (2011) 72–75. https://doi.org/10.3201/eid1701.100051.
[66] V.N. Bakhvalova, G.S. Chicherina, O.F. Potapova, V.V. Panov, V.V. Glupov, M.A. Potapov, S.J.
Seligman, O.V. Morozova, Tick-Borne Encephalitis Virus Diversity in Ixodid Ticks and Small
Mammals in South-Western Siberia, Russia, Vector-Borne Zoonotic Dis. 16 (2016) 541–549.
https://doi.org/10.1089/vbz.2015.1834.
[67] X. Dai, G. Shang, S. Lu, J. Yang, J. Xu, A new subtype of eastern tick-borne encephalitis virus
discovered in Qinghai-Tibet Plateau, China, Emerg. Microbes Infect. 7 (2018) 1–9.
https://doi.org/10.1038/s41426-018-0081-6.
[68] M. Bestehorn, S. Weigold, W.V. Kern, L. Chitimia-Dobler, U. Mackenstedt, G. Dobler, J.P. Borde,
Phylogenetics of tick-borne encephalitis virus in endemic foci in the upper Rhine region in France
and Germany, PLOS ONE 13 (2018) e0204790. https://doi.org/10.1371/journal.pone.0204790.
[69] H. Holzmann, Diagnosis of tick-borne encephalitis, Vaccine 21 (2003) S36–S40.
https://doi.org/10.1016/S0264-410X(02)00819-8.
[70] E. Tonteri, A. Kipar, L. Voutilainen, S. Vene, A. Vaheri, O. Vapalahti, Å. Lundkvist, The Three
Subtypes of Tick-Borne Encephalitis Virus Induce Encephalitis in a Natural Host, the Bank Vole
(Myodes glareolus), PLoS ONE 8 (2013) e81214. https://doi.org/10.1371/journal.pone.0081214.
[71] A.J. Kucharski, M. Kama, C.H. Watson, M. Aubry, S. Funk, A.D. Henderson, O.J. Brady, J.
Vanhomwegen, J.-C. Manuguerra, C.L. Lau, W.J. Edmunds, J. Aaskov, E.J. Nilles, V.-M. Cao-
Lormeau, S. Hué, M.L. Hibberd, Using paired serology and surveillance data to quantify dengue
transmission and control during a large outbreak in Fiji, eLife 7 (2018) e34848.
https://doi.org/10.7554/eLife.34848.
[72] T. Vilibic-Cavlek, T. Ferenc, M. Vujica Ferenc, M. Bogdanic, T. Potocnik-Hunjadi, D. Sabadi, V. Savic,
L. Barbic, V. Stevanovic, F. Monaco, E. Listes, G. Savini, Cross-Reactive Antibodies in Tick-Borne
Encephalitis: Case Report and Literature Review, Antibodies 11 (2022) 72.
https://doi.org/10.3390/antib11040072.
[73] I. García-Bocanegra, J. Paniagua, A.V. Gutiérrez-Guzmán, S. Lecollinet, M. Boadella, A. Arenas-
Montes, D. Cano-Terriza, S. Lowenski, C. Gortázar, U. Höfle, Spatio-temporal trends and risk
factors affecting West Nile virus and related flavivirus exposure in Spanish wild ruminants, BMC
Vet. Res. 12 (2016) 249. https://doi.org/10.1186/s12917-016-0876-4.
[74] T.S. Gritsun, V.A. Lashkevich, E.A. Gould, Tick-borne encephalitis, Antiviral Res. 57 (2003) 129–
146. https://doi.org/10.1016/S0166-3542(02)00206-1.
[75] A.J. Main, W.G. Downs, R.E. Shope, R.C. Wallis, Avian arboviruses of the Witless Bay Seabird
Sanctuary, Newfoundland, Canada, J. Wildl. Dis. 12 (1976) 182–194.
https://doi.org/10.7589/0090-3558-12.2.182.
[76] R. Garnier, R. Ramos, A. Sanz-Aguilar, M. Poisbleau, H. Weimerskirch, S. Burthe, J. Tornos, T.
Boulinier, Interpreting ELISA analyses from wild animal samples: Some recurrent issues and
solutions, Funct. Ecol. 31 (2017) 2255–2262. https://doi.org/10.1111/1365-2435.12942.
[77] L. Paillard, K.L. Jones, A.L. Evans, J. Berret, M. Jacquet, R. Lienhard, M. Bouzelboudjen, J.M.
Arnemo, J.E. Swenson, M.J. Voordouw, Serological signature of tick-borne pathogens in

Scandinavian brown bears over two decades, Parasit. Vectors 8 (2015) 398.
https://doi.org/10.1186/s13071-015-0967-2.

[78] A. Arnal, E. Gómez-Díaz, M. Cerdà-Cuéllar, S. Lecollinet, J. Pearce-Duvet, N. Busquets, I. García-
Bocanegra, N. Pagès, M. Vittecoq, A. Hammouda, B. Samraoui, R. Garnier, R. Ramos, S. Selmi, J.
González-Solís, E. Jourdain, T. Boulinier, Circulation of a Meaban-Like Virus in Yellow-Legged Gulls
and Seabird Ticks in the Western Mediterranean Basin, PLoS ONE 9 (2014) e89601.
https://doi.org/10.1371/journal.pone.0089601.

[79] E. Jurado-Tarifa, S. Napp, S. Lecollinet, A. Arenas, C. Beck, M. Cerdà-Cuéllar, M. Fernández-
Morente, I. García-Bocanegra, Monitoring of West Nile virus, Usutu virus and Meaban virus in
waterfowl used as decoys and wild raptors in southern Spain, Comp. Immunol. Microbiol. Infect.
Dis. 49 (2016) 58–64. https://doi.org/10.1016/j.cimid.2016.10.001.

[80] F.H. Weissbach, H.H. Hirsch, Comparison of Two Commercial Tick-Borne Encephalitis Virus IgG
Enzyme-Linked Immunosorbent Assays, Clin. Vaccine Immunol. 22 (2015) 754–760.
https://doi.org/10.1128/CVI.00096-15.

[81] B. Ytrehus, K. Vainio, S.G. Dudman, J. Gilray, K. Willoughby, Tick-Borne Encephalitis Virus and
Louping-Ill Virus May Co-Circulate in Southern Norway, Vector-Borne Zoonotic Dis. 13 (2013) 762–
768. https://doi.org/10.1089/vbz.2012.1023.

[82] L. Bournez, G. Umhang, E. Faure, J.-M. Boucher, F. Boué, E. Jourdain, M. Sarasa, F. Llorente, M.A.
Jiménez-Clavero, S. Moutailler, S.A. Lacour, S. Lecollinet, C. Beck, Exposure of Wild Ungulates to
the Usutu and Tick-Borne Encephalitis Viruses in France in 2009–2014: Evidence of Undetected
Flavivirus Circulation a Decade Ago, Viruses 12 (2019) 10. https://doi.org/10.3390/v12010010.

[83] D. Fish, A.G. Hoen, J.I. Tsao, A.G. Barbour, M.A. Diuk-Wasser, J. Bunikis, B. Travinsky, Niche
Partitioning of Borrelia burgdorferi and Borrelia miyamotoi in the Same Tick Vector and
Mammalian Reservoir Species, Am. J. Trop. Med. Hyg. 81 (2009) 1120–1131.
https://doi.org/10.4269/ajtmh.2009.09-0208.

[84] X.-A. Zhang, F. Tian, Y. Li, X.-L. Zhang, B.-G. Jiang, B.-C. Liu, J.-T. Zhang, S. Tian, H. Ding, S. Li, H. Li,
L.-Q. Fang, W. Liu, Molecular detection and identification of relapsing fever *Borrelia* in ticks and
wild small mammals in China, Emerg. Microbes Infect. 11 (2022) 2632–2635.
https://doi.org/10.1080/22221751.2022.2134054.

[85] M.C. Scott, M.E. Rosen, S.A. Hamer, E. Baker, H. Edwards, C. Crowder, J.I. Tsao, G.J. Hickling, High-
Prevalence Borrelia miyamotoi Infection Among Wild Turkeys (Meleagris gallopavo) in Tennessee,
J. Med. Entomol. 47 (2010) 1238–1242. https://doi.org/10.1603/ME10075.

[86] J.C. Burtis, J.B. Yavitt, T.J. Fahey, R.S. Ostfeld, Ticks as Soil-Dwelling Arthropods: An Intersection
Between Disease and Soil Ecology, J. Med. Entomol. 56 (2019) 1555–1564.
https://doi.org/10.1093/jme/tjz116.

[87] S. Szekeres, E.C. Coipan, K. Rigó, G. Majoros, S. Jahfari, H. Sprong, G. Földvári, Eco-epidemiology of
Borrelia miyamotoi and Lyme borreliosis spirochetes in a popular hunting and recreational forest
area in Hungary, Parasit. Vectors 8 (2015) 309. https://doi.org/10.1186/s13071-015-0922-2.

[88] A. Obiegala, N. Król, C. Oltersdorf, J. Nader, M. Pfeffer, The enzootic life-cycle of Borrelia
burgdorferi (sensu lato) and tick-borne rickettsiae: an epidemiological study on wild-living small
mammals and their ticks from Saxony, Germany, Parasit. Vectors 10 (2017) 115.
https://doi.org/10.1186/s13071-017-2053-4.

[89] C. Chastel, C. Guiguen, G. Le Lay, J.Y. Monnat, E. Hardy, G. Kerdraon, J.C. Beaucournu, [Arbovirus
serological survey among marine and non-marine birds of Brittany], Bull. Soc. Pathol. Exot. Filiales
78 (1985) 594–605.

[90] M.K. Krzysiak, K. Anusz, A. Konieczny, J. Rola, J. Salat, P. Strakova, W. Olech, M. Larska, The
European bison (Bison bonasus) as an indicatory species for the circulation of tick-borne

encephalitis virus (TBEV) in natural foci in Poland, Ticks Tick-Borne Dis. 12 (2021) 101799.
https://doi.org/10.1016/j.ttbdis.2021.101799.

[91] M.R. Holbrook, Kyasanur forest disease, Antiviral Res. 96 (2012) 353–362.
https://doi.org/10.1016/j.antiviral.2012.10.005.

[92] V.N. Bakhvalova, A.K. Dobrotvorsky, V.V. Panov, V.A. Matveeva, S.E. Tkachev, O.V. Morozova,
Natural Tick-Borne Encephalitis Virus Infection among Wild Small Mammals in the Southeastern
Part of Western Siberia, Russia, Vector-Borne Zoonotic Dis. 6 (2006) 32–41.
https://doi.org/10.1089/vbz.2006.6.32.

[93] V. Ollivier, R. Choquet, A. Gamble, M. Bastien, B. Combes, E. Gilot-Fromont, M. Pellerin, J.
Gaillard, J. Lemaître, H. Verheyden, T. Boulinier, Temporal dynamics of antibody level against
Lyme disease bacteria in roe deer: Tale of a sentinel?, Ecol. Evol. 13 (2023) e10414.
https://doi.org/10.1002/ece3.10414.

[94] S. Lambert, E. Gilot-Fromont, C. Toïgo, P. Marchand, E. Petit, S. Rossi, A. Thébault, Combining
seroprevalence and capture-mark-recapture data to estimate the force of infection of brucellosis
in a managed population of Alpine ibex, Epidemics 38 (2022) 100542.
https://doi.org/10.1016/j.epidem.2022.100542.

[95] A.R. Sweeny, G.F. Albery, Exposure and susceptibility: The Twin Pillars of infection, Funct. Ecol. 36
(2022) 1713–1726. https://doi.org/10.1111/1365-2435.14065.

[96] N.G. Yoccoz, J.D. Nichols, T. Boulinier, Monitoring of biological diversity in space and time, Trends
Ecol. Evol. 16 (2001) 446–453. https://doi.org/10.1016/S0169-5347(01)02205-4.

[97] R.S. Ostfeld, T. Levi, A.E. Jolles, L.B. Martin, P.R. Hosseini, F. Keesing, Life History and Demographic
Drivers of Reservoir Competence for Three Tick-Borne Zoonotic Pathogens, PLoS ONE 9 (2014)
e107387. https://doi.org/10.1371/journal.pone.0107387.

[98] J.G. Donahue, J. Piesman, A. Spielman, Reservoir Competence of White-Footed Mice for Lyme
Disease Spirochetes, Am. J. Trop. Med. Hyg. 36 (1987) 92–96.
https://doi.org/10.4269/ajtmh.1987.36.92.

[99] J.L. Panetta, R. Šíma, N.E.D. Calvani, O. Hajdušek, S. Chandra, J. Panuccio, J. Šlapeta, Reptile-
associated Borrelia species in the goanna tick (Bothriocroton undatum) from Sydney, Australia,
Parasit. Vectors 10 (2017) 616. https://doi.org/10.1186/s13071-017-2579-5.

[100] C. Larsson, M. Andersson, B.P. Guo, A. Nordstrand, I. Hägerstrand, S. Carlsson, S. Bergström,
Complications of Pregnancy and Transplacental Transmission of Relapsing-Fever Borreliosis, J.
Infect. Dis. 194 (2006) 1367–1374. https://doi.org/10.1086/508425.
662

Acknowledgements

This work was funded by Agence Nationale de la Recherche (ANR) under the ANRJCJC MoZArt project, grant number ANR-22-CE35-0003. Support is also acknowledged from ANR ECOPATHS project, grant number ANR-21-CE35-0016, and from IPEV ECOPATH-1151 project.

Author information

Authors and affiliations

Sorbonne Université, INSERM, Institut Pierre Louis d'Epidémiologie et de Santé Publique, UMRS 1136, Paris, France

Raphaëlle Métras, Armelle Poisson

Centre d'Ecologie Fonctionnelle et Evolutive, UMR 5175, CNRS, Université de Montpellier, IRD, EPHE, Montpellier, France

Thierry Boulinier, Armelle Poisson

676
Biogéosciences, UMR 6282, CNRS, Université Bourgogne, EPHE, Dijon, France

Bruno Faivre

ANSES, INRAE, Ecole Nationale Vétérinaire d'Alfort, UMR Virologie, Laboratoire de Santé Animale,
Maisons-Alfort, France

Gaëlle Gonzalez, Camille V. Migné

ANSES, INRAE, Ecole Nationale Vétérinaire d'Alfort, UMR BIPAR, Laboratoire de Santé Animale,
Maisons-Alfort, France

Sara Moutailler

Laboratoire de la Rage et de la Faune Sauvage, ANSES, Malzéville, France

Laure Bournez

Contributions

691 AP, TB and RM conceived the paper. AP conducted the analysis and designed the figures. AP and RM wrote
692 the paper. TB, GG, CM, LB, BF and SM revised the paper. All authors approved the final version of the
693 paper.

Competing interests

695 Authors declare no competing interests.

Corresponding authors

697 Correspondence to Armelle Poisson (armelle.poisson@laposte.net) and Raphaëlle Métras
698 (raphaelle.metras@inserm.fr).