Tick-borne zoonotic flaviviruses and *Borrelia* infections in wildlife hosts: what have field studies contributed?

Armelle Poisson[1], Thierry Boulinier[2], Laure Bournez[3], Gaëlle Gonzalez[4], Camille V. Migné[4], Sara Moutailler[5], Bruno Faivre[6], Raphaëlle Métras[1]

[1]Sorbonne Université, INSERM, Institut Pierre Louis d'Epidémiologie et de Santé Publique (IPLESP, UMRS 1136), Paris, France.

[2]Centre d'Ecologie Fonctionnelle et Evolutive, UMR 5175, CNRS, Université de Montpellier, IRD, EPHE, Montpellier, France.

[3]Laboratoire de la Rage et de la Faune Sauvage, ANSES, Malzéville, France.

[4]ANSES, INRAE, Ecole Nationale Vétérinaire d'Alfort, UMR Virologie, Laboratoire de Santé Animale, Maisons-Alfort, France.

[5]ANSES, INRAE, Ecole Nationale Vétérinaire d'Alfort, UMR BIPAR, Laboratoire de Santé Animale, Maisons-Alfort, France.

[6]Biogéosciences, UMR 6282, CNRS, Université de Bourgogne, EPHE, Dijon, France.

Abstract

Tick-borne flaviviruses and *Borrelia* spp. are globally spread pathogens of zoonotic potential which are maintained by a transmission cycle at the interface between ticks and vertebrate hosts, mainly wild animals. We reviewed how those infections have been studied in wildlife species in the field to discuss how collected data provided relevant epidemiological information and identify needs for further studies. The literature was screened for populational studies on direct or antibody detection for tick-borne *Borrelia* spp. and flaviviruses in animals, in the wild. Overall, *Borrelia* spp. were more studied (73% of case studies, representing 297 host species) than flaviviruses (27% of case studies, representing 114 host species). Studies on both *Borrelia* spp. and flaviviruses focused mainly on the same species (namely bank vole and yellow-necked mouse). Species with potential to act as reservoir hosts were overlooked, notably passerine birds. Most studies were order-specific and cross-sectional. We highlight the necessity of collecting both demographics and infection data in wildlife studies, and to consider communities of species, to better estimate zoonotic risk potential.

Keywords

Tick-borne diseases, wildlife, *Borrelia*, flavivirus, reservoir, sentinel, host.

Introduction

Vector-borne and zoonotic infections are emerging threats to public health[1]. Tick-borne pathogens (TBP) are maintained by multi-hosts and complex epidemiological cycles at the interface between ticks and animal hosts[2]. Transmission to humans occur on specific occasions, and understanding and quantifying it in a One Health framework is necessary to prevent and control disease. Persistence of tick-borne and zoonotic pathogens is allowed at the interface between the tick vector and a large variety of animal hosts, allowing ticks to feed and complete their life cycles. In those zoonotic tick-host systems, the early tick stages (larvae and nymph) usually feed on small mammals and birds, while mature tick stages mainly feed on larger animals (such as deer, wild boar, cattle)[3].

Beyond their feeding role for the tick, animal species can act as pathogen reservoir or sentinel hosts. Infection and exposure data from both types of hosts give precious information on the dynamics of the studied TBP[4]. Whilst reservoir hosts allow the pathogen to multiply and transmit it further, sentinel species are characterized by their ability to reflect an underlying epidemiological phenomenon and by being of easy access[5]. For instance, for zoonotic *Toxoplasma gondii*, wild red and Arctic foxes have been suggested as sentinels for human and animal toxoplasma risk in Northern Canada, based on serology and direct detection of the pathogen on carcasses[6]. Also, animal ecology can drive variations in risk of transmission to human[7]. Therefore, because understanding pathogens' transmission and distribution in wildlife is prerequisite to addressing the risk of transmission to humans, quantitively tracking infection and exposure to pathogens in those hosts is necessary.

Tick-borne *Borreliaceae* and flaviviruses are common tick-borne pathogens of zoonotic potential which are the most prevalent in the temperate regions of the world[8] and are thus of interest for both human and veterinary public health. Tick-borne encephalitis virus (TBEV), causative agent for a disease known from the 18[th] century, has first been described in Austria in 1931[9] and then isolated in Far Eastern Russia in 1937[10]. It is the most common tick-borne Flavivirus (TBFV) in Europe, with several endemic foci in Europe and Asia and growing number of cases in human populations[11,12]. Its ecology and epidemiology have early been studied in Europe, especially in Czech Republic[13]. Other TBFV have later been isolated from wild animals and their ticks such as Meaban virus (MEAV) in Brittany, France[14] or from human cases such as Powassan virus (POWV) in Powassan town, Canada[15]. Vectors for TBFV are hard ticks *Ixodes ricinus* in Western and Central Europe and *I. persulcatus* in Eastern Europe as well as local tick species in other parts of the world, such as *I. scapularis* and *I. cookei* in North America[16–18]. Similarly transmitted bacterial genospecies from the *Borrelia burgdorferi* sensu lato (Bbsl) complex are the causative agents for Lyme disease, one of the most common TBD in the Northern Hemisphere. Other tick-borne *Borrelia* spp. outside of this complex are identified to cause relapsing fever in humans[19,20].

Recent reviews on tick-borne *Borrelia* spp. and TBFV have focused on getting information on the biology of infection in vertebrate hosts[21–28], reporting prevalence and clinical cases in human[29–33] and discussing diagnostic methodologies[34–37], the role of non-vector transmissions for TBFV[38–40], or on the importance of modelling TBP regarding climate change consequences[41,42]. Some reviews have focused on wildlife hosts for TBP but only at a national or continental scale[43–46], or on reservoirs associated with the flavivirus genus in general, with little focus on TBFV[47]. Finally, a lot of research effort has focused on studying the infectious agents in the tick vectors[48–51].

Despite their major importance in transmission, no study has so far reviewed and analysed the body of evidence in wildlife hosts from field data at a global scale. Yet, data on the level of infection or exposure

82 in sampled free ranging populations are critical to model the dynamics of infectious diseases[52]. Strong
83 inference about the transmission processes underlying those dynamics often require a combination of
84 approaches, notably experiments to ascertain the role of reservoirs[53], but quantitative data on levels of
85 infection and exposure of potential hosts are a prerequisite in most cases. When it comes to wildlife
86 species, a strong heterogeneity in the type of data gathered is nevertheless expected, from local cross-
87 sectional sampling of particular host species, to broad spatial surveys or surveillance data over series of
88 years. Estimating epidemiological parameters may also require to account for potential biases in detection
89 probabilities and host population parameters[54,55], which may require specific study designs.

90 The aim of this review was thus to investigate how tick-borne *Borrelia* spp. and flaviviruses infection
91 burden has been studied in free-ranging wildlife species (that is, excluding the tick vector) to discuss how
92 collected data provided relevant epidemiological information and to identify perspectives and needs for
93 further studies. We predicted that a strong heterogeneity would appear in the types of hosts investigated,
94 with a high number of studies conducted on common species sharing landscape use with human in the
95 Northern Hemisphere. We expected few studies including whole communities of local potential hosts and
96 studies providing estimates of TBP burden to allow exploring host, temporal and spatial variation while
97 accounting for uncertainty. We anticipated that monitoring of wildlife diseases would have been difficult
98 and that knowledge of what is precisely measured would have been uncertain.

Studied hosts and their potential role in transmission cycle

100 It was possible to reconstruct 947 "cases studies" (unique combinations of species-pathogen-area-study
101 period) studying overall 347 host species, from the 314 articles matching the selection criteria. Detailed
102 metrics on the results of the literature search are available in SuppInfo1. Figure 1 links the order of tested
103 species and the investigated pathogen. Species from Rodentia order were the more studied for both
104 *Borrelia* spp. and TBFV, followed by Artiodactyla order for TBFV and Passeriformes for *Borrelia* spp. (Figure
105 1).

Historical papers shaped research on Rodents as reservoir hosts

107 In our review, we identified that Rodentia was the most investigated order with more than half (363/689)
108 of the *Borrelia* cases and more than a third (91/258) of the TBFV study cases. The white-footed mouse
109 (*Peromyscus leucopus*) has been regarded as the most common reservoir host in wildlife in Northern
110 America for tick-borne *Borrelia* spp. at least since the eighties[56,57], and therefore was investigated in the
111 first published studies from Northern America. Concerning TBFV, bank vole (*Myodes glareolus*) and yellow-
112 necked mouse (*Apodemus flavicollis*) were recognized as reservoirs for TBEV in Europe[58]. Historical aspect
113 is important to explain realized works. Taking rodents as example, studies included first the *Peromyscus*
114 family, only present in Northern America, and secondly the *Microtus* and *Mus* families. Even earlier, in the
115 sixties in Canada, a series of studies from the same authors detected POWV in *Tamias*, *Urocitellus* and
116 *Callospermophilus* chipmunks and *Tamiasciurus* and *Sciurus* squirrels, with an average seroprevalence of
117 15%[59–67]. Groundhogs (*Marmota monax*) sampled in the same studies, reported the highest prevalence of
118 anti-POWV antibodies (around 43%)[63–65]. Publications investigated as well striped skunk (*Mephitis*
119 *mephitis*) and snowshoe hare (*Lepus americanus*) with respectively 41% (121/299) and 11% (143/1301) of
120 seropositive animals[64,66]. Rodents as squirrels and chipmunks were then presented as primary reservoirs
121 for POWV on the basis of these studies. Several studies on the eco-epidemiology of TBFV were parallelly
122 conducted in the USSR but were not published in English (see limitations in SuppInfo1). Rodents were later
123 sampled on other continents and TBFV detections were several times successful in *Apodemus*, *Microtus*,

Myodes families in Europe, and in these families as well as in *Micromys*, *Mus* and *Sciurus* families for
Borrelia. In addition, Bbsl has been detected in *Tamias*, *Tamiasciurus* and *Neotoma* spp.. Furthermore,
Gerbillus, *Mastomys*, *Praomys* and *Apodemus*, *Cricetulus*, *Eothenomys*, *Meriones*, *Mus*, *Niviventer* and
Rattus spp. bore relapsing fever causative *Borrelia spp.* respectively in Africa and in Asia. It is notable that
Eastern grey squirrels (*Sciurus carolinensis*) similarly carried *Borrelia* DNA, both in their native countries
and in Europe where they are invasive, as raccoon (*Procyon lotor*) and raccoon dog (*Nyctereutes*
procyonoides) do.

Overall, in all studied areas, rodent species evidenced similar levels and ranges of pathogen or serological
detection. For case studies testing over 50 individuals, the median proportion of pathogen detection was
lower than the proportion of seropositivity, yet both of these values were higher for *Borrelia* than TBFV.
Specifically, the median proportion of *Borrelia* detection was 5.30% (range 0.2%-76%) and seropositivity
was three times higher (median at 15%, range 0.7%-89%), whilst the median value for TBFV detection was
1.8% (range 0.8%-62%) with seropositivity median value at 4.95% (range 0.4 - 44%). Variations were
remarked along study sites and periods, regardless of taxonomy. Beyond that such TBP might be
widespread, this could be explained by the ecology broadly shared by rodents, living close to the ground
where foraging infected ticks can be encountered[68], and therefore being similarly exposed. Variations in
detection rate across studies were observed, notably for a same species, such as for Bbsl prevalence in
yellow-necked mouse tissues varying from a proportion of 5.8% in Hungary in 2014-2015[69] to 23% in
Germany in 2012-2014[70]. No studies looking at Kyanasur Forest Disease virus (KFDV), MEAV, Saumarez
Reef Encephalitis virus (SREV), Tyuleni virus (TYUV) and Yamaguchi virus (YV) in rodents were recorded.
Whilst KFDV and YV have terrestrial hosts[71], MEAV, SREV, TYUV are known to be seabird viruses.

Deer, broadly studied as sentinel hosts

Other important investigated hosts were Artiodactyls. This second most important order by the number
of tested individuals was mainly represented by the *Cervidae* family as in terms of number of tested
animals (82%, 26,090/31,862) and in terms of number of species investigated (56%, 9/16) (Figure 2B and
2D). The first fieldwork setting targeting *Borrelia* in deer identified in this review dates back from 1971.
Deer are also considered as incompetent reservoir for *Borrelia* spp. by some authors because of
borreliacidal properties of their sera[72]. Yet, studies showed that the complement borreliacidal activity
depends on the strain and also exists on other *Borrelia* strains for other vertebrates including rodents[73,74].
Despite their poor competence as reservoir, deer are frequently met all across the world and can therefore
be infected by feeding adult ticks they commonly host[75]. *Cervidae* was widespread in this review, with
locally present species from America, Europe and Asia seeming to show similar level of TBP contact (with
an average seroprevalence of 18%). For *Borrelia*, average proportion was of 15.6% and average
seroprevalence of 27.3%, and were respectively of 1.1% and 14.4% for TBFV. As such, deer could help
identifying endemic or emerging risk areas and give information on underlying processes. In addition,
monitoring displayed restraint home range sizes of no more than 10km² among the different deer
species[76–79], which allowed studying the infection process at a rather small spatial scale. Moreover,
surveillance activities and pathogen detection may be facilitated through deer hunting activities, since
such reviewed studies allowed planning widespread sampling program for reasonable costs and working
effort on a yearly or at least regular basis[80–83]. Furthermore, because of that, deer were repeatedly
considered as potential sentinels of tick-borne diseases in the wild. Other game Artiodactyls identified in
this review such as wild boar (*Sus scrofa*) offer the same kind of possibilities for such sentinel surveying
approaches[84].

Birds, reservoir hosts, have been overlooked

Birds were also reported in the literature as reservoirs[85]. Contrary to rodents, birds may have been overlooked. In our review, the average seroprevalence of anti-TBFV in birds was 35%, corresponding to an overall 24% (783/3273) of anti-TBFV positive individuals all cases taken together (Figure 2C). We hypothesized that the limited availability of data on birds might be due to the logistical difficulties related to blood sampling in the wild (logistical difficulties, training and legal requirements, ethical concerns for endangered species). However, it is notable that antibodies against Gadgets Gully and Saumarez Reef viruses were extensively studied in a collection of Australian seabirds' sera[86]. This constitutes a unique occurrence of rare TBFV in the reviewed articles, but sadly no clarification was made on which species of the collection retrieved seropositivity.

Although birds represented about 19% (129/689) of the *Borrelia* cases, they originated from only 12% (28/228) of the selected publications. In those, *Borrelia* spp. presence was often screened in a banding study configuration. For instance, Newman *et al.* caught birds on the campus of the University of California in 2003 and 2004[87]. Their wide-range study allowed to detect Bbsl in 23 species for which at least two individuals were caught and one positive, out of 53 caught species. Mistnetting increases the possibility of capturing more birds and therefore more potential reservoirs, and underlines the interest of also surveying birds. However, this kind of study design is mostly favourable for Passeriformes, and passerines were the most studied avian order to get insight on the potential of reservoir for TBP they have. In addition, for migratory species or those exploiting very large areas during their life cycle, the infection status might not always reflect the underlying local infection processes. However, combined with movement information, they can unravel critical information on pathogen dissemination over long distances and between various ecosystems[88,89].

Least studied species with recent growing interest

Other vertebrate species of interest have been less studied. Recently however new species of tick-borne *Borrelia* spp. have been sought for and encountered in animals from other orders. For example, although quite unusual, Squamates as host for *Borrelia* were investigated in America, Africa and Europe, with 18 species associated with either *B. burgdorferi* sensu lato or *B. lusitaniae*, the latter being also found in rodents in Mediterranean Europe[90,91]. Primates interestingly were found positive to KFDV antibody in India, a TBFV member of the mammal group close to POWV and DTV[92]. In the UK, an isolated clinical case of lethal borreliosis was found in a *Pipistrellus* spp. bat[93]. Neotropical bats have been identified to carry particular strains of relapsing fever *Borrelia* spp. and Bbsl in Central and Southern America. Studies in Central America have been conducted as a follow-back of a paper published in 1968 by Marinkelle & Grose[94]. Actually, some areas are currently investigated more in depth or monitored. For instance, Columbian cave mentioned above has been resampled for neotropical bats, and Tribeč mountain region in Czech Republic has been screened for TBEV since the seventies and a research presence on TBP is still maintained there.

Infection data in host species

The average *Borrelia* spp. proportion of detection, all species confounded, tended to uniformize between 15% and 30% as the number of tested individuals increased (Figure 3A). For TBFV, average prevalence ranged from 8% for MEAV to 51% for TYUV (Figure 3B).

For *Borrelia*, 297 wild animal species (including mammals, birds and lizards) were investigated. Pathogen detection proportions in studies with over 50 tested individuals ranged from 0.1% for *B. lusitaniae* in the Eastern grey squirrel *Sciurus carolinensis* to 76% for Bbsl in the white-footed mouse *Peromyscus leucopus*[95], and from 0.2% for relapsing fever causing species in the house mouse *Mus musculus*[96] to 50% for *B. miyamotoi* in the wild turkey *Meleagris gallopavo*[97]. Mammals were more studied than birds, and for those mammalian species with over 100 specimens tested, Artiodactyls, Rodents and Carnivores showed level of prevalence between 20 and 25%. Chiropters, little studied, presented both seropositivity (36%) and pathogen prevalence (25%) of rather high level (Figure 2A). In birds, Columbiformes, Passeriformes, Charadriiforms and Galliformes were the most studied in terms of number of tested individuals, with Charadriiforms and Passeriformes reaching 29% of seroprevalence (Figure 2B).

TBFV or antibodies were searched for and detected in 114 terrestrial mammal and bird species (Figure 2C and 2D). Serological detection proportions in studies with more than 50 tested individuals ranged from 0.07% for MEAV in the red deer *Cervus elaphus*[98] to 68% for TYUV/MEAV/SREV-like in the European herring gull *Larus argentatus*[99], and for mammalian TBFV, from 0,2% for TBEV/LIV in the roe deer *Capreolus capreolus*[100] to 63% for TBEV in the European bison *Bison bonasus*[101]. Most studied avian orders on TBFV were Passeriformes, Charadriiforms, Galliformes and Procellariforms, with highest percentage of seropositivity for Passeriformes (about 40%), followed by Charadriiforms (27%) and Galliformes (21%) (Figure 2C and 2D) when Procellariforms tended to have less seropositive or viremic individuals. In mammals, Eulipothyphla were well-studied, only by PCR techniques, and showed high rates of viremia (of about 50%). Rodents, Artiodactyls, Lagomorphs were also well investigated but had lower seropositivity rates. For Pelecaniformes, 80% of European shag *Phalacrocorax aristotelis* had antibodies to TYUV, but only 8 birds were tested, and on a single location in the eighties[99].

Diagnostic tests: current versus past infections

Direct and indirect detection methods were at different use according to the type of pathogens. Whilst most Borrelia studies (71%) looked for the genome of the pathogen and used molecular techniques to identify an ongoing infection, 89% of flaviviruses studies used serological detection methods based on antibody detection. Long-lasting in the organisms *Borrelia* spp., were often sought for and found by PCR (60% of case studies overall). Skin and blood were materials used for spirochetes PCR detection, allowing positive samples to be sequenced and species identified, although the use of blood to directly detect *Borrelia* is discussed[102]. The determination of Borrelia subspecies was used in understanding the role of hosts in disease cycles[103]. Different *Borrelia* spp. have been preferentially detected in different hosts, such as *B. lusitaniae* which was the only Bbsl genospecies found in lizards in Europe[104–107].

In contrast, the short viremia noted after flaviviruses infection as demonstrated by experimental infections[108,109] makes viral detection challenging, especially in wildlife. Flaviviruses detection by PCR was therefore rarely used (only 11% of TBFV studies used PCR), but allowed sequencing of TBEV subtypes that clustered with TBEV-Eur (European) in Croatia, Slovakia and in the Netherlands[110–112], TBEV-Eur and TBEV-Sib (Siberian) in Finland[113], TBEV-Eur, TBEV-Sib and TBEV-FE (Far Eastern) in Russia[114] and TBEV-Sib in China[115]. Bakhvalova *et al.* evidenced co-infection of several subtypes in a same individual[114]. Dai *et al.* called the strain they studied in China TBEV-Him (Himalayan)[116]. Whilst identifying precise strains will help understanding further the ecology and dynamics of pathogens, it can also assist disease surveillance, with phylogenetics providing information on circulation and evolution[117]. In addition, viral identification is also an asset for public health management. For example, TBEV, a burden for human health, could be

sometimes mistaken for Looping Ill virus (LIV), causing severe conditions mainly in sheep[82]. Bai *et al.*, were
led to reconsider high values of TBEV seroprevalence because of the discovery of Karshi virus' circulation
in China[118].

Flaviviruses infections induce an immune response with neutralizing antibodies that could be detected in
blood samples lifelong in human[119] and up to 168 days after experimental infection in bank voles[120]. The
use of serological approaches was preferred through ELISA techniques (89% of flaviviruses cases, and 38%
of overall cases). Serological information, in combination with metadata (such as host age) can provide
precious information on pathogen transmission[121–123]. Yet, cross-reactivity in wide-range immunoassays
may hamper the interpretation of data[124]. For instance, MEAV antibodies were detected in *Cervidae*[98]. This
detection is particularly surprising because MEAV is part of the Seabird TBFV group in phylogeny[125]. As
unusual, antibodies against LIV have been identified in seabirds *Fratercula arctica* and *Hydrobates*
leucorhous[126]. Both of these identifications however might be misclassification or cross-reactions, which is
known as a possible weakness of serological studies[127]. Other issue with immunoassays is the choice of
cut-off in data analysis, which is generally reproduced from the manufacturer. However some authors
proposed other methodologies for cut-off determination for ELISA results, such as mathematical fit on
observed data[128]. Paillard *et al.* even chose to present raw ELISA results as a signal of contact with TBFV
and therefore not to indicate cut-off, nor prevalence[129].

Publications about TBFV outside of TBEV and POWV were marginal and specific detection for these TBFV
antibodies was not always commercially available. Arnal *et al.* found MEAV-seropositivity in *Larus*
michaellis michaellis egg yolks in France using flavivirus-ELISA cross-reactivity properties, with the ID-Vet
ID Screen® kit designed for the mosquito-borne flavivirus West Nile virus (WNV)[130]. Similarly, through both
COMPAC® WNV-ELISA and an exclusion via serum neutralization tests (SNT), Jurado-Tarifa *et al.*
investigated flaviviruses in Spanish birds of prey and detected evidence of exposure to flaviviruses
different from the mosquito-borne WNV and Usutu (USUV) viruses and from MEAV in *Circus pygargus*,
Falco tinnunculus, *Hieraaetus pennatus* and *Tyto alba*[131]. This relies on lack of specificity of commercially
available anti-WNV antibodies detection kits. Some commercial TBEV ELISA have also been compared for
their reliability according to the TBEV-strain used and for the at-risk cross-reactivity[132]. In the UK, Holding
et al. recorded similar LIV and TBEV seropositivity results in cervids[82] although TBEV was though not to be
present on the British Isles[133]. Holding et *al.* used the FSME/TBE All-Species Progen® ELISA kit and a
hemagglutination assay specific to LIV. For confirmation, Holding *et al.* sequenced TBEV in a tick itself.
Ytrehus *et al.* also showed presence of both LIV and TBEV antibodies in Southern Norway in *Alces alces*
and *Capreolus capreolus* using ELISA kits for TBEV and hemagglutination for LIV as well as SNT for
verification[134]. However, their results appeared differentiable for the two pathogens. On the opposite,
Bournez *et al.* chose to deliver their final TBEV micro-neutralization test serology results under the
TBEV/LIV label, after a first antibodies screening in *Capreolus capreolus* and *Sus scrofa* with the ID-Vet ID
Screen® WNV-ELISA[100]. TBEV and LIV as well as the Turkish Sheep Encephalomyelitis virus (TSEV) and the
Spanish Sheep Encephalomyelitis virus (SSEV) belong to the TBEV serocomplex[92], hence can particularly
easily be mistaken in serology.

Order   Pathogen

Figure 1: **Sankey diagram pairing study cases by pathogen and order.** Width of the ribbon is proportional to the number of cases included in this order-pathogen couple. Data are available in Supplementary Dataset 1.

Figure 2A-D: **Investigation efforts for TBP by Orders.** Top row shows for *Borrelia* spp. **(A)** the percentage of animals positive to tick-borne by order, **(B)** the number of tested animals in each order. Bottom row shows for TBFV **(C)** the percentage of animals positive by order, **(D)** the number of tested animals in each order. Most publications regarding TBFV used an indirect methodology (antibody detection), direct detection was used in only 11% of TBFV cases. **(B)** and **(D)** plots shows the number of tested individuals on a log10 scale as well as the number of different host species in each order on the inside. Orders are grouped by class, Mammalia are shown to the left, Aves to the right, and for *Borrelia* spp., Lepidosauria at the bottom. Data are available in Supplementary Dataset 1.

Figure 3A-B: **Average detection rates of pathogens and antibodies in the tested individuals, rounded to 0 digit.** The x-axis
represents the log10 of the sum of individuals that have been tested for **(A)** *Borrelia* spp. and **(B) Flaviviruses** in the selected
publications, the y-axis represents the % of average detection. DTV in oldest publications might have been detected under the
name of POWV, because DTV formerly corresponded to POWV lineage II. Data are available in Supplementary Dataset 1.

Study designs and eco-epidemiological parameters

Proposed study designs varied in spatial and temporal sampling, resulting in varying quality in data
acquisition. Most studies were cross-sectional (180), and 20 did not inform a study period. A total of 142
fieldworks were reported on two years or more (average study period = 2.9 years, median = 1 year) and
few studies (N = 56) were conducted over a 5-year time. Some studies conducted repeated sampling
without taking into consideration the period of sampling in data presentation, providing limited
epidemiological information. The longest studies recorded lasted 34 years to detect TBEV in striped field
mouse (*Apodemus agrarius*) and northern red-backed vole (*Myodes rutilus*) in Russia[114] and 30 years to
detect anti-POWV antibodies in white-tailed deer (*Odocoileus virginianus*)[135]. The spatial resolution varied
from locality to country scale, with more often province chosen as sampling location. Fieldwork areas were
described with environmental characteristic or administrative nomenclature. Inconstance came from
different focus either on a local targeted area or on over a broader-scale exposure to the pathogens. Sizes
of datasets were often small. The median sample size was 40 and the maximum sample size was 3186 rock
doves (*Columbia livia*) from an urban setting in Italy for *B. anserina* detection[136].

A total of 41 publications tagged captured individuals. But tags were not always specific and trustable,
some authors simply relying on scars made by biopsy or on feather clipping. The aim of marking was
sometimes only to avoid resampling, with immediate release in case of recapture. Only 24 publications
used individual tags and repeated capture occasions, and analysed for pathogen detection. In addition,
capture-mark-recapture data was not systematically analysed using relevant techniques. Capture-mark-
recapture is however an important specific sampling method to try to account for detection uncertainty.
For instance, Lambert *et al.* used seroprevalence and capture-mark-recapture data to model the force of
infection of brucellosis in the Alpine ibex (*Capra ibex*)[137]. Yet, in studies with no recapture, necropsies were
possible, allowing to search for the pathogen in specific organs. Infection and exposure can be evidenced
differently depending on tested biological material[138,139].

Moreover, metadata informing basic demographics such as sex and age were lacking in almost every case.
Whilst there could be a variability in pathogen susceptibility depending on age, these data are key in
understanding pathogens' transmission processes[140]. Mammals and birds or squamates were not studied
together in same areas. Designs in the selected publications tended to investigate a single order by setting
field material only fit for this order, whilst theorical definitions of reservoir and sentinel suggest that
reservoirs are communities rather than species[53]. The underlying population of hosts is seldomly discussed
and considered. Furthermore, although convenient, getting material from hunted, road-killed or rescued
animals implies bias in detection that should be accounted for. Monitoring in the wild should integrate
demographics. Indeed, recommendations of Yoccoz *et al.* is to use information on spatial scale and hosts
detectability to estimate unbiased demographic parameters[141]. Thereby, acquiring infection and exposure
data in wildlife should be designed using longitudinal studies targeting community of species with capture-
marking-recapture or patch-occupancy modelling.

Preceding remarks lead to interrogate the way species are targeted. Dogs have already been suggested as
possible sentinels for *Borrelia burgdorferi* and TBEV for humans[142,143]. Although easier to study from a
logistical perspective, these domestic species do not capture the information on the early dynamics of
transmission, mainly occurring in wildlife for tick-borne pathogens[75]. In addition, one gap in literature is
that for many investigated TBP, pathogenicity in vertebrate hosts is disregarded. For instance, while
Coxellia burnetii causes human Q fever, others *Coxellia*-like spp. are symbiotic in ticks and are not
pathogenic in vertebrates[144]. Therefore, combining data on pathogenicity from the field and the laboratory
is still needed.

The complementarity of laboratory and tick data

In the literature, models and experimental infections have attempted to characterize reservoir
competence[145–147]. Especially, xenodiagnoses are used to detect reservoir capacities in experimental
laboratory settings[148]. This consists in feeding non-infected ticks on a TBP laboratory-infected host, and
then record infection rates on uninfected hosts brought in contact with the ticks. In this way, vector
competence of ticks can also be assayed. Such experimental methodologies however do not account for
the infection in their ecosystems. Furthermore, ticks from the field have extensively been searched for
pathogens. For instance, from the 827 articles matching our literature search, 224 publications were
directly looking into ticks for pathogens with methods ranging from PCR to metagenomics[149,150].
Concerning birds, a literature review reported that most field studies were on host-attached ticks rather
than on samples from the birds themselves[151]. Studies on ticks can be complementary to data from hosts
insofar as they analyse tick-host interactions. Their deployment is less invasive on hosts than blood or
tissue sampling. They need less expertise in the field and are subject to less strict ethical protocols. In
addition, identification of ticks' life stages allows to infer ticks' contamination capacity, although other
ways of transmission such as vertical transmission in vertebrate hosts have been reported for some
agents[152]. It is also faster to collect a large number of individuals. Although it can be logistically easier to
collect ticks in the field to get an insight of TBP circulation and diversity, accounting for the whole local
host community and the infection status remains as important.

Conclusion

367   TBP are present in a wide range of vertebrate species, and rodents and artiodactyls were the most
368   investigated. Some infectious agents, rare TBFV or particular strains of *Borrelia* spp., were little studied
369   and information on their infecting capacities are still needed. Diagnostic detection methods varied with

the targeted pathogens. Selected publications showed that study designs, both in data collection and in data analysis, were heterogenous. Key information on host age and precise location was often unavailable; 58% (200/342) of the studies conducted fieldwork at one time point, and very few studies used quantitative methods to analyse the collected data. In combination with experimental data, quantifying transmission needs to gather empirical infection data from populational studies, including hosts demographics and allowing to identify what species act as reservoir in the local host community, and which one can be used as sentinel. Whilst we acknowledge that gathering all this information is difficult, this review highlights the necessity of including eco-epidemiological parameters in wildlife infection studies. The subject discussed in this review concerns a multidisciplinary scientific community (namely ecology, epidemiology, microbiology, parasitology, modelling, public health), emphasizing that getting and interpreting zoonotic infection data from wild animals in a rigorous way is important to build workable One Health programs.

Data availability

The data that support this study are available as Supplementary Dataset 1. Source data are provided with this paper.

Reference

1. Woolhouse, M. E. J. Population biology of emerging and re-emerging pathogens. *Trends Microbiol.* **10**, s3–s7 (2002).
2. Estrada-Peña, A. & De La Fuente, J. The ecology of ticks and epidemiology of tick-borne viral diseases. *Antiviral Res.* **108**, 104–128 (2014).
3. McCoy, K. D. & Boulanger, N. *Tiques et maladies à tiques: biologie, écologie évolutive, épidémiologie*. (IRD, 2015).
4. Becker, D. J., Seifert, S. N. & Carlson, C. J. Beyond Infection: Integrating Competence into Reservoir Host Prediction. *Trends Ecol. Evol.* **35**, 1062–1065 (2020).
5. Halliday, J. E. B. *et al.* A framework for evaluating animals as sentinels for infectious disease surveillance. *J. R. Soc. Interface* **4**, 973–984 (2007).
6. Bouchard, É. *et al.* Are foxes (Vulpes spp.) good sentinel species for Toxoplasma gondii in northern Canada? *Parasit. Vectors* **15**, 115 (2022).
7. Eby, P. *et al.* Pathogen spillover driven by rapid changes in bat ecology. *Nature* **613**, 340–344 (2023).
8. Riccardi, N. *et al.* Tick-borne encephalitis in Europe: a brief update on epidemiology, diagnosis, prevention, and treatment. *Eur. J. Intern. Med.* **62**, 1–6 (2019).
9. Varlacher, J.-F. *et al.* Tick-borne encephalitis. *Rev. Sci. Tech. OIE* **34**, 453–466 (2015).
10. Zlobin, V. I., Pogodina, V. V. & Kahl, O. A brief history of the discovery of tick-borne encephalitis virus in the late 1930s (based on reminiscences of members of the expeditions, their colleagues, and relatives). *Ticks Tick-Borne Dis.* **8**, 813–820 (2017).
11. Jaenson, T. G., Hjertqvist, M., Bergström, T. & Lundkvist, Å. Why is tick-borne encephalitis increasing? A review of the key factors causing the increasing incidence of human TBE in Sweden. *Parasit. Vectors* **5**, 184 (2012).
12. Amicizia, D. *et al.* Epidemiology of tick-borne encephalitis (TBE) in Europe and its prevention by available vaccines. *Hum. Vaccines Immunother.* **9**, 1163–1171 (2013).
13. Hubálek, Z. History of Arbovirus Research in the Czech Republic. *Viruses* **13**, 2334 (2021).
14. Chastel, C. *et al.* The isolation of Meaban virus, a new Flavivirus from the seabird tick Ornithodoros (Alectorobius) maritimus in France. *Arch. Virol.* **83**, 129–140 (1985).

15. McLean, D. M. & Donohue, W. L. Powassan virus: isolation of virus from a fatal case of encephalitis.
Can. Med. Assoc. J. **80**, 708–711 (1959).
16. Hermance, M. E. & Thangamani, S. Powassan Virus: An Emerging Arbovirus of Public Health Concern
in North America. *Vector-Borne Zoonotic Dis.* **17**, 453–462 (2017).
17. Eisen, L. Vector competence studies with hard ticks and Borrelia burgdorferi sensu lato spirochetes:
A review. *Ticks Tick-Borne Dis.* **11**, 101359 (2020).
18. Kahl, O. & Gray, J. S. The biology of Ixodes ricinus with emphasis on its ecology. *Ticks Tick-Borne Dis.*
14, 102114 (2023).
19. Rebaudet, S. & Parola, P. Epidemiology of relapsing fever borreliosis in Europe. *FEMS Immunol. Med.*
Microbiol. **48**, 11–15 (2006).
20. Kubiak, K., Szczotko, M. & Dmitryjuk, M. Borrelia miyamotoi—An Emerging Human Tick-Borne
Pathogen in Europe. *Microorganisms* **9**, 154 (2021).
21. Abi, M.-E. *et al.* Molecular Interactions During Borrelia burgdorferi Migration from the Vector to the
Mammalian Nervous System. *Curr. Protein Pept. Sci.* **21**, 517–526 (2020).
22. Kerstholt, M., Netea, M. G. & Joosten, L. A. B. Borrelia burgdorferi hijacks cellular metabolism of
immune cells: Consequences for host defense. *Ticks Tick-Borne Dis.* **11**, 101386 (2020).
23. Kurokawa, C. *et al.* Interactions between Borrelia burgdorferi and ticks. *Nat. Rev. Microbiol.* **18**, 587–
600 (2020).
24. Anderson, C. & Brissette, C. A. The Brilliance of Borrelia: Mechanisms of Host Immune Evasion by
Lyme Disease-Causing Spirochetes. *Pathogens* **10**, 281 (2021).
25. Helble, J. D., McCarthy, J. E. & Hu, L. T. Interactions between *Borrelia burgdorferi* and its hosts across
the enzootic cycle. *Parasite Immunol.* **43**, (2021).
26. Bogovič, P. *et al.* Clinical and Laboratory Characteristics and Outcome of Illness Caused by Tick-Borne
Encephalitis Virus without Central Nervous System Involvement. *Emerg. Infect. Dis.* **28**, 291–301
(2022).
27. Coburn, J. *et al.* Lyme Disease Pathogenesis. *Curr. Issues Mol. Biol.* 473–518 (2022)
doi:10.21775/cimb.042.473.
28. Stone, E. T. & Pinto, A. K. T Cells in Tick-Borne Flavivirus Encephalitis: A Review of Current Paradigms
in Protection and Disease Pathology. *Viruses* **15**, 958 (2023).
29. Dong, Y. *et al.* Global seroprevalence and sociodemographic characteristics of *Borrelia burgdorferi*
sensu lato in human populations: a systematic review and meta-analysis. *BMJ Glob. Health* **7**,
e007744 (2022).
30. Hoornstra, D. *et al.* Prevalence and clinical manifestation of Borrelia miyamotoi in Ixodes ticks and
humans in the northern hemisphere: a systematic review and meta-analysis. *Lancet Microbe* **3**,
e772–e786 (2022).
31. Wondim, M. A. *et al.* Epidemiological Trends of Trans-Boundary Tick-Borne Encephalitis in Europe,
2000–2019. *Pathogens* **11**, 704 (2022).
32. Burn, L. *et al.* Seroprevalence of Lyme Borreliosis in Europe: Results from a Systematic Literature
Review (2005–2020). *Vector-Borne Zoonotic Dis.* **23**, 195–220 (2023).
33. Teh, K. J. *et al.* Mini review: surveillance of Lyme borreliosis in Southeast Asia and method of
diagnosis. *Eur. Rev. Med. Pharmacol. Sci.* **27**, 4378–4385 (2023).
34. Raffetin, A. *et al.* Unconventional diagnostic tests for Lyme borreliosis: a systematic review. *Clin.*
Microbiol. Infect. **26**, 51–59 (2020).
35. Chou, E., Minor, A. & Cady, N. C. Quantitative multiplexed strategies for human Lyme disease
serological testing. *Exp. Biol. Med.* **246**, 1388–1399 (2021).
36. Gudowska-Sawczuk, M. & Mroczko, B. Selected Biomarkers of Tick-Borne Encephalitis: A Review. *Int.*
J. Mol. Sci. **22**, 10615 (2021).

37.   Henningsson, A. J. *et al.* Laboratory Methods for Detection of Infectious Agents and Serological
462   Response in Humans With Tick-Borne Infections: A Systematic Review of Evaluations Based on
463   Clinical Patient Samples. *Front. Public Health* **9**, 580102 (2021).
464   38.   Buczek, A. M., Buczek, W., Buczek, A. & Wysokińska-Miszczuk, J. Food-Borne Transmission of Tick-
465   Borne Encephalitis Virus—Spread, Consequences, and Prophylaxis. *Int. J. Environ. Res. Public. Health*
466   **19**, 1812 (2022).
467   39.   Elbaz, M. *et al.* Systematic Review and Meta-analysis of Foodborne Tickborne Encephalitis, Europe,
468   1980–2021. *Emerg. Infect. Dis.* **28**, (2022).
469   40.   Martello, E. *et al.* Systematic review on the non-vectorial transmission of Tick-borne encephalitis
470   virus (TBEv). *Ticks Tick-Borne Dis.* **13**, 102028 (2022).
471   41.   Zannou, O. M. *et al.* Models for Studying the Distribution of Ticks and Tick-Borne Diseases in Animals:
472   A Systematic Review and a Meta-Analysis with a Focus on Africa. *Pathogens* **10**, 893 (2021).
473   42.   Voyiatzaki, C. *et al.* Climate Changes Exacerbate the Spread of Ixodes ricinus and the Occurrence of
474   Lyme Borreliosis and Tick-Borne Encephalitis in Europe—How Climate Models Are Used as a Risk
475   Assessment Approach for Tick-Borne Diseases. *Int. J. Environ. Res. Public. Health* **19**, 6516 (2022).
476   43.   Colunga-Salas, P., Sánchez-Montes, S., Volkow, P., Ruíz-Remigio, A. & Becker, I. Lyme disease and
477   relapsing fever in Mexico: An overview of human and wildlife infections. *PLOS ONE* **15**, e0238496
478   (2020).
479   44.   Tsao, J. I., Hamer, S. A., Han, S., Sidge, J. L. & Hickling, G. J. The Contribution of Wildlife Hosts to the
480   Rise of Ticks and Tick-Borne Diseases in North America. *J. Med. Entomol.* **58**, 1565–1587 (2021).
481   45.   Defaye, B., Moutailler, S., Pasqualini, V. & Quilichini, Y. A Systematic Review of the Distribution of
482   Tick-Borne Pathogens in Wild Animals and Their Ticks in the Mediterranean Rim between 2000 and
483   2021. *Microorganisms* **10**, 1858 (2022).
484   46.   Mubemba, B. *et al.* Current knowledge of vector-borne zoonotic pathogens in Zambia: A clarion call
485   to scaling-up "One Health" research in the wake of emerging and re-emerging infectious diseases.
486   *PLoS Negl. Trop. Dis.* **16**, e0010193 (2022).
487   47.   Pandit, P. S. *et al.* Predicting wildlife reservoirs and global vulnerability to zoonotic Flaviviruses. *Nat.*
488   *Commun.* **9**, 5425 (2018).
489   48.   Greay, T. L. *et al.* Recent insights into the tick microbiome gained through next-generation
490   sequencing. *Parasit. Vectors* **11**, 12 (2018).
491   49.   Hussain, S. *et al.* The Role of Ticks in the Emergence of Borrelia burgdorferi as a Zoonotic Pathogen
492   and Its Vector Control: A Global Systemic Review. *Microorganisms* **9**, 2412 (2021).
493   50.   Hansford, K. M., Wheeler, B. W., Tschirren, B. & Medlock, J. M. Questing *Ixodes ricinus* ticks and
494   *Borrelia* spp. in urban green space across Europe: A review. *Zoonoses Public Health* **69**, 153–166
495   (2022).
496   51.   McVicar, M., Rivera, I., Reyes, J. B. & Gulia-Nuss, M. Ecology of Ixodes pacificus Ticks and Associated
497   Pathogens in the Western United States. *Pathogens* **11**, 89 (2022).
498   52.   Keeling, M. J. & Rohani, P. *Modeling Infectious Diseases in Humans and Animals*. (Princeton
499   University Press, 2011). doi:10.2307/j.ctvcm4gk0.
500   53.   Viana, M. *et al.* Assembling evidence for identifying reservoirs of infection. *Trends Ecol. Evol.* **29**, 270–
501   279 (2014).
502   54.   McClintock, B. T. *et al.* Seeking a second opinion: uncertainty in disease ecology: Uncertainty in
503   disease ecology. *Ecol. Lett.* **13**, 659–674 (2010).
504   55.   Sentenac, H. *et al.* Accounting for bias in prevalence estimation: The case of a globally emerging
505   pathogen. *J. Appl. Ecol.* 1365-2664.14457 (2023) doi:10.1111/1365-2664.14457.
506   56.   Spielman, A., Wilson, M. L., Levine, J. F. & Piesman, J. Ecology of Ixodes Dammini-Borne Human
507   Babesiosis and Lyme Disease. *Annu. Rev. Entomol.* **30**, 439–460 (1985).

508   57.   LoGiudice, K., Ostfeld, R. S., Schmidt, K. A. & Keesing, F. The ecology of infectious disease: Effects of
509   host diversity and community composition on Lyme disease risk. *Proc. Natl. Acad. Sci.* **100**, 567–571
510   (2003).
511   58.   Michelitsch, A., Wernike, K., Klaus, C., Dobler, G. & Beer, M. Exploring the Reservoir Hosts of Tick-
512   Borne Encephalitis Virus. *Viruses* **11**, 669 (2019).
513   59.   McLean, D. M., MacPherson, L. W., Walker, S. J. & Funk, G. Powassan virus: surveys of human and
514   animal sera. *Am. J. Public Health Nation Health* **50**, 1539–1544 (1960).
515   60.   McLean, D. M. *et al.* Powassan Virus: Investigations of Possible Natural Cycles of Infection. *J. Infect.*
516   *Dis.* **109**, 19–23 (1961).
517   61.   McLean, D. M. *et al.* Powassan Virus: Field Investigations in Northern Ontario, 1959 to 1961. *Can.*
518   *Med. Assoc. J.* **86**, 971–974 (1962).
519   62.   Mclean, D. M. & Larke, R. P. Powassan and Silverwater viruses: ecology of two Ontario arboviruses.
520   *Can. Med. Assoc. J.* **88**, 182–185 (1963).
521   63.   Mclean, D. M., Best, J. M., Mahalingam, S., Chernesky, M. A. & Wilson, W. E. Powassan virus: Summer
522   infection cycle, 1964. *Can. Med. Assoc. J.* **91**, 1360–1362 (1964).
523   64.   McLean, D. M., Smith, P. A., Livingstone, S. E., Wilson, W. E. & Wilson, A. G. Powassan virus: vernal
524   spread during 1965. *Can. Med. Assoc. J.* **94**, 532–536 (1966).
525   65.   McLean, D. M. *et al.* Powassan virus: persistence of virus activity during 1966. *Can. Med. Assoc. J.* **96**,
526   660–664 (1967).
527   66.   McLean, D. M., Ladyman, S. R. & Purvin-Good, K. W. Westward extension of Powassan virus
528   prevalence. *Can. Med. Assoc. J.* **98**, 946–949 (1968).
529   67.   McLean, D. M. *et al.* Arbovirus prevalence in the East Kootenay Region, 1968. *Can. Med. Assoc. J.*
530   **100**, 320–326 (1969).
531   68.   Burtis, J. C., Yavitt, J. B., Fahey, T. J. & Ostfeld, R. S. Ticks as Soil-Dwelling Arthropods: An Intersection
532   Between Disease and Soil Ecology. *J. Med. Entomol.* **56**, 1555–1564 (2019).
533   69.   Szekeres, S. *et al.* Eco-epidemiology of Borrelia miyamotoi and Lyme borreliosis spirochetes in a
534   popular hunting and recreational forest area in Hungary. *Parasit. Vectors* **8**, 309 (2015).
535   70.   Obiegala, A., Król, N., Oltersdorf, C., Nader, J. & Pfeffer, M. The enzootic life-cycle of Borrelia
536   burgdorferi (sensu lato) and tick-borne rickettsiae: an epidemiological study on wild-living small
537   mammals and their ticks from Saxony, Germany. *Parasit. Vectors* **10**, 115 (2017).
538   71.   Holbrook, M. R. Kyasanur forest disease. *Antiviral Res.* **96**, 353–362 (2012).
539   72.   Moore, S. I., Wilson, M. L., Spielman, A., Telford, S. R. & Mather, T. N. Incompetence of Deer as
540   Reservoirs of the Lyme Disease Spirochete. *Am. J. Trop. Med. Hyg.* **39**, 105–109 (1988).
541   73.   Kurtenbach, K., Sewell, H.-S., Ogden, N. H., Randolph, S. E. & Nuttall, P. A. Serum Complement
542   Sensitivity as a Key Factor in Lyme Disease Ecology. *Infect. Immun.* **66**, 1248–1251 (1998).
543   74.   Pearson, P., Rich, C., Feehan, M. J. R., Ditchkoff, S. S. & Rich, S. M. White-Tailed Deer Serum Kills the
544   Lyme Disease Spirochete, *Borrelia burgdorferi*. *Vector-Borne Zoonotic Dis.* vbz.2022.0095 (2023)
545   doi:10.1089/vbz.2022.0095.
546   75.   Radolf, J. D., Caimano, M. J., Stevenson, B. & Hu, L. T. Of ticks, mice and men: understanding the
547   dual-host lifestyle of Lyme disease spirochaetes. *Nat. Rev. Microbiol.* **10**, 87–99 (2012).
548   76.   Georgii, B. & Schröder, W. Home range and activity patterns of male red deer (Cervus elaphus L.) in
549   the alps. *Oecologia* **58**, 238–248 (1983).
550   77.   Kjellander, P. *et al.* Experimental evidence for density-dependence of home-range size in roe deer (
551   Capreolus capreolus L.): a comparison of two long-term studies. *Oecologia* **139**, 478–485 (2004).
552   78.   Morellet, N. *et al.* Seasonality, weather and climate affect home range size in roe deer across a wide
553   latitudinal gradient within Europe. *J. Anim. Ecol.* **82**, 1326–1339 (2013).
554   79.   Walter, W. D. *et al.* Heterogeneity of a landscape influences size of home range in a North American
555   cervid. *Sci. Rep.* **8**, 14667 (2018).

80. Tavernier, P. *et al.* Serologic screening for 13 infectious agents in roe deer (*Capreolus capreolus*) in Flanders. *Infect. Ecol. Epidemiol.* **5**, 29862 (2015).

81. Rijks, J. M. *et al.* Tick-Borne Encephalitis Virus Antibodies in Roe Deer, the Netherlands. *Emerg. Infect. Dis.* **25**, 342–345 (2019).

82. Holding, M. *et al.* Tick-Borne Encephalitis Virus, United Kingdom. *Emerg. Infect. Dis.* **26**, 90–96 (2020).

83. Paulsen, K. M. *et al.* Cervids as sentinel-species for tick-borne encephalitis virus in Norway - A serological study. *Zoonoses Public Health* **67**, 342–351 (2020).

84. Roelandt, S. *et al.* First TBEV serological screening in Flemish wild boar. *Infect. Ecol. Epidemiol.* **6**, 31099 (2016).

85. Gern, L. *et al.* European Reservoir Hosts of Borrelia burgdorferi sensu lato. *Zentralblatt Für Bakteriol.* **287**, 196–204 (1998).

86. Humphery-Smith, I., Cybinski, D. H., Byrnes, K. A. & St George, T. D. Seroepidemiology of arboviruses among seabirds and island residents of the Great Barrier Reef and Coral Sea. *Epidemiol. Infect.* **107**, 435–440 (1991).

87. Newman, E. A. *et al.* Borrelia burgdorferi Sensu Lato Spirochetes in Wild Birds in Northwestern California: Associations with Ecological Factors, Bird Behavior and Tick Infestation. *PLOS ONE* **10**, e0118146 (2015).

88. Bauer, S. & Hoye, B. J. Migratory Animals Couple Biodiversity and Ecosystem Functioning Worldwide. *Science* **344**, 1242552 (2014).

89. Boulinier, T. *et al.* Migration, Prospecting, Dispersal? What Host Movement Matters for Infectious Agent Circulation? *Integr. Comp. Biol.* **56**, 330–342 (2016).

90. De Carvalho, I. L. *et al.* Molecular Characterization of a New Isolate of *Borrelia lusitaniae* Derived from *Apodemus sylvaticus* in Portugal. *Vector-Borne Zoonotic Dis.* **10**, 531–534 (2010).

91. Cruciani, D. *et al.* Health Status of the Eastern Grey Squirrel (Sciurus carolinensis) Population in Umbria: Results of the LIFE Project 'U-SAVEREDS'. *Animals* **12**, 2741 (2022).

92. Carpio, K. L. *et al.* Differences in Genetic Diversity of Mammalian Tick-Borne Flaviviruses. *Viruses* **15**, 281 (2023).

93. Evans, N. J., Bown, K., Timofte, D., Simpson, V. R. & Birtles, R. J. Fatal Borreliosis in Bat Caused by Relapsing Fever Spirochete, United Kingdom. *Emerg. Infect. Dis.* **15**, 1331–1333 (2009).

94. Marinkelle, C. J. & Grose, E. S. Species of Borrelia from a Colombian Bat (Natalus tumidirostris). *Nature* **218**, 487–487 (1968).

95. Fish, D. *et al.* Niche Partitioning of Borrelia burgdorferi and Borrelia miyamotoi in the Same Tick Vector and Mammalian Reservoir Species. *Am. J. Trop. Med. Hyg.* **81**, 1120–1131 (2009).

96. Zhang, X.-A. *et al.* Molecular detection and identification of relapsing fever *Borrelia* in ticks and wild small mammals in China. *Emerg. Microbes Infect.* **11**, 2632–2635 (2022).

97. Scott, M. C. *et al.* High-Prevalence Borrelia miyamotoi Infection Among Wild Turkeys (Meleagris gallopavo) in Tennessee. *J. Med. Entomol.* **47**, 1238–1242 (2010).

98. García-Bocanegra, I. *et al.* Spatio-temporal trends and risk factors affecting West Nile virus and related flavivirus exposure in Spanish wild ruminants. *BMC Vet. Res.* **12**, 249 (2016).

99. Chastel, C. *et al.* [Arbovirus serological survey among marine and non-marine birds of Brittany]. *Bull. Soc. Pathol. Exot. Filiales* **78**, 594–605 (1985).

100. Bournez, L. *et al.* Exposure of Wild Ungulates to the Usutu and Tick-Borne Encephalitis Viruses in France in 2009–2014: Evidence of Undetected Flavivirus Circulation a Decade Ago. *Viruses* **12**, 10 (2019).

101. Krzysiak, M. K. *et al.* The European bison (Bison bonasus) as an indicatory species for the circulation of tick-borne encephalitis virus (TBEV) in natural foci in Poland. *Ticks Tick-Borne Dis.* **12**, 101799 (2021).

102. Norte, A. C. *et al.* Getting under the birds' skin: tissue tropism of Borrelia burgdorferi s.l. in naturally
and experimentally infected avian hosts. *Microb. Ecol.* **79**, 756–769 (2020).

103. O'Keeffe, K. R., Oppler, Z. J. & Brisson, D. Evolutionary ecology of Lyme Borrelia. *Infect. Genet. Evol.*
85, 104570 (2020).

104. Amore, G. *et al. Borrelia lusitaniae* in Immature *Ixodes ricinus* (Acari: Ixodidae) Feeding on Common
Wall Lizards in Tuscany, Central Italy. *J. Med. Entomol.* **44**, 303–307 (2007).

105. Ekner, A. *et al.* Anaplasmataceae and Borrelia burgdorferi sensu lato in the sand lizard Lacerta agilis
and co-infection of these bacteria in hosted Ixodes ricinus ticks. *Parasit. Vectors* **4**, 182 (2011).

106. De Sousa, R. *et al.* Role of the Lizard Teira dugesii as a Potential Host for Ixodes ricinus Tick-Borne
Pathogens. *Appl. Environ. Microbiol.* **78**, 3767–3769 (2012).

107. Musilová, L., Kybicová, K., Fialová, A., Richtrová, E. & Kulma, M. First isolation of Borrelia lusitaniae
DNA from green lizards (Lacerta viridis) and Ixodes ricinus ticks in the Czech Republic. *Ticks Tick-*
Borne Dis. **13**, 101887 (2022).

108. Van Tongeren, H. A. E. Viraemia and antibody response of the mallard (Anas platyrhynchos) to
infection with tick-borne encephalitis virus. *J. Comp. Pathol.* **93**, 521–530 (1983).

109. Salat, J. *et al.* Experimental and Natural Infections of Tick-Borne Encephalitis Virus in Dogs. *Viruses*
13, 2039 (2021).

110. Jemeršić, L. *et al.* Detection and genetic characterization of tick-borne encephalitis virus (TBEV)
derived from ticks removed from red foxes (Vulpes vulpes) and isolated from spleen samples of red
deer (Cervus elaphus) in Croatia. *Ticks Tick-Borne Dis.* **5**, 7–13 (2014).

111. Csank, T. *et al.* Detection of West Nile virus and tick-borne encephalitis virus in birds in Slovakia, using
a universal primer set. *Arch. Virol.* **161**, 1679–1683 (2016).

112. Esser, H. J. *et al.* Continued Circulation of Tick-Borne Encephalitis Virus Variants and Detection of
Novel Transmission Foci, the Netherlands. *Emerg. Infect. Dis.* **28**, 2416–2424 (2022).

113. Tonteri, E. *et al.* Tick-borne Encephalitis Virus in Wild Rodents in Winter, Finland, 2008–2009. *Emerg.*
Infect. Dis. **17**, 72–75 (2011).

114. Bakhvalova, V. N. *et al.* Tick-Borne Encephalitis Virus Diversity in Ixodid Ticks and Small Mammals in
South-Western Siberia, Russia. *Vector-Borne Zoonotic Dis.* **16**, 541–549 (2016).

115. Wu, Z. *et al.* Comparative analysis of rodent and small mammal viromes to better understand the
wildlife origin of emerging infectious diseases. *Microbiome* **6**, 178 (2018).

116. Dai, X., Shang, G., Lu, S., Yang, J. & Xu, J. A new subtype of eastern tick-borne encephalitis virus
discovered in Qinghai-Tibet Plateau, China. *Emerg. Microbes Infect.* **7**, 1–9 (2018).

117. Bestehorn, M. *et al.* Phylogenetics of tick-borne encephalitis virus in endemic foci in the upper Rhine
region in France and Germany. *PLOS ONE* **13**, e0204790 (2018).

118. Bai, Y. *et al.* Discovery of Tick-Borne Karshi Virus Implies Misinterpretation of the Tick-Borne
Encephalitis Virus Seroprevalence in Northwest China. *Front. Microbiol.* **13**, 872067 (2022).

119. Holzmann, H. Diagnosis of tick-borne encephalitis. *Vaccine* **21**, S36–S40 (2003).

120. Tonteri, E. *et al.* The Three Subtypes of Tick-Borne Encephalitis Virus Induce Encephalitis in a Natural
Host, the Bank Vole (Myodes glareolus). *PLoS ONE* **8**, e81214 (2013).

121. Benavides, J. A. *et al.* Estimating Loss of Brucella Abortus Antibodies from Age-Specific Serological
Data In Elk. *EcoHealth* **14**, 234–243 (2017).

122. Métras, R. *et al.* Drivers for Rift Valley fever emergence in Mayotte: A Bayesian modelling approach.
PLoS Negl. Trop. Dis. **11**, e0005767 (2017).

123. Kucharski, A. J. *et al.* Using paired serology and surveillance data to quantify dengue transmission
and control during a large outbreak in Fiji. *eLife* **7**, e34848 (2018).

124. Vilibic-Cavlek, T. *et al.* Cross-Reactive Antibodies in Tick-Borne Encephalitis: Case Report and
Literature Review. *Antibodies* **11**, 72 (2022).

125. Gritsun, T. S., Lashkevich, V. A. & Gould, E. A. Tick-borne encephalitis. *Antiviral Res.* **57**, 129–146
(2003).
126. Main, A. J., Downs, W. G., Shope, R. E. & Wallis, R. C. Avian arboviruses of the Witless Bay Seabird
Sanctuary, Newfoundland, Canada. *J. Wildl. Dis.* **12**, 182–194 (1976).
127. Gilbert, A. T. *et al.* Deciphering Serology to Understand the Ecology of Infectious Diseases in Wildlife.
EcoHealth **10**, 298–313 (2013).
128. Garnier, R. *et al.* Interpreting ELISA analyses from wild animal samples: Some recurrent issues and
solutions. *Funct. Ecol.* **31**, 2255–2262 (2017).
129. Paillard, L. *et al.* Serological signature of tick-borne pathogens in Scandinavian brown bears over two
decades. *Parasit. Vectors* **8**, 398 (2015).
130. Arnal, A. *et al.* Circulation of a Meaban-Like Virus in Yellow-Legged Gulls and Seabird Ticks in the
Western Mediterranean Basin. *PLoS ONE* **9**, e89601 (2014).
131. Jurado-Tarifa, E. *et al.* Monitoring of West Nile virus, Usutu virus and Meaban virus in waterfowl used
as decoys and wild raptors in southern Spain. *Comp. Immunol. Microbiol. Infect. Dis.* **49**, 58–64
(2016).
132. Weissbach, F. H. & Hirsch, H. H. Comparison of Two Commercial Tick-Borne Encephalitis Virus IgG
Enzyme-Linked Immunosorbent Assays. *Clin. Vaccine Immunol.* **22**, 754–760 (2015).
133. Randolph, S. E. & Rogers, D. J. Fragile transmission cycles of tick-borne encephalitis virus may be
disrupted by predicted climate change. *Proc. R. Soc. Lond. B Biol. Sci.* **267**, 1741–1744 (2000).
134. Ytrehus, B., Vainio, K., Dudman, S. G., Gilray, J. & Willoughby, K. Tick-Borne Encephalitis Virus and
Louping-Ill Virus May Co-Circulate in Southern Norway. *Vector-Borne Zoonotic Dis.* **13**, 762–768
(2013).
135. Bosco-Lauth, A. *et al.* Seroprevalence of Powassan Virus in New England Deer, 1979–2010. *Am. J.*
Trop. Med. Hyg. **88**, 1159–1162 (2013).
136. Fabbi, M. *et al.* Borrelia in Pigeons: no Serological Evidence of Borrelia burgdorferi Infection. *J. Vet.*
Med. Ser. B **42**, 503–507 (1995).
137. Lambert, S. *et al.* Combining seroprevalence and capture-mark-recapture data to estimate the force
of infection of brucellosis in a managed population of Alpine ibex. *Epidemics* **38**, 100542 (2022).
138. Hofmeister, E. K. & Childs, J. E. Eear biopsy location influences detection of Borrelia burgdorferi by
PCR, but not by culture in naturally infected Peromyscus leucopus. *J. Wildl. Dis.* **31**, 345–351 (1995).
139. Pascoe, E. L., De Vries, A., Esser, H. J., Koenraadt, C. J. M. & Sprong, H. Detection of tick-borne
encephalitis virus in ear tissue and dried blood spots from naturally infected wild rodents. *Parasit.*
Vectors **16**, 103 (2023).
140. Sweeny, A. R. & Albery, G. F. Exposure and susceptibility: The Twin Pillars of infection. *Funct. Ecol.*
36, 1713–1726 (2022).
141. Yoccoz, N. G., Nichols, J. D. & Boulinier, T. Monitoring of biological diversity in space and time. *Trends*
Ecol. Evol. **16**, 446–453 (2001).
142. Roelandt, S. *et al.* Tick-Borne Encephalitis Virus Seropositive Dog Detected in Belgium: Screening of
the Canine Population as Sentinels for Public Health. *Vector-Borne Zoonotic Dis.* **11**, 1371–1376
(2011).
143. Bowser, N. & Anderson, N. Dogs (Canis familiaris) as Sentinels for Human Infectious Disease and
Application to Canadian Populations: A Systematic Review. *Vet. Sci.* **5**, 83 (2018).
144. Brenner, A. E., Muñoz-Leal, S., Sachan, M., Labruna, M. B. & Raghavan, R. *Coxiella burnetii* and
Related Tick Endosymbionts Evolved from Pathogenic Ancestors. *Genome Biol. Evol.* **13**, evab108
(2021).
145. Giardina, A. R., Schmidt, K. A., Schauber, E. M. & Ostfeld, R. S. Modeling the role of songbirds and
rodents in the ecology of Lyme disease. *Can. J. Zool.* **78**, 2184–2197 (2000).

146. Ginsberg, H. S. *et al.* Reservoir Competence of Native North American Birds for the Lyme Disease
Spirochete, *Borrelia burgdorferi*. *J. Med. Entomol.* **42**, 445–449 (2005).
147. Ostfeld, R. S. *et al.* Life History and Demographic Drivers of Reservoir Competence for Three Tick-
Borne Zoonotic Pathogens. *PLoS ONE* **9**, e107387 (2014).
148. Donahue, J. G., Piesman, J. & Spielman, A. Reservoir Competence of White-Footed Mice for Lyme
Disease Spirochetes. *Am. J. Trop. Med. Hyg.* **36**, 92–96 (1987).
149. Harvey, E. *et al.* Extensive Diversity of RNA Viruses in Australian Ticks. *J. Virol.* **93**, e01358-18 (2019).
150. Binetruy, F. *et al.* A novel Borrelia species, intermediate between Lyme disease and relapsing fever
groups, in neotropical passerine-associated ticks. *Sci. Rep.* **10**, 10596 (2020).
151. Rataud, A., Henry, P., Moutailler, S. & Marsot, M. Research effort on birds' reservoir host potential
for Lyme borreliosis: A systematic review and perspectives. *Transbound. Emerg. Dis.* **69**, 2512–2522
(2022).
152. Larsson, C. *et al.* Complications of Pregnancy and Transplacental Transmission of Relapsing-Fever
Borreliosis. *J. Infect. Dis.* **194**, 1367–1374 (2006).
711

Acknowledgements

713 This work was funded by Agence Nationale de la Recherche under the ANRJCJC MoZArt project, grant
714 number ANR-22-CE35-0003.

715
716

Supplementary information

Literature metrics

A total of 827 articles were retrieved from the literature search, and 314 articles matched the inclusion criteria (Figure S1). The reference of the article, the study area, the study period, the pathogen, the species, the testing material and methodologies were recorded, as well as the prevalence of the pathogens and the number of tested individuals (see Supplementary Dataset 1). Those 314 articles were published between 1959 and 2022, with 2006 and 2008 as median start and end dates for fieldwork efforts in year-referenced publications (Figure S-2A).

Case studies were conducted over a 63-years span (1959-2022), with almost three-quarters (71%, N = 678) done in Europe and Northern America. 73% looked at *Borrelia* and 27% at flaviviruses. Finally, from the 347 vertebrate species investigated, 86% (N = 297) were searched for *Borrelia* spp., 32% (N = 114) for flaviviruses, and 7% (N = 64) were investigated for both.

From these, a total of 947 unique combinations (or "case studies") were identified, defined by one species, studied for one infectious agent, on a defined study area, and over a specific time period. 52 countries were represented in the working dataset. Most case studies were reporting fieldwork data from Europe (37%, N = 352) and Northern America (34%, N = 326), whilst 21% were conducted in Asia (N = 201), 5% in Central and Southern America (N = 50), 2% in Africa (N = 17) and less than 1% in Oceania (N = 1) (Figure S-2B). 69% of case studies (N = 689) looked at *Borrelia* spp. and 31% (N = 258) looked at TBFV. Out of 314 publications, only 3 looked at both kinds of pathogen. Over two-third of Borrelia studies (71%, N = 490) searched for the pathogen itself, whilst the others searched for antibodies. In contrast, the large majority of TBFV studies (89%, N = 230) performed antibody detection, whilst the remaining 11% (N = 28) searched directly for the pathogen.

Among tick-borne *Borrelia* spp., the most searched pathogens were *B. burgdorferi* sensu lato (Bbsl), but a large diversity of species and genospecies were studied. They included, ordered from greatest to smallest amount of tested individuals, *B. burgdorferi* sensu stricto (Bbss), unclassified tick-borne *B.* spp., relapsing fever group tick-borne *B.* spp., *B. afzelii*, *B. miyamotoi*, *B. anserina*, *B. garinii*, *B. lonestari*, *B. hermsii*, *B. lusitaniae*, *B. persica*, *B. turicatae*, *B. bissettiae*, *B. japonica*, *Candidatus* B. fainii, *B. tanukii*, *B. theileri*, *B.* strain HP3, *Candidatus* B. paulista, *B. carolinensis*, *B. turdi*, *B. andersonii*, *B. valaisiana* and *B. venezuelensis*. More than half of TBFV cases looked at antibody detection against Tick-Borne Encephalitis virus (TBEV) or Powassan virus (POWV). TBEV and POWV were well-investigated for some localities, with large datasets: out of the 12 largest datasets (i.e. with more than 1000 tested individuals), 8 were collected for anti-TBEV or anti-POWV antibodies detection and 2 for TBEV-targeted PCR. Yet, the diversity of TBF investigated also included Meaban virus (MEAV), Louping Ill virus (LIV), Deer Tick virus (DTV), Omsk Haemorrhagic Fever virus (OHFV), Tyuleniy virus (TYUV), Kyansanur Forest Disease virus (KFDV), Yamaguchi virus (YV) and Saumarez Reef virus (SREV).

About 40% of the studies used a form of PCR (such as nested-PCR, rt-PCR, q-PCR) for pathogen detection and about 20% ELISA to get seroprevalence (Figure S2C). Detection methods changed with time and availability of new techniques. Oldest articles used mainly microscopy on biological material cultivation on Barbour-Stoenner-Kelly medium or blood smears, hemagglutination assay and immunofluorescence assay on blood instead of PCR on nucleic acid extracted from blood and other tissues and ELISA on whole blood or serum. Inoculation of serum in mice in order to survey infection used to be applied for detection. Spirochetes cultivation used to be central in identifying *Borrelia* spp. and progressively was validated and

759 then substituted by the use of PCR. PCR, like the use of ELISA detection before a validation with serum
760 neutralization test, is now frequently used as a screening tool to optimize sequencing use.

Limitations due to paper access

Effectiveness in TBFV investigation can also particularly be questioned, with some flaviviruses having a
status of neglected diseases making difficult to find relevant work on them (Bhatia *et al.* 2020). In addition,
some publications on rare TBFV were not available online (without abstract and not in English). TYUV for
instance is a Siberian seabird TBFV that has been known for a long time. During the seventies, publications
started to clarify which species could host the virus but today traces of them are lacking. Effort in virus
investigation has also later been less intensive, before a new surge of interest today.

768
769 *Figure S1:* **Diagram of inclusion and exclusion in database screening.**

771
Figure S2A-C: **Location, period and method in selected publications. (A) Country location of selected publications' study area,**
categorized by continent. About a third of the publications (34%) were from the US or Canada, 37% related to European countries,
and 21% to Asia. AU = Australia; AT =Austria; BE = Belgium; CA = Canada; CH = Switzerland; CN = China; CO = Colombia; CZ = Czech
Republic; DE = Germany; DK = Denmark; ES = Spain; FI = Finland; FO = Faeroe Islands; FR = France; GB = Great Britain and Northern
Ireland; HU = Hungary; IL =Israel; IS = Iceland; IT = Italy; JP = Japan; KR = Republic of Korea; KZ = Kazakhstan; MA = Morocco; MG
= Madagascar; ML = Mali; MN = Mongolia; MR = Mauritania; MX = Mexico; MY = Malaysia; NA = Namibia; NL = the Netherlands;
NO = Norway; PL = Poland; PT = Portugal; SI = Slovenia; SK = Slovakia; TF = French Southern Territory; TH = Thailand; TN = Tunisia;
TR = Turkey; RO = Romania; RU = Russia; US = United States of America; ZM = Zambia. **(B) Distribution of the study years of**
fieldwork effort. From data on start and end dates, a count of publications putting fieldwork effort was implemented by year.
Unfortunately, 20 publications did not inform their time period of fieldwork. **(C) Detection methodologies.** PCR = Polymerase in
Chain Reaction; ELISA = Enzyme-Linked Immunosorbent Assay; IFA = Immunofluorescence Assay; μ = microscopy; HA =
Hemagglutination Assay; SNT = Serum Neutralisation Titer; SNAP = commercial veterinary Ig detection kit; WB = Western Blot; IP
= Indirect Immunoperoxidase; NI = Not Informed; SB = Southern Blot; seq = Sequencing; SIA = Strip Immunoblot Assay; CFR =
Complement Fixation Reaction; eph = electrophoresis; inj = Injection in mice. Data are available in Supplementary Dataset 1.