7 **Niche-specific macrophage loss promotes skin capillary aging**

14 Kailin R. Mesa[1*], Kevin A. O'Connor[1], Charles Ng[2], Steven P. Salvatore[2], Alexandra
15 Dolynuk[1], Michelle Rivera Lomeli[1], Dan R. Littman[1,3,4*]
16
17
[1] Department of Cell Biology, New York University School of Medicine, New York, NY
10016, USA
[2] Department of Pathology and Laboratory Medicine, Weill Cornell Medicine, New York-
Presbyterian Hospital, New York, NY 10021, USA
[3] Perlmutter Cancer Center, New York University Langone Health, New York, NY
10016, USA
[4] Howard Hughes Medical Institute, New York, NY 10016, USA
30 * To whom correspondence should be addressed
31
32 Email: kai.mesa@med.nyu.edu or dan.littman@med.nyu.edu
33 Tel: 212-263-7520

41 Key Words: Aging / Resident Macrophages / Blood Capillaries / Tissue Repair /
42 Intravital Microscopy
43
44
45
46 Running Title: Niche-specific macrophage loss promotes capillary aging

47 **Summary**

48
All mammalian organs depend upon resident macrophage populations to coordinate

repair processes and facilitate tissue-specific functions[1–3]. Recent work has

established that functionally distinct macrophage populations reside in discrete tissue

niches and are replenished through some combination of local proliferation and

monocyte recruitment[4,5]. Moreover, decline in macrophage abundance and function in

tissues has been shown to contribute to many age-associated pathologies, such as

atherosclerosis, cancer, and neurodegeneration[6–8]. Despite these advances, the

cellular mechanisms that coordinate macrophage organization and replenishment

within an aging tissue niche remain largely unknown. Here we show that capillary-

associated macrophages (CAMs) are selectively lost over time, which contributes to

impaired vascular repair and tissue perfusion in older mice. To investigate resident

macrophage behavior *in vivo*, we have employed intravital two-photon microscopy to

non-invasively image in live mice the skin capillary plexus, a spatially well-defined

model of niche aging that undergoes rarefication and functional decline with age. We

find that CAMs are lost with age at a rate that outpaces that of capillary loss, leading to

the progressive accumulation of capillary niches without an associated macrophage in

both mice and humans. Phagocytic activity of CAMs was locally required to repair

obstructed capillary blood flow, leaving macrophage-less niches selectively vulnerable

to both homeostatic and injury-induced loss in blood flow. Our work demonstrates that

homeostatic renewal of resident macrophages is not as finely tuned as has been

previously suggested[9–11]. Specifically, we found that neighboring macrophages do not

proliferate or reorganize sufficiently to maintain an optimal population across the skin

capillary niche in the absence of additional cues from acute tissue damage or increased

abundance of growth factors, such as colony stimulating factor 1 (CSF1). Such

limitations in homeostatic renewal and organization of various niche-resident cell types

are potentially early contributors to tissue aging, which may provide novel opportunities

for future therapeutic interventions.

79   **Main Text**

Tissue homeostasis is dependent on multiple macrophage populations that

reside in distinct sub-tissue compartments or niches, such as epithelium, blood vessels

or nerves, and are thought to support specialized tissue functions[4,12,13]. Recent work

has suggested that functional decline and rarefication of vascular niches may contribute

to various age-associated tissue pathologies (including sarcopenia, chronic wounds,

and Alzheimer's disease)[14–16]. It is not yet known how tissue-resident macrophages

resist or potentiate such niche-specific aging processes[6,17].

Capillary-associated macrophage decline with age correlates with impaired blood

flow.

To model mammalian tissue aging, we adapted an intravital microscopy

technique to visualize skin resident macrophage populations non-invasively in live mice

throughout the lifetime of the organism[18,19] (Figure 1a, Supplemental Videos 1 and 2).

Unexpectedly, longitudinal imaging of skin macrophages (marked by *Csf1r-EGFP*)

revealed niche-specific decline in macrophage populations. Subdividing the skin into

three anatomical layers, epidermis, upper (papillary) and lower (reticulated) dermis, we

observed that macrophages of the upper dermis were lost with age at a greater rate

than macrophages from both epidermis and lower dermis (Figure 1b-c). Further

characterization of this upper dermal CSF1R[+] population revealed expression of

additional macrophage markers[20–22] , including the chemokine receptor CX_3CR1,

lysozyme M (LysM), and the mannose receptor CD206 (Extended Data Fig. 1a,b,g,h). A

major component of the upper dermal niche is the superficial capillary plexus, which

supplies nutrient exchange for the overlying epidermis. To visualize this structure, we

utilized third harmonic generation (3[rd] Harmonic) from our imaging to track red blood cell

(RBC) flow through capillary vessels[23,24], which was comparable to conventional

rhodamine dextran labeling (Extended Data Fig. 2 and Supplemental Video 3) and not

significantly altered when utilizing a coverslip during our imaging sessions (Extended

Data Fig. 3a,b) . With this *in vivo* marker of RBC flow, we found that these macrophages

were closely associated with blood capillaries of the superficial plexus, suggesting that

they may provide support for this capillary niche (Extended Data Fig. 1c-e and

Supplemental Video 4). To investigate if these macrophages play a role in capillary

function, we assessed if RBC blood flow was altered in the presence of capillary-

associated macrophages (CAMs). Performing timelapse recordings of fluorescently

labeled CAMs in 2 month old mice with a cre-dependent dual reporter system (*Cx3cr1-*

CreERT2; R26-mTmG), we found that capillaries lacking an associated macrophage

had a higher rate of obstructed RBC flow (Figure 1d,e and Supplemental Video 5). We

next performed longitudinal analysis across multiple ages (1-18 months) to assess CAM

coverage and capillary function during physiological aging. We found across all ages

tested a significant loss in RBC blood flow in capillaries lacking associated

macrophages (Extended Data Fig. 3c). Furthermore, we found the fraction of capillaries

with an associated macrophage significantly decreased with age (Figure 1f). This

decrease in CAMs and coverage outpaced the loss of capillaries (Extended Data Fig.

3e,f), which was previously shown to be an early hallmark of mouse and human aging in

multiple tissues, including central nervous system, lung, kidney, and skin[14,15,25–33].

Therefore, to assess if this phenomenon also occurs in humans, we obtained both

young and old human patient skin samples. Consistent with our observations in mice,

human capillary-associated macrophages also displayed a decline with age. Moreover,

CAM decline also outpaced capillary loss with age, suggesting a similar loss in

macrophage coverage of the capillary niche (Extended Data Fig. 3g-j). Therefore, these

observations suggest that local macrophage loss in both mice and human may

contribute to impaired capillary function with age. To assess any functional role of CAMs

in maintaining capillary blood flow, we performed chemical and genetic ablation of

CAMs, via clodronate liposomes and *Cx3cr1-DTR* depletion, respectively, and observed

both acute CAM loss as well as loss in capillary flow (Figure 1 g,h and Extended Data

Fig. 4). Collectively, this work highlights an evolutionarily conserved loss in skin

capillary-associated macrophages with age, which correlates with impaired homeostatic

capillary perfusion.

Local recruitment of CAMs is required to restore capillary blood flow.

Given the link between CAMs and capillary blood flow, we next assessed the

long-term fate of vessels that lack an associated macrophage. To this end, we

performed a 6-month time course in *Cx3cr1-GFP*;*R26-mTmG* mice from 1 to 7 months

of age (Figure 2a) and found that capillaries fated for pruning in the 6 month time course

had decreased CAM coverage as compared to capillaries that were maintained (Figure

2b-c). This finding suggests that CAMs might be locally required to maintain proper

capillary function and preservation with age. Therefore, to assess the cellular

mechanism(s) by which CAMs support capillary function, we utilized a laser-induced

blood clotting model to precisely target and stop blood flow in individual capillary

segments (Figure 2d, Extended Data Fig. 5 and Supplemental Video 6). To assess

macrophage involvement, we tracked the daily displacement of surrounding CAMs to

laser-induced clots. These data showed that CAM recruitment to sites of capillary

damage as well as RBC engulfment are locally restricted to within approximately 80μm,

and largely occur within the first two days after injury (Figure 2e,f). This is consistent

with previous work describing macrophage cloaking as an acute behavioral response to

laser-induced tissue damage[34,35]. Multiple signaling pathways are involved in sensing

tissue damage[34,36,37], including the chemokine receptor CX3CR1, which is

homeostatically expressed by CAMs (Extended Data Fig. 1a-f). Therefore, we

interrogated the role of CX3CR1 signaling in recruiting nearby CAMs to capillary

damage. We performed laser-induced clotting and found significant impairment in CAM

recruitment in *Cx3cr1-gfp/gfp* mice as compared to *Cx3cr1-gfp/+* controls (Figure 2g-h).

We also found that CX3CR1-deficiency led to a significant delay and overall reduction in

capillary reperfusion by Day 7 post clot induction (Figure 2i). From these findings, we

assessed capillary RBC flow during physiological aging and found a significant loss in

flowing capillaries in *Cx3cr1-gfp/gfp* mice, as compared to *Cx3cr1-gfp/+* controls, by 6

months of age (Figure 2j). Together, our results support a model where local CAM

recruitment, in part through CX3CR1-signaling, is critical for both capillary repair and the

long-term preservation of the vascular network during physiological aging.

CAMs are required to clear vascular damage and preserve skin capillaries during

aging

To determine if diminished capillary function with age is directly related to

macrophage loss, we took advantage of the inherent variability in CAM density in older

mice. Specifically, we performed the laser-induced clotting of aged skin capillaries with

or without local CAMs (within 75μm from clot) (Figure 3a). Remarkably, capillaries that

retained local CAMs in aged mice were significantly better at reestablishing blood flow

as compared to capillaries without local CAMs in the same mice (Figure 3b). This result

suggests that local macrophage loss drives age-associated capillary dysfunction. To

directly test the role of macrophages in capillary repair, we performed laser-induced

ablation of local CAMs immediately prior to capillary clot-induction. Indeed, in regions

where CAMs were ablated, repair was significantly impaired and blood flow was not

properly reestablished (Figure 3c,d). We targeted other perivascular cells with the same

laser ablation conditions and found no impairment to capillary repair, nor did we detect

neutrophil swarming as has been described for much larger areas of laser-induced

damage[34,35,38–44] (Extended Data Fig. 6 and 7).

From our serial imaging of capillary repair, we found nearby CAMs often

surrounded by and containing RBC debris. To mechanistically understand if CAM

uptake and clearance of this vascular debris is functionally important for capillary repair,

we acutely impaired CAM phagocytosis through inducible cre-dependent knockout of

Rac1, a critical component of the phagocytic machinery[45], one week prior to laser-

induced clot formation. While there was no significant change in CAM density (Extended

Data Fig. 8), there was significant impairment in clearance of RBC debris and capillary

reperfusion in *Cx3cr1^{CreER};Rac1$^{fl/fl}$* mice compared to *Cx3cr1^{CreER};Rac1$^{fl/+}$* littermate

controls (Figure 3e-g), suggesting that Rac1-dependent phagocytic clearance is critical

for proper capillary repair and tissue reperfusion.

To test the long-term vascular effects of Rac1-deficiency in CAMs, we assessed

capillary density three months after tamoxifen administration in *Cx3cr1^{CreER};Rac1$^{fl/fl}$*

mice. We found an acceleration in rate of capillary pruning after three months of

physiological aging in the skin of *Cx3cr1^{CreER};Rac1$^{fl/fl}$* mice compared to

Cx3cr1^{CreER};Rac1$^{fl/+}$ littermate controls (Figure 3h-j). Therefore, it is likely that loss of

Rac-1 dependent behaviors, such as phagocytosis of vascular debris, directly

contributes to impaired recovery and preservation of the skin microvascular network.

Together, our results support a model in which local CAM recruitment and

phagocytic clearance of vascular debris is critical to maintain capillary function.

Therefore, as CAM density declines with age, so does capillary perfusion of the tissue

(Figure 3k).

Dermal macrophages utilize niche-specific self-renewal strategies leading to

selective CAM loss with age

Maintenance of tissue resident macrophage populations in the skin is thought to

be mediated through a combination of local proliferation and systemic replacement by

bone marrow-derived blood monocytes[13,46,47]. To understand how CAMs are

replenished under physiological conditions, we generated bone marrow chimeras with

Csf1r-GFP;*CAG-dsRed* bone marrow transferred into lethally irradiated *Csf1r-GFP*

mice. Importantly, the hind paws of these mice were lead-shielded to prevent loss of

resident macrophage populations from our imaging area (Figure 4a). Tracking

macrophage populations from all anatomical layers of the skin for 10 weeks post

transplantation showed that while most lower dermal macrophages are replaced by

monocytes, <5% of upper dermal macrophages were bone marrow-derived (Figure

4b,c). Additionally, capillaries with either bone marrow-derived (GFP+/dsRed+) or host-

derived (GFP+/dsRed-) CAMs showed no obvious differences in RBC blood flow

(Extended Data Fig. 3d), suggesting that CAM ontogeny may not play a major role in

homeostatic capillary function.

Given that nearly all CAMs remained host-derived, we next performed single-

macrophage lineage tracing to monitor local proliferation of dermal macrophages.

Specifically, we induced sparse cre-recombination in 1 month old *Cx3cr1-GFP*;*R26-*

mTmG mice to label and track individual macrophages from both the upper and lower

dermis over weekly revisits (Figure 4d-e). From these serial revisits, we observed

striking positional stability of macrophages on the vascular network as well as local cell

division (Figure 4f). Analysis of monthly rates of macrophage loss and division revealed

that lower dermal macrophages have a significantly lower division rate as well as

significantly higher loss rate, as compared to CAMs (upper dermal macrophages)

(Figure 4g).

Focusing on CAM turnover, we also found a significant skew toward cell loss

over division during the 4-month time course (Figure 4h). Consistent with these data,

there was a significant decline in the fraction of fate-mapped CAMs over the 20-week

time course (Figure 4i). These results demonstrate that CX3CR1+ dermal macrophage

self-renewal strategies are niche-specific, where blood monocyte recruitment is utilized

in the lower dermis and local cell division is utilized in the upper dermis. Importantly, we

find that CAMs from the upper dermis are insufficiently replenished by local proliferation,

which contributes to their progressive decline in this tissue niche (Figure 4j).

Macrophage loss without local tissue damage is not sufficient to promote CAM

renewal

The ability of resident macrophages to locally self-renew has largely been

studied through methods of near-total macrophage depletion, which have limited niche-

specificity and often generate tissue-wide inflammation[9–11]. To directly interrogate the

steps of macrophage self-renewal over time, we tracked both the replacement of

individual macrophages after loss and the redistribution of sister macrophages after

division. First, to track local macrophage replacement after CAM loss, we performed

laser-induced ablation of all CAMs within a defined $500\mu m^2$ region. Serial revisits up to

two weeks after ablation revealed minimal repopulation from adjacent capillary regions

that retained intact CAM populations (Figure 5a). We found a similar lack in

repopulation following partial CAM depletion in *Cx3cr1^DTR* mice following low dose

diphtheria toxin administration (Figure 5b). To avoid any non-physiological effects from

these cell depletion models, we developed a dual fluorescent macrophage reporter

mouse, *Cx3cr1-CreERT2; Rosa26-dsRed; Csf1r-EGFP*, allowing for cre-dependent

recombination to differentially label a small fraction of macrophages and track their

homeostatic replacement. Examination on serial weekly revisits indicated that, as we

observed in depletion models, most capillary niches did not recruit a new macrophage

for as long as two weeks following CAM loss (Figure 5c,d). In contrast, when we

employed large laser-induced damage ($500\mu m^2$ region) in either the upper dermis or

overlaying epidermis (Figure 5e and Extended Data Fig. 9a) we found CAMs were

readily replenished (Figure 5e-g), in a CCR2-dependent manner, suggesting at least

partial monocyte repopulation of CAMs after tissue damage. We also found that local

CAM proliferation was significantly increased one week following laser-induced damage

in either the upper dermis or epidermis (Figure 5h,i and Extended Data Fig. 9a-c).

Taken together, these results suggest that CAM loss alone is not a sufficient trigger to

promote macrophage replenishment and requires additional signals elicited by tissue

damage.

Second, to precisely track if proliferating CAMs readily redistribute across the

capillary network following cell division, we utilized mice with a dual fluorescent nuclear

reporter, *Cx3cr1-CreERT2; R26-nTnG*, and administered a low or high dose of

tamoxifen to label either a sparse subset or all CAMs, respectively (Extended Data Fig.

9d). Compared to the average distance between all nearest neighboring macrophages,

sister CAMs remained significantly closer (<15μm) to each other for at least 2 weeks

after division (Extended Data Fig. 9e,f). These findings further support the notion that

CAM division and loss are not spatiotemporally coupled, which we predict would

progressively lead to disorganized patterning and the accumulation of both empty and

crowded capillary regions. We tested this prediction by looking at the distribution of

neighboring CAMs in both young (2-month-old) and old (10-month-old) mice. In young

mice, the majority of macrophages were within 50μm of each other. In contrast, old mice

had a biphasic distribution of macrophage patterning, with most CAMs either within

25μm or further than 75μm apart (Extended Data Fig. 9g). These results highlight two

distinct cellular features that contribute to reduced CAM coverage with age: 1)

insufficient macrophage repopulation following CAM loss, and 2) insufficient

redistribution of these cells along the capillary niche, which may promote progressive

erosion of the vascular network (Extended Data Fig. 9h).

Local CAM expansion in old mice is sufficient to rejuvenate capillary repair and

tissue reperfusion

While homeostatic CAM proliferation was insufficient to maintain a stable

population density, we asked if extrinsic cues such as large tissue damage could

increase CAM density long-term to improve capillary function in older mice. To this end,

we found that large laser-induced epidermal damage ($500μm^2$ region) resulted in a

lasting increase in CAMs below the damaged regions compared to neighboring control

regions (Extended Data Fig. 9i-k). Furthermore, we found a significant improvement in

capillary repair in these same regions following laser-induced clotting (Extended Data

Fig. 9l).

Our results suggest that CAMs in old mice can be stably expanded following

environmental changes, such as tissue damage. Therefore, we next assessed if directly

increasing CAM density, without local damage, would also be sufficient to improve

future capillary repair and reperfusion. To this end, we utilized a fusion protein of the

canonical macrophage growth factor, CSF1, with the Fc region of porcine IgG (CSF1-

Fc), as it has been reported to robustly increase macrophage density in multiple tissues,

including skin[48–50]. We performed daily intradermal injections of either CSF1-Fc or PBS

in the left or right hind paws, respectively, of the same mice (Figure 6a). There was a

significant increase in CAMs in the CSF1-treated paws compared to contralateral PBS

controls, which had no significant change from pretreatment (Figure 6b,c).

To assess if CSF1-treatment modulated local CAM survival/proliferation or simply

recruited new bone marrow-derived macrophages from blood monocytes, we performed

CSF1-treatment in chimeric bone marrow mice where we could track the relative

expansion of local and recruited populations (Extended Data Fig. 10a,b). Consistent

with our previous experiments, we found that PBS injected paws showed no significant

increase in host- or BM-derived macrophages in our imaging areas (Extended Data Fig.

10c-d). Importantly we found that CSF1-induced CAM expansion did not alter the ratio

of resident (GFP+) and recruited (GFP+/dsRed+) CAMs (Extended Data Fig. 10e), as a

relative increase in GFP+/dsRed+ CAMs over GFP+ CAMs would have suggested BM-

derived monocyte recruitment. Therefore, this strongly suggests that CSF1 drives local

proliferation of the existing CAM population.

Strikingly, we found that CSF1-treatment was sufficient to improve homeostatic

capillary blood flow in old mice, as compared to PBS controls which had significantly

more obstructed capillary segments (Figure 6d,e). Utilizing the same aged mice, we

next tested if this increase in CAM density would be sufficient to improve capillary repair

rates. Following laser-induced clotting, there was a significant improvement in capillary

repair and reperfusion in CSF1-treated mice compared to PBS controls (Figure 6f,g),

demonstrating that restoring dermal macrophage density in old mice can improve age-

associated vascular dysfunction.

Discussion

Macrophage renewal has largely been studied in non-physiological settings, such

as through *in vitro* cell culture or severe depletion models that often are accompanied

by acute inflammation[9,10,51–53]. Our work clearly demonstrates that the homeostatic

renewal strategies of resident macrophages is niche-specific and not as finely tuned as

has been previously suggested. Specifically, we found that macrophages of the upper

dermis do not proliferate or redistribute sufficiently to maintain an optimal coverage

across the skin capillary network unless they receive additional cues from acute tissue

damage or increased growth factor abundance (Figure 6h). Interestingly, we confirmed

previous findings that show epidermal Langerhans cell density also declines with age,

which has been associated with impaired epidermal function[54–57]. This raises the

possibility that age-associated loss in macrophage density is a more general

phenomenon in populations that rely on local self-renewal.

In addition to self-renewal, we also found that CAM recruitment to repair tissue

damage was spatially restricted. To our knowledge, the long-term size and stability of

resident macrophage territories or niches *in vivo* has not been reported. Additionally,

our work provides strong evidence that injury-induced dermal macrophage recruitment

is restricted to approximately 80μm, as has been demonstrated in other tissues[34,35]. In

young mice, CAM density is high enough to provide substantial niche/territory overlap

between neighbors. However, with declining CAM density with age, we show that a

significant fraction of the skin capillary network is no longer within a CAM's territory

range. It will be important to understand how this property of resident macrophages is

influenced by other aspects of regional heterogeneity, such as innate immune

imprinting[13], to shape local immune responses in tissues.

Collectively, this work demonstrates that loss in CAMs: 1) begins within the first

few months of life, 2) is progressive throughout life, and 3) is functionally detrimental to

vascular function and preservation, which has been shown to be a primary driver of

age-associated tissue impairments[15,58]. Furthermore, this work provides a novel

platform to investigate age-associated deviations in tissue homeostasis at the single-cell

level in a living mammal.

References

1. Nobs, S. P. & Kopf, M. Tissue-resident macrophages: guardians of organ homeostasis. *Trends Immunol* **42**, 495–507 (2021).

2. Minutti, C. M., Knipper, J. A., Allen, J. E. & Zaiss, D. Tissue-specific contribution of macrophages to wound healing. *Semin Cell Dev Biol* **61**, 3–11 (2017).

3. Vannella, K. M. & Wynn, T. A. Mechanisms of Organ Injury and Repair by Macrophages. *Annual Review of Physiology* **79**, (2016).

4. Chakarov, S. *et al.* Two distinct interstitial macrophage populations coexist across tissues in specific subtissular niches. *Science* **363**, eaau0964 (2019).

5. Wu, Y. & Hirschi, K. K. Tissue-Resident Macrophage Development and Function. *Frontiers Cell Dev Biology* **8**, 617879 (2021).

6. Beek, A. A. van, Bossche, J. den, Mastroberardino, P. G., Winther, M. de & Leenen, P. Metabolic Alterations in Aging Macrophages: Ingredients for Inflammaging? *Trends Immunol* (2019) doi:10.1016/j.it.2018.12.007.

7. Franceschi, C., Garagnani, P., Vitale, G., Capri, M. & Salvioli, S. Inflammaging and 'Garb-aging.' *Trends Endocrinol Metabolism* **28**, 199–212 (2017).

8. Mass, E., Nimmerjahn, F., Kierdorf, K. & Schlitzer, A. Tissue-specific macrophages: how they develop and choreograph tissue biology. *Nat. Rev. Immunol.* 1–17 (2023) doi:10.1038/s41577-023-00848-y.

9. Bruttger, J. *et al.* Genetic Cell Ablation Reveals Clusters of Local Self-Renewing Microglia in the Mammalian Central Nervous System. *Immunity* **43**, 92–106 (2015).

10. Sakai, M. *et al.* Liver-Derived Signals Sequentially Reprogram Myeloid Enhancers to Initiate and Maintain Kupffer Cell Identity. *Immunity* **51**, 655-670.e8 (2019).

11. Hashimoto, D. *et al.* Tissue-Resident Macrophages Self-Maintain Locally throughout Adult Life with Minimal Contribution from Circulating Monocytes. *Immunity* **38**, 792–804 (2013).

12. Okabe, Y. & Medzhitov, R. Tissue biology perspective on macrophages. *Nature Immunology* **17**, 9–17 (2016).

13. Guilliams, M., Thierry, G. R., Bonnardel, J. & Bajenoff, M. Establishment and Maintenance of the Macrophage Niche. *Immunity* **52**, 434–451 (2020).

14. Fukada, K. & Kajiya, K. Age-related structural alterations of skeletal muscles and
associated capillaries. *Angiogenesis* **23**, 79–82 (2020).

15. Grunewald, M. *et al.* Counteracting age-related VEGF signaling insufficiency
promotes healthy aging and extends life span. *Science* **373**, (2021).

16. Pluvinage, J. V. & Wyss-Coray, T. Systemic factors as mediators of brain
homeostasis, ageing and neurodegeneration. *Nat Rev Neurosci* **21**, 93–102 (2020).

17. Shaw, A. C., Goldstein, D. R. & Montgomery, R. R. Age-dependent dysregulation of
innate immunity. *Nature reviews. Immunology* (2013) doi:10.1038/nri3547.

18. Pineda, C. M. *et al.* Intravital imaging of hair follicle regeneration in the mouse.
Nature protocols **10**, 1116–30 (2015).

19. Mesa, K. R. *et al.* Homeostatic Epidermal Stem Cell Self-Renewal Is Driven by
Local Differentiation. *Cell Stem Cell* (2018) doi:10.1016/j.stem.2018.09.005.

20. Dick, S. A. *et al.* Three tissue resident macrophage subsets coexist across organs
with conserved origins and life cycles. *Sci. Immunol.* **7**, eabf7777 (2022).

21. Siret, C. *et al.* Deciphering the heterogeneity of the Lyve1+ perivascular
macrophages in the mouse brain. *Nat. Commun.* **13**, 7366 (2022).

22. A-Gonzalez, N. *et al.* Phagocytosis imprints heterogeneity in tissue-resident
macrophages. *J. Exp. Med.* **214**, 1281–1296 (2017).

23. Dietzel, S. *et al.* Label-Free Determination of Hemodynamic Parameters in the
Microcirculaton with Third Harmonic Generation Microscopy. *PLoS ONE* **9**, e99615
(2014).

24. Saytashev, I. *et al.* Multiphoton excited hemoglobin fluorescence and third harmonic
generation for non-invasive microscopy of stored blood. *Biomedical Optics Express* **7**,
3449 (2016).

25. Bentov, I. & Reed, M. J. The effect of aging on the cutaneous microvasculature.
Microvasc. Res. **100**, 25–31 (2015).

26. Smith, L. Histopathologic characteristics and ultrastructure of aging skin. *Cutis* **43**,
414–24 (1989).

27. Li, L. *et al.* Age-Related Changes of the Cutaneous Microcirculation in vivo.
Gerontology **52**, 142–153 (2006).

28. Reeson, P., Choi, K. & Brown, C. E. VEGF signaling regulates the fate of obstructed
capillaries in mouse cortex. *eLife* **7**, e33670 (2018).

29. Das, A. *et al.* Impairment of an Endothelial NAD+-H2S Signaling Network Is a
Reversible Cause of Vascular Aging. *Cell* **173**, 74-89.e20 (2018).

30. Tsuchida, Y. The effect of aging and arteriosclerosis on human skin blood flow. *J.*
Dermatol. Sci. **5**, 175–181 (1993).

31. Rivard, A. *et al.* Age-Dependent Impairment of Angiogenesis. *Circulation* **99**, 111–
120 (1999).

32. Shimada, T. *et al.* Angiogenesis and Vasculogenesis Are Impaired in the
Precocious-Aging klotho Mouse. *Circulation* **110**, 1148–1155 (2004).

33. Abdellatif, M., Rainer, P. P., Sedej, S. & Kroemer, G. Hallmarks of cardiovascular
ageing. *Nat. Rev. Cardiol.* **20**, 754–777 (2023).

34. Uderhardt, S., Martins, A. J., Tsang, J. S., Lämmermann, T. & Germain, R. N.
Resident Macrophages Cloak Tissue Microlesions to Prevent Neutrophil-Driven
Inflammatory Damage. *Cell* **177**, 541-555.e17 (2019).

35. Freeman, S. A. *et al.* Lipid-gated monovalent ion fluxes regulate endocytic traffic
and support immune surveillance. *Science* **367**, 301–305 (2020).

36. Arandjelovic, S. & Ravichandran, K. S. Phagocytosis of apoptotic cells in
homeostasis. *Nat. Immunol.* **16**, 907–917 (2015).

37. Westman, J., Grinstein, S. & Marques, P. E. Phagocytosis of Necrotic Debris at
Sites of Injury and Inflammation. *Front. Immunol.* **10**, 3030 (2020).

38. Lämmermann, T. *et al.* Neutrophil swarms require LTB4 and integrins at sites of cell
death in vivo. *Nature* **498**, 371–375 (2013).

39. Sasmono, R. T. *et al.* Mouse neutrophilic granulocytes express mRNA encoding the
macrophage colony-stimulating factor receptor (CSF-1R) as well as many other
macrophage-specific transcripts and can transdifferentiate into macrophages in vitro in
response to CSF-1. *J. Leucoc. Biol.* **82**, 111–123 (2007).

40. Stackowicz, J., Jönsson, F. & Reber, L. L. Mouse Models and Tools for the in vivo
Study of Neutrophils. *Front. Immunol.* **10**, 3130 (2020).

41. Clausen, B. E., Burkhardt, C., Reith, W., Renkawitz, R. & Förster, I. Conditional
gene targeting in macrophages and granulocytes using LysMcre mice. *Transgenic Res.*
8, 265–277 (1999).

42. Lim, K. *et al.* In situ neutrophil efferocytosis shapes T cell immunity to influenza
infection. *Nat. Immunol.* **21**, 1046–1057 (2020).

43. Egen, J. G. *et al.* Macrophage and T Cell Dynamics during the Development and
Disintegration of Mycobacterial Granulomas. *Immunity* **28**, 271–284 (2008).

44. Abram, C. L., Roberge, G. L., Hu, Y. & Lowell, C. A. Comparative analysis of the
efficiency and specificity of myeloid-Cre deleting strains using ROSA-EYFP reporter
mice. *J. Immunol. Methods* **408**, 89–100 (2014).

45. Cox, D. *et al.* Requirements for Both Rac1 and Cdc42 in Membrane Ruffling and
Phagocytosis in Leukocytes. *J Exp Medicine* **186**, 1487–1494 (1997).

46. Perdiguero, E. & Geissmann, F. The development and maintenance of resident
macrophages. *Nat Immunol* **17**, 2–8 (2015).

47. Blériot, C., Chakarov, S. & Ginhoux, F. Determinants of Resident Tissue
Macrophage Identity and Function. *Immunity* **52**, 957–970 (2020).

48. Alfituri, O. A., Mararo, E. M., Steketee, P. C., Morrison, L. J. & Mabbott, N. A.
Dermal bacterial LPS-stimulation reduces susceptibility to intradermal Trypanosoma
brucei infection. *Sci Rep-uk* **11**, 9856 (2021).

49. Gow, D. J. *et al.* Characterisation of a Novel Fc Conjugate of Macrophage Colony-
stimulating Factor. *Mol Ther* **22**, 1580–1592 (2014).

50. Keshvari, S. *et al.* Therapeutic potential of macrophage colony-stimulating factor in
chronic liver disease. *Dis Model Mech* **15**, dmm049387 (2022).

51. Zhou, X. *et al.* Circuit Design Features of a Stable Two-Cell System. *Cell* **172**, 744-
757.e17 (2018).

52. Nicolás-Ávila, J. A. *et al.* A Network of Macrophages Supports Mitochondrial
Homeostasis in the Heart. *Cell* **183**, 94-109.e23 (2020).

53. Ferrer, I. R. *et al.* A wave of monocytes is recruited to replenish the long-term
Langerhans cell network after immune injury. *Sci Immunol* **4**, eaax8704 (2019).

54. Gilchrest, B. A., Murphy, G. F. & Soter, N. A. Effect of Chronologic Aging and
Ultraviolet Irradiation on Langerhans Cells in Human Epidermis. *J Invest Dermatol* **79**,
85–88 (1982).

55. Cruchley, A. T. *et al.* Langerhans cell density in normal human oral mucosa and
skin: relationship to age, smoking and alcohol consumption. *J Oral Pathol Med* **23**, 55–
59 (1994).

56. Hasegawa, T. *et al.* Reduction in human epidermal Langerhans cells with age is
associated with decline in CXCL14-mediated recruitment of CD14+ monocytes. *J Invest*
Dermatol (2019) doi:10.1016/j.jid.2019.11.017.

57. Fenske, N. A. & Lober, C. W. Structural and functional changes of normal aging
skin. *J Am Acad Dermatol* **15**, 571–585 (1986).

58. Cai, C. *et al.* Impaired dynamics of precapillary sphincters and pericytes at first-
order capillaries predict reduced neurovascular function in the aging mouse brain. *Nat*
Aging **3**, 173–184 (2023).

Acknowledgements

We thank members of the Littman lab for valuable discussion and critical reading of the manuscript. We thank Michael Cammer from the NYU Microscopy Core for valuable discussion, training and technical support. The Microscopy Core is partially supported by NYU Cancer Center Support Grant NIH/NCI P30CA016087 at the Laura and Isaac Perlmutter Cancer Center, S10 RR023704-01A1 and NIH S10 ODO019974-01A1. This work was supported by a Jane Coffin Childs Fund fellowship (K.R.M.), Kirschstein-NRSA training grant T32AR64184 (K.R.M.), Kirschstein-NRSA individual postdoctoral fellowship F32AG071336 (K.R.M.), a Charles H. Revson Senior Fellowship in Biomedical Science (K.R.M.), the Helen and Martin Kimmel Center for Biology and Medicine (D.R.L.), NIH grant R01AI158687 (D.R.L.), and the Howard Hughes Medical Institute (D.R.L.).

Author Contributions

K.R.M. and D.R.L. designed the study and analyzed the data; K.R.M., K.A.O., and A.D. performed mouse experiments. K.R.M. performed intravital multiphoton imaging. K.R.M and M.R.L. performed image analysis and quantification. C.N. and S.P.S. provided human biological samples and related quantitative analysis. K.R.M. and D.R.L. wrote the manuscript, with input from the other authors. D.R.L. supervised the research.

Figures

Figure 1

Figure 1 –**Niche-specific macrophage loss with age correlates with impaired skin capillary blood flow. a,** Schematic of intravital imaging of resident macrophage populations in mouse skin, indicating epidermal and dermal populations, using the macrophage reporter, *Csf1r-EGFP,* in combination with an *Actb*-driven universal tdTomato reporter, *R26-mTmG*. **b,** Representative optical sections of distinct resident macrophage populations in young (2-month-old) and old (18-month-old) mice. **c,** Quantification of niche-specific macrophage density change in 1, 2, 4, 10, and 18 month

old mice (n = 4 mice in each age group; two 500µm^2 regions per mouse; macrophage

density in each skin niche was compared across age groups by one-way ANOVA; mean

± SD). **d,** Skin resident macrophage labeling using *Cx3cr1-CreERT2;R26-mTmG* mice

was performed following a single high-dose intraperitoneal injection of tamoxifen (2mg)

in 1 month old mice. Single optical sections at successive time points 5 min apart

showing red blood cell (RBC) flow (white) in capillaries (red) with or without nearby

CAMs (green). Yellow arrowheads indicate obstructed RBC capillary flow. **e,**

Quantification of capillaries with blood flow as measured by stalled RBCs as described

in Extended Data Figure 2 (n = 226 CAM+ capillary segments, n = 27 CAM- capillary

segments; n = 4 mice; capillary blood flow (CAM+ vs CAM-) was compared by paired

Student's t test; mean ± SD). **f,** Percentage of capillary segments with at least one

associated macrophage (n = 4 mice in each age group; two 500µm^2 regions per mouse;

macrophage association with capillary segments was compared across age groups by

one-way ANOVA; mean ± SD). **g,** Representative images demonstrate macrophage

depletion following intradermal injections of clodronate-liposomes every 3 days.

Repeated intravital imaging of the vascular niche was performed to visualize

macrophages (*Csf1r-GFP*), capillaries (*R26-mTmG*) and RBC flow (Third Harmonic). **h,**

Percentage of capillaries with blood flow following macrophage depletion (n = 194

capillary segments in clodronate group, n = 199 capillary segments in PBS group; 3

mice in each group; capillary flow (clodronate vs PBS) was compared by unpaired

Student's t test; mean ± SD). Scale bar, 50 µm.

Figure 2

Figure 2 – Local recruitment of CAMs is required to restore capillary blood flow.

a, Scheme of long-term serial imaging of capillary niche in *Cx3cr1-GFP;R26-mTmG*

mice. **b,** Sequential revisits reveal progressive pruning of capillary niche over a 6-month

period. Yellow arrowheads indicate capillaries that will undergo pruning. **c,**

Quantification of capillary-associated macrophage coverage at 1 and 7 months of age

(n = 3 mice; three 500µm^2 regions per mouse; Frequency of macrophage association

with capillary segments was compared between capillaries fated for pruning and all

capillaries at 1- and 7-month-old time-points by paired Student's t test; mean ± SD). **d,**

Scheme of laser-induced capillary clot experiment. **e,** Sequential revisits of damaged

capillary segment (red dashed lines) after laser-induced clot formation in *Cx3cr1-*

CreERT2;R26-mTmG mice. Yellow lightning bolt indicates site of laser-induced capillary

clot. Yellow arrowheads indicate extra-luminal vascular debris. **f,** Quantification of

capillary-associated macrophage extension toward damaged niche at Day 1, 2 and 7

after laser-induced clotting (n = 50 CAMs, in 3 mice; mean ± SD). **g,** Sequential revisits

of damaged capillary niche after laser-induced clot formation (yellow lightning bolt) in

Cx3cr$^{gfp/+}$ and *Cx3cr$^{gfp/gfp}$* mice. CAMs (green), capillaries (red), RBC (white). Yellow

arrowheads indicate extra-luminal vascular debris. **h,** Quantification of capillary-

associated macrophage extension toward damaged niche at Day 7 after laser-induced

clotting (n = 99 CAMs in *Cx3cr$^{gfp/+}$* group; n = 65 CAMs in *Cx3cr$^{gfp/gfp}$* group; n = 3 mice

in each group; CAM extension (*Cx3cr$^{gfp/+}$* vs *Cx3cr$^{gfp/gfp}$*) was compared by unpaired

Student's t test; mean ± SD). **i,** Quantification of capillary reperfusion at Day 1, 2 and 7

after laser-induced clotting (n = 20 clots in *Cx3cr$^{gfp/+}$* group; n = 21 clots in *Cx3cr$^{gfp/gfp}$*

group; n = 3 mice in each group; capillary reperfusion (*Cx3cr$^{gfp/+}$* vs *Cx3cr$^{gfp/gfp}$*) was

compared by unpaired Student's t test; mean ± SD). **j,** Quantification of capillary blood

flow (n = 4 mice in each group; two 500µm^2 regions per mouse; capillary blood flow

(*Cx3cr$^{gfp/+}$* vs *Cx3cr$^{gfp/gfp}$*) was compared at 1 and 6 months of age by unpaired

Student's t test; mean ± SD). Scale bar, 50µm.

Figure 3

Figure 3 – CAMs are required to clear vascular damage and preserve skin

capillaries during aging. a, Sequential revisits of damaged capillary niche after laser-

induced clot formation in *Csf1r-EGFP;R26-mTmG* mice. Capillary clot formation (yellow

lightning bolt) was performed at 940nm for 1s in 10-month-old mice. Yellow arrowheads

indicate extra-luminal vascular debris. **b,** Quantification of capillary reperfusion at day 1

and 7 after laser-induced clotting (n = 16 capillary clots in regions with CAMs (<75μm

from clot), n = 16 capillary clots in regions without CAMs (>75μm from clot; 3 mice in

total; capillary reperfusion (with CAMs vs without CAMs) was compared at Day 1 and

Day 7 by paired Student's t test; mean ± SD). Scale bar, 50 μm. **c,** Sequential revisits of

damaged capillary niche after laser-induced CAM ablation and clot formation in *Csf1r-*

EGFP mice. Macrophage laser ablation (red lightning bolt) and capillary clot formation

(yellow lightning bolt) were both performed at 940nm for 1s. **d,** Quantification of capillary

reperfusion at Day 1 and 7 after laser-induced clotting and macrophage ablation (n = 19

capillary clots in CAM ablated regions, n = 16 capillary clots in control regions; 3 mice in

total; capillary reperfusion (CAM ablated vs control) was compared by paired Student's t

test; mean ± SD). **e,** Sequential revisits of damaged capillary niche after laser-induced

clot formation in $Cx3cr1^{CreER}$;$Rac1^{fl/fl}$ and $Cx3cr1^{CreER}$;$Rac1^{fl/+}$ mice. CAMs (green),

capillaries (red), RBC (white). **f,** Quantification of perivascular red blood cell debris at

Day 3 after laser-induced clotting (n = 52 clots in $Rac1^{fl/+}$ group; n = 48 clots in $Rac1^{fl/fl}$

group; n = 3 mice in each group; capillary reperfusion ($Rac1^{fl/+}$ vs $Rac1^{fl/fl}$) was

compared by unpaired Student's t test; mean ± SD). **g,** Quantification of capillary

reperfusion at Day 1, 3 and 7 after laser-induced clotting (n = 67 clots in $Rac1^{fl/+}$ group;

n = 82 clots in $Rac1^{fl/fl}$ group; n = 3 mice in each group; capillary reperfusion ($Rac1^{fl/+}$ vs

$Rac1^{fl/fl}$) was compared by unpaired Student's t test; mean ± SD). **h,** Sequential revisits

reveal pruning of capillaries over a 3-month period. Yellow arrowheads indicate

capillaries that will undergo pruning. **i,** Quantification of capillary density (n = 4 mice in

$Rac1^{fl/}$ group and 5 mice in $Rac1^{fl/fl}$ group; two 500µm^2 regions per mouse; capillary

density ($Rac1^{fl/+}$ vs $Rac1^{fl/fl}$) was compared at 3 months post tamoxifen treatment by

unpaired Student's t test; mean ± SD). **j,** Quantification of capillary density (n = 4 mice in

$Rac1^{fl/}$ group and 5 mice in $Rac1^{fl/fl}$ group; two 500µm^2 regions per mouse; capillary

density (1 week vs 3 months) was compared in $Rac1^{fl/+}$ and $Rac1^{fl/fl}$ mice by paired

Student's t test; mean ± SD). **k,** Scheme of age-associated decline of skin capillary

niche. Capillary-associated macrophage density declines with age, which predisposes

aged capillaries to impaired repair and sustained tissue perfusion. Scale bar, 50μm.

Figure 4

Figure 4 – Dermal macrophages utilize niche-specific self-renewal strategies

leading to selective CAM loss with age. a, Scheme of serial imaging of skin resident

macrophage repopulation in bone marrow chimeras with *Csf1r-GFP*;*CAG-dsRed* bone

marrow transferred into lethally irradiated *Csf1r-GFP* mice. Note hind paws of these

mice were lead-shielded to prevent irradiation-induced loss of resident macrophage

populations from our imaging area. **b,** Representative optical sections of dermal

macrophage populations in mice prior to BM transfer and 4 weeks post BM transfer.

c, Quantification of macrophage repopulation at 4- and 10-weeks post bone marrow

transplantation (n = 4 mice; two 500µm^2 regions of each skin compartment per mouse;

bone marrow-derived macrophage replacement (percentage of GFP+/dsRed+

macrophages) was compared between skin compartments at 10 weeks post BM

transplantation by paired Student's t test; mean ± SD). **d,** Scheme of long-term tracking

of dermal macrophage self-renewal experiment. **e,** Representative images in upper and

lower dermis of single macrophage tracing in *Cx3cr1-CreERT2*; *R26-mTmG* mice.

Weekly revisits were performed during homeostatic conditions for 20 weeks following a

single low-dose intraperitoneal injection of tamoxifen (50μg). **f,** Representative serial

revisit images of single macrophage tracing in *Cx3cr1-CreERT2*; *R26-mTmG* mice. **g,**

Quantification of monthly rate of dermal macrophage loss and division in upper (n = 262

macrophages; 8 mice in total) and lower (n = 42 macrophages; 4 mice in total) dermis;

macrophage monthly division and loss rates (upper vs lower dermis) were compared by

unpaired Student's t test; mean ± SD. **h,i,** Quantification of (h) monthly rate of CAM loss

and division (n = 262 CAMs; 8 mice in total; CAM turnover rates (CAM loss vs division)

was compared by paired Student's t test; mean ± SD) and of (i) maintenance of the

labeled CAM population over 20wk lineage tracing (n = 59 CAMs; 3 mice in total;

number of fluorescently-labeled CAMs was compared across each time point in the

same mice by one-way repeated measures ANOVA with Geisser-Greenhouse

correction; mean ± SD). **j,** Scheme of local self-renewal imbalance drives CAM loss

with age. In contrast to lower dermal macrophages, which are continuously replenished

by monocytes, CAMs of the upper dermis largely rely of local proliferation. However, the

rate of CAM division is insufficient to stably maintain this population with age. Scale bar,

50μm.

Figure 5

Figure 5 – Macrophage loss without local tissue damage is not sufficient to

promote CAM renewal. a, Representative revisits of CAM replacement after laser-

induced CAM ablation in *Csf1r-EGFP;R26-mTmG* mice. Macrophage laser ablations

(yellow asterisk) were performed in a 500µm² region (yellow dashed line square), **b,**

Representative revisits of CAM replacement after a single intraperitoneal injection of

diphtheria toxin (25ng/g body weight in PBS) in *Cx3cr1-DTR* mice. **c,** Representative

revisits of single lineage-tracked macrophages in *Cx3cr1-CreERT2*; *R26-dsRed*; *Csf1r-*

EGFP mice. Weekly revisits were performed during homeostatic conditions for 2 weeks

following a single low-dose intraperitoneal injection of tamoxifen (50µg). **d,**

Quantification of replenishment rate of empty CAM niche under laser-induced loss (n =

44 CAMs; 4 mice in total), DT-induced loss (n = 53 CAMs; 3 mice in total), homeostatic

loss (n = 29 CAMs; 3 mice in total), or laser-induced CAM loss with capillary clot (n = 25

CAMs; 4 mice in total). Frequency of CAM replacement after forced and homeostatic

CAM loss were compared by unpaired Student's t test; mean ± SD. **e,** Scheme of CAM

replacement after capillary niche damage in *Csf1r-EGFP*; *Ccr2^{RFP/+}* or *Ccr2^{RFP/RFP}* mice.

f, Quantification of replenishment rate (*Csf1r-EGFP+* cells) of empty CAM niche after

laser-induced CAM loss and niche damage (n = 4 mice per group; two 500µm^2 regions

per mouse; CAM density (*Ccr2^{RFP/+}* vs *Ccr2^{RFP/RFP}*) was compared by unpaired

Student's t test; mean ± SD). **g,** Representative revisits of CAM replacement after a

regional laser-induced damage to the capillary niche. **h,** Representative revisits of single

macrophage lineage tracing in *Cx3cr1-CreERT2*; *R26-nTnG* mice following capillary

damage. **i,** Quantification of CAM proliferation based on proximity to capillary damage

(n = 25 CAMs tracked in capillary damage regions, 4 mice in each group; CAM

proliferation (based on damage proximity) was compared at day 7 by paired Student's t

test; mean ± SD). Scale bar, 50µm.

Figure 6

Figure 6 – **Local CAM replenishment in old mice is sufficient to rejuvenate**

capillary repair and tissue reperfusion. a, Scheme of CSF1-induced rejuvenation of

aged capillary niche. **b,** Representative images of CAM density in *Csf1r-EGFP* mice

nine days following daily intradermal injections (4 days) of CSF1-Fc (porcine CSF1 and

IgG1a Fc region fusion protein) and PBS in contralateral hind paws of 20–24-month-old

mice. **c,** Quantification of CAM density following CSF1-Fc or PBS treatment (n = 4 mice

in total; two 500µm^2 regions of each treatment condition per mouse; Percentage of CAM

density change from Day -9 (CSF1-Fc vs PBS) was compared for Day -4 and Day 0

time points by paired Student's t test; mean ± SD). **d,** Representative images of

capillary (red dashed lines) blood flow in *Csf1r-EGFP* mice nine days following daily

intradermal injections of CSF1-Fc and PBS in contralateral hind paws of 20–24-month-

old mice. Magenta arrowheads indicate obstructed RBC capillary flow. **e,** Quantification

of capillary blood flow following CSF1-Fc or PBS treatment (n = 214 capillary segments

in CSF1-Fc treated regions (grey bar graph), n = 214 capillary segments in PBS treated

regions (white bar graph); n = 4 mice in total; capillary blood flow (CSF1-Fc vs PBS)

was compared for Day -9 and Day 0 time points by paired Student's t test; mean ± SD).

f, Sequential revisits of damaged capillary niche after laser-induced clot formation

(yellow lightning bolt) in *Csf1r-EGFP* mice. Yellow arrowheads indicate extra-luminal

vascular debris. **g,** Quantification of capillary reperfusion at Day 1 and 7 after laser-

induced clotting (n = 18 clots in CSF1-Fc group (grey bar graph); n = 20 clots in PBS

group (white bar graph); n = 4 mice in total; capillary reperfusion (CSF1-Fc vs PBS) was

compared by paired Student's t test; mean ± SD). **h,** Working model of resident

macrophage aging in the skin capillary niche. Age-associated impairment in capillary

repair can be rejuvenated following local expansion of resident macrophage population.

Scale bar, 50μm.

Methods

Mice

Mice were bred and maintained in the Alexandria Center for the Life Sciences animal

facility of the New York University School of Medicine, in specific pathogen-free

conditions. Albino B6 (B6(Cg)-*Tyr^{c-2J}*/J, Jax 000058), Csf1r^{EGFP} (B6.Cg-Tg(Csf1r-

EGFP)1Hume/J, Jax 018549), Ccr2^{RFP} (B6.129(Cg)-Ccr2tm2.1Ifc/J, Jax 017586),

R26^{mTmG} (B6.129(Cg)-Gt(ROSA)26Sortm4(ACTB-tdTomato,-EGFP)Luo/J, Jax 007676),

R26^{nTnG} (B6N.129S6-Gt(ROSA)26Sortm1(CAG-tdTomato*,-EGFP*)Ees/J, Jax 023537),

LysM^{Cre} (B6.129P2-Lyz2tm1(cre)Ifo/J, Jax 004781), Rac1$^{f/f}$ (Rac1tm1Djk/J, Jax

005550), and CAG-dsRed (B6.Cg-Tg(CAG-DsRed*MST)1Nagy/J, JAX 006051) mice

were purchased from Jackson Laboratories. *R26^{dsRed}* mice were described previously

(Luche et al., 2007) and were obtained from the laboratory of Dr. Gordon Fishell.

Cx3cr1^{CreER}, *Cx3cr1^{GFP}, and Cx3cr1^{DTR}* were generated in our laboratory and have been

described (Parkhurst, et al., 2013; Jung, et al., 2000; Diehl, et al., 2013). All

experimental mice for this study were albino (homozygous for *Tyr^{c-2J}*) as is required for

intravital imaging in the skin. Cre-induction for the lineage tracing or total CAM labeling

experiments was induced with a single intraperitoneal injection of Tamoxifen (Sigma-

Aldrich; T5648) (100µg or 4mg in corn oil, respectively) in 1 month old mice. *Rac1$^{f/f}$*

recombination was induced with two intraperitoneal injections of Tamoxifen (2mg in corn

oil) 48hrs apart in 1 month old mice. All imaging and experimental manipulations were

performed on non-hairy mouse plantar (hind paw) skin. Preparation of skin for intravital

imaging were performed as described below. Briefly, mice were anesthetized with

intraperitoneal injection of ketamine/xylazine (15 mg/ml and 1 mg/ml, respectively in

PBS). After imaging, mice were returned to their housing facility. For subsequent

revisits, the same mice were processed again with injectable anesthesia. The plantar

epidermal regions were briefly cleaned with PBS pH 7.2, mounted on a custom-made

stage, and a glass coverslip was placed directly against the skin. Anesthesia was

maintained throughout the course of the experiment with vaporized isoflurane delivered

by a nose cone. Mice from experimental and control groups were randomly selected for

live imaging experiments. All lineage tracing and ablation experiments were repeated in

at least three different mice. All animal procedures were performed in accordance with

protocols approved by the Institutional Animal Care and Usage Committee of New York

University School of Medicine.

Intravital microscopy and laser ablation

Image stacks were acquired with an Olympus multiphoton FVMPE-RS system equipped

with both InSight X3 and Mai Tai Deepsee (Spectra-Physics) tunable Ti:Sapphire

lasers, using Fluoview software. For collection of serial optical sections, a laser beam

(860nm for Hoechst 33342, 940nm for GFP/tdTomato/dsRed/RFP/Rhodamine/Second

Harmonic Generation, 1200nm for Alexa Fluor 647, and 1300nm for Third Harmonic

Generation, respectively) was focused through a water immersion lens (N.A. 1.05;

Olympus) and scanned with a field of view of 0.5mm^2, at 600 Hz. Z-stacks were

acquired in 1-2μm steps for a ~50-100μm range, covering the epidermis and dermis.

For all animal imaging, 1mm x 2 mm imaging fields (ROIs) were acquired with only the

second harmonic signal (collagen) as a reference guide to the same anatomical position

(1mm proximal of the most proximal walking pad on the mouse paw plantar skin.

Capillary blood flow was visualized in some experiments through intravenous injection

with 18 mg/kg of dextran-rhodamine 70 kD (Sigma-Aldrich; R9379). Cell tracking

analysis was performed by re-visiting the same area of the dermis in separate imaging

experiments through using inherent landmarks of the skin to navigate back to the

original region, including the distinct organization of the superficial vasculature

networks. Cells that were unambiguously separated (by at least 250μm) from another

were sampled to ensure the identity of individual lineages. For time-lapse recordings,

serial optical sections were obtained between 5–10-minute intervals, depending on the

experimental setup. Laser-induced cell ablation, capillary clot, or tissue damage was

carried out with the same optics as used for acquisition. An 940nm laser beam was

used to scan the target area (1-500μm^2) and ablation was achieved using 50-70% laser

power for ~1sec. Ablation parameters were adjusted according to the depth of the target

(10-50μm). Mice from experimental and control groups were randomly selected for live

imaging experiments. All lineage tracing and ablation experiments were repeated in at

least three different mice.

In situ staining of neutrophils for intravital microscopy

Anesthetized mice were given fluorescently labeled antibodies via intravenous

retroorbital injection immediately prior to imaging. Neutrophils were identified with 4μg of

anti-Gr1-AF647 (BioLegend; clone RB6-8C5; Cat#108418).

Drug treatments

To induce macrophage depletion, mice received intradermal injections of either

Clodronate-liposomes or PBS-liposomes (stock concentration 5mg/ml; Liposoma; CP-

005-005) (5µl per paw) every 3 days. Depending on experimental details, Cx3cr1[DTR]

mice received either intraperitoneal (IP) injection of diphtheria toxin (Sigma-Aldrich;

D0564) every other day or a single low dose at 25ng/g body weight in PBS. To induce

macrophage expansion, mice received daily intradermal injections of CSF1-FC (Bio-

Rad; PPP031) or PBS in contralateral hind paws (5µl per paw) for 4 days.

Generation of bone marrow chimeric reconstituted mice

Bone-marrow mononuclear cells were isolated from *Csf1r-GFP*;*CAG-dsRed* mice by

flushing the long bones. Red blood cells were lysed with ACK lysing buffer and the

remaining cells were resuspended in PBS for retroorbital injection. 4×10^6 cells were

then injected intravenously into 6- to 8-week-old *Csf1r-GFP* mice that were irradiated

4hr before reconstitution using 1,000 rads per mouse (2×500 rads, at an interval of

2hr, at X-RAD 320 X-Ray Irradiator). During irradiation, hind paws of recipient mice

were lead-shielded to prevent any irradiation-induced loss of resident macrophage

populations from our imaging area. At 1- and 10-months post irradiation, peripheral

blood samples were collected from the submandibular (facial) vein in tubes containing

EDTA (BD Biosciences; Dipotassium EDTA Microtainer; Cat# 365972). Red blood cells

were lysed with ACK lysing buffer and the remaining cells were subjected flow

cytometric analysis on a LSR II with FACSDiva and FlowJo 10.10.1 software (BD

Biosciences) to check for reconstitution (Extended Data Fig. 10b).

Skin whole mount staining

Whole skin was collected from the hind paw and fixed in 4% paraformaldehyde in PBS

overnight at 4°C, washed in PBS, permeabilized and blocked for 1hr (2% Triton-X, 5%

normal donkey serum and 1% BSA in PBS). For CD206 staining, blocked tissue was

incubated in Alexa Fluor 647 rat anti-CD206 (1:500; Biolegend C068C2) overnight at

4°C, washed in PBS with 2% Triton-X, washed with PBS, and then mounted on a slide

with Prolong Gold antifade mounting medium (Invitrogen) with a #1.5 coverslip. Nuclear

counterstaining was achieved by performing a single intravenous injection of Hoechst

33342 (15 mg/kg) 30 minutes prior to mouse euthanasia. Whole mount skin samples

were imaged with the same imaging conditions and setup used for intravital microscopy.

Human skin samples

Written informed consent was obtained for postmortem examination from next of kin for

all patients. Clinical information and laboratory data were obtained from the electronic

medical record. Sex and gender information was not used. Patients in young group

were below 40 years of age. Patients in old group were above 40 years of age. Patients

with skin or vascular pathologies were excluded. Skin samples were obtained from the

anterolateral chest and fixed in 10% formalin for at least 24h prior to processing. Slides

were stained with hematoxylin and eosin, CD68 (Clone 514H12) and ERG (Clone

EPR3864). Macrophages and capillaries were identified using a combination of

morphology, CD68 and ERG staining. The age group of samples were blinded to the

researcher and counting was performed on at least eight high-power fields (40x) within

100um of the epidermis.

Image Analysis

Raw image stacks were imported into Fiji (NIH, USA) or Imaris software

(Bitplane/Perkin Elmer) for further analysis. Provided images and supplementary videos

are typically presented as a maximal projection of 4-8µm optical sections. For

visualizing individual labeled cells expressing the dsRed or tdTomato Cre reporters, the

brightness and contrast were adjusted accordingly for the green (GFP) and red

(dsRed/tdTomato) channels and composite serial image sequences were assembled as

previously described. Images were obtained as large, tiled image stacks at roughly the

same positions and then manually aligned over the experimental time course in Imaris

(Bitplane/Perkin Elmer) by using data from all channels. Random ROI were selected for

image analysis. Quantification of CAM coverage and capillary blood flow was performed

blinded to minimize researcher bias.

Statistical Analysis

Data are expressed as mean ± SD. An unpaired Student's *t*-test was used to analyze

data sets with two groups. One-way ANOVA was used to analyze data sets with three

or more groups. $p < 0.05$ to $p < 0.0001$ indicated a significant difference. Statistical

calculations were performed using the Prism software package (GraphPad, USA).

Extended Data Figure Legends

Extended Data Figure 1

Extended Data Figure 1. Upper dermal macrophages are capillary-associated and decline in density with age. a, Representative images of cells expressing *Ccr2-RFP*, *Csf1r-EGFP*, or *Cx3cr1-GFP* in the epidermis, upper dermis, and lower dermis of 1 month old mice. **b,** Quantifications of cells expressing *Csf1r-EGFP*, *Cx3cr1-GFP, or Ccr2-RFP*, in the upper dermis (n = 3 mice in each group; two 500µm² regions per

mouse; Cell number (*Csf1r-EGFP* vs *Cx3cr1-GFP; Csf1r-EGFP* vs *Ccr2-RFP*) was

compared by unpaired Student's t test; mean ± SD). **c,** Visualization of CX3CR1-

expressing cells in all skin layers: epidermis, upper dermis, and lower dermis (*Cx3cr1-*

GFP;R26-mTmG) was performed during homeostatic conditions. **d,** Representative

image of labeled upper dermal macrophages in contact with capillary superficial plexus

(red) in 1 month old mice. **e,** Quantifications reveal most upper dermal macrophages

are capillary-associated macrophages (CAMs) (n = 3 mice in total; three 500µm^2

regions per mouse; mean ± SD). **f,** Quantifications of cells expressing *Cx3cr1-GFP* in

the upper dermis (n = 3 mice in each group; two 500µm^2 regions per mouse; CAM

density (1 vs 7-month-old) was compared by unpaired Student's t test; mean ± SD). **g,**

Representative optical sections of CAMs in young (1-month-old) and old (6-month-old)

LysM-Cre;R26-mTmG mice. Quantifications of membrane-GFP+ cells in the upper

dermis (n = 4 mice in each group; two 500µm^2 regions per mouse; CAM density (1 vs 6-

month-old) was compared by unpaired Student's t test; mean ± SD). **h,** Representative

upper dermal optical sections from whole mount skin samples of young (1-month-old)

and old (18-month-old) *Csf1r-EGFP* mice stained with AF647 anti-mouse CD206 (clone

C068C2) antibody. Quantifications for CD206 and CSF1R co-expression in the upper

dermis (n = 3 mice in each group; two 500µm^2 regions per mouse; CAM density (1 vs

18-month-old) was compared by unpaired Student's t test; mean ± SD). Scale bar, 50

µm.

Extended Data Figure 2. Label-free *in vivo* visualization of capillary blood flow via third harmonic effect generated by red blood cells. a-b, Simultaneous visualization of blood flow through the superficial capillary plexus via third harmonic generation (white) from red blood cells (RBC) and intravenous rhodamine dextran (RhDex) (red). **c**, Quantifications of RBC and RhDex labeling of the upper dermal superficial capillary plexus (n = 3 mice in total; four 500µm^2 regions per mouse; mean ± SD). **d,** Scheme of

red blood cell (RBC) flow through a segment of the superficial skin capillary network.

During three-dimensional image acquisition, flowing red blood cells are captured at

different x,y positions for each z-section along the capillary segment. Pseudo coloring

each z-step through a capillary segment distinguishes flowing RBCs as a multicolor

patchwork or rainbow-effect and leaves obstructed/non-flowing RBCs as white (full

overlap of all colors). **e,** Representative image of RBC visualization in the upper dermis

in *Csf1r-EGFP; R26-mTmG* mice. **f,** Representative optical z-sections through upper

dermal capillaries. Third harmonic signal is pseudo colored differently for each z-step to

visualize RBC movement along the capillary segments during image acquisition. Scale

50µm.

Extended Data Figure 3

Extended Data Figure 3. Capillary flow and associated macrophage loss in mice and humans. a, Representative images of capillary blood flow in WT mice before and after placement of a coverslip on top of hind paw skin of 6 month old mice. **b,** Quantification of capillary blood flow (n = 3 mice in total; two 500μm^2 regions per mouse; capillary blood flow (Before Coverslip vs With Coverslip) was compared in the same imaging areas by paired Student's t test; mean ± SD). **c,** Quantification of

capillaries with blood flow as measured by stalled RBCs as described in Extended Data

Figure 2 in 1, 2-, 4-, 10-, and 18-month-old mice (n = 878 CAM+ capillary segments, n =

215 CAM- capillary segments; n = 4 mice in each age group; capillary blood flow (CAM+

vs CAM-) was compared by paired Student's t test; mean ± SD). **d,** Quantification of

capillaries with blood flow in in bone marrow chimeras with *Csf1r-GFP*;*CAG-dsRed*

bone marrow transferred into lethally irradiated *Csf1r-GFP* mice (n = 207 host-derived

CAM+ capillary segments, n = 79 donor-derived CAM+ capillary segments; n = 3 mice

in total; CAM+ capillary blood flow (Host-derived vs Donor-derived) was compared by

paired Student's t test; mean ± SD). **e,f** Quantification of capillary niche age-associated

changes, (e) Capillary density and (f) CAM / Capillary segment ratio (n = 4 mice in each

age group; two 500µm^2 regions per mouse; comparison across age groups was by one-

way ANOVA; mean ± SD). **g,** Representative immunohistochemistry of CD68+

macrophage and Erg+ capillary density in the upper dermis of both young (<40y) and

old (>40y) human samples. **h-j,** Quantification of capillary niche age-associated

changes: (h) CD68+ macrophage density, (i) Erg+ capillary density, (j) Upper dermal

macrophage / Capillary endothelium ratio (n = 5 patient samples in each group; three

imaging regions per sample; Comparison by unpaired Student's t test; mean ± SD).

Scale bar, 50µm.

Extended Data Figure 4. Impaired skin capillary blood flow following acute

macrophage depletion. a, Scheme of capillary blood flow tracking following

intraperitoneal injection every other day with diphtheria toxin (DT) (25ng/g body weight

in PBS) in both *Cx3cr1-DTR*; *R26-mTmG* and WT control (*R26-mTmG*) mice. **b,**

Representative revisits of the same upper dermal capillary niches to visualize CAMs

(*Csf1r-EGFP*) and dermal collagen (Second Harmonic). **c,** Quantifications reveal a

significant reduction in CAM density following DT-treatment (n = 3 mice in each group;

two 500µm^2 regions per mouse; CAM density (WT vs *Cx3cr1-DTR*) was compared for

Day 0 and Day 5 time points by unpaired Student's t test; mean ± SD). **d,**

Representative revisits of the same capillary network to visualize capillaries (*R26-*

mTmG) and RBC flow (Third Harmonic). **e,** Quantifications reveal a significant reduction

in the percentage of capillaries with blood flow following DT-induced cell depletion (n = 3

mice in each group; two 500µm^2 regions per mouse; obstructed capillary flow (WT vs

Cx3cr1-DTR) was compared for Day 0 and Day 5 time points by unpaired Student's t

test; mean ± SD). **f,** Representative images demonstrate macrophage depletion

following intradermal injections of clodronate-liposomes every 3 days. Repeated

intravital imaging of the vascular niche was performed to visualize macrophages (*Csf1r-*

GFP), capillaries (*R26-mTmG*) and RBC flow (Third Harmonic). **g,** Quantification of

CAM density following macrophage depletion (n = 3 mice in each group; two 500µm^2

regions per mouse; CAM density (clodronate vs PBS) was compared by unpaired

Student's t test; mean ± SD). Scale bar, 50 µm.

1000

1001

Extended Data Figure 5

1002

Extended Data Figure 5. Laser-induced model of acute capillary clot formation

and repair. a, Scheme of laser-induced capillary clot experiment. **b,** Sequential revisits

of damaged capillary niche after laser-induced clot formation in *Csf1r-EGFP* mice.

Simultaneous visualization of blood flow before and after clot formation via third

harmonic generation (white) from red blood cells (RBC) and intravenous rhodamine

dextran (RhDex) (red). Yellow lightning bolt indicates site of laser-induced capillary clot.

Yellow arrowheads indicate extra-luminal vascular debris. **c,** Sequential revisits of

damaged capillary niche after laser-induced clot formation in *Csf1r-EGFP;R26-mTmG*

mice. Capillary clot formation (yellow lightning bolt) was performed at 940nm for 1s in 4-

month-old mice. **d,** Quantification of capillary lumen diameter before and immediately

after laser-induced clotting (n = 24 capillary clots; 3 mice in total; capillary luminal

diameter (pre-ablation vs post-ablation) was compared by paired Student's t test; mean

± SD). Scale bar, 50 µm.

1016

1017

1018

1019

1020

1021

1022

1023

1024

1025

1026

1027

1028

1029

1030

1031

1032

Extended Data Figure 6. Laser-induced clots and CAM ablations do not recruit

neutrophil swarms. a, Scheme of tracking neutrophil recruitment and swarming after

laser-induced capillary clot experiment. **b-c,** (b) Sequential revisits of damaged capillary

niche after laser-induced CAM ablation and clot formation or (c) large non-sterile tissue

damage (28-gauge needle stick) in *Csf1r-EGFP* mice with intravenous neutrophil

antibody labeling (Gr-1, clone RB6-8C5). Macrophage laser ablation (red lightning bolt)

and capillary clot formation (yellow lightning bolt) were both performed at 940nm for 1s.

Needle injury (yellow dashed line) was performed through epidermis and dermis. **d,**

Quantification of neutrophil swarming at 6 hours tissue damage (n = 16 capillary clots in

CAM ablated regions, n = 9 needle injuries; 3 mice in each group; mean ± SD). **e,**

Quantification of intravascular recruitment of neutrophil and monocyte populations at 0,

6, and 24 hours post laser-induced capillary clotting in CAM ablated regions (n = 16

capillary clots in CAM ablated regions, 3 mice in total; mean ± SD). **f,** Scheme of

tracking neutrophil recruitment and swarming after laser-induced capillary clot

experiment. **g,** Sequential revisits of damaged capillary niche after laser-induced CAM

ablation and clot formation in *LysMCre;R26-dsRed* mice. Macrophage laser ablation

(red lightning bolt) and capillary clot formation (yellow lightning bolt) were both

performed at 940nm for 1s. **h,** Quantification of neutrophil swarming at 6 hours tissue

damage (n = 32 capillary clots in CAM ablated regions; 3 mice in total; mean ± SD).

Scale bar, 50 μm.

1053

1054

1055

1056

1057

1058

1059

1060

1061

1062

Extended Data Figure 7. Laser ablation of non-macrophage perivascular dermal

cells does not impair capillary clot repair and reperfusion. a, Scheme of laser-

induced capillary clot experiment. **b,** Sequential revisits of damaged capillary niche after

laser-induced perivascular dermal cell ablation and clot formation in *Csf1r-EGFP*;*CAG-*

dsRed mice. Perivascular dermal cell laser ablation (blue lightning bolt) and capillary

clot formation (yellow lightning bolt) were both performed at 940nm for 1s. **c,**

Quantification of capillary reperfusion at Day 1 and 7 after laser-induced clotting and

perivascular cell ablation (n = 34 capillary clots in CAM ablated regions, n = 37 capillary

clots in control regions; 3 mice in total; capillary reperfusion (perivascular dermal cell

ablation vs control) was compared by paired Student's t test; mean ± SD). Scale bar, 50

μm.

1074

1075

1076

1077

1078

1079

1080

1081

1082

1083

1084

1085

1086

1087

1088

1089

1090

1091

Extended Data Figure 8

1092

Extended Data Figure 8. CAM density in mice with deficiencies in macrophage

phagocytic machinery. a, Scheme of laser-induced capillary clot in

Cx3cr1^{CreER};Rac1$^{fl/fl}$ and *Cx3cr1^{CreER};Rac1$^{fl/+}$* mice. **b,** Quantification of CAM density at

Day 7 after laser-induced clotting (n = 3 mice in each group; two 500μm^2 regions per

mouse; CAM density (*Rac1$^{fl/+}$* vs *Rac1$^{fl/fl}$*) was compared at day 0 (same day of clot

induction) by unpaired Student's t test; mean ± SD).

Extended Data Figure 9

1099

Extended Data Figure 9. CAM loss and disorganization is partially restored by

local epidermal damage to improve capillary repair in old mice. a, Scheme of

tracking of CAM proliferation following laser-induced damage to nearby capillary or

epidermal niches. **b,** Representative revisits of single macrophage lineage tracing in

Cx3cr1-CreERT2; *R26-nTnG* mice following epidermal damage. **c,** Quantification of

CAM proliferation based on proximity to epidermal damage (n = 25 CAMs tracked in

capillary damage regions, n = 56 CAMs tracked in epidermal damage regions; 4 mice in

each group; CAM proliferation (based on damage proximity) was compared at day 7 by

paired Student's t test; mean ± SD). **d,** Scheme of long-term tracking of CAM migration

following cell division. **e,** Representative revisits of single macrophage lineage tracing in

Cx3cr1-CreERT2; *R26-nTnG* mice. Weekly revisits were performed during homeostatic

conditions for 5 weeks following a single low-dose intraperitoneal injection of tamoxifen

(50µg). **f,** Quantification of neighboring CAM distance of recently divided sister CAMs

was compared to total CAM neighboring distance from *Cx3cr1-CreERT2*; *R26-nTnG*

mice given a single high-dose intraperitoneal injection of tamoxifen (4mg) (n = 26 sister

CAM pairs, from 5 mice in low-dose group; n = 188 CAMs, from 4 mice in high-dose

group; Average nearest neighbor distance (2 vs 10 month old) was compared at 0-25,

26-50, 51-75, and >75µm intervals by unpaired Student's t test mean ± SD). **g,**

Quantification of distance between nearest CAM neighbors in 2- and 10-month-old

Csf1r-EGFP mice (n = 90 CAMs in 2-month-old mice, n = 101 CAMs in 10-month-old

mice; 3 mice in each age group; Frequency of CAM distribution (2 vs 10 month old) was

compared at 0-25, 26-50, 51-75, and >75µm intervals by unpaired Student's t test;

mean ± SD). **h,** Working model of macrophage renewal and organization in the skin

aging capillary niche. **i,** Scheme of local damage-induced expansion of CAMs in the

aged capillary niche. **j,** Representative images of CAM density in *Csf1r-EGFP; R26-*

mTmG mice 28 days following overlaying epidermal laser damage. **k,** Quantification of

CAM density following overlaying epidermal damage (n = 4 mice in total; two 500µm^2

regions for both epidermal damage and control conditions per mouse; CAM density

(Epidermal damage vs Control) was compared by paired Student's t test; mean ± SD). **l,**

Quantification of capillary reperfusion at day 1 and 7 after laser-induced clotting (n = 17

capillary clots in epidermal damage regions, n = 17 capillary clots in neighboring control

regions; 4 mice in total; capillary reperfusion (with epidermal damage vs control) was

compared at day 1 and day 7 by paired Student's t test; mean ± SD). following laser-

induced damage to overlaying epidermal niche. Scale bar, 50µm.

1134

1135

1136

1137

1138

1139

1140

1141

1142

1143

1144

1145

1146

1147

1148

1149

1150

1151

Extended Data Figure 10

1152

Extended Data Figure 10. Intradermal CSF1 treatment locally expands CAM

populations in old mice. a, Scheme of serial imaging of skin resident macrophage

repopulation in bone marrow chimeras with *Csf1r-GFP*;*CAG-dsRed* bone marrow

transferred into lethally irradiated *Csf1r-GFP* mice following CSF1-treatment. Note hind

paws of these mice were lead-shielded to prevent irradiation-induced loss of resident

macrophage populations from our imaging area. **b,** Quantification of blood chimerism 10

months after *Csf1r-GFP*;*CAG-dsRed* bone marrow transfer into lethally irradiated *Csf1r-*

GFP mice (n = 3 mice in each group; Percentage of cells from blood (*Csf1r-GFP*;*CAG-*

dsRed mice vs bone marrow transferred mice) was compared for dsRed-/GFP+,

dsRed+/GFP+, dsRed+/GFP-, dsRed-/GFP- populations by unpaired Student's t test;

mean ± SD). **c,** Representative images of CAM density 14 days following daily

intradermal injections (4 days) of CSF1-Fc (porcine CSF1 and IgG1a Fc region fusion

protein) and PBS in contralateral hind paws of 12-month-old mice. **d,** Quantification of

capillary-associated macrophage density following CSF1-Fc or PBS treatment (n = 3

mice in total; two 500µm^2 regions of each treatment condition per mouse; Percentage of

CAM density change (Day 0 vs 14) was compared for CSF1-Fc and PBS regions by

paired Student's t test; mean ± SD). **e,** Quantification of donor bone marrow-derived

(GFP+/dsRed+) CAM density following CSF1-Fc or PBS treatment (n = 3 mice in total;

two 500µm^2 regions of each treatment condition per mouse; Percentage of donor bone

marrow-derived CAM density change (Day 0 vs 14) was compared for CSF1-Fc and

PBS regions by paired Student's t test; mean ± SD). Scale bar, 50µm.

1174

1175

1176

1177

1178

1179

1180

1181

1182   **Supplemental Video Legends**

1183   **Supplemental Video 1.** Serial optical sections through mouse plantar skin, including

1184   epidermis (0-25µm), upper dermis (26-50µm), and lower dermis (51-100µm) in *R26-*

1185   *mTmG* (red) mice. Note Second and Third Harmonic Generation illuminates dermal

1186   collagen (blue) and red blood cells (white), respectively. Scale bar, 50µm.

1187

1188   **Supplemental Video 2.** Serial optical sections through mouse plantar skin, including

1189   epidermis (0-15µm), upper dermis (16-40µm), and lower dermis (41-52µm) in *Csf1r-*

1190   *EGFP* mice. Note distinct macrophage (green) density and morphology in each tissue

1191   niche. Scale bar, 50µm.

1192

1193   **Supplemental Video 3.** Time-lapse recording of capillary blood flow via third harmonic

1194   generation (white) from red blood cells and intravenous rhodamine dextran (red) in

1195   *Csf1r-EGFP* mice. Note some capillary segments show obstructed red blood cell and

1196   rhodamine dextran flow. Scale bar, 50µm.

1197

1198   **Supplemental Video 4.** Serial optical sections through upper dermal capillary niche in

1199   *Cx3cr1-GFP; R26-mTmG* mice. Note macrophage (green) localization and morphology

1200   around capillary (red) network. Scale bar, 50µm.

1201

1202   **Supplemental Video 5.** Time-lapse recording of capillary blood flow in *Cx3cr1-CreER;*

1203   *R26-mTmG* mice. Note inconsistent and obstructed blood flow (white) in capillary (red)

1204   segments without associated macrophages (green). Scale bar, 50µm.

1205

1206   **Supplemental Video 6.** Time-lapse recording following laser-induced capillary clot

1207   formation in *Csf1r-GFP; R26-mTmG* mice. Note rapid migration of neighboring capillary-

1208   associated macrophages (green) toward site of clot formation (yellow arrowhead). Scale

1209   bar, 50μm.

1210

1211

1212

1213

1214

1215

1216

1217

1218

1219

1220

1221

1222

1223

1224

1225

1226

1227