Niche-specific macrophage loss promotes skin capillary aging

14 Kailin R. Mesa[1*], Kevin A. O'Connor[1], Charles Ng[2], Steven P. Salvatore[2], Dan R.
15 Littman[1,3,4*]
16
17
[1] Department of Cell Biology, New York University School of Medicine, New York, NY
10016, USA
[2] Department of Pathology and Laboratory Medicine, Weill Cornell Medicine, New York-
Presbyterian Hospital, New York, NY 10021, USA
[3] Perlmutter Cancer Center, New York University Langone Health, New York, NY
10016, USA
[4] Howard Hughes Medical Institute, New York, NY 10016, USA
30 * To whom correspondence should be addressed
31
32 Email: kai.mesa@med.nyu.edu or dan.littman@med.nyu.edu
33 Tel: 212-263-7520

41 Key Words: Aging / Resident Macrophages / Blood Capillaries / Tissue Repair /
42 Intravital Microscopy
43
44
45
46 Running Title: Niche-specific macrophage loss promotes capillary aging

47   **Summary**

48
All mammalian organs depend upon resident macrophage populations to coordinate

repair processes and facilitate tissue-specific functions[1–3]. Recent work has

established that functionally distinct macrophage populations reside in discrete tissue

niches and are replenished through some combination of local proliferation and

monocyte recruitment[4,5]. Moreover, decline in macrophage abundance and function in

tissues has been shown to contribute to many age-associated pathologies, such as

atherosclerosis, cancer, and neurodegeneration[6–8]. Despite these advances, the

cellular mechanisms that coordinate macrophage organization and replenishment

within an aging tissue niche remain largely unknown. Here we show that capillary-

associated macrophages (CAMs) are selectively lost over time, which contributes to

impaired vascular repair and tissue perfusion in older mice. To investigate resident

macrophage behavior *in vivo*, we have employed intravital two-photon microscopy to

non-invasively image in live mice the skin capillary plexus, a spatially well-defined

model of niche aging that undergoes rarefication and functional decline with age. We

find that CAMs are lost with age at a rate that outpaces that of capillary loss, leading to

the progressive accumulation of capillary niches without an associated macrophage in

both mice and humans. Phagocytic activity of CAMs was locally required to repair

obstructed capillary blood flow, leaving macrophage-less niches selectively vulnerable

to both homeostatic and injury-induced loss in blood flow. Our work demonstrates that

homeostatic renewal of resident macrophages is not as finely tuned as has been

previously suggested[9–11]. Specifically, we found that neighboring macrophages do not

proliferate or reorganize sufficiently to maintain an optimal population across the skin

capillary niche in the absence of additional cues from acute tissue damage or increased

abundance of growth factors, such as colony stimulating factor 1 (CSF1). Such

limitations in homeostatic renewal and organization of various niche-resident cell types

are potentially early contributors to tissue aging, which may provide novel opportunities

for future therapeutic interventions.

79   **Main Text**

Tissue homeostasis is dependent on multiple macrophage populations that

reside in distinct sub-tissue compartments or niches, such as epithelium, blood vessels

or nerves, and are thought to support specialized tissue functions[4,12,13]. Recent work

has suggested that functional decline and rarefication of vascular niches may contribute

to various age-associated tissue pathologies (including sarcopenia, chonic wounds, and

Alzheimer's disease)[14–16]. It is not yet known how tissue-resident macrophages resist or

potentiate such niche specific aging processes[6,17].

Capillary-associated macrophage decline with age correlates with impaired blood

flow.

To model mammalian tissue aging, we adapted an intravital microscopy

technique to visualize skin resident macrophage populations non-invasively in live mice

throughout the lifetime of the organism[18,19] (Figure 1a, Supplemental Videos 1 and 2).

Unexpectedly, longitudinal imaging of skin macrophages (marked by *Csf1r-EGFP*)

revealed niche-specific decline in macrophage populations. Subdividing the skin into

three anatomical layers, epidermis, upper (papillary) and lower (reticulated) dermis, we

observed that macrophages of the upper dermis were lost with age at a greater rate

than macrophages from both epidermis and lower dermis (Figure 1b-c). Further

characterization of this population with additional myeloid fluorescent reporters revealed

that most macrophages of the upper dermis express the chemokine receptor CX_3CR1,

but not CCR2 (Extended Data Fig. 1a,b). A major component of the upper dermal niche

is the superficial capillary plexus, which supplies nutrient exchange for the overlying

epidermis. To visualize this structure, we utilized third harmonic generation from our

imaging to track red blood cell (RBC) flow[20,21], which was consistent with conventional

labeling methods such as rhodamine dextran injection (Extended Data Fig. 1c-e and

Supplemental Video 3). With this *in vivo* marker of blood flow, we found that the

macrophages were closely associated with blood capillaries of the superficial plexus,

suggesting that they may provide support for this capillary niche (Extended Data Fig. 1f-

h and Supplemental Video 4). To investigate if these macrophages play a role in

capillary function, we assessed if blood flow, via RBC flow (Extended Data Fig. 2), was

altered in the presence of capillary-associated macrophages (CAMs). Performing

timelapse recordings of fluorescently labeled CAMs in mice with a cre-dependent dual

reporter system (*Cx3cr1-CreERT2; R26-mTmG*), we found that capillaries lacking an

associated macrophage had a higher rate of obstructed blood flow (Figure 1d,e and

Supplemental Video 5). Longitudinal imaging across multiple ages showed that the

fraction of capillaries with CAMs significantly decreased with age (Figure 1f). This

decrease in CAMs and coverage outpaced the loss of capillaries (Extended Data Fig.

3a,b), which was previously shown to be an early hallmark of aging in multiple tissues,

including central nervous system, lung, kidney, and skin[14,22–26]. To assess if this

phenomenon also occurs in humans, we obtained both young and old human patient

skin samples. Consistent with our observations in mice, human capillary-associated

macrophages also displayed a decline with age. Moreover, CAM decline also outpaced

capillary loss with age, suggesting a similar loss in macrophage coverage of the

capillary niche (Extended Data Fig. 3c-f). To assess any functional role of CAMs in

maintaining capillary blood flow, we performed chemical and genetic ablation of CAMs,

via clodronate liposomes and *Cx3cr1-DTR* depletion, respectively, and observed an

acute loss in capillary flow (Figure 1 g,h and Extended Data Fig. 4). Collectively, this

work highlights an evolutionarily conserved loss in skin capillary-associated

macrophages with age, which correlates with impaired homeostatic capillary perfusion.

Local CAM recruitment and phagocytosis is required to restore blood flow.

We next aimed to assess the cellular mechanism(s) by which CAMs support

capillary blood flow, using a laser-induced blood clotting model to precisely target and

stop blood flow in individual capillary segments (Figure 2a, Extended Data Fig. 5a,b and

Supplemental Video 6). To assess any macrophage involvement, we tracked the daily

displacement of surrounding CAMs to laser-induced clots. These data showed that

CAM recruitment to sites of capillary damage as well as RBC engulfment are locally

restricted to approximately 80-100μm, and largely occur within the first two days after

injury (Figure 2b,c). Under homeostatic conditions, neighboring CAMs were similarly

involved locally, taking up RBC debris from naturally occurring capillary clots (Figure

2d).

Given these findings, we next tested if capillary blood flow was preferentially

repaired when at least one resident CAM was in close proximity (<75μm) with the

injured capillary segment. The results showed a significant improvement in capillary

repair for vessels with an associated macrophage (Extended Data Fig. 5c-e),

suggesting that the loss of CAMs could lead to compromised capillary blood flow with

age. To rule out an indirect association of capillary repair and CAM proximity, we

performed laser-induced ablation of CAMs adjacent to capillary segments immediately

prior to inducing capillary clots. Indeed, in regions with CAM ablation within 75μm of the

capillary clot, repair was significantly impaired and blood flow was not properly

reestablished (Figure 2e,f).

Our serial imaging revealed that CAMs at capillary repair sites often contained

RBC debris shortly after clot formation. To understand if CAM uptake and clearance of

this cellular debris is functionally important for capillary repair, we acutely impaired CAM

phagocytosis through inducible cre-dependent knockout of Rac1, a critical component

of the phagocytic machinery[27], one week prior to laser-induced clot formation. While

there was no significant change in CAM density (Extended Data Fig. 6a,b), we found

significant impairment in capillary repair in *Cx3cr1[CreER];Rac1[fl/fl]* mice compared to

Cx3cr1[CreER];Rac1[fl/+] littermate controls (Figure 2g,h), suggesting that Rac1-dependent

phagocytic clearance is critical for proper capillary repair and tissue reperfusion.

Together, our results support a model in which local CAM recruitment and phagocytic

clearance of capillary debris is critical to maintain capillary function. Therefore, as CAM

density declines with age, so does capillary perfusion of the tissue (Figure 2i).

CAMs do not replenish lost neighbors, resulting in population loss with age

Maintenance of tissue resident macrophage populations in the skin is thought to

be mediated through a combination of local proliferation and systemic replacement by

blood monocytes[13,28,29]. To understand how CAM density declines with age, we

performed single-macrophage lineage tracing, by inducing sparse cre-recombination to

label and tracking of individual macrophages over weekly revisits (Extended Data Fig.

7a-c). In doing so, we observed discrete behaviors, including macrophage loss and

proliferation (Figure 3a). Analysis of weekly rate of CAM loss and division revealed a

significant skew toward more CAM loss than division in the first 4 months of age (Figure

3b). Consistent with these data, there was a significant decline in the fraction of fate-

mapped CAMs over a 20-week time course (Figure 3c). These results demonstrate that

the CAMs are insufficiently replenished by local proliferation, which contributes to their

progressive decline in this tissue niche.

The ability of resident macrophages to locally self-renew has largely been

studied through methods of near-total macrophage depletion, which have limited niche-

specificity and often generate tissue-wide inflammation[9–11]. To directly interrogate the

steps of macrophage self-renewal over time, we tracked both the replacement of

individual macrophages after loss and the distribution of sister macrophages after

division. First, to track local macrophage replacement after CAM loss, we performed

laser-induced ablation of all CAMs within a defined 500µm^2 region. Serial revisits up to

two weeks after ablation revealed minimal repopulation from adjacent capillary regions

that retained intact CAM populations (Extended Data Fig. 8a and Figure 3d). We found

a similar lack in repopulation following partial CAM depletion in *Cx3cr1^{DTR}* mice

following low dose diphtheria toxin administration (Figure 3e). To avoid any non-

physiological effects from these cell depletion models, we developed a dual fluorescent

macrophage reporter mouse, *Cx3cr1-CreERT2; Rosa26-dsRed; Csf1r-EGFP*, allowing

for cre-dependent recombination to differentially label a small fraction of macrophages

and track their homeostatic replacement. Examination on serial weekly revisits indicated

that, as we observed in the depletion models, most capillary niches did not recruit a new

macrophage for as long as two weeks following CAM loss (Figure 3f). In contrast,

emptied capillary niches were readily replenished with new CAMs when laser-induced

macrophage loss was accompanied by laser-induced capillary damage (Figure 3g).

Such replenishment was partially CCR2-dependent, suggestive that monocytes may

participate in repopulating the capillary niche after injury (Extended Data Fig. 8b-d and

Figure 3g). Taken together, these results suggest that CAM loss is not a sufficient

trigger to promote neighboring macrophages into the emptied niche.

Second, to precisely track sister macrophage migration following cell division, we

studied mice with a dual fluorescent nuclear reporter, *Cx3cr1-CreERT2; R26-nTnG*,

where we can use low or high doses of tamoxifen to either label individual or all CAMs,

respectively (Extended Data Fig. 9a). Compared to the average distance between all

neighboring macrophages, sister CAMs remained significantly closer to each other even

2 weeks after division (Extended Data Fig. 9b,c). These findings further support the

notion that CAM division and loss are not spatiotemporally coupled, which we predict

would progressively lead to disorganized patterning and the accumulation of both empty

and crowded capillary niches. We tested this prediction by looking at the distribution of

neighboring CAMs in both young (2-month old) and old (10-month old) mice. In young

mice, the majority of macrophages were within 50μm of each other. In contrast, old mice

had a biphasic distribution of macrophage patterning, with most CAMs either within

25μm or further than 75μm apart (Extended Data Fig. 9d). These results highlight two

distinct cellular features that contribute to reduced CAM coverage with age: 1)

insufficient macrophage repopulation following CAM loss, and 2) insufficient distribution

of these cells along the capillary niche, which may promote progressive erosion of the

vascular bed (Extended Data Fig. 9e).

Local CAM replenishment in old mice is sufficient to rejuvenate capillary repair

and tissue reperfusion

While CAM proliferation was insufficient for maintenance of the population during

homeostasis, we asked if extrinsic cues such as broader tissue damage could enhance

division rates. We employed large laser-induced wounds in both the epidermal and

upper dermal niches (Extended Data Fig. 10a) and found that local CAM proliferation

was significantly increased one week following tissue damage in both layers (Extended

Data Fig. 10b-e). Given these results we hypothesized that damage-induced CAM

proliferation in the aged capillary niche would provide lasting increases in macrophage

density. Indeed, broad epidermal damage resulted in a lasting increase in CAMs in

damaged regions compared to neighboring control regions (Extended Data Fig. 10f,h).

Our results suggest that the capillary-associated macrophage population in old

mice can be stably expanded following environmental changes, such as injury.

Therefore, we next assessed if increased CAM density, without previous injury, would

be sufficient to improve future capillary repair and reperfusion. To this end, we utilized a

fusion protein of the canonical macrophage growth factor, CSF1, with the Fc region of

porcine IgG, as it has been reported to robustly increase macrophage density in multiple

tissues, including skin[30–32]. We performed daily intradermal injections of either CSF1-Fc

or PBS in the left or right hind paws, respectively, of the same mice (Figure 4a). There

was a significant increase in CAMs in the CSF1-treated paws compared to contralateral

PBS controls, which had no significant change from pretreatment (Figure 4b,c).

Strikingly, we found that CSF1 treatment was sufficient to improve homeostatic capillary

blood flow in old mice, as compared to PBS controls which had significantly more

obstructed capillary segments (Figure 4d,e). Utilizing the same aged mice, we next

tested if this increase in CAM density would be sufficient to improve capillary repair

rates. Following laser-induced clotting, there was a significant improvement in capillary

repair and reperfusion in CSF1-treated mice compared to PBS controls (Figure 4 f,g).

Consistent with this result, there was also significant improvement in capillary repair and

tissue reperfusion in old mice following damage-induced CAM expansion (Extended

Data Fig. 10i).

Discussion

Macrophage renewal has largely been studied in non-physiological settings, such

as through *in vitro* cell culture or severe depletion models that often are accompanied

by acute inflammation[9,10,33–35]. Our work clearly demonstrates that homeostatic renewal

of resident macrophages is not as finely tuned as has been previously suggested.

Specifically, we found that neighboring macrophages do not proliferate or reorganize

sufficiently to maintain an optimal population across the skin capillary niche unless they

receive additional cues from acute tissue damage or increased growth factor

abundance. Interestingly, we confirmed previous findings that show epidermal

Langerhans cell density also declines with age, which has been associated with

impaired epidermal function[36–39]. This raises the possibility that age-associated loss in

macrophage density is a more general phenomenon in populations that rely on local

self-renewal.

In addition to self-renewal, we also found that CAM recruitment to repair tissue

damage was spatially restricted. To our knowledge, the long-term size and stability of

resident macrophage territories or niches *in vivo* has not been reported. Our work

provides strong evidence that injury-induced macrophage recruitment is restricted to

approximately 100μm. In young mice, CAM density is high enough to provide

substantial niche/territory overlap between neighbors. However, with declining CAM

density with age, we show that a significant fraction of the skin capillary network is no

longer within a CAM's territory range. It will be important to understand how this

property of CAMs is influenced by other aspects of regional heterogeneity, such as

innate immune imprinting[13], to shape local immune responses in tissues.

Collectively, this work demonstrates that loss in CAMs: 1) begins within the first

few months of life, 2) is progressive throughout life, and 3) is functionally detrimental to

vascular function, which has been shown to be a primary driver of age-associated tissue

impairments[15,40]. Furthermore, this work provides a novel platform to investigate age-

associated deviations in tissue homeostasis at the single-cell level in a living mammal.

References

1. Nobs, S. P. & Kopf, M. Tissue-resident macrophages: guardians of organ homeostasis. *Trends Immunol* **42**, 495–507 (2021).

2. Minutti, C. M., Knipper, J. A., Allen, J. E. & Zaiss, D. Tissue-specific contribution of macrophages to wound healing. *Semin Cell Dev Biol* **61**, 3–11 (2017).

3. Vannella, K. M. & Wynn, T. A. Mechanisms of Organ Injury and Repair by Macrophages. *Annual Review of Physiology* **79**, (2016).

4. Chakarov, S. *et al.* Two distinct interstitial macrophage populations coexist across tissues in specific subtissular niches. *Science* **363**, eaau0964 (2019).

5. Wu, Y. & Hirschi, K. K. Tissue-Resident Macrophage Development and Function. *Frontiers Cell Dev Biology* **8**, 617879 (2021).

6. Beek, A. A. van, Bossche, J. den, Mastroberardino, P. G., Winther, M. de & Leenen, P. Metabolic Alterations in Aging Macrophages: Ingredients for Inflammaging? *Trends Immunol* (2019) doi:10.1016/j.it.2018.12.007.

7. Franceschi, C., Garagnani, P., Vitale, G., Capri, M. & Salvioli, S. Inflammaging and 'Garb-aging.' *Trends Endocrinol Metabolism* **28**, 199–212 (2017).

8. Mass, E., Nimmerjahn, F., Kierdorf, K. & Schlitzer, A. Tissue-specific macrophages: how they develop and choreograph tissue biology. *Nat. Rev. Immunol.* 1–17 (2023) doi:10.1038/s41577-023-00848-y.

9. Bruttger, J. *et al.* Genetic Cell Ablation Reveals Clusters of Local Self-Renewing Microglia in the Mammalian Central Nervous System. *Immunity* **43**, 92–106 (2015).

10. Sakai, M. *et al.* Liver-Derived Signals Sequentially Reprogram Myeloid Enhancers to Initiate and Maintain Kupffer Cell Identity. *Immunity* **51**, 655-670.e8 (2019).

11. Hashimoto, D. *et al.* Tissue-Resident Macrophages Self-Maintain Locally throughout Adult Life with Minimal Contribution from Circulating Monocytes. *Immunity* **38**, 792–804 (2013).

12. Okabe, Y. & Medzhitov, R. Tissue biology perspective on macrophages. *Nature Immunology* **17**, 9–17 (2016).

13. Guilliams, M., Thierry, G. R., Bonnardel, J. & Bajenoff, M. Establishment and Maintenance of the Macrophage Niche. *Immunity* **52**, 434–451 (2020).

14. Fukada, K. & Kajiya, K. Age-related structural alterations of skeletal muscles and
associated capillaries. *Angiogenesis* **23**, 79–82 (2020).

15. Grunewald, M. *et al.* Counteracting age-related VEGF signaling insufficiency
promotes healthy aging and extends life span. *Science* **373**, eabc8479 (2021).

16. Pluvinage, J. V. & Wyss-Coray, T. Systemic factors as mediators of brain
homeostasis, ageing and neurodegeneration. *Nat Rev Neurosci* **21**, 93–102 (2020).

17. Shaw, A. C., Goldstein, D. R. & Montgomery, R. R. Age-dependent dysregulation of
innate immunity. *Nature reviews. Immunology* (2013) doi:10.1038/nri3547.

18. Pineda, C. M. *et al.* Intravital imaging of hair follicle regeneration in the mouse.
Nature protocols **10**, 1116–30 (2015).

19. Mesa, K. R. *et al.* Homeostatic Epidermal Stem Cell Self-Renewal Is Driven by
Local Differentiation. *Cell Stem Cell* (2018) doi:10.1016/j.stem.2018.09.005.

20. Dietzel, S. *et al.* Label-Free Determination of Hemodynamic Parameters in the
Microcirculaton with Third Harmonic Generation Microscopy. *PLoS ONE* **9**, e99615
(2014).

21. Saytashev, I. *et al.* Multiphoton excited hemoglobin fluorescence and third harmonic
generation for non-invasive microscopy of stored blood. *Biomedical Optics Express* **7**,
3449 (2016).

22. Bentov, I. & Reed, M. J. The effect of aging on the cutaneous microvasculature.
Microvasc Res **100**, 25–31 (2015).

23. Smith, L. Histopathologic characteristics and ultrastructure of aging skin. *Cutis* **43**,
414–24 (1989).

24. Li, L. *et al.* Age-Related Changes of the Cutaneous Microcirculation in vivo.
Gerontology **52**, 142–153 (2006).

25. Reeson, P., Choi, K. & Brown, C. E. VEGF signaling regulates the fate of obstructed
capillaries in mouse cortex. *eLife* **7**, e33670 (2018).

26. Das, A. *et al.* Impairment of an Endothelial NAD+-H2S Signaling Network Is a
Reversible Cause of Vascular Aging. *Cell* **173**, 74-89.e20 (2018).

27. Cox, D. *et al.* Requirements for Both Rac1 and Cdc42 in Membrane Ruffling and
Phagocytosis in Leukocytes. *J Exp Medicine* **186**, 1487–1494 (1997).

28. Perdiguero, E. & Geissmann, F. The development and maintenance of resident
macrophages. *Nat Immunol* **17**, 2–8 (2015).

29. Blériot, C., Chakarov, S. & Ginhoux, F. Determinants of Resident Tissue
Macrophage Identity and Function. *Immunity* **52**, 957–970 (2020).

30. Alfituri, O. A., Mararo, E. M., Steketee, P. C., Morrison, L. J. & Mabbott, N. A.
Dermal bacterial LPS-stimulation reduces susceptibility to intradermal Trypanosoma
brucei infection. *Sci Rep-uk* **11**, 9856 (2021).

31. Gow, D. J. *et al.* Characterisation of a Novel Fc Conjugate of Macrophage Colony-
stimulating Factor. *Mol Ther* **22**, 1580–1592 (2014).

32. Keshvari, S. *et al.* Therapeutic potential of macrophage colony-stimulating factor in
chronic liver disease. *Dis Model Mech* **15**, dmm049387 (2022).

33. Zhou, X. *et al.* Circuit Design Features of a Stable Two-Cell System. *Cell* **172**, 744-
757.e17 (2018).

34. Nicolás-Ávila, J. A. *et al.* A Network of Macrophages Supports Mitochondrial
Homeostasis in the Heart. *Cell* **183**, 94-109.e23 (2020).

35. Ferrer, I. R. *et al.* A wave of monocytes is recruited to replenish the long-term
Langerhans cell network after immune injury. *Sci Immunol* **4**, eaax8704 (2019).

36. Gilchrest, B. A., Murphy, G. F. & Soter, N. A. Effect of Chronologic Aging and
Ultraviolet Irradiation on Langerhans Cells in Human Epidermis. *J Invest Dermatol* **79**,
85–88 (1982).

37. Cruchley, A. T. *et al.* Langerhans cell density in normal human oral mucosa and
skin: relationship to age, smoking and alcohol consumption. *J Oral Pathol Med* **23**, 55–
59 (1994).

38. Hasegawa, T. *et al.* Reduction in human epidermal Langerhans cells with age is
associated with decline in CXCL14-mediated recruitment of CD14+ monocytes. *J Invest
Dermatol* (2019) doi:10.1016/j.jid.2019.11.017.

39. Fenske, N. A. & Lober, C. W. Structural and functional changes of normal aging
skin. *J Am Acad Dermatol* **15**, 571–585 (1986).

40. Cai, C. *et al.* Impaired dynamics of precapillary sphincters and pericytes at first-
order capillaries predict reduced neurovascular function in the aging mouse brain. *Nat
Aging* **3**, 173–184 (2023).

371
372

Acknowledgements

We thank members of the Littman lab for valuable discussion and critical reading of the manuscript. We thank Michael Cammer from the NYU Microscopy Core for valuable discussion, training and technical support. The Microscopy Core is partially supported by NYU Cancer Center Support Grant NIH/NCI P30CA016087 at the Laura and Isaac Perlmutter Cancer Center, S10 RR023704-01A1 and NIH S10 ODO019974-01A1. This work was supported by a Jane Coffin Childs Fund fellowship (K.R.M.), Kirschstein-NRSA training grant T32AR64184 (K.R.M.), Kirschstein-NRSA individual postdoctoral fellowship F32AG071336 (K.R.M.), a Charles H. Revson Senior Fellowship in Biomedical Science (K.R.M.), the Helen and Martin Kimmel Center for Biology and Medicine (D.R.L.), NIH grant R01AI158687 (D.R.L.), and the Howard Hughes Medical Institute (D.R.L.).

Author Contributions

K.R.M. and D.R.L. designed the study and analyzed the data; K.R.M. and K.A.O. performed mouse experiments. K.R.M. performed intravital multiphoton imaging. C.N. and S.P.S. provided human biological samples and related quantitative analysis. K.R.M. and D.R.L. wrote the manuscript, with input from the other authors. D.R.L. supervised the research.

398   **Figures**

Figure 1

Figure 1 –**Niche-specific macrophage loss with age correlates with impaired skin**

capillary blood flow. a, Schematic of intravital imaging of resident macrophage

populations in mouse skin, indicating epidermal and dermal populations, using the

macrophage reporter, *Csf1r-EGFP,* in combination with an *Actb*-driven universal

tdTomato reporter, *R26-mTmG*. **b,** Representative optical sections of distinct resident

macrophage populations in young (2-month-old) and old (18-month-old) mice. **c,**

Quantification of niche-specific macrophage density change in 1, 2, 4, 10, and 18 month

old mice (n = 4 mice in each age group; two 500μm^2 regions per mouse; macrophage

density in each skin niche was compared across age groups by one-way ANOVA; mean

± SD). **d,** Skin resident macrophage labeling using *Cx3cr1-CreERT2;R26-mTmG* mice

was performed following a single high-dose intraperitoneal injection of tamoxifen (2mg)

in 1 month old mice. Single optical sections at successive time points 5 min apart

showing red blood cell (RBC) flow (white) in capillaries (red) with or without nearby

CAMs (green). Yellow arrowheads indicate obstructed RBC capillary flow. **e,**

Quantification of obstructed capillary flow as measured by stalled RBCs as described in

Extended Data Figure 2 (n = 226 CAM+ capillary segments, n = 27 CAM- capillary

segments; n = 4 mice; obstructed capillary flow (CAM+ vs CAM-) was compared by

paired Student's t test; mean ± SD). **f,** Percentage of capillary segments with at least

one associated macrophage (n = 4 mice in each age group; two 500µm^2 regions per

mouse; macrophage association with capillary segments was compared across age

groups by one-way ANOVA; mean ± SD). **g,** Representative images demonstrate

macrophage depletion following intradermal injections of clodronate-liposomes every 3

days. Repeated intravital imaging of the vascular niche was performed to visualize

macrophages (*Csf1r-GFP*), capillaries (*R26-mTmG*) and RBC flow (Third Harmonic). **h,**

Percentage of capillaries with blood flow following macrophage depletion (n = 194

capillary segments in clodronate group, n = 199 capillary segments in PBS group; 3

mice in each group; capillary flow (clodronate vs PBS) was compared by unpaired

Student's t test; mean ± SD). Scale bar, 50 µm.

Figure 2 – Local CAM recruitment and phagocytosis are required to restore

impaired blood flow. a, Scheme of laser-induced capillary clot experiment. **b,**

Sequential revisits of damaged capillary segment (red dashed lines) after laser-induced

clot formation in *Cx3cr1-CreERT2;R26-mTmG* mice. Yellow lightning bolt indicates site

of laser-induced capillary clot. Yellow arrowheads indicate extra-luminal vascular debris. **c,**

Quantification of capillary-associated macrophage extension toward damaged niche at

Day 1, 2 and 7 after laser-induced clotting (n = 50 CAMs, in 3 mice; mean ± SD). **d,**

Sequential revisits of naturally forming capillary clots in *Csf1r-EGFP* mice. Simultaneous

visualization of blood flow during clot repair via third harmonic generation (white) from

red blood cells (RBC) and intravenous rhodamine dextran (RhDex) (red). **e,** Sequential

revisits of damaged capillary niche after laser-induced CAM ablation and clot formation

in *Csf1r-EGFP* mice. Macrophage laser ablation (red lightning bolt) and capillary clot

formation (yellow lightning bolt) were both performed at 940nm for 1s. **f,** Quantification

of capillary reperfusion at Day 1 and 7 after laser-induced clotting and macrophage

ablation (n = 19 capillary clots in CAM ablated regions, n = 16 capillary clots in control

regions; 3 mice in total; capillary reperfusion (CAM ablated vs control) was compared by

paired Student's t test; mean ± SD). **g,** Sequential revisits of damaged capillary niche

after laser-induced clot formation in $Cx3cr1^{CreER};Rac1^{fl/fl}$ and $Cx3cr1^{CreER};Rac1^{fl/+}$ mice.

CAMs (green), capillaries (red), RBC (white). **h,** Quantification of capillary reperfusion at

Day 1, 3 and 7 after laser-induced clotting (n = 67 clots in $Rac1^{fl/+}$ group; n = 82 clots in

$Rac1^{fl/fl}$ group; n = 3 mice in each group; capillary reperfusion ($Rac1^{fl/+}$ vs $Rac1^{fl/fl}$) was

compared by unpaired Student's t test; mean ± SD). **i,** Scheme of age-associated

decline of skin capillary niche. Capillary-associated macrophage density declines with

age, which predisposes aged capillaries to impaired repair and sustained tissue

perfusion. Scale bar, 50μm.

Figure 3 – Capillary-associated macrophages do not replenish lost neighbors,

resulting in population loss with age. a, Representative revisits of single macrophage

lineage tracing (blue arrowhead) in *Cx3cr1-CreERT2*; *R26-mTmG* mice. Weekly revisits

were performed during homeostatic conditions for 20 weeks following a single low-dose

intraperitoneal injection of tamoxifen (50µg). **b,c,** Quantification of (b) weekly rate of

CAM loss and division (n = 262 CAMs; 8 mice in total; CAM turnover rates (CAM loss vs

division) was compared by paired Student's t test; mean ± SD) and of (c) maintenance

of the labeled CAM population over 20wk lineage tracing (n = 59 CAMs; 3 mice in total;

number of fluorescently-labeled CAMs was compared across each time point in the

same mice by one-way repeated measures ANOVA with Geisser-Greenhouse

correction; mean ± SD). **d,** Representative revisits of CAM replacement after laser-

induced CAM ablation in *Csf1r-EGFP;R26-mTmG* mice. Macrophage laser ablations

(yellow asterisk) were performed in a 500µm^2 region (yellow dashed line square), **e,**

Representative revisits of CAM replacement after a single intraperitoneal injection

diphtheria toxin (10ng/g body weight in PBS) in *Cx3cr1-DTR* mice. **f,** Representative

revisits of single lineage-tracked macrophages in *Cx3cr1-CreERT2*; *R26-dsRed*; *Csf1r-*

EGFP mice. Weekly revisits were performed during homeostatic conditions for 2 weeks

following a single low-dose intraperitoneal injection of tamoxifen (50µg). **g,**

Quantification of replenishment rate of empty CAM niche under laser-induced loss (n =

44 CAMs; 4 mice in total), DT-induced loss (n = 53 CAMs; 3 mice in total), homeostatic

loss (n = 29 CAMs; 3 mice in total), or laser-induced CAM loss with capillary clot (n = 25

CAMs; 4 mice in total). Frequency of CAM replacement after forced and homeostatic

CAM loss were compared by unpaired Student's t test; mean ± SD. Scale bar, 50µm.

Figure 4

Figure 4 –**Local CAM replenishment in old mice is sufficient to rejuvenate**

capillary repair and tissue reperfusion. a, Scheme of CSF1-induced rejuvenation of

aged capillary niche. **b,** Representative images of CAM density in *Csf1r-EGFP* mice

nine days following daily intradermal injections (4 days) of CSF1-Fc (porcine CSF1 and

IgG1a Fc region fusion protein) and PBS in contralateral hind paws of 20-24 month old

mice. **c,** Quantification of CAM density following CSF1-Fc or PBS treatment (n = 4 mice

in total; two 500µm^2 regions of each treatment condition per mouse; Percentage of CAM

density change from Day -9 (CSF1-Fc vs PBS) was compared for Day -4 and Day 0

time points by paired Student's t test; mean ± SD). **d,** Representative images of

capillary (red dashed lines) blood flow in *Csf1r-EGFP* mice nine days following daily

intradermal injections of CSF1-Fc and PBS in contralateral hind paws of 20-24 month

old mice. Yellow arrowheads indicate obstructed RBC capillary flow. **e,** Quantification of

obstructed capillary blood flow following CSF1-Fc or PBS treatment (n = 214 capillary

segments in CSF1-Fc treated regions, n = 214 capillary segments in PBS treated

regions; n = 4 mice in total; obstructed capillary flow (CSF1-Fc vs PBS) was compared

for Day -9 and Day 0 time points by paired Student's t test; mean ± SD). **f,** Sequential

revisits of damaged capillary niche after laser-induced clot formation (yellow lightning

bolt) in *Csf1r-EGFP* mice. Yellow arrowheads indicate extra-luminal vascular debris. **g,**

Quantification of capillary reperfusion at Day 1 and 7 after laser-induced clotting (n = 18

clots in CSF1-Fc group; n = 20 clots in PBS group; n = 4 mice in total; capillary

reperfusion (CSF1-Fc vs PBS) was compared by paired Student's t test; mean ± SD). **h,**

Working model of resident macrophage aging in the skin capillary niche. Age-

associated impairment in capillary repair can be rejuvenated following local expansion

of resident macrophage population. Scale bar, 50µm.

Methods

Mice

Mice were bred and maintained in the Alexandria Center for the Life Sciences animal

facility of the New York University School of Medicine, in specific pathogen-free

conditions. Albino B6 (B6(Cg)-Tyr^{c-2J}/J, Jax 000058), Csf1r^{EGFP} (B6.Cg-Tg(Csf1r-

EGFP)1Hume/J, Jax 018549), Ccr2^{RFP} (B6.129(Cg)-Ccr2tm2.1Ifc/J, Jax 017586),

R26^{mTmG} (B6.129(Cg)-Gt(ROSA)26Sortm4(ACTB-tdTomato,-EGFP)Luo/J, Jax 007676),

R26^{nTnG} (B6N.129S6-Gt(ROSA)26Sortm1(CAG-tdTomato*,-EGFP*)Ees/J, Jax 023537),

LysM^{Cre} (B6.129P2-Lyz2tm1(cre)Ifo/J, Jax 004781), Rac1$^{f/f}$ (Rac1tm1Djk/J, Jax 005550)

mice were purchased from Jackson Laboratories. *R26^{dsRed}* mice were described

previously (Luche et al., 2007) and were obtained from the laboratory of Dr. Gordon

Fishell. *Cx3cr1^{CreER}*, *Cx3cr1^{GFP}, and Cx3cr1^{DTR}* were generated in our laboratory and

have been described (3 ref). Cre-induction for the lineage tracing or total CAM labeling

experiments was induced with a single intraperitoneal injection of Tamoxifen (Sigma-

Aldrich; T5648) (100μg or 4mg in corn oil, respectively) in 1 month old mice. *Rac1$^{f/f}$*

recombination was induced with two intraperitoneal injections of Tamoxifen (2mg in corn

oil) 48hrs apart in 1 month old mice. All imaging and experimental manipulation are

performed on non-hairy mouse plantar (hind paw) skin. Preparation of skin for intravital

imaging are performed as described below. Briefly, mice are anesthetized with

intraperitoneal injection of ketamine/xylazine (15 mg/ml and 1 mg/ml, respectively in

PBS). After imaging, mice are returned to their housing facility. For subsequent revisits,

the same mice are processed again with injectable anesthesia. The plantar epidermal

regions are briefly cleaned with PBS pH 7.2, mounted on a custom-made stage and a

glass coverslip are placed directly against the skin. Anesthesia is maintained throughout

the course of the experiment with vaporized isoflurane delivered by a nose cone. Mice

from experimental and control groups were randomly selected for live imaging

experiments. No blinding was done. All lineage tracing and ablation experiments were

repeated in at least three different mice. All animal procedures were performed in

accordance with protocols approved by the Institutional Animal Care and Usage

Committee of New York University School of Medicine.

In vivo imaging and laser ablation

Image stacks were acquired with an Olympus multiphoton FVMPE-RS system equipped

with both InSight X3 and Mai Tai Deepsee (Spectra-Physics) tunable Ti:Sapphire

lasers, using Fluoview software. For collection of serial optical sections, a laser beam

(940nm for GFP/tdTomato/dsRed/RFP/Rhodamine/Second Harmonic Generation and

1300nm for Third Harmonic Generation, respectively) was focused through a water

immersion lens (N.A. 1.05; Olympus) and scanned with a field of view of 0.5mm^2, at

600 Hz. Z-stacks were acquired in 1-2µm steps for a ~50-100µm range, covering the

epidermis and dermis. Capillary blood flow was visualized in some experiments through

intravenous injection with 18 mg/kg of dextran-rhodamine 70 kD (Sigma-Aldrich;

R9379). Cell tracking analysis was performed by re-visiting the same area of the dermis

in separate imaging experiments through using inherent landmarks of the skin to

navigate back to the original region, including the distinct organization of the superficial

vasculature networks. Cells that were unambiguously separated (by at least 250µm)

from another were sampled to ensure the identity of individual lineages. For time-lapse

recordings, serial optical sections were obtained between 5-10 minute intervals,

depending on the experimental setup. Laser-induced cell ablation, capillary clot, or

tissue damage was carried out with the same optics as used for acquisition. An 940nm

laser beam was used to scan the target area (1-500µm^2) and ablation was achieved

using 50-70% laser power for ~1sec. Ablation parameters were adjusted according to

the depth of the target (10-50µm). Mice from experimental and control groups were

randomly selected for live imaging experiments. No blinding was done. All lineage

tracing and ablation experiments were repeated in at least three different mice.

Drug treatments

To induce macrophage depletion, mice received intradermal injections of either

Clodronate-liposomes or PBS-liposomes (stock concentration 5mg/ml; Liposoma; CP-

005-005) (5µl per paw) every 3 days. Depending on experimental details, Cx3cr1^{DTR}

mice received either intraperitoneal (IP) injection of diphtheria toxin (Sigma-Aldrich;

D0564) every other day at 25ng/g body weight in PBS or a single low dose IP injection

at 10ng/g body weight in PBS. To induce macrophage expansion, mice receive daily

intradermal injections of CSF1-FC (Bio-Rad; PPP031) or PBS in contralateral hind paws

(5µl per paw) for 4 days.

Human skin samples

Written informed consent was obtained for postmortem examination from next of kin for

10 patients. Clinical information and laboratory data were obtained from the electronic

medical record. Skin samples were obtained from the anterolateral chest and fixed in

10% formalin for at least 24h prior to processing. Slides were stained with hematoxylin

and eosin, CD68 (Clone 514H12) and ERG (Clone EPR3864). Macrophages and

capillaries were identified using a combination of morphology, CD68 and ERG staining.

Counting was performed on at least eight high-power fields (40x) within 100um of the

epidermis.

Image Analysis

Raw image stacks were imported into Fiji (NIH, USA) or Imaris software (Bitplane/Perkin

Elmer) for further analysis. Provided images and supplementary videos are typically

presented as a maximal projection of 4-8μm optical sections. For visualizing individual

labeled cells expressing the dsRed or tdTomato Cre reporters, the brightness and

contrast were adjusted accordingly for the green (GFP) and red (dsRed/tdTomato)

channels and composite serial image sequences were assembled as previously

described. Images were obtained as large tiled image stacks at roughly the same

positions and then manually aligned over the experimental time course in Imaris

(Bitplane/Perkin Elmer) by using data from all channels.

Statistical Analysis

Data are expressed as mean ± SD. An unpaired Student's *t*-test was used to analyze

data sets with two groups. One-way ANOVA was used to analyze data sets with three

or more groups. $p < 0.05$ to $p < 0.0001$ indicated a significant difference. Statistical

calculations were performed using the Prism software package (GraphPad, USA).

Extended Data Figure Legends

Extended Data Figure 1

Extended Data Figure 1. CX3C-chemokine receptor 1 (CX3CR1) expression in the

upper dermis defines a resident population of capillary-associated macrophages

(CAMs). a) Representative images of cells expressing *Ccr2-RFP*, *Csf1r-EGFP*, or

Cx3cr1-GFP in the epidermis, upper dermis, and lower dermis of 1 month old mice. b)

Quantifications of cells expressing *Csf1r-EGFP*, *Cx3cr1-GFP, or Ccr2-RFP*, in the upper

dermis (n = 3 mice in each group; two 500μm^2 regions per mouse; Cell number (*Csf1r-*

EGFP vs *Cx3cr1-GFP; Csf1r-EGFP* vs *Ccr2-RFP*) was compared by unpaired Student's

t test; mean ± SD). c-d) Simultaneous visualization of blood flow through the superficial

capillary plexus via third harmonic generation (white) from red blood cells (RBC) and

intravenous rhodamine dextran (RhDex) (red). e) Quantifications of RBC and RhDex

labeling of the upper dermal superficial capillary plexus (n = 3 mice in total; four 500μm^2

regions per mouse; mean ± SD). f) Visualization of CX3CR1-expressing cells in all skin

layers: epidermis, upper dermis, and lower dermis (*Cx3cr1-GFP;R26-mTmG*) was

performed during homeostatic conditions. g) Representative image of labeled upper

dermal macrophages in contact with capillary superficial plexus (red) in 1 month old

mice. h) Quantifications reveal most upper dermal macrophages are capillary-

associated macrophages (CAMs) (n = 3 mice in total; three 500μm^2 regions per mouse;

mean ± SD) . Scale bar, 50 μm.

Extended Data Figure 2. Label-free *in vivo* visualization of capillary blood flow via third harmonic effect generated by red blood cells. a, Scheme of red blood cell (RBC) flow through a segment of the superficial skin capillary network. During three-dimensional image acquisition, flowing red blood cells are captured at different x,y positions for each z-section along the capillary segment. Pseudocoloring each z-step through a capillary segment distinguishes flowing RBCs as a multicolor patchwork or

rainbow-effect and leaves obstructed/non-flowing RBCs as white (full overlap of all

colors). **b,** Representative image of RBC visualization in the upper dermis in *Csf1r-*

EGFP; R26-mTmG mice. **c,** Representative optical z-sections through upper dermal

capillaries. Third harmonic signal is pseudocolored differently for each z-step to

visualize RBC movement along the capillary segments during image acquisition. Scale

50µm.

Extended Data Figure 3

Extended Data Figure 3. Capillary and associated macrophage loss in mice and

humans. a,b Quantification of capillary niche age-associated changes, (a) Capillary

density and (b) CAM / Capillary segment ratio (n = 4 mice in each age group; two

500μm^2 regions per mouse; comparison across age groups was by one-way ANOVA;

mean ± SD). **c,** Representative immunohistochemistry of CD68+ macrophage and Erg+

capillary density in the upper dermis of both young (<40y) and old (>40y) human

samples. **d-f,** Quantification of capillary niche age-associated changes: (d) CD68+

macrophage density, (e) Erg+ capillary density, (f) Upper dermal macrophage / Capillary

endothelium ratio (n = 5 patient samples in each group; three imaging regions per sample;

Comparison by unpaired Student's t test; mean ± SD). Scale bar, 50μm.

Extended Data Figure 4

Extended Data Figure 4. Impaired skin capillary blood flow following acute

macrophage depletion. a, Scheme of long-term capillary blood flow tracking following

intraperitoneal injection every other day with diphtheria toxin (DT) (25ng/g body weight

in PBS) in both *Cx3cr1-DTR*; *R26-mTmG* and WT control (*R26-mTmG*) mice. **b,**

Representative revisits of the same capillary network to visualize capillaries (*R26-*

mTmG) and RBC flow (Third Harmonic). **c,** Quantifications reveal a significant reduction

in the percentage of capillaries with blood flow following DT-induced cell depletion (n =

647 capillary segments in WT control regions, n = 695 capillary segments in *Cx3cr1-*

DTR regions; n = 3 mice in each group; obstructed capillary flow (WT vs *Cx3cr1-DTR*)

was compared for Day 0 and Day 5 time points by unpaired Student's t test; mean ±

SD). Scale bar, 50 µm.

Extended Data Figure 5

724 Extended Data Figure 5. Laser-induced model of acute capillary clot formation

725 and repair. a, Scheme of laser-induced capillary clot experiment. **b,** Sequential revisits

of damaged capillary niche after laser-induced clot formation in *Csf1r-EGFP* mice.

Simultaneous visualization of blood flow before and after clot formation via third

harmonic generation (white) from red blood cells (RBC) and intravenous rhodamine

dextran (RhDex) (red). Yellow lightning bolt indicates site of laser-induced capillary clot.

Yellow arrowheads indicate extra-luminal vascular debris. **c,** Scheme of laser-induced

capillary clot in niches with or without CAMs. **d,** Sequential revisits of damaged capillary

niche after laser-induced clot formation in *Csf1r-EGFP;R26-mTmG* mice. Capillary clot

formation (yellow lightning bolt) was performed at 940nm for 1s in 10 month old mice.

Yellow arrowheads indicate extra-luminal vascular debris. **e,** Quantification of capillary

reperfusion at day 1 and 7 after laser-induced clotting (n = 16 capillary clots in regions

with CAMs (<75μm from clot), n = 16 capillary clots in regions without CAMs (>75μm

from clot; 3 mice in total; capillary reperfusion (with CAMs vs without CAMs) was

compared at Day 1 and Day 7 by paired Student's t test; mean ± SD). Scale bar, 50 μm.

Extended Data Figure 6

Extended Data Figure 6. CAM density in mice with deficiencies in macrophage

phagocytic machinery. a, Scheme of laser-induced capillary clot in

Cx3cr1^{CreER};Rac1^{fl/fl} and *Cx3cr1^{CreER};Rac1^{fl/+}* mice. **b,** Quantification of CAM density at

Day 7 after laser-induced clotting (n = 3 mice in each group; two 500µm^2 regions per

mouse; CAM density (*Rac1^{fl/+}* vs *Rac1^{fl/fl}*) was compared at day 0 (same day of clot

induction) by unpaired Student's t test; mean ± SD).

Extended Data Figure 7

Extended Data Figure 7. Single-cell *in vivo* lineage tracing of resident capillary-

asssociated macrophages. a, Scheme of long-term tracking of CAM self-renewal

experiment. **b,** Representative image of single macrophage tracing in *Cx3cr1-CreERT2*;

R26-mTmG mice. **c,** Representative serial revisit images of single macrophage tracing

in *Cx3cr1-CreERT2*; *R26-mTmG* mice.

Extended Data Figure 8

Extended Data Figure 8. Injury-induced replenishment of CAM population is

partially CCR2-dependent. a, Representative revisits of CAM replacement after laser-

induced CAM ablation in *Csf1r-EGFP* mice. Macrophage laser ablations (yellow

lightning bolt) were performed to leave an empty capillary niche (orange dashed line). **b,**

Scheme of CAM replacement after capillary niche damage in *Csf1r-EGFP*; *Ccr2^RFP/+^* or

*Ccr2^RFP/RFP^* mice. **c,** Representative revisits of CAM replacement after a regional laser-

induced damage to the capillary niche. **d,** Quantification of replenishment rate (*Csf1r-*

793 *EGFP+* cells) of empty CAM niche after laser-induced CAM loss and niche damage (n =

794 4 mice per group; two 500µm^2 regions per mouse; CAM density (*Ccr2$^{RFP/+}$* vs *Ccr2$^{RFP/RFP}$*)

795 was compared by unpaired Student's t test; mean ± SD). Scale bar, 50µm.

Extended Data Figure 9

Extended Data Figure 9. Spatially uncoupled proliferation and loss progressively lead to disorganized macrophage niche patterning. a, Scheme of long-term tracking of CAM migration following cell division. **b,** Representative revisits of single macrophage lineage tracing in *Cx3cr1-CreERT2*; *R26-nTnG* mice. Weekly revisits were performed during homeostatic conditions for 5 weeks following a single low-dose intraperitoneal injection of tamoxifen (50µg). **c,** Quantification of neighboring CAM distance of recently divided sister CAMs was compared to total CAM neighboring distance from *Cx3cr1-CreERT2*; *R26-nTnG* mice given a single high-dose intraperitoneal injection of tamoxifen (4mg) (n = 26 sister CAM pairs, from 5 mice in low-dose group; n = 188 CAMs, from 4 mice in high-dose group; Average nearest neighbor distance (2 vs 10 month old) was compared at 0-25, 26-50, 51-75, and >75µm intervals by unpaired Student's t test mean ± SD). **d,** Quantification of distance between nearest CAM

829   neighbors in 2 and 10 month old *Csf1r-EGFP* mice (n = 90 CAMs in 2 month old mice, n

830   = 101 CAMs in 10 month old mice; 3 mice in each age group; Frequency of CAM

831   distribution (2 vs 10 month old) was compared at 0-25, 26-50, 51-75, and >75µm

832   intervals by unpaired Student's t test; mean ± SD). **e,** Working model of macrophage

833   renewal and organization in the skin aging capillary niche. Scale bar, 50µm.

Extended Data Figure 10

Extended Data Figure 10. Large niche damage expands CAM population and

improves capillary repair in old mice. a, Scheme of tracking of CAM proliferation

following laser-induced damage to nearby capillary or epidermal niches. **b,c,**

Representative revisits of single macrophage lineage tracing in *Cx3cr1-CreERT2*; *R26-*

nTnG mice following (b) capillary and (c) epidermal damage. **d,e,** Quantification of CAM

proliferation based on proximity to (d) capillary and (e) epidermal damage (n = 25 CAMs

tracked in capillary damage regions, n = 56 CAMs tracked in epidermal damage

regions; 4 mice in each group; CAM proliferation (based on damage proximity) was

compared at day 7 by paired Student's t test; mean ± SD). **f,** Scheme of local damage-

induced expansion of CAMs in the aged capillary niche. **g,** Representative images of

CAM density in *Csf1r-EGFP; R26-mTmG* mice 28 days following overlaying epidermal

laser damage. **h,** Quantification of CAM density following overlaying epidermal damage

(n = 4 mice in total; two 500µm^2 regions for both epidermal damage and control

conditions per mouse; CAM density (Epidermal damage vs Control) was compared by

paired Student's t test; mean ± SD). **i,** Quantification of capillary reperfusion at day 1

and 7 after laser-induced clotting (n = 17 capillary clots in epidermal damage regions, n

= 17 capillary clots in neighboring control regions; 4 mice in total; capillary reperfusion

(with epidermal damage vs control) was compared at day 1 and day 7 by paired

Student's t test; mean ± SD). following laser-induced damage to overlaying epidermal

niche. Scale bar, 50µm.

Supplemental Video Legends

Supplemental Video 1. Serial optical sections through mouse plantar skin, including

epidermis (0-25µm), upper dermis (26-50µm), and lower dermis (51-100µm) in *R26-*

mTmG (red) mice. Note Second and Third Harmonic Generation illuminates dermal

collagen (blue) and red blood cells (white), respectively. Scale bar, 50µm.

Supplemental Video 2. Serial optical sections through mouse plantar skin, including

epidermis (0-15µm), upper dermis (16-40µm), and lower dermis (41-52µm) in *Csf1r-*

EGFP mice. Note distinct macrophage (green) density and morphology in each tissue

niche. Scale bar, 50µm.

Supplemental Video 3. Time-lapse recording of capillary blood flow via third harmonic

generation (white) from red blood cells and intravenous rhodamine dextran (red) in

Csf1r-EGFP mice. Note some capillary segments show obstructed red blood cell and

rhodamine dextran flow. Scale bar, 50µm.

Supplemental Video 4. Serial optical sections through upper dermal capillary niche in

Cx3cr1-GFP; R26-mTmG mice. Note macrophage (green) localization and morphology

around capillary (red) network. Scale bar, 50µm.

Supplemental Video 5. Time-lapse recording of capillary blood flow in *Cx3cr1-CreER;*

R26-mTmG mice. Note inconsistent and obstructed blood flow (white) in capillary (red)

segments without associated macrophages (green). Scale bar, 50μm.

Supplemental Video 6. Time-lapse recording following laser-induced capillary clot

formation in *Csf1r-GFP; R26-mTmG* mice. Note rapid migration of neighboring capillary-

associated macrophages (green) toward site of clot formation (yellow arrowhead). Scale

bar, 50μm.