Mutational order and epistasis determine the consequences of *FBXW7* mutations during colorectal cancer evolution

Dedrick Kok Hong Chan[1,2], Amit Mandal[1], Yi Zhou[1], Scott David Collins[1], Richard Owen[3], James Bundred[1], Hannah Fuchs[3], Sabrina James[3], Iolanda Vendrell[4,5], Sarah Flannery[4], David Fawkner-Corbett[1,6], Jacob Househam[7], Trevor A Graham[7], Roman Fischer[4,5], Alison Simmons[6], Xin Lu[3], Simon James Alexander Buczacki*[1]

[1] Nuffield Department of Surgical Sciences, University of Oxford, Oxford, United Kingdom

[2] NUS Centre for Cancer Research, Yong Loo Lin School of Medicine, Singapore

[3] Ludwig Institute for Cancer Research, Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom

[4] Target Discovery Institute, Centre for Medicines Discovery, Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom

[5] Chinese Academy of Medical Sciences Oxford Institute, Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom

[6] MRC Translational Immune Discovery Unit, Weatherall Institute of Molecular Medicine, University of Oxford, John Radcliffe Hospital, Headington, Oxford. United Kingdom

[7] Centre for Evolution and Cancer, The Institute of Cancer Research, London, United Kingdom.

Running title: *FBXW7* and mutational order in CRC evolution

Correspondence to:

✉ simon.buczacki@nds.ox.ac.uk

☎ +44 (0) 7885 433458

Summary

Somatic driver mutations, in genes such as *FBXW7,* have been discovered in phenotypically normal

colonic tissue, however their role in cancer initiation remains elusive. Using normal and gene-edited

patient-derived human colon organoids as models of early tumour evolution we observed that

FBXW7$^{-/-}$ mutations exerted an epistatic effect on subsequent transcription depending on the

mutational background of the cell. Specifically, the timing of acquiring an *FBXW7$^{-/-}$* mutation

respective to an *APC* mutation, led to profound phenotypic and transcriptomic differences. When

FBXW7 was mutated before *APC*, a near-normal cell state was maintained alongside repression of the

APC transcriptional response. However, when *APC* was mutated before *FBXW7*, cells acquired

classical cancer-stem cell features. Single-cell RNA sequencing revealed that mutation of *FBXW7* in

normal tissue subtly switched cells from adult to a foetal/regenerative stem cell state. Further analysis

using transposase-accessible chromatin sequencing identified this cellular plasticity was driven by

changes in chromatin accessibility of transcriptional start site regions associated with TEAD, SNAI1

and AP-1 motifs, which in turn activate the foetal-like state. Taken together, we demonstrate a critical

role of *FBXW7* mutations in preventing colorectal cancer initiation and provide exemplar evidence for

the importance of epistasis and mutational order in cancer biology.

Introduction

Somatic mutations have been detected in a range of normal human tissues, including the liver [1], oesophagus [2], skin [3], and endometrium [4]. Similarly, a catalogue of somatic mutations in normal human colonic epithelium using whole genome sequencing of two thousand crypts, strikingly found mutated cancer genes in phenotypically normal colonic epithelium [5]. In healthy colon such mutations form a distinct category from initiating cancer driving mutations in genes such as *TP53* or *APC*, which are ubiquitously found in colorectal cancer. *FBXW7* is a gene within the catalogue of mutations observed in normal colonic epithelium but its role in cancer initiation has not yet been investigated. FBXW7 is a key component of the Skp1-Cdc53/Cullin-F-box-protein complex (SCF/β-TrCP), an E3 ubiquitin ligase responsible for marking proteins by ubiquitination and targeting them for proteasomal degradation [6]. Substrates of FBXW7 are regulators of gene transcription, cell cycle progression or activators of signalling pathways, thus making *FBXW7* a pivotal tumour suppressor gene. Given *FBXW7* appears to acquire mutations before the widely accepted initiating mutations associated with colorectal cancer such as *APC*, *TP53*, *SMAD4* or *KRAS*, raises the possibility that mutated *FBXW7* could play a critical role in early cancer evolution. Furthermore, the timing of the *FBXW7* mutation suggests that the order of mutation acquisition may have important effects on phenotype.

Recently, there has also been significant interest in the effects of epigenetic plasticity in cancer evolution. In models of early pancreatic cancer, *KRAS*-mutant cells were shown to demonstrate distinct chromatin accessibility patterns which could predict divergence into either a benign or malignant fate [7]. In colorectal cancer, the impact of epigenetic changes on tumour initiation are beginning to be uncovered. Johnstone et al. highlighted the importance of the three-dimensional architecture, in the form of chromatin loops, topologically activating domains and large-scale compartments, of DNA in restricting malignant progression [8]. A more recent study by Heide et al. found chromatin modifier genes are actively mutated in the early initiation of colorectal cancer and can result in the accumulation of further genetic mutations which herald carcinogenesis [9].

In this study, we used normal human adult and foetal colonic organoids, applying CRISPR-Cas9 gene

editing to explore the cell autonomous and gene-interactive effects of an *FBXW7* mutation.

Combining gene editing with organoid culture fashions a powerful tool ideal for studying the

functional consequences of driver mutations in normal tissue for it allows specific mutations to be

introduced into tissue devoid of genetic perturbation. By sequentially introducing multiple distinct

mutations into the same cell, we also aimed to determine relationships arising between cells with

mutant *FBXW7* and the acquisition of subsequent cancer driver mutations. We found that cells with

mutant *FBXW7* were transcriptionally like wildtype cells but primed towards a foetal stem cell state.

Further, we observed that an initial *FBXW7* mutation repressed the transcriptional effect of a

subsequent *APC* mutation, resulting in retention of the near-wildtype phenotype. Whereas when we

reversed this order of mutations, despite identical genotype, we generated cancer stem cell organoids.

We uncovered this profound mutational order-based cellular plasticity to be driven by epigenetic

differences in both global and local chromatin accessibility. Here we provide direct evidence that

epistasis, and the order of mutations, as much as the mutations themselves, are key in determining

phenotypes.

Results

Mutant *FBXW7*[-/-] recapitulate known effects on downstream targets

Patient-derived adult stem cell wildtype organoids (W) were generated from normal colon tissue of

patients undergoing surgery for colorectal cancer. CRISPR-Cas9 editing was used to generate

FBXW7[-/-] organoids (F) [10]. Our gene editing strategy utilizes three sgRNAs to introduce large

deletions at exon 2 of the *FBXW7* gene. F and W organoids possessed similar morphology (Fig. 1a).

We validated knockout of the *FBXW7* gene using western blot (Fig. 1b), and immunofluorescence

(Fig. 1c). Western blotting of known FBXW7 targets, found upregulation of the expected substrates

upon loss of FBXW7 (Fig. 1d).

To characterise transcriptional differences between F and W organoids, we performed bulk-RNA

sequencing (bulk-RNAseq) (Fig. 1e). Using a cut-off of $\log_2 FC \leq -1.5$ or ≥ 1.5, and adjusted p-value <

0.05, F and W were transcriptionally similar with only two genes found differentially expressed.

Proteomic comparison of F and W organoids showed a larger number of differentially expressed

proteins (n=1097, adjusted p-value < 0.05) however most of those identified showed only small fold

changes. When applying the same fold change cut-off as used in bulk-RNAseq, this dropped to 31

proteins ($\log_2 FC \leq -1.5$ or ≥ 1.5, and adjusted p-value < 0.05) (Fig. 1f). Gene set enrichment analysis

(GSEA) of the top 200 upregulated proteins in F organoids and the ranked bulk-RNAseq gene list

comparing F and W organoids showed strong enrichment (Fig. 1g). Pathway analysis of the ranked

proteins revealed 42 significant Reactome pathways (5 enriched and 37 depleted: adjusted p-value <

0.01 and NES $\leq$ -2 or $\geq$ 2). Interestingly, histone-related proteins contributed significantly (Mann-

Whitney p-value = 1.64e-07) to the depleted Reactome pathways (Fig. 1h,i). The average recurrence

of proteins H2AC21, H2AC6, H2AC7, H2AX, H2AZ1, H2BC11, H2BC17, H2BC26, H2BC3,

H2BC5, H3-3A and H4C1 was 24.9, whereas it was 1.7 only for the remaining set of total 270

proteins contributing to the leading-edge of the enriched/ depleted pathways (Extended Data Table 1).

In summary these analyses confirm $FBXW7^{-/-}$ mutations on a wildtype background have a subtle effect

on phenotype and transcriptome although proteomic changes were apparent.

$FBXW7^{-/-}$ mutants do not exhibit a competitive proliferative advantage over wildtype organoids

Given the transcriptional similarities between W and F organoids, we interrogated the possibility for

competitive or cooperative effects by co-culturing F organoids with W organoids, analogous to the

recently described supercompetitor effect seen with $Apc^{-/-}$ intestinal organoids [11]. $APC^{-/-}$ (A)

organoids were also generated by targeting the hotspot region using CRISPR-Cas9 editing to

introducing a biallelic frameshift indel at codon 1499. A were co-cultured with W organoids and A

were found to have a significant proliferative advantage in concordance with previous murine data

(Extended Data Fig. 1a,c,d). F organoids however did not demonstrate any advantage when co-

cultured with W organoids and transcriptional profiling showed minimal differences between W:W-

cocultured and F:Fcocultured organoids (Extended Data Fig. 1b, e-i) in contrast to our previous

findings in colorectal cancer cell lines [12].

FBXW7 **mutation provides evidence of epistasis**

In light of the lack of a cooperative/competitive interaction with wildtype cells and building on our

earlier finding that F organoids were transcriptionally similar to W organoids, we generated an

organoid model mirroring the polypoidal and cancerous phases of CRC carcinogenesis (Fig. 2a). A

organoids represent the polypoidal, precancerous phase, while a $TP53^{-/-}$ mutation introduced into A

organoids, making a double knockout $APC^{-/-} / TP53^{-/-}$ mutation, generated AT organoids, representing

the cancerous phase. Successful targeting was confirmed using Sanger sequencing and western

blotting (Extended Data Fig. 2a). On each of these A and AT organoids, an *FBXW7* mutation was then

introduced, generating AF and ATF organoids respectively (Fig. 2b) (Extended Data Table 2).

As previously described, A organoids were morphologically distinct from W organoids, growing large

and cystic [13, 14]. The subsequent acquisition of *TP53* mutation preserved this phenotype.

Importantly, the additional mutation of *FBXW7* on both the A and AT organoids also maintained this

cystic morphology. To further characterise transcriptional differences between these different

organoids bulk-RNAseq of A organoids were compared with AF, while AT organoids were compared

with ATF organoids. A organoids were compared with AF organoids, while AT organoids were

compared with ATF organoids. Together with our earlier comparison between F and W organoids

which found only 2 differentially expressed genes, we observed 87 differentially expressed genes

between A and AF, and 294 between AT and ATF (Fig. 2c, and Extended Data Fig. 2b and 2c).

Hierarchical clustering of these different groups showed that F and W organoids clustered together,

while A, AF and AT, ATF organoids formed separate clusters (Fig. 2d). These results demonstrate an

epistatic effect with the loss of *FBXW7* having varying effects on the cells depending on the

mutational background. More importantly, it suggests that an *FBXW7* mutation can influence different

cellular phenotypes depending on when the mutation was acquired, and that an early mutation will

have fewer effects compared with a later mutation.

To ensure our observations were not an artefact of repeated iterations of gene editing, we targeted the

AAVS1 safe harbour locus three separate times to replicate the triple-mutant ATF organoids.

Morphologically, *AAVS1*-targetted organoids resembled wildtype organoids (Extended Data Fig. 3a).

Bulk-RNAseq of *AAVS1*-targetted organoids compared with W organoids showed only two

differentially expressed genes with a cut-off of $\log_2$FC $\leq$ -1.5 or $\geq$ 1.5, and adjusted *p*-value < 0.05

(Extended Data Fig. 3b), greatly differing from the 493 differentially expressed genes between ATF

and W organoids (Extended Data Fig. 3c). These findings confirm that the epistatic effects described

earlier were not simply a function of repeated iterations of CRISPR-Cas9 gene editing.

Order of *FBXW7* mutation determines effect on transcriptional profile

To ascertain the transcriptional effect of order of mutation, we designed an experiment in which

organoids possessed the same mutations, but differed in the order in which the mutations were

introduced. As previously described, AF organoids possessed $APC^{-/-}$ followed by $FBXW7^{-/-}$. We

further generated FA organoids, which acquired $FBXW7^{-/-}$ mutation prior to the $APC^{-/-}$ (Fig. 3a).

Morphologically, FA organoids did not acquire the cystic phenotype characteristic of A or AF

organoids (Fig. 3b). Principal component analysis (PCA) of AF and FA organoids processed for bulk-

RNAseq, revealed that together with W, A and F organoids gene expression of FA organoids did not

cluster with AF organoids (Fig. 3c). Instead, FA organoids clustered with F and W organoids, while

AF organoids clustered with A organoids. Organoid clustering for all samples was independent of

patient donor (Extended Data Fig. 3d). To analyse the transcriptional effect mutational order, we used

a cut-off of $\log_2$FC $\leq$ -0.7 or $\geq$ 0.7 to identify differentially expressed genes when AF and FA

organoids groups were compared with W organoids. This cut-off selected for 3335 genes in the AF

group and 861 genes in the FA group (Fig. 3d). There was a significant overlap of the FA genes where

628 (72.9%) were also differentially expressed in the AF organoids compared to W. Further analyses

of the gene expression changes for these 628 genes suggested a repressive effect after an initial

FBXW7 mutation, such that the magnitude of the transcriptional perturbation was rendered less

positive or negative (Fig. 3e). Using GSEA, we also observed that some key signalling pathways such

as the EMT pathway were significantly more prevalent in FA compared with AF organoids

(normalised enrichment score (NES) 2.21, adjusted *p*-value= 1.70×10^{-10}). Collectively, these findings

promulgate our observation that mutational background alters the transcriptional effect of a mutation.

Moreover, our findings also suggest that early loss of *FBXW7* in normal tissue may protect against the

future effect of *APC* loss through transcriptional repression, and by extension could be protective

against the initiation of CRC.

scRNAseq data shows de-differentiation towards a foetal state in *FBXW7* mutant organoids

We endeavoured to explain our finding where an early *FBXW7* mutation was protective to

consequences of a later acquisition of an *APC* mutation, considering our observation that mutational

order affects transcription. We hypothesised that subtle transcriptomic changes could be present in F

organoids, in small subpopulations of cells, which might be masked by bulking during RNAseq. To

overcome this, we performed single-cell RNA sequencing (scRNAseq) on F and W organoids.

Differential gene expression analysis using pseudobulked F and W cells showed a significant

enrichment of upregulated F specific genes between the scRNAseq samples and our previously

generated F and W bulk-RNAseq (p<0.005) (Extended Data Fig. 4a). Clustering by Uniform Manifold

Approximation and Projection (UMAP) uncovered 14 clusters (Fig. 4a). Notably cluster 13 was

unique to *FBXW7* mutant cells (Fig. 4b). This cluster was marked by genes such as *TAGLN* ($\log_2$FC=

-1.33, *p*= 2.81×10^{-35}) and *MMP7* ($\log_2$FC= -1.47, *p*= 7.70×10^{-34}) and possessed an EMT signature

analogous to that recently demonstrated in a separate analysis of murine foetal organoids (Extended

Data Fig. 4b) [15]. Importantly, we observed that W cells expressed *FBXW7*, indicating that this gene

is expressed in normal epithelium (Fig. 4c). Analysis by RNA velocity indicated that cluster 13 was an

important terminal differentiation state, which could only have occurred in *FBXW7* mutant cells (Fig.

4d). As expected, quantitative analysis between UMAP plots from F and W organoids demonstrated a

statistically significant upregulation of the FA_up gene signature in F organoids relative to W

organoids (Fig. 4e). We also observed increased expression of the foetal gene signature [16] (Fig. 4f),

and the YAP pathway gene signature [17] (Fig. 4g) in F organoids (Extended Data Table 3). There was

also a decreased expression of the adult intestinal stem cell signature (ISC) [18] (Fig. 4h), and

decreased expression of the adult crypt-based columnar (CBC) stem cell signature [19] (Fig. 4i) in F

organoids. Reanalysis of our bulk-RNAseq data comparing F and W organoids using GSEA found

that foetal (Extended Data Fig 5a) and YAP (Extended Data Fig. 5b) signatures correlated with our

scRNAseq results. Similarly, overlaying our bulk-RNAseq F signature on a previously published

scRNAseq analysis of foetal and adult colonic epithelia demonstrated expression of F-specific genes

in foetal cells and W specific genes such as *OLFM4* in adult cells (Extended Data Fig. 5c) [20].

Immunofluorescence staining for LY6D validated it as a novel foetal marker; having been found

consistently upregulated in F using bulk bulk-RNAseq, scRNAseq and proteomics. (Extended Data

Fig. 5d).

In F vs W organoids we also observed a downregulation of the adult-ISC (Extended Data Fig. 5e) and

adult-CBC (Extended Data Fig. 5f) signatures in bulk-RNAseq as in scRNAseq, though this did not

achieve statistical significance. To further explore the YAP pathway, we performed additional GSEA

on FA vs W and AF vs W organoids. Intriguingly, this showed greater enrichment of the YAP

signature in FA vs W (ES= 0.58) (Extended Data Fig. 6a) compared to AF vs W (ES= 0.36) (Extended

Data Fig. 6b). Taken together, these findings point towards a plasticity in stem cell identity, favouring

pathways which recall a foetal regenerative/YAP active stem cell state following *FBXW7* mutation.

In contrast to F organoids acquiring a foetal stem cell state, bulk-RNAseq comparing A vs W showed

loss of the adult ISC signature and acquisition of the recently described proliferative colon stem cell

(proCSC) signature in line with findings of others [21] (Extended Data Fig 7a,b). Further comparison

of FA and AF transcriptomes showed that AF organoids retained a greater enrichment for the proCSC

state whereas FA organoids developed even stronger enrichment for the foetal stem cell state than

seen with the initial F mutation (NES Foetal: FA vs W = 2.03; F vs W= 1.99, p<0.0001) (Extended

Data Fig 7c and d). To better dissect stem cell changes associated with acquisition of an *APC* mutation

we performed scRNAseq on W and A organoids. scRNAseq recapitulated bulk-RNAseq findings

demonstrating clear plasticity in stem cell populations with the proCSC population absent in W

organoids but a dominant population in A organoids (Extended Data Fig 7e). In summary these data

demonstrate that stem cell identity can be determined by order of mutations.

Transcriptional effects of mutational order are driven by local and global changes in chromatin

accessibility

We next sought to uncover the mechanism which resulted in FA and AF organoids exhibiting such

profound transcriptional differences despite their identical genotypes. We hypothesised that changes

in chromatin accessibility induced by a preceding *FBXW7* mutation could influence the effects of a

subsequent *APC* mutation. To interrogate this, we performed assay for transposase-accessible

chromatin with sequencing (ATACseq) on W, F, A, FA, AF and human foetal (Fo) colon organoids

(17pcw and 21pcw), as an *in vitro* epigenomic model of tumour evolution. Initially we compared W, F,

A (matched for donor) and foetal organoids finding whole genome accessibility to be greater in F, A

and foetal organoids when compared to W (Fig. 5a). Foetal and F organoids possessed most

euchromatin, compatible with the profound loss of histones seen in our proteomic analysis. We

performed a *de novo* transcription factor motif scan using the GADEM algorithm on ATACseq peaks

for all comparisons. Accessibility to TEAD, AP-1 and SNAI1 motifs, previously shown to be

associated with the foetal/YAP state, were highly enriched in both foetal and F organoids (Extended

Data Table 4) (Fig. 5b and c) [15]. However, the genomic loci of these motif sites were distinct

between foetal and F organoids suggesting epigenetic differences between induced oncogenic (onco-

foetal) and intrinsic foetal states. As expected, the F vs WT and FA vs AF transcriptional signatures

were also found more accessible in F and foetal organoids (Extended Data Fig. 8a and b). In line with

our bulk-RNAseq, scRNAseq and proetomics data we also found accessibility to *LY6D* to be greater

in F organoids (Extended Data Fig. 8c).

Next, we compared W, FA and AF organoids (matched for donor). In total, 14 543 peaks differed

between AF and W organoids, 16 990 peaks differed between FA and W organoids, and 2 156 peaks

differed between AF and FA organoids. Reassuringly, these findings suggest that there was much

similarity in the effect of mutational order on chromosome accessibility between AF and FA. We

noted that overall whole genome accessibility was reduced in FA vs W organoids compared with AF

vs W organoids (Fig. 5d). Concordant with scRNAseq analysis between F and W organoids, we

observed that there was increased accessibility to genes in the crypt base columnar (CBC) signature in

AF vs W organoids compared with FA vs W organoids (Fig. 5e). De novo motif identification was

performed as previous, and two motifs found enriched in the FA vs W organoid comparison included

"nCwGCmCwGn" (Fig. 5f) and "rGCAGGTGn" (Fig. 5g). The top 3 closest matches to each of these

motifs as per the JASPAR database, were *ZNF449*, *TEAD2*, and *TEAD1, SNAI1*, *SCRT2*, and *CTCFL*

respectively. Intriguingly, TEAD1 and TEAD2 motifs were also more accessible in FA vs W

organoids relative to AF vs W organoids (Fig. 5h). Finally, we observed increased chromatin

accessibility of FA vs W organoids compared with AF vs W organoids at the SNAI1 transcription

factor motif sites (Fig. 5i), which mediates EMT by repressing E-cadherin expression, therefore

concurring with results observed in bulk-RNAseq and scRNAseq analysis of differentially expressed

pathways. Taken together, our findings strongly implicate an epigenetic mechanism in which an early

FBXW7 mutation results in transcriptional repression of the consequences of a later *APC* mutation.

Effects of mutation order are recapitulated in patient data

To perform human *in vivo* validation of our findings, we turned towards two datasets to infer the

transcriptional impact of mutation order. First, we analysed transcriptional data from the recently

published EPICC dataset [9]. Here, the authors performed matched single-crypt RNAseq and whole-

genome sequencing on a small number of phenotypically normal crypts. We reanalysed transcriptional

and genomic data from normal glands and performed PCA (Fig. 6a). Notably, one gland had an

FBXW7-R578 mutation in the absence of an APC mutation, while another gland had an APC-S1346

mutation in the absence of an FBXW7 mutation. Two glands from the same patient possessed an

identical FBXW7-R578 mutation as well as a separate APC-G1288 mutation; which can only be

explained by the *FBXW7* mutation arising first, clonally expanding and then acquiring an *APC*

mutation subsequently. Intriguingly, the gland with an *APC* mutation alone was located on PCA

analysis distinct from all other glands. Notably, the two glands possessing both *APC* and *FBXW7*

mutations also clustered closer to the wildtype glands and the single *FBXW7* mutant gland, analogous

to our findings in gene-edited organoids. Whilst these inferences have been drawn based on a limited

number of glands it is evident that FA mutations can also be found in phenotypically normal colon

crypts.

Next, we adapted a newly described bioinformatic approach to infer mutational order from targeted

sequencing data and applied this to a dataset (S:CORT) of 91 pre-malignant human colonic adenomas

that had undergone targeted sequencing and transcriptional profiling [22] (Fig. 6b). Modified

ASCETIC analysis showed that the invariant mode of adenoma development is an initial *APC*

mutation (p<0.05). *FBXW7* mutations were never found to occur before *APC* in this cohort. In

summary, whilst FA normal crypts can be observed in patient samples, our analysis of adenomas

found that AF is the dominant mode of adenoma formation.

The foetal stem cell state in advanced malignancy has previously been associated with diminished

response to chemotherapy [23, 24]. Chemotherapy can create somatic mutations in cancer and normal

cells [25]. Given our findings that induced foetal states appear resistant to effects of further oncogenic

transformation we hypothesised that the FA defined onco-foetal signature may also identify tumours

resistant to chemotherapy. We analysed gene expression data from the Fluoropyrimidine, Oxaliplatin

& Targeted Receptor pre-Operative Therapy for colon cancer study (FOxTROT), where patients were

randomised to receive neoadjuvant chemotherapy prior to surgery [26]. Here, tumours were biopsied

before and after treatment to identify transcriptional biomarkers of response. We generated differential

gene expression lists by segregating patients into responders and non-responders with pre (n=95) and

post (n=80) treatment biopsies. Next, we performed GSEA using the MSigDB Hallmarks and foetal

gene signatures (n=73) unexpectedly finding the FA_up signature outperformed all previously

published foetal and Hallmarks signatures in identifying non-responders to neo-adjuvant

chemotherapy on pre-treatment biopsies (Fig. 6c and Extended Data Table 5). The AF signature as

well as interferon response signatures on the other hand were strongly associated with improved

responsiveness on pre-treatment biposies. Finally, ssGSEA of TCGA COAD/READ samples showed

an inverse relationship between FA_up and FA_down gene signatures (correlation score: -0.55, $p<2.2$

$\times 10^{-16}$) (Fig. 6e) [27]. Thus, whilst FA order of mutations in pre-neoplasia is an evolutionary dead end,

the FA-defined onco-foetal signature in advanced malignancy is highly predictive of chemotherapy

resistance.

Discussion

Recent work has uncovered that molecular background in intestinal tumours influences a complex

three-way interplay between revival/regenerative stem cell (foetal/YAP), homeostatic adult stem cell

(CBC/ISC) and cancer stem cell populations (proCSC) [19, 21]. Whether similar processes play a role

in pre-neoplasia and how this changes temporally remain unexplored. Using *FBXW7* as an example,

our analyses demonstrate that epigenetics underlie the context-specific cellular response to somatic

driver mutations, progressing the recent notion that tumour genotype alone does not equate to

phenotype [28].

In our study, we found introducing an *FBXW7*$^{-/-}$ mutation in normal colon organoids generated little

transcriptomic or proteomic response and organoids remained broadly 'wildtype' although primed

towards a foetal state. These findings are compatible with the observation *in vivo* where *FBXW7*

mutations are found in histologically normal crypts but failed to explain how *FBXW7* mutations are

drivers in established CRC. In answer to this we uncovered an epistatic effect of the mutation when

acquired on increasingly complex mutational backgrounds.

Building on our demonstration of epistasis we tested whether reordering mutations generated similar

results. To our surprise not only were FA mutants phenotypically distinct from AF mutants but

transcriptionally they also remained closer to the wildtype profiles, despite having lost two major

tumour suppressor genes. Mutational order effects have been described once before in *JAK2/TET2*

mutant haematological malignancies [29]. In a highly analogous manner, the authors found that blood

cancers with switched order of mutations had alterations in stem cell state and transcriptomes.

To understand the link between mutation order, epistasis and stem cell state we tested whether an

epigenetic mechanism could underly our data. Chromatin accessibility analyses showed that whilst

both FA and AF organoids possessed more permissive chromatin than wildtype, broadly FA chromatin

was less accessible than AF accounting for the general repression in the transcriptional consequences

of an *APC* mutation. However, we found FA organoids did have increased accessibility to

TEAD/SNAI1/AP-1 motifs. These data explain the foetal/YAP active state seen with F and FA

mutations. Two very recent studies have explored chromatin landscapes in murine foetal and adult

intestinal organoids [15, 30]. In highly concordant findings, these studies show similar TEAD/AP-

1/SNAI1 family accessibility in foetal organoids and most importantly find *Fbxw7* to be a top hit in a

CRISPR screen for regulators of the adult/foetal intestinal transition. Ultimately, the mechanism by

which *FBXW7* mutations effect changes in chromatin accessibility remains cryptic, though a similar

observation has been observed previously [31]. Given FBXW7's important role in the E3 ubiquitin

ligase complex, we speculate that decreased substrate degradation of one or a number of FBXW7

targets could result in this change in chromatin accessibility, though further work to elucidate this

mechanism will need to be performed.

Our study develops on recent work describing Waddington-like plasticity in intestinal stem cell states

[21]. We propose that whilst the normal colon crypt adopts predominantly the adult homeostatic stem

cell state, altering the order of mutations affects subsequent fate trajectory. *FBXW7* mutation first

cause cells to acquire a foetal state whereas an *APC* mutation first induces cells to acquire an

additional proliferative/cancer stem cell state (Fig. 6f). These states remain epigenetically fixed and

subsequent mutations prevent further plasticity between stem cell states/valleys. Overall, our study

explains the basis for why *FBXW7* mutations in normal colonic epithelia fail to induce malignant

transformation and provide the rationale for testing the translational impact of mutational order across

a wider mutational spectrum and different cancers.

Methods

Human material for organoid cultures

Ethics approval for the retrieval of human colon specimens was accorded by the University of Oxford – Translation Gastroenterology Unit (16/YH/0247). Foetal samples were acquired under the HBDR project 200462, REC: 18/LO/0822. Written consent was obtained from all patients/donors.

Organoid culture

Foetal organoids were derived from two donors (17pcw, Male, Proximal Colon) and (21pcw, Male, Proximal Colon). Adult colonic tissue was obtained from patients who were undergoing surgery for colorectal cancer. A 1 x 1 cm piece of colon from a region at least 5cm distant from the tumour, and appearing phenotypically normal was resected. Organoids were derived based on a previously published protocol [10]. Organoid culture growth media consists of advanced DMEM/F12 (Gibco), and supplemented with penicillin/streptomycin (Gibco), 10mM HEPES buffer solution (Gibco), 2mM GlutaMAX (Gibco), 50% Wnt3a conditioned medium (produced from ATCC CRL-2647 cell line), 25% R-spondin conditioned medium (produced from Cultrex HA-R-Spondin1-Fc 293T cells), 1X B-27 plus supplement (Gibco), 10μM SB 202190 (Tocris), 0.5μM SB 431542 (Tocris), 1μM prostaglandin E2 (Tocris), 50ng/ml Noggin (Peprotech), 50ng/ml EGF (Peprotech), 10mM nicotinamide (Sigma-Aldrich), and 1.25mM N-acetylcysteine (Sigma-Aldrich). For selection of APC mutants, Wnt3a and R-spondin conditioned media were withdrawn from the growth media for at least 3 weeks. For selection of TP53 mutants, growth media was supplemented with 10μM of Nutlin-3 for at least 1 month (Sigma-Aldrich).

CRISPR/Cas9 gene editing of human organoids

sgRNAs were designed using Synthego's CRISPR Design Tool (Extended Data Table 2). For *FBXW7*, a multiguide approach utilising three different guide RNAs in equal concentration was used. This increased the targeting efficacy given the lack of a selecting agent which can be used to enrich for *FBXW7* mutants. The *AAVS1* site is a commonly used safe harbour locus which served as a control to ensure studied effects were not a function of CRISPR gene editing.

Our protocol for gene editing human organoids has been described previously [10]. An

electroporation-based approach was used to maximise gene editing efficacy. Cells were first

dissociated into a single-cell suspension using TrypLe (Gibco) by incubating in a 37°C water bath for

10min. triturating at regular intervals. Organoids were then pelleted with centrifugation at 400g for 6

min. Pelleted organoids were resuspended in advanced DMEM, triturated and pelleted by

centrifugation once more. A ribonucleoprotein (RNP) complex comprising Cas9 (Sigma-Aldrich) and

guide RNA in a molar ratio of 1:4 was resuspended in Buffer R (Invitrogen) and left to stand at room

temperature for 10min. Pelleted organoids were resuspended in Buffer T (Invitrogen). The RNP

complex and resuspended organoids were then mixed. Electroporation was performed using a Neon

Transfection System (ThermoFisher) with the following settings – voltage 1350V, width 20ms, and

pulses 2. After electroporation, organoids are transferred into advanced DMEM and left to recover for

10min. Centrifugation was performed to pellet organoids, which were then resuspended in BME and

transferred to cell culture plates as described above. Following electroporation, organoids were

incubated for one week and allowed to expand. Primers which generated amplicons of approximately

500bp in length were designed on Benchling (Extended Data Table 2). Amplicons were Sanger

sequenced by an external vendor (Genewiz). Efficiency of targeting was carried out using Synthego

Performance Analysis, ICE Analysis. 2019. V3.0 Synthego.

Human colon organoids routinely senesce ~ P7/8 necessitating continual re-derivation, targeting and

expansion. In all experiments each replicate represents a distinct clone generated from an individual

patient unless stated otherwise. Each edited clone was used for assays once targeting efficiency

through selection (APC/TP53) or repeated targeting (FBXW7) was >80%.

Protein lysates and western blot

Proteins were harvested using RIPA (ThermoFisher) with addition of protease and phosphatase

inhibitor cocktails (Sigma-Aldrich). Quantification of protein concentration was performed using

BCA assay (ThermoFisher).

For western blot, 30µg of protein was used for each lane. Proteins were separated using a NuPage 4 –

12% Bis-Tris gel (Invitrogen) and transferred onto a 0.45µm nitrocellulose membrane. For

phosphorylated proteins, 5% BSA was used. For non-phosphorylated proteins, 5% low-fat milk was

used. The membrane was blocked for 1 hour before overnight incubation with primary antibodies at

4°C. Rabbit anti-human FBXW7 antibody (1:2500, BS-8394R, Bioss, USA), rabbit anti-human

phosphor-CJUN antibody (1:2500, PA5-40193, Invitrogen, USA), rabbit anti-human CJUN antibody

(1:2500, ab40766, Abcam, USA), rabbit anti-human phosphor-CCNE1 antibody (1:2500, ab52195,

Abcam, USA), rabbit anti-human CCNE1 antibody (1:2500, ab33911, Abcam, USA), rabbit anti-

human phosphor-CMYC (1:2500, ab185655 and ab185656, Abcam, USA), rabbit anti-human CMYC

(1:2500, ab32072, Abcam, USA), rabbit anti-human NICD (1:2500, #4147, Cell signalling technology,

USA), rabbit anti-human NOTCH1 (1:2500, ab52627, Abcam, USA), rabbit anti-human beta-catenin

(1:1000, #8480, Cell signalling technology, USA), rabbit anti-human p53 (1:1000, #9282, Cell

signalling technology, USA) and rabbit anti-human GAPDH (1:2500, ab52627, Abcam, USA) was

used. The membrane was then incubated with goat anti-rabbit antibody (1:5000, ab6721, Abcam,

USA) at room temperature for 1 hour. The membranes were then imaged with a ChemiDoc XRS+

system (Bio-Rad).

Immunofluorescence and imaging

Immunofluorescence on organoids was previously described in our protocol [10]. Briefly, media from

the wells of organoids was removed and fixation with 2% paraformaldehyde was performed. This was

incubated at room temperature for 30min. Organoids were then washed with ice-cold DPBS gently

and allowed to settle by gravity. The supernatant was then removed and blocked with Organoid

Blocking Solution comprising DPBS supplemented with 3% BSA, 1% Triton X-100, 1% saponin and

1% secondary antibody animal serum. Blocking was performed for 3 hours at room temperature.

Organoids were again allowed to settle by gravity, and the supernatant was removed. Primary

antibody diluted in organoid blocking solution was then added to the organoids for 24 hours at 4°C.

Organoids were then washed with DPBS supplemented with 3% BSA, 1% Triton X-100, and 1%

saponin. Secondary antibody diluted in the organoid washing solution was then added to the

organoids for 2 hours at 4°C. After 2 hours, organoids were again washed with organoid washing

solution. Phalloidin stain was added to the organoids for 60min, before DAPI stain was added to the

organoids for 5min. Imaging was performed with the Andor Dragonfly High-Speed Confocal

Microscope Systems (Oxford Instruments). Antibodies used were mouse anti-human FBXW7

antibody (1:100, H00055294-M02. Novus Biologicals, USA), rabbit anti-LY6D goat anti-mouse

(1:200, 17361-1-AP, Proteintech, USA), IgG H&L (Dylight 550) (1:500, ab96872, Abcam, USA), IgG

H&L (Alexa Fluor 488) (1:1000, Abcam, USA), Phalloidin iFluor 647 (1:400, ab176759, Abcam,

USA), and DAPI (300nM, D1306, Invitrogen, USA).

Co-culture of F, A and W organoids

Transduction of organoids was performed to achieve stable expression of GFP or mCherry

fluorescence. Lentiviral particles were first generated by transfection of HEK293T cells with

pMDLg/pRRE, pRSV-Rev, pSF-CMV-VSVG (Sigma-Aldrich) and either pCDH-EF1-copGFP-T2A-

Puro for GFP fluorescence or pCDH-EF1-copGFP-T2A-Puro for mCherry fluorescence.

pMDLg/pRRE, and pRSV-Rev were gifts from Didier Trono (Addgene plasmid # 12251;

http://n2t.net/addgene:12251 ; RRID:Addgene_12251) [32], pCDH-EF1-copGFP-T2A-Puro and

pCDH-CMV-mCherry-T2A-Puro were gifts from Kazuhiro Oka (Addgene plasmid # 72263 ;

http://n2t.net/addgene:72263 ; RRID:Addgene_72263).

To transduce organoids with the lentivirus particle, the organoids were first dissociated into single

cells by TrypLe for 10min in a 37°C water bath, with trituration at regular intervals. Single cells were

centrifuged at 400g for 6 min to pellet cells, which were then passed through a 50μm filter. The

number of cells was then counted using a hemocytometer, and the required number of cells was

aspirated, centrifuged and then pelleted. These cells were then resuspended in organoid growth media,

and viral particles were added to achieve a multiplicity of infection (MOI) of 20. Cells were infiltrated

over wells which had been overlaid with a pre-polymerised BME layer and left in a humidified 37°C

incubator for 24 hours. After 24 hours, dead cells were found floating above the BME, while live cells

would have become embedded in the BME. The media and dead cells were aspirated and fresh

organoid growth media was added. Fluorescence was checked using an EVOS M5000 cell imaging

system (ThermoFisher) after 72 hours.

For the co-culture of organoids, a GFP-labelled population was co-cultured with an mCherry-labelled

population in equal amounts to ascertain differences in proliferation. Cell culture wells were imaged

on day 0, 1, 3, 5 and 7 using an EVOS M5000 cell imaging system (ThermoFisher). ImageJ was used

to determine fluorescent intensity on each of these days.

Bulk RNA sequencing and analysis

Extraction of total RNA was performed using the RNeasy Mini Kit (Qiagen) as per manufacturer's

protocol. The quantity and quality of the RNA was ascertained using the Agilent 2100 Bioanalyzer.

Sequencing was performed by the Oxford Genomics Centre, Oxford using an Illumina NextSeq500

instrument using the standard paired-end protocol with a read length of 150bp. Fastq reads were

processed to clip low quality leading and trailing edges and to remove any adapter content using

Cutadapt (v.3.5). Quality checked FASTQ reads were aligned to the GRCh38 human genome and

gene annotation (Ensembl release 105) using STAR aligner (v.2.7.3a) two-pass mode to generate

gene-level quantification. Raw counts were processed in R (v.4.1.3) for all statistical testing and

plotting purposes. Normalisation and differential expression were performed using limma (v.3.50.3).

This included the generation of heatmaps using gplots (v.3.1.1) , volcano plots using ggplot2 (v3.3.5)

and PCA plots using ggfortify (v.0.4.14). Gene set enrichment analysis was performed using fgsea

(v.1.20.0) and GSVA (v.1.42.0).

Single-cell RNA sequencing and analysis

scRNA-seq was conducted using a 5' scRNA-seq gene expression workflow (10x Genomics). Cells

were loaded onto a 10x Chromium Controller for GEM generation followed by single cell library

construction using 10x Chromium Next GEM Single Cell 5' Library and gel bead kit v1.1 following

manufacturer's instructions. Size profiles of amplified cDNA and final libraries for sequencing were

verified by electrophoresis (Agilent 2100 Bioanalyzer system, High Sensitivity DNA Kit), and the

concentration of final libraries was measured with Qubit (Thermo Fisher Scientific). Libraries were

sequenced on an Illumina NextSeq 2000 (26 cycles read 1, 8 cycles i7 index, 98 cycles read 2),

achieving a mean of 39,412 reads per cell. Gene by cell barcode quantification matrix were obtained

using Cell Ranger (v.7.0.1) and Seurat (v.4.2.0) was used to perform normalisation and UMAP based

cluster assignment of pass cells.

Single sample gene set enrichment analyses (ssgsea) method from GSVA (v.1.42.0) was used on a

custom list of gene signature sets defined by editing the MSigDb (v.7.5.1) pathways to leave out

SPERMATOGENSIS, MYOGENESIS and PANCREAS_BETA_CELLS, and replacing the default

WNT_BETA_CATENIN_SIGNALING set by WNT_SIGNALING gene set (MSigDb systematic

name M5493), as per Househam et al. [33]. Additional gene sets included were – genes up- or down-

regulated in bulk RNA-seq comparison of 'FBXW7-APC' versus 'APC-FBXW7' organoids (adjusted

p-value < 0.05 and logFC >= |1.5|; n= 248 up- and n= 166 down-regulated genes), mouse foetal

gastric epithelium gene set from Vallone et al. 2016 (n= 122 human homologues), intestinal stem cell

signature from Merlos-Suárez et al. 2011 (n= 74), YAP pathway signature from Gregorieff et al. 2015

(n= 213 human homologues), crypt-base columnar (CBC) and regenerative stem cell (RSC) signature

sets from Vazquez et al. 2022 (n= 340 and 206 human homologues respectively)(Extended Data Table

3). Enrichment scores obtained per gene set was z-score transformed, and cells with scores greater

than 3^{rd} quartile or less than 1^{st} quartile of the score distribution were termed as enriched or depleted

respectively, for that gene set. To assess statistical enrichment or depletion of a given gene set in

FBXW7 mutant versus wildtype categories, chi-square test was applied on contingency table for cell

counts from enriched or depleted classes. Pearson residuals were visualised using mosaic plot from R

package vcd (v.1.4-11). For RNA velocity analysis, separate count matrices for spliced and unspliced

transcripts were created using Kallisto-Bustools (v.0.27.3) with the La Manno et al. 2018 strategy,

while using the same reference annotation from Cell Ranger (GRCh38, v.2020-A). Spliced/ unspliced

count data were combined with Seurat based UMAP clustering using scVelo (v.0.2.5) under python

v.3.10.6. RNA velocity graph computed was overlaid on UMAP cluster embeddings to infer trajectory

direction.

Proteomics

Four replicates per condition, each containing 50 µg protein, were solubilised in 5% SDS then

processed by S-Trap micro (Protifi) protocol according to the manufacturer's instructions. Digestion

was performed overnight with a 1:25 ratio of trypsin (Sequencing Grade, Promega) to protein. Tryptic

peptides were dried by vacuum centrifugation, then reconstituted prior to MS analysis in 0.1% formic

acid. LC-MS/MS analysis was performed using the Orbitrap Ascend Tribrid instrument (Thermo

Scientific) connected to a Thermo Scientific Vanquish[TM] Neo UHPLC system interfaced using a

nano-EASY spray source. The Vanquish[TM] Neo was operated in Trap and Elute mode using 0.1 %

Formic acid in water as solvent A, 0.1% formic acid in acetonitrile as solvent B and strong wash

buffer and 0.1% trifluoroacetic acid as weak wash. Tryptic peptides were loaded onto a PepMap Neo

C18 Trap (Thermo Scientific; S/N 174500) at 8 ul/min (total volume of 24ul) and separated on a 50C

heated EasySpray Pep Map Neo (Thermo Scientific; S/N ES75500) column using a multistep

gradient going from 2% to 18% solvent B in 40 min and from 18% to 35% solvent B in 20 min at

300nl/min flow rate. The column was washed for 14 min with 99% B, followed by the fast

equilibration on the Vanquish Neo (combine control mode, upper pressure to 1000bar). In parallel, the

trap was subjected to the zebra wash and fast equilibrated.

MS data were acquired in data-independent mode (DIA) with minor changes from previously

described method [34-36]. Briefly, MS1 scans were collected in the orbitrap at a resolving power of

45K at m/z 200 over m/z range of 350 – 1650m/z. The MS1 normalised AGC was set at 125%

(5e5ions) with a maximum injection time of 91 ms and a RF lens at 30%. DIA MS2 scans were then

acquired using the tMSn scan function at 30K orbitrap resolution over 40 scan windows with variable

width, with a normalized AGC target of 1000%, maximum injection time set to auto and a 30 %

collision energy.

Raw data were searched in DIA-NN v1.8.1 in library-free mode against the UniProt human proteome

database (UP000005640, downloaded 18[th] May 2023), plus common contaminants [37]. A maximum

of one missed cleavage was permitted for Trypsin/P digestion, with cysteine carbamidomethylation

set as a fixed modification. 'Match between runs' and retention time-dependent cross-run

normalisation were enabled, with mass accuracy settings inferred from the data. The DIA-NN neural

network classifier was set to double-pass mode. Further data analysis was performed in Perseus

v2.0.11, where protein groups were filtered to include only those identified in at least three replicates

of one experimental condition. Protein intensity values were log2-transformed, then missing values

were imputed with random values generated from a downshifted normal distribution (width 0.3,

downshift 1.8). Data are available via ProteomeXchange with identifier PXD052352.

To identify differential protein expression Student's t-test was applied on replicate level data from

FBXW7 mutant and WT samples, for each of the detected proteins using the "t.test" function. The p-

values thus obtained were corrected for multiple-testing using the Benjamini and Hochberg (i.e. FDR)

method by using the "p.adjust" function. All calculations were done in R ver. 4.3.3.

Assay for transposase-accessible chromatin sequencing (ATACseq) and analysis

Human colon organoids were generated as previously described and suspended in 40ul of BME in

400ul of complete organoid media. At Passage 2, organoids were subjected to electroporation based

CRISPR using either no guide, multiguide RNA targeting FBXW7 or single guide RNA targeting

APC (ref STAR protocols). Organoids targeted with guide RNA against APC were subsequently

grown in complete organoid media (also in STAR paper) omitting WNT3a conditioned media until 7

days prior to ATAC-sequencing. Organoids targeted with FBXW7 guide RNA underwent a second

round of electroporation based CRISPR at passage 3 to achieve >80% knockout of FBXW7. Wildtype

and knockout organoids were subsequently passaged to achieve 1×10^{12} cells, grown in identical

complete organoid media for 7 days and submitted for library preparation and ATAC-sequencing

(Genewiz/Azenta). ATAC experiments were carried out separately for W, A, F and Foetal (1) and W,

AF and FA (2). To ensure that background epigenetic status was corrected for, all CRISPR-edited

organoids for these experiments were generated from the same donor W organoids for each of the two

experiments.

Library preparation for W/FA/AF ATACseq was performed using the ATAC-Seq Kit (Active Motif) as

per the manufacturer's instructions. Multiplexing was performed using unique i7 and i5 indexed

primers. The quantity and quality of DNA was assessed using the Agilent High Sensitivity DNA Kit,

ensuring that transposed DNA fragments were between 200 and 1000bp, with a periodicity of ~200bp.

Sequencing was performed by the Oxford Genomics Centre, Oxford using an Illumina NextSeq2000

instrument.

Quality checked fastq reads were aligned to the human genome (GRCh38) using bwa aligner

(v.0.7.17). Alignment output (BAM files) were filtered, and TA (Tag Alignment) format data created

using samtools (v.1.14), picard (v.2.6.0) and bedtools (v.2.30.0). Further, MACS3 (v.3.0.0b2) was

used to perform peak calling which were then annotated using ChIPseeker (v.1.30.3). MACS3

identified peaks for the 'AF vs W' and 'FA vs W' comparisons were thresholded to select those with

score (i.e., -10*log(q-value)) >median of the respective distribution, overlapping regions merged and

the 50bp span from the centre of the peak region were extracted as input for *de novo* transcription

factor motif enrichment analysis using rGADEM (v.2.46.0). Identified motifs were plotted using

TFBSTools (v.1.36.0) and annotated for the closest matching core human transcription factors using

JASPAR2020 (v.0.99.0). Gene/ motif-set level visualisation of ATAC-seq signal enrichment was

performed using deeptools (v.3.5.2). Comparative bigwig files were created for 'AF vs W', 'FA vs W'

and 'FA vs AF' with BPM (Bins per million) normalisation, non-covered regions skipped, ENCODE

blacklist regions filtered out and effective genome size adjusted to reflect the filtering. The bigwig

files were then profiled for ATAC-seq signal enrichment with reference-point at the TSS

(Transcription Start Site) and upstream/ downstream 5kb region. Signal enrichments were profiled for

each of these sets –all genes in the genome (gencode v.38; 'Whole Genome'), the CBC gene signature

set ('CBC') and the 50bp genomic regions where de novo motif search identified TEAD1/2 motifs

(542 regions; 'TEAD1/2 motif') or SNAI1 motifs (491 regions; 'SNAI1 motif').

Comparative bigwig files were created from the final BAM files for 'F', 'A' and 'Foetal' organoids by

comparing each to the 'WT' organoid BAM, using 'bamCompare' from deeptools (v.3.5.2) with

parameters: binSize 10, normalisation BPM, ENCODE blacklist regions filtered out and effective

genome size adjusted to reflect the filtering. The bigwig files were then used to create the accessibility

tracks by extracting UCSC Genome Browser data (ver. GRCh38) and plotting it in R (v.4.3.3) using

bwtool (v.1.0) and trackplot (v1.5.10).

Validation of results in published patient datasets

Three independent published datasets were used to validate our findings. Somatic mutation and gene expression for the TCGA-COAD cohort were obtained from NCI-GDC data portal [27], to create a dataset of colon adenocarcinoma cases only from 320 unique patients. Per-sample enrichment scores for the gene sets was calculated using 'ssgsea' method in GSVA (v.1.42.0). The second validation made use of transcriptomic results derived from the EPICC dataset [9]. Here, individual gland-level transcriptomic results were obtained. Generation of PCA plots using ggplot2 (v3.3.5). The third S:CORT study provided clinical and molecular data for both ASCETIC analysis of adenomas and transcriptional analysis of predictors of neo-adjuvant chemotherapy (FOxTROT) [26].

Modified ASCETIC analysis

The Agony-based cancer evolution inference (ASCETIC) framework was employed to identify conserved patterns of driver gene mutations across a cohort of bulk sequenced adenomas [22]. For 91 adenomas, copy number and ploidy estimate were used to estimate cancer cell fraction of each single nucleotide variant at the single sample level. Functions from the ASCETIC package were used to calculate a partial ordered set of genes based on minimising agony across the entire cohort and generate a ranking estimate of genes containing single nucleotide variants. Genes containing the most commonly occurring single nucleotide variants were displayed graphically, with the y-axis representing overall probability of early mutation, based on recurrent high confidence ($p<0.05$) evolutionary steps and arrow thickness, representing recurrent evolutionary step number (Fig. 6b).

Acknowledgements

S.J.A.B. was supported by The Pharsalia Trust, UK & a Cancer Research UK Advanced Clinician Scientist Fellowship (C14094/A27178). D.K.H.C. was supported by Singapore Ministry of Health's National Medical Research Council under a research training fellowship (MOH-FLWSHP10may-0001). A.S. was supported by the UK Medical Research Council, Wellcome Investigator Award (219523/Z/19/Z). D.F-C was supported by an NIHR Academic Clinical Lectureship Award. The

Stratification in Colorectal Cancer Consortium (S:CORT) was funded by the Medical Research

Council and Cancer Research UK (MR/M016587/1). The human foetal material was provided by the

Joint MRC/Wellcome Trust (MR/R006237/1) Human Developmental Biology Resource

(www.hdbr.org) and we thank B. Crespo, Professor Copp and the entire HDBR team.

Author contributions

D.K.H.C. and S.J.A.B. conceived the overall experimental questions and design. D.K.H.C. performed

experiments, wrote the initial draft of the manuscript, and was supervised by S.J.A.B. A.M. and

S.J.A.B. performed bioinformatic analysis for the project. S.D.C. assisted with immunofluorescence

of organoids. Y. Z. generated plasmids for lentiviral labelling of organoids, performed the proteomics

experiment and W vs APC scRNAseq experiment. R.O. H.F. S.J. and X.L. assisted with scRNAseq.

J.H. and T.G. contributed the EPICC dataset and assisted with its analysis. J.B. performed the

ASCETIC and EPICC analysis and carried out the W, F and A ATACseq experiment. I.V. S.F. and R.F.

carried out proteomic analysis. D.F.C and A.S. assisted with foetal organoid experiments. S.J.A.B.

was overall in-charge of the project. All authors contributed to drafting and final approval of the

manuscript.

Competing interests

The authors declare no competing interests.

Materials and correspondence

Raw sequence data would be made available via the European Genome-Phenome Archive (EGA) and

proteomics via PRIDE, on publication of the manuscript. The mass spectrometry proteomics data

have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository with the

dataset identifier PXD052352. Please relay correspondence to Simon Buczacki.

Inclusion and ethics in global research

Research was conducted in accordance with principles of inclusion and good ethics in global research.

Code availability

All relevant scripts to replicate the analysis would be made available on Github on publication of the

manuscript.

References

1. Brunner, S.F., et al., *Somatic mutations and clonal dynamics in healthy and cirrhotic human liver.* Nature, 2019. **574**(7779): p. 538-542.

2. Colom, B., et al., *Spatial competition shapes the dynamic mutational landscape of normal esophageal epithelium.* Nat Genet, 2020. **52**(6): p. 604-614.

3. Martincorena, I., et al., *Tumor evolution. High burden and pervasive positive selection of somatic mutations in normal human skin.* Science, 2015. **348**(6237): p. 880-6.

4. Moore, L., et al., *The mutational landscape of normal human endometrial epithelium.* Nature, 2020. **580**(7805): p. 640-646.

5. Lee-Six, H., et al., *The landscape of somatic mutation in normal colorectal epithelial cells.* Nature, 2019. **574**(7779): p. 532-537.

6. Yeh, C.H., M. Bellon, and C. Nicot, *FBXW7: a critical tumor suppressor of human cancers.* Mol Cancer, 2018. **17**(1): p. 115.

7. Burdziak, C., et al., *Epigenetic plasticity cooperates with cell-cell interactions to direct pancreatic tumorigenesis.* Science, 2023. **380**(6645): p. eadd5327.

8. Johnstone, S.E., et al., *Large-Scale Topological Changes Restrain Malignant Progression in Colorectal Cancer.* Cell, 2020. **182**(6): p. 1474-1489.e23.

9. Heide, T., et al., *The co-evolution of the genome and epigenome in colorectal cancer.* Nature, 2022. **611**(7937): p. 733-743.

10. Chan, D.K.H., S.D. Collins, and S.J.A. Buczacki, *Generation and immunofluorescent validation of gene knockouts in adult human colonic organoids using multi-guide RNA CRISPR-Cas9.* STAR Protoc, 2023. **4**(1): p. 101978.

11. van Neerven, S.M., et al., *Apc-mutant cells act as supercompetitors in intestinal tumour initiation.* Nature, 2021. **594**(7863): p. 436-441.

12. Chan, D.K.H., et al., *Biallelic FBXW7 knockout induces AKAP8-mediated DNA damage in neighbouring wildtype cells.* Cell Death Discov, 2023. **9**(1): p. 200.

13. Drost, J., et al., *Sequential cancer mutations in cultured human intestinal stem cells.* Nature, 2015. **521**(7550): p. 43-7.

14.   Matano, M., et al., *Modeling colorectal cancer using CRISPR-Cas9-mediated engineering of*

human intestinal organoids. Nat Med, 2015. **21**(3): p. 256-62.

15.   Pikkupeura, L.M., et al., *Transcriptional and epigenomic profiling identifies YAP signaling as*

a key regulator of intestinal epithelium maturation. Sci Adv, 2023. **9**(28): p. eadf9460.

16.   Fernandez Vallone, V., et al., *Trop2 marks transient gastric fetal epithelium and adult*

regenerating cells after epithelial damage. Development, 2016. **143**(9): p. 1452-63.

17.   Gregorieff, A., et al., *Yap-dependent reprogramming of Lgr5(+) stem cells drives intestinal*

regeneration and cancer. Nature, 2015. **526**(7575): p. 715-8.

18.   Merlos-Suárez, A., et al., *The intestinal stem cell signature identifies colorectal cancer stem*

cells and predicts disease relapse. Cell Stem Cell, 2011. **8**(5): p. 511-24.

19.   Vasquez, E.G., et al., *Dynamic and adaptive cancer stem cell population admixture in*

colorectal neoplasia. Cell Stem Cell, 2022. **29**(8): p. 1213-1228 e8.

20.   Elmentaite, R., et al., *Cells of the human intestinal tract mapped across space and time.*

Nature, 2021. **597**(7875): p. 250-255.

21.   Qin, X., et al., *An oncogenic phenoscape of colonic stem cell polarization.* Cell, 2023.

186(25): p. 5554-5568 e18.

22.   Fontana, D., et al., *Evolutionary signatures of human cancers revealed via genomic analysis*

of over 35,000 patients. Nat Commun, 2023. **14**(1): p. 5982.

23.   Sole, L., et al., *p53 wild-type colorectal cancer cells that express a fetal gene signature are*

associated with metastasis and poor prognosis. Nat Commun, 2022. **13**(1): p. 2866.

24.   Ramos Zapatero, M., et al., *Trellis tree-based analysis reveals stromal regulation of patient-*

derived organoid drug responses. Cell, 2023. **186**(25): p. 5606-5619 e24.

25.   Pich, O., et al., *The mutational footprints of cancer therapies.* Nat Genet, 2019. **51**(12): p.

1732-1740.

26.   Morton, D., et al., *Preoperative Chemotherapy for Operable Colon Cancer: Mature Results of*

an International Randomized Controlled Trial. J Clin Oncol, 2023. **41**(8): p. 1541-1552.

27.   *TCGA-COAD.* 7 March 2023]; Available from: https://portal.gdc.cancer.gov/projects/TCGA-

COAD.

28.   Gardner, E.E., et al., *Lineage-specific intolerance to oncogenic drivers restricts histological*
706   *transformation.* Science, 2024. **383**(6683): p. eadj1415.

707   29.   Ortmann, C.A., et al., *Effect of mutation order on myeloproliferative neoplasms.* N Engl J
708   Med, 2015. **372**(7): p. 601-612.

709   30.   Hansen, S.L., et al., *An organoid-based CRISPR-Cas9 screen for regulators of intestinal*
710   *epithelial maturation and cell fate.* Sci Adv, 2023. **9**(28): p. eadg4055.

711   31.   Thirimanne, H.N., et al., *Global and context-specific transcriptional consequences of*
712   *oncogenic Fbw7 mutations.* Elife, 2022. **11**.

713   32.   Dull, T., et al., *A third-generation lentivirus vector with a conditional packaging system.* J
714   Virol, 1998. **72**(11): p. 8463-71.

715   33.   Househam, J., et al., *Phenotypic plasticity and genetic control in colorectal cancer evolution.*
716   Nature, 2022. **611**(7937): p. 744-753.

717   34.   Muntel, J., et al., *Comparison of Protein Quantification in a Complex Background by DIA*
718   *and TMT Workflows with Fixed Instrument Time.* J Proteome Res, 2019. **18**(3): p. 1340-1351.

719   35.   Dellar, E.R., et al., *Data-independent acquisition proteomics of cerebrospinal fluid implicates*
720   *endoplasmic reticulum and inflammatory mechanisms in amyotrophic lateral sclerosis.* J
721   Neurochem, 2024. **168**(2): p. 115-127.

722   36.   O'Brien, D.P., et al., *Structural Premise of Selective Deubiquitinase USP30 Inhibition by*
723   *Small-Molecule Benzosulfonamides.* Mol Cell Proteomics, 2023. **22**(8): p. 100609.

724   37.   Mellacheruvu, D., et al., *The CRAPome: a contaminant repository for affinity purification-*
725   *mass spectrometry data.* Nat Methods, 2013. **10**(8): p. 730-6.

 Extended data figure legends

Extended data figure 1: *FBXW7$^{-/-}$* organoids do not influence growth of neighbouring wildtype

organoids.

APC$^{-/-}$ cells act as supercompetitors which suppress the growth of neighbouring wildtype cells,

evidenced by proliferative advantage of A organoids over neighbouring W organoids (**a, c, d**). A

similar effect was not seen in F organoids co-cultured with W organoids (**b, e, f**). **g,** EdU assay for W

and F organoids showed no significant differences in proliferation rate. Volcano plot of co-cultured W

versus non-co-cultured W (**h**) and co-cultured F versus non-co-cultured F (**i**) organoids showed that

the effect of co-culture did not significantly influence the transcriptome. Positive fold change

represents upregulated in co-cultured. The values in this figure are mean ± sd, and statistical

significance was measured by unpaired t-test. All experiments were performed with n=3 biological

replicates.

Extended data figure 2: Mutational background influences the transcriptomic effect of an

***FBXW7* mutation.**

a, Western blots for beta-catenin and p53 in wildtype and CRISPR-edited organoid. Sanger traces for

edited and wildtype organoids as analysed by ICE tool (Synthego). **b**, RNAseq volcano plot

comparing transcriptomes of AF vs A organoids. Positive fold change represents upregulated in AF. **c**,

RNAseq volcano plot comparing transcriptomes of ATF vs AT organoids. Positive fold change

represents upregulated in ATF. All experiments were performed with n=3 biological replicates.

Extended data figure 3: Effect of epistasis not an artefact of CRISPR/Cas9 gene editing.

a, Brightfield microscopy of organoids which had been gene edited at the AAVS1 loci three times to

mimic triple-mutant ATF organoids. AAVS1 organoids did not adopt a cystic phenotype. Scale bar =

100um. **b**, RNAseq volcano plot comparing transcriptomes of AAVS1 vs W organoids. Positive fold

change represents upregulated in AAVS1 samples **c**, RNAseq volcano plot comparing transcriptomes

of ATF vs W organoids. Positive fold change is upregulated in ATF samples. All experiments were

performed with n=3 biological replicates. **d**, Principal component analysis of transcriptomes from all

gene-edited models identified by genotype and patient donor.

Extended data figure 4: Cluster 13 in F organoids was marked by EMT signature.

a, GSEA plot of the enrichment of the top 100 F specific genes identified through pseudobulk analysis

of scRNAseq data compared to ranked bulk RNAseq data derived from F vs W comparison. **b**, UMAP

scRNAseq plots comparing F and W organoids with the EMT signature colour coded. Red = enriched.

Blue = de-enriched

Extended data figure 5: GSEA of bulk RNAseq data of F vs W organoids using key pathways

derived from scRNAseq analysis.

GSEA plots for enrichment of the foetal signature [16] (**a**), and YAP pathway (**b**) in F organoids

compared with W organoids. **c**, UMAP scRNAseq plots from Elmentaite et al of human, colon adult

and foetal epithelial cells overlaid with enrichment for FvW signature, *ANXA13* and *OLFM4*

expression. Blue = enriched. **d**, Immunofluorescence for LY6D in F and W organoids (Blue = DAPI,

Green = anti-LY6D) Scale bars = 50um. GSEA plots for enrichments of the ISC (**e**) and CBC (**f**)

signatures in F compared to W organoids.

Extended data figure 6: GSEA showing greater enrichment of YAP signalling in FA compared to

AF organoids.

GSEA plots of bulk RNAseq data of FA vs W (**a**) and AF vs W (**b**) for the YAP pathway signature.

Extended data figure 7: GSEA and scRNAseq analysis of A organoids compared to W

GSEA plots of bulk RNAseq comparisons between A vs W for the adult intestinal stem cell signature

(**a**) and proliferative cancer stem cell signature (**b**). GSEA plots of bulk RNAseq comparisons

between FA vs AF for the proliferative cancer stem cell signature (**c**) and the foetal stem cell signature

(**d**). **e**, UMAP single cell RNAseq plots of A and W organoids overlaid with enrichment for the

proCSC signature (degree of red is equivalent to degree of enrichment).

Extended data figure 8: ATACseq profiles for accessibility of F vs W and FA vs AF signatures in W, A, F and Foetal organoids

a, ATAC accessibility profiles for genomic loci associated with gene in the FvW RNAseq signature. (blue = increased accessibility, red = decreased accessibility). **b**, ATAC accessibility profiles for genomic loci associated with top 200 genes in the FavAF RNAseq comparison. (blue = increased accessibility, red = decreased accessibility). **c**, Chromatin accessibility for the *LY6D* gene region (chr8:142781535-142792411) in 'A.vs.WT', 'F.vs.WT' and 'Foetal.vs.WT' comparisons from ATAC-seq, visualised as track plots after group autoscaling of the input data. The Y-axis notes the track label and the range of the values within the region visualised. The vertical red bar on the ideogram of the chr8 indicates the location of the region. The X-axis indicates the direction of the genome, and the numbers (bottom-most track) are 2kb-spaced positions on the chr8. The canonical protein-coding transcript of LY6D gene (NM_003695) is represented as the dark-blue boxes (exons, n=3) connected by arrows ('<') on horizontal bar (introns).

Extended Data Table 1: Reactome pathway analysis of F vs W proteomics

Extended Data Table 2: Primer and guide sequences for gene editing

Extended Data Table 3: Gene signatures used for GSEA and ssGSEA

Extended Data Table 4: Table of de novo transcription factor identification results in F, A and W organoids

Extended Data Table 5: GSEA results for Hallmark and stem cell signatures on pre-operative biopsies from the FOxTROT study

Fig. 1 Characterisation of W and F organoids
a, Brightfield microscopy of W and F organoids revealed no observable phenotypic differences. Scale bars represent 500µm. **b**, Western blot validation showing loss of FBXW7 in F organoids. **c**, Immunofluorescence of W and F organoids with DAPI nuclear stain (blue), F-actin (red), FBXW7 (orange). Scale bars represent 100µm. **d**, Western blot analysis of known ubqiutiination substrates of FBXW7. **e**, Volcano plot of bulk RNAseq of F vs W organoids revealed minimal differential expression of genes. Positive fold change = up in F. **f**, Volcano plot of proteomic MS analysis of F and W organoids. Positive fold change = up in F. **g**, GSEA comparing the top 200 proteins over expressed in F compared to the ranked gene list of differentially expressed transcripts between F and W organoids. **h**, Box and whisker plot for the recurrence of histone pathways in depleted Reactome pathways in the F vs W proteomics. Median +/- IQR. **i**, Chart detailing the enrichment scores for the top 10 Reactome pathways altered in F vs W proteomics. All experiments were performed with n=3 biological replicates (proteomics n=4).

Fig. 2 Effect of FBXW7 mutation is dependent on mutational background
a, Schematic showing the organoid models used. W, A and AT organoids were generated to reflect
normal, adenomatous and carcinomatous lesions. On each of these organoid types, an FBXW7
mutation was introduced to generate F, AF and ATF organoids respectively. **b**, Brightfield microsco-
py of A, AF, AT and ATF organoids. A organoids adopt a cystic phenotype which is representative of
increased proliferation. AF organoids appear smaller in size compared with A organoids. AT orga-
noids also recapitulate the cystic appearance of A organoids. ATF organoids appear less cystic
than AT organoids. Scale bars represent 100μm. **c**, Histogram showing the number of differentially
expressed genes on the addition of an FBXW7 mutation changes based on the mutational back-
ground. While 2 genes were differentially expressed between W and F, 87 and 294 genes were
differentially expressed between A and AF, and AT and ATF respectively. **d**, Hierarchical clustering,
based on gene expression data, of the different organoid types showed that W and F organoids
clustered most similarly together. All experiments were performed with n=3 biological replicates.

Fig. 3 Effect of FBXW7 mutation is dependent on the order of mutation
a, Schematic showing the organoid models used in this analysis. A and F organoids were generat-
ed separately. On the A organoid, an *FBXW7* mutation was introduced, generating AF organoids.
On the F organoid, an *APC* mutation was introduced, generating FA organoids. AF and FA orga-
noids possessed the same mutations but acquired the mutations in an inverse order. **b**, Brightfield
microscopy of A, AF, F and FA organoids. Although FA organoids acquired an *APC* mutation, this
did not recapitulate the cystic phenotype observed in A or AF organoids. Scale bars represent 100μ
m. **c**, Principal component analysis (PCA) showed not only that FA organoids clustered separately
from AF organoids, FA organoids clustered together with W and F organoids, while AF organoids
clustered closer to A organoids. **d**, Venn diagram depicting the overlap of differentially expressed
genes at cut-off log2FC ≤ -0.7 or ≥ 0.7 between AF vs W and FA vs W comparisons. 628 genes
were differentially expressed in both AF and FA organoids. **e**, Linked column graph of AFvW and
FAvW overlap genes (n=628) showed there was a reduction in the magnitude by which the genes
were differentially expressed. All experiments were performed with n=3 biological replicates.

Fig. 4 scRNAseq shows FBXW7 mutation induces a switch from an adult to foetal stem cell state

a, Uniform manifold approximation and projection (UMAP) analysis from scRNAseq of W and F organoids yielded 14 clusters. **b**, Cluster 13 was unique to F organoids. **c**, *FBXW7* is actively expressed in wildtype organoids. Intensity of blue reflects *FBXW7* level. **d**, RNA velocity analysis demonstrated that clusters 1, 8 and 13 were terminally differentiating subpopulations. Chi-square analysis of W and F organoids demonstrated that F organoids were enriched for **(e)** FA_up gene signature ($p < 2.22 \times 10^{-16}$), **(f)** foetal signature ($p = 9.4754 \times 10^{-11}$), **(g)** YAP pathway signature ($p = 3.2134 \times 10^{-7}$), and depleted for **(h)** intestinal stem cell signature ($p = 1.4887 \times 10{-6}$) and **(i)** crypt-based columnar stem cell signature ($p < 2.22 \times 10^{-16}$) Red = increased expression, Blue = decreased expresion. Data presented represents two biological replicates for each condition (n=4).

Fig. 5 ATACseq analysis demonstrates cell state changes are asscoiated with global and specific alterations in chromatin accessibility

a, Whole genome chromatin accessibility profiles for W, A, F and Fo organoids (blue = increased accessability, red = less accessability) **b,** Chromatin accessibility profiles of TEAD 1/2 motifs for W, A, F and Fo organoids (blue = increased accessability, red = less accessability). **c,** Chromatin accessibility profiles of SNAI1 motifs for W, A, F and Fo organoids (blue = increased accessibility, red = less accessibility). **d,** Whole genome chromatin accessibility profiles for F, AF and FA organoids (blue = increased accessibility, red = less accessibility). **e,** Chromatin accessibility profiles for CBC signature genes for W, AF and FA organoids (blue = increased accessability, red = less accessability). Motif plots of the two de novo motifs found enriched in the FA vs W organoid comparison were **(f)** "nCwGCmCwGn" and **(g)** "rGCAGGTGn". Chromatin accessibility profiles of TEAD1/2 **(h)** and SNAI1 **(i)** motifs between W, AF and FA organoids. All experiments were performed with n=3 biological replicates with n=4 for foetal samples.

Fig. 6 Patient data confirms FBXW7 mutation order-specific effects and identifies a novel chemotherapy predictive onco-foetal signature

a, PCA plot based on transcriptional grouping of 'normal' crypts containing F, A and no mutations. Mutations highlighted on plot. **b**, Plot representing ASCETIC analysis of 65 colonic adenomas from the S:CORT study. Arrow thickness represents strength of likelihood of mutation order for seven cancer genes sequenced. **c**, Linked column graph of GSEA NES for 73 Hallmark and stem cell signatures on FOxTROT trial pre and post treatment biopsies. Highlighted signatures significant on pre-treatment biopsies (p<0.05). **d**, Dot plot of enrichment scores in TCGA-COAD samples (n=320) for FA_UP and FA_DOWN signatures showing an inverse relationship (correlation score: -0.55, p< 2.2 x 10^{-16}). **e**, Waddington schematic of the consequences of mutation order on stem cell state. Adult stem cells (ISC/CBC) are driven down two axes: a proliferative cancer stem cell (proCSC) following APC mutation or an onco-foetal route following FBXW7 mutation.