Divergent Folding-Mediated Epistasis Among Unstable Membrane Protein Variants

Laura M. Chamness,[1] Charles P. Kuntz,[2] Andrew G. McKee,[1] Wesley D. Penn,[1] Christopher M. Hemmerich,[3] Douglas B. Rusch,[3] Hope Woods,[4,5] Dyotima,[1] Jens Meiler,[4,6] and Jonathan P. Schlebach[1,2*]

[1]*Department of Chemistry, Indiana University, Bloomington, Indiana, USA*
[2]*Department of Chemistry, Purdue University, West Lafayette, Indiana, USA*
[3]*Center for Genomics and Bioinformatics, Indiana University, Bloomington, Indiana, USA*
[4]*Department of Chemistry, Vanderbilt University, Nashville, Tennessee, USA*
[5]*Chemical and Physical Biology Program, Vanderbilt University, Nashville, Tennessee, USA*
[6]*Institute for Drug Development, Leipzig University, Leipzig, SAC, Germany*

Corresponding author E-mail address: jschleba@purdue.edu

Keywords: Epistasis, GnRHR, Membrane Protein Folding, Misfolding, Proteostasis

Abstract

Many membrane proteins are prone to misfolding, which compromises their functional expression at the plasma membrane. This is particularly true for the mammalian gonadotropin-releasing hormone receptor GPCRs (GnRHR). We recently demonstrated that evolutionary GnRHR modifications appear to have coincided with adaptive changes in cotranslational folding efficiency. Though protein stability is known to shape evolution, it is unclear how cotranslational folding constraints modulate the synergistic, epistatic interactions between mutations. We therefore compared the pairwise interactions formed by mutations that disrupt the membrane topology (V276T) or tertiary structure (W107A) of GnRHR. Using deep mutational scanning, we evaluated how the plasma membrane expression of these variants is modified by hundreds of secondary mutations. An analysis of 251 mutants in three genetic backgrounds reveals that V276T and W107A form distinct epistatic interactions that depend on both the severity and the mechanism of destabilization. V276T forms predominantly negative epistatic interactions with destabilizing mutations in soluble loops. In contrast, W107A forms positive interactions with mutations in both loops and transmembrane domains that reflect the diminishing impacts of the destabilizing mutations in variants that are already unstable. These findings reveal how epistasis is remodeled by conformational defects in membrane proteins and in unstable proteins more generally.

Impact Statement

Many of the synergistic interactions between mutations within a gene are modified in the context of unstable proteins in a manner that depends on how the variants promote misfolding in the cell.

Introduction

Mutational effects on protein stability have important consequences for evolution. Destabilized proteins misfold more often, which can attenuate their ability to function in the cell (Bloom et al., 2007; Marinko et al., 2019). Though natural selection typically maximizes fitness by incorporating mutations that produce new or improved functions, this optimization process is often hindered by the destabilizing effects of most random mutations (Bloom et al., 2007; Tokuriki et al., 2007). The cumulative energetic effects of these mutations on protein stability are generally capable of creating non-additive, context-dependent epistatic interactions (Gong et al., 2013; Olson et al., 2014; Starr and Thornton, 2016). Nevertheless, there are many different aspects of protein synthesis, folding, and assembly that are governed by distinct energetic constraints. While epistatic interactions arising from changes in protein stability have been previously characterized in soluble proteins (Gong et al. 2013; Olson et al. 2014; Faber et al. 2019; Nedrud et al. 2021), the impact of epistasis on membrane protein folding and stability has yet to be explored. Although many of the mechanistic underpinnings of pairwise epistasis in soluble proteins are likely to be generalizable to all proteins, membrane proteins undergo additional cotranslational folding reactions that are governed by mechanistically distinct kinetic and energetic constraints (Marinko et al., 2019; Schlebach and Sanders, 2016). Based on these considerations, we suspect that mutations in membrane proteins could potentially modify cotranslational processes in a manner that generates distinct epistatic interactions that bias their evolutionary pathways in unique ways.

Unlike water-soluble proteins, membrane proteins must be cotranslationally inserted into lipid bilayers (Stage I folding) in order to fold into their native structure (Stage II folding) (Popot and Engelman, 1990). In eukaryotes, Stage I folding is primarily facilitated by Sec61 and a spectrum of various other translocon complexes (Marinko

et al., 2019; Smalinskaite and Hegde, 2022). Ribosomes translating membrane proteins are targeted to this
translocon complex at the endoplasmic reticulum (ER) membrane, where the nascent transmembrane domains
(TMDs) partition into the lipid bilayer and establish their native topology relative to the membrane (Stage I)
(Pfeffer et al., 2016; Popot and Engelman, 1990; van den Berg et al., 2003). After translation is complete,
membrane proteins fold and assemble (Stage II) in a manner that is coupled to their passage through the
secretory pathway to the plasma membrane and/ or other destination organelles (Marinko et al., 2019; Popot
and Engelman, 1990; Wiseman et al., 2007). Though they are governed by distinct physicochemical constraints,
failures in either Stage I or Stage II folding are capable of increasing the proportion of nascent membrane proteins
that are retained in the ER and prematurely degraded (Marinko et al., 2019; Wiseman et al., 2007). We therefore
expect that epistatic interactions can arise from the cumulative energetic effects of mutations on either of these
processes. Given that three-dimensional protein structures are stabilized by similar energetic principles in water
and lipid bilayers, mutations that modify the fidelity of Stage II folding energetics should potentially generate
long-range epistatic interactions that are comparable to those observed in soluble proteins (Marinko et al., 2019).
By comparison, the topological transitions that happen during the early stages of membrane protein synthesis
may be coupled to neighboring loops and helices in a manner that can play a decisive role in the formation of
the native topology (Hessa et al., 2007; Öjemalm et al., 2012; White and Von Heijne, 2008). Based on these
mechanistic distinctions and their association with different components of the proteostasis network, we reason
that the effects of mutations on these two processes could potentially create complex and/ or high-order epistatic
interactions (Morrison et al., 2021; Sailer and Harms, 2017).
81

To gain insights into the mechanistic basis of epistatic interactions in membrane proteins, we surveyed the effects
of single and double mutants on the plasma membrane expression (PME) of the gonadotropin-releasing
hormone receptor (GnRHR), a G-protein coupled receptor (GPCR) involved in reproductive steroidogenesis
across many species (Janovick et al., 2013, 2006). We have previously demonstrated that various evolutionary
sequence modifications within mammalian GnRHRs have tuned its fitness by compromising the fidelity of Stage
I folding (Chamness et al., 2021). Nevertheless, it remains unclear how the inefficient cotranslational folding of
GnRHR reshapes its tolerance of secondary mutations. Here, we utilize deep mutational scanning (DMS) to
carry out a focused analysis of 251 mutations in the background of two mouse GnRHR (mGnRHR) variants that
selectively compromise either Stage I or Stage II folding. Our results show that mutations which generate Stage
I and Stage II folding defects form distinct epistatic interactions throughout this receptor. An unsupervised
learning analysis reveals how these interactions depend on both changes in stability and the topological context
of the mutation. Together, these findings suggest that the distinct biosynthetic mechanisms of integral membrane

Figure 1. Structural context of GnRHR background mutations. A homology model of *M. musculus* GnRHR (mGnRHR) and a crystal structure of *H. sapiens* GnRHR (hGnRHR) provide structural context for two mutants which disrupt folding. A) The homology model of mGnRHR (green) structurally aligns with the crystal structure of hGnRHR (gray). B) A side view of hGnRHR shows that T277, which is analogous to V276 in mGnRHR, is located in transmembrane domain 6 and is exposed to the lipid bilayer. C) A top view of hGnRHR shows that W107 in extracellular loop 1 forms a hydrogen bond network with two other residues.

94 proteins may differentially shape their fitness landscapes. The general
95 implications of our findings for the role of protein folding and stability
96 in protein evolution are discussed.
97

Results
99

Mutational Library Design
101

To compare how perturbations of Stage I and Stage II folding shape
epistatic interactions, we first identified and characterized two
individual mutations that are likely to selectively disrupt either the
cotranslational membrane integration (V276T) or the native tertiary
structure (W107A) of mGnRHR. We previously showed that T277 in
TMD6 of human GnRHR (hGnRHR) contributes to the inefficient
translocon-mediated membrane integration of TMD6, which
decreases the receptor's PME (Chamness et al., 2021). A recent
crystallographic structure of hGnRHR confirms that the polar side
chain of T277, which corresponds to V276 in the mouse receptor
(89.3% sequence identity), is exposed to the lipid bilayer and does not
appear to make interhelical contacts that stabilize the native structure
(Fig. 1 A & B) (Yan et al., 2020). Thus, mutations at this position
modify the hydrophobicity of TMD6 without perturbing the native
tertiary contacts that stabilize the native fold. Indeed, replacing the
native valine residue in TMD6 of the mouse receptor with the
threonine residue found in hGnRHR reduces the efficiency of its
translocon-mediated membrane integration *in vitro* (Fig. S1, Table
S1). Furthermore, this substitution decreases the PME of the mouse
receptor by 65 ± 4% relative to that of wild-type mGnRHR (WT, Fig.

Figure 2. Plasma Membrane Expression of mGnRHR variants. Surface immunostaining and flow cytometry were used to compare the plasma membrane expression of select mGnRHR variants expressed in HEK293T cells. A histogram from one representative biological replicate depicts the distribution of plasma membrane expression intensities among cells expressing WT (black), V276T (red), or W107A (blue) mGnRHR.

2). Consistent with previous findings, these results confirm that the inefficient translocation of this helix promotes
cotranslational GnRHR misfolding (Chamness et al., 2021). In contrast, substitutions at W107 within extracellular
loop 1 (ECL1) preserve the hydrophobicity of TMDs and their topological energetics while disrupting a conserved
hydrogen bond network that is found in a wide array of GPCR structures (Fig. 1C) (Jones et al., 2020). Mutating
this conserved tryptophan to alanine (W107A) reduces the PME of mGnRHR by 88 ± 4% relative to WT (Fig. 2),
which suggests that disrupting this hydrogen bond network destabilizes mGnRHR in a manner that promotes its
cellular misfolding. Comparing how these mutations modify the proteostatic effects of secondary mutations will
therefore reveal how changes in the fidelity of Stage I (V276T) and Stage II (W107A) folding differentially impact
mutational epistasis.

To survey the epistatic interactions formed by these mutations, we generated a series of genetic libraries
consisting of 1,615 missense variants in the background of V276T, W107A, and WT mGnRHR. To ensure
adequate dynamic range in the downstream assay, we created these variants in the cDNA of mGnRHR, which
exhibits intermediate, tunable expression (Chamness et al., 2021). Briefly, we used a structural homology model
of mGnRHR to select 85 residues distributed across the loops and helices of mGnRHR. We included all 19
amino acid substitutions at the 3 most solvent-accessible and the 3 most buried residues in each TMD (Fig. S2
and Table S2). We also included mutations encoding all 19 amino acid substitutions at positions that are evenly
distributed across each soluble domain. We then generated an array of mutagenic oligonucleotides encoding
each of these mutations (Table S3), which was used to introduce these mutations in the context of the V276T,
W107A, and WT mGnRHR cDNAs by nicking mutagenesis (Medina-Cucurella et al., 2019). We generated each
of these libraries in the context of plasmids containing a randomized ten-base region within the backbone. Unique
10-base plasmid identifiers that could be matched to specific mGnRHR variants of interests were used to score
variants in the downstream assay using Illumina sequencing. Using whole plasmid PacBio sequencing of each
library, we matched 1,383 of the 1,615 possible variants to one or more ten-base unique molecular identifier
(UMI) in at least one of the three libraries, 320 of which were found in all three genetic backgrounds (see *Methods*
for details).

Plasma Membrane Expression of GnRHR Mutant Libraries

Figure 3. Plasma membrane expression of mGnRHR cellular libraries. mGnRHR mutant libraries (red), as well as their respective controls (WT or V276T/W107A single mutants, black) were expressed in mammalian cells, and the amount of mGnRHR expressed at the plasma membrane was measured during fluorescence-activated cell sorting. Histograms show the distributions of plasma membrane expression for mGnRHR mutants in the (A) WT library, (B) V276T library, and (C) W107A library.

To identify secondary mutations that form epistatic interactions with V276T and W107A, we measured the
relative PME of single and double mutants within each genetic library by DMS as previously described (Penn et
al., 2020). Briefly, we first used the genetic libraries described above to generate three recombinant, pooled
HEK293T cell lines in which the individual cells each express one of the single or double mGnRHR mutants of
interest. A comparison of the mGnRHR surface immunostaining profiles of these cellular populations that
recombinantly express this collection of variants in context of the V276T, W107A, or an otherwise WT genetic
background reveals striking context-dependent differences in their proteostatic effects. Recombinant cells
expressing these variants in the WT background exhibit a bimodal distribution of mGnRHR surface
immunostaining, where 59% of cells exhibit immunostaining that is comparable to WT and 41% of cells exhibit
reduced expression relative to WT (Fig. 3A). Although cells expressing this same collection of variants in the
V276T background also exhibit bimodal surface immunostaining, 48% of the population exhibit diminished
expression relative to the V276T variant (Fig. 3B). This observation suggests many of these variants
synergistically decrease mGnRHR expression in combination with the V276T mutation. Unlike the WT and
V276T libraries, cells expressing this same collection of variants in combination with the W107A mutation exhibit
a continuous range of surface immunostaining intensities that are only slightly lower, on average, relative to cells
expressing the W107A single mutant (Fig. 3C). The lack of dispersion likely reflects a compression of the
distribution that arises as the folding efficiency of these variants approaches zero (see *Discussion*). Cells
expressing this library of W107A double mutants do, however, still contain cellular subpopulations with surface
immunostaining intensities that are well above or below that of the W107A single mutant, which suggests that
this fluorescence signal is sensitive enough to detect subtle differences in the PME of these variants (Fig. 3C).
Indeed, the mean fluorescence intensity associated with the surface immunostaining of the WT, V276T, and
W107A mutant libraries were respectively 33.5-fold, 13.5-fold, and 2.6-fold higher than the background, on
average, across the two biological replicates. Overall, these results suggest that none of the secondary mutations
can fully compensate for the proteostatic effects of the W107A mutation.
175
To compare the PME of these variants in each genetic background, we fractionated each of the three mixed cell
populations into quartiles based on the relative surface immunostaining of their expressed mGnRHR variants
using fluorescence-activated cell sorting (FACS), then extracted the genomic DNA from each cellular fraction.
We then used Illumina sequencing to track the relative abundance of the recombined UMIs and their associated
variants within each fraction and used these measurements to estimate the corresponding PME of each variant.
A series of filters relating to the quality of the reads, the sampling of each mutation, and the similarity of variant
scores across replicates were used to remove poorly sampled variants with unreliable scores (see *Methods*).
Briefly, to restrict our analysis to variants that can be reliably scored, we removed variants that did not have at
least 50 counts across the four bins in each replicate. Additionally, we compared the percentile rank of each
variant across replicates and discarded variant scores for which the difference in percentile rank across the two
replicates was greater than 25%. The relative PME of the remaining variants was averaged across two biological
replicates. Overall, we measured the relative PME of 1,004 missense variants in the WT background, 338

Figure 4. Relative plasma membrane expression of mGnRHR mutants. A) A histogram shows the distribution of plasma membrane expression measurements relative to their respective controls (WT or V276T/W107A single mutants) for the WT (black), V276T (red) and W107A (blue) mutational libraries. A dashed line at the normalized plasma membrane expression value of 1.0, representing no mutational effect relative to the control, is displayed for reference. B & C) The average relative intensities across each domain are compared between the V276T (B, red) and W107A (C, blue) double mutant libraries and the single mutant library (black). The size of each dot represents the number of variants measured. D – F) Average relative intensities for the (D) WT library, (E) V276T library, and (F) W107A library were projected onto a homology model of mGnRHR. Residues 276 and 107 are displayed in green for reference. Individual variant scores can be found in Table S4.

missense variants in the V276T background, and 1,179 missense variants in the W107A background (Table S4).
Out of the 320 variants identified in all three plasmid libraries by PacBio sequencing, 251 mutations passed these
quality filters in the Illumina sequencing of the cellular isolates (Table S4). The intensity values for these variants
were highly correlated across independent biological replicates, which demonstrates that these measurements
exhibit precision despite variations in the signal-to-noise associated with their surface immunostaining (Fig. S3).
In the following, we will focus on the comparison of the effects of these 251 mutations in each background in
order to determine how their proteostatic effects are modified by secondary mutations.
195
To facilitate the comparison of estimated immunostaining intensities for each variant across the three genetic
backgrounds, we normalized the raw immunostaining intensity values for each variant relative to that of the
measured intensity values for the corresponding reference sequence (V276T, W107A, or WT mGnRHR).
Relative intensity values of 1.0 correspond to no effect, whereas values over 1.0 correspond to mutations that
enhance surface expression in that genetic background, and vice versa. A histogram of the relative intensity
values for the 251 variants measured in all three backgrounds generally recapitulates the trends in the cellular
immunostaining histograms (Fig. 3 & 4A). Most of these mutations decrease the PME of mGnRHR in all three
backgrounds, which reflects the limited mutational tolerance of membrane proteins (Fig. 4A) (Telenti et al., 2016).
Notably, only a modest fraction of mutations measurably enhance the surface expression of WT mGnRHR (51
mutations, 20%) and V276T mGnRHR (21 mutations, 8% Fig. 4A). Mutations in the V276T background tend to
decrease surface expression more than they do in the WT background, and this trend persists across most
domains of the protein (Fig. 4 A & B). In contrast, a larger proportion of these mutations (100 mutants, 40%)
enhance the surface expression of W107A (Fig. 4A), though these increases in immunostaining are relatively

modest (Fig. 3C). Furthermore, the mutations tend to be better tolerated
with respect to surface expression in combination with W107A relative
to their effects on expression in the WT background (Fig. 4C).
212
Projecting the average relative intensities for the mutations at each
residue onto the structure reveals that most positions exhibit similar
trends in the mutational tolerance in each background, with loop
residues being generally more permissive than those within TMDs (Fig.
4 D – F). Nevertheless, there are subdomains where the quantitative
mutagenic effects deviate in a context- dependent manner. Variants
bearing mutations within the C-terminal regions including ICL3, TMD6,
and TMD7 fare consistently worse in the V276T background relative to
WT (paired Wilcoxon-Signed Rank Test *p*-values of 0.0001, 0.02, and
0.005, respectively) (Fig. 4 B & E). Given that V276T perturbs the
cotranslational membrane integration of TMD6 (Fig. S1, Table S1), this
directional bias potentially suggests that the apparent interactions
between these mutations manifest during the late stages of
cotranslational folding. In contrast, mutations that are better tolerated in

Figure 5. Epistasis in the mGnRHR double mutant libraries. A histogram depicts the distribution of epistasis scores (Ɛ) for interactions the subset of 251 high quality secondary mutations form with V276T (red) and W107A (blue).

227 the context of W107A mGnRHR are located throughout the structure but are particularly abundant among
228 residues in the middle of the primary structure that form ICL2, TMD4, and ECL2 (paired Wilcoxon-Signed Rank
229 Test *p*-values of 0.0005, 0.0001, and 0.004, respectively) (Fig. 4 C & F). Together, these observations show how
230 the proteostatic effects of mGnRHR mutations are modified by secondary mutations that differentially affect the
231 fidelities of Stage I and Stage II folding.
232

Distinct Pairwise Epistasis in V276T and W107A

To compare epistatic trends in these libraries, we calculated the magnitude of the epistatic interactions these
251 mutations form with V276T and W107A by comparing their relative effects on PME of the WT, V276T, and
W107A variants using a previously described relative epistasis model (product model, Olson et al. 2014). Briefly,
we defined fitness as the ratio of the single/ double mutant PME to that of WT mGnRHR. Epistasis scores (Ɛ)
were then determined by taking the log of the double mutant fitness value divided by the difference between the
single mutant fitness values (see *Methods*). Positive Ɛ values denote double mutants that have greater PME
than would be expected based on the effects of single mutants. Negative Ɛ values denote double mutants that
have lower PME than would be expected based on the effects of single mutants. Pairs of mutations with Ɛ values
near zero have additive effects on PME. For most double mutants, epistasis scores are near zero, suggesting
that many of the 251 random mutations have additive effects on PME in each background. This result is
consistent with evidence that epistatic interactions are typically rare in the context of stable proteins (Starr and
Thornton, 2016). Nevertheless, the distributions of epistasis scores for the two double mutant libraries are shifted
in opposite directions. The mutations generally form positive epistatic interactions with W107A and negative
epistatic interactions with V276T (Fig. 5). Interestingly, the difference in the epistasis scores for the interactions
these variants form with W107A and V276T was at least 1.0 for 98 of the 251 variants (Table S4). Notably, these
98 variants are enriched with TMD variants (65% TMD) relative to the overall set of 251 variants (45% TMD,
Fisher's Exact Test *p* = 0.0019). These findings suggest random mutations form epistatic interactions in the
context of unstable mGnRHR variants in a manner that depends on the specific folding defect (V276T vs.
W107A) and topological context.

Molecular Basis for the Observed Epistatic Interactions

To identify general trends in the observed epistatic interactions, we utilized unsupervised learning to elucidate
patterns among the relative PME values of variants across each genetic background. We first utilized uniform
manifold approximation projection (UMAP) (McInnes et al., 2018) to identify mutations that have similar
expression profiles across these conditions. A projection of the variants onto a resulting two-dimensional
coordinate reveals that most variants fall into one of approximately three groups (Fig. 6A). Using a density-based
hierarchical clustering analysis (HDBSCAN), we unambiguously assigned 182 variants into three distinct clusters
based on their expression profiles alone (Fig. 6A, Table S4) (Campello et al., 2013). An analysis of the structural
context of these mutants reveals that one of the largest clusters (Cluster 3, n = 76) primarily consists of mutations
of soluble loop residues that have little impact on expression and exhibit minimal epistasis (Fig. 6 A-C). This
analysis also identified a smaller cluster of loop mutations (Cluster 2, n = 30) that moderately decrease

Figure 6. General trends in mGnRHR epistasis. Trends associated with the observed pairwise epistasis within mGnRHR are identified using unsupervised learning. A) Uniform manifold approximation projection (UMAP) was used to differentiate variants based on differences in their relative expression in the V276T, W107A, or WT background. Variants are projected onto an arbitrary two-dimensional coordinate based on the results and are colored according to whether they were assigned to cluster 1 (green), cluster 2 (purple), cluster 3 (orange), or were designated as outliers (gray) by HDBSCAN. The percentage of the mutations that fall within TMDs or loops are shown for reference. B) A box and whisker plot depicts the statistical distributions of relative PME values among variants within each cluster in the context of V276T (red), W107A (blue), or WT (Gray) mGnRHR. Select clusters of variants that exhibit statistically different expression profiles according to a Mann-Whitney U-test are indicated (*, $p < 0.001$). A value of 1 corresponds to mutations that have no effect on the PME of mGnRHR in the indicated genetic background. C) A box and whisker plot depicts the statistical distribution of epistasis scores associated with the interactions between the mutations within each cluster and either V276T (red) or W107A (blue). *P*-values for select Mann-Whitney U-tests comparing the interactions of these mutations with V276T and W107A are indicated. A value of 0 indicates that the effects of the two mutations are additive. D) A box and whisker plot depicts the statistical distribution of Rosetta ΔΔG values among mutations within each cluster. *P*-values for select Mann-Whitney U-tests comparing the ΔΔG values across clusters are shown for reference. For panels B-D, the edges of the boxes correspond to the 75th and 25th percentile values while the whiskers reflect the values of the 90th and 10th percentile. The central hash and square within the box represent the average and median values, respectively. These analyses were carried out on a subset of 243 variants with high-quality expression measurements with calculable Rosetta ΔΔG values.

expression and exhibit a greater propensity to form epistatic interactions with V276T and/ or W107A relative to
the neutral loop mutations in Cluster 3 (Fig. 6 A-C). Nevertheless, most of the mutations that exhibit strong
positive epistatic interactions with W107A fell into Cluster 1 (n = 76), which primarily consists of mutations within
TMDs that significantly decrease expression in all three genetic backgrounds (Fig. 6 A-C). A comparison across
these clusters suggests positive epistatic interactions with W107A primarily stems from the fact that mutations
that compromise expression can only cause a relatively modest reduction in the PME of W107A GnRHR (Fig. 6
B & C), which is already predominantly misfolded. In contrast, the moderately disruptive loop mutations in Cluster
2 tend to exhibit more pronounced negative epistatic interactions with V276T than do the highly disruptive TMD
mutations in Cluster 1 (Fig. 6C). The divergent epistatic interactions that disruptive loop and TMD mutations form
with V276T potentially arise from differences in the mechanistic basis for the destabilization caused by these two
classes of mutations (see *Discussion*).
278

Many epistatic interactions arise from the additive energetic effects of mutations on protein folding (Olson et al.,
2014; Tokuriki and Tawfik, 2009). To determine how these trends relate to the effects of mutations on
thermodynamic stability, we utilized Rosetta CM to generate structural models of 243 of the 251 variants that
were scored in all three backgrounds and fall within the structured regions of the receptor (Song et al., 2013).
We then utilized a membrane-optimized Rosetta energy scoring function to approximate the change in the
energy of the native structural ensemble of each variant (Alford et al., 2015). The cluster of loop mutations with
lower expression (Cluster 2, avg. ΔΔG = + 2.2 REU) generally contains more destabilizing mutations relative to
the cluster of loop mutations with WT-like expression (Cluster 3, avg. ΔΔG = + 0.8 REU), which affirms the
general relationship between stability and PME in this system (p = 0.003, Fig. 6 B & D). However, the cluster of
TMD variants (Cluster 1, avg. ΔΔG = + 3.4 REU) is generally more destabilizing than either of the loop clusters
(Fig. 6D). A comparison across the clusters suggests that the degree of destabilization generally tracks with the
magnitude of positive epistatic interactions that form with W107A, regardless of structural context (Fig. 6 C & D).

This uptick in positive epistasis among mutations that destabilize
the native fold and severely reduce PME likely reflects the
attenuated effect of destabilizing mutations as the equilibrium
fraction of folded protein approaches zero (see *Discussion*, Fig. 7).
In contrast to W107A, epistatic interactions with V276T are
predominantly negative and are most pronounced among
moderately destabilizing loop mutations (Fig. 6 C & D). The
abundance of negative interactions with V276T parallels the
observed pairwise epistasis in other folded proteins (Olson et al.,
2014), and likely reflects the higher baseline expression and/ or
stability of this variant relative to W107A (Figs. 2 & 7).
Nevertheless, stability-mediated epistasis cannot account for the
attenuation of the epistatic interactions that V276T forms with the
highly destabilizing mutations within Cluster 1 relative to the
moderately destabilizing mutations within Cluster 2 (Fig. 6 C & D).
These observations suggest that the magnitude of the epistatic
interactions formed by V276T is dependent on not only the change
in stability, but also the topological context of the secondary
mutation.
310

Discussion

Many epistatic interactions between mutations within genes
encoding soluble proteins arise from their cumulative effects on
protein folding energetics (Nedrud et al., 2021; Olson et al., 2014).
Folded proteins can only tolerate a limited number of destabilizing
mutations before their cumulative energetic effects cause a
cooperative decrease in the fraction of folded protein (Fig. 7A).
Although this general principle should apply to all proteins, it is
unclear how the distinct energetic processes involved in membrane
protein biosynthesis and folding may alter such evolutionary
couplings. In this study, we utilize DMS to compare the epistatic
interactions formed by two mutations that reduce the expression of
a misfolding-prone GPCR by selectively destabilizing either its
cotranslational membrane integration (V276T) or its native tertiary
structure (W107A). We note that the introduction of the V276T
mutation into the mouse receptor utilized herein mimics one of the
key destabilizing modifications that reduces the expression of
hGnRHR relative to mGnRHR (Chamness et al., 2021). Though the
quantitative energetic effects of these mutations on cotranslational
and/ or post-translations folding processes remains unclear in the
absence of a suitable biophysical assay (see *Response to the*
Reviewers), they each markedly reduce the plasma membrane
expression of mGnRHR and it seems highly unlikely that they
should elicit identical quantitative effects on folding and assembly
(Fig. 2). Thus, a comparison of the epistatic interactions formed by
these two mutations should reveal the extent to which the

Figure 7. Context-dependent impacts of stabilizing and destabilizing mutations. Line plots depict the relationship between the equilibrium fraction of folded protein (f_{fold}) and the free energy of unfolding (ΔG_{unf}). A) A moderately stable protein (ΔG_{unf} = +2 kcal/mol) exhibits a greater change in f_{fold} after acquiring one or more destabilizing mutations (total $\Delta\Delta G_{unf}$ = -2 kcal/mol, red) than after acquiring one or more mutations that stabilize the protein to the same extent (total $\Delta\Delta G_{unf}$ = +2 kcal/mol, blue). B) A moderately unstable protein (ΔG_{unf} = -2 kcal/mol) exhibits a greater change in f_{fold} after acquiring one or more stabilizing mutations (total $\Delta\Delta G_{unf}$ = +2 kcal/mol, blue) than after acquiring one or more mutations that destabilize the protein to the same extent (total $\Delta\Delta G_{unf}$ = -2 kcal/mol, red).

mechanism of misfolding impacts the propensity for genetic interactions between mutations. Through a focused
analysis of 753 total single and double mutants, we find that these mutations give rise to divergent epistatic
interactions that modify the PME of mGnRHR (Fig. 5A). Mutations that form epistatic interactions with V276T
tend to synergistically decrease mGnRHR expression (Figs. 3 A & B, 4B, and 5A)- a typical epistatic trend that
likely arises from the combined destabilizing effects of both mutations. Surprisingly, these interactions are most
prominent among moderately destabilizing loop mutations (Fig. 6C). In contrast, many of these same secondary
mutations instead form positive epistatic interactions with W107A mGnRHR, regardless of their topological
context (Figs. 5A & 6C). As this comparison reflects the epistatic interactions formed by an identical set of
secondary mutations, we conclude that these context-dependent epistatic interactions arise from differences in
the magnitude and/ or the mechanism of the mGnRHR destabilization caused by the V276T and W107A

mutations. These observed trends potentially reflect a previously undescribed manifestation of "ensemble
epistasis" (Morrison et al., 2021) in which genetic interactions arise from the differential effects of mutations on
the accumulation of distinct misfolded conformations that vary with respect to their recognition and/ or
degradation by cellular quality control pathways.
352

In the context of stable proteins, stability-mediated epistasis tends to magnify the effects of deleterious mutations
due to the cumulative destabilization that arises from sequential random substitutions (Tokuriki et al., 2007). This
is a fundamental consequence of the cooperative dependence of the fraction of folded protein on the free energy
of unfolding. Indeed, most of the random mGnRHR mutations characterized herein are predicted to destabilize
the native structure in a manner that coincides with a measurable decrease in the PME in the context of all three
genetic backgrounds (Figs. 4A & 6D). Whereas these destabilizing mutations exhibit the expected negative
epistatic couplings with V276T, their interactions with W107A skew positive (Fig. 5A). This anomalous positive
epistasis is potentially a byproduct of the severe destabilizing effects of W107A relative to V276T (Fig. 2). While
destabilizing secondary mutations can cause significant decreases in the yield of folded WT or V276T mGnRHR,
the W107A variant already exhibits marginal surface immunostaining. In the context of this predominantly
misfolded, poorly expressed variant, even highly destabilizing secondary mutations can only cause a relatively
small decrease in the yield of folded protein. This observation can be more generally related to the relatively
shallow dependence of the fraction of folded protein on the free energy of folding under conditions where folding
becomes unfavorable- unfolding transitions become shallow as the fraction of folded protein approaches zero
(Fig. 7B). Stabilizing mutations can also generate potent positive epistatic couplings in this context given that
even small increases in stability can push the system into the steepest portion of the transition zone (Fig. 7B).
Paradoxically, the asymmetry of the folding curve in this regime suggests destabilizing mutations are more
tolerable in the context of an unstable protein- the effects of these variants exhibit "diminishing returns" (Otto and
Feldman, 1997) because folding efficiency effectively has nowhere to go but upwards. Essentially, the nature of
this sigmoidal transition suggests the typical trends associated with stability-mediated epistasis should become
inverted as stability decreases. Such considerations could potentially expand the accessible sequence space in
metastable proteins, which constitute a significant portion of the proteome (Ghosh and Dill, 2010; Zeldovich et
al., 2007). Combinations of mutations that would not generally be tolerated in stable proteins may also occur
through such mechanisms in the context of the numerous misfolded secondary alleles that are carried within the
genomes of humans and other diploids with no appreciable fitness cost (e.g. the ΔF508 variant of *CFTR*).
378

Unlike W107A, the epistatic interactions formed by V276T appear to depend on their topological context-
negative epistasis is more prevalent among moderately destabilizing loop variants than highly destabilizing TMD
variants (Fig. 6 C & D). This observation suggests stability-mediated epistasis between mutations that destabilize
the native tertiary structure of integral membrane proteins (Stage II folding) may be distinct from the interactions
that arise from the disruption of their translocon-mediated cotranslational membrane integration (Stage I folding).
This divergence could potentially reflect the distinction between the energetic principles of protein folding relative
to those that guide the insertion of proteins into membranes, which is fundamentally driven by a membrane
depth-dependent solvation energetics (Hessa et al., 2007; Moon and Fleming, 2011). Counterintuitively, polar
mutations that disrupt the membrane integration of individual TMDs also have the potential to form interhelical
hydrogen bonds that can stabilize the topological orientation of neighboring TMDs, which can give rise to long-
range topological couplings. (Hermansson and von Heijne, 2003; Meindl-Beinker et al., 2006; Öjemalm et al.,
2012). The propensity of such contacts to form may in some cases depend on the length and hydrophobicity of
the loops that connect them (Woodall et al., 2017). Finally, we should note that complex epistatic interactions
between residues within TMDs could potentially arise through their propensity to remodel the translocon complex
and/ or other quality control interactions- polar residues within TMDs may affect the recruitment of secondary
insertases and/ or intramembrane chaperones such as TRAP, the PAT complex, the BOS complex, the GEL
complex, and the ER membrane protein complex (Chen et al., 2023; Gemmer et al., 2023; Matreyek et al., 2017;
Pleiner et al., 2020; Shurtleff et al., 2018; Smalinskaitė et al., 2022; Sundaram et al., 2022). Additional insights
into the mechanistic basis for the recruitment of such complexes may therefore be needed to fully rationalize
evolutionary couplings between TMDs. We note that, more generally, we suspect that synergistic genetic
interactions between mutations could potentially arise from changes in the energetics of many other biosynthetic
processes such as cofactor binding, oligomer formation, and chaperone interactions (Bershtein et al., 2013;
Rodrigues et al., 2016).
402

Together, our results reveal divergent, mechanism-dependent patterns of pairwise epistasis in the context of a
misfolding-prone integral membrane protein. These findings suggest the folding-mediated epistasis is likely to
vary among different classes of destabilizing mutations in a manner that should also depend on folding efficiency

and/ or the mechanism(s) of misfolding in the cell. These findings provide novel insights as to how the unique
biosynthetic constraints of integral membrane proteins can potentially lead to unique evolutionary patterns.
Moreover, our results generally suggest that the deleterious effects of destabilizing mutations are partially
attenuated in the context of unstable proteins. Additional investigations are needed to explore how stability-
mediated epistasis is modified by other proteostatic constraints, such as the interactions of nascent proteins with
cofactors and/ or molecular chaperones.
412

Materials & Methods

413
414

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
gene (*Mus musculus*)	GnRHR	GenBank	L01119	-
gene (*Homo sapiens*)	GnRHR	GenBank	L03380	-
strain, strain background (*Escherichia coli*)	NEB 10-β	New England Biolabs	C3020K	Electrocompetent
cell line (*Homo sapiens*)	HEK293T	ATCC	CRL-3216	-
transfected construct (*Homo sapiens*)	Tet-Bxb1-BFP HEK293T	Laboratory of Doug Fowler	-	Clone 37 described in Ref. 18
biological sample (*Canis familiarus*)	Pancreatic rough microsomes	tRNA probes	-	-
biological sample (*Oryctolagus cuniculus*)	Reticulocyte lysate	Promega	L4960	Nuclease-treated

antibody	Anti-Hemagglutinin Antibody (Mouse monoclonal)	Invitrogen	26183-D550	2-2.2.14, DyLight™ 550 conjugate
recombinant DNA reagent	pGEM- Lep-TMD6	Ref. 6	-	-
recombinant DNA reagent	pcDNA5 CMV-HA-mGnRHR-IRES-eGFP	Ref. 6	-	-
recombinant DNA reagent	pcDNA5 attB-HA-mGnRHR-IRES-eGFP	This paper	-	-
sequence-based reagent	Custom mGnRHR primer pool	Agilent Technologies Inc.	-	See Table S3 for sequences
sequence-based reagent	Custom 10-base randomization primer	Integrated DNA Technologies	-	gcatgaagaatctgcttagggttaggcgnnnnnnnnnnncttcgc gatgtacgggccagat
peptide, recombinant protein	PrimeSTAR HS DNA Polymerase	Takara Bio Inc.	R010B	-
peptide, recombinant protein	KAPA HiFi HotStart Ready Mix	Roche Diagnostics	07958927001	-
commercial assay or kit	InFusion HD Cloning	Takara Bio Inc.	638944	-
commercial assay or kit	GenElute Mammalian gDNA Miniprep Kit	Sigma Aldrich	G1N70	-
software, algorithm	Flowjo X	Treestar	-	-
software, algorithm	OriginLab 2023	OriginLab	-	-

415
Plasmid Preparation and Characterization of Genetic Libraries

The expression of transiently expressed single mutants was carried out using a previously described pcDNA5
FRT expression vector containing mGnRHR cDNA with an N-terminal hemagglutinin (HA) epitope and bicistronic
eGFP (Fig. 2) (Chamness et al., 2021). Mutations were introduced by site-directed mutagenesis with PrimeSTAR
HS DNA Polymerase (Takara Bio, Shiga, Japan). Biochemical measurements of the membrane integration of
TMD6 were carried out using a previously described pGEM vector containing modified leader peptidase (Lep),
with TMDs of interest cloned into the H-segment (Fig. S1) (Chamness et al., 2021; Hessa et al., 2007). Mutations
were introduced into the H-segment by site-directed mutagenesis with PrimeSTAR HS DNA Polymerase (Takara
Bio, Shiga, Japan). Plasmids were purified using a ZymoPURE Midiprep Kit (Zymo Research, Irvine, CA). DMS
experiments were carried out using a previously described pcDNA5 FRT vector containing mGnRHR cDNA
bearing an N-terminal influenza hemagglutinin (HA) epitope, followed by an internal ribosome entry site (IRES)
and eGFP sequence, which was further modified to be compatible with recombination-based DMS approaches
(Chamness et al., 2021). First, an attB recombination site was inserted in place of the CMV promoter by InFusion
HD Cloning (Takara Bio, Shiga, Japan). To facilitate the creation of these libraries using nicking mutagenesis, a
BbvCI restriction site was introduced by site-directed mutagenesis using PrimeSTAR HS DNA Polymerase
(Takara Bio, Shiga, Japan). The V276T and W107A mutations were also introduced by site-directed mutagenesis
with PrimeSTAR HS DNA Polymerase (Takara Bio, Shiga, Japan).
434

To generate mutational libraries, a 10N unique molecular identifier (UMI), or "barcode," was first inserted into the
plasmid backbone using a previously described nicking mutagenesis method (Wrenbeck et al., 2016). Nicking
mutagenesis plasmid products were transformed into NEB 10-beta electrocompetent cells (New England
Biolabs, Ipswich, MA) and purified using a ZymoPURE Midiprep Kit (Zymo Research, Irvine, CA). A programmed
oligo pool (Agilent, Santa Clara, CA) encoding all 19 amino acid substitutions at 85 residues throughout the
mGnRHR sequence was then used to generate pools of single and double mutants in the context of WT, V276T,
and W107A mGnRHR cDNA using a previously described nicking mutagenesis method (Medina-Cucurella et
al., 2019). The resulting plasmid products were transformed into NEB 10-beta electrocompetent cells (New
England Biolabs, Ipswich, MA) and purified using a ZymoPURE Midiprep Kit (Zymo Research, Irvine, CA). To
limit the number of mGnRHR variants per UMI, these libraries were then bottlenecked through an additional
transformation in NEB Turbo electrocompetent cells (New England Biolabs, Ipswich, MA), and were again
purified using a ZymoPURE Midiprep Kit (Zymo Research, Irvine, CA). The resulting plasmid preparations
contained detectable levels of a concatemer product, which were removed from the mutational libraries by gel
purification using a Zymoclean Gel DNA Recovery Kit (Zymo Research, Irvine, CA). Purified products were again
transformed into either NEB 10-beta (New England Biolabs, Ipswich, MA) or SIG10 (Sigma-Aldrich, St. Louis,
MO) electrocompetent cells, and purified with a ZymoPURE Midiprep Kit (Zymo Research, Irvine, CA).
451

To associate each UMI with its corresponding GnRHR variant, the final plasmid libraries were also sequenced
using PacBio SMRT sequencing. Briefly, plasmid libraries were first double-digested with PmlI and MfeI-HF
restriction enzymes (New England Biolabs, Ipswich, MA), then purified using a Zymo DNA Clean & Concentrator-
5 kit (Zymo Research, Irvine, CA). Complete digestion was verified by analyzing the products on an agarose gel,
purity was assessed with a Synergy Neo2 microplate reader (BioTek, Winooski, VT), and the final yield was
quantified using a Qubit4 fluorometer (ThermoFisher, Waltham, MA). The 1537 bp fragment for each library,
containing the 10N UMI and GnRHR open reading frame, was isolated on a BluePippin instrument (Sage
Science, Beverly, MA). These fragments were then assembled into PacBio libraries and sequenced on the
Sequel II system with a 30-hour runtime.
461

The sequence of each plasmid-derived fragment containing mGnRHR variants and their corresponding UMIs
was determined by reconstructing the circular consensus sequences (CCS) for each independent well in the
PacBio readout. mGnRHR libraries were sequenced to a depth of 1,825,010 wells for WT, 1,545,151 wells for
V276T, and 1,497,359 wells for W107A. A minimum coverage of five passes around each circular fragment
(subread) was required for incorporation of sequencing data from an individual well into the analysis. CCS were
mapped to the reference open reading frame using minimap2. CCS that did not contain a complete 10-base UMI
sequence or fully cover the target ORF were excluded from the analysis. Variations in the mGnRHR sequence
of individual fragments were called using SAMtools. Reads were then grouped according to their UMI sequences.
Given that insertion and deletion (indel) artifacts are prevalent within PacBio sequencing data, we only removed
UMIs from the analysis if indels were detected at a specific position within the UMI in at least 10% of the
subreads. Indels observed within the open reading frame were only called if their occurrence was judged to be
statistically significant relative to the background indel rate for the reads according to a binomial test ($p <= .05$).
Codon substitutions encoded in our primer library (Table S3) were called within the mGnRHR open reading

frame only if they were detected in at least 75% of the individual subreads. Only UMIs that were found to be associated with a single mGnRHR variant were included in the final analysis. Our final "dictionaries" for the WT, V276T, and W107A libraries contained 3,261, 531, and 4,962 unique UMIs, respectively, that could be indexed to a corresponding mGnRHR variant.

Preparation of Cellular Libraries and Cell Sorting

A previously described HEK293T cell line with a genomic Tet-Bxb1-BFP landing pad was obtained from D. Fowler (Matreyek et al., 2017). Cells were grown at 37 °C and 5.0% CO_2 in complete media, made of Dulbecco's Modified Eagle's medium (DMEM, Gibco, Grand Island, NY) supplemented with 10% fetal bovine serum (FBS, Gibco, Grand Island, NY), 0.5% penicillin (Gibco, Grand Island, NY), and 0.5% streptomycin (Gibco, Grand Island, NY). 2 million cells were plated in 10-cm tissue culture plates and co-transfected the next day with 475 ng of plasmid encoding bxb1 recombinase and 7125 ng of the plasmid library using Lipofectamine 3000 (Invitrogen, Carlsbad, CA). Cells were grown at 33 °C for 4 days after transfection. 2 days after transfection, the cells were induced with 2 µg/mL doxycycline. All cells were expanded into 15-cm tissue culture plates 6 days after transfection, and 10 million cells were re-plated into new 15-cm tissue culture plates 3 days later.

Twelve days after transfection, the cells were washed with 1X phosphate-buffered saline (PBS, Gibco, Grand Island, NY), and harvested with TrypLE Express protease (Gibco, Grand Island, NY). Cells were washed twice with 2% fetal bovine serum in PBS (wash buffer), resuspended in PBS containing 2% FBS to 10M cells/ mL, and then sorted on a BD FACS Aria II (BD Biosciences, Franklin Lakes, NJ). Recombined cells were isolated based on their characteristic gain of bicistronic eGFP and loss of BFP expression (Matreyek et al., 2017). Forward and side scatter profiles were first used to gate for intact live cells, and cells with eGFP-positive (488 nm laser, 530/30 nm emission filter) and BFP-negative (405 nm laser, 450/50 nm emission filter) fluorescence profiles were collected in complete media supplemented with 10% FBS (Gibco, Grand Island, NY) and then plated in 10-cm tissue culture plates. Cells were induced with 2 µg/mL doxycycline 24 hours after sorting. 4 days after sorting, 10 million cells were passaged into 15-cm tissue culture plates.

7 days after sorting for eGFP-positive cells, cells were sorted again according to the surface expression of GnRHR variants. Briefly, cells were first washed with 1X phosphate-buffered saline (PBS, Gibco, Grand Island, NY), and harvested with TrypLE Express protease (Gibco, Grand Island, NY). GnRHRs expressed at the cell surface were immunostained for 30 minutes in the dark with DyLight 550-conjugated anti-HA antibody (Invitrogen, Carlsbad, CA). Cells were then washed twice with PBS containing 2% FBS, and sorted on a BD FACS Aria II (BD Biosciences, Franklin Lakes, NJ). Forward and side scatter profiles were used to gate for intact live cells, and eGFP fluorescence was used to gate for recombined cells. Cells were then sorted into even quartiles based on the amount of DyLight 550 fluorescence (561 nm laser, 585/15 nm emission filter). Sorted cells were collected in complete media supplemented with 10% FBS (Gibco, Grand Island, NY) and plated in 10-cm tissue culture plates. Cells were allowed to grow to confluency and then were washed with 1X phosphate-buffered saline (PBS, Gibco, Grand Island, NY) and harvested with TrypLE Express protease (Gibco, Grand Island, NY). Cell pellets were frozen for subsequent analysis.

Extraction of Genomic DNA and Next Generation Sequencing

Genomic DNA (gDNA) was extracted from cell pellets using the Sigma GenElute Mammalian gDNA Miniprep Kit (Sigma-Aldrich, St. Louis, MO). DNA amplicons for Illumina sequencing were produced from the gDNA using a previously described semi-nested polymerase chain reaction (PCR) technique (Matreyek et al., 2017). In order to selectively amplify recombined DNA, the first PCR utilized a primer which anneals upstream of the 10N UMI and a primer which anneals in the BFP-encoding region of the landing pad. Eight replicate PCRs were carried out with 2.5 µg of gDNA and HiFi HotStart Ready Mix (Kapa Biosystems, Wilmington, MA). These PCRs were limited to 7 cycles to avoid PCR bias. PCR products were purified using a ZR-96 DNA Clean & Concentrator-5 kit (Zymo Research, Irvine, CA), and replicate products were then pooled. 10 µL of this first PCR product was then used as the template for a second PCR, which utilized primers that incorporate Illumina adapter sequences to both ends of the 10N UMI. To avoid PCR bias, these reactions were monitored by real-time PCR on a StepOne RT-PCR instrument (Applied Biosystems, Waltham, MA) and terminated during mid-log amplification (17 or 18 cycles). Four replicate PCRs were carried out with HiFi HotStart Ready Mix (Kapa Biosystems, Wilmington, MA) and SYBR Green fluorescent dye (Applied Biosystems, Waltham, MA). Replicate PCR products were then combined and purified with a Zymoclean Gel DNA Recovery Kit (Zymo Research, Irvine, CA). To further improve the yield and quality of the amplicons, an additional 6-cycle PCR was carried out with the HiFi HotStart Ready Mix (Kapa Biosystems, Wilmington, MA) and primers that preserve the DNA sequence. These PCR products

were then purified using a Zymo DNA Clean & Concentrator-5 kit (Zymo Research, Irvine, CA). Finally, amplicon yield and quality were assessed using an Agilent 2200 TapeStation with D1000 tape (Agilent Technologies, Santa Clara, CA) and a Synergy Neo2 microplate reader (BioTek, Winooski, VT). Amplicons were sequenced using an Illumina NextSeq150 flow cell, paired end, with 50% PhiX content.

Plasma Membrane Expression and Epistasis Measurements

Illumina sequencing of the recombined UMIs within each sorted cell fraction were matched to the corresponding GnRHR mutants using the PacBio sequencing data and subjected to previously described quality filtering (Penn et al., 2020). Estimations of PME for each mutant were then calculated as previously described (Penn et al., 2020). PME measurements were averaged across two biological replicates. Within each library, the average fluorescence intensity values of the mutants were ranked and converted into a percentile. Mutants which differed in their percentile values by more than 25% across the two replicates were excluded from analysis. The use of percentile-based filtering overcomes intrinsic limitations associated with the distinct dynamic range constraints that occur because of the distinct surface immunostaining profiles of these three cellular libraries (Fig. 3). Intensity values that meet these criteria are highly correlated across the two biological replicates (average Spearman's ρ=0.936 across the three libraries, Supplementary Figs. S3 & S4). To facilitate the comparison of epistatic interactions within a common set of secondary mutations, we excluded mutants that did not pass quality filtering in all three libraries from our analysis. After data filtering, we obtained PME measurements for 251 mutants in all three libraries (Table S 4-6). These PME measurements were then utilized to calculate epistasis scores for the double mutants in the V276T and W107A libraries using the following equation:

$$\varepsilon = \ln(\omega_{ab}) - \ln(\omega_a) - \ln(\omega_b)$$

where $\mathcal{E}$ is the epistasis score, ω_{ab} is the fluorescence intensity of the double mutant relative to WT mGnRHR, ω_a is the fluorescence intensity of the single mutant in the WT library relative to WT mGnRHR, and ω_b is the fluorescence intensity of the background mutation (either V276T or W107A) relative to WT mGnRHR, as previously described (Olson et al., 2014).

Expression Measurements of Individual GnRHR Variants

HEK283T cells were obtained from ATCC and grown under the same conditions described above. GnRHR variants were transiently expressed in these cells, and the plasma membrane and intracellular expression of these variants was measured by flow cytometry, as previously described (Chamness et al., 2021). The reported expression levels represent measurements from three biological replicates.

In vitro Translation of Chimeric Lep Proteins

Messenger RNA (mRNA) of chimeric Lep proteins was generated and used for *in vitro* translation reactions as previously described (Chamness et al., 2021). Translation products were then analyzed by SDS-PAGE as previously described (Chamness et al., 2021). The intensities of singly (*G1*) and doubly (*G2*) glycosylated protein were quantified by densitometry in ImageJ software. Apparent transfer free energy values representing transfer of the TMDs of interest from the translocon to the membrane were then calculated using the following equation:

$$\Delta G_{app} = -RTln(K_{app}) = -RTln(\frac{G1}{G2})$$

where ΔG_{app} is the free energy for transfer of the TMD into the membrane, R is the universal gas constant, T is the temperature, K_{app} is the equilibrium constant for transfer of the TMD into the membrane, $G1$ is the intensity of the singly glycosylated band, and $G2$ is the intensity of the doubly glycosylated band, as previously described (Hessa et al., 2005). The reported transfer free energy values represent the average of three experimental replicates.

Structural Modeling of M. musculus GnRHR

The sequence of mGnRHR was obtained from Uniprot (accession number Q01776) and a homology model in the inactive state was generated as previously described (Chamness et al., 2021). Images were generated by structurally aligning this mGnRHR model to an experimentally determined inactive-state human GnRHR crystal structure (GNRHR, PDB 7BR3, 2.79 Å) using the Super command in PyMol (Schrödinger, Inc., New York, NY) (Yan et al., 2020).

Computational Estimates for the Stability of mGnRHR Variants

Structural estimates for the effects of mutations on the stability of mGnRHR were determined using a previously described Rosetta based protocol featuring a membrane protein optimized energy function (Alford et al., 2015; Barlow et al., 2018). Briefly, the homology model of mGnRHR described above was used as the starting structure for computational stability estimates. A spanfile describing the transmembrane regions was created for mGnRHR using OCTOPUS predictions from Topcons (Bernsel et al., 2009; Viklund and Elofsson, 2008). The homology model was transformed into membrane coordinates using the mp_transform application. An ensemble of structures of each variant and wildtype (50 iterations) was generated from the homology model. For variants containing two mutations, the mutations were both introduced at the same time. The $\Delta\Delta G$ for each variant was calculated by subtracting the average score of the three lowest scoring variant models from the average score of the three lowest scoring WT models ($\Delta\Delta G = \Delta G_{mut} - \Delta G_{wt}$).

Unsupervised Learning

To classify mGnRHR variants based on their expression profiles, we first standardized their scores using the Standard Scaler tool from Python's scikit-learn library in order to ensure that the clustering algorithms are not affected by the scale of the data. Next, we then reduced the dimensionality of the standardized data using UMAP (McInnes et al., 2018). We then used a density-based clustering algorithm (HDBSCAN) to cluster variants based on their expression profiles (McInnes et al., 2017). Unsupervised learning analyses were carried out based on deep mutational scanning measurements alone, and did not incorporate any structural and/ or energetic parameters.

Quantification and Analysis

Flow cytometry data were analyzed in FlowJo software (Treestar, Ashland, OR), and *in vitro* translation data were analyzed in ImageJ software. PME measurements were analyzed in Origin software (OriginLab, Northampton, MA) and mapped onto the mGnRHR homology model in PyMol (Schrödinger, Inc., New York, NY).

Data Availability

Code for the analysis of sequencing data for deep mutational scanning experiments can be found on the Schlebach Lab GitHub page (https://github.com/schebachlab/gnrhr_epistasis_dms). Illumina sequencing data is freely available through NCBI (https://www.ncbi.nlm.nih.gov/bioproject/PRJNA1062836). Fluorescence intensity values for mutants in all three mutational libraries are included as Excel files in the Supplementary Materials. Additional data is available upon request. All materials described herein will be made freely available by the corresponding author upon request.

Acknowledgements

We thank Christiane Hassel and the Indiana University Bloomington Flow Cytometry Core Facility, as well as Luke Tallon, Xuechu Zhao, Lisa Sadzewicz, and the University of Maryland Institute for Genome Sciences for their experimental support. This work was supported by a National Science Foundation (NSF) Graduate Research Fellowship (1342962 to L. M. C.) and by the National Institutes of Health (R01GM129261 to J.P.S). Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the NSF.

References

Alford RF, Koehler Leman J, Weitzner BD, Duran AM, Tilley DC, Elazar A, Gray JJ. 2015. An Integrated Framework Advancing Membrane Protein Modeling and Design. *PLoS Comput Biol* **11**. doi:10.1371/journal.pcbi.1004398

Barlow KA, Ó Conchúir S, Thompson S, Suresh P, Lucas JE, Heinonen M, Kortemme T. 2018. Flex ddG: Rosetta Ensemble-Based Estimation of Changes in Protein–Protein Binding Affinity upon Mutation. *J Phys Chem B* **122**:5389–5399. doi:10.1021/acs.jpcb.7b11367

Bernsel A, Viklund H, Hennerdal A, Elofsson A. 2009. TOPCONS: consensus prediction of membrane protein topology. *Nucleic Acids Res* **37**:W465-8. doi:10.1093/nar/gkp363

Bershtein S, Mu W, Serohijos AWR, Zhou J, Shakhnovich EI. 2013. Protein Quality Control Acts on Folding Intermediates to Shape the Effects of Mutations on Organismal Fitness. *Mol Cell* **49**:133–144. doi:10.1016/j.molcel.2012.11.004

659 Bloom JD, Raval A, Wilke CO. 2007. Thermodynamics of neutral protein evolution. *Genetics* **175**:255–266.
660 doi:10.1534/genetics.106.061754
661

662 Campello RJGB, Moulavi D, Sander J. 2013. Density-Based Clustering Based on Hierarchical Density Estimates
663 In: Advances in Knowledge Discovery and Data Mining. Springer. pp. 160–172. doi:10.1007/978-3-642-
664 37456-2
665

666 Chamness LM, Zelt NB, Harrington HR, Kuntz CP, Bender BJ, Penn WD, Ziarek JJ, Meiler J, Schlebach JP.
667 2021. Molecular basis for the evolved instability of a human G-protein coupled receptor. *Cell Rep* **37**.
668 doi:10.1016/J.CELREP.2021.110046
669

670 Chen Z, Mondal A, Abderemane-Ali F, Jang S, Niranjan S, Montaño JL, Zaro BW, Minor DL. 2023. EMC
671 chaperone–CaV structure reveals an ion channel assembly intermediate. *Nature* **619**:410–419.
672 doi:10.1038/s41586-023-06175-5
673

674 Faber MS, Wrenbeck EE, Azouz LR, Steiner PJ, Whitehead TA. 2019. Impact of in Vivo Protein Folding
675 Probability on Local Fitness Landscapes. *Mol Biol Evol* **36**:2764–2777. doi:10.1093/molbev/msz184
676

677 Gemmer M, Chaillet ML, van Loenhout J, Cuevas Arenas R, Vismpas D, Gröllers-Mulderij M, Koh FA, Albanese
678 P, Scheltema RA, Howes SC, Kotecha A, Fedry J, Förster F. 2023. Visualization of translation and protein
679 biogenesis at the ER membrane. *Nature* **614**:160–167. doi:10.1038/s41586-022-05638-5
680

681 Ghosh K, Dill K. 2010. Cellular proteomes have broad distributions of protein stability. *Biophys J* **99**:3996–4002.
682 doi:10.1016/j.bpj.2010.10.036
683

684 Gong LI, Suchard MA, Bloom JD. 2013. Stability-mediated epistasis constrains the evolution of an influenza
685 protein. *Elife* **2013**:1–19. doi:10.7554/eLife.00631
686

687 Hermansson M, von Heijne G. 2003. Inter-helical hydrogen bond formation during membrane protein integration
688 into the ER membrane. *J Mol Biol* **334**:803–9. doi:10.1016/j.jmb.2003.10.019
689

690 Hessa T, Kim H, Bihlmaier K, Lundin C, Boekel J, Andersson H, Nilsson I, White SH, von Heijne G. 2005.
691 Recognition of transmembrane helices by the endoplasmic reticulum translocon. *Nature* **433**: 377–381. doi:
692 10.1038/nature03216
693

694 Hessa T, Meindl-Beinker NM, Bernsel A, Kim H, Sato Y, Lerch-Bader M, Nilsson I, White SH, Gunnar Von Heijne
695 &. 2007. Molecular code for transmembrane-helix recognition by the Sec61 translocon. *Nature* **450**.
696 doi:10.1038/nature06387
697

698 Janovick JA, David Stewart M, Jacob D, Martin LD, Deng JM, Stewart A, Wang Y, Cornea A, Chavali L, Lopez
699 S, Mitalipov S, Kang E, Lee H-S, Manna PR, Stocco DM, Behringer RR, Conn PM. 2013. Restoration of
700 testis function in hypogonadotropic hypogonadal mice harboring a misfolded GnRHR mutant by
701 pharmacoperone drug therapy. *Proc Natl Acad Sci U S A* **110**:21030–21035. doi:10.1073/pnas.1315194110
702

703 Janovick JA, Knollman PE, Brothers SP, Ayala-Yáñ Ez R, Aziz AS, Conn PM. 2006. Regulation of G Protein-
704 coupled Receptor Trafficking by Inefficient Plasma Membrane Expression Molecular Basis of an Evolved
705 Strategy. *J Biol Chem* **281**:8417–8425. doi:10.1074/jbc.M510601200
706

707 Jones EM, Lubock NB, Venkatakrishnan AJ, Wang J, Tseng AM, Paggi JM, Latorraca NR, Cancilla D, Satyadi
708 M, Davis JE, Babu MM, Dror RO, Kosuri S. 2020. Structural and functional characterization of G protein-
709 coupled receptors with deep mutational scanning. *Elife* **9**:1–52. doi:10.7554/ELIFE.54895
710

711 Marinko JT, Huang H, Penn WD, Capra JA, Schlebach JP, Sanders CR. 2019. Folding and Misfolding of Human
712 Membrane Proteins in Health and Disease: From Single Molecules to Cellular Proteostasis. *Chem Rev*
713 **119**:5537–5606. doi:10.1021/acs.chemrev.8b00532
714

715 Matreyek KA, Stephany JJ, Fowler DM. 2017. A platform for functional assessment of large variant libraries in
716 mammalian cells. *Nucleic Acids Res* **45**:1–12. doi:10.1093/nar/gkx183
717

718 McInnes L, Healy J, Astels S. 2017. hdbscan: Hierarchical density based clustering. *J Open Source Softw* **2**:205.
719 doi:10.21105/joss.00205
720

721 McInnes L, Healy J, Saul N, Großberger L. 2018. UMAP: Uniform Manifold Approximation and Projection. *J
722 Open Source Softw* **3**:861. doi:10.21105/joss.00861

Medina-Cucurella A v., Steiner PJ, Faber MS, Beltrán J, Borelli AN, Kirby MB, Cutler SR, Whitehead TA. 2019. User-defined single pot mutagenesis using unamplified oligo pools. *Protein Eng Des Sel* **32**:41–45. doi:10.1093/protein/gzz013

Meindl-Beinker NM, Lundin C, Nilsson I, White SH, von Heijne G. 2006. Asn- and Asp-mediated interactions between transmembrane helices during translocon-mediated membrane protein assembly. *EMBO Rep* **7**:1111–6. doi:10.1038/sj.embor.7400818

Moon CP, Fleming KG. 2011. Side-chain hydrophobicity scale derived from transmembrane protein folding into lipid bilayers. *Proc Natl Acad Sci U S A* **108**:10174–7. doi:10.1073/pnas.1103979108

Morrison AJ, Wonderlick DR, Harms MJ. 2021. Ensemble epistasis: Thermodynamic origins of nonadditivity between mutations. *Genetics* **219**:iyab105. doi:10.1093/GENETICS/IYAB105

Nedrud D, Coyote-Maestas W, Schmidt D. 2021. A large-scale survey of pairwise epistasis reveals a mechanism for evolutionary expansion and specialization of PDZ domains. *Proteins* **89**:899–914. doi:10.1002/PROT.26067

Öjemalm K, Halling KK, Nilsson IM, von Heijne G. 2012. Orientational Preferences of Neighboring Helices Can Drive ER Insertion of a Marginally Hydrophobic Transmembrane Helix. *Mol Cell* **45**:529–540. doi:10.1016/j.molcel.2011.12.024

Olson CA, Wu NC, Sun R. 2014. A comprehensive biophysical description of pairwise epistasis throughout an entire protein domain. *Curr Biol* **24**:2643–2651. doi:10.1016/j.cub.2014.09.072

Otto SP, Feldman MW. 1997. Deleterious mutations, variable epistatic interactions, and the evolution of recombination. *Theor Popul Biol* **51**:134–147.

Penn WD, McKee AG, Kuntz CP, Woods H, Nash V, Gruenhagen TC, Roushar FJ, Chandak M, Hemmerich C, Rusch DB, Meiler J, Schlebach JP. 2020. Probing Biophysical Sequence Constraints within the Transmembrane Domains of Rhodopsin by Deep Mutational Scanning. *Sci Adv* **6**:aay7505. doi:10.1126/sciadv.aay7505

Pfeffer S, Dudek J, Zimmermann R, Förster F. 2016. Organization of the native ribosome–translocon complex at the mammalian endoplasmic reticulum membrane. *Biochim Biophys Acta* **1860**:2122–2129. doi:10.1016/j.bbagen.2016.06.024

Pleiner T, Tomaleri GP, Januszyk K, Inglis AJ, Hazu M, Voorhees RM. 2020. Structural basis for membrane insertion by the human ER membrane protein complex. *Science* **369**:433–436. doi:10.1126/science.abb5008

Popot JL, Engelman DM. 1990. Membrane Protein Folding and Oligomerization. *Biochemistry* **29**:4031–4037. doi:10.1021/bi00469a001

Rodrigues J V., Bershtein S, Li A, Lozovsky ER, Hartl DL, Shakhnovich EI. 2016. Biophysical principles predict fitness landscapes of drug resistance. *Proc Natl Acad Sci U S A* **113**:E1470–E1478. doi:10.1073/pnas.1601441113

Sailer ZR, Harms MJ. 2017. Detecting high-order epistasis in nonlinear genotype-phenotype maps. *Genetics* **205**:1079–1088. doi:10.1534/genetics.116.195214

Schlebach JP, Sanders CR. 2016. The Safety Dance: Biophysics of Membrane Protein Folding and Misfolding in a Cellular Context. *Q Rev Biophys* **48**:1–34. doi:10.1017/S0033583514000110.

Shurtleff MJ, Itzhak DN, Hussmann JA, Schirle Oakdale NT, Costa EA, Jonikas M, Weibezahn J, Popova KD, Jan CH, Sinitcyn P, Vembar SS, Hernandez H, Cox J, Burlingame AL, Brodsky JL, Frost A, Borner GH, Weissman JS. 2018. The ER membrane protein complex interacts cotranslationally to enable biogenesis of multipass membrane proteins. *Elife* **7**:e37018. doi:10.7554/eLife.37018

Smalinskaite L, Hegde RS. 2022. The Biogenesis of Multipass Membrane Proteins. *Cold Springs Harb Persp Biol* **15**:a041251. doi:10.1101/cshperspect.a041251

Smalinskaitė L, Kim MK, Lewis AJO, Keenan RJ, Hegde RS. 2022. Mechanism of an intramembrane chaperone for multipass membrane proteins. *Nature* **611**:161-166. doi:10.1038/s41586-022-05336-2

Song Y, DiMaio F, Yu-Ruei Wang R, Kim D, Miles C, Brunette T, Thompson J, Baker D. 2013. High resolution comparative modeling with RosettaCM. *Structure* **21**:1735-1742. doi:10.1016/j.str.2013.08.005

Starr TN, Thornton JW. 2016. Epistasis in protein evolution. *Protein Sci* **25**:1204–1218. doi:10.1002/pro.2897

Sundaram A, Yamsek M, Zhong F, Hooda Y, Hegde RS, Keenan RJ. 2022. Substrate-driven assembly of a translocon for multipass membrane proteins. *Nature* **611**:167–172. doi:10.1038/s41586-022-05330-8

Telenti A, Pierce LCT, Biggs WH, di Iulio J, Wong EHM, Fabani MM, Kirkness EF, Moustafa A, Shah N, Xie C, Brewerton SC, Bulsara N, Garner C, Metzker G, Sandoval E, Perkins BA, Och FJ, Turpaz Y, Venter JC. 2016. Deep sequencing of 10,000 human genomes. *Proc Natl Acad Sci U S A* **113**:11901–11906. doi:10.1073/pnas.1613365113

Tokuriki N, Stricher F, Schymkowitz J, Serrano L, Tawfik DS. 2007. The Stability Effects of Protein Mutations Appear to be Universally Distributed. *J Mol Biol* **369**:1318–1332. doi:10.1016/j.jmb.2007.03.069

Tokuriki N, Tawfik DS. 2009. Stability effects of mutations and protein evolvability. *Curr Opin Struct Biol* **19**:596–604. doi:10.1016/j.sbi.2009.08.003

van den Berg B, Clemons WM, Collinson I, Modis Y, Hartmann E, Harrison SC, Rapoport TA. 2003. X-ray structure of a protein-conducting channel. *Nature* **427**:36–44. doi: 10.1038/nature02218

Viklund H, Elofsson A. 2008. OCTOPUS: improving topology prediction by two-track ANN-based preference scores and an extended topological grammar. *Bioinformatics* **24**:1662–1668. doi:10.1093/bioinformatics/btn221

White SH, Von Heijne G. 2008. How Translocons Select Transmembrane Helices. *Annu Rev Biophys* **37**:23-42. doi:10.1146/annurev.biophys.37.032807.125904

Wiseman RL, Koulov A, Powers E, Kelly JW, Balch WE. 2007. Protein energetics in maturation of the early secretory pathway. *Curr Opin Cell Biol* **19**:359. doi:10.1016/J.CEB.2007.05.005

Woodall NB, Hadley S, Yin Y, Bowie JU. 2017. Complete topology inversion can be part of normal membrane protein biogenesis. *Protein Sci* **26**:824–833. doi:10.1002/pro.3131

Wrenbeck EE, Klesmith JR, Stapleton JA, Adeniran A, Tyo KEJ, Whitehead TA. 2016. Plasmid-based one-pot saturation mutagenesis. *Nat Methods* **13**:928–930. doi:10.1038/nmeth.4029

Yan W, Cheng L, Wang W, Wu C, Yang X, Du X, Ma L, Qi S, Wei Y, Lu Z, Yang S, Shao Z. 2020. Structure of the human gonadotropin-releasing hormone receptor GnRH1R reveals an unusual ligand binding mode. *Nat Commun* **11**:5287. doi:10.1038/s41467-020-19109-w

Zeldovich KB, Chen P, Shakhnovich EI. 2007. Protein stability imposes limits on organism complexity and speed of molecular evolution. *Proc Natl Acad Sci U S A* **104**:16152-16157. doi:10.1073/pnas.0705366104